Role of Snow Depth in the Influence of El Niño on Summer Climate Anomalies over East Asia

Xuguang Sun[1], Yanfeng Wang[2], and Xiu-Qun Yang[1]

[1]Nanjing University
[2]Institute of Atmospheric Physics

November 24, 2022

Abstract

With datasets of Global Land Data Assimilation System (GLDAS) NOAH land surface model, GPCC monthly mean rainfall and NCAR/NCEP global monthly mean reanalysis from 1948 to 2010, by using methods of filtering, composite and linear regression and correlation, characteristics of Eurasian snow depth anomalies in El Niño mature winter, its influences on soil moisture after snow melting, and finally on East Asian summer monsoon are investigated, and the main conclusions are as follows: In El Niño mature winter, snow depth in regions of the Iranian Plateau, the northeast of Lake Balkhash and the southern Tibetan Plateau increase remarkably, so are the related snow melting and soil moisture. The above-mentioned three regions are identified as the key regions for snow depth to store and extend the El Niño signals. In spring, the snow begins to melt, and the soil moisture increases correspondingly, thus the El Niño signals are transmitted from winter snow depth to soil moisture in spring. As a result, sensible heat flux decreases and latent heat flux increases, and the atmospheric circulations are greatly influenced. The anomalous soil moisture in the Iranian plateau is most important for the East Asian summer monsoon in El Niño decaying summer, since it has similar impact pattern on the anomalous summer precipitation as the El Niño composite. The spring and summer soil moisture in both the southern Tibetan plateau and the northeast of Lake Balkhash increase simultaneously, which significantly contribute to the increased precipitation in North China. Therefore, to investigate and predict the East Asian summer monsoon variabilities by using El Niño signal, the roles of snow depth in storing and modulating El Niño impacts in those key regions should be considered.

Role of snow depth in the influence of El Niño on summer climate anomalies over East Asia

Yanfeng Wang, **Xuguang Sun*** and Xiu-Qun Yang

*LCPS, School of Atmospheric Sciences, **Nanjing University**, Nanjing 210023, China * Contact E-mail: **xgsun@nju.edu.cn***

1. Motivation



Questions:

- What are the key regions of snow depth anomalies in response to the winter El Niño forcing?
- How do the El Niño-related winter snow depth anomalies in key regions influence the East Asian summer climate anomalies?

2. Data and Method

SST dataset: NOAA ERSSTv3 2°×2° 1948-2010

Definition of an El Niño event:

niño3 index is greater than or equal to 0.5 °C for any five months during the period from September to the next May

List of 18 El Niño events:

1951-1952, 1953-1954, 1957-1958, 1963-1964, 1965-1966, 1968-1969, 1969-1970, 1972-1973, 1976-1977, 1982-1983, 1986-1987, 1987-1988, 1991-1992, 1997-1998, 2002-2003, 2004-2005, 2006-2007, 2009-2010

Land dataset: GLDAS_NOAH025 0.25°×0.25° 1948-2010

GLADS (Global Land Data Assimilation System) data is a product of data assimilation of satellite data, land surface model and ground observational data *(Zhu and Shi, 2014)*, including 4 soil layers: 0-10 cm, 10-40 cm, 40 cm-1m and 1m-2m.

Variables: soil moisture, soil temperature, snow depth, snow melting, sensible heat flux, latent heat flux

Precipitation dataset: GPCCv7 0.5°×0.5° 1948-2010

Atmosphere Reanalysis dataset: NCEP/NCAR reanalysis 2.5°×2.5° 1948-2010

Methodology:

band-pass filtering at interannual time scale (13-month to 8-year)

Composite, linear regression/correlation

3. Results



Fig.1: Composites anomalies in El Niño mature winter (DJF) based on the 18 El Niño events. The crossed areas indicate significant correlation at 0.1 significance level. Black boxes indicate the three key regions, i.e., **Iranian Plateau**, **northeast of Lake Balkhash** and **southern Tibetan Plateau**.

Three key regions of snow depth anomalies in El Nino mature winter (DJF)

Key regions	Correlation with niño3	Cases in El Niño events
Iranian Plateau	0.42	9/18 (50%)
Northeast of Lake Balkhash	0.38	7/18 (39%)
Southern Tibetan Plateau	0.37	5/18 (28%)



Fig.2: Evolutions of composite area-averaged anomalies of snow depth (snowd, unit: kg/m^2), snow melting (snowm, unit: kg/m^2), soil moisture (unit: kg/m^2), sensible heat flux (shflx, unit: Wm^2) and latent heat flux (lhflx, unit: Wm^2) for Iranian plateau (a), northeast of Lake Balkhash (b) and southern Tibetan plateau regions in El Niño life time.



Fig.3: El Niño composite and regressed anomalies of summer atmospheric variables on the standardized time series of simultaneous area-averaged second-layer soil moisture anomalies in the three key regions.

4. Summary

- Three key regions, i.e., Iranian Plateau, northeast of Lake Balkhash and southern Tibetan Plateau, of snow depth anomalies are identified associated with El Niño events.
- Major processes: El Niño → increased snow depth (DJF) → increased soil moisture due to snow melting (DJF-to-JJA) → increased local precipitation due to positive feedback between soil moisture and rainfall *(Pal and Eltahir 2003)* → atmospheric circulation and precipitation anomalies over East Asia due to the key-region rainfall-induced diabatic heating forcing.

Wang YF, **XG Sun***, X-Q Yang, 2017: Role of snow depth in the influence of El Niño on summer climate anomalies over East Asia. *Chinese Journal of Geophysics* (in Chinese), 60(9), 3325-3337. doi: 10.6038/cjg20170905.