Behavior of phengite at high temperature and high pressure: In situ IR and Raman spectroscopic studies

WENDI Liu¹, Yan Yang², Zeming Qi³, Zhongping Wang³, Weihua Huang¹, and Qunke Xia¹

¹Zhejiang University ²School of Earth Science, Zhejiang University ³University of Science and Technology of China

November 22, 2022

Abstract

Water release in subduction zones is not only an important part of the deep Earth's water cycle, but also plays an essential role in the physical and chemical properties of rocks constituting the deep Earth. To understand water release processes, it is important to know properties of dehydration in hydrous phases of the downgoing slab. Although it is widely accepted that phengite can be stable to greater depth in subduction environment, behavior of hydroxyl and lattice of it at high temperature and high pressure are less investigated in contrast to other hydrous phases. Here, using IR and Raman spectroscopy, we characterize hydroxyl and lattice of ammonium-bearing and ammonium-free phengite at high temperature and high pressure. No proton transferring and structural phase transition in phengite were observed over the measured temperature and pressure range. Both pressure and temperature induce hydroxyl band shifting to lower frequencies, and pressure has a greater impact. The band width of hydroxyl increases with temperature and pressure. Hydroxyl bond weakening and hydrogen disordering at high temperature and high pressure should be responsible for the spectra variations. On the other hand, the lattice modes soften with increasing temperature whereas stiffen under compression, and ammonium plays an important role in the Grüneisen parameters of the lattice modes, especially the K-O mode. These features of hydroxyl and lattice at high temperature and high pressure could benefit for further understanding dehydration, thermodynamic properties and stability of phengite in subduction zones.

Hosted file

agusupporting-information_word_template.docx available at https://authorea.com/users/535924/ articles/598898-behavior-of-phengite-at-high-temperature-and-high-pressure-in-situ-irand-raman-spectroscopic-studies

1	AGU Word Manuscript Template
2 3 4	Behavior of phengite at high temperature and high pressure: In situ IR and Raman spectroscopic studies
5	
6 7	Wendi Liu ¹ , Yan Yang ¹ *, Zeming Qi ² , Zhongping Wang ³ , Weihua Huang ¹ , Qun-Ke Xia ¹ ¹ School of Earth Sciences, Zhejiang University, Hangzhou 310027, China.
8 9	² National Synchrotron Radiation Laboratory, University of Science and Technology of China, Hefei, Anhui 230026, China.
10	³ Department of Physics, University of Science and Technology of China, Hefei, Anhui 230026, China.
11	Corresponding author: Yan Yang (<u>yanyang2005@zju.edu.cn)</u>
12	Key Points:
13 14	• Vibrations of hydroxyl and lattice of phengite at high temperature and high pressure were analyzed.
15	• High pressure induces O-H bond weakening and hydrogen disordering.
16 17	• Ammonium incorporated has potential effects on thermodynamic properties and stability of phengite.

20 Abstract

Water release in subduction zones is not only an important part of the deep Earth's water cycle, 21 but also plays an essential role in the physical and chemical properties of rocks constituting the 22 23 deep Earth. To understand water release processes, it is important to know properties of dehydration in hydrous phases of the downgoing slab. Although it is widely accepted that 24 phengite can be stable to greater depth in subduction environment, behavior of hydroxyl and 25 lattice of it at high temperature and high pressure are less investigated in contrast to other 26 27 hydrous phases. Here, using in situ IR and Raman spectroscopy, we characterize hydroxyl and lattice of ammonium-bearing and ammonium-free phengite at high temperature and high 28 pressure. No proton transferring and structural phase transition in phengite were observed over 29 the measured temperature and pressure range. Both pressure and temperature induce hydroxyl 30 band shifting to lower frequencies, and pressure has a greater impact. The band width of 31 hydroxyl increases with temperature and pressure. Hydroxyl bond weakening and hydrogen 32 disordering at high temperature and high pressure should be responsible for the spectra 33 variations. On the other hand, the lattice modes soften with increasing temperature whereas 34 stiffen under compression, and ammonium plays an important role in the Grüneisen parameters 35 of the lattice modes, especially the K-O mode. These features of hydroxyl and lattice at high 36 temperature and high pressure could benefit for further understanding dehydration, 37 thermodynamic properties and stability of phengite in subduction zones. 38

39 **1 Introduction**

Dehydration of hydrous minerals in subduction zones not only determines the transport 40 efficiency of water to the deep Earth, but also can trigger magmatism and earthquakes, cause 41 oxidization of the sub-arc mantle wedge, and lead to rheological heterogeneities and large 42 43 conductivity anomalies (e.g., Green et al., 2010; Grove et al., 2012; Schmidt & Poli, 2014; Debret et al., 2015; Manthilake et al., 2016; Okazaki & Hirth, 2016; Chen et al., 2017; Ferrand et 44 al., 2017; Behr et al., 2018). Dehydration of hydrous minerals has therefore been extensively 45 investigated over the years (Ono, 1998; Mibe et al., 1999; Hattori & Guillot, 2003; Wallace, 46 2005; Tokiwa & Nakashima, 2010; Van Keken et al., 2011; Hu et al., 2017; Hwang et al., 2017; 47 Liu et al., 2019). Several recent studies provide new insights into dehydration of hydrous 48 minerals in subduction zones. For example, Hu et al. (2017) reported dehydrogenation rather 49

than dehydration of goethite, revisiting the general concept of hydrogen circulation by hydration and dehydration. Our recent study (Liu et al., 2019) reported intimate links between other volatiles such as nitrogen and dehydration of phengite, highlighting the need to re-evaluate dehydration of hydrous minerals, in particular phengite. Most importantly, the role of superhydrated kaolinite formed under high pressures in the deep Earth's water cycle was recently evaluated by Hwang et al. (2017). They suggest that other hydrous minerals in the oceanic sediment and crust may undergo similar super-hydration during subduction.

Phengite is the most common high-pressure white mica observed in subducted metasediments, 57 58 being stable in subduction zones to depths of about 300 km (Schmidt, 1996). It is a good candidate for transporting both hydrogen and nitrogen into the deep Earth (Busigny et al., 2003; 59 Watenphul et al., 2009). Noteworthily, residing as ammonium (NH₄⁺) in the interlayer sites in 60 phengite, nitrogen is intimately linked to the kinetics and mechanism of dehydration in phengite 61 at high temperatures (Yang et al., 2017, Liu et al., 2019). Most interestingly, in contrast to other 62 hydrous minerals (Hwang et al., 2017), there is no hydration of phengite compressed in water, as 63 64 indicated by in situ Raman spectroscopic studies (Goryainov et al., 2017). To understand the stability and dehydration of phengite in subduction zones, the behavior of hydroxyl in phengite 65 and the lattice properties of phengite at high temperature and pressure should be examined in 66 detail. However, in contrast to other hydrous minerals in subduction zones, in situ high-T-P 67 vibrational spectroscopy has been rarely applied to phengite. In this study, we present new data 68 of the structural stability of hydroxyl and variation of lattice feature in phengite at high 69 70 temperature and pressure.

71 2 Materials and Methods

72 **2.1. Samples**

73 Single crystals of natural ammonium-bearing and ammonium-free phengite from a previous

study (Liu et al., 2019) were selected for this study. The chemical composition and unit-cell

- 75 parameters are listed in Table S1.
- 76 2.2. *In situ* high temperature IR and Raman spectroscopy

For *in situ* high temperature IR spectra measurements, an Instec HS1300 heating stage with CaF_2 windows was used to generate high temperatures. The samples were heated in the stage purged

with Ar. IR spectra in the frequency range 4000-1000 cm⁻¹ were collected at every temperature at 79 Zhejiang University, China, using a Nicolet iS50 FTIR spectrometer coupled with a Continuum 80 81 microscope. A KBr beam-splitter and a liquid nitrogen-cooled MCT-A detector were applied. A total of 128 scans were accumulated for each spectrum at a 4 cm⁻¹ resolution. The aperture size 82 was set to be 50×50 µm. For *in situ* Raman spectra measurements, a Linkam 600 heating stage 83 was used to generate high temperatures. Raman spectra were collected from a range of 50 cm⁻¹ to 84 1200 cm⁻¹ at every temperature employing a HPRIBA LABRAM-HR spectrometer at Zhejiang 85 University, China. Single-crystal silicon was used as a reference. The sample was excited by the 86 514.5 nm green line of a Spectra Physics Ar laser. A 50× objective was used to focus the incident 87 laser light on the sample and to collect the scattered light. The diameter of the focused laser light 88 spot was estimated to be 1 µm. 89

90 2.3. In situ high pressure IR and Raman spectroscopy

For in situ high pressure IR and Raman spectra measurements, hydrostatic high-pressure 91 92 conditions were generated in a symmetric piston-cylinder diamond anvil cell (DAC) with diamonds of 300 µm culet size. Prior to sample loading, pieces of stainless steel foil or Re gasket 93 94 were pre-indented in the cell down to a thickness of about 20-30 µm. A 100 to150 µm diameter 95 hole was then drilled into the gasket. A thin sample section was loaded into the sample chamber between two diamond anvils together with ruby for pressure calibration. The sample chamber 96 was then filled with KBr as a pressure medium. During the experiments, pressure variations 97 within the samples were monitored with the multiple ruby clips located near the sample. At the 98 99 peak pressures on compression, these variations did not exceed 10% of the total pressure.

Using a Bruker VERTEX 70v spectrometer coupled with a Bruker Hyperion 3000 microscope, 100 IR spectra from 4000-1000 cm⁻¹ were collected at every pressure at the Infrared spectroscopy 101 and microscopic imaging beamline (BL01B) of the National Synchrotron Radiation Laboratory 102 (University of Science and Technology of China). A KBr beam-splitter and a liquid nitrogen-103 cooled MCT-A detector were applied. The aperture size was set to 30×30 µm. A total of 128 104 scans were accumulated for each spectrum at a resolution of 4 cm⁻¹. Raman spectra were 105 collected from 50 to 1200 cm⁻¹ and from 3000 to 4000 cm⁻¹ at every pressure using a HPRIBA 106 LABRAM-HR spectrometer. Single-crystal silicon was used as a reference. The sample was 107 108 excited by the 514.5 nm green line of a Spectra Physics Ar laser. A 50× objective was used to

focus the incident laser light on the sample and to collect the scattered light. The diameter of thefocused laser light spot was estimated to be 1 µm.

- All the parameters of IR and Raman spectra at elevated temperatures and pressures are listed in
- 112 Tables S2-S5.

113 **3 Results and discussion**

114 **3.1. Hydroxyl in phengite at ambient conditions**

Phengite is a common high pressure white mica, with chemical compositions between muscovite 115 end-member and celadonite end-member. Muscovite is comprised of an octahedral sheet 116 sandwiched between two tetrahedral sheets (SiO₄), forming tetrahedral-octahedral-tetrahedral (T-117 118 O-T) layer structure. Although structure of muscovite has been extensively investigated, hydrogen sites in muscovite lattice are still not well known. Liang et al. (1998) reported two 119 independent but closely separated hydrogen sites within muscovite based on the neutron 120 spectroscopy at -261 °C, which is consistent with the IR spectrum of muscovite and electrostatic 121 calculations (Serratosa & Bradley, 1958; Rouxhet, 1970; Abbott et al., 1989; Liang & 122 Hawthorne, 1998). However, Gatta et al. (2011) argued for the presence of a single hydrogen site 123 in muscovite based on neutron diffraction at -261 °C. Hydrogen disordering in muscovite at high 124 pressure, indicated by IR spectra, also suggests multiple hydrogen sites (Williams et al., 2012). 125 Because of these discrepancies, only one band around 3620 cm⁻¹ in the IR spectra of hydroxyl in 126 muscovite is generally considered. Recently, Tuladhar et al. (2019) detected multiple distinct 127 peaks in the O-H stretching vibration region using vibrational sum frequency generation 128 spectroscopy. In this study, we also find the broad and asymmetric O-H stretching bands of the 129 phengite samples at ambient conditions (Fig. 1). Band deconvolution indicates two OH bands 130 around 3600 and 3620 cm⁻¹. Therefore, the broad and asymmetric OH bands should be assigned 131 to hydrogen sites with different environments. 132



135 Figure 1 IR spectra o

Figure 1. IR spectra of hydroxyl in the phengite at ambient conditions. The crystal structures of the two samples are
 illustrated.

137 **3.2. Hydroxyl in phengite at high temperatures**

Evolutions of hydroxyl in ammonium-bearing and ammonium-free phengite samples at elevated 138 temperatures to 800 °C are displayed in Figure 2. The frequency of hydroxyl appears insensitive 139 to temperature for both phengite types. The hydroxyl band gradually broadens with increasing 140 temperature, with a discontinuous broadening taking place at about 400 °C, indicating 141 disordering of the different hydrogen sites. Comparing the spectra at 20 °C before and after 142 heating, it is apparent that the integral absorbance decreases for the ammonium-bearing phengite 143 but does not change for the ammonium-free phengite. This is in accordance with our previous 144 study showing that ammonium promotes dehydration of phengite at high temperatures (Liu et al., 145 2019). Figure 3 compares temperature-induced frequency shifts of some hydrous phases in 146 subduction zones. Hydroxyl bands in these phases generally have negative frequency shifts with 147 increasing temperature. In contrast to nominally anhydrous minerals (Yang et al., 2019), there 148 149 appears no relationship between the frequency shift and room-temperature frequency of these hydrous phases. However, it is obvious that hydroxyl in phengite is most insensitive to 150 temperature, which is distinct to other hydrous minerals especially muscovite and pyrophyllite. 151 The frequency of a hydroxyl reflects strength of the O-H bond. Thus, negative frequency shifts 152 are caused by lengthening of the O-H bond during heating. The contrasting behavior of O-H 153 154 bond between phengite and other hydrous minerals at high temperatures is compatible with the better thermal stability of the former. 155



159 Figure 2. Temperature evolutions of hydroxyl in phengite: (a) and (b) in situ high temperature IR spectra of ammonium-

160 bearing phengite and ammonium-free phengite, respectively; (c) variation of the hydroxyl frequency with increasing

161 temperature; (d) variation of the width of hydroxyl band the full width at half-maximum, FWHM) with increasing temperature.

162

163



164

165 Figure 3. Temperature-induced frequency shifts of the hydroxyl band in hydrous minerals.

166 **3.3. Hydroxyl in phengite at high pressures**

167 Evolutions of hydroxyl in ammonium-bearing and ammonium-free phengite under compression are shown in Figure 4. We did not find pressure-induced hydrogen transferring similar to that 168 169 observed for brucite by Shinoda et al. (2002), but pressure can indeed induce dramatic variations of hydroxyl. The hydroxyl band broadens with increasing pressure. Interestingly, there is a 170 discontinuity at around 10 GPa in the evolution of band width with pressure, for both IR band 171 and Raman band. The band broadening indicates pressure-induced hydrogen disordering, which 172 173 is similar to the behavior of hydroxyl in muscovite (Williams et al., 2012). In addition, pressure also induces hydroxyl band shifting to lower frequencies, and the frequency shift is much greater 174

than temperature-induced shift. Hydroxyl in phengite continuously shifts to lower frequencies 175 under compression, without displaying a discontinuity in the pressure-induced frequency shift as 176 hydroxyl in lawsonite (Scott & Williams, 1999). Figure 5 includes pressure-induced frequency 177 shifts of hydroxyl in hydrous phases of subduction zones. Some bands shift to higher 178 frequencies, while others shift to lower frequencies. There seems to be no general relationship 179 between frequency shift and room-pressure frequency for the minerals. However, this 180 relationship indeed exists within the same mineral. For example, there is a positive relationship 181 between the frequency shift and room-pressure frequency for most of the minerals except for 182 lawsonite and phase D displaying a negative relationship. The pressure-induced negative 183 hydroxyl frequency shift is generally ascribed to shortening and strengthening of the hydrogen 184 bonding (an attractive interaction between a hydrogen atom from a hydroxyl and a neighboring 185 oxygen atom, O-H...O), thereby lengthening and weakening of the O-H bond with increasing 186 pressure (e.g., Pawley & Jones, 2011; Williams et al., 2012). Therefore, the negative frequency 187 shift of hydroxyl in phengite under compression indicates weakening of the hydroxyl bonds at 188 high pressures. The frequency shift range of hydroxyl in phengite is relatively small compared to 189 190 other minerals, especially lawsonite and brucite.



Figure 4. Evolutions of hydroxyl in phengite under compression: (a) and (b) *in situ* high pressure IR spectra of ammonium-bearing phengite and ammonium-free phengite, respectively; (c) and (d) *in situ* high pressure Raman spectra

ammonium-bearing phengite and ammonium-free phengite, respectively; (c) and (d) *in situ* high pressure Raman spectra of ammonium-bearing phengite and ammonium-free phengite, respectively; (e) variation of the hydroxyl frequency with

increasing pressure; (f) variation of the width of hydroxyl IR band with increasing pressure; (g) variation of the width of hydroxyl Raman band with increasing pressure.



198 Figure 5. Pressure-induced frequency shifts of hydroxyl bands in hydrous minerals.

3.4. Lattice of phengite at high temperature/high pressure

200 There are about five prominent bands around 95, 190, 260, 420 and 706 cm⁻¹ in the spectra of

201 phengite at ambient conditions. McKeown et al. (1999) suggested that the modes between 800

and 360 cm^{-1} had internal tetrahedral sheet motions mixed with K and octahedral Al 202 displacements, and the modes at frequencies less than 360 cm^{-1} had lattice motions. The 203 evolutions of lattice vibrations with increasing temperature to 500 °C are shown in Figure 6. The 204 spectra at high temperatures suggest no disordering of the lattice within this temperature range. 205 The frequencies of most of the lattice modes slightly shift to lower frequencies with increasing 206 temperature. Inversely, the 435 cm⁻¹ mode in the ammonium-bearing phengite slightly shifts to 207 higher frequencies. In addition, the frequency shift of the 95 cm⁻¹ mode is much smaller for the 208 ammonium-bearing phengite than the ammonium-free phengite. According to the band 209 assignment of previous studies (Holtz et al., 1993, McKeown et al., 1999, Mookherjee & 210 Redfern, 2002, Zhang et al., 2010), the band at 95 cm⁻¹ in the present study should be due to K-O 211 stretching vibration. The temperature-induced lower frequency shift of the lattice modes 212 indicates expansion of the microstructure with increasing temperature. When the isobaric 213 Grüneisen parameter of the five modes of the ammonium-bearing phengite, the ammonium-free 214 phengite and muscovite are compared, the isobaric mode Grüneisen parameters of muscovite are 215 obviously different from those of phengite. Furthermore, the distinct behavior of some modes, 216 especially the 95 cm⁻¹ mode, between the ammonium-bearing phengite and the ammonium-free 217 phengite may suggest some effects of ammonium on the thermodynamic properties and stability 218 219 of the phengite lattice.

220



Figure 6. Temperature evolutions of phengite lattice: (a) and (b) *in situ* high temperature Raman spectra of ammoniumbearing phengite and ammonium-free phengite, respectively; (c)-(g) variations of Raman shifts with increasing temperature; (h) isobaric Grüneisen parameters of the Raman modes. The isobaric Grüneisen parameters of phengite were calculated using the equation of Fujimori et al. (2002), and the thermal expansion coefficient was taken from Gemmi et al. (2008).

Based on high pressure Raman and XRD spectroscopic studies of muscovite, lattice disordering 227 of muscovite occurs at pressures around 15-18 GPa (Faust & Knittle, 1994; Williams et al., 228 2012). However, the Raman spectra of phengite suggest no lattice disordering within the 229 pressure range of this study (Fig. 7), which is in agreement with the observations of phengite 230 made by Goryainov et al. (2017). In contrast to temperature, pressure causes modes significantly 231 shifting to higher frequencies for both the ammonium-bearing and ammonium-free phengite 232 samples. There is little difference between the pressure-induced mode shifts of the two phengite 233 species. The pressure-induced stiffening of the Raman modes indicates strengthening of the K-O, 234 M-O and T-O bonds. The shortening of these bonds is compatible with the compressed structure 235 of phengite (Gatta et al., 2009, 2010). The isothermal mode Grüneisen parameters of muscovite 236 are similar to those of phengite, except for the 112 cm⁻¹ mode assigned to K-translation 237 238 (Williams et al., 2012). Comparison between the ammonium-bearing phengite and the ammonium-free phengite indicates that ammonium has little effect on the isothermal mode 239 Grüneisen parameters. Nevertheless, ammonium effects on the compression of phengite still 240 need to be investigated. 241

242



Figure 7. Evolutions of phengite lattice under compression: (a) and (b) *in situ* high pressure Raman spectra of ammonium-bearing phengite and ammonium-free phengite, respectively; (c)-(f) variations of Raman shifts with increasing pressure; (h) isothermal Grüneisen parameter of the Raman modes. The isobaric Grüneisen parameters of

247 phengite were calculated using the equation of Fujimori et al. (2002), and the isothermal bulk modulus was taken from

248 Curetti et al. (2006).

249 **4. Implications**

250 4.1. Dehydration before phengite breakdown

Water release in subduction zones provides the main source for fluids triggering flux melting in 251 252 the mantle wedge, formation of volcanoes and earthquakes (e.g., Van KeKen et al., 2011). In studying dehydration of hydrous minerals in subducting slabs, much more attention has been 253 paid to mineral breakdown than water diffusion in the mineral lattice. However, it was 254 recognized by Gaines & Vedder (1964) that the rate of muscovite breakdown may be limited by 255 hydroxyl diffusion from its lattice. Indeed, many subsequent experimental studies focused on 256 dehydroxylation of muscovite without breakdown during heating at room pressure (e.g., Gaines 257 & Vedder, 1964; Mazzuccato et al., 1999; Tokiwai & Nakashima, 2010; Zhang et al., 2010; 258 Heller-Kallai & Lapides, 2015; Liu et al., 2019). Whether dehydroxylation without mineral 259 breakdown can happen at high pressures need to be further clarified. Our study concerns the 260 behavior of hydroxyl in phengite at high pressure and high temperature, revealing hydroxyl 261 bonds weakening and hydrogen disordering. For brucite, any modulation of the O...O distance 262 as a result of thermal motions will effectively lower the height of the potential barrier between 263 proton sites of hydroxyl (Martens & Freund, 1976). On the other hand, pressure-induced 264 hydrogen disordering in hydroxide has been extensively reported. It can trigger destabilization of 265 the oxygen substructure, and enhancement of electrical conductivity (e.g., Duffy et al., 1995; 266 Guo & Yoshino, 2014). According to Mazzuccato et al. (1999), the process of dehydration in 267 muscovite encompasses the following three steps: condensation of two adjacent hydroxyls in the 268 octahedral layer to form a water molecule, diffusion of the water molecule throughout the 269 tetrahedral layer, and diffusion of the water molecule in the interlayer region to be releases. 270 Since the formation of water from structural hydroxyl in the lattice is the primary step, 271 dehydroxylation in the lattice is vital for the whole dehydration process. We propose that 272 273 pressure-induced shortening of the O...O distance (weakening of hydroxyl bond) and hydrogen disordering in phengite should favor condensation of two adjacent hydroxyls in the octahedral 274 275 layer to form water molecules. These results may shed new light on dehydration in phengite. Water may be continuously released from the lattice before the phengite breakdown with 276 277 increasing temperature and pressure. To further compare the properties of hydroxyl of the

hydrous phases under compression in subduction zones, we display the available isothermal 278 Grüneisen parameters of hydroxyl in several hydrous minerals in Figure 8. It is clear that the 279 isothermal Grüneisen parameters for hydroxyl in most phases in subduction zones are positive, 280 except for phengite, phase E and phase D showing negative isothermal Grüneisen parameters. 281 Although water content in phengite is not very high relative to other hydrous phases in 282 subduction zones, phengite can be stable at a wide range of pressure in the subduction 283 environment. Therefore, dehydroxylation from the lattice before its breakdown can not be 284 285 neglected in the evaluation of water release in subduction zones.





287

288 Figure 8. Comparison of isothermal Grüneisen parameters of hydroxyl between different minerals in subduction zones: 289 (a) isothermal Grüneisen parameters of hydroxyl in hydrous minerals. The parameters of muscovite, phase E, phase D 290 and lawsonite were taken from the references listed in the figure. The parameters of the other minerals were calculated 291 using the equation of Fujimori et al. (2002). The frequency shifts were adopted for phlogopite from Williams et al. (2012), 292 for chlorite from Kleppe et al. (2003), and for clinohumite and chondrodite from Liu et al. (2003). The isothermal bulk 293 modulus values were taken for phengite from Curetti et al. (2006), for phlogopite from Comodi et al. (2004), for chlorite 294 from Welch and Crichton (2002), and for clinohumite and chondrodite from Ross and Crichton (2001). (b) average 295 isothermal Grüneisen parameters of hydroxyl in minerals that are stable to different depths, with the maximum stability 296 depths taken from Litasov and Ohtani (2007).

4.2. Potential effects of ammonium on thermodynamic properties and stability of phengite

Mineral stability under equilibrium conditions is governed by the lowest free energy. Lattice vibrations are related to thermodynamic properties of minerals, thus, mineral stabilities can be studied from lattice vibrations (Kieffer, 1979; Ross, 1992). Our study found that temperature and pressure have opposite impacts on the properties of the phengite lattice. Temperature induces softening of the framework, while pressure makes the framework stiffer. Based on the mode Grüneisen parameters of the ammonium-bearing and ammonium-free phengite samples (Figs. 6 and 7), the absolute value of the average isobaric Grüneisen parameter is much smaller for ammonium-bearing phengite (0.67) than the ammonium-free phengite (1.30), while the absolute value of the average isothermal Grüneisen parameter is not much different between the two samples. It should be noted that the mode Grüneisen parameters in this study were calculated on the basis of the thermal expansion coefficient and isothermal bulk modulus of ammonium-free phengite (Curetti et al., 2006, Gemmi et al., 2008). Even so, ammonium impacts on the mode Grüneisen parameters can still be observed. Therefore, the effects of ammonium on the thermodynamic properties and stability of phengite deserve careful consideration.

312

313 ACKNOWLEDGEMENTS

This work is supported by the National Natural Science Foundation of China (41972038)

and the Zhejiang Province Natural Science Foundation of China (LY18D020001). All data
presented in this manuscript are available at Figshare
http://doi.org/10.6084/m9.figshare.11339546.

- 318 **References**
- Abbott, J. R. N., Post, J. E., & Burnham, C. W. (1989). Treatment of the hydroxyl in structure energy calculations. American Mineralogist, 74(1-2), 141-150.
- Auzende, A.L., Daniel, I., Reynard, B., Lemaire, C., and Guyot, F. (2004) High-pressure
 behaviour of serpentine minerals: A Raman spectroscopic study. Physics and Chemistry of
 Minerals, 31, 269–277. https://doi.org/10.1007/s00269-004-0384-0.
- Behr, W. M., Kotowski, A. J., & Ashley, K. T. (2018). Dehydration-induced rheological
 heterogeneity and the deep tremor source in warm subduction zones. Geology, 46(5), 475478. https://doi.org/10.1130/G40105.1.
- Busigny, V., Cartigny, P., Philippot, P., & Javoy, M. (2003). Ammonium quantification in
 muscovite by infrared spectroscopy. Chemical Geology, 198(1), 21-31.
 https://doi.org/10.1016/S0009-2541(02)00420-5.
- Chen, S. B., Guo, X. Z., Yoshino, T., Jin, Z. M., & Ping, L. (2017). Dehydration of phengite
 inferred by electrical conductivity measurements: Implication for the high conductivity
 anomalies relevant to the subduction zones. Geology, 46(1).
 https://doi.org/10.1130/G39716.1.
- Comodi, P., Fumagalli, P., Montagnoli, M., and Zanazzi, P.F. (2004) A single-crystal study on
 the pressure behavior of phlogopite and petrological implications. American Mineralogist,
 89, 647–653. https://doi.org/10.2138/am-2004-0420.
- Comodi, P., Cera, F., Dubrovinsky, L., & Nazzareni, S. (2006) The high-pressure behaviour of
 the 10 Å phase: A spectroscopic and diffractometric study up to 42 GPa. Earth and
 Planetary Science Letters, 246, 444-457. https://doi.org/10.1016/j.epsl.2006.03.046.

Curetti N, Levy D, Pavese A, Ivaldi G (2006) Elastic properties and stability of coexisting 3T
 and 2M1 phengite polytypes. Phys Chem Minerals 32:670-678.
 https://doi.org/10.1007/s00269-005-0044-z.

- Debret, B., Bolfan-Casanova, N., Padrón-Navarta, J. A., Martin-Hernandez, F., Andreani, M.,
 Garrido, C. J., Sánchez-Vizcaíno, V. L., Gómez-Pugnaire, M. T., Muñoz, M., & Trcera, N.
 (2015). Redox state of iron during high-pressure serpentinite dehydration. Contributions to
 Mineralogy and Petrology, 169(4), 36. https://doi.org/10.1007/s00410-015-1130-y.
- Duffy, T.S., Meade, C., Fei, Y.W., Mao, H.K., and Hemley, R.J. (1995) High-pressure phase
 transition in brucite, Mg(OH)2. American Mineralogist, 80, 222–230.
 https://doi.org/10.2138/am-1995-3-404.
- Faust, J., & Knittle, E. (1994). The equation of state, amorphization, and high-pressure phase
 diagram of muscovite. Journal of Geophysical Research, 99(B10), 19785.
 https://doi.org/10.1029/94jb01185.
- Ferrand, T. P., Hilairet, N., Incel, S., Deldicque, D., Labrousse, L., Gasc, J., Renner, J., Wang, Y.,
 Green Ii, H. W., & Schubnel, A. (2017). Dehydration-driven stress transfer triggers
 intermediate-depth earthquakes. Nature Communications, 8, 15247.
 https://doi.org/10.1038/ncomms15247.
- Fujimori, H., Komatsu, H., Ioku, K., and Goto, S. (2002) Anharmonic lattice mode of Ca2SiO4:
 Ultraviolet laser Raman spectroscopy at high temperatures. Physical Review B, 66, 064306.
 https://doi.org/10.1103/PhysRevB.66.064306.
- Gaines, G. L., & Vedder, W. (1964). Dehydroxylation of Muscovite. Nature, 201(4918), 495-495.
 https://doi.org/10.1038/201495a0.
- Gatta, G. D., Mcintyre, G. J., Sassi, R., Rotiroti, N., & Pavese, A. (2011). Hydrogen-bond and
 cation partitioning in muscovite: A single-crystal neutron-diffraction study at 295 and 20 K.
 American Mineralogist, 96(1), 34-41. https://doi.org/10.2138/am.2011.3595.
- Gatta, G. D., Rotiroti, N., Pavese, A., Lotti, P., & Curetti, N. (2009). Structural evolution of a 3T
 phengite mica up to 10 GPa: an in-situ single-crystal X-ray diffraction study. Zeitschrift für
 Kristallographie, 224(5-6), 302-310. https://doi.org/10.1524/zkri.2009.1131.
- Gatta, G. D., Rotiroti, N., Lotti, P., Pavese, A., & Curetti, N. (2010). Structural evolution of a 2
 M1 phengite mica up to 11 GPa: an in situ single-crystal X-ray diffraction study. Physics
 and Chemistry of Minerals, 37(8), 581-591. https://doi.org/10.1007/s00269-010-0359-2.
- Gemmi, M., Merlini, M., Pavese, A., and Curetti, N. (2008) Thermal expansion and
 dehydroxylation of phengite micas. Physics and Chemistry of Minerals, 35: 367-379.
 https://doi.org/10.1007/s00269-008-0230-x.
- Goryainov, S. V., Krylov, A. S., Polyansky, O. P., & Vtyurin, A. N. (2017). In-situ Raman study
 of phengite compressed in water medium under simultaneously high P-T parameters.
 Journal of Raman Spectroscopy, 48(11), 1431-1437. https://doi.org/10.1002/jrs.5112.
- Green, D. H., Hibberson, W. O., István, K., & Anja, R. (2010). Water and its influence on the
 lithosphere-asthenosphere boundary. Nature, 467(7314), 448-451.
 https://doi.org/10.1038/nature09948.

- Grove, T. L., Till, C. B., & Krawczynski, M. J. (2012). The Role of H2O in Subduction Zone
 Magmatism. Annual Review of Earth and Planetary Sciences, 40(1), 413-439.
 https://doi.org/10.1146/annurev-earth-042711-105310.
- Guo, X., and Yoshino, T., 2014, Pressure-induced enhancement of proton conduction in brucite:
 Geophysical Research Letters, 41, 813-819. https://doi.org/10.1002/2013GL058627.
- Hattori, K. H., & Guillot, S. (2003). Volcanic fronts form as a consequence of serpentinite
 dehydration in the forearc mantle wedge. Geology, 31(6), 525-528.
 https://doi.org/10.1130/0091-7613(2003)031<0525:VFFAAC>2.0.CO;2.
- Heller-Kallai, L., & Lapides, I. (2015). Dehydroxylation of muscovite: study of quenched
 samples. Physics and Chemistry of Minerals, 42(10), 835-845.
 https://doi.org/10.1007/s00269-015-0767-4.
- Holtz, M., Solin, S.A., and Pinnavaia, T.J. (1993) Effect of pressure on the Raman vibrational
 modes of layered aluminosilicate compounds. Physical Review B, 13313-13317.
 https://doi.org/10.1103/PhysRevB.48.13312.
- Hu, Q. Y., Kim, D. Y., Liu, J., Meng, Y., Yang, L. X., Zhang, D. Z., Mao, W. L., & Mao, H.-K.
 (2017). Dehydrogenation of goethite in Earth's deep lower mantle. Proceedings of the
 National Academy of Sciences, 114(7), 1498-1501.
 https://doi.org/10.1073/pnas.1620644114.
- Hwang, H. J., Seoung, D., Lee, Y., Liu, Z., Liermann, H. P., Cynn, H., Vogt, T., Kao, C. C., &
 Mao, H. K. (2017). A role for subducted super-hydrated kaolinite in Earth's deep water
 cycle. Nature Geoscience, 10(12), 947-953. https://doi.org/10.1038/s41561-017-0008-1.
- Kieffer, S. W. (1979). Thermodynamics and lattice vibrations of minerals: 3. Lattice dynamics
 and an approximation for minerals with application to simple substances and framework
 silicates. Reviews of Geophysics, 17(1), 35-59. https://doi.org/10.1029/RG017i001p00035.
- Kleppe, A.K., Jephcoat, A.P., & Welch, M.D. (2003) The effect of pressure upon hydrogen
 bonding in chlorite: A Raman spectroscopic study of clinochlore to 26.5 GPa. American
 Mineralogist, 88, 567-573. https://doi.org/10.2138/am-2003-0410.
- Liang, J.J., & Hawthorne, F.C. (1998) Calculated H-atom positions in micas and clay minerals.
 Canadian Mineralogist, 36, 1577-1585.
- Liang, J. J., & Hawthorne, F. C. (1998). Calculated H-atom positions in micas and clay minerals.
 The Canadian Mineralogist, 36(6), 1577-1585.
- Litasov, K.D., & Ohtani, E. (2007) Effect of water on the phase relations in Earth's mantle and
 deep water cycle, in Ohtani, E., ed., Advances in High-Pressure Mineralogy: Geological
 Society of America Special Paper 421, p. 115-156, doi: 10.1130/2007.2421(08).
- Liu, W. D., Yang, Y., Busigny, V., & Xia, Q.-K. (2019). Intimate link between ammonium loss
 of phengite and the deep Earth's water cycle. Earth and Planetary Science Letters, 513(1),
 95-102. https://doi.org/10.1016/j.epsl.2019.02.022.
- Liu, Z., Lager, G.A., Hemley, R.J., & Ross, N.L. (2003) Synchrotron infrared spectroscopy of
 OH-chondrodite and OH-clinohumite at high pressure. American Mineralogist, 88, 14121415. https://doi.org/10.2138/am-2003-1003.

- Manthilake, G., Bolfan-Casanova, N., Novella, D., Mookherjee, M., & Andrault, D. (2016).
 Dehydration of chlorite explains anomalously high electrical conductivity in the mantle
 wedges. Science Advances, 2(5), e1501631. https://doi.org/10.1126/sciadv.1501631.
- Martens, R., & Freund, F. (1976) The potential energy curve of the proton and dissociation
 energy of the OH- ion in Mg(OH)2. Phys Status Solidi A, 37(1): 97-104.
 https://doi.org/10.1002/pssa.2210370112.
- Mazzucato, E., Artioli, G., & Gualtieri, A. (1999). High temperature dehydroxylation of
 muscovite-2M1: a kinetic study by in situ XRPD. Physics and Chemistry of Minerals, 26(5),
 375-381. https://doi.org/10.1007/s002690050197.
- McKeown, D.A., Bell, M.L., and Etz, E. (1999) Vibrational analysis of the dioctahedral mica:
 2M1 muscovite. American Mineralogist, 84: 1041-1048. https://doi.org/10.2138/am-19997-806.
- Mibe, K., Fujii, T., & Yasuda, A. (1999). Control of the location of the volcanic from in island
 arcs by aqueous fluid connectivity in the mantle wedge. Nature, 401(6750), 259-262.
 https://doi.org/10.1038/45762.
- Mookherjee, M. and Redfern, S.A.T. (2002) A high-temperature Fourier transform infrared study
 of the interlayer and Si-O-stretching region in phengite-2M1. Clay Minerals, 37: 323-336.
 https://doi.org/10.1180/0009855023720036.
- Okazaki, K., & Hirth, G. (2016). Dehydration of lawsonite could directly trigger earthquakes in
 subducting oceanic crust. Nature, 530(7588), 81-84. https://doi.org/10.1038/nature16501.
- 440 Ono, S. (1998). Stability limits of hydrous minerals in sediment and mid-ocean ridge basalt
 441 compositions: Implications for water transport in subduction zones. Journal of Geophysical
 442 Research, 103(B8), 18253-18267. https://doi.org/10.1029/98JB01351.
- Pawley, A. R., & Jones, R. L. (2011). Hydroxyl stretching in phyllosilicates at high pressures
 and temperatures: an infrared spectroscopic study. Physics and Chemistry of Minerals,
 38(10), 753-765. https://doi.org/10.1007/s00269-011-0448-x.
- Ross, N.L. (1992) Lattice vibration and mineral stability. In: Price G.D., Ross N.L. (eds) The
 Stability of Minerals. The Mineralogical Society Series, vol 3. Springer, Dordrecht.
- Ross, N.L. & Crichton, W.A. (2001) Compression of synthetic hydroxylclinohumite
 [Mg9Si4O16(OH)2] and hydroxylchondrodite [Mg5Si2O8(OH)2]. American Mineralogist,
 86, 990–996. https://doi.org/10.2138/am-2001-8-905.
- Rouxhet, P. G. (1970). Hydroxyl stretching bands in micas: a quantitative interpretation. Clay
 Minerals, 8(4), 375-388. https://doi.org/10.1180/claymin.1970.008.4.02.
- Scott, H.P., & Williams, Q. (1999) An infrared spectroscopic study of lawsonite to 20 GPa.
 Physics and Chemistry of Minerals, 26, 437-445.
- Scott, H.P., Liu, Z., Hemley, R.J., and Williams, Q. (2007) High-pressure infrared spectra of talc
 and lawsonite. American Mineralogist, 92, 1814–1820.
 https://doi.org/10.2138/am.2007.2430.

- Schmidt, M. W. (1996). Experimental constraints on recycling of potassium from subducted
 oceanic crust. Science, 272(5270), 1927-1930.
 https://doi.org/10.1126/science.272.5270.1927.
- Schmidt, M. W., & Poli, S. (2014), Devolatilization during subduction, in Treatise on
 geochemistry (second edition), edited by H. D. Holland and K. K. Turekian, pp. 669-701,
 Elsevier, Amsterdam.
- Serratosa, J. M., & Bradley, W. F. (1958). Determination of the Orientation of OH Bond Axes in
 Layer Silicates by Infrared Absorption. Journal of Physical Chemistry, 62(10), 1164-1167.
 https://doi.org/10.1021/j150568a003
- Shieh, S.R., Duffy, T.S., Liu, Z., and Ohtani, E. (2009) High-pressure infrared spectroscopy of
 the dense hydrous magnesium silicates phase D and phase E. Physics of the Earth and
 Planetary Interiors, 175, 106–114. https://doi.org/10.1016/j.pepi.2009.02.002.
- 470 Shinoda, K., Yamakata, M., Nanba, T., Kimura, H., Moriwaki, T., Kondo, Y., Kawamoto, T., Niimi, N., Miyoshi, N., and Aikawa, N. (2002) High-pressure phase transition and 471 472 behaviour of protons in brucite Mg(OH)2: a high-pressure-temperature study using IR radiation. and Chemistry synchrotron Physics of Minerals, 29. 396-402. 473 https://doi.org/10.1007/s00269-002-0243-9. 474
- Thompson, E.C., Campbell, A.J., & Liu, Z. (2016) In-situ infrared spectroscopic studies of
 hydroxyl in amphiboles at high pressure. American Mineralogist, 101, 706-712.
 https://doi.org/10.2138/am-2016-5465.
- Tokiwai, K., & Nakashima, S. (2010). Dehydration kinetics of muscovite by in situ infrared
 microspectroscopy. Physics and Chemistry of Minerals, 37(2), 91-101.
 https://doi.org/10.1007/s00269-009-0313-3.
- 481 Trittschack, R, Grobéty, B., & Koch-Müller, M. (2012) In situ high-temperature Raman and
 482 FTIR spectroscopy of the phase transformation of lizardite. American Mineralogist, 97,
 483 1965-1976. https://doi.org/10.2138/am.2012.4162.
- Tuladhar, A., et al. (2019). Direct Observation of the Orientational Anisotropy of Buried
 Hydroxyl Groups inside Muscovite Mica. Journal of the American Chemical Society,
 141(5), 2135-2142. https://doi.org/10.1021/jacs.8b12483.
- Van Keken, P. E., Hacker, B. R., Syracuse, E. M., & Abers, G. A. (2011). Subduction factory: 4.
 Depth-dependent flux of H2O from subducting slabs worldwide. Journal of Geophysical Research, 116, B01401. https://doi.org/10.1029/2010JB007922.
- Welch, M.D. and Crichton, W.A. (2002) Compressibility of clinochlore to 8 GPa at 298 K and a
 comparison with micas. European Journal of Mineralogy, 14, 561-565.
 https://doi.org/10.1127/0935-1221/2002/0014-0561.
- Wallace, P. J. (2005). Volatiles in subduction zone magmas: concentrations and fluxes based on
 melt inclusion and volcanic gas data. Journal of Volcanology and Geothermal Research,
 140(1-3), 217-240. https://doi.org/10.1016/j.jvolgeores.2004.07.023.
- Watenphul, A., Wunder, B., & Heinrich, W. (2009). High-pressure ammonium-bearing silicates:
 implications for nitrogen and hydrogen storage in the Earth's mantle. American Mineralogist, 94(2-3), 283-292. https://doi.org/10.2138/am.2009.2995.

- Williams, Q., Knittle, E., Scott, H. P., & Liu, Z. X. (2012). The high-pressure behavior of micas:
 Vibrational spectra of muscovite, biotite, and phlogopite to 30 GPa. American Mineralogist,
 97(1), 241-252. https://doi.org/10.2138/am.2012.3824.
- Yang, Y., Busigny, V., Wang, Z. P., & Xia, Q.-K. (2017). The fate of ammonium in phengite at
 high temperature. American Mineralogist, 102(11), 2244-2253. https://doi.org/10.2138/am2017-6094.
- Yang, Y., Ingrin, J., Xia, Q.K., & Liu, W.D. (2019) Nature of hydrogen defects in clinopyroxenes from room temperature up to 1000 oC: Implication for the preservation of hydrogen in the upper mantle and impact on electrical conductivity. American Mineralogist, 104, 79-93. https://doi.org/10.2138/am-2019-6661.
- Zhang M, Hui Q, Lou XJ, Redfern SAT, Salje EKH, Tarantino SC (2006) Dehydroxylation,
 proton migration, and structural changes in heated talc: an infrared spectroscopic study.
 American Mineralogist 91:816–825. https://doi.org/10.2138/am.2006.1945.
- Zhang M, Salje EKH, Carpenter MA, Wang JY, Groat LA, Lager GA, Wang L, Beran A,
 Bismayer U (2007) Temperature dependence of IR absorption of hydrous/hydroxyl species
 in minerals and synthetic materials. American Mineralogist 92:1502-1517.
 https://doi.org/10.2138/am.2007.2586.
- Zhang, M., Redfern, S. A. T., Salje, E. K. H., Carpenter, M. A., & Hayward, C. L. (2010).
 Thermal behavior of vibrational phonons and hydroxyls of muscovite in dehydroxylation:
 In situ high-temperature infrared spectroscopic investigations. American Mineralogist,
 95(10), 1444-1457. https://doi.org/10.2138/am.2010.3472
- 520
- 521