

Time Dependent 2.5D Model of the Chromosphere

Qusai Al Shidi¹, Ofer Cohen², Paul Song², and Jiannan Tu²

¹University of Michigan

²University of Massachusetts–Lowell

November 24, 2022

Abstract

The Sun's chromosphere is a highly dynamic, partially ionized region where spicules (hot jets of plasma) form. Here we present a two-fluid MHD model to study the chromosphere, which includes ion–neutral interaction and frictional heating. Our simulation recovers a magnetic-canopy shape that forms quickly, but it is also quickly disrupted by the formation of a jet. Our simulation produces a shock self-consistently, where the jet is driven by the frictional heating, which is much greater than the ohmic heating. Thus, our simulation demonstrates that the jet could be driven purely by thermal effects due to ion–neutral collisions and not by magnetic reconnection. We plan to improve the model to include photo-chemical effects and radiation.




1. The Chromosphere

The sun's atmosphere is called the chromosphere. Compared to the sun's corona we don't have many observations of the chromosphere. We do know that, strangely, the chromosphere is much colder than the corona. The plasma in the chromosphere is therefore partially ionized—it has about a thousand times more neutral atoms than ions or electrons

2. The Problem


The chromosphere is a difficult region to simulate. The plasma behaves very differently depending on the height of interest. The plasma beta changes several orders of magnitude and the plasma changes from being dominated by magnetic forces near the top and dominated by thermal forces near the bottom.



Analytical solution of the steady-state chromosphere. (Song et al 2014)

4. Results


Preliminary results show a jet forming in the lower chromosphere between the network and internetwork regions of the sun. This is caused by frictional heating, the heating due to collisions between ions and neutrals. The strong magnetic gradient between the network and inter-network regions causes a speed difference between the two fluids and consequently causing friction. This then causes a shock to form and the shock steepens making large thermal pressure within the jet driving the jet upward and then out of the chromosphere. It was thought that chromospheric jets were only caused by reconnection events and in this it's a thermal effect due to collisions.



3. The Simulation

Collisional Multi-Fluid Ion (CoMFi) (github: qalshidi/comfi) is written from scratch and uses a semi-implicit TVD-MUSCL scheme. It is also GPU-accelerated taking advantage of the frontiers of scientific computing. The model is a two-fluid (plasma, neutrals) collisional MHD model.

$$\begin{aligned} \frac{\partial n_{i,n}}{\partial t} + \nabla \cdot n_{i,n} \mathbf{V} &= 0 \\ \frac{\partial \rho_i \mathbf{V}}{\partial t} + v_{in} \rho_i (\mathbf{V} - \mathbf{U}) &= -\nabla \cdot (\rho_i \mathbf{V} \mathbf{V} + n_i k T_i + \frac{B^2}{2\mu_0} - \frac{\mathbf{B} \mathbf{B}}{\mu_0}) - \rho_i g + \frac{m_e}{e} (v_{en} - v_{in}) \mathbf{J} \\ \frac{\partial \rho_n \mathbf{U}}{\partial t} - v_{in} \rho_i (\mathbf{V} - \mathbf{U}) &= -\nabla \cdot (\rho_n \mathbf{U} \mathbf{U} + n_n k T_n) - \rho_n g - \frac{m_e}{e} (v_{en} - v_{in}) \mathbf{J} \\ \frac{\partial k T_i}{\partial t} - \frac{2}{3} \frac{m_i}{m_i + m_n} v_{in} [m_n |\mathbf{V} - \mathbf{U}|^2 + 3(T_n - T_i)] &= -\nabla \cdot (k T_i \mathbf{V}) + \frac{1}{3} k T_i \nabla \cdot \mathbf{V} - \frac{2}{3 n_i} \nabla \cdot (\mathbf{q}_i + \mathbf{B} \times \eta \mathbf{J}) \\ \frac{\partial k T_n}{\partial t} - \frac{2}{3} \frac{m_n}{m_i + m_n} v_{in} [m_i |\mathbf{V} - \mathbf{U}|^2 + 3(T_i - T_n)] &= -\nabla \cdot (k T_n \mathbf{U}) + \frac{1}{3} k T_n \nabla \cdot \mathbf{U} - \frac{2}{3 n_n} \nabla \cdot (\mathbf{q}_n) \\ \frac{\partial \mathbf{B}}{\partial t} &= -\nabla \cdot (\mathbf{V} \mathbf{B} - \mathbf{B} \mathbf{V}) - \nabla \times (\eta \mathbf{J}) \end{aligned}$$



Top: Vertical slice of shock front of jet. You can see that the frictional heating (red dashed line) is enhanced behind the shockfront.

Left: Time series mosaic of jet shooting from the near the photosphere. The left mosaic is of ion density and the mosaic on the right is that of frictional heating.