Asymmetric internal tide generation in the presence of a steady flow

Yvan Dossmann¹, Callum J. Shakespeare², Kial Douglas Stewart³, and Andrew McC. $\rm Hogg^3$

¹LEMTA - Université de Lorraine ²Autralian National University ³Australian National University

November 21, 2022

Abstract

The generation of topographic internal waves (IWs) by the sum of an oscillatory and a steady flow is investigated experimentally and with a linear model. The two forcing flows represent the combination of a tidal constituent and a weaker quasi-steady flow interacting with an abyssal hill. The combined forcings cause a coupling between internal tides and lee waves that impacts their dynamics of internal waves as well as the energy carried away. An asymmetry is observed in the structure of upstream and downstream internal wave beams due to a Doppler shift effect. This asymmetry is enhanced for the narrowest ridge on which a super-buoyancy (ω >N) downstream beam and an evanescent upstream beam are measured.

Energy fluxes are measured and compared with the linear model, that has been extended to account for the coupling mechanism. The structure and amplitude of energy fluxes match well in most regimes, showing the relevance of the linear prediction for IW wave energy budgets, while the energy flux toward IW beams is limited by the generation of periodic vortices in a particular experiment. The upstream-bias energy flux - and consequently net horizontal momentum - described in Shakespeare [2020] is measured in the experiments.

The coupling mechanism plays an important role in the pathway to IW induced mixing, that has previously been quantified independently for lee waves and internal tides. Hence, future parameterizations of IW processes ought to include the coupling mechanism to quantify its impact on the global distribution of mixing.

Asymmetric internal tide generation in the presence of a steady flow

Yvan Dossmann^{1,2}, Callum Shakespeare², Kial Stewart², Andrew McC.

 $\mathrm{Hogg}^{2,3}$

Corresponding author: Yvan Dossmann, LEMTA, Université de Lorraine, CNRS, France. (yvan.dossmann@univ-lorraine.fr)

¹LEMTA, UMR 7563, Université de

Lorraine, CNRS, F-54500

Vandoeuvre-les-Nancy, France

²Research School of Earth Sciences, The

Australian National University, Canberra

2601, Australia

³ARC Centre of Excellence for Climate

Extremes

Key Points.

- The dynamics of topographic internal waves generated by the sum of a steady and oscillatory flow is studied for varying ridge shape.
- A topographically controlled asymmetry is measured in the energy fluxes carried by the upstream and downstream IW beams.
- The asymmetry and the generation of nonlinear structures impact the distribution of IW induced mixing in the ocean interior.
- ³ Abstract.

⁴ The generation of topographic internal waves (IWs) by the sum of an os-

 $_{\scriptscriptstyle 5}$ cillatory and a steady flow is investigated experimentally and with a linear

6 model. The two forcing flows represent the combination of a tidal constituent

 $_{\scriptscriptstyle 7}~$ and a weaker quasi-steady flow interacting with an abyssal hill. The com-

⁸ bined forcings cause a coupling between internal tides and lee waves that im-

⁹ pacts their dynamics of internal waves as well as the energy carried away.

¹⁰ An asymmetry is observed in the structure of upstream and downstream in-

¹¹ ternal wave beams due to a Doppler shift effect. This asymmetry is enhanced

¹² for the narrowest ridge on which a super-buoyancy ($\omega > N$) downstream

¹³ beam and an evanescent upstream beam are measured.

Energy fluxes are measured and compared with the linear model, that has been extended to account for the coupling mechanism. The structure and amplitude of energy fluxes match well in most regimes, showing the relevance of the linear prediction for IW wave energy budgets, while the energy flux toward IW beams is limited by the generation of periodic vortices in a particular experiment. The upstream-bias energy flux - and consequently net

²⁰ horizontal momentum - described in *Shakespeare* [2020] is measured in the
²¹ experiments.

The coupling mechanism plays an important role in the pathway to IW induced mixing, that has previously been quantified independently for lee waves and internal tides. Hence, future parameterizations of IW processes ought to include the coupling mechanism to quantify its impact on the global distribution of mixing.

Plain language summary

When tides and currents interact with abyssal topographies, such as ridges and hills, they generate internal waves that propagate in the ocean interior. The energy transported by these waves sustains the largest scale oceanic motions. To improve our understanding of how and where energy is transferred to oceanic currents, an important step is to describe the fate of internal waves, from their generation to their breaking.

Previous studies have independently described the dynamics of internal waves generated by tides, or by a steady current. Here we combine the two types of currents - a situation that is met at many oceanic sites - using laboratory experiments and a linear model. The combined currents cause an asymmetry in the internal wave structure. Internal waves are more energetic on the upstream side of the ridge, a phenomenon that is amplified when decreasing the ridge width. The measured energy matches the model prediction in all but one experiment. This gap is likely related to the formation of a vertical swirl close to the ridge that limits energy transfers to internal waves.

These results contributes to improve our understanding of the internal wave dynamics and to better represent their effects in oceanic models.

1. Introduction

Internal waves play an important role in sustaining the Meridional Overturning Circulation (MOC) by inducing a wide range of turbulent mixing processes [Sarkar and Scotti, 2017; MacKinnon et al., 2017]. Internal waves are predominantly generated either at the surface by wind fluctuations or at the ocean bottom by the interaction of flow with topography. Two seemingly independent classes of topographic internal waves have been distinguished in previous studies. In regions where strong barotropic tidal flows interact

DRAFT

June 15, 2020, 11:47am

DRAFT

X - 4

with large scale topographies, such as the Mid-Atlantic Ridge, internal waves are usu-48 ally organized as propagating beams called internal tides [Polzin et al., 1997; Gregg and 49 Ozsoy, 1999; Melet et al., 2013a; Nycander, 2005, 2006; Buijsman et al., 2014]. On the 50 other hand, in regions where a vigorous geostrophic eddy fields interact with smaller scale 51 topographies such as in the Antarctic Circumpolar Current, internal waves are organized 52 as quasi-steady lee waves [Nikurashin and Ferrari, 2010a, b; Dossmann et al., 2016a; 53 Yang et al., 2018; Zheng and Nikurashin, 2019]. The respective contributions of these two 54 classes of internal waves to water-masses transformation have been assessed independently 55 using the linear theory of Bell [1975] and semi-empirical parameterizations [St. Laurent 56 et al., 2002; Polzin, 2009] leading to global estimates maps of internal wave mixing (e.g. 57 Nikurashin and Ferrari [2011, 2013]; Melet et al. [2013b]). 58

Barotropic tides coexist with quasi-steady flows at many internal wave generation sites. 59 This observation motivated *Shakespeare* [2020] to revisit the classical linear internal wave 60 generation theory of *Bell* [1975]. The combination of a barotropic tidal component and a 61 steady flow causes a linear coupling between internal tides and lee waves that has not been 62 accounted for in previous internal wave energy budgets. Bell [1975]'s theory was extended 63 to include the coupling between multiple flow constituents. A new nomenclature was 64 introduced to distinguish internal waves propagating as 'free beams' from 'flow-trapped' 65 waves that can only propagate in the presence of the flow that supports them. 66

The extended linear theory predicts major changes in the internal wave dynamics owing to this coupling that affects:

DRAFT

• The energy fluxes to lee waves. Lee wave generation and induced mixing are largely cancelled (by up to 90%- in some key areas, such as the Drake Passage) and is overestimated by 13 - 19% on a global scale using the classical approach.

The generation of internal tides beyond the critical latitudes. Doppler-shifted internal
wave beams can propagate in regions where they would dissipate in the absence of a steady
flow, hence modifying the spatical distribution of internal waves and subsequent mixing.

• The existence of a net wave stress above abyssal hills. The linear coupling causes an asymmetry between the upstream and downstream internal tide beams generated over hills, which induces a net wave stress of $0.01 - 0.1 \text{ N/m}^2$ (comparable to the local wind stress).

⁷⁹ Several limitations have been put forward by *Shakespeare* [2020] regarding the appli-⁸⁰ cation of the classical and extended theory of *Bell* [1975]. First, the model is linear and ⁸¹ does not account for finite amplitude effects that induce flow blocking and saturation in ⁸² the supercritical regime. Second, the large uncertainties on the deep-ocean stratification ⁸³ and geostrophic flows lead to an important variability of internal wave energy, momentum ⁸⁴ and mixing budgets between studies.

The use of laboratory experiments permits exploration of the dynamics of internal waves and the pathway to mixing in controlled parameter ranges that are relevant to real oceanic processes [Aguilar et al., 2006; Zhang et al., 2007; Bourget et al., 2013; Dossmann et al., 2011, 2016b]. The turbulent regime is sufficient for the development of nonlinear processes leading to the direct cascade to mixing scales. In a previous experimental study, both the linear lee wave field predicted by Bell's theory and a complex nonlinear dynamics in the boundary layer were captured using the light attenuation technique [Dossmann et al.,

DRAFT

⁹² 2016a]. Mixing diagnostics were performed and showed that the nonlinear processes close
⁹³ to the topography overcome internal wave induced mixing in ocean relevant parameter
⁹⁴ regimes, highlighting the need to account for these effects in the mass and energy budgets
⁹⁵ of the Meridional Overturning Circulation [*Todd*, 2017; *MacKinnon et al.*, 2017; *Zheng*⁹⁶ and Nikurashin, 2019].

The present work aims to describe the linear dynamics of topographic internal waves 97 forced by the combination of an oscillatory flow and a steady flow, and to accurately 98 quantify the associated energy flux. It relies on an experimental campaign for which the 99 flow, topography and background stratification are precisely controlled and the flow dy-100 namics is measured at high space-time resolution. This idealized approach permits direct 101 comparison of the linear energy flux estimated from the extended theory of Bell [1975] 102 against measurements in an identical configuration, allowing the generation of internal 103 waves as well as non-linear structures that also participate to mixing. 104

The manuscript is organized as follows. In section 2, the extended linear theory is described and the different internal wave regimes are introduced. The experimental setup is presented in section 3. The internal wave dynamics and energy flux fields are described in sections 4 and 5. Implications for IW energy budgets in the ocean are discussed in section 6.

2. Linear theory

2.1. Extended linear theory

In order to make meaningful comparisons with our experimental results, it is necessary to slightly extend previous models for linear internal wave generation by flow over topography to simultaneously include: (i) finite fluid depth, (ii) both steady and oscillatory

¹¹³ background flow, (iii) non-negligible excursion parameters, (iv) non-hydrostatic dynam-¹¹⁴ ics, and (v) a mechanism of dissipation for the waves. As such, the theory here can be ¹¹⁵ thought of as generalising that of *Bell* [1975] and *Shakespeare* [2020] to include finite-depth ¹¹⁶ and wave dissipation, or the theory of *Shakespeare et al.* [2020a] to include non-negligible ¹¹⁷ excursion parameters.

Here, the background flow of interest is a combined steady and oscillatory flow: $U = U_c + U_0 \cos \omega t$, where ω is the forcing frequency and t is time. For completeness, we will include a non-zero Coriolis parameter f and three-dimensionality in the following. The simplest starting point for our derivation is Equation (21)-(25) in *Shakespeare et al.* [2020a] which describes the vertical velocity w(x, y, z, t) and pressure p(x, y, z, t) associated with wave generation at the ocean bottom (z = -H) and a rigid-lid boundary condition at the ocean surface (z = 0):

$$\widehat{w}_n = \widehat{w}_n|_{-H} \frac{e^{-imz - \gamma z} - e^{imz + \gamma z}}{e^{imH + \gamma H} - e^{-imH - \gamma H}},$$
(1a)

and,

X - 8

$$\widehat{p}_n = -\frac{N^2 - \omega_n^2}{m\omega_n} \widehat{w}_n|_{-H} \frac{e^{-imz - \gamma z} + e^{imz + \gamma z}}{e^{imH + \gamma H} - e^{-imH - \gamma H}},$$
(1b)

where

$$m = \operatorname{sgn}(\omega_n) K \sqrt{\frac{N^2 - \omega_n^2}{\omega_n^2 - f^2}},$$
(1c)

is the vertical wavenumber, $K = \sqrt{k^2 + l^2}$ is the horizontal wavenumber modulus, and

$$\gamma = \alpha m \frac{\omega_n (2N^2 - \omega_n^2 - f^2)}{2(N^2 - \omega_n^2)(\omega_n^2 - f^2)},$$
(1d)

is the decay rate per unit wavelength, with $\gamma \ll m$ for valid solutions. The magnitude of this decay rate is determined by the Rayleigh drag α (in inverse seconds) — this damping D R A F T June 15, 2020, 11:47am D R A F T ¹²⁷ is used as a purely numerical tool herein, as will be explained below. Equation (1) may be
¹²⁸ generalised to include steady and tidal flows, and large excursion distances, by summing
¹²⁹ over intrinsic frequencies for each harmonic

$$\omega_n = n\omega + \boldsymbol{k} \cdot \boldsymbol{U}_c. \tag{2}$$

where n is the harmonic number, and the Fourier transform in x and y of the total vertical velocity is thus

$$\widehat{w}(k,l,z,t) = \sum_{n=-\infty}^{\infty} \widehat{w}_n e^{i\omega_n t},$$
(3)

and similarly for pressure. The bottom boundary condition on the vertical velocity, associated with the no-normal flow condition (i.e. $w = U \cdot \nabla h$ for topographic height h in the small amplitude approximation) gives

$$\widehat{w}_n|_{-H} = i\omega_n \widehat{h} J_n\left(\frac{\boldsymbol{k} \cdot \boldsymbol{U}_0}{\omega}\right),\tag{4}$$

where J_n is the order *n* Bessel function of the first kind (see, for example, *Bell* [1975]; *Shakespeare* [2020]).

The solution in this form is expressed in a coordinate frame x_0 following the total flow U (that is, $x = x_0 + \int_0^t U \, dt'$ where x is the Eulerian position, and x_0 the flow-following coordinate). We use the above solutions to compute the theoretical energy flux as a function of time and space via

$$F(x, y, z, t) = \mathcal{F}^{-1}[\widehat{w}e^{-i\boldsymbol{k}\cdot\boldsymbol{U}_{c}t}]\mathcal{F}^{-1}[\widehat{p}e^{-i\boldsymbol{k}\cdot\boldsymbol{U}_{c}t}],$$
(5)

where $\mathcal{F}^{-1} = 1/(4\pi^2) \int \int e^{-i(kx+ly)} dk \, dl$ denotes the inverse Fourier transform. To calculate these inverse Fourier transforms numerically, the vertical velocity and pressure

fields must decay towards the boundaries of the (numerical) domain; otherwise, the waves continuously wrap around and the numerical solution is meaningless. The decay of the wave fields is ensured by turning the Rayleigh drag α in (1d) to a sensible value (interested readers are referred to *Shakespeare et al.* [2020b] for a discussion of the impact α on the wave field).

2.2. Internal wave regimes

Four internal wave regimes are expected due to the combination of a steady and an oscillatory forcing flow. The linearization of Euler's equation under the Boussinesq approximation leads to the following dispersion relation for internal waves:

$$\tilde{\omega}^2 = N^2 \sin^2 \theta^2 + f^2 \cos^2 \theta^2, \tag{6}$$

with $\tilde{\omega} = \omega - \mathbf{k} \cdot \mathbf{U}_{c}$, N and f the intrinsic internal wave frequency, the Brunt-Väisälä 149 frequency and the local Coriolis parameter. Here, ω is the frequency of the oscillatory 150 flow, $k \cdot U_c$ is a quasi Doppler frequency shift induced by the geostrophic flow U_c onto 151 the internal wave of horizontal wavenumber k, and θ is the angle of the group velocity 152 with respect to the horizontal. Equation (6) implies that $f < |\tilde{\omega}| < N$ for propagating 153 internal waves. Depending on the signs of $|\mathbf{k} \cdot \mathbf{U}_{\mathbf{c}}|/\omega$ (the dot-product of the inverse wave 154 phase speed and the steady flow velocity) and $\tilde{\omega}$ four non-evanescent bands are allowed: 155 lee waves, upstream and downstream beams, and tidal lee waves. The four bands are 156 represented in Fig. 1a. in the non-rotating case. 157

The first non-evanescent band, indicated by the solid red line in Fig. 1a. corresponds to the classical lee wave generation by the steady forcing ($\omega = 0, f < |\mathbf{k} \cdot \mathbf{U}_{c}| < N$) downstream of the steady flow.

DRAFT

There are two non-evanescent bands (blue shaded areas in Fig. 1a) with positive intrinsic 161 frequencies $\tilde{\omega} > 0$, one in the upstream direction $(\mathbf{k} \cdot \mathbf{U}_{c}/\omega < 0, f - |\mathbf{k} \cdot \mathbf{U}_{c}| < \omega < N - |\mathbf{k} \cdot \mathbf{U}_{c}|$ 162 $\boldsymbol{U_c}|), \, \text{and one in the downstream direction } (\boldsymbol{k} \cdot \boldsymbol{U_c}/\omega > 0, \, f + |\boldsymbol{k} \cdot \boldsymbol{U_c}| < \omega < N + |\boldsymbol{k} \cdot \boldsymbol{U_c}|).$ 163 In these two bands, the internal wave field is organized as beams propagating upstream 164 and downstream of the flow. The dispersion relation for the downstream and upstream 165 beams are shown in Figs 1b and c for the three experimental ridges. In particular internal 166 waves are able to propagate in regions where $\omega > N$ (superbuoyancy downstream beam) 167 or $\omega < f$ (subinertial upstream beam). 168

In the downstream direction, non-evanescent IWs with $\tilde{\omega} < 0$ are allowed in the band $|\mathbf{k} \cdot \mathbf{U}_{c}| - N < \omega < |\mathbf{k} \cdot \mathbf{U}_{c}| - f$ (red shaded areas in Fig. 1a). In this band, IWs are organized as a steady lee wave field at larger wavenumber than classical lee waves and which only exists in the presence of a barotropic forcing. The term tidal lee waves was introduced by *Shakespeare* [2020] to refer to this IW regime.

Three of the four internal wave regimes generated by the combination of a steady and 174 an oscillatory flow are shown in Fig.2 using the extended linear model described in the 175 previous section. The two forcing flows are artificially uncoupled in Fig.2a, in which 176 one identifies the generation of symmetrical internal tidal beams on either side of the 177 ridge as well as lee waves. Turning on the coupling leads to more complex dynamics. 178 The symmetry between the upstream beam and the downstream beam is broken by the 179 presence of the steady flow as will be discussed in section 4. Lee waves are still generated 180 on the downstream side, but their amplitude is weakened by the coupling mechanism. For 181 the chosen forcing parameters, the generation of tidal lee waves is limited by the relatively 182 small value of the steady flow, as will be the case in the experiments. 183

DRAFT

June 15, 2020, 11:47am

In this work experiments are performed in a non-rotating, strongly stratified flow. The 184 effects of rotation on the IW dynamics are neglected on the experiment timescale. The 185 dispersion relation (6) can therefore be simplified to $\omega = N \sin(\theta) - |\mathbf{k} \cdot \mathbf{U}_c|$ for the 186 upstream beam and $\omega = N \sin(\theta) + |\mathbf{k} \cdot \mathbf{U_c}|$ for the downstream beam, as shown in 187 Fig. 1. The mean excursion distance parameter $\bar{\epsilon} = |\mathbf{k} \cdot \mathbf{U}_{\mathbf{c}}|/\omega$ is introduced to quantify 188 the relative magnitude of the quasi-Doppler shift, and hence the asymmetry between the 189 upstream and downstream beams. The beam horizontal wavenumber is not prescribed 190 by the dispersion relation, but scaled by the topographic width. Hence, the asymmetry 191 between the upstream and downstream beams is expected to be enhanced for strong 192 steady flows impinging on smaller scale topographies. 193

The experimental campaign aims to describe the internal wave field induced by the combined forcings and the effect of the asymmetry on the transported energy fluxes.

3. Experimental set-up

3.1. Apparatus

Experiments have been performed in a 5.26 m long, 20 cm wide and 40 cm high perspex 196 tank in the Geophysical Fluid Dynamics laboratory at ANU. The air temperature is 197 controlled at $20 \pm 1^{\circ}$ C. The experimental setup is sketched in Fig. 3. The tank is filled 198 with the double-bucket technique using two drums, one initially filled with brine and 199 the other with freshwater. The drums are filled with hot water to allow an efficient salt 200 dissolution. A fixed volume of 10 mL of red food dye is added to the drum filled with 201 brine. It is used as a passive tracer for salinity as described in the following subsection. 202 The tank is filled to a height of $H \approx 27$ cm in approximately three hours using two 203 floating devices. The slow flow rate allows a linear stratification to be obtained over the 204

DRAFT

June 15, 2020, 11:47am

fluid depth, while shallow mixed layers of less than 1 cm are observed close to the tank bottom and free surface, due to boundary exchanges. In most fills the two drums have been prepared 24 hours before the start of the fill to achieve temperature equilibrium between water and air. Several fills have been carried out with water hotter than air, which led to the formation of double diffusive layers. These layers had no visible impact on the internal wave dynamics and permitted observation of the interfacial displacement using the shadowgraph technique in qualitative experiments.

For most experiments, a constant Brunt-Väisälä frequency $N_0 = 0.7$ rad/s is set, with $N_0 = \sqrt{-g/\rho_0 \times d\bar{\rho}/dz}$. g = 9.81 m/s², $\rho_0 = 1000$ kg/m³, $\bar{\rho}$ and z are the gravity, the reference density, the density profile and the vertical coordinate with the origin at the tank bottom, respectively. In total, 14 fills have been performed during the experimental campaign, for which the forcing parameters were varied.

The forcing is performed by towing a bidimensional ridge of Gaussian shape over the length of the tank, at the free surface of the fluid. The ridge profile is given by

$$h(x) = h_0 \exp\left(-\left(k_r x\right)^2\right),\tag{7}$$

where x is the horizontal coordinate with the origin at the center of the ridge base, 219 $h_0 = 3.5$ cm is the ridge height and k_r is the characteric horizontal wavenumber. Ex-220 periments have been performed with three different ridges of varying e-folding width 221 $l_r = 1/k_r = 1.5$ cm (ridge A), $l_r = 5$ cm (ridge B) and $l_r = 16$ cm (ridge C). The 222 ridges are attached to a 80 cm long plate to allow for boundary layers to develop during 223 motion. The weight of the plate is adjusted so that the ridge is completely submerged, 224 and the flat section of the plate lies a few millimeters below the free surface. The ridge 225 motion is controlled by a Smartmotor interface via a wire attached to a rotating cylinder. 226

DRAFT June 15, 2020, 11:47am DRAFT

The program independently controls a steady and an oscillatory rotation of the cylinder, which respectively impose a constant velocity U_c and an oscillatory velocity $U_0 \cos(\omega t)$ to the ridge. The ridge transits 1 to 20 times over the length of the tank during an experiment. The flow is decelerated/accelerated over 30 cms close to the end walls to ensure a continuous forcing velocity.

Hence, this configuration permits simulation of the combined effects of a geostrophic and a barotropic forcing interacting with an oceanic ridge. The relative magnitude of the two forcings as well as the tidal period can be adjusted to reach a parameter space relevant for oceanic configurations. The initially linear stratification represents the regular density gradient in the deep ocean.

Five selected experiments showing different aspects of the asymmetric generation of free beams - which are the main source of IW energy transport over hills apart from the Southern Ocean - are discussed in the following sections. Table 1 summarizes the relevant parameters of the experiments.

3.2. Light attenuation technique

The light attenuation technique infers local density from absorption measurements at high spatial resolution [*Allgayer and Hunt*, 1991; *Hacker et al.*, 1996; *Sutherland et al.*, 2012]. This method has previously been used to describe the dynamics of internal waves generated in the lee of a moving topography as well as the induced mixing [*Dossmann et al.*, 2016a].

The light absorption $A(x, z, t) = \log(I_0(x, z)/I(x, z, t))$ due to the red food dye is measured using the intensity field from a reference image $(I_0(x, z))$ and from an image recorded in the course of the experiment I(x, z, t). The initial concentration of red dye is chosen

DRAFT June 15, 2020, 11:47am DRAFT

²⁴⁹ such the absorption law is in the linear (Beer-Lambert) regime: $A = \epsilon_{\lambda} W c$, where ϵ_r , W²⁵⁰ and c are the extinction coefficient for the red component of the camera sensor, the tank ²⁵¹ width and the dye concentration.

A calibration is applied using two density samples from the flow to eventually convert 252 absorption values to cross-tank averaged density anomalies $\Delta \rho(x, z, t)$ at each pixel. A 253 median filter is applied over 10 pixels in the two directions to remove measurement noise. 254 The reader can refer to *Dossmann et al.* [2016a] for a detailed description of the light 255 attenuation technique. A LED panel of dimensions $100 \text{ cm} \times 60 \text{ cm}$ is placed to illuminate 256 to central part of the tank. Images are recorded in RAW format at 14 bits using a 257 Nikon D300 DSLR camera equipped with a 200 mm telephoto lens. The camera is placed 258 approximately 6 m away from the tank to minimize parallax effects. 259

4. Internal wave dynamics

In this section we investigate the dynamics induced by the flow/ridge interaction for varying the mean excursion distance parameter $\bar{\epsilon}$ and forcing frequency ω/N . In experiments II to V, the ridge is moving rightwards at a constant velocity $U_c = 2 \text{ mm/s}$, which represents a leftwards steady flow. Hence the downstream (upstream) region is at the left (right) side of the ridge.

In the absence of the steady forcing flow ($\bar{\epsilon} = 0$) the generation of symmetrical internal wave beams on the two sides of the ridge is measured in experiment I. (Fig. 4). The beams are in phase opposition and their angle matches the expected angle from the linear dispersion relation. The high mode internal wave field is superimposed on a background wave field likely due to fast propagating low modes reflecting at the tank walls.

DRAFT

X - 16

The steady flow is then turned on in experiment II ($\bar{\epsilon} = 0.09$), for the same ridge and 270 forcing frequency as in experiment I (5). The generation of sharp beams occurs on both 271 sides of the ridge with an important asymmetry between the upstream and downstream 272 beams. The upstream beam shows a larger amplitude and is steeper than the downstream 273 beam. The linear prediction of the beam angles including the steady flow is shown by 274 the dashed lines in Fig. 5. Both beam angles match the linear dispersion relation. The 275 slow rightwards ridge motion induces the periodic formation of a vortex on the lee side of 276 the ridge as seen at t = 0.71T (circled). This vortex only forms when the oscillatory and 277 steady forcings are combined and is likely to contribute to mixing in the vicinity of the 278 ridge. 279

Decreasing the ridge width in experiment III leads to a larger value of $\bar{\epsilon} = 0.36$ (see Fig. 6). Again sharp internal wave beams form periodically on the two sides of the ridge, with a larger angle difference between the upstream and downstreams beams than in experiment II. Contrary to experiment II, no vortex is observed close to the ridge. This indicates that the generation of this non-linear structure in the boundary layer is dependent on the shape of the topography.

The Doppler shift induced by the steady flow interacting with a sufficiently narrow ridge allows for the generation of a superbuoyancy IW beam in experiment IV ($\bar{\epsilon} = 0.20$ and $\omega/N = 1.21$, Fig. 6)). The propagating beam is generated on the downstream side on the ridge, while evanescent waves remain on the upstream side as predicted by the regime diagram in Fig. 1a. The superbuoyancy beam interferes with propagating subharmonic internal waves with a similar amplitude.

DRAFT

The asymmetry between the upstream and downstream dynamics is illustrated with on horizontal spacetime diagrams measured at mid-depth for exp II-IV in Fig. 8. The larger amplitude and shorter wavelength of the upstream beam are clearly visible in exps II-III. In exp IV, the periodic density anomalies do not propagate on the upstream side (as shown by the absence of slope in the spacetime diagram) while the subcritical downstream beam and a supercritical subharmonic beam propagate on the downstream side.

Finally, the ridge width is increased to the largest value of 16 cm in experiment V, 298 while keeping the same steady velocity and forcing frequency as in exp I-III (Fig. 9). 299 Hence the mean excursion parameter $\epsilon = 0.03$ is small relative to previous experiments 300 and one expects a weak asymmetry between the downstream and upstream sides. In fact, 301 periodic density anomalies with a horizontal scale comparable to l_r are generated by the 302 ridge motion. These anomalies appear to be dominated by propagating low modes - by 303 constrast with the sharper beams in exp I-IV. No asymmetry in the beams is measured 304 in this experiment. 305

5. Energy Flux

The measured asymmetry between the upstream and downstream beams impacts the energy fluxes, $\mathbf{J}(x, z, t) = p\mathbf{v}$, carried by internal waves and eventually the induced mixing. The direct measure of energy fluxes remains an experimental challenge as it relies on the simultaneous measurements of the pressure and velocity fields. While it is possible to perform simultaneous measurements of velocity fields with other fields related to density and their derivatives to assess buoyancy fluxes, the authors are not aware of an experimental technique to directly measure the pressure anomaly field.

DRAFT

X - 18

The method developed by Allshouse et al. [2016]; Lee et al. [2018] is applied to extract 313 the linear energy flux radiated away by internal wave beams. In this method the pressure 314 field is calculating from a Green's function method using the density anomaly field while 315 the velocity field is obtained by integrating the continuity equation using the incompress-316 ibility assumption. This method was tested on experimental Schlieren data which provides 317 density gradient fields that are then integrated to obtain density anomalies. Agreement 318 with Navier-Stokes simulations was obtained for small-amplitude internal wave field. In 319 the present work the density anomalies are directly measured from the light attenuation 320 technique, which suppresses the spatial integration step. Instantaneous energy fluxes are 321 calculated using the Matlab GUI EnergyFlux introduced by Allshouse et al. [2016]. Their 322 mean contribution is post-processed and compared with the energy fluxes calculated from 323 the extended linear theory described in 2.1 in Fig. 10. 324

When no steady flow is present, the energy flux is focussed in the symmetrical beams 325 generated on either side of the ridge (exp I, Fig.10a. and d.). The predicted and ex-326 perimental flux fields closely match in amplitude and direction. Adding a steady flow 327 leads to an asymmetry in both the shape and the amplitudes of the energy carried away 328 by the beams as described in section 4. In both experiments II ($\bar{\epsilon} = 0.36$, Fig10c. and 329 f.) and III ($\bar{\epsilon} = 0.09$ Fig. 10b. and e.), the steady flow increases the energy flux in the 330 upstream beam relative to the downstream beam. The experimental beam amplitudes are 331 twice as large in exp II (ridge B) as in exp III (ridge C), showing the strong topographic 332 control exerted by the ridge slope on the energy fluxes. The theoretical and experimental 333 amplitudes are similar in the narrower ridge experiment III, however the linear energy 334 fluxes are overestimated by a typical factor of 2 in exp II. A plausible explanation for 335

DRAFT

this overestimation is that other nonlinear processes are at play in exp II that lead to a saturation in the energy fluxes carried away by the linear IW beams. This explanation is consistent with the observation of a vertical vortex generated every tidal period in exp II (Fig. 5).

A visible beam doubling appears in both the predicted and experimental mean fluxes when measured in the Eulerian frame (exp II and III) owing to the relatively large value of the tidal excursion parameter. When measured in the co-tidal frame, the mean energy flux is focussed in a single beam as shown for exp I.

6. Discussion

The combination of a steady flow and an oscillatory flow impinging up on a topography 344 leads to the generation of internal waves on both sides of the ridge in all selected experi-345 ments, except for exp IV in which the Doppler-shift prevents the generation of upstream 346 internal waves. The latter case is one where downstream internal waves are 'flow-trapped' 347 as they can only propagate in the presence of the steady flow see the discussion of Shake-348 speare, 2020]. In experiments I, II, and III internal waves are organized as propagating 349 'free beams' that are modified by the steady flow but would still propagate if the steady 350 flow is cancelled. These waves are asymmetric in their direction of propagation, their am-351 plitude and their wavelength. The asymmetry is increased by increasing the magnitude of 352 the steady flow and by decreasing the ridge width. The beam angles agree well with the 353 extended dispersion relation, including a Doppler-shift, showing that these IWs propagate 354 in a linear fashion. All experiments were performed in regimes where the internal wave 355 field is dominated by propagating upstream and downstream beams relative to lee waves 356

DRAFT

X - 20

³⁵⁷ owing to the small values of the steady flow. Hence, the signature of lee waves is not ³⁵⁸ visible in the density anomalies, nor in the energy fluxes.

The processed energy flux fields are successfully compared against the extended linear 359 theory described in section 2.1. The presence of a steady flow induces a larger amplitude 360 and greater energy flux in the upstream beam compared with the downstream beam. 361 The Doppler-shift leads to a more efficient barotropic to baroclinic conversion as the 362 upstream beam approaches the near-critical regime described in *Dossmann et al.* [2011] 363 while the downstream beam slope flattens. The measured asymmetry confirms the larger 364 integrated energy fluxes measured on the upstream side of ocean ridges and consequently 365 the existence of a net wave stress over abyssal hills proposed in *Shakespeare* [2020], noting 366 that these experiments are performed in a limited framework compared with the ocean 367 (non-rotating, single tidal forcing). Partial cancellation of the energy flux is expected 368 when including multi-tidal forcing. Future experiments will focus on quantifying this 369 cancellation by superimposing several tidal components in the oscillatory flow. 370

The steady flow amplitude is too weak to observe the generation of non-linear boundary layer structures (hydraulic jump, topographic blocking) described in *Dossmann et al.* [2016a]. However the coupling between the steady and oscillatory flows leads to the generation of periodic vortices that are likely to participate to mixing close to the topography and limit the energy transfer toward IW beams. The conditions for the generation of these vortices and their contribution to mixing relative to internal waves will be investigated in a companion experimental study.

The pathway to internal wave induced mixing in the ocean is likely to be affected by the coupling between the steady flow and the oscillatory flow. Internal wave mixing

DRAFT

³⁸⁰ processes are classified as either far-field or near-field, depending whether the internal wave ³⁸¹ energy escapes the generation site as propagating low-modes and dissipates elsewhere, or ³⁸² is dissipated in the vicinity of the topography through the breaking of higher mode internal ³⁸³ waves. The various routes for far-field and near-field internal wave dissipation are reviewed ³⁸⁴ by *MacKinnon et al.* [2017].

Low mode internal waves, generated by large scale topograpy (order 100 kms) are appar-385 ently unaffected by the presence of a steady flow, as illustrated in experiment V and con-386 sistent with theory predicting a weak Doppler shift at small wavenumber k_r . However, the 387 remote breaking process of these waves involves a direct cascade of energy through scat-388 tering by topography, breaking at continental slopes, interactions with mesoscale steady 389 flows and eddies *Dunphy and Lamb* [2014], and wave-wave interaction through Paramet-390 ric Subharmonic Instability [MacKinnon et al., 2013]. These mechanisms induce internal 391 waves at larger wavenumber that can be subject to the Doppler shift in the presence of a 392 steady flow or other tidal constituents. The role of the Doppler shift in the pathway to 393 mixing for low mode internal tides is yet to be investigated. 394

For high mode internal waves generated at smaller scale abyssal hills of order 10 kms, the 395 nearfield pathway to mixing is directly affected by the Doppler shift as seen in experiment 396 II-IV. Several processes are likely to be involved in transfer of energy to the scales of 397 mixing directly above the topography, such as interactions with a background internal 398 wave field, convective instability Buhler and Muller [2009] and the PSI mechanism. The 399 mixing efficiency of these mechanisms are controlled by the amplitude and wavenumber 400 of the internal wave beams [Joubaud et al., 2012; Bourget et al., 2013]. In particular, the 401 dominant sites for generation of high modes internal tides at abyssal hills are the mid-402

DRAFT

X - 22

ocean ridges in the Atlantic and western Indian Oceans. At these sites, the asymmetry 403 is particularly favored by the combination of the large topographic roughness of order 10 404 kms, an efficient barotropic to baroclinic conversion of the M2 tide, and the existence of 405 a steady flow. In fact, a global map of the mean excursion distance parameter shows that 406 $\bar{\epsilon} \approx 0.1$ around mid-ocean ridges (Fig. 11), which is the range for asymmetric internal 407 wave beam in the present experiments. The present work and *Shakespeare* [2020]) (their 408 Fig. 11) suggest that nearfield diapycnal mixing processes are enhanced on the upstream 409 side of abyssal hills. On the other hand, in regions where strong quasi-steady flows interact 410 with abyssal roughness, as shown by the red patches in Fig. 11, the generation of lee waves 411 is likely to be locally cancelled in specific areas due to the interaction with the M2 forcing 412 flow as in the Drake Passage. 413

In recent works, new process dependent parameterizations of mixing have been pro-414 posed to represent the 3D distribution of mixing both in the near-field and in the far-field 415 (relative to internal wave generation sites) using Lagrangian beam tracing from sources 416 to sinks [de Lavergne et al., 2019, 2020]. Four dissipative processes have been identified, 417 including the generation of high mode internal waves at abyssal hills. Global maps of in-418 ternal tide dissipation have been produced by summing the independent contributions of 419 the three main tidal components (M2, S2 and K1). The coupling between these tidal con-420 stituents and with the steady flow is not accounted for but is likely to affect the geography 421 of internal wave sources - by the partial energy transfers between different tidal compo-422 nents - and sinks - by modifying the path taken by internal wave beams to their breaking 423 site. Consequently, the role played by the coupling process on the global distribution of 424 mixing is a key property to investigate with the proposed parameterizations. 425

DRAFT

June 15, 2020, 11:47am

426 Acknowledgments.

This work was supported partly by the french PIA project <u>Lorraine Université d' Excellence</u>, reference ANR-15-IDEX-04-LUE. Yvan Dossmann acknowledges support from the Embassy of France in Australia. Reader can access the supporting data and linear model via an online repository [*Dossmann et al.*, 2020].

References

- Aguilar, D. A., B. R. Sutherland, and D. J. Muraki (2006), Laboratory generation of
 internal waves from sinusoidal topography, *Deep Sea Research Part II: Topical Studies in Oceanography*, 53(1-2), 96–115.
- ⁴³⁴ Allgayer, D. M., and G. R. Hunt (1991), On the application of the light-attenuation
 ⁴³⁵ technique as a tool for non-intrusive buoyancy measurements, *Experimental Thermal*⁴³⁶ and Fluid Science, 38, 257–261.
- ⁴³⁷ Allshouse, M. R., F. M. Lee, P. J. Morrison, and H. L. Swinney (2016), Internal wave
 ⁴³⁸ pressure, velocity, and energy flux from density perturbations, *Physical Review Fluids*,
 ⁴³⁹ 1(1), 014,301.
- Bell, T. H. (1975), Topographically generated internal waves in open ocean, Journal of
 Geophysical Research, 80, 320–327.
- Bourget, B., T. Dauxois, S. Joubaud, and P. Odier (2013), Experimental study of parametric subharmonic instability for internal plane waves, *Journal of Fluid Mechanics*,
 723, 1–20.
- ⁴⁴⁵ Buhler, O., and C. Muller (2009), Saturation of the Internal Tides and Induced Mixing
 ⁴⁴⁶ in the Abyssal Ocean, American Meteorological Society, 39.

DRAFT

- ⁴⁴⁷ Buijsman, M. C., J. M. Klymak, S. Legg, M. H. Alford, D. Farmer, J. A. MacKinnon,
- J. D. Nash, J.-H. Park, A. Pickering, and H. Simmons (2014), Three-dimensional double-
- ridge internal tide resonance in luzon strait, Journal of Physical Oceanography, 44(3),
 850–869.
- ⁴⁵¹ de Lavergne, C., S. Falahat, G. Madec, F. Roquet, J. Nycander, and C. Vic (2019), Toward ⁴⁵² global maps of internal tide energy sinks, *Ocean Modelling*, 137, 52–75.
- 453 de Lavergne, C., C. Vic, G. Madec, F. Roquet, A. F. Waterhouse, C. B. Whalen,
- 454 Y. Cuypers, P. Bouruet Aubertot, B. Ferron, and T. Hibiya (2020), A Parameterization
- of Local and Remote Tidal Mixing, Journal of Advances in Modeling Earth Systems,
 12(5), e2020MS002,065.
- ⁴⁵⁷ Dossmann, Y., A. Paci, F. Auclair, and J. W. Floor (2011), Simultaneous velocity and ⁴⁵⁸ density measurements for an energy-based approach to internal waves generated over a ⁴⁵⁹ ridge, *Experiments in Fluids*, 51(4), 1013–1028.
- ⁴⁶⁰ Dossmann, Y., M. G Rosevear, R. W. Griffiths, G. O. Hughes, M. Copeland, et al.
 (2016a), Experiments with mixing in stratified flow over a topographic ridge, *Journal*⁴⁶² of Geophysical Research: Oceans, 121(9), 6961–6977.
- ⁴⁶³ Dossmann, Y., B. Bourget, C. Brouzet, T. Dauxois, S. Joubaud, and P. Odier (2016b),
 ⁴⁶⁴ Mixing by internal waves quantified using combined piv/plif technique, *Experiments in*⁴⁶⁵ Fluids, 57(8), 132.
- ⁴⁶⁶ Dossmann, Y., C. Shakespeare, and A. M. Hogg (2020), Experimental data and linear
 ⁴⁶⁷ model for "Asymmetric internal tide generation in the presence of a steady flow", doi:
 ⁴⁶⁸ 10.5281/zenodo.3889612.

- ⁴⁷⁰ mode internal tide by mesoscale eddy interaction, Journal of Geophysical Research:
 ⁴⁷¹ Oceans, 119, 523–536.
- 472 Goff, J. A. (2010), Global prediction of abyssal hill root-mean-square heights from
- small-scale altimetric gravity variability, Journal of Geophysical Research: Solid Earth,
 115(B12).
- Gregg, M. C., and E. Ozsoy (1999), Mixing on the black sea shelf north of the bosphorus, *Geophysical Research Letters*, 26(13), 1869–1872.
- Hacker, J., P. F. Linden, and S. B. Dalziel (1996), Mixing in lock-release gravity currents,
 Dyn. Atmos Oceans, 24, 183–195.
- Joubaud, S., J. Munroe, P. Odier, and T. Dauxois (2012), Experimental parametric subharmonic instability in stratified fluids, *Physics of Fluids*, 24 (4), 041703.
- 481 Kiss, A. E., A. M. Hogg, N. Hannah, F. Boeira Dias, G. B. Brassington, M. A. Cham-
- berlain, C. Chapman, P. Dobrohotoff, C. M. Domingues, E. R. Duran, et al. (2020),
- Access-om2 v1. 0: a global ocean-sea ice model at three resolutions, Geoscientific Model
 Development, 13(2), 401-442.
- Lee, F. M., M. R. Allshouse, H. L. Swinney, and P. J. Morrison (2018), Internal wave energy flux from density perturbations in nonlinear stratifications, *Journal of Fluid Mechanics*, 856, 898–920.
- 488 MacKinnon, J. A., M. H. Alford, O. Sun, R. Pinkel, Z. Zhao, and J. Klymak (2013),
- Parametric subharmonic instability of the internal tide at 29 n, Journal of Physical
 Oceanography, 43(1), 17–28.

469

June 15, 2020, 11:47am

X - 25

- ⁴⁹¹ MacKinnon, J. A., Z. Zhao, C. B. Whalen, A. F. Waterhouse, D. S. Trossman, O. M. Sun,
- ⁴⁹² L. C. St. Laurent, H. L. Simmons, K. Polzin, R. Pinkel, et al. (2017), Climate process
- team on internal wave-driven ocean mixing, Bulletin of the American Meteorological
 Society, 98(11), 2429-2454.
- ⁴⁹⁵ Melet, A., M. Nikurashin, C. Muller, S. Falahat, J. Nycander, P. G. Timko, B. K. Arbic,
 ⁴⁹⁶ and J. A. Goff (2013a), Internal tide generation by abyssal hills using analytical theory,
- ⁴⁹⁷ Journal of Geophysical Research: Oceans, 118(11), 6303–6318.
- Melet, A., R. Hallberg, S. Legg, and K. Polzin (2013b), Sensitivity of the ocean state to the
 vertical distribution of internal-tide-driven mixing, *Journal of Physical Oceanography*,
 43(3), 602–615.
- Nikurashin, M., and R. Ferrari (2010a), Radiation and dissipation of internal waves gen erated by geostrophic motions impinging on small-scale topography: Theory, *Journal* of Physical Oceanography, 40, 1055–1074.
- Nikurashin, M., and R. Ferrari (2010b), Radiation and dissipation of internal waves gen erated by geostrophic motions impinging on small-scale topography: Application to the
 southern ocean, *Journal of Physical Oceanography*, 40, 2025–2042.
- Nikurashin, M., and R. Ferrari (2011), Global energy conversion rate from geostrophic
 flows into internal lee waves in the deep ocean, *Geophysical Research Letters*, 38,
 L08,610.
- Nikurashin, M., and R. Ferrari (2013), Global energy conversion rate from geostrophic
 flows into internal lee waves in the deep ocean, *Geophysical Research Letters*, 40, 3133–
 3137.

- Nycander, J. (2005), Generation of internal waves in the deep ocean by tides, J. Geophys. 513 *Res*, *110*, C10,028. 514
- Nycander, J. (2006), Tidal generation of internal waves from a periodic array of steep 515 ridges, Journal of Fluid Mechanics, 567, 415. 516
- Polzin, K. L. (2009), An abyssal recipe, Ocean Modelling, 30(4), 298–309. 517
- Polzin, K. L., J. M. Toole, J. R. Ledwell, and R. W. Schmitt (1997), Spatial variability 518 of turbulent mixing in the abyssal ocean, Science, 276(5309), 93–96. 519
- Sarkar, S., and A. Scotti (2017), From topographic internal gravity waves to turbulence, 520 Annual Review of Fluid Mechanics, 49, 195–220. 521
- Shakespeare, C. J. (2020), Interdependence of internal tide and lee wave generation at 522 abyssal hills: global calculations, Journal of Physical Oceanography, 50(3), 655-677.
- Shakespeare, C. J., B. K. Arbic, and A. M. Hogg (2020a), The drag on the barotropic 524 tide due to baroclinic motions, Journal of Physical Oceanography, in review. 525
- Shakespeare, C. J., A. F. Waterhouse, B. K. Arbic, A. M. Hogg, J. A. MacKinnon, and 526
- H. L. Simmons (2020b), A simple model for dissipating and reflecting internal waves, 527
- and its application to oceanographic observations, Journal of Physical Oceanography, 528 in review. 529
- St. Laurent, L. C., H. L. Simmons, and S. R. Jayne (2002), Estimating tidally driven 530 mixing in the deep ocean, Geophysical Research Letters, 29, 2106. 531
- Sutherland, B. R., B. Lee, and J. K. Ansong (2012), Light attenuation experiments on 532 double diffusive plumes and fountains, Physics of Fluids, 24, 066,605. 533
- Todd, R. E. (2017), High-frequency internal waves and thick bottom mixed layers observed 534 by gliders in the gulf stream, Geophysical Research Letters, 44(12), 6316–6325. 535

523

- Yang, L., M. Nikurashin, A. M. Hogg, and B. M. Sloyan (2018), Energy loss from transient
 eddies due to lee wave generation in the southern ocean, *Journal of Physical Oceanog- raphy*, 48(12), 2867–2885.
- ⁵³⁹ Zhang, H. P., B. King, and H. L. Swinney (2007), Experimental study of internal gravity
 ⁵⁴⁰ waves generated by supercritical topography, *Physics of Fluids*, 19(9), 096,602.
- ⁵⁴¹ Zheng, K., and M. Nikurashin (2019), Downstream propagation and remote dissipation of
- internal waves in the southern ocean, Journal of Physical Oceanography, 49(7), 1873–

543 1887.

Exp	$U_c \ (cm/s)$	$U_0 \ (cm/s)$	ridge	ω/N	$\overline{\epsilon}$
Ι	0	2	В	0.66	0
II	0.2	2	В	0.66	0.09
III	0.2	2	A	0.70	0.36
IV	0.2	2	A	1.21	0.20
V	0.2	2	С	0.76	0.03

Table 1. Experimental parameters. The experiment is carried out several times to vary the duration of forcing sequences. The latter are separated by resting sequences to allow for measurements of the background stratification.



Figure 1. a. Regimes diagram for internal waves in the presence of an oscillatory and a steady forcing flow. The blue line and the blue shaded area are the non-evanescent bands for pure internal tides and free beams. The red line and the red shaded area are the non-evanescent bands for lee waves and tidal lee waves. The symbols \Diamond , \bigcirc and + show the experiments with ridge A, B and C, respectively.

b. and c. Linear dispersion relations for internal waves in the presence of a steady flow for the downstream (b) and upstream beams (c) for ridges A (solid), B (dashed) and C (dotted).



Figure 2. Vertical velocity (m/s) field for internal wave generation over a Gaussian ridge using the extended linear theory of *Bell* [1975] for independent (a.) and coupled (b.) forcings. The solid arrows show the relative magnitude of the oscillatory (2 cm/s) and steady (0,4 cm/s) flows. The dashed and dotted-dashed arrows indicate the phase propagation in the upstream and downstream beams. The topography is shown shaded in red.



Figure 3. Sketch of the experimental apparatus. The tank is filled using the double-bucket technique with a linearly stratified fluid. The Step motor is computer-controlled to impose a steady (blue arrow) and oscillatory (red arrow) forcing motion to the ridge in Exps II-V, which generates internal wave beams in the body of the flow. The steady motion is set to zero in Exp I. The blue rectangle indicates the measurement area for the light attenuation technique.



Figure 4. Exp I: Density anomaly field $\Delta \rho$ (kg/m)³ ($U_c = 0$ cm/s, $U_0 = 2$ cm/s, $\omega/N = 0.66$, ridge B). The dashed lines show the beam angles predicted by linear theory.



Figure 5. Exp II: Density anomaly field $\Delta \rho \, (\text{kg/m})^3 \, (U_c = 0.2 \text{ cm/s}, U_0 = 2 \text{ cm/s}, \omega/N = 0.66,$ ridge B). The steady ridge motion is to the right of the field. The dashed lines show the beam angles predicted by linear theory. The formation of a vortex on the downstream side of the ridge is shown by the purple dashed circle at t = 0.71 T.

June 15, 2020, 11:47am



Figure 6. Exp III: Density anomaly field $\Delta \rho \ (\text{kg/m})^3 \ (U_c = 0.2 \text{ cm/s}, U_0 = 2 \text{ cm/s}, \omega/N = 0.70$, ridge A). The dashed lines show the beam angles predicted by linear theory.

June 15, 2020, 11:47am



Figure 7. Exp IV: Density anomaly field $\Delta \rho$ (kg/m)³ ($U_c = 0.2 \text{ cm/s}$, $U_0 = 2 \text{ cm/s}$, $\omega/N = 1.21$, ridge A).



Figure 8. Density anomaly space-time diagrams at mid-depth for Exps III (a.), II (b.) and IV (c.). The dashed lines indicates the movement of the ridge due to the steady motion. The whitespace in (c) is where data are unavailable due to the limited camera buffer.



Figure 9. Exp V: Density anomaly field $\Delta \rho \, (\text{kg/m})^3 \, (U_c = 0.2 \text{ cm/s}, U_o = 2 \text{ cm/s}, \omega/N = 0.76,$ ridge C).



Figure 10. Mean energy fluxes measured in the experiments (top panels) and assessed from the extended Bell's theory (bottom panels) for Exp I (a, d), Exp II (b, e) and Exp III (c, f)



Figure 11. Mean excursion distance parameter $\bar{\epsilon} = k \times U_c/\omega_{M2}$ for the M2 tidal frequency, where k is the abyssal hill cut-off wavenumber. The steady near-bottom velocity U_c is obtained from 5-day averaged near-bottom velocities issued from the ACCESS-OM2-01 0.1^o resolution global ocean-ice model [*Kiss et al.*, 2020]. Abyssal hill topography is sourced from *Goff* [2010]

June 15, 2020, 11:47am