

# The Diet of Early Birds Based on Modern and Fossil Evidence and a New Framework for its Reconstruction

Case Miller<sup>1,1</sup> and Michael Pittman<sup>1,1</sup>

<sup>1</sup>The University of Hong Kong

November 30, 2022

## Abstract

Birds are some of the most diverse organisms on Earth, with species inhabiting nearly every conceivable niche in every major biome. As such, birds are vital to our understanding of modern ecosystems. Unfortunately, this is hampered by knowledge gaps relating to the origin of this modern diversity and its role in ecosystems. A crucial part of addressing these shortcomings is improving our understanding of the earliest birds, the non-avian avialans i.e. non-crown birds. The diet of non-avian avialans has been a matter of substantial debate, partly related to some of the ambiguous qualitative approaches that have been used to reconstruct it. Here we review the methods of determining diet in both modern avians and fossil avian and non-avian theropods, and comment on their usefulness when applied to non-avian avialans. We use this to propose a set of comparable, quantitative approaches to ascertain fossil bird diet and on this basis provide a consensus of what we currently know about fossil bird diet. While no single approach can precisely predict diet in birds, each can exclude some diets and narrow the dietary possibilities. We recommend combining [1] dental microwear, [2] landmark-based muscular reconstruction, [3] stable isotope geochemistry, [4] body mass estimations, [5] traditional and/or geometric morphometric analysis, and [6] finite element analysis to accurately reconstruct fossil bird diet. Our review provides specific methodologies to implement each approach and discusses complications future researchers should keep in mind. On this basis we report the current state of knowledge of non-avian avialan diet which remains very incomplete. The ancestral dietary condition in non-avian avialans remains unclear due to a scarcity of data and contradictory evidence in *Archaeopteryx*. Among early non-avian pygostylians, *Confuciusornis* has finite element analysis and mechanical advantage evidence pointing to herbivory, whilst *Sapeornis* only has mechanical advantage evidence indicating granivory, which agrees with fossilised ingested material known for this taxon. The enantiornithine ornithothoracine *Shenqiornis* has mechanical advantage and pedal morphometric evidence pointing to carnivory. In the hongshanornithid ornithuromorph *Hongshanornis*, only mechanical advantage evidence indicates granivory, but this is congruent with evidence of fossilised ingested material in this taxon. The same is true for the songlingornithid ornithuromorph *Yanornis* and its inferred carnivorous diet. Due to the sparsity of robust dietary assignments, no clear trends in non-avian avialan dietary evolution have yet emerged. Dietary diversity may seem to increase through time, but this is a preservational bias associated with a predominance of data from the Early Cretaceous Jehol Lagerstätte. With this new framework and our current synthesis of current knowledge of non-avian non-avian diet, we expect dietary knowledge and evolutionary trends to become much clearer[...]

# The Diet of Early Birds Based on Modern and Fossil Evidence and a New Framework for its Reconstruction

Case V. Miller<sup>1,\*</sup> & Michael Pittman<sup>1,\*</sup>

<sup>1</sup>Vertebrate Palaeontology Laboratory, Division of Earth and Planetary Science, The University of Hong Kong, Pokfulam, Hong Kong SAR, China

\*Authors for correspondence (e-mail: [case.miller@connect.hku.hk](mailto:case.miller@connect.hku.hk); [mpittman@hku.hk](mailto:mpittman@hku.hk))

## ABSTRACT

Birds are some of the most diverse organisms on Earth, with species inhabiting nearly every conceivable niche in every major biome. As such, birds are vital to our understanding of modern ecosystems. Unfortunately, this is hampered by knowledge gaps relating to the origin of this modern diversity and its role in ecosystems. A crucial part of addressing these shortcomings is improving our understanding of the earliest birds, the non-avian avialans i.e. non-crown birds. The diet of non-avian avialans has been a matter of substantial debate, partly related to some of the ambiguous qualitative approaches that have been used to reconstruct it. Here we review the methods of determining diet in both modern avians and fossil avian and non-avian theropods, and comment on their usefulness when applied to non-avian avialans. We use this to propose a set of comparable, quantitative approaches to ascertain fossil bird diet and on this basis provide a consensus of what we currently know about fossil bird diet. While no single approach can precisely predict diet in birds, each can exclude some diets and narrow the dietary possibilities. We recommend combining [1] dental microwear, [2] landmark-based muscular reconstruction, [3] stable isotope geochemistry, [4] body mass estimations, [5] traditional and/or geometric morphometric analysis, and [6] finite element analysis to accurately reconstruct fossil bird diet. Our review provides specific methodologies to implement each approach and discusses complications future researchers should keep in mind. On this basis we report the current state of knowledge of non-avian avialan diet which remains very incomplete. The ancestral dietary condition in non-avian avialans remains unclear due to a scarcity of data and contradictory evidence in *Archaeopteryx*. Among early non-avian pygostylians, *Confuciusornis* has finite element analysis and mechanical advantage evidence pointing to herbivory, whilst *Sapeornis* only has mechanical advantage evidence indicating granivory, which agrees with fossilised ingested material known for this taxon. The enantiornithine ornithothoracine *Shenqiornis* has mechanical advantage and pedal morphometric evidence pointing to carnivory. In the hongshanornithid ornithuromorph *Hongshanornis*, only mechanical advantage evidence indicates granivory, but this is congruent with evidence of fossilised ingested material in this taxon. The same is true for the songlingornithid ornithuromorph *Yanornis* and its inferred carnivorous diet. Due to the sparsity of robust dietary assignments, no clear trends in non-avian avialan dietary evolution have yet emerged. Dietary diversity may seem to increase through time, but this is a preservational bias associated

with a predominance of data from the Early Cretaceous Jehol Lagerstätte. With this new framework and our current synthesis of current knowledge of non-avian non-avian diet, we expect dietary knowledge and evolutionary trends to become much clearer in the coming years, especially as fossils from other locations and climates are found. This will allow us to gain a deeper and more robust understanding of the role birds played in Mesozoic ecosystems and how this led to their pivotal role in modern ecosystems.

*Key words:* Avialae, birds, dental microwear, diet, dinosaurs, finite element analysis, fossil, mechanical advantage, morphometrics, stable isotopes

## Contents

I. Introduction .....	4
(1) Modern and Ancient Bird Diet .....	4
(2) Diet and Morphology in Aves .....	4
(3) Techniques for Determining Avian Diet .....	5
(4) Fossil Birds and the Focus of This Review .....	6
II. Direct Evidence .....	7
(1) Preserved Meals .....	7
(a) Introduction .....	7
(b) Avialan Consumulites .....	8
(2) Dental Wear.....	9
(a) Introduction .....	9
(b) Mesowear .....	9
(c) Microwear.....	10
(3) Dental Ultrastructure .....	12
(4) Soft Tissue.....	13
(a) Landmark-Based Cervical Reconstructions in Fossil Theropods.....	13
(5) Discussion .....	14
III. Stable Isotopes .....	15
(1) Introduction .....	15
(a) Carbon Isotopes .....	15
(b) Oxygen Isotopes .....	15
(c) Calcium Isotopes .....	16

(d) Hydrogen Isotopes.....	16
(e) Nitrogen Isotopes.....	16
(f) Sulphur Isotopes.....	16
(g) Heavy Metal Isotopes.....	16
(2) Stable Isotopes in Extant Birds .....	17
(3) Stable Isotopes in Fossil Theropods.....	17
(4) Discussion .....	17
IV. Body Mass .....	18
(1) Introduction .....	18
(2) Discussion .....	18
V. Physical Approaches .....	19
(1) Skull Reconstruction .....	19
(a) Existing Reconstructions .....	19
(b) Avenues for Improvement.....	20
(2) Morphometrics .....	21
(a) Introduction .....	21
(b) Traditional Morphometrics.....	22
(c) Geometric Morphometrics.....	26
(3) Functional Studies .....	30
(a) Introduction .....	30
(b) Lever Modelling .....	31
(c) Finite Element Analysis.....	34
(4) Complications Applicable across Physical Approaches.....	40
(a) Phylogenetic Signal .....	40
(b) Allometric Signal .....	41
(c) Many-To-One Mapping.....	41
(d) Liem's Paradox .....	41
(e) Integration and Modularity .....	42
(f) Behavioural Signals .....	42
(5) Discussion .....	44
VI. The Framework and Current Knowledge of Non-avian Avialan Diet.....	44
VII. Conclusions .....	47

VIII. References.....	47
Figure Captions.....	87
Tables.....	98

## I. Introduction

### (1) Modern and Ancient Bird Diet

Living birds [Aves: defined herein as crown birds (see Pittman, O'Connor, Field *et al.*, 2020a)] have been studied more than most any other organisms, and are at the forefront of human efforts to understand global ecology (Tietze, 2018). In large part this is because birds display some of the most varied diets in the animal kingdom. Many are familiar with their neighbourhood songbirds which feed on worms and seeds. However, birds are able to thrive in aquatic, terrestrial, and aerial environments around the world (Rico-Guevara, Sustaita, Gussekloo *et al.*, 2019) and consume nearly every source of nutrition imaginable therein. Rodents, fruit, fish, leaves, plankton, blood, beeswax, and organic mud are just a few of the food sources living birds may subsist off of (Lopes, Fernandes, Medeiros *et al.*, 2016). This rich diversity has also evolved in incredible ways. The ancestral avian has been proposed as an aquatic predator (Brusatte, O'Connor & Jarvis, 2015), granivore (Larson, Brown & Evans, 2016), or omnivore (Felice & Goswami, 2018) with a myriad of dietary radiations occurring during avian evolution including at least three origins of nectarivory, seven origins of aquatic predation, and 18 origins of frugivory (Felice, Tobias, Pigot *et al.*, 2019a).

Dietary diversification outside of Aves among the non-avian avialan birds [Avialae: defined herein as crown birds plus relatives as distant as *Archaeopteryx* (Gauthier, 1986; and see Pittman *et al.*, 2020a)] is much less well understood. Two species of early-diverging avialans preserve evidence of granivory, a single enantiornithine preserves an ingested invertebrate, and ten ornithuromorph species preserve evidence of granivory or piscivory (O'Connor, 2019). Beyond these, the diet of non-avian avialans is virtually unknown, and accordingly an ancestral avialan diet has not been proposed. It remains unclear if the vast dietary breath of living birds is unique or has deeper roots in the avialan tree, and if birds played the same unique ecological roles during the Late Jurassic and Cretaceous periods that they do in modern ecosystems.

### (2) Diet and Morphology in Aves

While avian diet itself has been well-reported, few correlations between diet and morphology are known in living birds, and fewer still from more than a single quantitative study. We provide a convenient glossary for the various descriptors of diet used in this review in Table 1. Invertivorous birds possess skulls with a low mechanical advantage (Corbin, Lowenberger & Gray, 2015; Olsen, 2017), while probing feeders [e.g. sandpipers (Pettigrew & Frost, 1985), ibises (Frederick & Bildstein, 1992), kiwis (Cunningham, Corfield, Iwaniuk *et al.*, 2013), and some songbirds (Adamík & Kornan, 2004; Lockie, 1956)] have particularly elongate rostra (Barbosa & Moreno, 1999; Kulemeyer, Asbahr, Gunz *et al.*, 2009). Granivorous birds tend to have ventrodorsally tall beaks (Soons, Herrel, Genbrugge *et al.*, 2010; van der Meij, 2004) exhibiting high mechanical advantage (Corbin *et al.*, 2015; Navalón, Bright, Marugán-Lobón *et al.*, 2018a) and a high strength (Soons, Genbrugge, Podos *et al.*, 2015; Soons *et al.*, 2010) [i.e. low peak Von Mises stresses when loaded (Dumont, Grosse & Slater, 2009)]. Raptorial birds

possess talons that, on average, are hypertrophied at digit I (Csermely, Rossi & Nasi, 2012; Fowler, Freedman & Scannella, 2009) and are more recurved (Csermely & Rossi, 2006; Tsang, Wilson, Ledogar *et al.*, 2019) than in non-raptorial birds. Among raptors, specialists in hunting other birds have longer toes and (Csermely *et al.*, 2006; Tsang *et al.*, 2019) and a wider skull (Hertel, 1995; Sun, Si, Wang *et al.*, 2018) [presumably for housing a larger cerebellum, the part of the brain that processes spatial orientation (Sun *et al.*, 2018)] while those that specialise in hunting fish tend to have all four talons enlarged (Einoder & Richardson, 2007; Fowler *et al.*, 2009). Scavenging raptors appear to be the most morphologically diagnostic group, characterised by large body size (Einoder *et al.*, 2007; Fowler *et al.*, 2009) and a narrow, shallow, (Bright, Marugán-Lobón, Cobbe *et al.*, 2016; Navalón *et al.*, 2018a; Sun *et al.*, 2018) and long (Hertel, 1995; Kulemeyer *et al.*, 2009; Pecsics, Laczi, Nagy *et al.*, 2019; Si, Dong, Ma *et al.*, 2015; Sun *et al.*, 2018) skull with a highly recurved rostrum (Hertel, 1995; Kulemeyer *et al.*, 2009).

Beyond these, diet/morphology correlations are at best known from a single study [e.g. small body size as indicative of nectarivory (Pigot, Sheard, Miller *et al.*, 2020)] and at worst contradicted between studies [e.g. bill curvature has been positively (Kulemeyer *et al.*, 2009), negatively (Navalón *et al.*, 2018a), or not (Barbosa *et al.*, 1999) correlated with probing behaviour]. Characters known from a single study may be awaiting corroboration by future studies, but contradictions between studies suggest that we do not fully understand some aspects of avian diet/morphology relations yet. If living birds are to be used as proxies for fossil birds, then, further work on understanding what they eat and why is imperative.

### (3) Techniques for Determining Avian Diet

When an ornithologist wants to know what a bird eats, the most straightforward technique is to directly observe them feeding and record what they ate (reviewed in Rosenberg & Cooper, 1990 alongside most of the following methods). Proxies for direct dietary observation include remote observation (detailed in Sullivan, Wood, Iliff *et al.*, 2009; Zhang, Rayner, Vickers *et al.*, 2019) and examination of faeces (critiqued in Carlisle & Holberton, 2006; augmented in Jarman, McInnes, Faux *et al.*, 2013; detailed in Ralph, Nagata & Ralph, 1985), pellets (critiqued in Votier, Bearhop, Ratcliffe *et al.*, 2001), uneaten prey remains near the nest (critiqued in Tornberg & Reif, 2007), or stomach contents [from dead birds (reviewed in Duffy & Jackson, 1986), those captured and forced to regurgitate (critiqued in Carlisle *et al.*, 2006; Gales, 1987), or collected natural non-pellet regurgitations (detailed in Oro, Ruiz, Jover *et al.*, 1997)]. These ‘direct evidence’ data provide an unambiguous association between an organism and a certain diet. Unfortunately, these techniques require an animal to be alive or recently deceased, and the closest fossil equivalents to these forms of direct evidence, preserved meals, rarely fossilise and are prone to a variety of preservational biases that makes them inadequate to reconstruct the diet of non-avian birds alone. Chemical analysis of stable isotopes in the soft tissues of living birds are commonly used to reconstruct trophic webs (detailed in Kloskowski, Trembaczowski & Filipiuk, 2019), and similar methods have been applied to bioapatite and amino acids preserved in fossil birds (see Section III.2 Stable Isotopes in Extant Birds), but the wide variety of factors controlling them leave them as more ambiguous lines of evidence.

While not used to determine diet *per se*, several physical approaches have been used to study modern birds in order to explain observed dietary trends and apply them to extinct organisms with unknown diets. Body mass has recently been found to predict a large amount of the variance in the diet of modern birds (Navalón *et al.*, 2018a; Pigot *et al.*, 2020), and so its estimation may represent a key metric

that has been relatively unexplored. With that said, most past studies have adopted ‘physical approaches’, those grounded in morphology and/or mechanics, to the study of avian diet. Dietary studies tend to focus on the skull, living birds’ most important tool for feeding (Rico-Guevara *et al.*, 2019), and the pes, their most important tool for manipulation of food prior to feeding (Clark, 1973; Sustaita, Pouydebat, Manzano *et al.*, 2013) and the primary tool for killing prey in raptorial birds (Fowler *et al.*, 2009). Traditional (detailed in Hertel, 1994) and geometric (detailed in Bright *et al.*, 2016) morphometrics both seek to quantify the shape of body parts, under the assumption that form will reflect function. Other studies investigate functional capacity directly. Lever model simplifications of skulls (detailed in Corbin *et al.*, 2015) describe the efficiency of force production and speed of the jaw while finite element analysis (detailed in Soons *et al.*, 2010) models the response of body parts to loading in order to compare their relative strength between organisms. Each of these techniques have been applied to living birds and non-avian theropods, but only the two forms of morphometrics have included non-avian avialan taxa. While physical approaches have some of the broadest applications to fossil organisms, they also introduce a variety of complications (see Section V.4 Complications Applicable across Physical Approaches).

There are also a select few lines of direct evidence that have been applied to fossil taxa but never to living and fossil birds. Dental wear analysis is commonly applied to living and fossil mammals (Green & Croft, 2018) with a recent application in theropods (Bestwick, Unwin, Butler *et al.*, 2018; Torices, Wilkinson, Arbour *et al.*, 2018). While inapplicable to avians in its current form because they are all toothless, it is of potential value in the study of toothed non-avian avialans. The same holds true for dentine ultrastructure analysis, which has only been applied to crocodylians and non-avian dinosaurs (Brink, Chen, Wu *et al.*, 2016). Ultrastructure of enamel has recently been investigated in toothed avialans and provides promising, if still preliminary, results (Li, Wang, Wang *et al.*, 2020). Finally, neck musculature has been proposed to inform food disassembly behaviours in non-avian theropods (Snively & Russell, 2007b), although the study of food disassembly in living birds has focused on the head and pes exclusively (Fowler *et al.*, 2009; Sustaita, 2008; Sustaita, Gloumakov, Tsang *et al.*, 2019). Each of these techniques is also addressed due to their potential application in fossil birds.

#### (4) Fossil Birds and the Focus of This Review

Avialans appear in the fossil record as early as the Late Jurassic, and by the Middle Cretaceous inhabited tropical to polar latitudes and were present on every continent (Pittman, O'Connor, Tse *et al.*, 2020b). While most Mesozoic birds are considered to fall somewhere along a continuum of arboreal and terrestrial lifestyles (Cobb & Sellers, 2020; Mayr, 2017; Serrano, Palmqvist, Chiappe *et al.*, 2017), *Ichthyornis* and Hesperornithiformes are undisputedly aquatic (Hinić-Frlog & Motani, 2010; Rees & Lindgren, 2005). Enantiornithes are the most widespread and speciose Mesozoic birds. They comprise roughly 60% of all non-avian avialan genera followed by non-avian ornithuromorph avialans (~25%) with the remainder made up by *Archaeopteryx*, Jeholornithiformes, Confuciusornithidae, Jinguoortidae, and *incertae sedis* taxa (Table 5 in Pittman *et al.*, 2020b; uncertainty based on their notes of taxa possibly referable to non-avian clades). Anchiornithinae and Scansoriopterygidae have been placed in Avialae previously, but their inclusion is controversial (Pittman *et al.*, 2020a). This review will err on the side of inclusivity and discuss these two clades as ‘avialans’ in addition to well-established Avialae [though personally we only see anchiornithines as avialans (see Pittman *et al.*, 2020a)].

A variety of dietary habits have been proposed for non-avian avialans. The vast majority are based on qualitative methodologies vulnerable to individual interpretation (Dumont, Tafforeau, Bertin *et*

*et al.*, 2016; O'Connor, Zhang, Chiappe *et al.*, 2013; Thulborn & Hamley, 1985; Zinoviev, 2009) or a few preserved meals which provide only a small glimpse of diet (O'Connor, 2019; O'Connor & Zhou, 2019). More comparable, quantitative approaches to fossil avialan diet have been made (Attard, Wilson, Worthy *et al.*, 2016; Navalón, 2014; Wang, Zhou, O'Connor *et al.*, 2014c) but these are few and far between. This review seeks to establish techniques that have proven effective at discriminating diet in living birds as well as fossil dinosaurs and to construct a framework for studying non-avian avialan diet. Parts of this framework not relying on teeth can also be applied to fossil avians, which have a narrower and better-constrained extant phylogenetic bracket. On this basis, we then present a consensus of what we currently know about non-avian avialan diet and how this can be improved upon moving forward. With this, future studies will make strides both in understanding Mesozoic ecosystems and tracing the evolution of one of the most important groups of living organisms.

## II. Direct Evidence

### (1) Preserved Meals

#### (a) Introduction

Preserved meals may take the form of food preserved in the digestive system (consumulites) or as closely-associated excretions (coprolites) or egestions (regurgitilites) (all *sensu* Hunt, Lucas, Milan *et al.*, 2012). Identification of consumulites may be problematic as accessing them often requires destroying the overlying remains, and misdiagnosis of ingested material may lead to erroneous dietary inferences (e.g. Nesbitt, Turner, Erickson *et al.*, 2006). Coprolites, while ostensibly simpler to analyse, are often taxonomically indeterminate (e.g. Chin, Tokaryk, Erickson *et al.*, 1998; Hollocher, Chin, Hollocher *et al.*, 2001; Hunt *et al.*, 2012; Qvarnström, Wernstrom, Piechowski *et al.*, 2019) and of questionable association with surrounding body fossils (e.g. Hunt *et al.*, 2012; James & Burney, 1997; Wang, Zhou & Sullivan, 2016c; Wood, Rawlence, Rogers *et al.*, 2008). Regurgitilites are sparse in the fossil record, possibly due to collection biases and/or misdiagnosis as coprolites (Myhrvold, 2012), but otherwise provide similar information to coprolites with similar referral issues. Hunt *et al.* (2012) provide a detailed review of the study of coprolites, and Myhrvold (2012) provides one for regurgitilites [which he refers to as emetolites]. There appears to be no comprehensive review of consumulites across fossil taxa other than the two discussed in the following section. Reports are typically centred on individual specimens. Smith and Scanferla (2016) provide an example of a single specimen with both strong and weak candidates for being true consumulites (lizard and insect respectively).

A preserved meal sheds light on only a single meal in an organism's life. Extant organisms are known to consume a wide variety of food (Cortés, 1997; Vitt & Pianka, 2005; Wilman, Belmaker, Simpson *et al.*, 2014), so any preserved meal should be viewed as a single data point in reconstructing diet. Taphonomic effects should also be taken into account, as meals with elements that fossilise easier will be more likely to be preserved inside of their consumer (O'Connor, 2019). As the remains of deceased individuals, the meals associated with a fossil may be the thing that killed them rather than a normal food source. This is more likely in the context of consumulites or minimally-processed regurgitilites than coprolites, given that coprolites require time and digestion to produce. In short, the possibility of a preserved meal being atypical cannot be ruled out by a single specimen.



## (b) Avialan Consumulites

O'Connor (2019) reviews the consumulites known from traditional Cretaceous avialans which consist of seeds, fish, and invertebrate exoskeletons. O'Connor and Zhou (2019) expand this review to cover all paravians, attributing lizard and fish consumulites to anchiornithines. A coprolite containing fish bones has been associated with a specimen of *Baptornis* (Martin & Tate, 1976), and an indistinct coprolite is known from a *Sapeornis* specimen (Zheng, Martin, Zhou *et al.*, 2011). Four specimens record evidence of avialans themselves as prey of other organisms [ichthyosaurs (Kear, Boles & Smith, 2003), non-avialan theropods (O'Connor, Zhou & Xu, 2011a; Xing, Bell, Persons *et al.*, 2012), and an indeterminate pellet-producing animal (Sanz, Chiappe, Fernández-Jalvo *et al.*, 2001)].

There is particular controversy around enantiornithine consumulites, which merits discussion. Only a single uncontroversial consumulite is known from an enantiornithine: small [ $<5$  mm] sections of crustacean exoskeleton in the abdomen of the holotype of *Eoalulavis hoyasi* (Sanz, Chiappe, Pérez-Moreno *et al.*, 1996). The fish bones associated with the holotype of *Piscivorenanthornis inusitatus* (Wang & Zhou, 2017c; Wang *et al.*, 2016c) and the amber in the abdomen of the holotype of *Enantiophoenix electrophyla* (Cau & Arduini, 2008; Dalla Vecchia & Chiappe, 2002) were both rejected as consumulites by O'Connor (2019). The former is interpreted as a fish coprolite (but see Xu, Zhou, Wang *et al.*, 2020) and the latter as elements reworked from surrounding soil (O'Connor, 2019). Spherical inclusions in two *incertae sedis* enantiornithines [and a specimen of *Jeholornis*] have been proposed as consumed plant matter (Mayr, 2016; Mayr & Manegold, 2013; O'Connor *et al.*, 2019), though other studies have suggested that these inclusions are fossilised ovarian follicles (Bailleul, O'Connor, Zhang *et al.*, 2019; O'Connor, 2019; O'Connor, Zheng, Wang *et al.*, 2014; O'Connor *et al.*, 2019; Wang, Wang, O'Connor *et al.*, 2016d; Zheng, O'Connor, Huchzermeyer *et al.*, 2013).

Consumulite preservation is known from at least: four specimens of *Jeholornis*, eight specimens of *Sapeornis*, and 14 specimens of ornithuromorph birds [*Eogranivora edentulata*, *Piscivoravis lii*, and 12 specimens of *Yanornis martini*] (Table 1 in O'Connor, 2019). In contrast, confuciusornithid and enantiornithine consumulites have not been found so far, so they have been proposed as being generally molliphagous [feeding on soft things; opposite of durophagous] (O'Connor, 2019). However, several other factors may be at play. Preservation of a consumulite is dependent on the meal being inside the animal's body [i.e. retained in the gut] and for parts hard enough to fossilise to not be dissolved at time of death. Thus, a longer gut retention time and a lower gut pH favour the preservation of consumulites. Gut retention times vary among modern birds due both to long-term lifestyle differences and short-term events. Lifestyle differences include locomotor habits (Barton & Houston, 1992; Caviedes-Vidal, McWhorter, Lavin *et al.*, 2007; Frei, Ortmann, Reutlinger *et al.*, 2014; Hilton, Houston, Barton *et al.*, 1999; Jackson, 1992) and nutrient density of diets (Hilton, Furness & Houston, 2000a; Hilton, Houston & Furness, 1998; Levey & Del Rio, 2001; McWhorter & Martínez del Rio, 2000). Events include coincidental dietary switching (Hilton, Furness & Houston, 2000b), migration (McWilliams, Caviedes-Vidal & Karasov, 1999), and rearing young (Thouzeau, Peters, Le Bohec *et al.*, 2004). Stomach acidities are also known to vary among modern raptors, with less bone remaining in the pellets of raptors with lower stomach pH (Duke, 1997). To our knowledge explanations of differing stomach pH have not been explored. Of these complicating factors, locomotor habits are of particular note as enantiornithines have been viewed as more arboreally inclined than contemporary ornithuromorphs (Field, Bercovici, Berv *et al.*, 2018a). In living birds, fully terrestrial species tend to have much longer gut retention times than flighted species (Frei *et al.*, 2014). Among flighted species, those that are more active in flight tend to

have shorter gut retention times (Caviedes-Vidal *et al.*, 2007; Hilton *et al.*, 1999; Jackson, 1992). If enantiornithines were more active fliers than contemporary ornithuromorphs they likely also had shorter gut retention times. This would contribute to a lower incidence of consumulite preservation. Other possibilities, such as diets with higher nutrient density or a generally higher stomach pH than contemporary ornithuromorphs, can only be tested by future discoveries of consumulites or potentially corroborated by the other methods explored below.

## (2) Dental Wear

### (a) Introduction

Teeth are dynamic systems worn continuously both by contact with ingested material and other occluding teeth (Green *et al.*, 2018). As such, the way teeth are worn directly reflects the diet of an animal. Dental wear analysis traditionally occurs at two distinct scales: mesowear, visible to the naked eye and reflecting periods of months to years (Green *et al.*, 2018); and microwear, visible only under magnification and reflecting the animal's final days before death (Ungar, 2019). Green and Croft (2018) provide a review of both scales while Ungar (2015; 2018; 2019) provides more in-depth reviews of microwear in particular. Studies of larger-scale wear [e.g. fracture and erasure of denticles] have been used to examine patterns of occlusion (Farlow & Brinkman, 1994; Lambe, 1917; Sankey, Brinkman, Guenther *et al.*, 2002), penetration angle (D'Amore, 2009; Farlow *et al.*, 1994), and potentially grooming (Currie & Evans, 2019) in dinosaurs. All of these studies have remained purely qualitative and uninterested in diet.

### (b) Mesowear

Dental mesowear has only been analysed in herbivorous mammals in order to distinguish between browsers and grazers (Green *et al.*, 2018). Its underlying principle is worth unpacking for potential broader use. Teeth experience two distinct types of mesowear: attrition, from contact with occluding teeth; and abrasion, from contact with ingested materials (Fortelius & Solounias, 2000). In mammalian herbivores, these sharpen and dull the teeth respectively (Fortelius *et al.*, 2000; Green *et al.*, 2018). Dental mesowear can be seen as the interaction between these two forces: in softer food diets attrition dominates and cusps are sharper, in tougher food diets abrasion dominates and cusps are rounder and are eventually completely flattened (Fortelius *et al.*, 2000; Green *et al.*, 2018; supporting online material of Muhlbachler, Rivals, Solounias *et al.*, 2011).

This approach is likely applicable to hadrosaurian dinosaurs whose dentition is reminiscent of equid mammals (Carrano, Janis & Sepkoski, 1999) and possibly to ceratopsian dinosaurs where attrition is believed to play a more complicated role (Erickson, Sidebottom, Kay *et al.*, 2015). Both possess teeth with tight occlusion upon which similar interactions between attrition and abrasion to modern herbivorous mammals may have occurred. The teeth of known theropods occlude only slightly, though more than the unoccluding teeth of extant saurians (Schubert & Ungar, 2005). Because of this, the interpretation of mesowear in theropods is obscure. Schubert and Ungar (2005) propose “wear facets” on tyrannosaurid teeth to be the product of attrition while “enamel spalling” is the product of abrasion; the former completely obliterates the latter over time. Candeiro *et al.* (2017) identify two additional attritional features [vertical and perpendicular attritional surfaces] and one additional abrasional feature [apical grooves]. They also broaden the phylogenetic bracket of these features to Theropoda. All of attritional/abrasional features may or may not overwrite one another based on chance, and so cannot be quantified as simply as the antagonistic mesowear seen in herbivorous mammals. Mesowear analysis as it currently exists, then, is inapplicable to the currently known fossil birds. Instead, a new system would

need to be constructed. Because theropod teeth occlude on only one surface [the lingual surface of upper teeth and labial surface of lower teeth], the difference in wear between the two surfaces may provide similar information to mesowear analysis. The mesowear paradigms of attrition and abrasion balance can, instead of being quantified from cusp shape, be quantified as a ratio between the number and depth of marks on occluding and non-occluding surfaces. Assuming Schubert and Ungar (2005) are correct and that attritional wear would overwrite abrasional wear, the non-occluding surface would provide a baseline for abrasion while the occluding surface would provide information on attrition. However, a lack of extant analogues with theropod-like occlusion renders the validity of such an approach dubious; digital or practical modelling of theropod jaws may provide a baseline for study.

### (c) Microwear

#### (i) Introduction

Dental microwear has been studied across vertebrates (Bestwick, Unwin & Purnell, 2019; Nevatte, Wueringer, Jacob *et al.*, 2017; Purnell, 1995; Purnell, Hart, Baines *et al.*, 2006; Ungar, 2019; Winkler, Schulz-Kornas, Kaiser *et al.*, 2019). Dental microwear describes the surface scarring of tooth enamel at a microscopic level, which can provide insight into the hardness [resistance to fracture] and toughness [resistance to tearing] of an animal's last meals, typically within the last few days of its life (Ungar, 2019).

Traditionally, microwear analysis involves directly counting surface features under light microscopy or from electron micrographs, with greater numbers of pits considered indicative of consuming harder foods while greater numbers of scratches indicative of consuming tougher foods (Ungar, 2019). This technique is inconsistent, with counting errors regularly reaching 10 percent among trained professionals (Grine, Ungar & Teaford, 2002; Mihlbachler, Beatty, Caldera-Siu *et al.*, 2012). In order to remove measurement noise, wear surfaces have more recently been imported as point clouds and analysed as fractal surfaces (Ungar, 2019) using techniques and software common in micro-scale manufacturing (e.g. Țălu, Stach, Méndez *et al.*, 2014). An increased area-scale fractal complexity is associated with harder foods in the diet, while an increased surface texture anisotropy is associated with tougher foods (Ungar, 2015). The only known source of error exclusive to fractal surface quantification is inter-microscope variability, which can be minimised by incorporating consistent automated treatments (Arman, Ungar, Brown *et al.*, 2016).

While microwear has been viewed as phylogenetically independent, this may be because it was studied among closely-related taxa. A recent study comparing disparate clades of herbivorous mammals found microwear to describe phylogeny better than diet, but to successfully distinguish diet within each phylogenetic group (Mihlbachler, Campbell, Ayoub *et al.*, 2016). It seems, then, that dental microwear comparisons should be restricted in taxonomic scope out of caution in order to avoid potential biases. Because the temporal coverage of dental microwear is so short, analyses also require a large sample size to properly encompass the full breadth of an animal's diet (Ungar, 2019). Green and Croft (2018) imply that a minimum of 10 individuals should be sampled. Individual tooth sets can still act as individual examples of diet similar to preserved meals, and differently worn teeth within a single jaw have been proposed to give insight into non-dietary behaviours such as grooming (see Section V.4.f.i Grooming).

#### (ii) *Rhamphotheca* Microwear

The possibility of applying techniques like those used in dental microwear to bird rhamphothecae [the horny covering of the beak] is interesting, but as of now uncertain in viability. Sload (2014) is the only

researcher to apply microwear techniques to structures other than teeth [claws of Florida stone crabs, *Menippe mercenaria*]. He notes that the lower hardness of the carapace [average microhardness of 1.33 GPa in melanised Florida stone crab carapace (Melnick, Chen & Mecholsky, 1996) versus 3.56 GPa in human enamel (Eimar, Ghadimi, Marelli *et al.*, 2012)] leads to atypical patterns of microwear, with many surfaces worn completely away (Sload, 2014 pg. 11). Reported averages for rhamphotheca hardness range from 1/4 [woodpecker (Lee, Horstemeyer, Rhee *et al.*, 2014)] to 1/11 [starling (Bonser & Witter, 1993)] that of Florida stone crab carapace, and so may experience even more extreme destruction of surface features. Rhamphotheca microwear is expected to reflect only very short periods of dietary input, requiring accordingly large sample sizes in order to acquire meaningful data about diet. However, hardness alone cannot predict wear resistance or patterns. Material behaviour and contact angle with abrading particles also play major roles (Zum Gahr, 1998). Enamel and crab carapace are both brittle ceramics for whom wear resistance increases near-linearly with hardness, but more flexible materials like metals and, potentially, keratin can display neutral or even negative correlation between hardness and wear resistance (Figure 4 in Zum Gahr, 1998). The combined differences in material type and contact angles of food particles [with a rhamphotheca likely straighter and smoother than any dental battery] means rhamphothecal microwear patterns will not resemble any known dental microwear patterns. A ground-up approach to the science will be necessary to make this application feasible: just as early mammal researchers identified scratches and pits to reflect tough and hard foods, diagnostic features of rhamphotheca wear have to be identified. The most straightforward method for identifying features would be laboratory experiments feeding birds known diets with differing mechanical properties. Such features can provide additional insight into living birds with obscure dietary habits as well as potentially being applied to fossilised rhamphothecae (see Section II.4 Soft Tissue).

### (iii) Microwear in Fossil Theropods

Owing to their lack of teeth, dental microwear studies have not been performed on any extant avians (see above). There are two studies of dental microwear in fossil theropods, both of which address diet. Candeiro *et al.* (2017) provide little detail of their methodology but stated that analysis was “undertaken with the support of a scanning electron microscope” (pg. 230). They observed the presence of an elongated groove worn into select teeth and proposed it as evidence of osteophagy (Candeiro *et al.*, 2017). Torices *et al.* (2018) combine qualitative analysis of dental microwear with finite element analysis (see Section V.3.c.iii Finite Element Analysis in Fossil Theropod Skulls). They found all theropod teeth from the area of study to be worn with only scratches and no pits (Torices *et al.*, 2018), indicating a diet of tough but soft material (Ungar, 2019). Torices *et al.* (2018) interpret this as a lack of bone-crushing behavior, where either flesh was removed selectively or prey was swallowed whole. They also cite the bimodal distribution of scratch directions as evidence of a puncture-and-pull feeding style. They propose scratches parallel to the tooth margin are formed while biting down while those oblique to the margin form when pulling back to disassemble prey (Torices *et al.*, 2018). Finally, they make commentary on possible omnivory in the troodontid *Troodon* based on finite element analysis. In their provided example micrographs, though, the scratches on the *Troodon* teeth appear longer, more numerous, and less parallel to the tooth margin than those in cf. *Pyroraptor* or *Gorgosaurus* (Figure 2 in Torices *et al.*, 2018). This may evince *Troodon* incorporating more abrasive foods into its diet than the contemporary theropods studied. If not an artefact of the small number of examples provided, quantification of microwear may provide additional insight into dietary differences among toothed theropods. Also of note is a conference abstract finding the dental microwear of *Archaeopteryx* to most closely resemble that of invertivorous

saurians (Bestwick *et al.*, 2018). These results are planned for full publication, but dietary conclusions for *Archaeopteryx* are tentative due to a small sample size (Jordan Bestwick pers. com. 2020).

#### (iv) Application to Fossil Avialans

Fractal quantification of microwear (Ungar, 2015) has not been applied to any theropod taxon, and may be of particular interest in testing the proposed durophagy of certain enantiornithines including *Shenqiornis* (O'Connor & Chiappe, 2011b; Wang, O'Connor, Zhao *et al.*, 2010b) and *Sulcavis* (O'Connor *et al.*, 2013). Successful application of the technique to fossil lepidosaurians of a similar size (Bestwick *et al.*, 2019) shows promise for success in toothed avialans. Should dental microwear of toothed avialans prove exclusively scratch-dominated as in non-avian theropods (Torices *et al.*, 2018), only durophagy can be effectively ruled out. Other techniques are necessary to further refine a dietary niche. It is worth noting that, while complications from swallowing prey whole have been raised (O'Connor *et al.*, 2019; Torices *et al.*, 2018), microwear in particular has been observed to reflect diet in extant lepidosaurs and archosaurs despite their limited use of the teeth in prey processing (Bestwick *et al.*, 2019; Winkler *et al.*, 2019). Purely tooth-based approaches avoid many of the issues of reconstructing skull material. However, the possibility of anterior or posterior rhamphothecae in toothed avialans (Wang, Stiegler, Wu *et al.*, 2020a) acting as an additional feeding surface may complicate conclusions drawn from dental analysis only.

Application of microwear to beaked fossil avialans is contingent on the preservation of the rhamphotheca and on validation studies in modern birds. Microwear represents only a short window of time in tooth enamel usage (Green *et al.*, 2018), and appears to turn over even faster in crab carapace (Sload, 2014). Thus rhamphothecae, which are even softer, will likely require large sample sizes for meaningful data. With less than a dozen rhamphothecae reported in the entire vertebrate fossil record (see Section II.4 Soft Tissue) this avenue requires the discovery of more specimens before it can be attempted. Procedures may also need to be devised to account for alterations to the microstructure of the keratin during burial, as in feathers (Figure 3 in Saitta, Kaye & Vinther, 2019).

### (3) Dental Ultrastructure

Brink *et al.* (2016) report that the tubule density of dentine in archosaur teeth [imaged via multiple harmonic generation microscopy] is able to discriminate between taxa proposed as hypercarnivorous and hyperherbivorous. Hypercarnivorous taxa, according to their study, possess a higher density of tubules within the dentine. While potentially promising in the future, the authors provide an incomplete explanation for the functional significance of this difference. The taxonomic breadth of the study leaves room for the differences observed to reflect phylogeny (as proposed by Wang, Song, Song *et al.*, 2015a) rather than function. Should future studies validate the dietary signal of this technique it would be applicable to any specimens in which dental wear analysis can be used.

Li *et al.* (2020) performed similar investigations into theropod enamel. They found loss of interglobular porous spaces and thinning of enamel at the avialan transition. These are proposed as reductions in tooth strength coincident with a dietary shift away from hard foods. However, while the authors propose interglobular porous spaces as restricting crack propagation, the mechanical differences between enamel and dentine are sufficient for this purpose in extant species (Bechtle, Fett, Rizzi *et al.*, 2010). Enamel thinning has alternatively been proposed as a byproduct of selection for rapid incubation (Yang & Sander, 2018). This does not mean that it could not have caused a dietary transition as another byproduct, and the thickening of enamel in enantiornithines proposed as durophagous (Li *et al.*, 2020 pg.

6) would seem to imply an ecological effect. This hypothesis will be worth examining against finite element models of non-avian avialans (see Section V.3.c Finite Element Analysis).

#### (4) Soft Tissue

While not direct evidence of diet, the preservation of muscular tissue can aid in determining the inputs for functional models of extinct animal feeding. Unfortunately, fossilised jaw musculature has only been reported in placoderm fish (Trinajstić, Marshall, Long *et al.*, 2007) and fossilised gular musculature in an ornithomimosaurian theropod (Briggs, Wilby, Perez-Moreno *et al.*, 1997). The more commonly preserved postcranial musculature (e.g. Dal Sasso & Signore, 1998; Kellner, 1996; Schultze, 1989) may become useful in reconstructing dietary habits as the alliance between cranial and postcranial systems in feeding becomes better understood (Montuelle & Kane, 2019). Similar can be said for body outlines (Wang, Pittman, Zheng *et al.*, 2017d) for corroborating landmark-based muscular reconstructions (see following section).

Fossilised rhamphothecae are invaluable when studying the diet of edentulous fossil taxa. Fossilised rhamphotheca impressions are known from a pterosaur (Frey, Martill & Buchy, 2003), hadrosaurid (Farke, Chok, Herrero *et al.*, 2013; Morris, 1970; Sternberg, 1935) and ceratopsian (Lingham-Soliar, 2008) ornithischians, and ornithomimosaurid (Barrett, 2005; Norell, Makovicky & Currie, 2001) and confuciusornithid (Chiappe & Meng, 2016 pg. 156; Falk, O'Connor, Wang *et al.*, 2019; Hou, Martin, Zhou *et al.*, 1999c; Miller, Pittman, Kaye *et al.*, In Press; Zhang, Zhou & Benton, 2008) theropods. While preservation of rhamphothecae appears to be rare from this small sample size, the fact that half of known confuciusornithid rhamphothecae are only visible with the use of UV or laser-stimulated fluorescence (LSF) imaging (Chiappe *et al.*, 2016 pg. 156; Falk *et al.*, 2019; Miller *et al.*, In Press) shows promise for modern imaging techniques revealing previously unknown rhamphothecae. These fossils allow construction of more accurate models of beaked organisms in the fossil record. They also narrow the phylogenetic bracket for studying taxa with rhamphothecae that were not preserved.

#### (a) Landmark-Based Cervical Reconstructions in Fossil Theropods

While the skull is often the first point of contact and/or the primary tool used in feeding, it functions only with the aid of postcranial systems (Montuelle *et al.*, 2019). Once the jaws have bitten down, the neck powers further disassembly of food by using the teeth and/or beak to tear material into a swallowable size [except in cases of chewing, which is not known in theropods (Zanno & Makovicky, 2011)]. In an animal that uses its neck for disassembly, selection is expected to favour an increase in size of those muscles that power disassembly. While not studied at length in living birds, comparisons of radiographs of *Gallus* and *Anas* appear to show greater muscle volume spanning the areas of most intense flexion during feeding (Figures 3 and 4 in van der Leeuw, Bout & Zweers, 2001). Thus, reconstruction of neck musculature can elucidate what way, if any, non-avian avialans disassembled their food before swallowing.

Among theropods, neck muscles have been reconstructed in ceratosaurids (Snively *et al.*, 2007b), allosaurids (Bakker, 1998; Snively, Cotton, Ridgely *et al.*, 2013; Snively *et al.*, 2007b), and tyrannosaurids (Bakker & Williams, 1988; Snively & Russell, 2007a; Snively *et al.*, 2007b; Tsuihiji, 2010) based primarily on the occipital region of the skull. The general consensus of these studies is that ceratosaurids and allosaurids could exhibit greater force in dorsiflexion while tyrannosaurids could exhibit greater lateroflexive force. This leads to reconstruction of ceratosaurids and allosaurids pulling their heads back to disassemble prey and tyrannosaurids shaking their heads side-to-side (Snively *et al.*, 2007b). Such reconstructions have yet to be attempted in paravian theropods.

#### (i) Application to Fossil Avialans

Tsuihiji (2005; 2007) compiled homologies of cervical muscles across extant diapsids, including crocodilians and birds. This provides an extant phylogenetic bracket for reconstruction of non-avian avialan cervical muscles, though typically two-dimensional fossil preservation may prove to be obstacle to the reconstruction process. Aside from the possibility of digital reconstruction (see Section V.1 Skull Reconstruction), only *Archaeopteryx* (Alonso, Milner, Ketcham *et al.*, 2004; Rauhut, 2014), *Neuquenornis* (Chiappe & Calvo, 1994), *Piscivorenanantiornis* (Wang *et al.*, 2017c), *Enaliornis* (Elzanowski & Galton, 1991), *Hesperornis* (Elzanowski, 1991), and *Ichthyornis* (Field, Hanson, Burnham *et al.*, 2018b) preserve the occipital region well enough to potentially identify muscular insertions. Of these, only *Neuquenornis*, *Hesperornis*, and *Ichthyornis* preserve any other regions of the skull. However, cervical muscles can be mapped on cervical vertebrae in lateral view (Snively *et al.*, 2007b; Tsuihiji, 2010), and the relative size of muscle insertions on the skull are consistent with those on the vertebrae (Snively *et al.*, 2007b). Muscles that in both crocodilians and birds contribute to dorsoventral flexion (e.g. *m. spinalis capitis*) and lateral flexion (e.g. *m. obliquus capitis*) can be identified from the cervical columns of avialan compression fossils preserved in lateral view [e.g. IVPP V13313 (Dalsätt, Zhou, Zhang *et al.*, 2006), IVPP V13558 (Zheng, O'Connor, Huchzermeyer *et al.*, 2014), STM 2-15 (O'Connor, Wang, Sullivan *et al.*, 2018), STM 29-11 (O'Connor, Wang, Zheng *et al.*, 2016c)]. Once identified, their relative areas can be compared following the methodology of Snively and Russell (2007b) to determine predispositions in cervical flexion and, in turn, methods of prey disassembly. We propose that prey disassembly method, in turn, can be extrapolated to inform the typical loading of the jaw. Their work implies that dorsiflexion shifts the muscular load vector cranially, ventroflexion shifts it rostrally, and lateroflexion shifts it laterally although they did not explicitly state this.

#### (5) Discussion

Lines of direct evidence are the most powerful and unequivocal data that we can obtain about avialan diet, but have a small scope for application. The study of dental mesowear and dentine ultrastructure are not recommended to investigate toothed avialan diet due to, respectively, the lack of occlusion in the clade and the lack of robust validation of dentine ultrastructure studies. While the information they provide is vital, the paucity of avialan consumulites in the fossil record prevents reliance on them for understanding diet in most specimens. Enamel ultrastructure has shown promising preliminary results, but physical approaches (Section V) are necessary to corroborate their findings. The lack of fossilised avialan musculature means reconstructions must currently rely on landmarks for their attachment. Landmarks on the cervical vertebrae in particular can inform habits of prey disassembly by presenting adaptations for cervical flexion (see also Section V.3.b.iv Discussion for notes on skull muscle reconstruction) in both toothed and beaked avialans. The most promising line of direct evidence of avialan diet is dental microwear. In particular, its utility in detecting the input of hard foods into animals' diets makes it ideal for investigations of possible durophagy in enantiornithines. The main drawback of this technique is the requirement of a large sample size, and so referral of unidentified specimens to known taxa may be necessary before such studies can be undertaken at phylogenetically meaningful levels. Rhamphotheca microwear requires a foundation in extant taxa before any application to beaked avialans can be attempted.

### III. Stable Isotopes

#### (1) Introduction

Natural abundances of stable isotopes [i.e. those not known to radioactively decay] vary both geographically and by the way they are preferentially incorporated into biomolecules. With knowledge of these variations observed in living communities, the abundance of stable isotopes in the tissues of extinct animals can be used to reconstruct various aspects of palaeobiology (Clementz, 2012). The preservation window of stable isotope systems cited in all publications post-2010 can be traced to those listed in Table 5.2 of Koch's (2007) review of the stable isotope chemistry in fossil vertebrates. Koch, however, does little to justify the ranges of these windows. He justifies the preservation window of bone, enamel, and soft tissues, but the provided temporal limits of individual isotope systems in these tissues are not justified. As such, the exclusion of an isotope system from analysis because the specimen is 'too old' is unfounded. Therefore, in addition to the traditional systems of carbon, oxygen and calcium isotopes commonly analysed in Mesozoic enamel and bone [bioapatite], we will also address hydrogen, nitrogen, and sulfur systems found in collagen [convincing evidence of preservation dating to the Early Jurassic (Lee, Chiang, Huang *et al.*, 2017)] as well as heavy metal [strontium, neodymium, lead, iron, copper, magnesium, and zinc] systems found in bioapatite. Keratin can theoretically preserve all isotope systems collagen does (Koch, 2007), but recent taphonomic evidence finds keratin chemistry to be lost too quickly for fossilisation (Saitta *et al.*, 2019; contra Schweitzer, Zheng, Moyer *et al.*, 2018).

#### (a) Carbon Isotopes

$^{13}\text{C}$  is enriched relative to  $^{12}\text{C}$  in plants utilising a C4 photosynthetic pathway relative to those using the C3 pathway (Park & Epstein, 1960). The isotopic ratio of the carbon contained in the  $\text{CO}_3$  components of bioapatite and within the amino acids of collagen can be used to determine what photosynthetic source(s) the nutrition of the animal in question ultimately came from. DNA and palynological evidence agree on an Oligocene origin of the C4 pathway (Sage, Sage & Kocacinar, 2012), and so this aspect of the isotope system is uninformative of diet in specimens older than roughly 30 Ma.

Marine ecosystems are known to be enriched in  $^{13}\text{C}$  relative to terrestrial ecosystems, though upper extremes of terrestrial species tend to overlap with those of marine species (Schoeninger & DeNiro, 1984) likely due to terrestrial input from C4 plants. Thus, prior to the emergence of C4 plants, we may expect a more bimodal distribution of  $^{13}\text{C}$  enrichment, with high enrichment of  $^{13}\text{C}$  indicating marine input into an organism's diet.

#### (b) Oxygen Isotopes

$^{18}\text{O}$  is enriched relative to  $^{16}\text{O}$  in the leaves of plants relative to their other tissues, with increasing enrichment the higher those leaves are in the canopy (Koch, 2007). So, an enrichment in  $^{18}\text{O}$  in the  $\text{CO}_3$  and  $\text{PO}_4$  of bioapatite or the amino acids of collagen may indicate a higher proportion of leaves in the diet or feeding on leaves higher in the canopy. While not directly indicative of foraging height,  $^{18}\text{O}$  could potentially be used as a proxy. However, this enrichment must be evaluated based on comparison with specimens from the same locality, as atmospheric temperature and water temperature are stronger controls on the enrichment of  $^{18}\text{O}$ .

More frequently,  $^{18}\text{O}$  enrichment is used to discern metabolic activity. Body fluids are known to fractionate  $^{18}\text{O}$  with temperature (Koch, 2007). Thus comparisons of the  $^{18}\text{O}$  enrichment between different bones within an organism (Barrick & Showers, 1994; Barrick, Showers & Fischer, 1996; but see



Kolodny, Luz, Sander *et al.*, 1996) and comparing global trends of  $^{18}\text{O}$  enrichment to that of known ectotherms (Amiot, Lécuyer, Buffetaut *et al.*, 2006; Fricke & Rogers, 2000) can both yield evidence of homeothermy. If used in dietary reconstruction, then, both locality and metabolic rate must be kept constant when making comparisons.

#### (c) Calcium Isotopes

$^{44}\text{Ca}$  is known to deplete relative to all other calcium isotopes with increasing trophic level (Clementz, 2012), and so ratios of  $^{44}\text{Ca}/\text{Ca}$  in bioapatite across a locality may allow for rough approximation of the trophic pyramid. However, carnivorous taxa that do not consume the mineralised tissues of prey [e.g. early hominids] appear to be at a lower trophic level from these ratios (Reynard, Henderson & Hedges, 2010). Osteophagous herbivorous taxa (Esque & Peters, 1994; Hutson, Burke & Haynes, 2013) will presumably appear to be at a higher trophic level.

#### (d) Hydrogen Isotopes

$^2\text{H}$  is enriched relative to  $^1\text{H}$  with similar trends in plant tissues to  $^{18}\text{O}/^{16}\text{O}$ , but with less contribution from evaporative conditions and greater contribution from differences in plant tissues (Koch, 2007). If collagen can be recovered, then this system appears more appropriate than oxygen for comparing between different localities and between organisms with differing metabolic rates.

#### (e) Nitrogen Isotopes

$^{15}\text{N}$  is enriched relative to  $^{14}\text{N}$  with increasing trophic level, and is used widely in analyses of modern food webs (e.g. Davenport & Bax, 2002; Gu, Schelske & Hoyer, 1996; Rau, Ainley, Bengtson *et al.*, 1992). The baseline enrichment of  $^{15}\text{N}$  varies based on locality, and so comparisons must be made within a given locality (Koch, 2007). Nitrogen is sourced in the amino acids of an organism which are more difficult to avoid consuming than bone, and so if collagen can be recovered nitrogen isotopes may provide a superior reflection of trophic level to calcium isotopes.

$^{15}\text{N}$  is also known to be enriched in marine ecosystems relative to terrestrial ecosystems, and with significant separation in levels between the two [except in reef fish] (Schoeninger *et al.*, 1984). As such, an ecosystem with organisms feeding on exclusively marine or terrestrial organisms should display a bimodal distribution and not affect signal for trophic level, but organisms taking from both sources may muddy the waters.

#### (f) Sulphur Isotopes

Sulphur isotopes are known to vary between plants in modern ecosystems, but not in any predictable manner (Connolly, Guest, Melville *et al.*, 2004; Koch, 2007). Sulphur extracted from collagen may provide evidence of consumers having different producers contribute to their diet. Without fossilised plant proteins, which taphonomic studies rule unlikely (Fogel & Tuross, 1999), greater precision appears impossible.

#### (g) Heavy Metal Isotopes

$^{87}\text{Sr}/^{86}\text{Sr}$  (Koch, 2007),  $^{144}\text{Nd}/^{143}\text{Nd}$  (van de Flierdt, Robinson, Adkins *et al.*, 2006), and  $^{207}\text{Pb}/^{206}\text{Pb}$  (Scheuhammer & Templeton, 1998) are not known to fractionate in biological systems, and so are typically used as indicators of location. In fossil terms, differences in these systems in bioapatite of organisms from a given locality would represent different migratory patterns of said organisms in life.

In extant mammals,  $^{56}\text{Fe}/^{54}\text{Fe}$  is higher in females than males while the reverse is true for  $^{65}\text{Cu}/^{63}\text{Cu}$  (Jaouen, Balter, Herrscher *et al.*, 2012; Martin, Tacail & Balter, 2017). While potentially useful for identifying sexual dimorphism from bioapatite, the trend is hypothesised to be linked to menstrual cycles (Martin *et al.*, 2017) and thus may not be applicable outside of Eutheria.

Enrichment of both  $^{26}\text{Mg}$  and  $^{66}\text{Zn}$  in bioapatite with increasing trophic level have also been reported, but these isotopes appear more vulnerable to small regional variations than calcium or nitrogen (Martin *et al.*, 2017). They may prove to be effective as secondary systems used to confirm predictions based on calcium and/or nitrogen.

## (2) Stable Isotopes in Extant Birds

The first record of stable isotopes analysed in extant birds comes from the study across vertebrates of Schoeninger and DeNiro (1984), followed by Hobson (1987) as the first to focus specifically on birds. Both studies focused on determining marine or terrestrial input to the diet via  $^{13}\text{C}$  and/or  $^{15}\text{N}$ . Hobson (1990) was the first to apply  $^{13}\text{C}$  towards determining trophic level and proposed  $^{15}\text{N}$  as a superior alternative. His following paper (Hobson, 1993) codified the role of  $^{15}\text{N}$  in determining trophic level and serves as a basis for all subsequent avian studies.  $^{18}\text{O}$  (Farmer, Rye, Landis *et al.*, 2003; Hobson, Bowen, Wassenaar *et al.*, 2004),  $^2\text{H}$  (Chamberlain, Blum, Holmes *et al.*, 1996; Farmer *et al.*, 2003; Hobson *et al.*, 2004; Lott, Meehan & Heath, 2003; Norris, Marra, Bowen *et al.*, 2006),  $^{34}\text{S}$  (Farmer *et al.*, 2003; Lott *et al.*, 2003; Sanpera, Ruiz, Moreno *et al.*, 2007),  $^{87}\text{Sr}$  (Blum, Taliaferro & Holmes, 2001; Chamberlain *et al.*, 1996), and  $^{207}\text{Pb}$  (Scheuhammer, Bond, Burgess *et al.*, 2003; Scheuhammer *et al.*, 1998; Svanberg, Mateo, Hillström *et al.*, 2006) have all only been used as indicators of locality in extant birds, typically by comparing sets of feathers from known localities to determine an isotopic signature without regard to the ecological drivers described above. To our knowledge no studies of calcium, neodymium, iron, copper, magnesium, or zinc stable isotopes in birds have been performed.

## (3) Stable Isotopes in Fossil Theropods

Among fossil avians, stable isotopes have been used in dietary reconstructions in recent (Hobson & Montevecchi, 1991; Miller, Fogel, Magee *et al.*, 2005) and Palaeocene-Eocene (Angst, Amiot, Buffetaut *et al.*, 2015; Angst, Lécuyer, Amiot *et al.*, 2014) taxa. Dietary studies older than this are restricted to non-paravian dinosaurs, primarily analyses of  $^{13}\text{C}$  determining ecosystems were based on C3 plants (Amiot, Wang, Lécuyer *et al.*, 2010; Amiot, Wang, Zhou *et al.*, 2015; Ghosh, Bhattacharya, Sahni *et al.*, 2003; Montanari & Norell, 2011; van Baal, Janssen, van der Lubbe *et al.*, 2013). The one exception is the work of Ostrom *et al.* (1993) who tabulated  $^{15}\text{N}$  enrichment in bulk bones and teeth of vertebrates from the Judith River Formation, and recovered expected trends of greater enrichment in proposed hypercarnivorous taxa [e.g. tyrannosaurs, plesiosaurs] than proposed herbivorous/omnivorous taxa [e.g. hadrosaur, sturgeon]. No additional attempts to analyse nitrogen stable isotopes of any Mesozoic vertebrate have been made, nor have stable isotopes of any paravian dinosaur tissue been analysed.

## (4) Discussion

Stable isotopes promise information with validity akin to that of direct evidence, but the variety of influences on their ratios complicates the signals they provide. As such, the range of tissues and elements used in dietary inference is inherently small. Any attempts at bioapatite stable isotope analysis should be performed on teeth rather than bone if possible to ensure the highest level of accuracy (Hollund, Arts, Jans *et al.*, 2015). The only locality with a large number of published avialan teeth is the Jehol Group

(Chiappe *et al.*, 2016; O'Connor *et al.*, 2011b). All evidence points to the Jehol Group being entirely terrestrial (Zhou, Barrett & Hilton, 2003a) so distinction of marine and terrestrial producer input via carbon isotopes is irrelevant. Use of oxygen isotopes to determine foraging height may be effective in the earliest-diverging avialans. However the variability of histological character in enantiornithines (Cabrera-Moo, Buscalioni, Cubo *et al.*, 2006; O'Connor, Wang, Zheng *et al.*, 2014) and early-diverging ornithuromorphs (Wang, Hao, Kundrát *et al.*, 2019a) would predict variation in metabolic rates acting as confounding factors. The only isotope systems of interest in all avialan bioapatite, then, would be calcium isotopes with potential secondary confirmation with magnesium and zinc. While calcium isotopes could reliably be recovered, the aforementioned exceptions to their trends make them less desirable than nitrogen. Reconstructing trophic levels via nitrogen isotopes after Ostrom *et al.* (1993) may be invaluable in future palaeoecological studies, but because no replication of this study has ever been published there is risk that specimens will be destroyed without yielding results. We recommend studies on fossils that are incomplete or otherwise of low scientific value to confirm nitrogen isotope trends can be consistently reproduced in Mesozoic food webs. In lieu of such a study, a comparison of calcium isotope ratios across a given fossil site will provide the most reliable insight into avialan diet. Evidence of paravian theropods swallowing prey whole (O'Connor *et al.*, 2019) is encouraging for such a study, as swallowing whole requires consumption of mineralised tissues by predators.

## IV. Body Mass

### (1) Introduction

While not traditionally used to determine diet in fossil organisms, recent studies of modern birds (Bright *et al.*, 2016; Navalón *et al.*, 2018a; Pigot *et al.*, 2020) have found body mass to explain more of the variance in diet than physical approaches. Invertebrate feeders tend to be smaller than those that scavenge or hunt vertebrates [separation near 300 g] (Navalón *et al.*, 2018a), and among raptorial birds scavengers are distinctly larger than active hunters or omnivores (Bright *et al.*, 2016). It is worth noting that this trend appears to apply only to feeding on animals, as herbivorous diet types are spread across the range of measured body masses (Figure 6 in Navalón *et al.*, 2018a). However, body mass alone can only consistently predict nectarivory in extant birds (Pigot *et al.*, 2020). Body mass has a major influence on feeding strategy in extant birds, but can be used only as a component in analysis [e.g. coupled with TM as in (Pigot *et al.*, 2020)] or as a secondary determinant. For instance, if other methods within the framework provide evidence of general carnivory mass may help specify prey to vertebrates or invertebrates.

### (2) Discussion

Body mass is a universal metric among animals, but the reasons behind its observed effects on bird diet remain unclear and thus its application in fossil avialans is questionable. The proposed reason for size having such influence in raptorial birds is tight integration of the rostrum and cranium, disallowing significant change of one without the other (Bright *et al.*, 2016). If this is true, then this means that the control of size on diet is developmental, not mechanical. Bird skulls have undergone extreme changes from the early-diverging avialan condition thought to be brought about by radical shifts in developmental controls (Bhullar, Hanson, Fabbri *et al.*, 2016). Certain features of non-ornithuromorph skulls [small premaxilla [except in confuciusornithids], large maxilla [except in confuciusornithids], robust nasal and lacrimal, prominent postorbital (Hu, O'Connor, McDonald *et al.*, 2020a; O'Connor *et al.*, 2011b; Rauhut, 2014)] more strongly resemble that of early-diverging theropods than extant birds. As such, developmental constraints on extant bird skulls are unlikely to be at play in groups diverging earlier than

Ornithuromorpha. However, size explaining the largest portion of dietary variance persists across extant birds (Navalón *et al.*, 2018a; Pigot *et al.*, 2020) despite differences in modularity between avian clades and a general decoupling of the rostrum and braincase shape across living birds as a whole (Felice *et al.*, 2018). Diet, then, may be under mechanical control rather than developmental control. For instance, invertebrate taxa tend to be smaller than vertebrate taxa, so larger birds are less likely to feed on them. With that assumption, it is recommended that mass calculations be factored into dietary reconstructions of fossil birds if possible.

Serrano *et al.* (2015) provide mass estimates of 43 Mesozoic birds based on extant bird skeletons (see their Table 8). Table 2 expands on their work, providing mass estimates based on photographs in the literature for 71 additional specimens of non-avian avialans representing 61 species. Combined with the estimates of (Serrano *et al.*, 2015), ~65% of non-avian avialan species likely fell below the 300g dietary transition observed by (Navalón *et al.*, 2018a). However, this is largely driven by ornithothoracine species. Among non-ornithothoracine taxa only three specimens of *Archaeopteryx* have an estimated mass range below 300g (Table 8 in Serrano *et al.*, 2015). There may be some taphonomic bias against preservation of large ornithothoracines, as the largest enantiornithine (Atterholt, Hutchison & O'Connor, 2018) and non-avian ornithuromorph (Buffetaut & Angst, 2016) taxa are only known from highly fragmentary material. Of taxa known from more complete material, it appears predatory ornithothoracines would be more likely to prey on invertebrates. Conversely, non-ornithothoracine avialans lacking distinct evidence of herbivory [*Archaeopteryx* and Confuciusornithidae] are more likely to have taken vertebrate prey. Some ornithothoracine groups [Bohaiornithidae, Pengornithidae, Songlingornithidae] also tend to have body masses above 300g (Table 1) and may represent a secondary adaptation to take vertebrate prey. 13 specimens of *Yanornis* [a songlingornithid] preserving ingested fish (O'Connor, 2019) support this premise.

## V. Physical Approaches

### (1) Skull Reconstruction

#### (a) Existing Reconstructions

All of the physical approaches described below, when applied to skulls, require reconstruction of the skull. A listing of all reported avialan skulls is provided in Table 3. No complete avialan skull is preserved in three dimensions. A composite reconstruction of the *Ichthyornis dispar* skull has been constructed in 3D (Field *et al.*, 2018b). A team has also assembled a full 3D model of *Archaeopteryx*, but it is unreleased and intended for public education so its accuracy is unclear (Carney, Kaplan, Kirk *et al.*, 2018). All other non-avian avialan skull reconstructions to date are 2D, owing to the flattened preservation of most avialans.

Over a dozen 2D skull reconstructions of *Archaeopteryx* exist (Elzanowski, 2001b; Rauhut, 2014). These tend to agree with one another, differing mostly in how bones contact at the antorbital fenestra and in the dorsocranial region. The same is true of *Hesperornis*, with some variation in the structure of the orbit (compare Bühler, Martin & Witmer, 1988; Gingerich, 1973) and the addition of the predentary to later restorations (Martin & Naples, 2008). Reconstructions of *Sapeornis* are less consistent, with the skull generally seen as more robust with a more downturned rostrum and mandible over time (compare Hu *et al.*, 2020a; Zhou & Zhang, 2003b). Skull reconstructions of *Confuciusornis* (Chiappe, Shu'an, Qiang *et al.*, 1999; Elzanowski, Peters & Mayr, 2018; Navalón, 2014; Zhou & Hou, 2002a) have

no clear trend in their variation. Most differences between reconstructions are in the length of the rostrum and height of the frontal, possibly representing intraspecific variation. *Anchiornis* sees the most variability in skull reconstruction, with noticeable differences in the size of fenestrae, shape of the mandible, placement of sutures in all reconstructions (Wang *et al.*, 2017d; Xu, You, Du *et al.*, 2011; Xu, Zhou, Dudley *et al.*, 2014). In each case the skull was merely illustrative with no record of reconstruction methods, and so a more intentional reconstruction of the skull of *Anchiornis* is necessary. Quite a few avialans have had only a single skull reconstruction produced. These include *Yi* (Xu, Zheng, Sullivan *et al.*, 2015) [probably a non-avian pennaraptoran (Pittman *et al.*, 2020a)], *Xiaotingia* (Xu *et al.*, 2011), *Jeholornis* (O'Connor, Wang, Sullivan *et al.*, 2013), *Gobipteryx* (Elzanowski, 1977), *Cathayornis* (Martin & Zhou, 1997), *Eoenantiornis* (Hou, D Martin, Zhou *et al.*, 1999b), *Shenqiornis*, *Rapaxavis*, *Pengornis* (O'Connor *et al.*, 2011b), *Piscivorenanantiornis* (Wang *et al.*, 2016c), an indeterminate enantiornithine hatchling (Sanz, Chiappe, Pérez-Moreno *et al.*, 1997), *Patagopteryx* (Chiappe, 2002), *Yanornis* (Huang, Wang, Hu *et al.*, 2016), and *Yixianornis* (Clarke, Zhou & Zhang, 2006).

### (b) Avenues for Improvement

Due to most avialan specimens being compression fossils, relatively little material lends itself to construction of 3D models. A 3D reconstruction of *Parahesperornis*, or at least a general hesperornithiform, should be possible with reported material. A nearly complete 3D skull of *Parahesperornis* is known, alongside well-preserved 3D pieces of the skull of *Hesperornis* (Bell & Chiappe, 2020) and *Pasquiaornis* (Sanchez, 2010). Excellent skulls of *Gobipteryx minuta* (Chiappe, Norell & Clark, 2001; Elzanowski, 1974; Elzanowski, 1977) are nearly complete and preserved in three dimensions, and represent the most complete picture of the enantiornithine skull. The skulls of *Gobipipus reshetovi* (Kurochkin *et al.*, 2013) and an unnamed gobipterygid (Lu, Xu, Zhang *et al.*, 2011) are more fragmentary, but are similar enough to *Gobipteryx* that complete reconstruction is possible. Unfortunately, the skulls of these taxa are highly derived (Hu, Sansalone, Wroe *et al.*, 2019) and are of dubious use in reconstructing the skulls of other enantiornithines (O'Connor *et al.*, 2011b). One complete enantiornithine skull is preserved in amber (Xing *et al.*, 2017), but is of an extremely early ontogenetic stage, damaged from preparation, and extremely difficult to image (Xing *et al.*, 2017 pg. 266). The remainder of three-dimensionally preserved non-avian avialan skull material (Table 2) is too fragmentary for reconstruction. While an avian, the holotype of the Mesozoic bird *Asteriornis* includes a well-preserved and nearly complete skull (Field, Benito, Chen *et al.*, 2020) which can also offer insight into avian ecosystems in the Late Cretaceous.

As the vast majority of avialan skulls are preserved two-dimensionally (Table 2), two-dimensional reconstructions will likely remain standard in the coming years. As such, valid approaches to 2D reconstruction are imperative for applying physical approaches to avialans. In an ideal scenario of a near-perfect skull (Sanz *et al.*, 1997) reconstruction is often just a matter of retrodeformation. When no one skull can supply all the information needed, multiple skulls must be combined into a chimera. O'Connor and Chiappe (2011b) set the standard for enantiornithines by restricting the phylogenetic bracket to only other members of the clade. Within Enantiornithes, however, there is still ample variation in skull morphology (Morschhauser, Varricchio, Chunling *et al.*, 2009; O'Connor *et al.*, 2013; Zhang, Ericson & Zhou, 2004). We therefore recommend a refinement of current 2D avialan reconstructions by narrowing the phylogenetic bracket of reconstructions further. Four distinct families are generally recognised within Enantiornithes [Avisauridae (Atterholt *et al.*, 2018; Brett-Surman & Paul, 1985), Bohaiornithidae (O'Connor, 2019; Shi & Li, 2019; Wang, Zhou, O'Connor *et al.*, 2014c; *contra* Chiappe,

Qingjin, Serrano *et al.*, 2019b) , Longipterygidae (O'Connor, 2019; O'Connor, Wang, Chiappe *et al.*, 2009; Pittman *et al.*, 2020a; Shi *et al.*, 2019), and Pengornithidae (O'Connor, 2019; Pittman *et al.*, 2020a; Wang, O'Connor, Zheng *et al.*, 2014e)], and so missing information can be preferentially filled by members of the family. Similar is true for the three widely recognised families of early-diverging ornithuromorphs [Hesperornithiformes (Bell & Chiappe, 2016; Clarke, 2004), Hongshanornithidae (O'Connor, Gao & Chiappe, 2010b; Pittman *et al.*, 2020a; Wang, Zheng, O'Connor *et al.*, 2015b), and Songlingornithidae (Clarke *et al.*, 2006; Hou, 1997; Pittman *et al.*, 2020a)]. In addition, the level of error introduced by chimerisation can be estimated by creating similar chimeras of modern bird skulls and comparing results of individuals to that of the composite. Skulls or individual teeth of lizards can help estimate the effects of chimeric dentition for toothed avialans. This will provide information on the accuracy of any analyses performed on inevitably chimeric avialan skull reconstructions.

Alternatively, technological advancements may allow for 3D reconstruction from 2D sources. 3D images of specimens can be taken using computed tomography [CT] (Abel, Laurini & Richter, 2012) or augmented laminography [AL] (Zuber, Laaß, Hamann *et al.*, 2017). Subsequently, broken parts can be segmented and reconstructed according to techniques described by Lautenschlager (2016). Unfortunately, in addition to the man hours necessary to segment and manipulate the numerous shards of a shattered skull, there are significant obstacles to collecting these data. Typically, avialan skulls are part of slab specimens. These slabs do not yield good results when imaged with standard scanners on site and are too large for higher-resolution scanners (Michael Pittman, pers. obs.). Preparation can resolve the size issue, but most museums are hesitant to approve preparation work, especially as slab specimens are generally beautifully articulated (Michael Pittman, pers. obs.). This means that to obtain CT data, whole slabs must be sent to far away specialist scanners [large chamber  $\mu$ CT scanner and synchrotron scanner] that involves time-consuming permit- and logistics-related paperwork, high transports cost [usually personal courier via air travel] and an elevated risk of damage to the specimen during its transport. Thus, 3D data has been difficult to collect from avialan skulls and will continue to be a challenge whilst these obstacles remain. The 2D approach described above is recommended to circumvent these issues and preserve fossils for future study, though fossils whose skulls are already disarticulated [e.g. *Longusunguis* (Hu, O'Connor, Wang *et al.*, 2020b; Wang *et al.*, 2014c) or *Eogranivora* (Zheng, O'Connor, Wang *et al.*, 2018a)] (see also Table 3) would be ideal for pilot studies of 3D reconstruction of avialan skulls.

## (2) Morphometrics

### (a) Introduction

Morphometrics is the study of shape and the quantification of shape change, typically in a biological context (Rohlf, 1990). Shapes are quantified by defining landmarks, “(1) homologous anatomical loci that (2) do not alter their topological positions relative to other landmarks, (3) provide adequate coverage of the morphology, [and] (4) can be found repeatedly and reliably...” (Zelditch, Swiderski, Sheets *et al.*, 2004 pg. 24). In studies that are concerned with only two dimensions of geometry, all landmarks must also lie in the same plane (Zelditch *et al.*, 2004 pg. 24). Once landmarks are placed on the structure in question, the methods in which they are analysed differentiate traditional morphometrics [TM], based on pre-selected distances and angles between landmarks (Marcus, 1990), and geometric morphometrics [GM], where the differences between landmark position in all models is quantified and significant variables are identified *a posteriori* (Zelditch *et al.*, 2004 pg. 24). Guillerme *et al.* (2020) review the collection, analysis, and interpretation of all morphometric data.

## (b) Traditional Morphometrics

### (i) Introduction

Comparing linear measurements of animals likely predates the formalisation of science itself. Traditional morphometrics [TM] are differentiated by rigorous statistical considerations of measurements taken, and selection of measurements believed to be relevant to the topic of study. The full range of statistical techniques used in TM are detailed by Marcus (1990).

The traditional appeal of TM has been the low computing power required. Measurements can be taken by hand and compiled into a spreadsheet for analysis of relatively low complexity. Now that computing power is no longer a limiting factor, the main appeals of TM are the lower investment of time and money into the project. TM does not necessitate creating two- or three-dimensional models nor placement of landmarks and semi-landmarks onto models (Rohlf, 1990). TM also allows for *a priori* selection of measures believed to have functional significance [though this may lead to arguments of cherry picking or p-hacking (e.g. Warton & Hui, 2011)]. TM is most recently utilised in studies of diet with positive results (Fraser & Theodor, 2011; Hertel, 1994; Hertel, 1995; McBrayer & Corbin, 2007; Surkov & Benton, 2008).

### (ii) Traditional Morphometrics in Extant Bird Skulls

Traditional morphometric approaches to understanding the avian skull trace their roots to the descriptive measurements of Shufeldt (1909 pg. 88, 181). The following decades were characterised by numerous univariate comparisons of skull measurements (e.g. Schoener, 1965). Baker and Moeed (1979) were the first to include measures of the skull in an avian morphometric study. Grant and Grant (1993) are the first to link avian morphometric results to dietary groups. Hertel (1994) is the first to study purely the morphometrics of the avian skull, and his follow-up study (Hertel, 1995) is the first to compare fossil and modern avians. Hertel (1995) defines 21 linear and 4 angular measurements to define the avian skull, though subsequent studies of the skull tend to focus exclusively on linear (Ladyguin, 2000; Li & Clarke, 2016) or angular (Button, 2018) morphometrics. All of the landmarks chosen in these studies make up some subset of Hertel's (1995). These studies, as well as those incorporating postcranial morphometrics (Barbosa *et al.*, 1999; Corbin, 2008; Herrel, Soons, Aerts *et al.*, 2010b; Pigot *et al.*, 2020) and mechanical properties of the skull (Corbin *et al.*, 2015; Herrel *et al.*, 2010b), have all been successful in predicting diet in extant birds. However, as is always true with TM, the *a priori* selection of measurements may limit their broader applicability. Beak curvature, for instance, is a major axis of the morphospace in vultures (Hertel, 1994) but considered irrelevant in shorebirds (Barbosa *et al.*, 1999) and flycatchers (Corbin, 2008). Pigot *et al.* (2020) successfully applied their morphometrics to 99.7% of extant bird species, but their reliance on soft tissue landmarks means they are inapplicable to most fossil taxa.

### (iv) Traditional Morphometrics in Fossil Theropod Skulls

While TM applied to non-avian theropods is applicable across a broad range of taxa, the diets of all extinct theropods are necessarily speculative. So, TM use has been mostly restricted to detecting inter-population differences (Smith, 1998) or niche partitioning (Henderson, 1998; Holtz Jr., 2008; Van Valkenburgh & Molnar, 2002) in fossil theropods. Holtz Jr. (2008) frames his study of tyrannosaurids in terms of hunting style, but the section therein focusing on skull morphometrics only shows a lack of separation of tyrannosaurids and non-tyrannosaurid morphospaces with no comments on diet. Button and Zanno (2020) incorporate many skull measurements typical of TM in a study of theropod diet, such as

skull length or premaxillary angle. However, because they investigate them in context of function rather than shape the study is covered in Section V.3.b.iii Lever Models of Fossil Theropods.

While most rigorously defined for phylogenetic use (Hendrickx & Mateus, 2014), TM has been applied twice to non-avian theropod teeth as an exploration of diet. Holtz Jr. *et al.* (Holtz Jr., Brinkman & Chandler, 1998) found a trend of generally coarser denticles in herbivorous vertebrates than carnivorous ones, with therizinosaurs and some troodontids plotting near the former. This technique is inapplicable to currently-known avians as all reported avian teeth are unserrated (O'Connor 2019; but see Dumont, Tafforeau, Bertin *et al.*, 2016 and Wang, Zhao, Shen *et al.*, 2015a). Holtz Jr. (2008) examined the height, length, and width of theropod tooth crowns, which can be applied to unserrated teeth. While tyrannosaurid teeth weakly separated in the morphospace from non-tyrannosaurids, they were only effectively distinguished when examining functional indices [with an implication of increased strength indicating osteophagy]. No comparisons to taxa with known diets were made.

#### (v) Traditional Morphometrics in Extant and Extinct Theropod Feet

Every TM study of the fossil theropod pes has included extant taxa for comparison, and few studies of extant avians exclude fossil taxa. It is, therefore, unhelpful to divide these studies by their subject, and so instead are divided by their focus on angular or linear measurements. In addition to studies concerned directly with dietary cues (Csermely *et al.*, 2006; Csermely *et al.*, 2012; Einoder *et al.*, 2007; Fowler *et al.*, 2009; Fowler, Freedman, Scannella *et al.*, 2011; Mosto & Tambussi, 2014; Wang *et al.*, 2014c) and grasping ability (Kambic, 2008), we also include studies concerned purely with locomotion (Clark, Hopson, Fastovsky *et al.*, 1998; Cobb *et al.*, 2020; Dececchi & Larsson, 2011; Feduccia, 1993; Glen & Bennett, 2007; Hopson, 2001; Morschhauser *et al.*, 2009; Peters & Görgner, 1992; Pike & Maitland, 2004; Tinius & Russell, 2017; Zhou, 1999). Locomotion can provide insight into what resources would be available to individuals to feed on and how free the hind limbs were to manipulate food while feeding.

#### 1. Angular Measures

The first application of TM, based on the curvature of avian claws, was proposed by Peters and Görgner (1992). However their method, ambiguously worded and requiring fitting of an idealised parabola, was less influential than that of Feduccia (1993). Feduccia modelled the ventral arc of the claw as that of a circle (Figure 1A) and proposed that a greater degree of curvature [i.e. larger central angle measure] of said arc indicated a greater degree of arboreal behavior within avians. Xu *et al.* (2000) briefly mention utilising this method in their description of *Microraptor zhaoianus*. Pike and Maitland (2004) would go on to formalise this circular arc methodology, preferring to measure the dorsal arc of the claw due to a less ambiguous proximal landmark (see Figure 1C). Csermely and Rossi (2006) utilised this method and were the first to apply it to feeding behaviour. They investigated if it could distinguish between raptorial and non-raptorial birds, as the former is known heavily utilise their feet in prey restraint and manipulation (e.g. Csermely & Gaibani, 1998b; Goslow, 1972; Sustaita *et al.*, 2013; Ward, Weigl & Conroy, 2002). Subsequent studies measuring curvature would either measure the dorsal arc exclusively (Csermely *et al.*, 2012; Glen *et al.*, 2007; Mosto *et al.*, 2014) or both the dorsal and ventral arcs (Birn-Jeffery, Miller, Naish *et al.*, 2012; Cobb *et al.*, 2020; Fowler *et al.*, 2009; Fowler *et al.*, 2011; Tinius *et al.*, 2017) [though note (Fowler *et al.*, 2009; Fowler *et al.*, 2011; Tinius *et al.*, 2017) use an alternate ventral arc proximal landmark (Figure 1B)]. Dececchi and Larsson (2011) code claws as either “straight”, “recurved”, or “highly recurved” without clarification of how these codings are determined. Of these methods, the angle from the dorsal arc as defined by (Pike *et al.*, 2004) or the ventral arc as defined by (Fowler *et al.*, 2009) appear to be the most informative as both utilise unambiguous landmarks that represent the proximal and



distal extents of the keratinous covering of the talon (Hedrick, Cordero, Zanno *et al.*, 2019a). Because the majority of studies incorporate other measurements in addition to curvature it is impossible to pinpoint which measure most effectively discriminates between groups.

## 2. Linear Measures

The second application of TM, based on linear measurements of the phalanges, traces its roots to the descriptive measurements of Shufeldt (1909 pg. 47, 93, 125, 196, 335). The first statistical analysis of these measurements as ecological indicators appears to be from Leisler and Thaler (1982). They and subsequent researchers, though, tended to utilise pedal measurements as a single component in larger whole-body morphometric analyses. Zhou (1999) included a chapter which introduced analysis of avian ecomorphology focused entirely on pedal proportions, though the paradigm was popularised by and is commonly attributed to Hopson (2001). Both studies followed similar methodologies to intuit locomotor behavior in extinct birds. Csermely and Rossi (2006) were the first to investigate signal of diet [specifically predatory behavior] with linear pes measurements. Kambic (2008) was the first to search for signals specific to grasping ability in the pes. Unlike angular measurements, linear measurements of the avian pes are far from standardised. Non-ungual phalanges may have measurements of length [whole toe (Abourachid, Fabre, Cornette *et al.*, 2017; Clark *et al.*, 1998; Dececchi *et al.*, 2011; Hopson, 2001; Kambic, 2008; Morschhauser *et al.*, 2009; Wang *et al.*, 2014c; Zhou, 1999) and individual phalanges (Abourachid *et al.*, 2017; Clark *et al.*, 1998; Dececchi *et al.*, 2011; Hopson, 2001; Kambic, 2008; Morschhauser *et al.*, 2009; Wang *et al.*, 2014c; Zhou, 1999)], width [whole toe (Csermely *et al.*, 2006; Csermely *et al.*, 2012) and individual phalanges (Abourachid *et al.*, 2017; Kambic, 2008)], and height [whole toe (Csermely *et al.*, 2006; Csermely *et al.*, 2012) and individual phalanges (Abourachid *et al.*, 2017; Kambic, 2008)] on record (Figure 2A–C). Ungual phalanges may have measures of length [chord (Csermely *et al.*, 2006; Csermely *et al.*, 2012; Fowler *et al.*, 2009; Fowler *et al.*, 2011; Hedrick *et al.*, 2019a), arc length (Abourachid *et al.*, 2017; Mosto *et al.*, 2014), and flexor tubercle chord (Abourachid *et al.*, 2017; Mosto *et al.*, 2014)], width [at the base of the claw (Csermely *et al.*, 2006; Csermely *et al.*, 2012) and of the flexor tubercle (Mosto *et al.*, 2014)], and height [at the base of the claw (Fowler *et al.*, 2009; Fowler *et al.*, 2011), at the midpoint of the arc (Fowler *et al.*, 2009; Fowler *et al.*, 2011), and of the flexor tubercle (Mosto *et al.*, 2014)] on record (Figure 2D–E). The only linear measurements comparable with the majority of studies of linear phalangeal measurements are ungual chord length and individual phalanx length [which closely approximates total toe length (Falk, 2014)]. Arguments have been made for the inclusion of certain measurements, e.g. the ungual chord length as a proxy for body size (Pike *et al.*, 2004) or phalangeal lengths indicating flexibility of the toes (Zhou 1999 citing Fisher 1946). However, the inconsistency of measures attests to how little justification there is for any given set of linear measurements to properly quantify the shape of avian feet.

This category includes two of the three quantitative studies focused on avialan diet, though both are directly examining locomotion and only refer to diet as a subsidiary of these results. Hou *et al.* (2004) hypothesised that *Longirostravis* was a probing feeder, and Morschhauser *et al.* (2009) tested the hypothesis through the closely-related *Rapaxavis*. Morschhauser *et al.* (2009) incorporated digit III phalanx measurements of *Rapaxavis* into the data set of Hopson (2001) and found it to fall into the arboreal morphospace. From this they propose that, instead of being ground probers comparable to charadriiforms, they may have instead been bark probers (Morschhauser *et al.*, 2009 pg. 553). Wang *et al.* (2014c) took a similar approach to Bohaiornithidae, adding their phalangeal measurements to the dataset of Hopson (2001) and adding claw curvature measurements of Pike and Maitland (2004) [it appears

comparisons to the latter are merely qualitative, referring only to “high” and “low” curvature]. Narrowing the dataset to raptorial birds due to high claw curvature and lack of specialised limb proportions associated with climbing, they found the closest phalangeal proportions to be those of *Pandion*. While the researchers propose a piscivorous diet in bohaiornithids because of this similarity, they attenuate the diagnosis with qualitative assertions of their teeth seeming to be more adapted for durophagy (Wang *et al.*, 2014c pg. 68).

(vi) *Discussion*

Pilot studies are necessary to examine the application of TM to non-avian avialan skulls to predict diet. All TM frameworks that rely on the curvature of the premaxilla (Barbosa *et al.*, 1999; Button, 2018; Hertel, 1994; Hertel, 1995) cannot be applied to toothed avialans as nearly all possess straight premaxillae. The exceptions are for *Sapeornis*, *Ichthyornis* and the Hesperornithiformes which are interpreted as possessing rhamphothecae which replaced the rostralmost teeth (Wang *et al.*, 2020a); premaxillary curvature is likely a product of edentulism with teeth no longer serving as a gripping surface for food. Whether the straight beaks of all known fully-edentulous avialans (Chiappe *et al.*, 2001; Elzanowski *et al.*, 2018; O'Connor, 2019; Zhou, Zhou & O'Connor, 2012; Zhou, Zhou & O'Connor, 2013; Zhou & Li, 2010) represents a dietary or developmental signal is unclear. Frameworks reliant on the width of the skull (Corbin, 2008; Corbin *et al.*, 2015; Li *et al.*, 2016; Pigot *et al.*, 2020) would require referral of more specimens with skulls preserved in dorsal or ventral views. Skull width is unknown in all described non-avian avialan genera except *Anchiornis*, *Confuciusornis*, *Sapeornis*, *Dalingheornis*, *Eopengornis*, *Fortunguavis*, *Longusunguis*, *Monoenantiornis*, *Archaeorhynchus*, *Yixianornis*, *Ichthyornis*, *Hesperornis*, and *Parahesperornis* (Table 2). Of these all but *Ichthyornis*, *Hesperornis*, and *Parahesperornis* are crushed skulls whose widths are dubious. TM seems more likely to prove useful when applied to dentition. Dental TM is already known to effectively determine the phylogenetic placement of most theropods (Hendrickx *et al.*, 2014; Hendrickx, Tschopp & d Ezcurra, 2020), with universal characters already defined in the literature (Hendrickx, Mateus & Araújo, 2015). While preliminary studies of lizard teeth have shown only a tenuous link with diet (Estes & Williams, 1984; Melstrom, 2017), they remain the best extant group to compare to due to the much more narrow range of tooth form in crocodilians (Erickson, Gignac, Stepan *et al.*, 2012). So, we recommend a study of extant lizard teeth using the existing TM framework (Hendrickx *et al.*, 2015) as a baseline dietary morphospace. Otherwise, the lack of effective application of skull TM to non-avian theropod diet does not bode well for the technique applying to non-avian avialans.

TM studies of the theropod pes have both a more consistent focus on dietary analogues and a more consistently applied set of measurements, and so a more thorough evaluation of methods is possible. Most TM studies of the pes concerned themselves with only digit III [DIII] (Clark *et al.*, 1998; Cobb *et al.*, 2020; Dececchi *et al.*, 2011; Feduccia, 1993; Glen *et al.*, 2007; Kambic, 2008; Morschhauser *et al.*, 2009; Peters *et al.*, 1992; Pike *et al.*, 2004; Tinius *et al.*, 2017; Wang *et al.*, 2014c), two studies with digit I [DI] and DIII (Csermely *et al.*, 2006; Csermely *et al.*, 2012), two studies with digit II [DII]–digit IV [DIV] (Hopson, 2001; Zhou, 1999), and the remainder with all four digits (Einoder *et al.*, 2007; Fowler *et al.*, 2009; Fowler *et al.*, 2011; Mosto *et al.*, 2014). Studies that rely solely on DIII justify this choice by DIII being the weight-bearing toe (Glen *et al.*, 2007; Hedrick *et al.*, 2019a), and all but (Kambic, 2008) are concerned with locomotion. Studies on DII–DIV (Hopson, 2001; Zhou, 1999) are similar in concept, disregarding DI as it rarely bears weight during forward motion. Zhou (1999) compared the phalangeal proportions of DII–IV individually and found each digit discriminated between locomotor categories

equally well (his Figures 38–42). Backus *et al.* (2015) provide a strong argument for the efficacy of the two-dimensional simplification of avian grasping [i.e. to the action of DI and DIII] as such simplification is adequate to create functioning artificial ‘hands’ for robots (Backus, Odhner & Dollar, 2014; Dollar & Howe, 2011). Abourachid *et al.* (2017), though, note that in non-ansiodactyl pedal arrangements DII and DIV supplement or supplant the role of DI and DIII in grasping. Measuring all digits of the pes acquires the most data possible about its structure but imposes additional complexity in the analysis and time for data collection. Significant interdigital variation is often cited as a reason to avoid measurements of the whole pes (Cobb *et al.*, 2020; Hedrick *et al.*, 2019a), but Fowler *et al.* (2009; 2011) argue that interdigital variation itself is the diagnostic which separates ecological niches. Given that all of the above-mentioned studies of all four digits were able to discriminate among their chosen ecological groups it seems unlikely that their inclusion is detrimental to analysis, but any benefit from their inclusion over the analysis of a single digit is equivocal.

It is recommended in TM [and GM] studies of ungual phalanges to compare the bony cores of claws rather than the keratin sheathes. Studies measuring both sheath and core found both to give similar information (Cobb *et al.*, 2020; Hedrick *et al.*, 2019a), and many fossils do not preserve the keratin sheath. In those that do the keratin stain is likely more deformed than the bone (Saitta *et al.*, 2019). Therefore empirical conversion factors (e.g. Glen *et al.*, 2007; Hedrick *et al.*, 2019a) or direct measurements of the stain give less valid data than measuring the core. Extant avian ungual bones can be found in many museum skeletal collections, and remaining keratin can be easily removed from the claws of macerated skeletons by soaking the claws in dilute ammonia (Stephen Rogers, Carnegie Museum of Natural History, pers. com. 2019). Among styles of applying TM, Fowler *et al.* (2009; 2011) produced the finest scale of discrimination. Their framework delineated not only arboreal/ground and raptor/nonraptor groups, but also the individual hunting guilds among raptors. However, these studies did not employ phylogenetic correction. Thus, they have been proposed as simply grouping talons based on familial relations (Hedrick *et al.*, 2019a), which so happen to correlate with predatory style in extant raptors (Csermely *et al.*, 1998b; Fowler *et al.*, 2009; Goslow, 1972; Sustaita, 2008). More fundamentally, the Fowler *et al.* studies appear to utilise improper statistical analysis. They apply correspondence analysis, designed to utilise categorical data. Numerical data can be binned and analysed as categorical data (Kim, 2011), but Fowler *et al.* (2009; 2011) do not make note of doing so in either study. When the more appropriate principal component analysis is applied to these data (see Figure S1) groups inhabit less distinct regions of the morphospace. With that said, they still show more distinction than that of the next most discriminating study, Csermely *et al.* (2012). It is recommended, then, that any pes-based TM investigation of non-avian avian diet follow the analytical procedures of Fowler *et al.* (2009) with phylogenetic correction (see Section V.4.a Phylogenetic Signal).

### (c) Geometric Morphometrics

#### (i) Introduction

Geometric morphometrics [GM] uses digital models, either full 3D models or 2D dorsal/lateral silhouettes, with landmarks placed just as one would with TM (Zelditch *et al.*, 2004). Landmarks are typically supplemented by sliding semi-landmarks, a digitisation of the curve[s] between landmarks (Perez, Bernal & Gonzalez, 2006). The shifts in the relative position of landmarks and semi-landmarks between models can be quantified and analysed digitally (Adams & Otárola-Castillo, 2013).

While GM can be significantly more time-consuming than TM (Rohlf, 1990), GM provides at least as much information about the shape of the element in question as every possible measurement taken between every pair of landmarks (Zelditch *et al.*, 2004 pg. 2–7). With the introduction of semi-landmarks the information increases beyond that possible in TM (Zelditch *et al.*, 2004 pg. 396). Perez *et al.* (2006) detail different methods of creating semi-landmarks. Note that at an inter-species or higher level, such as that typically employed in palaeontology, the differences between methods should be negligible. Because GM creates an unbiased representation of form it has proven effective in tracing evolutionary trends (Fernandez Blanco, Cassini & Bona, 2018; Figueirido, MacLeod, Krieger *et al.*, 2011; Miyashita, 2013; Openshaw, D'Amore, Vidal-García *et al.*, 2016; Polly, Stayton, Dumont *et al.*, 2016) but has returned only mixed results in feeding studies (Gailer, Calandra, Schulz-Kornas *et al.*, 2016; Kłaczko, Sherratt & Setz, 2016; Meloro, Hudson & Rook, 2015; Samuels, 2009; Tarquini, Chemisquy, Ladevèze *et al.*, 2019).

Another form of GM, outline-based GM, has been introduced as an alternate method of quantifying 2D shape. This method interprets and compares the outlines of a bodies as composites of trigonometric curves [Fourier analysis]. The primary advantage over landmark-based methods is applicability to bodies with few homologous landmarks (Bonhomme, Picq, Gaucherel *et al.*, 2014). While the concept dates back several decades (Kaesler & Waters, 1972) this method has been used very little in vertebrate palaeontology so far (Navarro, Martin-Silverstone & Stubbs, 2018; Schaeffer, Benton, Rayfield *et al.*, 2019) and so by default ‘GM’ will be used in this paper to refer to landmark-based GM. In phylogenetically broad studies in which homology is unclear this method may prove useful in coming years.

#### (ii) *Geometric Morphometrics in Extant Bird Skulls*

van der Meij (2004) and Marugán-Lobón and Buscalioni (2004) were the first to apply geometric morphometrics (GM) to bird skulls. van der Meij (2004) combined landmarked pictures of fringillid and estrildid finches into effectively 3D models of the skull. Shape differences in these models were analysed and compared to bite force. Bite force, in turn, was translated into seed husking efficiency and its effect on diet (van der Meij, 2004). This study, then, is also the first to apply GM to diet [albeit indirectly]. Marugán-Lobón and Buscalioni (2004) were the first to apply 2D GM to both extant birds and non-avian theropods, and Kulemeyer *et al.* (2009) the first to use true 3D models for bird skull GM.

GM Studies of extant avian diet can be divided into those that study only the bill (Button, 2018; Cooney, Bright, Capp *et al.*, 2017; Matsui, Hunt, Oberhofer *et al.*, 2016; Navalón *et al.*, 2018a; Olsen, 2017; Sustaita & Rubega, 2014; van der Meij, 2004), those that treat the beak and cranium as a single unit (Bright *et al.*, 2016; Bright, Marugán-Lobón, Rayfield *et al.*, 2019; Chávez-Hoffmeister, 2020; Felice *et al.*, 2019a; Navalón, 2014; Pecsics *et al.*, 2019; Si *et al.*, 2015; Sun *et al.*, 2018; Tokita, Yano, James *et al.*, 2017), and those that examine each individually (Bright *et al.*, 2016; Felice *et al.*, 2019a; Kulemeyer *et al.*, 2009). Landmarks for the bill tend to be consistent. The bill tip and the base of the frontal are used in nearly every study (Bright *et al.*, 2016; Bright *et al.*, 2019; Button, 2018; Chávez-Hoffmeister, 2020; Kulemeyer *et al.*, 2009; Matsui *et al.*, 2016; Navalón, 2014; Navalón *et al.*, 2018a; Si *et al.*, 2015; Sun *et al.*, 2018; Tokita *et al.*, 2017; van der Meij, 2004) alongside either the rostral extreme of the jugal (Bright *et al.*, 2019; Button, 2018; Chávez-Hoffmeister, 2020; Matsui *et al.*, 2016; Navalón, 2014; Navalón *et al.*, 2018a; Tokita *et al.*, 2017; van der Meij, 2004) or the ventrocranial extreme of the palatine (Kulemeyer *et al.*, 2009; Pecsics *et al.*, 2019; Si *et al.*, 2015; Sun *et al.*, 2018; Tokita *et al.*, 2017). Landmarks of the cranium are less consistent, with only the border of the orbit commonly mapped (Felice *et al.*, 2019a; Kulemeyer *et al.*, 2009; Si *et al.*, 2015; Sun *et al.*, 2018; Tokita *et al.*, 2017). Discrimination of diet is

inconsistent. GM can often discriminate diet in restricted phylogenetic groups [family- to superfamily-level] (Chávez-Hoffmeister, 2020; Kulemeyer *et al.*, 2009; Olsen, 2017; Pecsics *et al.*, 2019; Si *et al.*, 2015; Sun *et al.*, 2018; Sustaita *et al.*, 2014; van der Meij, 2004), but dietary morphospace tends to heavily overlap between groups (Bright *et al.*, 2016; Bright *et al.*, 2019; Tokita *et al.*, 2017) and across birds overall (Button, 2018; Felice *et al.*, 2019a; Navalón, 2014; Navalón *et al.*, 2018a). It seems likely that more distantly-related clades of birds face developmental constraints that either prevent them from attaining converging morphotypes or predisposes them towards differing solutions under similar environmental pressures (Gould, 2002). Similar levels of craniofacial integration within most families of land birds [Telluraves] (Navalón, Marugán-Lobón, Bright *et al.*, 2020) implies similar developmental patterns among them. Constraints preventing shape/diet correlation, then, are less expected within Telluraves.

### (iii) *Geometric Morphometrics in Fossil Theropod Skulls*

As stated above, Marugán-Lobón and Buscalioni (2004) were the first to apply GM to non-avian theropods. They focused purely on disparity [quantified difference in shape]. Shychoski and Snively (2008a) published an abstract that is the first use of GM to investigate non-avian theropod feeding. They reported juvenile tyrannosaurs to more resemble non-tyrannosaurid morphotypes than their adult forms, but a full publication of the data is not planned (Eric Snively pers. com. 2020). Brusatte *et al.* (2012) found oviraptorids and, to a lesser extent, ornithomimosaurians and alvarezsaurians [all proposed as herbivorous (Zanno *et al.*, 2011)] to cluster outside the ‘carnivorous’ morphospace. The lack of any confirmed diets in the dataset limits the validity of the results. Foth and Rauhut (2013) report similar results with a significant correlation between purported diet and shape. They saw more overlap in dietary morphospace and also did not include taxa with known diets. Recoding taxa based on more explicit evidence of diet (Figure S2) provides a division between herbivorous and carnivorous morphospace if *Anchiornis* and *Bambiraptor* are excluded [which inhabit a unique region of the morphospace overlapping with herbivores]. This study is the first to include early-diverging avian taxa in GM analysis (Table S1 in Foth *et al.*, 2013). Schaeffer *et al.* (2019) are the first to incorporate outline-based GM to study theropod diet, and compare it to both landmark-based GM and discrete characters. The authors found all three approaches to be of similar effectiveness, though it is noteworthy that separation of morphospace via discrete characters appears more vulnerable to changes in sample size (compare their Figures 3C and 7C). When diets are more conservatively assigned (Figure S3) resolution between herbivores/omnivores and small carnivores increases. Their graphs show landmark-based GM as less effective at discriminating between large and small carnivores than outline-based GM (their Figures 3 and 7), though this appears to be due to principal components defining different shapes between the two (their Figure 4). When plotted as PC1 vs PC3 a similar trend is seen in landmark-based GM to that shown for outline-based (their Figure S7 and supplemental files; our Figure S3E). Button (2018) and Navalón (2014) are the only researchers to combine GM analysis of extant avian taxa with known diets and early-diverging theropods. The dietary morphospace of most extant avians in these studies overlap and are often completely enveloped in another morphospace (Figure 33 in Button, 2018; Figure 8 in Navalón, 2014). In both studies almost all fossil taxa either plot in regions of heavy overlap or fall completely outside the dietary morphospace. The exception is two unidentified taxa in (Button, 2018), which plot in a region inhabited only by terrestrial herbivores.

All GM studies of early-diverging theropod diet are performed in two dimensions, all are concerned with the upper jaw except for (Schaeffer *et al.*, 2019; Shychoski *et al.*, 2008a) [on the lower

jaw], and all landmarks are placed in in only lateral view except for in (Button, 2018) [placed in lateral and dorsal view]. Landmarks of theropod skulls are variable. The only landmarks shared between the four studies of the upper jaw are the rostroventral extreme of the premaxilla and the contact between the jugal and maxilla (Brusatte *et al.*, 2012; Button, 2018; Foth *et al.*, 2013; Navalón, 2014). Landmarks used in both studies of the lower jaw (Schaeffer *et al.*, 2019; Shychoski & Snively, 2008b) are the anterodorsal and anteroventral corners of the dentary, the dentary-surangular suture, the dentary-angular suture, and the articular glenoid. None of the studies include any landmarks on teeth, allowing toothed organisms to be examined using the same landmarks as those with varying extents of edentulism (Button, 2018; Wang *et al.*, 2020a). The importance of tooth morphology in lepidosaur (Smith, 1993) and mammal (Bergqvist, 2003; Pineda-Munoz, Lazagabaster, Alroy *et al.*, 2017) diet raises questions of error introduced in this practice. A single study (D'Amore, 2009) has studied theropod teeth in isolation using GM, but focused on penetration angle independent of diet.

#### *(iv) Geometric Morphometrics in Extant Bird Feet*

Tinius and Russel (2017) were the first to apply GM methodology to bird claws, finding GM to be the only one of six claw angle measures capable of discriminating locomotor groups in the taxa studied [GM, the aforementioned methods of (Feduccia, 1993; Peters *et al.*, 1992) [though note their execution of (Feduccia, 1993) more closely resembles that of (Fowler *et al.*, 2009)], one method applied to insects (Petie & Muller, 2007), one to lizards (Zani, 2000), and one theorised in but never applied to amniotes (Thompson, 1942)]. Hedrick *et al.* (2019a) were the first group to investigate dietary signal with geometric morphometrics,. They found neither TM nor GM could discriminate between ecological groups. Abourachid *et al.* (2017) and Tsang *et al.* (2019) incorporate three-dimensional models and landmarks into their studies, but (Abourachid *et al.*, 2017) uses them primarily to automate the collection of linear measurements for TM. Tsang *et al.* (2019) are able to discriminate both between predatory and non-predatory taxa and between sizes of prey predators fed on. This level of precision presents a promising outlook for the future. This is only one study, though. While theoretically 3D GM provides the most accurate representation of claw shape, the merits of the technique have not yet been proven consistent in discrimination of avian ecology.

#### *(v) Discussion*

GM studies of theropod taxa both extant and extinct have been highly variable in both methodology and results. 3D GM studies of skulls, presumably capturing the largest amount of data about shape, tend to find strong relationships between form and diet at the family- to superfamily level (Kulemeyer *et al.*, 2009; Olsen, 2017) that disappear when comparing larger-scale trends (Bright *et al.*, 2016; Felice *et al.*, 2019a; Navalón *et al.*, 2018a). GM analysis of non-avian theropods is inhibited by the lack of known diets in the group. Certain groups, in particular oviraptorids, tend to cluster distinctly from other theropods (Brusatte *et al.*, 2012; Foth *et al.*, 2013). A study incorporating both oviraptorids and caenagnathids [both beaked but respectively hypothesised to be herbivorous and carnivorous (Ma, Pittman, Lautenschlager *et al.*, 2020)] may clarify if the segregation in these studies stems from edentulism or a dietary shift. Schaeffer *et al.* (2019) found considerable overlap of dentary morphospace in herbivorous and small carnivorous taxa. But, they coded all oviraptorosaurians as herbivorous [including caenagnathids] and all troodontids as small carnivorous [including those proposed as omnivorous (Holtz Jr. *et al.*, 1998; Torices *et al.*, 2018)] (Table S5 in Schaeffer *et al.*, 2019). This study highlights the necessity of defining dietary morphospaces with organisms whose diet is not controversial. When assumptions of herbivory in therizinosaurs and carnivory in avialans are discarded the

morphospaces become more distinct (Figure S3C–E). The study also shows landmark- and outline-based GM to provide similar information about skull shape, though comparisons between the two should be preceded by comparing thin spline plates to ensure the same shape changes are being modelled on each axis (Figure S3E). While Button (2018) successfully combines extant and extinct theropods in skull GM analysis, he reports only landmarks placed in dorsal/ventral view are effective at discriminating diet categories. Even then, his model appears to better separate terrestrial feeders from other groups rather than any particular diet categories (his Figure 33). Regardless, dorsal/ventral placement of skull landmarks excludes most published non-avian avialan specimens (Table 2). Navalón (2014) managed to assign non-avian avialans to a diet category using GM, but only in conjunction with mechanical advantage (see Section V.3.b.iii Lever Models of Fossil Theropods). In sum, skull shape tends to be a poor predictor of diet in extant avians. Studies of non-avian theropod morphospaces are limited by a lack of knowledge about the diet of taxa included, and still provide only mixed results. An attempt to combine avian and non-avian theropods show poor resolution in the dietary morphospace and has limited application to fossil avialans. Therefore, GM of the non-avian avialan skull is recommended as at most a compliment to a functional study (e.g. Navalón *et al.*, 2018a) if it is utilised at all.

Tsang *et al.* (2019) were able to distinguish nonpredatory taxa [both nonraptors and scavenging raptors] from predatory taxa, and partially distinguish between predators hunting large prey and small prey. Unfortunately, there are few fossil bird claws preserved in three dimensions. Large-scale 2D geometric morphometric analysis of bird claws has only been undertaken by Hedrick *et al.* (2019a) who failed to discriminate between any ecological groups. By their own admission, their ecological categories may have been too broad to allow for delineation (Hedrick *et al.*, 2019a pg. 9), and that the ratio between the lengths of keratin sheaths and bone cores was a major element of the first principal component in their analysis (Hedrick *et al.*, 2019a pg. 6). It is possible that a similar study landmarking solely bone cores and with a more diverse set of ecological groupings (e.g. the ecological groups of Fowler *et al.*, 2009; Glen *et al.*, 2007; Pigot *et al.*, 2020) may produce more useful results. Therefore, we recommend utilising the techniques of Tsang *et al.* (2019) where possible and an improved 2D GM framework where not possible.

### (3) Functional Studies

#### (a) Introduction

Of the two styles of functional study described herein, finite element analysis [FEA] is typically the go-to for fossil dietary inference. Lever models, which provide measures of mechanical advantage [MA], are more commonly used as broad approximations that can be calculated easily and quickly. However, recent questions of the validity of FEA results in small animals (see Section V.3.c.ii The Strength Criterion) may apply to non-avian avialans [in which the largest taxa other than *Hesperornis* (Martin *et al.*, 2008) and possibly *Gargantuavis* (Mayr, Codrea, Solomon *et al.*, 2020) have a skull length less than 9 cm (Field *et al.*, 2018b)]. Both techniques, then, should be taken into consideration.

Relevant to both techniques are concerns of dimensionality and comparison taxa. In both cases, researchers will likely be restricted to 2D analysis due to the reconstruction issues related to incomplete skull preservation (detailed in Section V.1 Skull Reconstruction). But, if possible, validation studies with smaller sample sizes using both 2D and 3D techniques should be performed. A 3D model of the skull of *Ichthyornis dispar* is already published (Field *et al.*, 2018b), and the phalanges of *Mirarce eatoni* are three-dimensionally preserved (Atterholt *et al.*, 2018). Both are prime candidates for such studies.

Both FEA and lever models of fossil taxa are only interpretable in comparison with other models. The choice of comparative taxa, then, is of vital importance. To create a phylogenetic bracket, beyond the obvious extant avians, it is recommended that early-diverging non-avian paravians with preserved meals (O'Connor *et al.*, 2019) are included. Certain lepidosaurian taxa that share similarities in dentition (Smith, 1993) with toothed avialans may be necessary to include as well.

## (b) Lever Modelling

### (i) Introduction

The jaw of most animals acts as a third class lever: the joint acts as a fulcrum about which a load - the distal bone - is rotated by the effort - a muscle - attached in between (Figure 3). Because jaws act as levers we can use known mathematical principles to predict their behaviour. Examples include models utilising known muscle vectors to calculate bite vectors (Santana, Dumont & Davis, 2010) or to compare torque generation (Kiltie, 1982). However, when studying fossils, muscle size can often only be very roughly approximated. Instead, they tend to rely on the concept of mechanical advantage [MA].

The basic principle of MA was outlined in antiquity by Archimedes of Syracuse (1897 pg. 192–194): the force exerted on a lever will be multiplied by the distance from it to the fulcrum [the inlever] and divided by the distance from the fulcrum to the load [the outlever]. MA, then, is the useful ratio defined as the inlever divided by the outlever. In practical terms, it can be viewed as the ratio of the output force to the input force of a system; or, as the factor by which the input force is multiplied to determine the output force. When mechanical advantage is lowered, the load is moved across a greater distance in the same amount of time and thus at a higher speed. As a trade-off, the output force is reduced.

In animal jaws, then, the inlever is the distance from the site of adductor muscle attachment to the joint between the cranium and mandible. The outlever is the distance from said joint to the point at which the animal is biting down. The bite point is typically operationalised as the rostralmost or cranialmost point of the occlusal surface [either of the rhamphotheca or apices of teeth] (Ma *et al.*, 2020; Navalón *et al.*, 2018a). More rarely MA will be calculated for each tooth in the row (Cox, 2017; Sakamoto, 2010; Therrien, Quinney, Tanaka *et al.*, 2016). Functionally, MA is seen as the trade-off between speed and power in skull architecture (Adams, Rayfield, Cox *et al.*, 2019; Corbin *et al.*, 2015; Dumont, Samadivam, Grosse *et al.*, 2014; Stayton, 2006), though this trade-off is a simplification and can be circumvented (Corbin *et al.*, 2015; McHenry & Summers, 2011). Organisms feeding on immobile sources, such as plant matter, experience selective pressure for high jaw forces [high MA] that can efficiently process food. In contrast, those hunting mobile prey, such as insects or small vertebrates, experience selective pressure for jaws able to open and close quickly [low MA] in order to capture prey (Adams *et al.*, 2019; Corbin *et al.*, 2015; Stayton, 2006). Some trophic specialisations involving elongation of the snout such as nectarivory (Dumont *et al.*, 2014) or probe-feeding (Navalón *et al.*, 2018a) necessarily also reduce MA.

### (ii) Lever Models of Extant Birds

Burger (1978) was the first to record MA in extant birds. They found that higher MA in several South African cormorant [*Phalacrocorax*] species correlates with capturing larger prey while lower MA correlates with hunting small, fast-moving prey. Similar trends have been found in suliform birds (Carlos, Alvarenga & Mazzochi, 2017). Most subsequent studies in the field focused on “force-velocity trade-offs” by measuring bite force and jaw speed directly (Herrel, Podos, Vanhooydonck *et al.*, 2009). Corbin *et al.* (2015) returned to mechanical advantage in order to examine trends in bite force across extant avians (especially among passerines), finding that bite force and velocity correlate positively and



negatively with MA respectively. They briefly and qualitatively comment on diet by predicting low MA in insectivorous birds and high MA in granivorous birds (Corbin *et al.*, 2015 pg. 813). When analysed quantitatively the data they provide shows only a weak correlation [ $R^2 \approx 0.05$ ] between MA and diet. The correlation increases to moderate strength [ $R^2 \approx 0.40$ ] when restricted to passerine taxa (Figure S4). Olsen (2017) was the first to quantitatively examine the relationship between MA and diet, finding increased MA correlates to increased consumption of leaves in Anseriformes. Navalón *et al.* (2018a) broadened the scope of this technique by applying it across extant avians, but found MA to only weakly explain dietary components. Their data shows MA better predicting the use of the beak during feeding [UBF], particularly when combined with beak curvature (Navalón *et al.*, 2018a). Further research may translate UBF into a set of possible dietary categories [taxa with tearing UBFs are often raptorial, cracking UBFs at least partly granivorous, etc.]. It is of note that all quantitative dietary analyses of avian skull MA have focused on the upper jaw, with those regarding the lower jaw (Burger, 1978; Carlos *et al.*, 2017) being purely qualitative (but see Figure S4).

### (iii) Lever Models of Fossil Theropods

Mazzetta *et al.* (2009) are the first to measure MA in non-avian theropods, reporting differences in MA between *Carnotaurus* and *Allosaurus* without addressing possible implications of these differences. Sakamoto (2010) and Brusatte *et al.* (2012) calculated MA across the tooth row for a variety of non-ornithomimidae tetanurans and found lower values of MA in smaller taxa. This was tentatively interpreted as smaller taxa being adapted for hunting more agile prey. Sakamoto (2010 pg. 3330), however, proposes the potential for saw-motion biting in dromaeosaurids feeding on large prey. Noticeable in their figures (Figure S6 in Brusatte *et al.*, 2012; Figure 2b in Sakamoto, 2010) but not commented on in the text is the high MA of oviraptorosaurians, studied in-depth by Ma *et al.* (2019; 2020; 2017). Oviraptorosaurians consist of two distinct groups, Oviraptoridae and Caenagnathidae (Funston, Mendonca, Currie *et al.*, 2018). Only the Oviraptoridae was included in previous studies (Brusatte *et al.*, 2012; Sakamoto, 2010). Ma *et al.*, first in passing (2017) and later with statistical rigor (2019), note that caenagnathids tend to have a lower MA than oviraptorids. This is proposed to reflect herbivory in oviraptorids and carnivory in caenagnathids. Caenagnathid MA indeed falls within ranges of other theropods, while oviraptorid MA is significantly higher (Ma *et al.*, 2020). Ma *et al.* (2020) introduce MA measurements of *Jeholornis* as well as anchiornithines and scansoriopterygids. Button and Zanno (2020) include six measures of MA, five additional mechanical indices, and twenty-two traditional morphometric ratios believed to have functional implications in diet. While they effectively delineate herbivory and carnivory as well as distinct herbivorous strategies, the lack of variance explained by the principal components used is problematic [further discussed in the following section]. Navalón (2014) is the only MA study to focus chiefly on non-avian avialans. Combining MA with GM, it recovers *Confuciusornis* and *Sapeornis* as herbivorous and *Eoconfuciusornis* as omnivorous. While it presents initial promise, additional MA measurements of the fossil taxa should be taken before broader comparisons are made (Guillermo Navalón pers. com. 2020) and a subsequent study discussed above which expands the avian taxa included (Navalón *et al.*, 2018a) resolved diets much less effectively. This highlights the importance of not only including extant taxa but a *wide range* of extant taxa for ecological comparisons.

### (iv) Discussion

While MA has proven an effective predictor of diet in lepidosaurs (Stayton, 2006) and small mammals (Adams *et al.*, 2019; Dumont *et al.*, 2014), Navalón *et al.* (2018a) report minimal association between diet

and MA across extant birds. This is likely due to the fact that, as they point out, similar diets can often be associated with radically different foraging strategies (Navalón *et al.*, 2018a pg. 423). In particular, only some granivorous birds crack open the hard outer coating of [de-husk] seeds (Prosser & Hart, 2005). Those that do not de-husk instead crush them in the muscular gizzard (Janzen, 1981). Confuciusornithids and enantiornithines are not believed to possess a gizzard (O'Connor *et al.*, 2019) meaning they would not be susceptible to this source of signal interference. With this trend accounted for, mechanical advantages above 0.15 in Navalón *et al.* (2018a)'s dataset appear to correspond to herbivory while those below 0.15 correspond to carnivory, nectarivory, and frugivory [though the latter is complicated by the inclusion of nuts as fruits rather than with seeds. For instance, *Anodorhynchus* has an MA near 0.35 (Fig. 6 in Navalón *et al.*, 2018a) and is coded as 70–100% frugivorous depending on species (Wilman *et al.*, 2014). However these 'fruits' are palm nuts (Faria, Guedes, Yamashita *et al.*, 2008) which are hard enough that a close relative has been used as a model for construction materials (Staufenberg, Graupner & Müssig, 2015)]. Together, these factors make MA promisingly informative of diet in confuciusornithids and enantiornithines. MA may still provide insight into other non-avian avialans, but low MA cannot definitely rule out herbivory in these other groups.

The broad scope of the study by Button and Zanno (2020), combining MA with TM and some less commonly-used functional indices, gives good evidence for diet being delineated. Taxa such as ornithomimids and sauropods that are well-established as herbivorous occupy a region of the morphospace distant from theropods well-established as carnivorous [e.g. tyrannosaurids]. Their methods, at first, appear promising to apply to non-avian avialans. While Button and Zanno cite their resolution as reason to incorporate large complexes of characters in dietary analyses (Button *et al.*, 2020 pg. 163), the small amount of variance explained by their graphs belies this fact. The authors perform statistical operations within the first three principal components of morphospace which only explains 45.7% of the total variance observed; nine principal components are required to pass 70% explanation (their Data S2) as is the common standard (Jolliffe & Cadima, 2016 pg. 4). PCA functions to reduce the number of dimensions worked within an analysis by creating axes that explain large amounts of the variance. But once data encompasses a large number of dimensions it is unlikely that principal components will be able to explain adequate amounts of variance in the three dimensions humans can easily work in (Brown, 2009). It is therefore recommended here that the number of variables investigated are reduced to reduce the dimensionality of the data (but see Guillerme *et al.*, 2020 for alternatives to dimension reduction). Of the 34 measurements taken by Button and Zanno (2020), only nine [C2–8, 22, 23] can be applied to compression fossils and have proven effective at discriminating diet among extant groups other than ungulates. When only these measurements are used the first three principal components explain more than 70% of the variance while maintaining the same general structure of the morphospace (Figure S5). Data can be further reduced as the mechanical advantages for each individual muscle group have similar influence on principal components (Figure S2A in Button *et al.*, 2020) and so represent redundant information. They can be merged into anterior and posterior jaw-closing MA. Ma *et al.* (2020) additionally define jaw-opening MA and two other functional indices proven discriminative of diet across a wide range extant taxa. We recommend combining these studies to, in total, three forms of MA and five accompanying functional indices (Figure 4). However, unlike these two studies which measure MA based on the lower jaw, we recommend measurements of the upper jaw after the sensitivity analysis of Brusatte *et al.* (2012). They found the upper jaw, not the lower jaw, has the greatest effect on the overall MA of the jaw system (their Figure S27). Transition from the lower to upper jaw will necessarily require adaptation of landmarks from which measurements are taken. Landmarks at the anterior and posterior of

the occlusal margin, attachment sites of the adductor muscles, and those for *m. depressor mandibulae* will refer to their locations on the upper jaw rather than the lower jaw. The articular glenoid of the surangular will be replaced with the articular condyle of the quadratojugal (Figure 4B–D). All other landmarks remain unchanged from those in Button and Zanno (2020) and Ma *et al.* (2020).

### (c) Finite Element Analysis

#### (i) Introduction

Finite Element Analysis [FEA] is a technique originally applied to structural engineering. In it, irregular bodies are partitioned into a mesh of simple shapes, the ‘elements’, in order to predict the response of said body to a given load (Bathe, 2014). Applications in palaeontology began with structural predictions of depth adaptations in cephalopods (Daniel, Helmuth, Saunders *et al.*, 1997). Soon after it was used in its most common palaeontological application today: vertebrate jaws (Rayfield, Norman, Horner *et al.*, 2001). Bright (2014) provides an effective review of techniques up until 2014, but several notable advances have taken place since. The rise of formalised digital reconstruction (Lautenschlager, 2016) augmented by quantification of skeletal asymmetry (Hedrick, Schachner, Rivera *et al.*, 2019c) has opened the door for analysis of specimens previously considered hopeless cases. These techniques may allow for a greater number of studies incorporating both the upper and lower jaw. Studies taking both into account (Adams *et al.*, 2019; Attard, Parr, Wilson *et al.*, 2014; Attard *et al.*, 2016; Lautenschlager, Witmer, Altangerel *et al.*, 2013; Moreno, Wroe, Clausen *et al.*, 2008; Wroe, 2008; Wroe, Clausen, McHenry *et al.*, 2007) find differing peak Von Mises [VM] stress [a summary statistic of distortion energy in a body (Ugural & Fenster, 2012 pg. 189–190)] and differing distribution of VM stresses between the upper and lower jaw. These differences could in theory cause overestimation of jaw strength by only using the upper or lower jaw if the unmeasured element is the limiting factor in jaw strength. To our knowledge no study has been undertaken to explore the potential significance of this error, but no study of the full jaw has reported differing dietary signals between the upper and lower jaws.

Formalisation of applying two-dimensional FEA techniques to organisms (Marcé-Nogué, DeMiguel, Fortuny Terricabras *et al.*, 2013; Neenan, Ruta, Clack *et al.*, 2014) allows FE models to be constructed and analysed much faster with a lower prerequisite of computing power. It also allows for analysis of compression fossils. Simplifying three-dimensional bodies into two-dimensional outlines inherently requires assumptions that will induce error. Most commonly in palaeontology these assumptions are of planar strain, where strain in the excluded dimension is assumed to be negligible, or planar stress, where the stress exerted on the excluded dimension is assumed to be negligible. Planar strain is known to introduce error into shear strain predictions in bone (Verner, Lehner, Lamas *et al.*, 2016). Planar stress requires a known thickness of material (Marcé-Nogué *et al.*, 2013) that will likely vary over models. When thickness is known, creating 3D extrusion models from outlines has proven more valid than 2D assumptions (Morales-García, Burgess, Hill *et al.*, 2019). Plane stress assumptions, then, have little reason for use. For compression fossils, in which thickness is unknown, it is recommended to use planar strain assumptions so that error will be similar across studies. The exception to this recommendation is in groups like ornithomimid dinosaurs where significant shear strain is theorised to be involved in feeding (Rybczynski, Tirabasso, Bloskie *et al.*, 2008).

Finally, the comparison of FEA outputs has undergone a major paradigm shift. In the past, FE models have been compared qualitatively (Rayfield, 2005), or their peak (Rayfield, Milner, Xuan *et al.*, 2007) and/or average (Rayfield, 2011b) strength criteria compared quantitatively. Both quantitative

methods, however, are sensitive to the ways the FE model is constructed and require considerable mathematical correction (Marcé-Nogué, de Esteban-Trivigno, Escrig Pérez *et al.*, 2016). The amount of data required for correction renders comparison across studies nearly impossible. Marcé-Nogué *et al.* (2017) introduced the intervals method as a quantitative comparison robust to model construction. Ultimately, it determines the percentage of the model area/volume which experiences a given interval of stress. These percentages can then be compared directly in a histogram or plotted into principal component space. The intervals method has proven effective in subsequent feeding studies (Coatham, Vinther, Rayfield *et al.*, 2020; Miller *et al.*, In Press; Zhou, Winkler, Fortuny *et al.*, 2019), and is recommended here.

#### (ii) *The Strength Criterion*

A strength criterion is the measure of a model by which a researcher judges it. Typically, the lower the value of a strength criterion, the stronger the model. The strength criterion for biological FEA has traditionally been the peak VM stress (Figure 5A), after the recommendation of Dumont *et al.* (2009). Dumont *et al.* (2009) justify the use of VM stress to predict failure with a textbook on machine component design (Juvinall & Marshek, 2011). In contrast, studies comparing failure criteria in models of human femora found that maximum principal strain, not VM stress, best predicted the location of and load required for fracture (Schileo, Taddei, Cristofolini *et al.*, 2008; Yosibash, Tal & Trabelsi, 2010). This criterion remains in use in the medical community as the most effective method of predicting fracture risk in patients (Dahan, Trabelsi, Safran *et al.*, 2019). This discrepancy likely originates in the use of a failure criteria for machine components, abiotic metallic objects, to describe the behaviour of bones, living composite ceramic (*sensu* Carter & Norton, 2007) objects. The textbook takes into account two forms of failure in machine parts, plastic distortion and fracture (Juvinall *et al.*, 2011 pg. 250). Unlike metal, when bones deform plastically they are often capable of quick repair and light use during rest (Vogel, 2013 pg. 342). Therefore, fracture is the more critical component of bone failure. The work of fracture of commercial steel (Tattersall & Tappin, 1966), as one might find in machine components, is nearly 20 times that of bovine femora and still an order of magnitude beyond that of even impact-resistant antler (Currey, 1999). This means that once cracks form in bone they propagate much more readily than in steel. Here it is proposed that principal strain of finite element models best predicts failure in bone because surficial cracks in bone, indicated by extreme principal strain at the surface, propagate readily to the point of failure under normal loading conditions. This allows principal strain-based FEA (Figure 5B) to predict areas of weakness more precisely (Schileo *et al.*, 2008) and with greater validity (Yosibash *et al.*, 2010) in bones. For these reasons we suggest studies going forward to evaluate the strength of models based on differences in maximum principal strain, not VM stress. The FEA software Abaqus (Dassault Systèmes, France), Ansys (Ansys, Inc., USA), COSMOSWorks (Dassault Systèmes, France), Optistruct (Altair Engineering, Inc., USA), Strand7 (Strand7 Pty. Ltd., Australia), and VOX-FE2 (Banglawala, Bethunel, Fagan *et al.*, 2015) are already capable of this.

It is worth noting that the maximum principal strain criterion may decrease in validity with decreasing body mass, due to the decreasing importance of fracture in smaller organisms. Work of fracture is a relationship between work [energy] and surface area. The effective work done by an animal and the surface area created by breaks scale differently with size. In an isometric scenario one would expect work output to increase linearly with mass [ $m^1$ ]. It is the product of distance [i.e. length], which scales with  $m^{.33}$ , and force, which is known to scale with muscle cross-sectional area at  $m^{.66}$  (Froese, 2006). The surface area of a crack in bone, like any other surface (Froese, 2006), should scale with  $m^{.66}$ .

As work scales at a higher rate than the surface area it creates in cracks, one would expect cracks in bones to more easily propagate at higher body masses. Conversely at low body masses cracks propagate less readily relative to loading. Thus, in smaller organisms, the formation of cracks via principal strain of the surface may impose only weak selective pressure. McIntosh and Cox (2016 pg. 8) point out a similar trend in VM stress. Small animals working far from the yield strength of bone tend to experience selection towards mechanical efficiency of biting over minimising VM stress in the skull. Mechanical advantage analysis of lever models (see Section V.3.b Lever Modelling) may be more appropriate for dietary inferences in small animals where these pressures hold sway. What size range[s] this shift in selective pressure affects, potential allometric complications [e.g. phylogenetic (Wroe, McHenry & Thomason, 2005) or dietary (van der Meij & Bout, 2004) influences on scaling], and what other measures of feeding efficiency may be applicable [e.g. total strain energy as proposed by Dumont *et al.* (2009)] all warrant further study.

### (iii) Finite Element Analysis in Fossil Theropod Skulls

Contrary to the other methods described herein, FEA of the skull was first codified in non-avian theropods and only later applied to extant birds. Rayfield *et al.* (2001) performed FEA on a 3D model of the upper jaw of *Allosaurus fragilis*. Seeing the skull could withstand stresses greater than predicted bite forces, they proposed the animal fed by slashing its jaws at high velocity into prey. Rayfield (2004) was the first to perform 2D FEA and to utilise the now ubiquitous ‘heat’ maps to chart stress on a single skull. Her follow-up study (Rayfield, 2005) was the first to use FEA to compare stress distributions across taxa, as it is most commonly used today. Rayfield *et al.* (2007) are the first to combine FEA of extinct and extant taxa, comparing spinosaurid theropods and extant crocodilians. They are also the first to make dietary predictions based on FEA [piscivory in *Baryonyx walkeri*]. Shychoski and Snively (2008a) are the first to publish preliminary FEA results on lower jaws, finding the mandibles of adult tyrannosaurids to be more resistant to stress than that of juvenile tyrannosaurids or non-tyrannosaurids. These results are not currently in preparation for full publication (Eric Snively pers. com. 2020). Torices *et al.* (2018) performed FEA on individual theropod teeth. They found *Troodon* teeth experienced higher stresses than other taxa when loaded nonoptimally [i.e. at an angle other than the scratches observed in dental microwear]. This is interpreted as its teeth being poorly-adapted for struggling prey, and thus *Troodon* more likely fed on plant matter and/or small animals (Torices *et al.*, 2018). Lautenschlager *et al.* (2013) are the first to study both the upper and lower jaws of a theropod, finding the lower jaw to display higher VM stress in all loading conditions than the upper jaw. They are also first to incorporate reconstructed rhamphothecae into fossil FEA studies and to apply postcranial forces to the skull, proposing deconstruction powered by the cervical muscles compensated for low bite forces. This was expanded on in the follow-up study (Lautenschlager, 2017) in which loadings were varied in orientation to compare skulls’ adaptation for specific feeding styles. This technique forgoes direct modelling of cervical action and instead simply investigates the resultant forces the jaw would experience in contact with food. Miller *et al.* (In Press) are the first to incorporate a true fossilised rhamphotheca into an FE model. They are also the first to apply the technique to non-avian avialans, finding *Confuciusornis sanctus* to be most similar to a modern sally-striking predator or general herbivore. Cost *et al.* (2019) present the most complicated dinosaur FE models to date, with beams given ligament properties to connect bones. These were used to compare the skull of *T. rex* to that of an extant avian and lepidosaur. This study was concerned with presence of absence of cranial kinesis, but the increased accuracy of reconstruction utilised by (Cost *et al.*, 2019) in a manner like (Lautenschlager, 2017; Rayfield *et al.*, 2007) could show promise for illuminating dietary preferences and feeding strategies.

Two FEA studies on the skulls of fossil avians have also been undertaken. Degrange *et al.* (2010) compared the stress distribution in the upper jaw of *Andalgalornis steulleti* to those of *Haliaeetus albicilla* and *Cariama cristata*. They found *Andalgalornis* to experience the lowest stresses in models of pull-back feeding, analogous to extant accipiters. Attard *et al.* (2016) compared five genera of moa [Dinornithiformes] to two extant ratites in a variety of loading conditions to see which they were most well-suited to. They found that loadings with the lowest stress reflected observed feeding styles of the extant ratites, and that those of the moa match consumulite evidence previously reported (Attard *et al.*, 2016 their Figures S6 and S7).

#### (iv) Finite Element Analysis in Extant Bird Skulls

The first use of FEA to examine extant birds is from Oda *et al.* (2006), studying the shock absorption of the woodpecker skull. Numerous FEAs have been performed on woodpeckers, reviewed by (Jung, 2019), but none address the topic of diet. Herrel *et al.* (2010b) and Soons *et al.* (2010) performed similar FE analyses on the upper jaws of Darwin's finches. Both found that finches that ate harder foods experienced lower peak stresses, and even that taxa known to feed using the tip or base of the beak experienced lower peak stresses when loaded there. Subsequent studies of Darwin's finches (Soons *et al.*, 2015) support these initial results and emphasise the role of keratin in stress dissipation and the necessity of including it in FE models. All other applications of FEA to extant birds have been validation studies performed on a Darwin's finch (Soons, Lava, Debruyne *et al.*, 2012b), toucan (Seki, Mackey & Meyers, 2012), and ostrich (Cuff, Bright & Rayfield, 2015; Rayfield, 2011a). The Darwin's finch study is qualitatively evaluated as having a "fairly good correspondence" between the model and physical specimen (Soons *et al.*, 2012b pg. 190). The same is said of the toucan despite the *ex vivo* stress/strain curves provided appearing disjointed from those predicted by the model (Figure 7 in Seki *et al.*, 2012). Those of the ostrich (Cuff *et al.*, 2015; Rayfield, 2011a) are more thorough in their criticism. In short, they determine that strain patterns [e.g. areas of lower and higher strain and the range of strains experienced] are reflected in FEA but absolute strain and strain angles are not, particularly in the cranial region of the skull. The conclusion these studies support, whether directly or by interpretation of their data in hindsight, is that FEA is effective at showing stress/strain distributions in bird skulls and reflecting dietary habits. However, modelling limitations prevent them from providing any absolute information about *in vivo* strain states. Similar conclusions when testing extant crocodilians (Porro, Holliday, Anapol *et al.*, 2011; Reed, Porro, Iriarte-Diaz *et al.*, 2011; Sellers, Middleton, Davis *et al.*, 2017) and mammals (Bright & Rayfield, 2011; Godinho, Toro-Ibacache, Fitton *et al.*, 2017; Kupczik, Dobson, Fagan *et al.*, 2007) imply this is true across amniotes.

#### (v) Finite Element Analysis in Extant Bird Claws

Birn-Jeffery and Rayfield (2009) are the first to apply FEA to bird claws, and report preliminary success with discriminating between both locomotor and predator/nonpredator categories with 2D FEA. A full study has not yet been published. Tsang *et al.* (2019) were the first to analyse 3D FE models of bird claws. FEA revealed differences in strength that, while not easily pinpointed as variables in geometric morphometric analysis, were diagnostic of the behaviours associated with raptorial predation (Csermely *et al.*, 1998b; Goslow, 1972; Ward *et al.*, 2002). The broad scope and robust results of this study are promising for detection of raptorial behaviour in fossil species. Unfortunately, its application may be limited by the large number of fossil birds known only from compression fossils.

## 1. Robotic Modelling

While not a form of FEA, the modelling techniques of Backus *et al.* (2015) more strongly resemble FEA than any other techniques commonly applied in palaeontology. Their technique is herein dubbed ‘robotic modelling’, due to its original application in construction of robotic hands (Backus *et al.*, 2014; Dollar *et al.*, 2011). Their approach models non-ungual phalanges as beams and ungual phalanges as semicircles, with actuators acting as the digital flexor tendons. They evaluate models based on the tensional force exerted in order to maintain a grip with given parameters. Backus *et al.* (2015) are primarily concerned with the differences in actuation between passerine and non-passerine feet [i.e. passerine birds have digital flexors inserted distally and proximally while non-passerines have only distal digital flexors]. They find both on average to be equally well-equipped for grasping but passerine actuation to be uniquely well-adapted for minimising forces required to perch. While used to compare perching and grasping behaviours in this study, one can easily see the expansion of this technique into more granular studies. Future avenues include comparing adaptations for raptorial behaviour in which prey is completely encircled in the toes [e.g. owls] and held in an open grip [e.g. accipiters] (Fowler *et al.*, 2009), with these results compared to those of fossil taxa.

### (vi) Finite Element Analysis in Fossil Theropod Claws

Birn-Jeffery and Rayfield (2009) incorporated maniraptorans - including the early-diverging avialan *Archaeopteryx* - into their 2D FEA analysis of bird claws. They only report results for *Archaeopteryx*, with stress regimes aligning with those of arboreal perching taxa. Manning *et al.* (2009) were the first full study to apply 3D FEA to a theropod claw. They tested an earlier hypothesis based on a practical model (Manning, Payne, Pennicott *et al.*, 2006) that dromaeosaurids utilised their recurved second digit unguals for climbing rather than tearing through flesh. Creating fixed points in a model of a *Velociraptor* claw to replicate use during climbing, the team found the claw to experience levels of stress well below the yield strength of bone under a loading of the estimated body mass of *Velociraptor*. Thus, they proposed the animal could have supported its body weight on the claw during climbing (Manning *et al.*, 2009). Unfortunately, validation studies of FEA in animals find that only patterns of stress/strain distribution, not absolute values, are predicted by FEA (Bright *et al.*, 2011; Cuff *et al.*, 2015; Rayfield, 2011a; Stansfield, Parker & O'Higgins, 2018). As such, this interpretation is called into doubt. Furthermore, it is unclear how their climbing load simulation differs from a theoretical simulation of slashing [both would be loaded at a point slightly proximal to the claw apex and in a direction subparallel to the chord of the claw arc]. This work, then, is regarded as inconclusive in its palaeobiological reconstruction. The work of Lautenschlager (2014) on therizinosaurian unguals provides a superior framework for modelling claw use. Manual claws were distinctly loaded as if digging, hooking and pulling, or piercing substrate. The lowest VM stresses were seen in piercing simulations in all but *Alxasaurus elesitaiensis*. While not linked directly to diet in the study, behavioural optimisation of claws may provide information regarding niche and food available [e.g. fossorial adaptations likely indicate consumption of arthropods (Smith, 1982) or tubers (Andersen, 1987)].

### (vii) Discussion

Reconstructions of non-avian avialan skulls are recommended to remain in 2D for the time being (see Section V.1 Skull Reconstruction), and so FE models will have to remain 2D as well. However, indications of significant lateroflexion in the neck (see Section II.4.a Landmark-Based Cervical Reconstructions in Fossil Theropods) would suggest that 2D models will not fully capture typical loading. Analysis of these models should, for reasons described above, incorporate planar strain assumptions and

examine principal strain as a failure criterion using the intervals method (Marcé-Nogué *et al.*, 2017). However, strength may be less strongly selected for than efficiency in smaller animals (Dumont *et al.*, 2014; McIntosh *et al.*, 2016). If true, total strain energy may better reflect the efficiency of jaws (Dumont *et al.*, 2009) and thus selection for a given diet. With that said, strength-based FEA of Darwin's finches, smaller than many non-avian avialans, was able to provide clear dietary signal (Herrel *et al.*, 2010b; Soons *et al.*, 2010). Size concerns, then, may be irrelevant in avialans. It is therefore suggested here that both principal strain and total strain energy be compared to determine which best explains diet among small extant avian taxa. The superior metric can then be applied to fossil avialans. Finally, images of extant avian skulls should ideally be taken as radiographs and modelled with their original keratin thickness so as to model the effects of the rhamphotheca (Soons, Herrel, Genbrugge *et al.*, 2012a). However the precise thickness of the rhamphotheca seems to have little effect on stress/strain distributions or magnitudes in bird mandibles (Figure 7). So, in lieu of radiographs, surface photographs may be used for model construction as well (Miller *et al.*, In Press). When possible, fossilised rhamphothecae (see Section II.4 Soft Tissue) should be modelled directly in FE models of extinct taxa (Miller *et al.*, In Press). Otherwise, hypothetical rhamphothecae can be crafted for extinct taxa as in *Erlikosaurus* (Lautenschlager, Witmer, Altangerel *et al.*, 2014b) with refinement by subsequent studies of beak shape in relation to the underlying bone (Button, 2018; Miller *et al.*, In Press; Urano, Tanoue, Matsumoto *et al.*, 2018). Note the sensitivity analyses of Lautenschlager (2017) and Soons *et al.* (2012a) find keratin inclusion to affect stress and strain magnitudes but not patterns. Thus, comparisons within a skull, e.g. comparing various theoretical loadings, should not need to incorporate rhamphothecae. They may also then be unnecessary when comparing beaked avialans to one another, and only needed when toothed avialans are examined.

Cranial kinesis plays a major role in feeding in Neognathae (Bhullar *et al.*, 2016; Bout & Zweers, 2001; Zusi, 1984; absent in palaeognaths Gussekloo & Bout, 2005) and thus excluding it will undoubtedly alter the principal strain or total strain energy modelled from the *in vivo* conditions. Conversely, levels of kinesis similar to those in Neognathae are believed to have never been reached in non-neognathous avialans (Bhullar *et al.*, 2016; Hu *et al.*, 2019) with the possible exception of *Gobipteryx* and *Ichthyornis* (Hu *et al.*, 2019 pg. 19576). Incorporating connective structures to allow kinesis to present itself as in (Cost *et al.*, 2019) will in theory increase the validity of models, though attempts to incorporate them into the skull of an enantiornithine have proven troublesome. Jointing of the skull imposes unreasonable dislocation of the jugal, quadrate, and quadratojugal (Figure 6A–B). This may be an inherent flaw either with 2D simplification of the skull or the reconstruction itself, though sensitivity analyses showed a considerable influence of the cross-section and Young's modulus used for connective tissue in the model (Figure 6C). This issue can only be addressed with further research of suture and ligament physical properties. Avoiding kinesis by loading models of the upper jaw posterior to the bending zone of extant birds would mean loading at the maxilla/jugal contact in many taxa (Figure 1 and Table 1 in Zusi, 1984), well cranial to the tooth row in most toothed avialans. The exclusion of kinesis in Darwin's finches seemed to have no major repercussions on FEA reflecting diet (Herrel *et al.*, 2010b; Soons *et al.*, 2010) but this likely stems from similar levels of cranial kinesis in all studied groups. To avoid issues in modelling kinesis, comparison between the akinetic lower jaws of taxa is recommended here as a simple solution. While studies including both jaws in FEA (Adams *et al.*, 2019; Attard *et al.*, 2014; Attard *et al.*, 2016; Lautenschlager *et al.*, 2013; Moreno *et al.*, 2008; Wroe, 2008; Wroe *et al.*, 2007) have recovered differing peak VM stresses between the jaws and different stress



distributions [thus making models of both jaws ideal] the information has not conflicted in terms of dietary interpretation.

Application of FEA for determining raptorial use of the pes is promising after the results of Tsang *et al.* (2019). However, because their predatory groupings differed markedly on dorsal and lateral curvatures (their Figure S2) [not accounted for in 2D FEA and not known in compression fossils], larger 2D FEA studies should be preceded by a comparative analysis of 2D and 3D FE models (see introduction of Section V.3.a Introduction). The reported success of Birn-Jeffery and Rayfield (2009) shows promise for their correspondence, but a lack of published data renders a full proof-of-concept necessary. Refinement of raptorial use type to more precise grips can be achieved by modifying the work of Backus *et al.* (2015). A pilot study is necessary, though, to determine in extant raptors if their method works using the curvature of ungual bones rather than claw sheathes. If not, reconstructions of keratin sheathes are necessary [from avialans preserving impressions of the sheathe, e.g. AGB-6997 (Wang, Huang, Kundrát *et al.*, 2020b), DNHM D2945/6 (Chiappe, Zhao, O'Connor *et al.*, 2014), GMV-2130 (Chiappe *et al.*, 1999), GSGM-05-CM-004 (O'Connor, Li, Lamanna *et al.*, 2016a), IVPP V18687 (Hu, O'Connor & Zhou, 2015)]. While Lautenschlager (2014) provides an outline for modelling the effectiveness of varying claw uses, the use of claws directly in acquiring food is rare among extant avians. Piscivorous raptors are known to pierce fish with their talons in order to maintain grip (Fowler *et al.*, 2009 pg. 7), so claw FEA may prove useful to test the hypothesis of similar habits in bohaiornithids (Wang *et al.*, 2014c). Digging incorporating the claws plays a major role in foraging in Megapodidae (Friedmann, 1931) and a more minor role in shorebirds (Jacobs, 1982). Beyond these cases, claws tend to play a minor role in prey manipulation compared to the whole toes (Clark, 1973; Csermely *et al.*, 1998b; Sustaita *et al.*, 2013), at most increasing traction (Backus *et al.*, 2015; Fowler *et al.*, 2009; Ramos & Walker, 1998) or elongating the toes to increase grasp reach (Csermely, Bertè & Camoni, 1998a; Fowler *et al.*, 2009). While general strength trends appear useful in parsing raptorial behaviour, the lack of functional importance of claws outside specialised groups renders more specific loading comparisons a lower priority in lieu of established hypotheses to test.

#### (4) Complications Applicable across Physical Approaches

While eating is a major part of an organism's survival, a myriad of unrelated factors play into the form and function of body parts. These factors are confounding variables in any palaeodietary reconstruction. Some factors can be corrected for, theoretically negating their influence, but all must be kept in mind when interpreting data from methods described herein. Table 4 provides a convenient reference for which approaches a given complication must be accounted for.

##### (a) Phylogenetic Signal

It is easy to see how phylogeny can affect shape. Genetics is one of the principal factors controlling shape, and thus in the absence of selection one would assume that the more similar the genetics of two organisms the more similar they will appear (Blomberg, Garland & Ives, 2003). At large enough phylogenetic scales, developmental pathways may even completely prohibit an optimised form, or predispose two organisms to find different functionally optimised forms (Gould, 2002). Thus, morphometric comparisons at any timescale over which evolution is a factor should undergo phylogenetic corrections (de Bello, Berg, Dias *et al.*, 2015). Uyeda *et al.* (2018) review how to effectively craft hypotheses so that phylogenetic corrections can be made. Adams and Collyer (2018) review the

mathematical methods and assumptions required for these corrections. Guillaume *et al.* (2020) provide general guidance for the timing of corrections and potential pitfalls of certain evolutionary models.

Over broad enough phylogenetic scales, homologous structures may eventually become functionally incomparable. This is particularly important in the theropod pes due to the evolution of flight. Among coelurosaurian theropods, grasping ability of the manus generally decreases the later a taxon diverges (Hutchinson & Allen, 2008). The proximally-fused metacarpals of *Sapeornis* and later-diverging avialans (Fig. 4 in Pittman *et al.*, 2020a) likely placed similar constraints of manipulation on them as in living avians, meaning the two groups would have had relied similarly on the pes for grasping. The more refined manual manipulation of some early-diverging paravians (Senter, 2006) means they faced weaker selective pressures for using the pes as a device for manipulation. In this example a comparison of pedal proportions of function among Jeholornithiformes and pygostylians should give meaningful data, but one at the paravian level may not.

#### (b) Allometric Signal

Different measures of organisms [such as length, surface area, and volume] have different dimensionality. This means they grow at different rates relative to one another. Most organisms would cease to function if they grew isometrically [i.e. with every part scaling the same way as every other part] both through ontogeny and evolution. Thus, nearly all organisms display some form of allometry [i.e. certain components scaling at different rates than others] (Froese, 2006). In other words, body size has an inherent effect on body shape. Klingenberg (2016) reviews methods to quantify and correct for this effect in morphometric studies. Functional effects of allometry have almost exclusively been derived empirically and are known to vary phylogenetically (van der Meij *et al.*, 2004; Wroe *et al.*, 2005). Thus, allometry presents a more confounding influence in functional studies. In addition to influencing the strength or efficiency of structures inherently, size may also determine if strength or efficiency is more strongly selected for (see Section V.3.c.ii The Strength Criterion).

#### (c) Many-To-One Mapping

Coined by Wainwright *et al.* (2005), many-to-one mapping describes the ability of systems with different forms to perform the same mechanical function. This means that elements which morphometrics classifies as very different may be operating essentially the same in practice. This aligns with the observation above that GM tends to provide mixed results in feeding studies. GM creates a level of abstraction between the data and how the animal interacts with the world. Identifying a case of many-to-one mapping [via comparison of morphometric and functional studies] can be useful. It can formulate hypotheses of evolutionary constraints on form (Button *et al.*, 2020; Ungar & Hlusko, 2016) or illuminate interference of common behavioural signals [e.g. herbivorous and carnivorous taxa engaging in similar bouts of intrasexual competition for mates, or diving predators and foragers facing similar pressures from long-term submersion].

#### (d) Liem's Paradox

Occasionally referred to as one-to-many mapping, Liem's paradox was originally coined to describe the peculiarity that a set of cichlid fishes with highly specialised jaws seemed to have no particular dietary specialty (Liem, 1980). The prevailing explanation has been that some specialist morphologies minimally inhibit acquiring 'easy' resources while aiding in obtaining 'difficult' resources [commonly referred to as 'fallback foods'] when others are scarce (Robinson & Wilson, 1998). Recent ecological evidence supports this theory (Golcher-Benavides & Wagner, 2019; Lambert, Chapman, Wrangham *et al.*, 2004; Wiseman,

Greene, Koo *et al.*, 2019). Essentially, this means that organisms that appear morphologically specialised for a certain diet cannot be precluded from being generalists [or specialists feeding on a different easy-to-acquire food source] overall. This is also the reason dental microwear studies emphasise large sample sizes for analysis, to capture signs of rare but important resource use (Ungar, 2018).

#### (e) Integration and Modularity

Put simply, integration is when otherwise distinct parts of an organism function and/or evolve as one unit. Modularity is when an organism displays distinct regions [modules] within which integration is high and between which integration is low (Klingenberg, 2014). Farina *et al.* (2019) provide more rigorous definitions and a review of the concepts.

Both extreme modularity and extreme integration can encourage diversification (Hedrick, Mutumi, Munteanu *et al.*, 2019b). In highly modular systems parts are free to evolve independently from one another. This should, in theory, increase adaptation to environmental changes and thus speciation. Highly integrated systems, on the other hand, restrict parts to evolving as a single unit. While limiting form to a single spectrum, it can allow for rapid diversification and speciation along that spectrum (Hedrick *et al.*, 2019b). However, each does not encourage diversification on the same scale. Modularity is associated with diversification overall in high-level clades [class- to subclass-level] (Felice *et al.*, 2018; Felice, Watanabe, Cuff *et al.*, 2019b; Hu, Ghigliotti, Vacchi *et al.*, 2016; Marroig, Shirai, Porto *et al.*, 2009) and at smaller scales in several evolutionary circumstances (Collar, Wainwright, Alfaro *et al.*, 2014; Drake & Klingenberg, 2008; Tokita, Kiyoshi & Armstrong, 2007; Young, Wagner & Hallgrímsson, 2010). Integration, in contrast, is known to lead to diversification when lineages invade habitats with a preponderance of unoccupied niches (Hedrick *et al.*, 2019b; Hu *et al.*, 2016; Navalón *et al.*, 2020).

Integration and modularity are of most concern in a morphometric context. Integration limits disparity to a single continuum (Hedrick *et al.*, 2019b; Navalón *et al.*, 2020). So, highly integrated structures are likely to cluster or spread on a single axis of shape. Modularity also can help prioritise functional studies. More modular structures, with more ability to create unique geometries, are more likely to exhibit many-to-one mapping. Thus, they should be checked for functional convergence with higher priority. Adams and Felice (2014) and Adams (2016) provide techniques for quantifying integration and modularity. Because of the complicated relationship they have with evolutionary trajectories no universal correction for their effects has yet been proposed.

#### (f) Behavioural Signals

##### (i) Grooming

The influence of grooming behaviour on morphology has been studied more thoroughly in birds than any other group (Bush & Clayton, 2018). Some anatomical work has been done on specialised dental (Asher, 1998; Gingerich & Rose, 1979; Rose, Walker & Jacobs, 1981) and manual (Bishop, 1962; Dunkel, 2019; Koenigswald, Habersetzer & Gingerich, 2011; Maiolino, Boyer & Rosenberger, 2011) grooming in primates in addition. In birds, destruction of ectoparasites is known to be aided by a short bill (Clayton & Cotgreave, 1994) with a rostral hook (Bush, Villa, Boves *et al.*, 2012; Clayton, Moyer, Bush *et al.*, 2005; Clayton & Walther, 2001). Intraspecific differences in bill shape are known to reflect ectoparasite load in communities (Bardwell, Benkman & Gould, 2001; Moyer, Peterson & Clayton, 2002). Some birds also have a pectinate [“comb”] claw with distinct serrations on the lateral surface thought to play a role in grooming. Studies of its effect, however, have yielded mixed results (Bush *et al.*, 2012; Clayton, Koop, Harbison *et al.*, 2010). It is worth noting that, of these variables, only bill length is always reflected in the

animal's skeletal morphology. The same is not true of mammalian toothcombs. Toothcombs convergently evolved in lemurs, flying lemurs, tree shrews, and the arctocyonid *Thryptacodon* (Rose *et al.*, 1981). In toothcombs the mandibular incisors and, variably, canines are deflected rostrally. During grooming they are brushed perpendicular to hair shafts to aid in ectoparasite removal (Gingerich *et al.*, 1979; Rose *et al.*, 1981; Schwartz, 1978). This role is augmented or replaced by a specialised, more robust and recurved grooming claw [or, 'toilet claw'] in non-simian primates and several polyphyletic simians (Koenigswald *et al.*, 2011; Maiolino *et al.*, 2011). Manual grooming has been reported thoroughly in simian primates, but almost exclusively in social (Schino, 2006; Xia, Kyes, Wang *et al.*, 2019) or spatial (Dunbar, 1991; Freeland, 1981) contexts, rather than its effect on morphology. Opposable thumbs have been proposed as the product of selective pressures related to manual grooming (Bishop, 1962), though this theory remains only one of many (Dunkel, 2019).

From these trends in birds and mammals, oral grooming may be expected to select for perpendicularly-oriented structures in the rostrum [beak hook, tooth comb] while manual grooming can be accomplished with more diverse structures [pectinate claw, grooming claw, possibly opposable thumbs]. Each of these structures accompanies grooming behaviour with stresses distinct from that of feeding. In turn, structures experience selective pressure to resist those stresses. To our knowledge only circumstantial notes have been made on how significant these stresses may be. There are notes of beak overhangs breaking (Bush *et al.*, 2012) and enamel microscopically wearing (Rose *et al.*, 1981) from grooming activities. Because wear patterns on tooth combs differ from those of the surrounding teeth, unique wear in teeth has been proposed as indicative of a grooming function in the dromaeosaurid *Saurornitholestes* (Currie *et al.*, 2019).

#### (ii) Thermoregulation

The size of structures is commonly assumed to relate to their strength or speed, but thermoregulation is known to influence the scale of elements. Joel Asaph Allen famously noted that animals living in warmer climates tend to have larger extremities and *vice versa* (Allen, 1877 pg. 112–119), a trend now referred to as Allen's rule. The rule is supported by several quantitative studies (Alho, Herczeg, Laugen *et al.*, 2011; Greenberg, Cadena, Danner *et al.*, 2012; Nudds & Oswald, 2007; Symonds & Tattersall, 2010; Tilkens, Wall-Scheffler, Weaver *et al.*, 2007; *contra* Stevenson, 1986). More dramatic structures such as goat horns (Taylor, 1966), elephant ears (Phillips & Heath, 1992), and toucan bills (Tattersall, Andrade & Abe, 2009) have all been proposed as tools for active thermoregulation. The size of body parts should then be interpreted with the climate inhabited by the organism in mind, and hypertrophy of body parts [particularly when not accompanied by significant increases in structural strength] may be a sign of active thermoregulation rather than dietary pressure.

#### (iii) Sensation

Sensory systems are paramount in both feeding (Montuelle *et al.*, 2019) and reproduction (Ptacek, 2000) in living animals. Thus, selective pressure on sensation can be expected to dramatically shape organisms. For instance, enhanced mechanoreception is often associated with elongation and extensive pitting of the skull in amniotes in general (Morhardt, 2009) and birds in particular (Cunningham *et al.*, 2013). In the same vein, an increase of the size of amniote eyes often creates a corresponding reduction in bite force (Fortuny, Marcé-Nogué, De Esteban-Trivigno *et al.*, 2011; Henderson, 2002). Sensory specialisations that parallel those in extant taxa can be understood and tested for by understanding the biology of those taxa. The possibility of novel forms of sensory augmentation present only in extinct taxa, however, renders sensation a true confounding variable.

#### (iv) Sexual Display

Animals have a variety of tools for communication and competition that improve survival and reproductive success. All of these may alter the body in unpredictable ways. Protuberances may serve as intersexual signals (Mayr, 2018) or as bases for intrasexual combat (Clutton-Brock, 1982; Rico-Guevara & Araya-Salas, 2014). Changes in skull morphology can lead to changes in vocalisation (Giraudeau, Nolan, Black *et al.*, 2014; Herrel *et al.*, 2009; Huber & Podos, 2006) or altered ability to detect chemical messages (Rouquier & Giorgi, 2007). Sexual dimorphism itself can lead to functional differences on small (Verwajen, Van Damme & Herrel, 2002) or large (Pietsch, 2005) scales. Sexual dimorphism may be able to be detected in the fossil record given an adequate number of samples (Plavcan, 1994), but other forms of sexual display can drive shape and affect function with little evidence left in the fossil record. Thus, sexual display also remains a true confounding variable.

#### (5) Discussion

The broad application of physical approaches across fossil taxa makes them ideal for comparative frameworks, but not all frameworks are appropriate approaches to dietary reconstruction. Neither TM nor GM of theropod skulls appears particularly effective at isolating features of diet across phylogenetically diverse groups. In addition, the lack of consistent landmarks in studies highlights how few measurements there are of the theropod skull that would have an intuitive effect on dietary choice. Both forms of morphometrics are more effective at revealing diet when applied to the pes. This is likely because the pes performs a smaller variety of roles in the organism (Montuelle *et al.*, 2019) and those other roles are often in service of obtaining food (Fowler *et al.*, 2009; Sustaita *et al.*, 2013). Results have been obtained with both TM and GM analysis of extant avian feet, and so neither is recommended over the other by this review. TM may also prove useful to find dietary signal in teeth due to the large disparity seen across theropods as a whole (Hendrickx *et al.*, 2014; Hendrickx *et al.*, 2020) and Avialae in particular (Huang *et al.*, 2016; O'Connor, 2019; O'Connor *et al.*, 2011b). A lack of prior study in this field leaves this recommendation only tentative. Lizards are the closest related extant taxon with similarly disparate teeth, and quantitative analyses of lizard teeth have already identified characters indicative of durophagy (Estes *et al.*, 1984) and herbivory (Melstrom, 2017) that could be applied to toothed avialans.

In general, this review recommends functional studies over morphometrics due to the fewer complications that influence them (see Table 4). 2D simplifications of functional models appear necessary for the time being, but if possible, comparisons of 2D and 3D models should be undertaken to confirm the former's validity. Whether more valid dietary signal can be gleaned from lever models of the upper jaw combined with functional indices from (Button *et al.*, 2020; Ma *et al.*, 2020) (Figure 4) or principal strain-based FEA of the lower jaw is still unclear. The size at which selection for strength transitions to selection for mechanical efficiency (Adams *et al.*, 2019), if such a transition truly exists, is not yet established. Ideally, a study including both approaches could directly compare the two. Separate studies, each focusing on one functional approach, should have results that can be compared nearly as effectively. Incorporating connective tissue into the skull to create kinetic structures is not recommended pending more precise understandings of the physical properties of said connective tissues. Discovery of fossilised rhamphothecae allows for them to be included in models, as is suggested here.

## VI. The Framework and Current Knowledge of Non-avian Avialan Diet

In sum, our recommended approaches in Sections II–V combine into a framework for narrowing fossil avialan diet. Figure 8 provides a summary of the techniques described herein, the body parts which need

to be preserved for their use, expected results from their application, and interpretation of said results. A general workflow for applying the framework is provided in Figure 9, from determining specimens of interest to synthesizing test results into a dietary assignment. We provide an example of applying our framework to the extant golden eagle (*Aquila chrysaetos*), a raptorial vertebrate predator. Figure 10 provides a graphical summary of this example. While *Aquila chrysaetos* does not have teeth, if it did they should have high surface anisotropy from the repeated tearing of tough meat. Its pull-back method of prey disassembly should select for enlarged attachment sites for dorsiflexive muscles, and finite element models of its lower jaw should have muscular loadings deflected cranially. Its bones should be depleted in  $^{44}\text{Ca}$  and its proteins enriched in  $^{15}\text{N}$  due to its high trophic level. Its body mass should be above 300g due to vertebrate predation; this is true even in unusually small subspecies (Watson, 2010 pg. 33). Its talons should be highly recurved with a hypertrophied DI due to their raptorial use; this has been reported (Fowler *et al.*, 2009). Its jaws should have a low MA, high tooth recurvature if it had teeth, and high values of the other recommended indices as it hunts agile prey; its jaw MA is low among avians (Navalón *et al.*, 2018a). Its lower jaw should experience relatively high principal strains when loaded in FEA due to the compliance of animal flesh. Finally, its talons should experience relatively low principal strains when loaded in FEA due to their raptorial use; this has been supported (Tsang *et al.*, 2019). Given the current studies which fit into our framework, we would say there are four lines of evidence which would lead to classifying the golden eagle as a raptorial vertebrate predator were it extinct.

While limited by a dearth of quantitative studies, our framework can be used to establish what we currently know of non-avian avialan diet. The scansoriopterygids [possibly early-diverging avialans or non-avian pennaraptorans (Pittman *et al.*, 2020a)], *Epidexipteryx* and *Yi* appear adapted for carnivory due to low values of mechanical advantage and most of the functional indices recommended herein (Ma *et al.*, 2020). Body mass estimates above 300g in *Yi* (Dececchi, Roy, Pittman *et al.*, Accepted) specifically point to it being a vertivore. To our knowledge, no quantitative study of anchiornithine [possibly early-diverging avialans or troodontids (Pittman *et al.*, 2020a)] diet has been performed. Preserved fish and lizard meals are known in *Anchiornis* (Zheng, Wang, Sullivan *et al.*, 2018b), but it is unclear if this represents a typical part of its diet. Evidence of diet in the earliest-diverging unequivocal avialan *Archaeopteryx* appears contradictory. Studies show dental microwear reminiscent of invertivores (Bestwick *et al.*, 2018), a body mass in the range of vertivores (Serrano *et al.*, 2015), and relatively high jaw MA (Navalón, 2014) expected in herbivores or durophages. The most likely source of this contradiction is the specimens studied. Bestwick (2018) studied microwear of the Munich specimen (Jordan Bestwick pers. com. 2020); Serrano *et al.* (2015) measured the London, Berlin, Eichstätt, and Thermopolis specimens; and the work of Navalón (2014) was likely based on the reconstruction of (Rauhut, 2014), which was in turn based primarily on the Eichstätt specimen. Considerable morphological disparity has been previously noted within *Archaeopteryx* (Rauhut, Foth & Tischlinger, 2018) which may yet indicate diverse diets within the genus. Alternatively, the contradictory evidence may be indicative of Liem's Paradox at work, with fallback food[s] not captured in the small sample size of dental microwear (see Lambert *et al.*, 2004 for a modern equivalent). Five preserved seed meals are currently known in the early-diverging avialan *Jeholornis* (O'Connor *et al.*, 2019; Zhou & Zhang, 2002b), but it is unknown if other major parts of its diet remain unaccounted for. Due to a sparsity of quantitative data, especially the seemingly contradictory evidence in *Archaeopteryx*, the ancestral dietary condition in avialans remains unclear.

The early-diverging pygostylian *Confuciusornis* has jaws with both strength (Miller *et al.*, In Press) and MA (Navalón, 2014) consistent with herbivorous avians (Miller *et al.*, In Press; Navalón, 2014). One study applying pedal morphometrics (Cobb *et al.*, 2020) recovered *Confuciusornis* as raptorial, and its body mass estimates are consistent with vertivory (Serrano *et al.*, 2015; Table 2). However, because the morphometric study measured only curvature of unguals and not their relative sizes, we consider that raptorial behaviour is not ruled out, but is not confirmed. This discrepancy can be directly addressed with stable isotope analysis, for which *Confuciusornis* is a prime candidate due to the large number of specimens known. Its close relative *Eoconfuciusornis* has only had MA measurements taken (Navalón, 2014), which we consider inadequate for dietary assignment. No dietary study has been conducted on any member of Jinguoformisidae to our knowledge. We also consider the single line of quantitative MA evidence (Navalón, 2014) favouring granivory in the early pygostylian *Sapeornis* to be inadequate for diet assignment, though it does agree with previously-reported ingested material (O'Connor, 2019; O'Connor *et al.*, 2019; Zheng *et al.*, 2011).

Among the enantiornithine ornithothoracines, *Shenqiornis* is tentatively proposed here as predatory due to low jaw MA (Navalón, 2014) and raptorial pes morphometrics in its close relatives (Wang *et al.*, 2014c). Its MA is of particular interest for future studies to attempt to replicate due to qualitative assertions of durophagy in the taxon (Wang *et al.*, 2010b). Navalón (2014) additionally reports intermediate values of MA for *Pengornis* and an indeterminate hatchling and low MA for *Rapaxavis*. Again, we consider this single line of evidence inadequate for dietary assignment. One specimen of *Zhouornis* [BMNHC Ph 756] has been reported as having claws as straight as modern ground birds (Cobb *et al.*, 2020), which would rule out raptorial behaviour in the taxon. However the claw measured, DIII, is aberrantly straight in this genus with the other claws highly recurved (Zhang, O'Connor, Di *et al.*, 2014; Zhang, Chiappe, Han *et al.*, 2013) so we do not consider raptorial behaviour ruled-out. The holotype of *Eoalulavis* preserves part of a crustacean in its stomach (Sanz *et al.*, 1996), but the lack of a skull or feet in the specimen inhibit investigation of the typical diet of this taxon within our framework.

Finally among non-avian ornithuromorphs, MA values reported in (Navalón, 2014) are congruent with direct evidence of granivory in *Hongshanornis* (Chiappe *et al.*, 2014) and piscivory in *Yanornis* (Zheng *et al.*, 2014; Zhou, Clarke, Zhang *et al.*, 2004). Additionally, *Eogranivora* and *Piscivoravis* have preserved meals that provide evidence of granivory (Zheng *et al.*, 2018a) and piscivory (Zhou, Zhou & O'Connor, 2014a) respectively, but determining whether these were normal parts of their diet requires further study.

The paucity of dietary assignments renders trends in avialan dietary evolution murky. Our framework supports a mixture of carnivory and herbivory/omnivory among both early-diverging non-avian avialans [*Archaeopteryx*, *Confuciusornis*, *Sapeornis*] and later-diverging ones [*Shenqiornis*, *Hongshanornis*, *Yanornis*]. Therefore, no particular macroevolutionary trends are currently clearly apparent. Dietary diversity seems to increase through time, but this is a preservational bias associated with the predominance of data from the Early Cretaceous Jehol Lagerstätte [all of these taxa save for *Archaeopteryx*]. This also means that relatively little geographic and climatic range is accounted for among currently known non-avian avialan diets. Thus, non-avian avialan material from a wider range of localities should also be prioritized for future study.

## VII. Conclusions

- (1) Our aim was to build a framework for studying non-avian avialan diet by reviewing techniques that have proven effective in both avians and non-avian theropods and the use this to summarise our current state of knowledge. With this framework in place, we hope this will generate progress in the reconstruction of Mesozoic ecosystems and in our understanding of the ecological history of birds.
- (2) Figure 8 provides a convenient summary of the techniques discussed in this review and our recommendations for applying them. Expected outcomes and their general interpretations are also provided. In short, we recommend combining direct evidence of diet, stable isotope geochemistry, body mass estimation, pes morphometrics, and functional analysis to obtain multiple lines of evidence relevant to diet.
- (3) Due to a dearth of quantitative studies, current knowledge of non-avian avialan diet is sparse. The ancestral avialan diet remains obscure, in large part due to contradictory evidence concerning the diet of *Archaeopteryx*. Both carnivory and herbivory/omnivory are present in early-diverging [*Archaeopteryx*, *Confuciusornis*, *Sapeornis*] and later-diverging [*Shenqiornis*, *Hongshanornis*, *Yanornis*] avialans, but no trends in the dietary evolution of non-avian avialans have presented themselves. We believe that new avialan specimens from a wider range of localities covering different geographies and climates will be instrumental to elucidating these trends in the future.
- (4) Our review demonstrates the need for establishing links between diet and morphology in avians, reconstructing the often-crushed remains of non-avian avialans, and combining the two in robust quantitative frameworks. Combined with a growing understanding of modern ecology, these will provide a new and exciting picture of earth during some of the most ground-breaking evolutionary transitions known.

## VIII. References

- ABEL R. L., LAURINI, C. R. & RICHTER, M. (2012). A palaeobiologist's guide to 'virtual' micro-CT preparation. *Palaeontologia Electronica* **15**, 6T.
- ABOURACHID A., FABRE, A. C., CORNETTE, R. & HOFLING, E. (2017). Foot shape in arboreal birds: two morphological patterns for the same pincer-like tool. *Journal of Anatomy* **231**, 1-11.
- ADAMÍK P. & KORNAN, M. (2004). Foraging ecology of two bark foraging passerine birds in an old-growth temperate forest. *Ornis Fennica* **81**, 13-22.
- ADAMS D. C. (2016). Evaluating modularity in morphometric data: challenges with the RV coefficient and a new test measure. *Methods in Ecology and Evolution* **7**, 565-572.
- ADAMS D. C. & COLLYER, M. L. (2018). Multivariate phylogenetic comparative methods: evaluations, comparisons, and recommendations. *Systematic Biology* **67**, 14-31.
- ADAMS D. C. & FELICE, R. N. (2014). Assessing trait covariation and morphological integration on phylogenies using evolutionary covariance matrices. *PLOS ONE* **9**, e94335.
- ADAMS D. C. & OTAROLA-CASTILLO, E. (2013). geomorph: an r package for the collection and analysis of geometric morphometric shape data. *Methods in Ecology and Evolution* **4**, 393-399.
- ADAMS N. F., RAYFIELD, E. J., COX, P. G., COBB, S. N. & CORFE, I. J. (2019). Functional tests of the competitive exclusion hypothesis for multituberculate extinction. *Royal Society Open Science* **6**, 181536.



- ALHO J. S., HERCZEG, G., LAUGEN, A. T., RÄSÄNEN, K., LAURILA, A. & MERILÄ, J. (2011). Allen's rule revisited: quantitative genetics of extremity length in the common frog along a latitudinal gradient. *Journal of Evolutionary Biology* **24**, 59-70.
- ALLEN J. A. (1877). The influence of physical conditions in the genesis of species. *Radical Review* **1**, 108-140.
- ALONSO P. D., MILNER, A. C., KETCHAM, R. A., COOKSON, M. J. & ROWE, T. B. (2004). The avian nature of the brain and inner ear of *Archaeopteryx*. *Nature* **430**, 666-669.
- AMiot R., LÉCUYER, C., BUFFETAUT, E., ESCARGUEL, G., FLUTEAU, F. & MARTINEAU, F. (2006). Oxygen isotopes from biogenic apatites suggest widespread endothermy in Cretaceous dinosaurs. *Earth and Planetary Science Letters* **246**, 41-54.
- AMiot R., WANG, X., LÉCUYER, C., BUFFETAUT, E., BOUDAD, L., CAVIN, L., DING, Z. L., FLUTEAU, F., KELLNER, A. W. A., TONG, H. Y. & ZHANG, F. S. (2010). Oxygen and carbon isotope compositions of middle Cretaceous vertebrates from North Africa and Brazil: ecological and environmental significance. *Palaeogeography, Palaeoclimatology, Palaeoecology* **297**, 439-451.
- AMiot R., WANG, X., ZHOU, Z., WANG, X., LÉCUYER, C., BUFFETAUT, E., FLUTEAU, F., DING, Z., KUSUHASHI, N. & MO, J. (2015). Environment and ecology of East Asian dinosaurs during the Early Cretaceous inferred from stable oxygen and carbon isotopes in apatite. *Journal of Asian Earth Sciences* **98**, 358-370.
- ANDERSEN D. C. (1987). Below-ground herbivory in natural communities: a review emphasizing fossorial animals. *The Quarterly Review of Biology* **62**, 261-286.
- ANGST D., AMiot R., BUFFETAUT, E., FOUREL, F., MARTINEAU, F., LAZZERINI, N. & LÉCUYER, C. (2015). Diet and climatic context of giant birds inferred from  $\delta^{13}\text{C}$  and  $\delta^{18}\text{O}$  values of Late Palaeocene and Early Eocene eggshells from southern France. *Palaeogeography, Palaeoclimatology, Palaeoecology* **435**, 210-221.
- ANGST D., LÉCUYER, C., AMiot R., BUFFETAUT, E., FOUREL, F., MARTINEAU, F., LEGENDRE, S., ABOURACHID, A. & HERREL, A. (2014). Isotopic and anatomical evidence of an herbivorous diet in the Early Tertiary giant bird *Gastornis*. Implications for the structure of Paleocene terrestrial ecosystems. *Naturwissenschaften* **101**, 313-322.
- ARCHIMEDES OF SYRACUSE. (1897). On the Equilibrium of Planes, Books I and II. In *The Works of Archimedes* (ed. T. L. Heath), pp. 189-220. Cambridge University Press, London, UK.
- ARMAN S. D., UNGAR, P. S., BROWN, C. A., DESANTIS, L. R. G., SCHMIDT, C. & PRIDEAUX, G. J. (2016). Minimizing inter-microscope variability in dental microwear texture analysis. *Surface Topography: Metrology and Properties* **4**, 024007.
- ASHER R. J. (1998). Morphological diversity of anatomical strepsirrhinism and the evolution of the lemuriform toothcomb. *American Journal of Physical Anthropology* **105**, 355-367.
- ATTARD M. R. G., PARR, W. C. H., WILSON, L. A. B., ARCHER, M., HAND, S. J., ROGERS, T. L. & WROE, S. (2014). Virtual reconstruction and prey size preference in the Mid Cenozoic thylacinid, *Nimbacinus dicksoni* (Thylacinidae, Marsupialia). *PLOS ONE* **9**, e93088.
- ATTARD M. R. G., WILSON, L. A. B., WORTHY, T. H., SCOFIELD, P., JOHNSTON, P., PARR, W. C. H. & WROE, S. (2016). Moa diet fits the bill: virtual reconstruction incorporating mummified remains and prediction of biomechanical performance in avian giants. *Proceedings of the Royal Society B: Biological Sciences* **283**, 20152043.
- ATTERHOLT J., HUTCHISON, J. H. & O'CONNOR, J. K. (2018). The most complete enantiornithine from North America and a phylogenetic analysis of the Avisauridae. *PeerJ* **6**, e5910.

- BACKUS S. B., ODHNER, L. U. & DOLLAR, A. M. (2014). Design of hands for aerial manipulation: actuator number and routing for grasping and perching. In *2014 IEEE/RSJ International Conference on Intelligent Robots and Systems*, pp. 34-40. IEEE.
- BACKUS S. B., SUSTAITA, D., ODHNER, L. U. & DOLLAR, A. M. (2015). Mechanical analysis of avian feet: multiarticular muscles in grasping and perching. *Royal Society Open Science* **2**.
- BAILLEUL A. M., O'CONNOR, J., ZHANG, S. K., LI, Z. H., WANG, Q., LAMANNA, M. C., ZHU, X. F. & ZHOU, Z. H. (2019). An Early Cretaceous enantiornithine (Aves) preserving an unlaidd egg and probable medullary bone. *Nature Communications* **10**, 1275.
- BAKER A. J. & MOEED, A. (1979). Evolution in the introduced New Zealand populations of the common myna, *Acridotheres tristis* (Aves: Sturnidae). *Canadian Journal of Zoology* **57**, 570-584.
- BAKKER R. T. (1998). Brontosaur killers: Late Jurassic allosaurids as sabre-tooth cat analogues. *Gaia* **15**, 145-158.
- BAKKER R. T. & WILLIAMS, M. (1988). *Nanotyrannus*, a new genus of pygmy tyrannosaur, from the latest Cretaceous of Montana. *Hunteria* **1**, 1.
- BANGLAWALA N., BETHUNEL, I., FAGAN, M. & HOLBREY, R. (2015). Voxel-based finite element modelling with VOX-FE2. *ARCHER White Papers*.
- BARBOSA A. & MORENO, E. (1999). Evolution of foraging strategies in shorebirds: an ecomorphological approach. *The Auk: Ornithological Advances* **116**, 712-725.
- BARDWELL E., BENKMAN, C. W. & GOULD, W. R. (2001). Adaptive geographic variation in Western Scrub-Jays. *Ecology* **82**, 2617-2627.
- BARRETT P. M. (2005). The diet of ostrich dinosaurs (Theropoda: Ornithomimosauria). *Palaeontology* **48**, 347-358.
- BARRICK R. E. & SHOWERS, W. J. (1994). Thermophysiology of *Tyrannosaurus rex*: evidence from oxygen isotopes. *Science* **265**, 222-224.
- BARRICK R. E., SHOWERS, W. J. & FISCHER, A. G. (1996). Comparison of thermoregulation of four ornithischian dinosaurs and a varanid lizard from the Cretaceous Two Medicine Formation: evidence from oxygen isotopes. *Palaaios*, 295-305.
- BARTON N. W. H. & HOUSTON, D. C. (1992). The influence of gut morphology on digestion time in raptors, University of Glasgow.
- BATHE K.-J. (2014). *Finite Element Procedures*, 2 edition. Bathe, Klaus-Jürgen, Watertown, MA, USA.
- BECHTLE S., FETT, T., RIZZI, G., HABELITZ, S., KLOCKE, A. & SCHNEIDER, G. A. (2010). Crack arrest within teeth at the dentinoenamel junction caused by elastic modulus mismatch. *Biomaterials* **31**, 4238-4247.
- BELL A. & CHIAPPE, L. M. (2016). A species-level phylogeny of the Cretaceous Hesperornithiformes (Aves: Ornithuromorpha): implications for body size evolution amongst the earliest diving birds. *Journal of Systematic Palaeontology* **14**, 239-251.
- BELL A. & CHIAPPE, L. M. (2020). Anatomy of *Paraheperornis*: evolutionary mosaicism in the Cretaceous Hesperornithiformes (Aves). *Life* **10**, 62.
- BERGQVIST L. P. (2003). The role of teeth in mammal history. *Brazilian Journal of Oral Sciences* **2**, 249-257.
- BESTWICK J., UNWIN, D. M., BUTLER, R. J., HENDERSON, D. H. & PURNELL, M. A. (2018). Dental microwear textural analysis: reconstructing diets of non-mammalian fossil taxa from the Solnhofen archipelago. In *The Palaeontological Association 62nd Annual Meeting*, pp. 68-69, University of Bristol, Bristol, UK.

- BESTWICK J., UNWIN, D. M. & PURNELL, M. A. (2019). Dietary differences in archosaur and lepidosaur reptiles revealed by dental microwear textural analysis. *Scientific Reports* **9**, 1-11.
- BHULLAR B. A. S., HANSON, M., FABBRI, M., PRITCHARD, A., BEVER, G. S. & HOFFMAN, E. (2016). How to make a bird skull: major transitions in the evolution of the avian cranium, paedomorphosis, and the beak as a surrogate hand. *Integrative and Comparative Biology* **56**, 389-403.
- BIRN-JEFFERY A. V., MILLER, C. E., NAISH, D., RAYFIELD, E. J. & HONE, D. W. E. (2012). Pedal claw curvature in birds, lizards and mesozoic dinosaurs - Complicated categories and compensating for mass-specific and phylogenetic control. *PLOS ONE* **7**, e50555.
- BIRN-JEFFERY A. V. & RAYFIELD, E. (2009). Finite element analysis of pedal claws to determine mode of life in birds, lizards and maniraptoran theropods. In *Sixty-Ninth Annual Meeting Society of Vertebrate Paleontology and the Fifty-Seventh Symposium of Vertebrate Palaeontology and Comparative Anatomy (SVPCA)* vol. 29, pp. 64A-64A. Society of Vertebrate Paleontology.
- BISHOP A. (1962). Control of the hand in lower primates. *Annals of the New York Academy of Sciences* **102**, 316-337.
- BLOMBERG S. P., GARLAND, T. & IVES, A. R. (2003). Testing for phylogenetic signal in comparative data: behavioral traits are more labile. *Evolution* **57**, 717-745.
- BLUM J. D., TALIAFERRO, E. H. & HOLMES, R. T. (2001). Determining the sources of calcium for migratory songbirds using stable strontium isotopes. *Oecologia* **126**, 569-574.
- BONHOMME V., PICQ, S., GAUCHEREL, C. & CLAUDE, J. (2014). Momocs: outline analysis using R. *Journal of Statistical Software* **56**, 1-24.
- BONSER R. H. C. & WITTER, M. S. (1993). Indentation hardness of the bill keratin of the European starling. *The Condor: Ornithological Applications*, 736-738.
- BOUT R. G. & ZWEERS, G. A. (2001). The role of cranial kinesis in birds. *Comparative Biochemistry and Physiology - Part A: Molecular & Integrative Physiology* **131**, 197-205.
- BRETT-SURMAN M. K. & PAUL, G. S. (1985). A new family of bird-like dinosaurs linking Laurasia and Gondwanaland. *Journal of Vertebrate Paleontology* **5**, 133-138.
- BRIGGS D. E. G., WILBY, P. R., PEREZ-MORENO, B. P., SANZ, J. L. & FREGENAL-MARTÍNEZ, M. (1997). The mineralization of dinosaur soft tissue in the Lower Cretaceous of Las Hoyas, Spain. *Journal of the Geological Society* **154**, 587-588.
- BRIGHT J. A. (2014). A review of paleontological finite element models and their validity. *Journal of Paleontology* **88**, 760-769.
- BRIGHT J. A., MARUGÁN-LOBÓN, J., COBBE, S. N. & RAYFIELD, E. J. (2016). The shapes of bird beaks are highly controlled by nondietary factors. *Proceedings of the National Academy of Sciences of the United States of America* **113**, 5352-5357.
- BRIGHT J. A., MARUGÁN-LOBÓN, J., RAYFIELD, E. J. & COBB, S. N. (2019). The multifactorial nature of beak and skull shape evolution in parrots and cockatoos (Psittaciformes). *BMC Evolutionary Biology* **19**, 104.
- BRIGHT J. A. & RAYFIELD, E. J. (2011). Sensitivity and ex vivo validation of finite element models of the domestic pig cranium. *Journal of Anatomy* **219**, 456-471.
- BRINK K. S., CHEN, Y. C., WU, Y. N., LIU, W. M., SHIEH, D. B., HUANG, T. D., SUN, C. K. & REISZ, R. R. (2016). Dietary adaptations in the ultrastructure of dinosaur dentine. *Journal of the Royal Society Interface* **13**.
- BROWN J. (2009). Choosing the right number of components or factors in PCA and EFA. *JALT Testing & Evaluation SIG Newsletter* **13**.

- BRUSATTE STEPHEN L., O'CONNOR, JINGMAI K. & JARVIS, ERICH D. (2015). The origin and diversification of birds. *Current Biology* **25**, R888-R898.
- BRUSATTE S. L., SAKAMOTO, M., MONTANARI, S. & HARCOURT SMITH, W. E. H. (2012). The evolution of cranial form and function in theropod dinosaurs: insights from geometric morphometrics. *Journal of Evolutionary Biology* **25**, 365-377.
- BUFFETAUT E. & ANGST, D. (2016). The giant flightless bird *Gargantuavis philoinos* from the Late Cretaceous of southwestern Europe: a review. In *Cretaceous Period: Biotic Diversity and Biogeography New Mexico Museum of Natural History and Science Bulletin* (ed. A. Khosla and S. G. Lucas), pp. 45-50. New Mexico Museum of Natural History and Science Albuquerque, NM, USA.
- BÜHLER P., MARTIN, L. D. & WITMER, L. M. (1988). Cranial kinesis in the Late Cretaceous birds Hesperornis and Parahesperornis. *The Auk: Ornithological Advances* **105**, 111-122.
- BURGER A. E. A. (1978). Functional anatomy of the feeding apparatus of four South African cormorants. *African Zoology* **13**, 81-102.
- BUSH S. E. & CLAYTON, D. H. (2018). Anti-parasite behaviour of birds. *Philosophical Transactions of the Royal Society B: Biological Sciences* **373**.
- BUSH S. E., VILLA, S. M., BOVES, T. J., BREWER, D. & BELTHOFF, J. R. (2012). Influence of bill and foot morphology on the ectoparasites of barn owls. *Journal of Parasitology* **98**, 256-261.
- BUTTON D. J. & ZANNO, L. E. (2020). Repeated evolution of divergent modes of herbivory in non-avian dinosaurs. *Current Biology* **30**, 158-168. e4.
- BUTTON K. A. (2018). Soft Tissue Reconstruction and Ecomorphology of Beaks in Extant and Extinct Theropod Dinosaurs, North Carolina State University.
- CAMBRA-MOO O., BUSCALIONI, Á. D., CUBO, J., CASTANET, J., LOTH, M.-M., DE MARGERIE, E. & DE RICQLÈS, A. (2006). Histological observations of enantiornithine bone (Saurischia, Aves) from the Lower Cretaceous of Las Hoyas (Spain). *Comptes Rendus Palevol* **5**, 685-691.
- CANDEIRO C. R. A., CURRIE, P. J., CANDEIRO, C. L. & BERGQVIST, L. P. (2017). Tooth wear and microwear of theropods from the Late Maastrichtian Marília Formation (Bauru Group), Minas Gerais State, Brazil. *Earth and Environmental Science Transactions of the Royal Society of Edinburgh* **106**, 229-233.
- CARLISLE J. D. & HOLBERTON, R. L. (2006). Relative efficiency of fecal versus regurgitated samples for assessing diet and the deleterious effects of a tartar emetic on migratory birds. *Journal of Field Ornithology* **77**, 126-135.
- CARLOS C. J., ALVARENGA, J. G. & MAZZOCHI, M. S. (2017). Osteology of the feeding apparatus of Magnificent Frigatebird *Fregata magnificens* and Brown Booby *Sula leucogaster* (Aves: Suliformes). *Papéis Avulsos de Zoologia* **57**, 265-274.
- CARNEY R., KAPLAN, H., KIRK, A., BAINES, A. & MASON, S. (2018). *Archaeopteryx holographica*: bringing the urvogel back to life with scientific animation and VR/AR. In *Society of Vertebrate Paleontology 78th Annual Meeting*, pp. 102, Albuquerque, New Mexico, U.S.A.
- CARRANO M. T., JANIS, C. M. & SEPKOSKI, J. J. (1999). Hadrosaurs as ungulate parallels: lost life styles and deficient data. *Acta Palaeontologica Polonica* **44**, 237-261.
- CARTER C. B. & NORTON, M. G. (2007). Ceramics in Biology and Medicine. In *Ceramic Materials: Science and Engineering*, vol. 716 (ed. C. B. Carter and M. G. Norton), pp. 659-676. Springer-Verlag New York, New York City, NY, USA.

- CARVALHO I. D. S., NOVAS, F. E., AGNOLÍN, F. L., ISASI, M. P., FREITAS, F. I. & ANDRADE, J. A. (2015). A new genus and species of enantiornithine bird from the Early Cretaceous of Brazil. *Brazilian Journal of Geology* **45**, 161-171.
- CAU A. & ARDUINI, P. (2008). *Enantiophoenix electrophyla* gen. et sp. nov. (Aves, Enantiornithes) from the Upper Cretaceous (Cenomanian) of Lebanon and its phylogenetic relationships. *Atti della Società Italiana di Scienze Naturali e del Museo Civico di Storia Naturale di Milano* **149**, 293-324.
- CAVIEDES-VIDAL E., MCWHORTER, T. J., LAVIN, S. R., CHEDIAK, J. G., TRACY, C. R. & KARASOV, W. H. (2007). The digestive adaptation of flying vertebrates: high intestinal paracellular absorption compensates for smaller guts. *Proceedings of the National Academy of Sciences* **104**, 19132-19137.
- CHAMBERLAIN C., BLUM, J., HOLMES, R. T., FENG, X., SHERRY, T. & GRAVES, G. R. (1996). The use of isotope tracers for identifying populations of migratory birds. *Oecologia* **109**, 132-141.
- CHÁVEZ-HOFFMEISTER M. (2020). Bill disparity and feeding strategies among fossil and modern penguins. *Paleobiology*, 1-17.
- CHIAPPE L. M. (2002). Osteology of the Flightless *Patagopteryx deferrariisi* from the Late Cretaceous of Patagonia (Argentina). In *Mesozoic Birds: Above the Heads of Dinosaurs* (ed. L. M. Chiappe and L. M. Witmer), pp. 281-316. University of California Press, Berkeley and Los Angeles, CA, USA.
- CHIAPPE L. M. & CALVO, J. O. (1994). *Neuquenornis volans*, a new Late Cretaceous bird (Enantiornithes: Avisauridae) from Patagonia, Argentina. *Journal of Vertebrate Paleontology* **14**, 230-246.
- CHIAPPE L. M., DI, L., SERRANO, F. J., YUGUANG, Z. & MENG, Q. (2019a). Anatomy and flight performance of the early enantiornithine bird *Protopteryx fengningensis*: information from new specimens of the Early Cretaceous Huajiyang Formation of China. *The Anatomical Record: Advances in Integrative Anatomy and Evolutionary Biology*, 1-16.
- CHIAPPE L. M. & MENG, Q. (2016). *Birds of Stone: Chinese Avian Fossils from the Age of Dinosaurs*. John Hopkins University Press, Baltimore, MD, USA.
- CHIAPPE L. M., NORELL, M. & CLARK, J. (2001). A new skull of *Gobipteryx minuta* (Aves: Enantiornithes) from the Cretaceous of the Gobi Desert. *American Museum Novitates* **1**, 1-16.
- CHIAPPE L. M., QINGJIN, M., SERRANO, F., SIGURDSEN, T., MIN, W., BELL, A. & DI, L. (2019b). New *Bohaiornis*-like bird from the Early Cretaceous of China: enantiornithine interrelationships and flight performance. *PeerJ* **7**, e7846.
- CHIAPPE L. M., SHU'AN, J., QIANG, J. & NORELL, M. A. (1999). Anatomy and systematics of the Confuciusornithidae (Theropoda : Aves) from the late mesozoic of northeastern China. *Bulletin of the American Museum of Natural History*, 3-89.
- CHIAPPE L. M. & WALKER, C. A. (2002). Skeletal Morphology and Systematics of the Cretaceous Euenantiornithes (Ornithothoraces: Enantiornithes). In *Mesozoic Birds: Above the Heads of Dinosaurs* (ed. L. M. Chiappe and L. M. Witmer), pp. 240-267. University of California Press, Berkeley and Los Angeles, CA, USA.
- CHIAPPE L. M., ZHAO, B., O'CONNOR, J. K., GAO, C. L., WANG, X. R., HABIB, M., MARUGÁN-LOBÓN, J., MENG, Q. J. & CHENG, X. D. (2014). A new specimen of the Early Cretaceous bird *Hongshanornis longicresta*: insights into the aerodynamics and diet of a basal ornithuromorph. *PeerJ* **2**, e234.

- CHIEN C. H., WU, Y. D., CHAO, Y.-J., CHEN, T., CHEN, W. F., YU, J. C. & LI, X. (2008). The effects of different cranial modules on mechanical properties of cranial suture in Lewis Rats and same-aged C57BL/6 Mice. *Strain* **44**, 272-277.
- CHIN K., TOKARYK, T. T., ERICKSON, G. M. & CALK, L. C. (1998). A king-sized theropod coprolite. *Nature* **393**, 680.
- CHINSAMY-TURAN A., CHIAPPE, L. M., MARUGÁN-LOBÓN, J., GAO, C. L. & ZHANG, F. J. (2013). Gender identification of the Mesozoic bird *Confuciusornis sanctus*. *Nature Communications* **4**, 1381.
- CHRISTIANSEN P. & BONDE, N. (2004). Body plumage in Archaeopteryx: a review, and new evidence from the Berlin specimen. *Comptes Rendus Palevol* **3**, 99-118.
- CLARK G. A. (1973). Holding food with the feet in passerines. *Bird-Banding* **44**, 91-99.
- CLARK J., HOPSON, J., FASTOVSKY, D. & MONTELLANO, M. (1998). Foot posture in a primitive pterosaur. *Nature* **391**, 886.
- CLARKE J. A. (2004). Morphology, phylogenetic taxonomy, and systematics of *Ichthyornis* and *Apatornis* (Avialae: Ornithurae). *Bulletin of the American Museum of Natural History* **2004**, 1-179.
- CLARKE J. A. & NORELL, M. A. (2002). The morphology and phylogenetic position of *Apsaravis ukhaana* from the Late Cretaceous of Mongolia. *American Museum Novitates* **2002**, 1-46.
- CLARKE J. A., ZHOU, Z. & ZHANG, F. (2006). Insight into the evolution of avian flight from a new clade of Early Cretaceous ornithurines from China and the morphology of *Yixianornis grabaui*. *Journal of Anatomy* **208**, 287-308.
- CLAYTON D. H. & COTGREAVE, P. (1994). Relationship of bill morphology to grooming behavior in birds. *Animal Behaviour* **47**, 195-201.
- CLAYTON D. H., KOOP, J. A., HARBISON, C. W., MOYER, B. R. & BUSH, S. E. (2010). How birds combat ectoparasites. *Open Ornithology Journal* **3**, 41-71.
- CLAYTON D. H., MOYER, B. R., BUSH, S. E., JONES, T. G., GARDINER, D. W., RHODES, B. B. & GOLLER, F. (2005). Adaptive significance of avian beak morphology for ectoparasite control. *Proceedings of the Royal Society B: Biological Sciences* **272**, 811-817.
- CLAYTON D. H. & WALTHER, B. A. (2001). Influence of host ecology and morphology on the diversity of Neotropical bird lice. *Oikos* **94**, 455-467.
- CLEMENTZ M. T. (2012). New insight from old bones: stable isotope analysis of fossil mammals. *Journal of Mammalogy* **93**, 368-380.
- CLUTTON-BROCK T. H. (1982). The functions of antlers. *Behaviour* **79**, 108-124.
- COATHAM S. J., VINTHER, J., RAYFIELD, E. J. & KLUG, C. (2020). Was the Devonian placoderm *Titanichthys* a suspension feeder? *Royal Society Open Science* **7**, 200272.
- COBB S. E. & SELLERS, W. I. (2020). Inferring lifestyle for Aves and Theropoda: a model based on curvatures of extant avian ungual bones. *PLOS ONE* **15**, e0211173.
- COLLAR D. C., WAINWRIGHT, P. C., ALFARO, M. E., REVELL, L. J. & MEHTA, R. S. (2014). Biting disrupts integration to spur skull evolution in eels. *Nature Communications* **5**, 5505.
- CONNOLLY R. M., GUEST, M. A., MELVILLE, A. J. & OAKES, J. M. (2004). Sulfur stable isotopes separate producers in marine food-web analysis. *Oecologia* **138**, 161-167.
- COONEY C. R., BRIGHT, J. A., CAPP, E. J. R., CHIRA, A. M., HUGHES, E. C., MOODY, C. J. A., NOURI, L. O., VARLEY, Z. K. & THOMAS, G. H. (2017). Mega-evolutionary dynamics of the adaptive radiation of birds. *Nature* **542**, 344-347.
- CORBIN C. E. (2008). Foraging ecomorphology within North American flycatchers and a test of concordance with southern African species. *Journal of Ornithology* **149**, 83-95.

- CORBIN C. E., LOWENBERGER, L. K. & GRAY, B. L. (2015). Linkage and trade-off in trophic morphology and behavioural performance of birds. *Functional Ecology* **29**, 808-815.
- CORTÉS E. (1997). A critical review of methods of studying fish feeding based on analysis of stomach contents: application to elasmobranch fishes. *Canadian Journal of Fisheries and Aquatic Sciences* **54**, 726-738.
- COST I. N., MIDDLETON, K. M., SELLERS, K. C., ECHOLS, M. S., WITMER, L. M., DAVIS, J. L. & HOLLIDAY, C. M. (2019). Palatal biomechanics and its significance for cranial kinesis in *Tyrannosaurus rex*. *The Anatomical Record: Advances in Integrative Anatomy and Evolutionary Biology*.
- COX P. G. (2017). The jaw is a second-class lever in *Pedetes capensis* (Rodentia: Pedetidae). *PeerJ* **5**, e3741.
- CROFTS S. B. & SUMMERS, A. P. (2014). How to best smash a snail: the effect of tooth shape on crushing load. *Journal of the Royal Society Interface* **11**, 20131053.
- CSERMELY D., BERTÈ, L. & CAMONI, R. (1998a). Prey killing by Eurasian Kestrels: the role of the foot and the significance of bill and talons. *Journal of Avian Biology* **29**, 10-16.
- CSERMELY D. & GAIBANI, G. (1998b). Is foot squeezing pressure by two raptor species sufficient to subdue their prey? *The Condor: Ornithological Applications* **100**, 757-763.
- CSERMELY D. & ROSSI, O. (2006). Bird claws and bird of prey talons: where is the difference? *Italian Journal of Zoology* **73**, 43-53.
- CSERMELY D., ROSSI, O. & NASI, F. (2012). Comparison of claw geometrical characteristics among birds of prey and non-raptorial birds. *Italian Journal of Zoology* **79**, 410-433.
- CUFF A. R., BRIGHT, J. A. & RAYFIELD, E. J. (2015). Validation experiments on finite element models of an ostrich (*Struthio camelus*) cranium. *PeerJ* **3**, e1294.
- CUNNINGHAM S. J., CORFIELD, J. R., IWANIUK, A. N., CASTRO, I., ALLEY, M. R., BIRKHEAD, T. R. & PARSONS, S. (2013). The anatomy of the bill tip of kiwi and associated somatosensory regions of the brain: comparisons with shorebirds. *PLOS ONE* **8**, e80036.
- CURREY J. D. (1999). The design of mineralised hard tissues for their mechanical functions. *Journal of Experimental Biology* **202**, 3285-3294.
- CURRIE P. J. & EVANS, D. C. (2019). Cranial anatomy of new specimens of *Saurornitholestes langstoni* (Dinosauria, Theropoda, Dromaeosauridae) from the Dinosaur Park Formation (Campanian) of Alberta. *The Anatomical Record: Advances in Integrative Anatomy and Evolutionary Biology*.
- CURTIS N., JONES, M. E. H., EVANS, S. E., O'HIGGINS, P. & FAGAN, M. J. (2013). Cranial sutures work collectively to distribute strain throughout the reptile skull. *Journal of the Royal Society Interface* **10**, 20130442.
- CZERKAS S. A. & FEDUCCIA, A. (2014). Jurassic archosaur is a non-dinosaurian bird. *Journal of Ornithology* **155**, 841-851.
- D'AMORE D. C. (2009). A functional explanation for denticulation in theropod dinosaur teeth. *The Anatomical Record: Advances in Integrative Anatomy and Evolutionary Biology* **292**, 1297-1314.
- DAHAN G., TRABELSI, N., SAFRAN, O. & YOSIBASH, Z. (2019). Finite element analyses for predicting anatomical neck fractures in the proximal humerus. *Clinical Biomechanics* **68**, 114-121.
- DAL SASSO C. & SIGNORE, M. (1998). Exceptional soft-tissue preservation in a theropod dinosaur from Italy. *Nature* **392**, 383.
- DALLA VECCHIA F. M. & CHIAPPE, L. M. (2002). First avian skeleton from the mesozoic of northern Gondwana. *Journal of Vertebrate Paleontology* **22**, 856-860.

- DALSÄTT J., ERICSON, P. G. P. & ZHOU, Z. G. (2014). A New Enantiornithes (Aves) from the Early Cretaceous of China. *Acta Geologica Sinica (English Edition)* **88**, 1034-1040.
- DALSÄTT J., ZHOU, Z., ZHANG, F. & ERICSON, P. G. P. (2006). Food remains in *Confuciusornis sanctus* suggest a fish diet. *Naturwissenschaften* **93**, 444.
- DANIEL T. L., HELMUTH, B. S., SAUNDERS, W. B. & WARD, P. D. (1997). Septal complexity in ammonoid cephalopods increased mechanical risk and limited depth. *Paleobiology* **23**, 470-481.
- DAVENPORT S. R. & BAX, N. J. (2002). A trophic study of a marine ecosystem off southeastern Australia using stable isotopes of carbon and nitrogen. *Canadian Journal of Fisheries and Aquatic Sciences* **59**, 514-530.
- DAVIS M. T., LOYD, A. M., SHEN, H.-Y. H., MULROY, M. H., NIGHTINGALE, R. W., MYERS, B. S. & BASS, C. D. (2012). The mechanical and morphological properties of 6 year-old cranial bone. *Journal of Biomechanics* **45**, 2493-2498.
- DE BELLO F., BERG, M. P., DIAS, A. T. C., DINIZ, J. A. F., GOTZENBERGER, L., HORTAL, J., LADLE, R. J. & LEPS, J. (2015). On the need for phylogenetic 'corrections' in functional trait-based approaches. *Folia Geobotanica* **50**, 349-357.
- DECECCHI T. A. & LARSSON, H. C. (2011). Assessing arboreal adaptations of bird antecedents: testing the ecological setting of the origin of the avian flight stroke. *PLOS ONE* **6**, e22292.
- DECECCHI T. A., ROY, A., PITTMAN, M., KAYE, T. G., XU, X., HABIB, M. B., LARSSON, H. C. E., WANG, X. & ZHENG, X. (Accepted). Aerodynamics show membranous-winged theropods were a poor gliding dead-end. *iScience*.
- DEGRANGE F. J., TAMBUSSI, C. P., MORENO, K., WITMER, L. M. & WROE, S. (2010). Mechanical analysis of feeding behavior in the extinct "terror bird" *Andalgalornis steulleti* (Gruiformes: Phorusrhacidae). *PLOS ONE* **5**, e11856.
- DIMERY N. J., ALEXANDER, R. M. & DEYST, K. A. (1985). Mechanics of the ligamentum nuchae of some artiodactyls. *Journal of Zoology* **206**, 341-351.
- DOLLAR A. M. & HOWE, R. D. (2011). Joint coupling design of underactuated hands for unstructured environments. *The International Journal of Robotics Research* **30**, 1157-1169.
- DRAKE A. G. & KLINGENBERG, C. P. (2008). The pace of morphological change: historical transformation of skull shape in St Bernard dogs. *Proceedings of the Royal Society B: Biological Sciences* **275**, 71-76.
- DUFFY D. C. & JACKSON, S. (1986). Diet studies of seabirds: a review of methods. *Colonial Waterbirds*, 1-17.
- DUKE G. E. (1997). Gastrointestinal physiology and nutrition in wild birds. *Proceedings of the Nutrition Society* **56**, 1049-1056.
- DUMONT E. R., GROSSE, I. R. & SLATER, G. J. (2009). Requirements for comparing the performance of finite element models of biological structures. *Journal of Theoretical Biology* **256**, 96-103.
- DUMONT E. R., SAMADEVAM, K., GROSSE, I., WARSI, O. M., BAIRD, B. & DAVALOS, L. M. (2014). Selection for mechanical advantage underlies multiple cranial optima in new world leaf-nosed bats. *Evolution* **68**, 1436-1449.
- DUMONT M., TAFFOREAU, P., BERTIN, T., BHULLAR, B. A., FIELD, D., SCHULP, A., STRILISKY, B., THIVICHON-PRINCE, B., VIRIOT, L. & LOUCHARTE, A. (2016). Synchrotron imaging of dentition provides insights into the biology of *Hesperornis* and *Ichthyornis*, the "last" toothed birds. *BMC Evolutionary Biology* **16**.



- DUNBAR R. I. M. (1991). Functional significance of social grooming in primates. *Folia Primatologica* **57**, 121-131.
- DUNKEL A. (2019). The evolution of grooming and hand use in primates: an interdisciplinary perspective, pp. 1-34, unpublished preprint.
- EIMAR H., GHADIMI, E., MARELLI, B., VALI, H., NAZHAT, S. N., AMIN, W. M., TORRES, J., CIOBANU, O., JUNIOR, R. F. A. & TAMIMI, F. (2012). Regulation of enamel hardness by its crystallographic dimensions. *Acta Biomaterialia* **8**, 3400-3410.
- EINODER L. D. & RICHARDSON, A. M. M. (2007). Aspects of the hindlimb morphology of some Australian birds of prey: a comparative and quantitative study. *The Auk: Ornithological Advances* **124**, 773-788.
- EKLÖV P. & DIEHL, S. (1994). Piscivore efficiency and refuging prey: the importance of predator search mode. *Oecologia* **98**, 344-353.
- ELZANOWSKI A. (1974). Preliminary note on the palaeognathous bird from the Upper Cretaceous of Mongolia. *Acta Palaeontologia Polonica* **30**, 103-109.
- ELZANOWSKI A. (1977). Skulls of *Gobipteryx* (Aves) from the Upper Cretaceous of Mongolia. *Palaeontologia Polonica* **37**, 153-165.
- ELZANOWSKI A. (1981). Embryonic bird skeletons from the Late Cretaceous of Mongolia. *Palaeontologia Polonica* **42**, 147-176.
- ELZANOWSKI A. (1991). New observations on the skull of *Hesperornis* with reconstructions of the bony palate and otic region. *Postilla* **207**, 1-20.
- ELZANOWSKI A. (2001a). A new genus and species for the largest specimen of Archaeopteryx. *Acta Palaeontologica Polonica* **46**.
- ELZANOWSKI A. (2001b). A novel reconstruction of the skull of Archaeopteryx. *Netherlands Journal of Zoology* **51**, 207-215.
- ELZANOWSKI A. & GALTON, P. M. (1991). Braincase of *Enaliornis*, an early Cretaceous bird from England. *Journal of Vertebrate Paleontology* **11**, 90-107.
- ELZANOWSKI A., PETERS, D. S. & MAYR, G. (2018). Cranial morphology of the Early Cretaceous bird *Confuciusornis*. *Journal of Vertebrate Paleontology* **38**, e1439832.
- ERICKSON G. M., GIGNAC, P. M., STEPPAN, S. J., LAPPIN, A. K., VLIET, K. A., BRUEGGEN, J. D., INOUE, B. D., KLEDZIK, D. & WEBB, G. J. W. (2012). Insights into the ecology and evolutionary success of crocodilians revealed through bite-force and tooth-pressure experimentation. *PLOS ONE* **7**, e31781.
- ERICKSON G. M., SIDEBOTTOM, M. A., KAY, D. I., TURNER, K. T., IP, N., NORELL, M. A., SAWYER, W. G. & KRICK, B. A. (2015). Wear biomechanics in the slicing dentition of the giant horned dinosaur *Triceratops*. *Science Advances* **1**, e1500055.
- ESQUE T. C. & PETERS, E. L. (1994). Ingestion of bones, stones, and soil by desert tortoises. *Fish and Wildlife Research* **13**, 73-84.
- ESTES R. & WILLIAMS, E. E. (1984). Ontogenetic variation in the molariform teeth of lizards. *Journal of Vertebrate Paleontology* **4**, 96-107.
- FALK A., O'CONNOR, J., WANG, M. & ZHOU, Z. (2019). On the preservation of the beak in *Confuciusornis* (Aves: Pygostylia). *Diversity* **11**, 212.
- FALK A. R. (2014). Foot and hindlimb morphology, soft tissues, and tracemaking behaviors of early Cretaceous birds from China and the Republic of Korea with a comparison to modern avian morphology and behavior, University of Kansas.

- FALK A. R., KAYE, T. G., ZHOU, Z. H. & BURNHAM, D. A. (2016). Laser fluorescence illuminates the soft tissue and life habits of the Early Cretaceous bird *Confuciusornis*. *PLOS ONE* **11**, e0167284.
- FARIA P. J., GUEDES, N. M. R., YAMASHITA, C., MARTUSCELLI, P. & MIYAKI, C. Y. (2008). Genetic variation and population structure of the endangered Hyacinth Macaw (*Anodorhynchus hyacinthinus*): implications for conservation. *Biodiversity and Conservation* **17**, 765-779.
- FARINA S. C., KANE, E. A. & HERNANDEZ, L. P. (2019). Multifunctional structures and multistructural functions: integration in the evolution of biomechanical systems. *Integrative and Comparative Biology* **59**, 338-345.
- FARKE A. A. (2008). Frontal sinuses and head-butting in goats: a finite element analysis. *Journal of Experimental Biology* **211**, 3085-3094.
- FARKE A. A., CHOK, D. J., HERRERO, A., SCOLIERI, B. & WERNING, S. (2013). Ontogeny in the tube-crested dinosaur *Parasaurolophus* (Hadrosauridae) and heterochrony in hadrosaurids. *PeerJ* **1**, e182.
- FARLOW J. O. & BRINKMAN, D. L. (1994). Wear surfaces on the teeth of *Tyrannosaurs*. *The Paleontological Society Special Publications* **7**, 165-176.
- FARMER A., RYE, R., LANDIS, G., BERN, C., KESTER, C. & RIDLEY, I. (2003). Tracing the pathways of Neotropical migratory shorebirds using stable isotopes: a pilot study. *Isotopes in Environmental and Health Studies* **39**, 169-177.
- FEDUCCIA A. (1993). Evidence from claw geometry indicating arboreal habits of *Archaeopteryx*. *Science* **259**, 790-793.
- FELICE R. N. & GOSWAMI, A. (2018). Developmental origins of mosaic evolution in the avian cranium. *Proceedings of the National Academy of Sciences of the United States of America* **115**, 555-560.
- FELICE R. N., TOBIAS, J. A., PIGOT, A. L. & GOSWAMI, A. (2019a). Dietary niche and the evolution of cranial morphology in birds. *Proceedings of the Royal Society B: Biological Sciences* **286**, 20182677.
- FELICE R. N., WATANABE, A., CUFF, A. R., NOIRAUT, E., POL, D., WITMER, L. M., NORELL, M. A., O'CONNOR, P. M. & GOSWAMI, A. (2019b). Evolutionary integration and modularity in the archosaur cranium. *Integrative and Comparative Biology* **59**, 371-382.
- FERNANDEZ BLANCO M. V., CASSINI, G. H. & BONA, P. (2018). Skull ontogeny of extant caimans: a three-dimensional geometric morphometric approach. *Zoology* **129**, 69-81.
- FIELD D. J., BENITO, J., CHEN, A., JAGT, J. W. M. & KSEPKA, D. T. (2020). Late Cretaceous neornithine from Europe illuminates the origins of crown birds. *Nature* **579**, 397-401.
- FIELD D. J., BERCOVICI, A., BERV, J. S., DUNN, R., FASTOVSKY, D. E., LYSON, T. R., VAJDA, V. & GAUTHIER, J. A. (2018a). Early Evolution of Modern Birds Structured by Global Forest Collapse at the End-Cretaceous Mass Extinction. *Current Biology* **28**, 1825-1831.
- FIELD D. J., HANSON, M., BURNHAM, D., WILSON, L. E., SUPER, K., EHRET, D., EBERSOLE, J. A. & BHULLAR, B.-A. S. (2018b). Complete *Ichthyornis* skull illuminates mosaic assembly of the avian head. *Nature* **557**, 96.
- FIGUEIRIDO B., MACLEOD, N., KRIEGER, J., DE RENZI, M., PEREZ-CLAROS, J. A. & PALMQVIST, P. (2011). Constraint and adaptation in the evolution of carnivorous skull shape. *Paleobiology* **37**, 490-518.
- FISHER H. I. (1946). Adaptations and comparative anatomy of the locomotor apparatus of New World vultures. *The American Midland Naturalist* **35**, 545-727.

- FOGEL M. L. & TUROSS, N. (1999). Transformation of plant biochemicals to geological macromolecules during early diagenesis. *Oecologia* **120**, 336-346.
- FORTELIUS M. & SOLOUNIAS, N. (2000). Functional characterization of ungulate molars using the abrasion-attrition wear gradient: a new method for reconstructing paleodiets. *American Museum Novitates* **2000**, 1-36.
- FORTUNY J., MARCÉ-NOGUÉ, J., DE ESTEBAN-TRIVIGNO, S., GIL, L. & GALOBART, A. (2011). Temnospondyli bite club: ecomorphological patterns of the most diverse group of early tetrapods. *Journal of Evolutionary Biology* **24**, 2040-2054.
- FOTH C. & RAUHUT, O. W. M. (2013). Macroevolutionary and morphofunctional patterns in theropod skulls: a morphometric approach. *Acta Palaeontologica Polonica* **58**, 1-16.
- FOTH C., TISCHLINGER, H. & RAUHUT, O. W. (2014). New specimen of *Archaeopteryx* provides insights into the evolution of pennaceous feathers. *Nature* **511**, 79-82.
- FOWLER D. W., FREEDMAN, E. A. & SCANNELLA, J. B. (2009). Predatory functional morphology in raptors: interdigital variation in talon size is related to prey restraint and immobilisation technique. *PLOS ONE* **4**, e7999.
- FOWLER D. W., FREEDMAN, E. A., SCANNELLA, J. B. & KAMBIC, R. E. (2011). The predatory ecology of *Deinonychus* and the origin of flapping in birds. *PLOS ONE* **6**, e28964.
- FRASER D. & THEODOR, J. M. (2011). Anterior dentary shape as an indicator of diet in ruminant artiodactyls. *Journal of Vertebrate Paleontology* **31**, 1366-1375.
- FREDERICK P. C. & BILDSTEIN, K. L. (1992). Foraging ecology of seven species of neotropical ibises (Threskiornithidae) during the dry season in the Llanos of Venezuela. *Wilson Bulletin* **104**, 1-21.
- FREELAND W. J. (1981). Functional aspects of primate grooming. *The Ohio Journal of Science* **81**, 173-177.
- FREI S., ORTMANN, S., REUTLINGER, C., KREUZER, M., HATT, J.-M. & CLAUSS, M. (2014). Comparative digesta retention patterns in ratites. *The Auk: Ornithological Advances* **132**, 119-131.
- FREY E., MARTILL, D. M. & BUCHY, M. (2003). A New Species of Tapejarid Pterosaur with Soft-Tissue Head Crest. In *Evolution and Palaeobiology of Pterosaurs*. Geological Society, London, Special Publications (ed. E. Buffetaut and J.-M. Mazin), pp. 1-3. The Geological Society London, London, UK.
- FRICKE H. C. & ROGERS, R. R. (2000). Multiple taxon–multiple locality approach to providing oxygen isotope evidence for warm-blooded theropod dinosaurs. *Geology* **28**, 799-802.
- FRIEDMANN H. (1931). Observations on the growth rate of the foot in the mound birds of the genus *Megapodius*. *Proceedings of the United States National Museum* **80**, 1-4.
- FROESE R. (2006). Cube law, condition factor and weight–length relationships: history, meta-analysis and recommendations. *Journal of Applied Ichthyology* **22**, 241-253.
- FUNSTON G. F., MENDONCA, S. E., CURRIE, P. J. & BARSBOLD, R. (2018). Oviraptorosaur anatomy, diversity and ecology in the Nemegt Basin. *Palaeogeography, Palaeoclimatology, Palaeoecology* **494**, 101-120.
- GAILER J. P., CALANDRA, I., SCHULZ-KORNAS, E. & KAISER, T. M. (2016). Morphology is not destiny: discrepancy between form, function and dietary adaptation in bovid cheek teeth. *Journal of Mammalian Evolution* **23**, 369-383.
- GALES R. P. (1987). Validation of the stomach-flushing technique for obtaining stomach contents of penguins. *IBIS* **129**, 335-343.

- GAO C., CHIAPPE, L. M., ZHANG, F., POMEROY, D. L., SHEN, C., CHINSAMY, A. & WALSH, M. O. (2012). A subadult specimen of the Early Cretaceous bird *Sapeornis chaoyangensis* and a taxonomic reassessment of sapeornithids. *Journal of Vertebrate Paleontology* **32**, 1103-1112.
- GARRARD G. E., MCCARTHY, M. A., VESK, P. A., RADFORD, J. Q. & BENNETT, A. F. (2012). A predictive model of avian natal dispersal distance provides prior information for investigating response to landscape change. *Journal of Animal Ecology* **81**, 14-23.
- GAUTHIER J. (1986). Saurischian monophyly and the origin of birds. *Memoirs of the California Academy of sciences* **8**, 1-55.
- GHOSH P., BHATTACHARYA, S. K., SAHNI, A., KAR, R. K., MOHABEY, D. M. & AMBWANI, K. (2003). Dinosaur coprolites from the Late Cretaceous (Maastrichtian) Lameta Formation of India: isotopic and other markers suggesting a C-3 plant diet. *Cretaceous Research* **24**, 743-750.
- GINGERICH P. D. (1973). Skull of *Hesperornis* and early evolution of birds. *Nature* **243**, 70-73.
- GINGERICH P. D. & ROSE, K. D. (1979). Anterior dentition of the eocene condylarth *Thryptacodon* - convergence with the tooth comb of lemurs. *Journal of Mammalogy* **60**, 16-22.
- GIRAudeau M., NOLAN, P. M., BLACK, C. E., EARL, S. R., HASEGAWA, M. & MCGRAW, K. J. (2014). Song characteristics track bill morphology along a gradient of urbanization in house finches (*Haemorrhous mexicanus*). *Frontiers in Zoology* **11**, 83.
- GLEN C. L. & BENNETT, M. B. (2007). Foraging modes of Mesozoic birds and non-avian theropods. *Current Biology* **17**, R911-R912.
- GODEFROIT P., CAU, A., DONG-YU, H., ESCUILLIÉ, F., WENHAO, W. & DYKE, G. (2013). A Jurassic avialan dinosaur from China resolves the early phylogenetic history of birds. *Nature* **498**, 359-362.
- GODEFROIT P., DEMUYNCK, H., DYKE, G., HU, D., ESCUILLIÉ, F. & CLAEYS, P. (2013). Reduced plumage and flight ability of a new Jurassic paravian theropod from China. *Nature Communications* **4**, 1394.
- GODINHO R. M., TORO-IBACACHE, V., FITTON, L. C. & O'HIGGINS, P. (2017). Finite element analysis of the cranium: validity, sensitivity and future directions. *Comptes Rendus Palevol* **16**, 600-612.
- GOLCHER-BENAVIDES J. & WAGNER, C. E. (2019). Playing out Liem's paradox: opportunistic piscivory across lake Tanganyikan cichlids. *American Naturalist* **194**, 260-267.
- GOSLOW G. E. (1972). Adaptive mechanisms of the raptor pelvic limb. *The Auk: Ornithological Advances* **89**, 47-64.
- GOULD S. J. (2002). The Integration and Adaptation (Structure and Function) in Ontogeny and Phylogeny: Historical Constraints and the Evolution of Development. In *The Structure of Evolutionary Theory*, pp. 1025-1177. Harvard University Press, Cambridge, MA, USA.
- GRANT B. R. & GRANT, P. R. (1993). Evolution of Darwin's finches caused by a rare climatic event. *Proceedings of the Royal Society B: Biological Sciences* **251**, 111-117.
- GRAU N., DAW, J. L., PATEL, R., EVANS, C., LEWIS, N. & MAO, J. J. (2006). Nanostructural and nanomechanical properties of synostosed postnatal human cranial sutures. *Journal of Craniofacial Surgery* **17**, 91-98.
- GREEN J. L. & CROFT, D. A. (2018). Using Dental Mesowear and Microwear for Dietary Inference: A Review of Current Techniques and Applications. In *Methods in Paleocology: Reconstructing Cenozoic Terrestrial Environments and Ecological Communities* (ed. D. A. Croft, D. F. Su and S. W. Simpson), pp. 53-73. Springer International Publishing, Gewerbesrasse, Switzerland.

- GREENBERG R., CADENA, V., DANNER, R. M. & TATTERSALL, G. (2012). Heat loss may explain bill size differences between birds occupying different habitats. *PLOS ONE* **7**, e40933.
- GRINE F. E., UNGAR, P. S. & TEAFORD, M. F. (2002). Error rates in dental microwear quantification using scanning electron microscopy. *Scanning: The Journal of Scanning Microscopies* **24**, 144-153.
- GU B., SCHELSKE, C. L. & HOYER, M. V. (1996). Stable isotopes of carbon and nitrogen as indicators of diet and trophic structure of the fish community in a shallow hypereutrophic lake. *Journal of Fish Biology* **49**, 1233-1243.
- GUILLERME T., JONES, M., LLOYD, G., O'REILLY, J., PATE, A., PUTTICK, M., RAYFIELD, E., SAUPE, E., SHERRATT, E. & SLATER, G. (2020). Disparities in the analysis of morphological disparity. *Biology Letters* **16**, 20200199.
- GUO X., XU, L. & JIA, S. (2018). Morphological and Phylogenetic study based on new materials of *Anchiornis huxleyi* (dinosauria, theropoda) from Jianchang, Western Liaoning, China. *Acta Geologica Sinica (English Edition)* **92**, 1-15.
- GUSSEKLOO S. W. & BOUT, R. G. (2005). Cranial kinesis in palaeognathous birds. *Journal of Experimental Biology* **208**, 3409-3419.
- HAUT R. C., LANCASTER, R. L. & DECAMP, C. E. (1992). Mechanical properties of the canine patellar tendon: some correlations with age and the content of collagen. *Journal of Biomechanics* **25**, 163-173.
- HEDRICK B. P., CORDERO, S. A., ZANNO, L. E., NOTO, C. & DODSON, P. (2019a). Quantifying shape and ecology in avian pedal claws: the relationship between the bony core and keratinous sheath. *Ecology and Evolution* **9**, 11545-11556.
- HEDRICK B. P., MUTUMI, G. L., MUNTEANU, V. D., SADIÉ, A., DAVIES, K. T., ROSSITER, S. J., SEARS, K. E., DÁVALOS, L. M. & DUMONT, E. (2019b). Morphological diversification under high integration in a hyper diverse mammal clade. *Journal of Mammalian Evolution*, 1-13.
- HEDRICK B. P., SCHACHNER, E. R., RIVERA, G., DODSON, P. & PIERCE, S. E. (2019c). The effects of skeletal asymmetry on interpreting biologic variation and taphonomy in the fossil record. *Paleobiology* **45**, 154-166.
- HENDERSON D. M. (1998). Skull and tooth morphology as indicators of niche partitioning in sympatric Morrison Formation theropods. *Gaia* **15**, 219-226.
- HENDERSON D. M. (2002). The eyes have it: the sizes, shapes, and orientations of theropod orbits as indicators of skull strength and bite force. *Journal of Vertebrate Paleontology* **22**, 766-778.
- HENDRICKX C. & MATEUS, O. (2014). Abelisauridae (Dinosauria: Theropoda) from the Late Jurassic of Portugal and dentition-based phylogeny as a contribution for the identification of isolated theropod teeth. *Zootaxa* **3759**, 1-74.
- HENDRICKX C., MATEUS, O. & ARAÚJO, R. (2015). A proposed terminology of theropod teeth (Dinosauria, Saurischia). *Journal of Vertebrate Paleontology* **35**, e982797.
- HENDRICKX C., TSCHOPP, E. & DEZCURRA, M. (2020). Taxonomic identification of isolated theropod teeth: the case of the shed tooth crown associated with *Aerosteon* (Theropoda: Megaraptora) and the dentition of Abelisauridae. *Cretaceous Research* **108**, 104312.
- HERREL A., MOORE, J. A., BREDEWEG, E. M. & NELSON, N. J. (2010a). Sexual dimorphism, body size, bite force and male mating success in tuatara. *Biological Journal of the Linnean Society* **100**, 287-292.

- HERREL A., PODOS, J., VANHOODYDONCK, B. & HENDRY, A. P. (2009). Force-velocity trade-off in Darwin's finch jaw function: a biomechanical basis for ecological speciation? *Functional Ecology* **23**, 119-125.
- HERREL A., SOONS, J., AERTS, P., DIRCKX, J., BOONE, M., JACOBS, P., ADRIAENS, D. & PODOS, J. (2010b). Adaptation and function of the bills of Darwin's finches: divergence by feeding type and sex. *Emu - Austral Ornithology* **110**, 39-47.
- HERTEL F. (1994). Diversity in body size and feeding morphology within past and present vulture assemblages. *Ecology* **75**, 1074-1084.
- HERTEL F. (1995). Ecomorphological indicators of feeding behavior in recent and fossil raptors. *The Auk: Ornithological Advances* **112**, 890-903.
- HILTON G. M., FURNESS, R. W. & HOUSTON, D. C. (2000a). A comparative study of digestion in North Atlantic seabirds. *Journal of Avian Biology* **31**, 36-46.
- HILTON G. M., FURNESS, R. W. & HOUSTON, D. C. (2000b). The effects of diet switching and mixing on digestion in seabirds. *Functional Ecology* **14**, 145-154.
- HILTON G. M., HOUSTON, D. C., BARTON, N. W. H., FURNESS, R. W. & RUXTON, G. D. (1999). Ecological constraints on digestive physiology in carnivorous and piscivorous birds. *Journal of Experimental Zoology Part A: Ecological Genetics and Physiology* **283**, 365-376.
- HILTON G. M., HOUSTON, D. C. & FURNESS, R. W. (1998). Which components of diet quality affect retention time of digesta in seabirds? *Functional Ecology* **12**, 929-939.
- HINIĆ-FRLOG S. & MOTANI, R. (2010). Relationship between osteology and aquatic locomotion in birds: determining modes of locomotion in extinct Ornithurae. *Journal of Evolutionary Biology* **23**, 372-385.
- HOBSON K. A. (1987). Use of stable-carbon isotope analysis to estimate marine and terrestrial protein content in gull diets. *Canadian Journal of Zoology* **65**, 1210-1213.
- HOBSON K. A. (1990). Stable isotope analysis of marbled murrelets: evidence for freshwater feeding and determination of trophic level. *The Condor: Ornithological Applications* **92**, 897-903.
- HOBSON K. A. (1993). Trophic relationships among high arctic seabirds - Insights from tissue-dependent stable-isotope models. *Marine Ecology Progress Series* **95**, 7-18.
- HOBSON K. A., BOWEN, G. J., WASSENAAR, L. I., FERRAND, Y. & LORMEE, H. (2004). Using stable hydrogen and oxygen isotope measurements of feathers to infer geographical origins of migrating European birds. *Oecologia* **141**, 477-488.
- HOBSON K. A. & MONTEVECCHI, W. A. (1991). Stable isotopic determinations of trophic relationships of great auks. *Oecologia* **87**, 528-531.
- HOLLIDAY C. M. (2009). New insights into dinosaur jaw muscle anatomy. *The Anatomical Record: Advances in Integrative Anatomy and Evolutionary Biology* **292**, 1246-1265.
- HOLLOCHER T. C., CHIN, K., HOLLOCHER, K. T. & KRUGE, M. A. (2001). Bacterial residues in coprolite of herbivorous dinosaurs: role of bacteria in mineralization of feces. *Palaios* **16**, 547-565.
- HOLLUND H. I., ARTS, N., JANS, M. M. E. & KARS, H. (2015). Are teeth better? Histological characterization of diagenesis in archaeological bone-tooth pairs and a discussion of the consequences for archaeometric sample selection and analyses. *International Journal of Osteoarchaeology* **25**, 901-911.
- HOLTZ JR. T. R. (2008). A critical reappraisal of the obligate scavenging hypothesis for *Tyrannosaurus rex* and other tyrant dinosaurs. In *Tyrannosaurus rex the Tyrant King* (ed. P. L. Larson and K. Carpenter), pp. 371-396. Indiana University Press, Bloomington, IN, USA.

- HOLTZ JR. T. R., BRINKMAN, D. L. & CHANDLER, C. L. (1998). Denticle morphometrics and a possibly omnivorous feeding habit for the theropod dinosaur *Troodon*. *Gaia* **15**, 159-166.
- HOPSON J. A. (2001). Ecomorphology of Avian and Nonavian Theropod Phalangeal Proportions: Implications for the Arboreal Versus Terrestrial Origin of Bird Flight. In *New Perspectives on the Origin and Early Evolution of Birds: Proceedings of the International Symposium in Honor of John H. Ostrom* (ed. J. H. Ostrom, L. F. Gall and J. Gauthier), pp. 211-235. Peabody Museum of Natural History, New Haven, CN, USA.
- HOU L. (1997). *Mesozoic Birds of China*. Phoenix Valley Provincial Aviary of Taiwan.
- HOU L. & CHEN, P. (1999a). *Liaoxiornis delicatus* gen. et sp. nov., the smallest Mesozoic bird. *Chinese Science Bulletin* **44**, 834-838.
- HOU L., D MARTIN, L., ZHOU, Z. & FEDUCCIA, A. (1999b). *Archaeopteryx* to opposite birds--Missing link from the Mesozoic of China. *Vertebrata Palasiatica* **37**, 88-95.
- HOU L., MARTIN, L. D., ZHOU, Z., FEDUCCIA, A. & ZHANG, F. (1999c). A diapsid skull in a new species of the primitive bird *Confuciusornis*. *Nature* **399**, 679.
- HOU L. H., CHIAPPE, L. M., ZHANG, F. C. & CHUONG, C. M. (2004). New Early Cretaceous fossil from China documents a novel trophic specialization for Mesozoic birds. *Naturwissenschaften* **91**, 22-25.
- HU D., LI, L., HOU, L. & XU, X. (2010). A new sapeornithid bird from China and its implication for early avian evolution. *Acta Geologica Sinica (English Edition)* **84**, 472-482.
- HU D., LI, L., HOU, L. & XU, X. (2011). A new enantiornithine bird from the Lower Cretaceous of western Liaoning, China. *Journal of Vertebrate Paleontology* **31**, 154-161.
- HU H. & O'CONNOR, J. K. (2017). First species of Enantiornithes from Sihedang elucidates skeletal development in Early Cretaceous enantiornithines. *Journal of Systematic Palaeontology* **15**, 909-926.
- HU H., O'CONNOR, J. K. & ZHOU, Z. H. (2015). A new species of Pengornithidae (Aves: Enantiornithes) from the Lower Cretaceous of China suggests a specialized scansorial habitat previously unknown in early birds. *PLOS ONE* **10**, e0126791.
- HU H., O'CONNOR, J. K., McDONALD, P. G. & WROE, S. (2020a). Cranial osteology of the Early Cretaceous Sapeornis chaoyangensis (Aves: Pygostylia). *Cretaceous Research*, 104496.
- HU H., O'CONNOR, J. K., WANG, M., WROE, S. & McDONALD, P. G. (2020b). New anatomical information on the bohaiornithid Longusunguis and the presence of a plesiomorphic diapsid skull in Enantiornithes. *Journal of Systematic Palaeontology*.
- HU H., SANSALONE, G., WROE, S., McDONALD, P. G., O'CONNOR, J. K., LI, Z. H., XU, X. & ZHOU, Z. H. (2019). Evolution of the vomer and its implications for cranial kinesis in Paraves. *Proceedings of the National Academy of Sciences of the United States of America* **116**, 19571-19578.
- HU Y., GHIGLIOTTI, L., VACCHI, M., PISANO, E., DETRICH, H. W. & ALBERTSON, R. C. (2016). Evolution in an extreme environment: developmental biases and phenotypic integration in the adaptive radiation of antarctic notothenioids. *BMC Evolutionary Biology* **16**, 142.
- HUANG J., WANG, X., HU, Y., LIU, J., PETEYA, J. A. & CLARKE, J. A. (2016). A new ornithurine from the Early Cretaceous of China sheds light on the evolution of early ecological and cranial diversity in birds. *PeerJ* **4**, e1765.
- HUBER S. K. & PODOS, J. (2006). Beak morphology and song features covary in a population of Darwin's finches (*Geospiza fortis*). *Biological Journal of the Linnean Society* **88**, 489-498.

- HULME P. E. & BENKMAN, C. W. (2002). Granivory. In *Plant–Animal Interactions: An Evolutionary Approach* (ed. O. Pellmyr and C. M. Herra), pp. 185-208. Blackwell Science Ltd, Malden, MA, USA.
- HUNT A. P., LUCAS, S. G., MILAN, J. & SPIELMANN, J. A. (2012). Vertebrate coprolite studies: status and prospectus. *New Mexico Museum of Natural History Science Bulletin* **57**, 5-24.
- HUTCHINSON J. R. & ALLEN, V. (2008). The evolutionary continuum of limb function from early theropods to birds. *Naturwissenschaften* **96**, 423-448.
- HUTSON J. M., BURKE, C. C. & HAYNES, G. (2013). Osteophagia and bone modifications by giraffe and other large ungulates. *Journal of Archaeological Science* **40**, 4139-4149.
- JACKSON S. (1992). Do seabird gut sizes and mean retention times reflect adaptation to diet and foraging method? *Physiological Zoology* **65**, 674-697.
- JACOBS R. P. W. M. (1982). Component Studies in Seagrass Ecosystems along West European Coasts, Radboud University Nijmegen.
- JAMES H. F. & BURNEY, D. A. (1997). The diet and ecology of Hawaii's extinct flightless waterfowl: evidence from coprolites. *Biological Journal of the Linnean Society* **62**, 279-297.
- JANZEN D. H. (1981). *Ficus ovalis* seed predation by an orange-chinned parakeet (*Brotogeris jugularis*) in Costa Rica. *The Auk: Ornithological Advances* **98**, 841-844.
- JAOUEN K., BALTER, V., HERRSCHER, E., LAMBOUX, A., TELOUK, P. & ALBAREDE, F. (2012). Fe and Cu stable isotopes in archeological human bones and their relationship to sex. *American Journal of Physical Anthropology* **148**, 334-340.
- JARMAN S. N., MCINNES, J. C., FAUX, C., POLANOWSKI, A. M., MARTHICK, J., DEAGLE, B. E., SOUTHWELL, C. & EMMERSON, L. (2013). Adelie Penguin Population Diet Monitoring by Analysis of Food DNA in Scats. *PLOS ONE* **8**, e82227.
- JI Q., JI, S., ZHANG, H., YOU, H., ZHANG, J., WANG, L., YUAN, C. & JI, X. (2002). A new avialian bird - *Jixiangornis orientalis* gen. et sp. nov. - from the Lower Cretaceous of western Liaoning, NE China. *Journal of Nanjing University Natural Sciences Edition* **38**, 723-736.
- JI Q., JI, S. A., YOU, H., ZHANG, J., ZHANG, H., ZHANG, N., YUAN, C. & JI, X. (2003). An early Cretaceous avialian bird, *Shenzhouraptor sinensis* from western Liaoning, China. *Acta Geologica Sinica (English Edition)* **77**, 21-27.
- JOLLIFFE I. T. & CADIMA, J. (2016). Principal component analysis: a review and recent developments. *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences* **374**, 20150202.
- JONES K. E., BIELBY, J., CARDILLO, M., FRITZ, S. A., O'DELL, J., ORME, C. D. L., SAFI, K., SECHREST, W., BOAKES, E. H., CARBONE, C., CONNOLLY, C., CUTTS, M. J., FOSTER, J. K., GRENYER, R., HABIB, M., PLASTER, C. A., PRICE, S. A., RIGBY, E. A., RIST, J., TEACHER, A., BININDA-EMONDS, O. R. P., GITTLEMAN, J. L., MACE, G. M. & PURVIS, A. (2009). PanTHERIA: a species-level database of life history, ecology, and geography of extant and recently extinct mammals. *Ecology* **90**, 2648-2648.
- JORDANO P. (2000). Fruits and Frugivory. In *Seeds: The Ecology of Regeneration in Plant Communities* (ed. M. Fenner), pp. 125-166. CABI Publishing, Wallingford, UK.
- JUNG J.-Y. (2019). Learning from Nature: An investigation of an impact-resistant system on the woodpecker head as a non-traumatic brain injury animal model, UC San Diego.



- JUVINALL R. C. & MARSHEK, K. M. (2011). Failure Theories, Safety Factors, and Reliability. In *Fundamentals of Machine Component Design*, pp. 248-282. Wiley New York, Hoboken, NJ, USA.
- KAESLER R. L. & WATERS, J. A. (1972). Fourier analysis of the ostracode margin. *Geological Society of America Bulletin* **83**, 1169-1178.
- KAMBIC R. E. (2008). Multivariate analysis of avian and non-avian theropod pedal phalanges, Montana State University-Bozeman, College of Letters & Science.
- KARBAN R. & AGRAWAL, A. A. (2002). Herbivore offense. *Annual Review of Ecology and Systematics* **33**, 641-664.
- KAYE T. G., PITTMAN, M., MARUGÁN-LOBÓN, J., MARTÍN-ABAD, H., SANZ, J. L. & BUSCALIONI, A. D. (2019). Fully fledged enantiornithine hatchling revealed by Laser-Stimulated Fluorescence supports precocial nesting behavior. *Scientific Reports* **9**, 5006.
- KEAR B. P., BOLES, W. E. & SMITH, E. T. (2003). Unusual gut contents in a Cretaceous ichthyosaur. *Proceedings of the Royal Society B: Biological Sciences* **270**, S206-S208.
- KELLNER A. W. (1996). Fossilized theropod soft tissue. *Nature* **379**, 32.
- KILTIE R. A. (1982). Bite force as a basis for niche differentiation between rain forest peccaries (*Tayassu tajacu* and *T. pecari*). *Biotropica*, 188-195.
- KIM S.-K. (2011). The Effect of Discretization of Continuous Data: Principal Component Analysis vs. Correspondence Analysis (Dual Scaling). In *American Educational Research Association*, New Orleans, Louisiana.
- KLACZKO J., SHERRATT, E. & SETZ, E. Z. F. (2016). Are diet preferences associated to skulls shape diversification in xenodontine snakes? *PLOS ONE* **11**, e0148375.
- KLINGENBERG C. P. (2014). Studying morphological integration and modularity at multiple levels: concepts and analysis. *Philosophical Transactions of the Royal Society B: Biological Sciences* **369**, 20130249.
- KLINGENBERG C. P. (2016). Size, shape, and form: concepts of allometry in geometric morphometrics. *Development genes and evolution* **226**, 113-137.
- KLOSKOWSKI J., TREMBACZOWSKI, A. & FILIPIUK, M. (2019). Stable isotope tracing of links between marine wintering and freshwater breeding habitats of Red-necked Grebes. *Journal of Ornithology* **160**, 593-605.
- KOCH P. L. (2007). Isotopic Study of the Biology of Modern and Fossil Vertebrates. In *Stable isotopes in ecology and environmental science*, vol. 2 (ed. R. H. Michener and K. Lajtha), pp. 99-154. Blackwell Publishing, Hoboken, NJ, USA.
- KOENIGSWALD W. V., HABERSETZER, J. & GINGERICH, P. D. (2011). Morphology and evolution of the distal phalanges in primates. In *The World at the Time of Messel: Puzzles in Palaeobiology, Palaeoenvironment, and the History of Early Primates* (ed. T. Lehmann and S. Schaal), pp. 91-94. International Senckenberg Conference, Frankfurt am Main, Germany.
- KOLODNY Y., LUZ, B., SANDER, M. & CLEMENS, W. (1996). Dinosaur bones: fossils or pseudomorphs? The pitfalls of physiology reconstruction from apatitic fossils. *Palaeogeography, Palaeoclimatology, Palaeoecology* **126**, 161-171.
- KULEMEYER C., ASBAHR, K., GUNZ, P., FRAHNERT, S. & BAIRLEIN, F. (2009). Functional morphology and integration of corvid skulls - A 3D geometric morphometric approach. *Frontiers in Zoology* **6**.

- KUNDRÁT M., NUDDS, J., KEAR, B. P., LÜ, J. & AHLBERG, P. (2019). The first specimen of *Archaeopteryx* from the upper Jurassic Mörnsheim Formation of Germany. *Historical Biology* **31**, 3-63.
- KUPCZIK K., DOBSON, C. A., FAGAN, M. J., CROMPTON, R. H., OXNARD, C. E. & O'HIGGINS, P. (2007). Assessing mechanical function of the zygomatic region in macaques: validation and sensitivity testing of finite element models. *Journal of Anatomy* **210**, 41-53.
- KUROCHKIN E. N., CHATTERJEE, S. & MIKHAILOV, K. E. (2013). An embryonic enantiornithine bird and associated eggs from the Cretaceous of Mongolia. *Paleontological Journal* **47**, 1252-1269.
- LADYGUIN A. (2000). The morphology of the bill apparatus in the Steller's Sea Eagle. In *First Symposium on Steller's and White-tailed Sea eagles in East Asia* (ed. M. Ueta and M. J. McGrady), pp. 1-10. Wild Bird Society of Japan, Tokyo, Japan.
- LAMBE L. M. (1917). *The Cretaceous theropod dinosaur Gorgosaurus*. Government Printing Bureau, Ottawa, Canada.
- LAMBERT J. E., CHAPMAN, C. A., WRANGHAM, R. W. & CONKLIN-BRITAIN, N. L. (2004). Hardness of cercopithecine foods: implications for the critical function of enamel thickness in exploiting fallback foods. *American Journal of Physical Anthropology: The Official Publication of the American Association of Physical Anthropologists* **125**, 363-368.
- LARSON D. W., BROWN, C. M. & EVANS, D. C. (2016). Dental Disparity and Ecological Stability in Bird-like Dinosaurs prior to the End-Cretaceous Mass Extinction. *Current Biology* **26**, 1325-1333.
- LAUTENSCHLAGER S. (2014). Morphological and functional diversity in therizinosaur claws and the implications for theropod claw evolution. *Proceedings of the Royal Society B: Biological Sciences* **281**, 20140497.
- LAUTENSCHLAGER S. (2016). Reconstructing the past: methods and techniques for the digital restoration of fossils. *Royal Society Open Science* **3**.
- LAUTENSCHLAGER S. (2017). Functional niche partitioning in Therizinosauria provides new insights into the evolution of theropod herbivory. *Palaeontology* **60**, 375-387.
- LAUTENSCHLAGER S., BRIGHT, J. A. & RAYFIELD, E. J. (2014a). Digital dissection—using contrast-enhanced computed tomography scanning to elucidate hard-and soft-tissue anatomy in the Common Buzzard *Buteo buteo*. *Journal of Anatomy* **224**, 412-431.
- LAUTENSCHLAGER S., WITMER, L. M., ALTANGEREL, P. & RAYFIELD, E. J. (2013). Edentulism, beaks, and biomechanical innovations in the evolution of theropod dinosaurs. *Proceedings of the National Academy of Sciences* **110**, 20657-20662.
- LAUTENSCHLAGER S., WITMER, L. M., ALTANGEREL, P., ZANNO, L. E. & RAYFIELD, E. J. (2014b). Cranial anatomy of *Erlikosaurus andrewsi* (Dinosauria, Therizinosauria): new insights based on digital reconstruction. *Journal of Vertebrate Paleontology* **34**, 1263-1291.
- LEE N., HORSTEMEYER, M., RHEE, H., NABORS, B., LIAO, J. & WILLIAMS, L. N. (2014). Hierarchical multiscale structure–property relationships of the red-bellied woodpecker (*Melanerpes carolinus*) beak. *Journal of the Royal Society Interface* **11**, 20140274.
- LEE Y.-C., CHIANG, C.-C., HUANG, P.-Y., CHUNG, C.-Y., HUANG, T. D., WANG, C.-C., CHEN, C.-L., CHANG, R.-S., LIAO, C.-H. & REISZ, R. R. (2017). Evidence of preserved collagen in an Early Jurassic sauropodomorph dinosaur revealed by synchrotron FTIR microspectroscopy. *Nature Communications* **8**, 14220.
- LEFÈVRE U., CAU, A., CINCOTTA, A., HU, D., CHINSAMY, A., ESCUILLIÉ, F. & GODEFROIT, P. (2017). A new Jurassic theropod from China documents a transitional step in the macrostructure of feathers. *The Science of Nature* **104**, 74.

- LEFÈVRE U., HU, D., ESCUILLIÉ, F., DYKE, G. & GODEFROIT, P. (2014). A new long-tailed basal bird from the Lower Cretaceous of north-eastern China. *Biological Journal of the Linnean Society* **113**, 790-804.
- LEISLER B. & THALER, E. (1982). Differences in morphology and foraging behaviour in the goldcrest *Regulus regulus* and firecrest *R. ignicapillus*. In *Annales Zoologici Fennici*, pp. 277-284. JSTOR.
- LEVEY D. J. & DEL RIO, C. M. (2001). It takes guts (and more) to eat fruit: lessons from avian nutritional ecology. *The Auk: Ornithological Advances* **118**, 819-831.
- LI L., DUAN, Y., HU, D. Y., WANG, L., CHENG, S. L. & HOU, L. H. (2006). New eoenantiornithid bird from the Early Cretaceous Jiufotang Formation of western Liaoning, China. *Acta Geologica Sinica (English Edition)* **80**, 38-41.
- LI L., GONG, E., ZHANG, L., YANG, Y. & HOU, L. (2010). A new enantiornithine bird (Aves) from the Early Cretaceous of Liaoning, China. *Acta Palaeontologica Sinica* **49**, 524-531.
- LI L. & HOU, S.-L. (2011). Discovery of a new bird (Enantiornithines) from Lower Cretaceous in western Liaoning, China. *Journal of Jilin University (Earth Science Edition)* **41**.
- LI L., HU, D., DUAN, Y., GONG, E. & HOU, L. (2007). Alethoalaornithidae fam. nov.: a new family of enantiornithine bird from the Lower Cretaceous of western Liaoning. *Acta Palaeontologica Sinica* **46**, 371.
- LI L., WANG, J., ZHANG, X. & HOU, S. (2012). A new enantiornithine bird from the Lower Cretaceous Jiufotang Formation in Jinzhou Area, Western Liaoning Province, China. *地质学报 (英文版)* **86**, 1039-1044.
- LI Q. G., CLARKE, J. A., GAO, K. Q., PETEYE, J. A. & SHAWKEY, M. D. (2018). Elaborate plumage patterning in a Cretaceous bird. *PeerJ* **6**, e5831.
- LI Z., WANG, C.-C., WANG, M., CHIANG, C.-C., WANG, Y., ZHENG, X., HUANG, E.-W., HSIAO, K. & ZHOU, Z. (2020). Ultramicrostructural reductions in teeth: implications for dietary transition from non-avian dinosaurs to birds. *BMC Evolutionary Biology* **20**, 1-8.
- LI Z. H. & CLARKE, J. A. (2016). The craniolingual morphology of waterfowl (Aves, Anseriformes) and its relationship with feeding mode revealed through contrast-enhanced x-ray computed tomography and 2D morphometrics. *Evolutionary Biology* **43**, 12-25.
- LI Z. H., ZHOU, Z. H., WANG, M. & CLARKE, J. A. (2014). A new specimen of large-bodied basal enantiornithine *Bohaiornis* from the Early Cretaceous of China and the inference of feeding ecology in Mesozoic birds. *Journal of Paleontology* **88**, 99-108.
- LIEM K. F. (1980). Adaptive significance of intra- and interspecific differences in the feeding repertoires of cichlid fishes. *American Zoologist* **20**, 295-314.
- LINGHAM-SOLIAR T. (2008). A unique cross section through the skin of the dinosaur *Psittacosaurus* from China showing a complex fibre architecture. *Proceedings of the Royal Society B: Biological Sciences* **275**, 775-780.
- LIU D., CHIAPPE, L. M., SERRANO, F., HABIB, M., ZHANG, Y. G. & MENG, Q. J. (2017). Flight aerodynamics in enantiornithines: information from a new Chinese Early Cretaceous bird. *PLOS ONE* **12**, e0184637.
- LIU D., CHIAPPE, L. M., ZHANG, Y. G., SERRANO, F. J. & MENG, Q. J. (2019). Soft tissue preservation in two new enantiornithine specimens (Aves) from the Lower Cretaceous Huajiying Formation of Hebei Province, China. *Cretaceous Research* **95**, 191-207.
- LOCKIE J. D. (1956). The food and feeding behaviour of the jackdaw, rook and carrion crow. *The Journal of Animal Ecology*, 421-428.

- LOPES L. E., FERNANDES, A. M., MEDEIROS, M. C. I. & MARINI, M. A. (2016). A classification scheme for avian diet types. *Journal of Field Ornithology* **87**, 309-322.
- LOTT C. A., MEEHAN, T. D. & HEATH, J. A. (2003). Estimating the latitudinal origins of migratory birds using hydrogen and sulfur stable isotopes in feathers: influence of marine prey base. *Oecologia* **134**, 505-510.
- LU J., XU, L., ZHANG, X., JIA, S. & CHANG, H. (2011). A new Gobipterygid bird from the Late Cretaceous Central China and its biogeographic implications. In *Journal of Vertebrate Paleontology*, vol. 31, pp. 147-147. Society of Vertebrate Paleontology, Las Vegas, NV, USA.
- MA W., BRUSATTE, S. L., LÜ, J. & SAKAMOTO, M. (2019). The skull evolution of oviraptorosaurian dinosaurs: the role of niche-partitioning in diversification. *Journal of Evolutionary Biology* **00**, 1-11.
- MA W., PITTMAN, M., LAUTENSCHLAGER, S., MEADE, L. E. & XU, X. (2020). Functional morphology of the oviraptorosaur and scansoriopterygid skull. In *Pennaraptoran Theropod Dinosaurs: Past Progress and New Frontiers* (ed. M. Pittman and X. Xu), pp. 229-249. Bulletin of the American Museum of Natural History.
- MA W., WANG, J., PITTMAN, M., TAN, Q., TAN, L., GUO, B. & XU, X. (2017). Functional anatomy of a giant toothless mandible from a bird-like dinosaur: *Gigantoraptor* and the evolution of the oviraptorosaurian jaw. *Scientific Reports* **7**, 1-15.
- MAIOLINO S., BOYER, D. M. & ROSENBERGER, A. (2011). Morphological correlates of the grooming claw in distal phalanges of platyrrhines and other primates: a preliminary study. *The Anatomical Record: Advances in Integrative Anatomy and Evolutionary Biology* **294**, 1975-1990.
- MANNING P. L., MARGETTS, L., JOHNSON, M. R., WITHERS, P. J., SELLERS, W. I., FALKINGHAM, P. L., MUMMERY, P. M., BARRETT, P. M. & RAYMONT, D. R. (2009). Biomechanics of dromaeosaurid dinosaur claws: application of X-ray microtomography, nanoindentation, and finite element analysis. *The Anatomical Record: Advances in Integrative Anatomy and Evolutionary Biology* **292**, 1397-1405.
- MANNING P. L., PAYNE, D., PENNICOTT, J., BARRETT, P. M. & ENNOS, R. A. (2006). Dinosaur killer claws or climbing crampons? *Biology Letters* **2**, 110-112.
- MARCÉ-NOGUÉ J., DE ESTEBAN-TRIVIGNO, S., ESCRIG PÉREZ, C. & GIL ESPERT, L. (2016). Accounting for differences in element size and homogeneity when comparing finite element models: armadillos as a case study. *Palaeontologia Electronica* **19**, 2T.
- MARCÉ-NOGUÉ J., DE ESTEBAN-TRIVIGNO, S., PÜSCHEL, T. A. & FORTUNY, J. (2017). The intervals method: a new approach to analyse finite element outputs using multivariate statistics. *PeerJ* **5**, e3793.
- MARCÉ-NOGUÉ J., DEMIGUEL, D., FORTUNY TERRICABRAS, J., DE ESTEBAN-TRIVIGNO, S. & GIL ESPERT, L. (2013). Quasi-homothetic transformation for comparing the mechanical performance of planar models in biological research. *Palaeontologia Electronica* **16**, 6T.
- MARCUS L. F. (1990). Traditional Morphometrics. In *Proceedings of the Michigan morphometrics workshop* (ed. F. J. Rohlf and F. L. Bookstein), pp. 77-122. University of Michigan Museum of Zoology, Ann Arbor, MI, USA.
- MARROIG G., SHIRAI, L. T., PORTO, A., DE OLIVEIRA, F. B. & DE CONTO, V. (2009). The evolution of modularity in the mammalian skull II: evolutionary consequences. *Evolutionary Biology* **36**, 136-148.

- MARTIN J. E., TACAIL, T. & BALTER, V. (2017). Non-traditional isotope perspectives in vertebrate palaeobiology. *Palaeontology* **60**, 485-502.
- MARTIN L. D. & NAPLES, V. L. (2008). Mandibular kinesis in Hesperornis. *Oryctos* **7**, 61-65.
- MARTIN L. D. & TATE, J. (1976). The skeleton of *Baptornis advenus* (Aves: Hesperornithiformes). *Smithsonian Contributions to Paleobiology* **27**, 35-66.
- MARTIN L. D. & ZHOU, Z. H. (1997). *Archaeopteryx*-like skull in Enantiornithine bird. *Nature* **389**, 556-556.
- MARUGÁN-LOBÓN J. & BUSCALIONI, Á. D. (2004). Geometric morphometrics in macroevolution: morphological diversity of the skull in modern avian forms in contrast to some theropod dinosaurs. In *Morphometrics* (ed. A. M. T. Elewa), pp. 157-173. Springer-Verlag, Berlin, Germany.
- MATSUI H., HUNT, G. R., OBERHOFER, K., OGIHARA, N., MCGOWAN, K. J., MITHRARATNE, K., YAMASAKI, T., GRAY, R. D. & IZAWA, E.-I. (2016). Adaptive bill morphology for enhanced tool manipulation in New Caledonian crows. *Scientific Reports* **6**, 1-11.
- MAYR G. (2016). *Avian Evolution: The Fossil Record of Birds and its Paleobiological Significance*. John Wiley & Sons, Oxford, UK.
- MAYR G. (2017). Evolution of avian breeding strategies and its relation to the habitat preferences of Mesozoic birds. *Evolutionary Ecology* **31**, 131-141.
- MAYR G. (2018). A survey of casques, frontal humps, and other extravagant bony cranial protuberances in birds. *Zoomorphology* **137**, 457-472.
- MAYR G., CODREA, V., SOLOMON, A., BORDEIANU, M. & SMITH, T. (2020). Reply to comments on “A well-preserved pelvis from the Maastrichtian of Romania suggests that the enigmatic *Gargantuavis* is neither an ornithurine bird nor an insular endemic”. *Cretaceous Research*, 104465.
- MAYR G. & MANEGOLD, A. (2013). Can ovarian follicles fossilize? *Nature* **499**, E1.
- MAZZETTA G. V., CISILINO, A. P., BLANCO, R. E. & CALVO, N. (2009). Cranial mechanics and functional interpretation of the horned carnivorous dinosaur *Carnotaurus sastrei*. *Journal of Vertebrate Paleontology* **29**, 822-830.
- MCBRAYER L. D. & CORBIN, C. E. (2007). Patterns of Head Shape Variation in Lizards: Morphological Correlates of Foraging Mode. In *Lizard ecology* (ed. D. Lance), pp. 271-301. Cambridge University Press, Cambridge, UK.
- MCHENRY M. & SUMMERS, A. (2011). A force–speed trade-off is not absolute. *Biology Letters* **7**, 880-881.
- MCINTOSH A. F. & COX, P. G. (2016). The impact of gape on the performance of the skull in chisel-tooth digging and scratch digging mole-rats (Rodentia: Bathyergidae). *Royal Society Open Science* **3**, 160568.
- MCLAUGHLIN E., ZHANG, Y. G., PASHLEY, D., BORKE, J. & YU, J. C. (2000). The load-displacement characteristics of neonatal rat cranial sutures. *The Cleft Palate-Craniofacial Journal* **37**, 590-595.
- MCWHORTER T. J. & MARTÍNEZ DEL RIO, C. (2000). Does gut function limit hummingbird food intake? *Physiological and Biochemical Zoology* **73**, 313-324.
- MCWILLIAMS S. R., CAVIEDES-VIDAL, E. & KARASOV, W. H. (1999). Digestive adjustments in cedar waxwings to high feeding rate. *Journal of Experimental Zoology* **283**, 394-407.
- MELNICK C. A., CHEN, Z. & MECHOLSKY, J. J. (1996). Hardness and toughness of exoskeleton material in the stone crab, *Menippe mercenaria*. *Journal of Materials Research* **11**, 2903-2907.

- MELORO C., HUDSON, A. & ROOK, L. (2015). Feeding habits of extant and fossil canids as determined by their skull geometry. *Journal of Zoology* **295**, 178-188.
- MELSTROM K. M. (2017). The relationship between diet and tooth complexity in living dentigerous saurians. *Journal of Morphology* **278**, 500-522.
- MIHLBACHLER M. C., BEATTY, B. L., CALDERA-SIU, A., CHAN, D. & LEE, R. (2012). Error rates and observer bias in dental microwear analysis using light microscopy. *Palaeontologia Electronica* **15**, 12A.
- MIHLBACHLER M. C., CAMPBELL, D., AYOUB, M., CHEN, C. & GHANI, I. (2016). Comparative dental microwear of ruminant and perissodactyl molars: implications for paleodietary analysis of rare and extinct ungulate clades. *Paleobiology* **42**, 98-116.
- MIHLBACHLER M. C., RIVALS, F., SOLOUNIAS, N. & SEMPREBON, G. M. (2011). Dietary change and evolution of horses in North America. *Science* **331**, 1178-1181.
- MILLER C. V., PITTMAN, M., KAYE, T. G., WANG, X. & ZHENG, X. (In Press). Disassociated rhamphotheca of fossil bird *Confuciusornis* informs early beak reconstruction, stress regime, and developmental patterns. *Communications Biology*.
- MILLER G. H., FOGEL, M. L., MAGEE, J. W., GAGAN, M. K., CLARKE, S. J. & JOHNSON, B. J. (2005). Ecosystem collapse in Pleistocene Australia and a human role in megafaunal extinction. *Science* **309**, 287-290.
- MIYASHITA T. (2013). Geometric and Developmental Perspectives on the Evolution of the Skull and Internal Carotid Circulation in Turtles. In *Morphology and Evolution of Turtles* (ed. D. B. Brinkman, P. A. Holroyd and J. D. Gardner), pp. 71-101. Springer Netherlands, Dordrecht, Netherlands.
- MONTANARI S. & NORELL, M. (2011). Dietary inferences of protoceratopsian dinosaurs from the Late Cretaceous of Mongolia based on stable isotope geochemistry. *Journal of Vertebrate Paleontology* **31**, 160-161.
- MONTUELLE S. J. & KANE, E. A. (2019). Food Capture in Vertebrates: A Complex Integrative Performance of the Cranial and Postcranial Systems. In *Feeding in Vertebrates: Evolution, Morphology, Behavior, Biomechanics* (ed. V. Bels), pp. 71-137. Springer International Publishing, Gewerbesrasse, Switzerland.
- MORALES-GARCÍA N. M., BURGESS, T. D., HILL, J. J., GILL, P. G. & RAYFIELD, E. J. (2019). The use of extruded finite-element models as a novel alternative to tomography-based models: a case study using early mammal jaws. *Journal of the Royal Society Interface* **16**, 20190674.
- MORENO K., WROE, S., CLAUSEN, P., MCHENRY, C., D'AMORE, D. C., RAYFIELD, E. J. & CUNNINGHAM, E. (2008). Cranial performance in the Komodo dragon (*Varanus komodoensis*) as revealed by high-resolution 3-D finite element analysis. *Journal of Anatomy* **212**, 736-746.
- MORHARDT A. C. (2009). Dinosaur Smiles: Do the Texture and Morphology of the Premaxilla, Maxilla, and Dentary Bones of Sauropsids Provide Osteological Correlates for Inferring Extra-Oral Structures Reliably in Dinosaurs?, Western Illinois University.
- MORRIS W. J. (1970). *Hadrosaurian Dinosaur Bills: Morphology and Function*. Los Angeles County Museum of Natural History, Los Angeles, CA, USA.
- MORSCHHAUSER E. M., VARRICCHIO, D. J., CHUNLING, G., JINYUAN, L., XURI, W., XIADONG, C. & QINGJIN, M. (2009). Anatomy of the Early Cretaceous bird *Rapaxavis pani*, a new species from Liaoning Province, China. *Journal of Vertebrate Paleontology* **29**, 545-554.

- MOSTO M. C. & TAMBUSSI, C. P. (2014). Qualitative and Quantitative Analysis of Talons of Diurnal Bird of Prey. *Anatomia Histologia Embryologia* **43**, 6-15.
- MOYER B. R., PETERSON, A. T. & CLAYTON, D. H. (2002). Influence of bill shape on ectoparasite load in Western Scrub-Jays. *The Condor: Ornithological Applications* **104**, 675-678.
- MYHRVOLD N. P. (2012). A call to search for fossilised gastric pellets. *Historical Biology* **24**, 505-517.
- NAVA W., ALVARENGA, H., CHIAPPE, L. & MARTINELLI, A. (2015). Three-dimensionally preserved cranial remains of enantiornithine birds from the Late Cretaceous of Brazil. In *V Congreso Latinoamericano de Paleontología de Vertebrados*, Colonia del Sacramento, Uruguay.
- NAVALÓN G. (2014). Reconstructing the palaeobiology of *Confuciusornis* and other Confuciusornithiformes, University of Bristol.
- NAVALÓN G., BRIGHT, J. A., MARUGÁN-LOBÓN, J. & RAYFIELD, E. J. (2018a). The evolutionary relationship among beak shape, mechanical advantage, and feeding ecology in modern birds. *Evolution* **73**, 422-435.
- NAVALÓN G., MARUGÁN-LOBÓN, J., BRIGHT, J. A., COONEY, C. R. & RAYFIELD, E. J. (2020). The consequences of craniofacial integration for the adaptive radiations of Darwin's finches and Hawaiian honeycreepers. *Nature Ecology & Evolution* **4**, 270-278.
- NAVALÓN G., MENG, Q., MARUGÁN-LOBÓN, J., ZHANG, Y., WANG, B., XING, H., LIU, D. & CHIAPPE, L. M. (2018b). Diversity and evolution of the Confuciusornithidae: evidence from a new 131-million-year-old specimen from the Huajiyi Formation in NE China. *Journal of Asian Earth Sciences* **152**, 12-22.
- NAVARRO C. A., MARTIN-SILVERSTONE, E. & STUBBS, T. L. (2018). Morphometric assessment of pterosaur jaw disparity. *Royal Society Open Science* **5**, 172130.
- NEENAN J. M., RUTA, M., CLACK, J. A. & RAYFIELD, E. J. (2014). Feeding biomechanics in *Acanthostega* and across the fish - tetrapod transition. *Proceedings of the Royal Society B: Biological Sciences* **281**.
- NESBITT S. J., TURNER, A. H., ERICKSON, G. M. & NORELL, M. A. (2006). Prey choice and cannibalistic behaviour in the theropod *Coelophysis*. *Biology Letters* **2**, 611-614.
- NEVATTE R. J., WUERINGER, B. E., JACOB, D. E., PARK, J. M. & WILLIAMSON, J. E. (2017). First insights into the function of the sawshark rostrum through examination of rostral tooth microwear. *Journal of Fish Biology* **91**, 1582-1602.
- NICOLSON S. W. & FLEMING, P. A. (2014). Drinking problems on a 'simple' diet: physiological convergence in nectar-feeding birds. *Journal of Experimental Biology* **217**, 1015-1023.
- NORELL M. A., MAKOVICKY, P. J. & CURRIE, P. J. (2001). The beaks of ostrich dinosaurs. *Nature* **412**, 873-874.
- NORRIS D. R., MARRA, P. P., BOWEN, G. J., RATCLIFFE, L. M., ROYLE, J. A. & KYSER, T. K. (2006). Migratory connectivity of a widely distributed songbird, the American Redstart (*Setophaga ruticilla*). *Ornithological Monographs*, 14-28.
- NOVELLI E. L. B., DINIZ, Y. S., GALHARDI, C. M., EBAID, G. M. X., RODRIGUES, H. G., MANI, F., FERNANDES, A. A. H., CICOGNA, A. C. & NOVELLI FILHO, J. L. V. B. (2007). Anthropometrical parameters and markers of obesity in rats. *Laboratory animals* **41**, 111-119.
- NUDDS R. L. & DYKE, G. J. (2010). Narrow primary feather rachises in *Confuciusornis* and *Archaeopteryx* suggest poor flight ability. *Science* **328**, 887-889.
- NUDDS R. L. & OSWALD, S. A. (2007). An interspecific test of Allen's rule: evolutionary implications for endothermic species. *Evolution: International Journal of Organic Evolution* **61**, 2839-2848.

- O'CONNOR J. & DYKE, G. (2010a). A reassessment of *Sinornis santensis* and *Cathayornis yandica* (Aves: Enantiornithes). *Records of the Australian Museum* **62**, 7-20.
- O'CONNOR J., WANG, X. L., SULLIVAN, C., WANG, Y., ZHENG, X. T., HU, H., ZHANG, X. M. & ZHOU, Z. H. (2018). First report of gastroliths in the Early Cretaceous basal bird *Jeholornis*. *Cretaceous Research* **84**, 200-208.
- O'CONNOR J., ZHOU, Z. & XU, X. (2011a). Additional specimen of *Microraptor* provides unique evidence of dinosaurs preying on birds. *Proceedings of the National Academy of Sciences* **108**, 19662-19665.
- O'CONNOR J. K. (2019). The trophic habits of early birds. *Palaeogeography, Palaeoclimatology, Palaeoecology* **513**, 178-195.
- O'CONNOR J. K. & CHIAPPE, L. M. (2011b). A revision of enantiornithine (Aves: Ornithothoraces) skull morphology. *Journal of Systematic Palaeontology* **9**, 135-157.
- O'CONNOR J. K., GAO, K.-Q. & CHIAPPE, L. M. (2010b). A new ornithuromorph (Aves: Ornithothoraces) bird from the Jehol Group indicative of higher-level diversity. *Journal of Vertebrate Paleontology* **30**, 311-321.
- O'CONNOR J. K., LI, D. Q., LAMANNA, M. C., WANG, M., HARRIS, J. D., ATTERHOLT, J. & YOU, H. L. (2016a). A new Early Cretaceous enantiornithine (Aves, Ornithothoraces) from northwestern China with elaborate tail ornamentation. *Journal of Vertebrate Paleontology* **36**.
- O'CONNOR J. K., SUN, C., XU, X., WANG, X. & ZHOU, Z. (2012). A new species of *Jeholornis* with complete caudal integument. *Historical Biology* **24**, 29-41.
- O'CONNOR J. K., WANG, M. & HU, H. (2016b). A new ornithuromorph (Aves) with an elongate rostrum from the Jehol Biota, and the early evolution of rostralization in birds. *Journal of Systematic Palaeontology* **14**, 939-948.
- O'CONNOR J. K., WANG, X. L., ZHENG, X. T., HU, H., ZHANG, X. M. & ZHOU, Z. H. (2016c). An enantiornithine with a fan-shaped tail, and the Evolution of the rectricial complex in early birds. *Current Biology* **26**, 114-119.
- O'CONNOR J. K., WANG, X. R., CHIAPPE, L. M., GAO, C. L., MENG, Q. J., CHENG, X. D. & LIU, J. Y. (2009). Phylogenetic support for a specialized clade of Cretaceous enantiornithine birds with information from a new species. *Journal of Vertebrate Paleontology* **29**, 188-204.
- O'CONNOR J. K., ZHANG, Y. G., CHIAPPE, L. M., MENG, Q. J., LI, Q. G. & DI, L. (2013). A new enantiornithine from the Yixian Formation with the first recognized avian enamel specialization. *Journal of Vertebrate Paleontology* **33**, 1-12.
- O'CONNOR J. K., ZHENG, X., WANG, X., WANG, Y. & ZHOU, Z. (2014). Ovarian follicles shed new light on dinosaur reproduction during the transition towards birds. *National Science Review* **1**, 15-17.
- O'CONNOR J. K. & ZHOU, Z. (2019). The evolution of the modern avian digestive system: insights from paravian fossils from the Yanliao and Jehol biotas. *Palaeontology*.
- O'CONNOR J. K., ZHOU, Z. H. & ZHANG, F. C. (2010c). A reappraisal of *Boluochia zhengi* (Aves: Enantiornithes) and a discussion of intraclade diversity in the Jehol avifauna, China. *Journal of Systematic Palaeontology* **9**, 51-63.
- O'CONNOR J. K., WANG, M., ZHENG, X.-T., WANG, X.-L. & ZHOU, Z.-H. (2014). The histology of two female Early Cretaceous birds. *Vertebrata Palasiatica* **52**, 112-128.
- O'CONNOR J. K., WANG, X., SULLIVAN, C., ZHENG, X., TUBARO, P., ZHANG, X. & ZHOU, Z. (2013). Unique caudal plumage of *Jeholornis* and complex tail evolution in early birds. *Proceedings of the National Academy of Sciences* **110**, 17404-17408.



- ODA J., SAKAMOTO, J. & SAKANO, K. (2006). Mechanical evaluation of the skeletal structure and tissue of the woodpecker and its shock absorbing system. *JSME International Journal Series A: Solid Mechanics and Material Engineering* **49**, 390-396.
- OLSEN A. M. (2017). Feeding ecology is the primary driver of beak shape diversification in waterfowl. *Functional Ecology* **31**, 1985-1995.
- OPENSHAW G. H., D'AMORE, D. C., VIDAL-GARCÍA, M. & KEOGH, J. S. (2016). Combining geometric morphometric analyses of multiple 2D observation views improves interpretation of evolutionary allometry and shape diversification in monitor lizard (*Varanus*) crania. *Biological Journal of the Linnean Society* **120**, 539-552.
- ORO D., RUIZ, X., JOVER, L., PEDROCCHI, V. & GONZALEZSOLIS, J. (1997). Diet and adult time budgets of Audouin's gull *Larus audouinii* in response to changes in commercial fisheries. *IBIS* **139**, 631-637.
- OSTROM J. H. (1976). *Archaeopteryx* and the origin of birds. *Biological Journal of the Linnean Society* **8**, 91-182.
- OSTROM P. H., MACKO, S. A., ENGEL, M. H. & RUSSELL, D. A. (1993). Assessment of trophic structure of Cretaceous communities based on stable nitrogen isotope analyses. *Geology* **21**, 491-494.
- PARK R. & EPSTEIN, S. (1960). Carbon isotope fractionation during photosynthesis. *Geochimica et Cosmochimica Acta* **21**, 110-126.
- PECSICS T., LACZI, M., NAGY, G., KONDOR, T. & CSÖRGÖ, T. (2019). Analysis of skull morphometric characters in diurnal raptors (Accipitriformes and Falconiformes). *Ornis Hungarica* **27**, 117-131.
- PEI R., LI, Q., MENG, Q., NORELL, M. A. & GAO, K.-Q. (2017). New specimens of *Anchiornis huxleyi* (Theropoda: Paraves) from the Late Jurassic of northeastern China. *Bulletin of the American Museum of Natural History* **2017**, 1-67.
- PEREZ S. I., BERNAL, V. & GONZALEZ, P. N. (2006). Differences between sliding semi-landmark methods in geometric morphometrics, with an application to human craniofacial and dental variation. *Journal of Anatomy* **208**, 769-784.
- PETERS D. S. & GÖRGNER, E. (1992). A Comparative Study on the Claws of *Archaeopteryx*. In *Papers in avian paleontology honoring Pierce Brodkorb*, vol. 36. *Natural History Museum of Los Angeles County Science Series* (ed. K. C. Campbell Jr.), pp. 29-37. Natural History Museum of Los Angeles County, Los Angeles, CA, USA.
- PETEYA J. A., CLARKE, J. A., LI, Q. G., GAO, K. Q. & SHAWKEY, M. D. (2017). The plumage and colouration of an enantiornithine bird from the Early Cretaceous of China. *Palaeontology* **60**, 55-71.
- PETIE R. & MULLER, M. (2007). Curvature facilitates prey fixation in predatory insect claws. *Journal of Theoretical Biology* **244**, 565-575.
- PETTIGREW J. D. & FROST, B. J. (1985). A tactile fovea in the Scolopacidae? *Brain, Behavior and Evolution* **26**, 185-195.
- PHILLIPS P. K. & HEATH, J. E. (1992). Heat exchange by the pinna of the African elephant (*Loxodonta africana*). *Comparative Biochemistry and Physiology - Part A: Physiology* **101**, 693-699.
- PIETSCH T. W. (2005). Dimorphism, parasitism, and sex revisited: modes of reproduction among deep-sea ceratioid anglerfishes (Teleostei: Lophiiformes). *Ichthyological Research* **52**, 207-236.
- PIGOT A. L., SHEARD, C., MILLER, E. T., BREGMAN, T. P., FREEMAN, B. G., ROLL, U., SEDDON, N., TRISOS, C. H., WEEKS, B. C. & TOBIAS, J. A. (2020). Macroevolutionary convergence connects morphological form to ecological function in birds. *Nature Ecology & Evolution*.

- PIKE A. V. L. & MAITLAND, D. P. (2004). Scaling of bird claws. *Journal of Zoology* **262**, 73-81.
- PINEDA-MUNOZ S., LAZAGABASTER, I. A., ALROY, J. & EVANS, A. R. (2017). Inferring diet from dental morphology in terrestrial mammals. *Methods in Ecology and Evolution* **8**, 481-491.
- PITTMAN M., O'CONNOR, J., FIELD, D. J., TURNER, A. H., MA, W., MAKOVICKY, P. J. & XU, X. (2020a). Pennaraptoran systematics. In *Pennaraptoran Theropod Dinosaurs: Past Progress and New Frontiers* (ed. M. Pittman and X. Xu), pp. 7-36. Bulletin of the American Museum of Natural History.
- PITTMAN M., O'CONNOR, J., TSE, E., MAKOVICKY, P. J., FIELD, D. J., MA, W., TURNER, A. H., NORELL, M. A. & XU, X. (2020b). The fossil record of Mesozoic and Paleocene pennaraptorans. In *Pennaraptoran Theropod Dinosaurs: Past Progress and New Frontiers* (ed. M. Pittman and X. Xu), pp. 37-96. Bulletin of the American Museum of Natural History.
- PLAVCAN J. M. (1994). Comparison of four simple methods for estimating sexual dimorphism in fossils. *American Journal of Physical Anthropology* **94**, 465-476.
- POLLY P. D., STAYTON, C. T., DUMONT, E. R., PIERCE, S. E., RAYFIELD, E. J. & ANGIELCZYK, K. D. (2016). Combining geometric morphometrics and finite element analysis with evolutionary modeling: towards a synthesis. *Journal of Vertebrate Paleontology* **36**.
- POMEROY D. L. (2013). A morphological and taxonomic revision of the Early Cretaceous Sapeornithidae (Aves: Pygostylia) of Liaoning Province, China, California State University, Long Beach.
- PORRO L. B., HOLLIDAY, C. M., ANAPOL, F., ONTIVEROS, L. C., ONTIVEROS, L. T. & ROSS, C. F. (2011). Free body analysis, beam mechanics, and finite element modeling of the mandible of *Alligator mississippiensis*. *Journal of Morphology*, 1-28.
- PROSSER P. & HART, A. D. M. (2005). Assessing potential exposure of birds to pesticide-treated seeds. *Ecotoxicology* **14**, 679-691.
- PROVINI P., ZHOU, Z. & ZHANG, F. (2009). A new species of the basal bird *Sapeornis* from the Early Cretaceous of Liaoning, China. *Vertebrata Palasiatica* **47**, 194-207.
- PTACEK M. B. (2000). The role of mating preferences in shaping interspecific divergence in mating signals in vertebrates. *Behavioural Processes* **51**, 111-134.
- PURNELL M. A. (1995). Microwear on conodont elements and macrophagy in the first vertebrates. *Nature* **374**, 798.
- PURNELL M. A., HART, P. J. B., BAINES, D. C. & BELL, M. A. (2006). Quantitative analysis of dental microwear in threespine stickleback: a new approach to analysis of trophic ecology in aquatic vertebrates. *Journal of Animal Ecology* **75**, 967-977.
- QVARNSTRÖM M., WERNSTROM, J. V., PIECHOWSKI, R., TALANDA, M., AHLBERG, P. E. & NIEDZWIEDZKI, G. (2019). Beetle-bearing coprolites possibly reveal the diet of a Late Triassic dinosauriform. *Royal Society Open Science* **6**.
- RADHAKRISHNAN P. & MAO, J. J. (2004). Nanomechanical properties of facial sutures and sutural mineralization front. *Journal of dental research* **83**, 470-475.
- RALPH C. P., NAGATA, S. E. & RALPH, C. J. (1985). Analysis of droppings to describe diets of small birds. *Journal of Field Ornithology*, 165-174.
- RAMOS A. M. & WALKER, I. D. (1998). Raptors—Inroads to multifingered grasping. In *Proceedings. 1998 IEEE/RSJ International Conference on Intelligent Robots and Systems. Innovations in Theory, Practice and Applications (Cat. No. 98CH36190)*, vol. 1, pp. 467-475. IEEE.

- RAU G. H., AINLEY, D. G., BENGTSON, J. L., TORRES, J. J. & HOPKINS, T. L. (1992).  $^{15}\text{N}/^{14}\text{N}$  and  $^{13}\text{C}/^{12}\text{C}$  in Weddell Sea birds, seals, and fish - implications for diet and trophic structure. *Marine Ecology Progress Series* **84**, 1-8.
- RAUHUT O. W. M. (2014). New observations on the skull of *Archaeopteryx*. *Paläontologische Zeitschrift* **88**, 211-221.
- RAUHUT O. W. M., FOTH, C. & TISCHLINGER, H. (2018). The oldest *Archaeopteryx* (Theropoda: Avialiae): a new specimen from the Kimmeridgian/Tithonian boundary of Schamhaupten, Bavaria. *PeerJ* **6**, e4191.
- RAYFIELD E. J. (2004). Cranial mechanics and feeding in *Tyrannosaurus rex*. *Proceedings of the Royal Society B: Biological Sciences* **271**, 1451-1459.
- RAYFIELD E. J. (2005). Aspects of comparative cranial mechanics in the theropod dinosaurs *Coelophysis*, *Allosaurus* and *Tyrannosaurus*. *Zoological Journal of the Linnean Society* **144**, 309-316.
- RAYFIELD E. J. (2011a). Strain in the ostrich mandible during simulated pecking and validation of specimen-specific finite element models. *Journal of Anatomy* **218**, 47-58.
- RAYFIELD E. J. (2011b). Structural performance of tetanuran theropod skulls, with emphasis on the Megalosauridae, Spinosauridae and Carcharodontosauridae. *Special Papers in Palaeontology* **86**, 241-253.
- RAYFIELD E. J., MILNER, A. C., XUAN, V. B. & YOUNG, P. G. (2007). Functional morphology of spinosaur 'crocodile-mimic' dinosaurs. *Journal of Vertebrate Paleontology* **27**, 892-901.
- RAYFIELD E. J., NORMAN, D. B., HORNER, C. C., HORNER, J. R., SMITH, P. M., THOMASON, J. J. & UPCHURCH, P. (2001). Cranial design and function in a large theropod dinosaur. *Nature* **409**, 1033.
- REED D. A., PORRO, L. B., IRIARTE-DIAZ, J., LEMBERG, J. B., HOLLIDAY, C. M., ANAPOL, F. & ROSS, C. F. (2011). The impact of bone and suture material properties on mandibular function in *Alligator mississippiensis*: testing theoretical phenotypes with finite element analysis. *Journal of Anatomy* **218**, 59-74.
- REES J. & LINDGREN, J. (2005). Aquatic birds from the Upper Cretaceous (Lower Campanian) of Sweden and the biology and distribution of hesperornithiforms. *Palaeontology* **48**, 1321-1329.
- REYNARD L. M., HENDERSON, G. M. & HEDGES, R. E. M. (2010). Calcium isotope ratios in animal and human bone. *Geochimica et Cosmochimica Acta* **74**, 3735-3750.
- RICO-GUEVARA A. & ARAYA-SALAS, M. (2014). Bills as daggers? A test for sexually dimorphic weapons in a lekking hummingbird. *Behavioral Ecology* **26**, 21-29.
- RICO-GUEVARA A., SUSTAITA, D., GUSSEKLOO, S., OLSEN, A., BRIGHT, J., CORBIN, C. & DUDLEY, R. (2019). Feeding in Birds: Thriving in Terrestrial, Aquatic, and Aerial Niches. In *Feeding in Vertebrates: Evolution, Morphology, Behavior, Biomechanics* (ed. V. Bels), pp. 643-693. Springer International Publishing, Gewerbesrasse, Switzerland.
- ROBINSON B. W. & WILSON, D. S. (1998). Optimal foraging, specialization, and a solution to Liem's paradox. *American Naturalist* **151**, 223-235.
- ROHLF F. J. (1990). Morphometrics. *Annual Review of Ecology and Systematics* **21**, 299-316.
- ROSE K. D., WALKER, A. & JACOBS, L. L. (1981). Function of the mandibular tooth comb in living and extinct mammals. *Nature* **289**, 583-585.
- ROSENBERG K. V. & COOPER, R. J. (1990). Approaches to avian diet analysis. *Studies in Avian Biology* **13**, 80-90.

- ROUQUIER S. & GIORGI, D. (2007). Olfactory receptor gene repertoires in mammals. *Mutation Research - Fundamental and Molecular Mechanisms of Mutagenesis* **616**, 95-102.
- RYBCZYNSKI N., TIRABASSO, A., BLOSKIE, P., CUTHBERTSON, R. & HOLLIDAY, C. (2008). A three-dimensional animation model of *Edmontosaurus* (Hadrosauridae) for testing chewing hypotheses. *Palaeontologia Electronica* **11**, 9A.
- SAGE R. F., SAGE, T. L. & KOCACINAR, F. (2012). Photorespiration and the evolution of C4 photosynthesis. *Annual review of Plant Biology* **63**, 19-47.
- SAITTA E. T., KAYE, T. G. & VINTHER, J. (2019). Sediment-encased maturation: a novel method for simulating diagenesis in organic fossil preservation. *Palaeontology* **62**, 135-150.
- SAKAMOTO M. (2010). Jaw biomechanics and the evolution of biting performance in theropod dinosaurs. *Proceedings of the Royal Society B: Biological Sciences* **277**, 3327-3333.
- SAMUELS J. X. (2009). Cranial morphology and dietary habits of rodents. *Zoological Journal of the Linnean Society* **156**, 864-888.
- SANCHEZ J. (2010). Late Cretaceous (Cenomanian) Hesperornithiformes from the Pasquia Hills, Saskatchewan, Canada, Carleton University.
- SANKEY J. T., BRINKMAN, D. B., GUENTHER, M. & CURRIE, P. J. (2002). Small theropod and bird teeth from the Judith River Group (late Campanian), Alberta. *Journal of Paleontology* **76**, 751-763.
- SANPERA C., RUIZ, X., MORENO, R., JOVER, L. & WALDRON, S. (2007). Mercury and stable isotopes in feathers of Audouin's Gulls as indicators of feeding habits and migratory connectivity. *The Condor: Ornithological Applications* **109**, 268-275.
- SANTANA S. E., DUMONT, E. R. & DAVIS, J. L. (2010). Mechanics of bite force production and its relationship to diet in bats. *Functional Ecology* **24**, 776-784.
- SANZ J. L., CHIAPPE, L. M., FERNÁNDEZ-JALVO, Y., ORTEGA, F., SÁNCHEZ-CHILLÓN, B., POYATO-ARIZA, F. J. & PÉREZ-MORENO, B. P. (2001). An early Cretaceous pellet. *Nature* **409**, 998-1000.
- SANZ J. L., CHIAPPE, L. M., PÉREZ-MORENO, B. P., BUSCALIONI, A. D., MORATALLA, J. J., ORTEGA, F. & POYATO-ARIZA, F. J. (1996). An Early Cretaceous bird from Spain and its implications for the evolution of avian flight. *Nature* **382**, 442.
- SANZ J. L., CHIAPPE, L. M., PÉREZ-MORENO, B. P., MORATALLA, J. J., HERNÁNDEZ-CARRASQUILLA, F., BUSCALIONI, A. D., ORTEGA, F., POYATO-ARIZA, F. J., RASSKINGUTMAN, D. & MARTÍNEZ-DELCLOS, X. (1997). A nestling bird from the Lower Cretaceous of Spain: implications for avian skull and neck evolution. *Science* **276**, 1543-1546.
- SAVOLDI F., XU, B., TSOI, J. K. H., PAGANELLI, C. & MATINLINNA, J. P. (2018). Anatomical and mechanical properties of swine midpalatal suture in the premaxillary, maxillary, and palatine region. *Scientific Reports* **8**, 1-12.
- SCHAEFFER J., BENTON, M. J., RAYFIELD, E. J. & STUBBS, T. L. (2019). Morphological disparity in theropod jaws: comparing discrete characters and geometric morphometrics. *Palaeontology* **63**, 283-299.
- SCHEUHAMMER A. M., BOND, D. E., BURGESS, N. M. & RODRIGUE, J. (2003). Lead and stable lead isotope ratios in soil, earthworms, and bones of American woodcock (*Scolopax minor*) from eastern Canada. *Environmental Toxicology and Chemistry: An International Journal* **22**, 2585-2591.
- SCHEUHAMMER A. M. & TEMPLETON, D. M. (1998). Use of stable isotope ratios to distinguish sources of lead exposure in wild birds. *Ecotoxicology* **7**, 37-42.

- SCHILEO E., TADDEI, F., CRISTOFOLINI, L. & VICECONTI, M. (2008). Subject-specific finite element models implementing a maximum principal strain criterion are able to estimate failure risk and fracture location on human femurs tested in vitro. *Journal of Biomechanics* **41**, 356-367.
- SCHINO G. (2006). Grooming and agonistic support: a meta-analysis of primate reciprocal altruism. *Behavioral Ecology* **18**, 115-120.
- SCHOENER T. W. (1965). The evolution of bill size differences among sympatric congeneric species of birds. *Evolution* **19**, 189-213.
- SCHOENINGER M. J. & DENIRO, M. J. (1984). Nitrogen and carbon isotopic composition of bone collagen from marine and terrestrial animals. *Geochimica et Cosmochimica Acta* **48**, 625-639.
- SCHUBERT B. W. & UNGAR, P. S. (2005). Wear facets and enamel spalling in tyrannosaurid dinosaurs. *Acta Palaeontologica Polonica* **50**, 93-99.
- SCHULTZE H.-P. (1989). Three-dimensional muscle preservation in Jurassic fishes of Chile. *Andean Geology* **16**, 183-215.
- SCHWARTZ J. H. (1978). Homologies of the toothcomb. *American Journal of Physical Anthropology* **49**, 23-29.
- SCHWEITZER M. H., ZHENG, W. X., MOYER, A. E., SJOVALL, P. & LINDGREN, J. (2018). Preservation potential of keratin in deep time. *PLOS ONE* **13**, e0206569.
- SEKI Y., MACKEY, M. & MEYERS, M. A. (2012). Structure and micro-computed tomography-based finite element modeling of Toucan beak. *Journal of the Mechanical Behavior of Biomedical Materials* **9**, 1-8.
- SELLERS K. C., MIDDLETON, K. M., DAVIS, J. L. & HOLLIDAY, C. M. (2017). Ontogeny of bite force in a validated biomechanical model of the American alligator. *Journal of Experimental Biology* **220**, 2036-2046.
- SENER P. (2006). Comparison of forelimb function between *Deinonychus* and *Bambiraptor* (Theropoda: Dromaeosauridae). *Journal of Vertebrate Paleontology* **26**, 897-906.
- SERRANO F. J. (2020). ERRATUM.
- SERRANO F. J., PALMQVIST, P., CHIAPPE, L. M. & SANZ, J. L. (2017). Inferring flight parameters of Mesozoic avians through multivariate analyses of forelimb elements in their living relatives. *Paleobiology* **43**, 144-169.
- SERRANO F. J., PALMQVIST, P. & SANZ, J. L. (2015). Multivariate analysis of neognath skeletal measurements: implications for body mass estimation in Mesozoic birds. *Zoological Journal of the Linnean Society* **173**, 929-955.
- SHI H. & LI, L. (2019). Advances in Cladistic Analysis of Enantiornithes from China. *Open Journal of Nature Science* **07**, 387-397.
- SHUFELDT R. W. (1909). *Osteology of Birds*. University of the State of New York, Albany, NY, USA.
- SHYCHOSKI L. & SNIVELY, E. (2008a). Ecological implications of tyrannosaurid lower jaw ontogeny, biomechanical scaling and bite function. *Journal of Vertebrate Paleontology* **28**, 142A-142A.
- SHYCHOSKI L. & SNIVELY, E. (2008b). Ecological implications of tyrannosaurid lower jaw ontogeny, biomechanical scaling and bite function. In *Sixty-Eighth Annual Meeting of the Society of Vertebrate Paleontology*. Poster IV 53, Cleveland, Ohio, USA.
- SI G., DONG, Y., MA, Y. & ZHANG, Z. (2015). Shape similarities and differences in the skulls of scavenging raptors. *Zoological science* **32**, 171-178.
- SLOAD E. J. (2014). Microwear Analysis of Crab Claw Fingers: A Functional Morphological Approach, Kent State University.

- SMITH C. R. (1982). Food resource partitioning of fossorial Florida reptiles. *Wildlife Research Report* **13**, 173-178.
- SMITH D. K. (1998). A morphometric analysis of *Allosaurus*. *Journal of Vertebrate Paleontology* **18**, 126-142.
- SMITH K. K. (1993). The Form of the Feeding Apparatus in Terrestrial Vertebrates: Studies of Adaptation and Constraint. In *The Skull, Volume 3: Functional and Evolutionary Mechanisms* (ed. J. Hanken and B. K. Hall), pp. 150-196. The University of Chicago Press, Chicago, IL, USA.
- SMITH K. T. & SCANFERLA, A. (2016). Fossil snake preserving three trophic levels and evidence for an ontogenetic dietary shift. *Palaeobiodiversity and Palaeoenvironments* **96**, 589-599.
- SMITH R. J. & JUNGERS, W. L. (1997). Body mass in comparative primatology. *Journal of Human Evolution* **32**, 523-559.
- SMITH S. A. & SMITH, B. J. (1990). Normal xeroradiographic and radiographic anatomy of the red-tailed hawk (*Buteo jamaicensis*), with reference to other diurnal raptors. *Veterinary Radiology* **31**, 301-312.
- SNIVELY E., COTTON, J. R., RIDGELY, R. & WITMER, L. M. (2013). Multibody dynamics model of head and neck function in *Allosaurus* (Dinosauria, Theropoda). *Palaeontologia Electronica* **16**, 11A.
- SNIVELY E. & RUSSELL, A. P. (2007a). Functional morphology of neck musculature in the Tyrannosauridae (Dinosauria, Theropoda) as determined via a hierarchical inferential approach. *Zoological Journal of the Linnean Society* **151**, 759-808.
- SNIVELY E. & RUSSELL, A. P. (2007b). Functional variation of neck muscles and their relation to feeding style in Tyrannosauridae and other large theropod dinosaurs. *The Anatomical Record: Advances in Integrative Anatomy and Evolutionary Biology* **290**, 934-957.
- SOONS J., GENBRUGGE, A., PODOS, J., ADRIAENS, D., AERTS, P., DIRCKX, J. & HERREL, A. (2015). Is beak morphology in Darwin's finches tuned to loading demands? *PLOS ONE* **10**, e0129479.
- SOONS J., HERREL, A., GENBRUGGE, A., ADRIAENS, D., AERTS, P. & DIRCKX, J. (2012a). Multi-layered bird beaks: a finite-element approach towards the role of keratin in stress dissipation. *Journal of the Royal Society Interface* **9**, 1787-1796.
- SOONS J., HERREL, A., GENBRUGGE, A., AERTS, P., PODOS, J., ADRIAENS, D., DE WITTE, Y., JACOBS, P. & DIRCKX, J. (2010). Mechanical stress, fracture risk and beak evolution in Darwin's ground finches (*Geospiza*). *Philosophical Transactions of the Royal Society B: Biological Sciences* **365**, 1093-1098.
- SOONS J., LAVA, P., DEBRUYNE, D. & DIRCKX, J. (2012b). Full-field optical deformation measurement in biomechanics: digital speckle pattern interferometry and 3D digital image correlation applied to bird beaks. *Journal of the Mechanical Behavior of Biomedical Materials* **14**, 186-191.
- STANSFIELD E., PARKER, J. & O'HIGGINS, P. (2018). A sensitivity study of human mandibular biting simulations using finite element analysis. *Journal of Archaeological Science-Reports* **22**, 420-432.
- STAUFENBERG G., GRAUPNER, N. & MÜSSIG, J. (2015). Impact and hardness optimisation of composite materials inspired by the babassu nut (*Orbignya speciosa*). *Bioinspiration & Biomimetics* **10**, 056006.
- STAYTON C. T. (2006). Testing hypotheses of convergence with multivariate data: morphological and functional convergence among herbivorous lizards. *Evolution* **60**, 824-841.

- STERNBERG C. M. (1935). Hooded Hadrosaurs of the Belly River Series of the Upper Cretaceous: A Comparison, With Descriptions of New Species. In *National Museum of Canada* (ed. W. H. Collins). Patenaude, J. O., I.S.O., Ottawa, Canada.
- STEVENSON R. D. (1986). Allen's rule in North American rabbits (*Sylvilagus*) and hares (*Lepus*) is an exception, not a rule. *Journal of Mammalogy* **67**, 312-316.
- SULLIVAN B. L., WOOD, C. L., ILIFF, M. J., BONNEY, R. E., FINK, D. & KELLING, S. (2009). eBird: a citizen-based bird observation network in the biological sciences. *Biological Conservation* **142**, 2282-2292.
- SUN Y., SI, G., WANG, X., WANG, K. & ZHANG, Z. (2018). Geometric morphometric analysis of skull shape in the Accipitridae. *Zoomorphology* **137**, 445-456.
- SURKOV M. V. & BENTON, M. J. (2008). Head kinematics and feeding adaptations of the Permian and Triassic dicynodonts. *Journal of Vertebrate Paleontology* **28**, 1120-1129.
- SUSTAITA D. (2008). Musculoskeletal underpinnings to differences in killing behavior between North American accipiters (Falconiformes : Accipitridae) and falcons (Falconidae). *Journal of Morphology* **269**, 283-301.
- SUSTAITA D., GLOUMAKOV, Y., TSANG, L. R. & DOLLAR, A. M. (2019). Behavioral correlates of semi-zygodactyly in Ospreys (*Pandion haliaetus*) based on analysis of internet images. *PeerJ* **7**, e6243.
- SUSTAITA D. & HERTEL, F. (2010). In vivo bite and grip forces, morphology and prey-killing behavior of North American accipiters (Accipitridae) and falcons (Falconidae). *Journal of Experimental Biology* **213**, 2617-2628.
- SUSTAITA D., POUYDEBAT, E., MANZANO, A., ABDALA, V., HERTEL, F. & HERREL, A. (2013). Getting a grip on tetrapod grasping: form, function, and evolution. *Biological Reviews* **88**, 380-405.
- SUSTAITA D. & RUBEGA, M. A. (2014). The anatomy of a shrike bite: bill shape and bite performance in loggerhead shrikes. *Biological Journal of the Linnean Society* **112**, 485-498.
- SVANBERG F., MATEO, R., HILLSTRÖM, L., GREEN, A. J., TAGGART, M. A., RAAB, A. & MEHARG, A. A. (2006). Lead isotopes and lead shot ingestion in the globally threatened marbled teal (*Marmaronetta angustirostris*) and white-headed duck (*Oxyura leucocephala*). *Science of the Total Environment* **370**, 416-424.
- SYMONDS M. R. E. & TATTERSALL, G. J. (2010). Geographical variation in bill size across bird species provides evidence for Allen's rule. *American Naturalist* **176**, 188-197.
- ȚĂLU Ș., STACH, S., MÉNDEZ, A., TREJO, G. & ȚĂLU, M. (2014). Multifractal characterization of nanostructure surfaces of electrodeposited Ni-P coatings. *Journal of The Electrochemical Society* **161**, D44-D47.
- TARQUINI S. D., CHEMISQUY, M. A., LADEVÈZE, S. & PREVOSTI, F. J. (2019). The scope of traditional and geometric morphometrics for inferences of diet in carnivorous fossil mammals. *Ameghiniana* **56**, 307-318.
- TATTERSALL G. J., ANDRADE, D. V. & ABE, A. S. (2009). Heat exchange from the toucan bill reveals a controllable vascular thermal radiator. *Science* **325**, 468-470.
- TATTERSALL H. G. & TAPPIN, G. (1966). The work of fracture and its measurement in metals, ceramics and other materials. *Journal of Materials Science* **1**, 296-301.
- TAYLOR C. R. (1966). The vascularity and possible thermoregulatory function of the horns in goats. *Physiological Zoology* **39**, 127-139.
- TAYLOR R. J. (2013). Predators and Predation. In *Predation*, pp. 1-5. Springer Science & Business Media, Berlin, Germany.

- THERRIEN F., QUINNEY, A., TANAKA, K. & ZELENITSKY, D. K. (2016). Accuracy of mandibular force profiles for bite force estimation and feeding behavior reconstruction in extant and extinct carnivorans. *Journal of Experimental Biology* **219**, 3738-3749.
- THOMAS M. A. (2014). Aspects of the Trophic Ecology of an Invertivorous Snake Community, Eastern Illinois University.
- THOMPSON D. W. (1942). *On growth and form*, 2nd ed. Macmillan, Cambridge University Pre, Oxford, UK.
- THOMPSON R. M., HEMBERG, M., STARZOMSKI, B. M. & SHURIN, J. B. (2007). Trophic levels and trophic tangles: the prevalence of omnivory in real food webs. *Ecology* **88**, 612-617.
- THOUZEAU C., PETERS, G., LE BOHEC, C. & LE MAHO, Y. (2004). Adjustments of gastric pH, motility and temperature during long-term preservation of stomach contents in free-ranging incubating king penguins. *Journal of Experimental Biology* **207**, 2715-2724.
- THULBORN R. A. & HAMLEY, T. L. (1985). A new palaeoecological role for *Archaeopteryx*. *The Beginnings of Birds*, 81-89.
- TIETZE D. T. (2018). Introduction: Studying Birds in Time and Space. In *Bird Species: How They Arise, Modify and Vanish* (ed. D. T. Tietze), pp. 1-7. Springer International Publishing, Gewerbesrasse, Switzerland.
- TILKENS M. J., WALL-SCHEFFLER, C., WEAVER, T. D. & STEUDEL-NUMBERS, K. (2007). The effects of body proportions on thermoregulation: an experimental assessment of Allen's rule. *Journal of Human Evolution* **53**, 286-291.
- TINIUS A. & RUSSELL, A. P. (2017). Points on the curve: an analysis of methods for assessing the shape of vertebrate claws. *Journal of Morphology* **278**, 150-169.
- TOKITA M., KIYOSHI, T. & ARMSTRONG, K. N. (2007). Evolution of craniofacial novelty in parrots through developmental modularity and heterochrony. *Evolution & Development* **9**, 590-601.
- TOKITA M., YANO, W., JAMES, H. F. & ABZHANOV, A. (2017). Cranial shape evolution in adaptive radiations of birds: comparative morphometrics of Darwin's finches and Hawaiian honeycreepers. *Philosophical Transactions of the Royal Society B: Biological Sciences* **372**, 20150481.
- TORICES A., WILKINSON, R., ARBOUR, V. M., RUIZ-OMENACA, J. I. & CURRIE, P. J. (2018). Puncture-and-pull biomechanics in the teeth of predatory coelurosaurian dinosaurs. *Current Biology* **28**, 1467-1474.
- TORNBERG R. & REIF, V. (2007). Assessing the diet of birds of prey: a comparison of prey items found in nests and images. *Ornis Fennica* **84**, 21.
- TRINAJSTIC K., MARSHALL, C., LONG, J. & BIFIELD, K. (2007). Exceptional preservation of nerve and muscle tissues in Late Devonian placoderm fish and their evolutionary implications. *Biology Letters* **3**, 197-200.
- TSANG L. R., WILSON, L. A. B., LEDOGAR, J., WROE, S., ATTARD, M. & SANSALONE, G. (2019). Raptor talon shape and biomechanical performance are controlled by relative prey size but not by allometry. *Scientific Reports* **9**.
- TSUIHJI T. (2005). Homologies of the transversospinalis muscles in the anterior presacral region of Sauria (crown Diapsida). *Journal of Morphology* **263**, 151-178.
- TSUIHJI T. (2007). Homologies of the Longissimus, Iliocostalis, and hypaxial muscles in the anterior presacral region of extant Diapsida. *Journal of Morphology* **268**, 986-1020.
- TSUIHJI T. (2010). Reconstructions of the axial muscle insertions in the occipital region of dinosaurs: evaluations of past hypotheses on Marginocephalia and Tyrannosauridae using the extant



- phylogenetic bracket approach. *The Anatomical Record: Advances in Integrative Anatomy and Evolutionary Biology* **293**, 1360-1386.
- TURNER K. L., ABERNETHY, E. F., CONNER, L. M., RHODES JR, O. E. & BEASLEY, J. C. (2017). Abiotic and biotic factors modulate carrion fate and vertebrate scavenging communities. *Ecology* **98**, 2413-2424.
- UGURAL A. C. & FENSTER, S. K. (2012). *Advanced Mechanics of Materials and Applied Elasticity*, 5th edition. Pearson Education, London, UK.
- ULLREY D. E. (2018). Nutrient Requirements: Carnivores. In *Encyclopedia of Animal Science* (ed. W. G. Pond, D. E. Ullrey and C. K. Baer), pp. 670-673. CRC Press, Boca Raton, FL, USA.
- UNGAR P. S. (2015). Mammalian dental function and wear: a review. *Biosurface and Biotribology* **1**, 25-41.
- UNGAR P. S. (2018). Tooth Surface Topography: A Scale-Sensitive Approach with Implications for Inferring Dental Adaptation and Diet. In *New Geospatial Approaches to the Anthropological Sciences* (ed. R. L. Anemone and G. C. Conroy). University of New Mexico Press, Albuquerque, NM, USA.
- UNGAR P. S. (2019). Inference of diets of early hominins from primate molar form and microwear. *Journal of dental research* **98**, 398-405.
- UNGAR P. S. & HLUSKO, L. J. (2016). The evolutionary path of least resistance. *Science* **353**, 29-30.
- URANO Y., TANOUE, K., MATSUMOTO, R., KAWABE, S., OHASHI, T. & FUJIWARA, S. (2018). How does the curvature of the upper beak bone reflect the overlying rhinotheca morphology? *Journal of Morphology* **279**, 636-647.
- UYEDA J. C., ZENIL-FERGUSON, R. & PENNELL, M. W. (2018). Rethinking phylogenetic comparative methods. *Systematic Biology* **67**, 1091-1109.
- VAN BAAL R. R., JANSSEN, R., VAN DER LUBBE, H. J. L., SCHULP, A. S., JAGT, J. W. M. & VONHOF, H. B. (2013). Oxygen and carbon stable isotope records of marine vertebrates from the type Maastrichtian, The Netherlands and northeast Belgium (Late Cretaceous). *Palaeogeography, Palaeoclimatology, Palaeoecology* **392**, 71-78.
- VAN DE FLIERDT T., ROBINSON, L. F., ADKINS, J. F., HEMMING, S. R. & GOLDSTEIN, S. L. (2006). Temporal stability of the neodymium isotope signature of the Holocene to glacial North Atlantic. *Paleoceanography* **21**, PA4102.
- VAN DER LEEUW A. H. J., BOUT, R. G. & ZWEERS, G. A. (2001). Control of the cranio-cervical system during feeding in birds. *American Zoologist* **41**, 1352-1363.
- VAN DER MEIJ M. A. A. (2004). A Tough Nut to Crack. Adaptations to seed cracking in finches, Leiden University.
- VAN DER MEIJ M. A. A. & BOUT, R. G. (2004). Scaling of jaw muscle size and maximal bite force in finches. *Journal of Experimental Biology* **207**, 2745-2753.
- VAN VALKENBURGH B. & MOLNAR, R. E. (2002). Dinosaurian and mammalian predators compared. *Paleobiology* **28**, 527-543.
- VERNER K. A., LEHNER, M., LAMAS, L. P. & MAIN, R. P. (2016). Experimental tests of planar strain theory for predicting bone cross-sectional longitudinal and shear strains. *Journal of Experimental Biology* **219**, 3082-3090.
- VERWAIJEN D., VAN DAMME, R. & HERREL, A. (2002). Relationships between head size, bite force, prey handling efficiency and diet in two sympatric lacertid lizards. *Functional Ecology* **16**, 842-850.

- VITT L. J. & PIANKA, E. R. (2005). Deep history impacts present-day ecology and biodiversity. *Proceedings of the National Academy of Sciences* **102**, 7877-7881.
- VOGEL S. (2013). Biological Materials: Cracks and Composites. In *Comparative Biomechanics: Life's Physical World*, pp. 329-346. Princeton University Press, Princeton, NJ, USA.
- VOTIER S. C., BEARHOP, S., RATCLIFFE, N. & FURNESS, R. W. (2001). Pellets as indicators of diet in Great Skuas *Catharacta skua*. *Bird Study* **48**, 373-376.
- WAINWRIGHT P. C., ALFARO, M. E., BOLNICK, D. I. & HULSEY, C. D. (2005). Many-to-one mapping of form to function: a general principle in organismal design? *Integrative and Comparative Biology* **45**, 256-262.
- WANG C., SONG, Y., SONG, S., JI, Q., CHIANG, C., MENG, Q., LI, H., HSIAO, K., LU, Y. & SHEW, B. (2015a). Evolution and function of dinosaur teeth at ultramicrostructural level revealed using synchrotron transmission X-ray microscopy. *Scientific Reports* **5**, 15202.
- WANG J., HAO, X., KUNDRÁT, M., LIU, Z., UESUGI, K., JURAŠEKOVÁ, Z., GUO, B., HOSHINO, M., LI, Y. & MONFROY, Q. (2019a). Bone tissue histology of the Early Cretaceous bird Yanornis: evidence for a diphyletic origin of modern avian growth strategies within Ornithuromorpha. *Historical Biology*, DOI: 10.1080/08912963.2019.1593405.
- WANG J., ZOU, D., LI, Z., HUANG, P., LI, D., SHAO, Y., WANG, H. & CHEN, Y. (2014a). Mechanical properties of cranial bones and sutures in 1–2-year-old infants. *Medical Science Monitor: International Medical Journal of Experimental and Clinical Research* **20**, 1808.
- WANG M., LI, Z. & ZHOU, Z. (2017a). Insight into the growth pattern and bone fusion of basal birds from an Early Cretaceous enantiornithine bird. *Proceedings of the National Academy of Sciences* **114**, 11470-11475.
- WANG M. & LIU, D. (2016a). Taxonomical reappraisal of Cathayornithidae (Aves: Enantiornithes). *Journal of Systematic Palaeontology* **14**, 29-47.
- WANG M., O'CONNOR, J. K. & ZHOU, Z. H. (2014b). A new robust enantiornithine bird from the Lower Cretaceous of China with scansorial adaptations. *Journal of Vertebrate Paleontology* **34**, 657-671.
- WANG M., O'CONNOR, J. & ZHOU, Z. (2019b). A taxonomical revision of the Confuciusornithiformes (Aves: Pygostylia). *Vertebrata Palasiatica* **10**, 1-37.
- WANG M., O'CONNOR, J. K., PAN, Y. & ZHOU, Z. (2017b). A bizarre Early Cretaceous enantiornithine bird with unique cranial feathers and an ornithuromorph plough-shaped pygostyle. *Nature Communications* **8**, 14141.
- WANG M., O'CONNOR, J. K., XU, X. & ZHOU, Z. (2019c). A new Jurassic scansoriopterygid and the loss of membranous wings in theropod dinosaurs. *Nature* **569**, 256-259.
- WANG M., STIDHAM, T. A. & ZHOU, Z. (2018). A new clade of basal Early Cretaceous pygostylid birds and developmental plasticity of the avian shoulder girdle. *Proceedings of the National Academy of Sciences* **115**, 10708-10713.
- WANG M., ZHENG, X., O'CONNOR, J. K., LLOYD, G. T., WANG, X., WANG, Y., ZHANG, X. & ZHOU, Z. (2015b). The oldest record of Ornithuromorpha from the Early Cretaceous of China. *Nature Communications* **6**, 6987.
- WANG M. & ZHOU, Z. (2016b). A new adult specimen of the basalmost ornithuromorph bird *Archaeorhynchus spathula* (Aves: Ornithuromorpha) and its implications for early avian ontogeny. *Journal of Systematic Palaeontology* **15**, 1-18.

- WANG M. & ZHOU, Z. (2017c). A morphological study of the first known piscivorous enantiornithine bird from the Early Cretaceous of China. *Journal of Vertebrate Paleontology* **37**, e1278702.
- WANG M. & ZHOU, Z. (2019d). A new confuciusornithid (Aves: Pygostylia) from the Early Cretaceous increases the morphological disparity of the Confuciusornithidae. *Zoological Journal of the Linnean Society* **185**, 417-430.
- WANG M., ZHOU, Z., O'CONNOR, J. K. & ZELENKOV, N. V. (2014c). A new diverse enantiornithine family (Bohaiornithidae fam. nov.) from the Lower Cretaceous of China with information from two new species. *Vertebrata Palasiatica* **52**, 31-76.
- WANG M. & ZHOU, Z. H. (2019e). A new enantiornithine (Aves: Ornithothoraces) with completely fused premaxillae from the Early Cretaceous of China. *Journal of Systematic Palaeontology* **17**, 1079-1092.
- WANG M., ZHOU, Z. H. & SULLIVAN, C. (2016c). A fish-eating enantiornithine bird from the Early Cretaceous of China provides evidence of modern avian digestive features. *Current Biology* **26**, 1170-1176.
- WANG M., ZHOU, Z. H. & XU, G. H. (2014d). The first enantiornithine bird from the Upper Cretaceous of China. *Journal of Vertebrate Paleontology* **34**, 135-145.
- WANG S., STIEGLER, J., WU, P. & CHUONG, C.-M. (2020a). Tooth vs beak: the evolutionary developmental control of the avian feeding apparatus. In *Pennaraptoran Theropod Dinosaurs: Past Progress and New Frontiers* (ed. M. Pittman and X. Xu), pp. 205-228. Bulletin of the American Museum of Natural History.
- WANG X., CHIAPPE, L. M., TENG, F. & JI, Q. (2013). *Xinghaiornis lini* (Aves: Ornithothoraces) from the Early Cretaceous of Liaoning: an example of evolutionary mosaic in early birds. *Acta Geologica Sinica (English Edition)* **87**, 686-689.
- WANG X., HUANG, J., KUNDRÁT, M., CAU, A., LIU, X., WANG, Y. & JU, S. (2020b). A new jeholornithiform exhibits the earliest appearance of the fused sternum and pelvis in the evolution of avialan dinosaurs. *Journal of Asian Earth Sciences*, 104401.
- WANG X., O'CONNOR, J. K., MAINA, J. N., PAN, Y., WANG, M., WANG, Y., ZHENG, X. & ZHOU, Z. (2018). *Archaeorhynchus* preserving significant soft tissue including probable fossilized lungs. *Proceedings of the National Academy of Sciences* **115**, 11555-11560.
- WANG X., PITTMAN, M., ZHENG, X., KAYE, T. G., FALK, A. R., HARTMAN, S. A. & XU, X. (2017d). Basal paravian functional anatomy illuminated by high-detail body outline. *Nature Communications* **8**, 14576.
- WANG X., ZHANG, Z., GAO, C., HOU, L., MENG, Q. & LIU, J. (2010a). A new enantiornithine bird from the Early Cretaceous of Western Liaoning, China. *The Condor: Ornithological Applications* **112**, 432-437.
- WANG X. L., O'CONNOR, J. K., ZHENG, X. T., WANG, M., HU, H. & ZHOU, Z. H. (2014e). Insights into the evolution of rachis dominated tail feathers from a new basal enantiornithine (Aves: Ornithothoraces). *Biological Journal of the Linnean Society* **113**, 805-819.
- WANG X. R., O'CONNOR, J. K., ZHAO, B., CHIAPPE, L. M., GAO, C. L. & CHENG, X. D. (2010b). New Species of Enantiornithes (Aves: Ornithothoraces) from the Qiaotou Formation in Northern Hebei, China. *Acta Geologica Sinica (English Edition)* **84**, 247-256.
- WANG X. R., ZHAO, B., SHEN, C. Z., LIU, S. Z., GAO, C. L., CHENG, X. D. & ZHANG, F. J. (2015c). New material of *Longipteryx* (Aves: Enantiornithes) from the Lower Cretaceous Yixian Formation of China with the first recognized avian tooth crenulations. *Zootaxa* **3941**, 565-578.

- WANG Y., HU, H., O'CONNOR, J. K., WANG, M., XU, X., ZHOU, Z., WANG, X. & ZHENG, X. (2017e). A previously undescribed specimen reveals new information on the dentition of *Sapeornis chaoyangensis*. *Cretaceous Research* **74**, 1-10.
- WANG Y., WANG, M., O'CONNOR, J. K., WANG, X. L., ZHENG, X. T. & ZHANG, X. M. (2016d). A new Jehol enantiornithine bird with three-dimensional preservation and ovarian follicles. *Journal of Vertebrate Paleontology* **36**.
- WARD A. B., WEIGL, P. D. & CONROY, R. M. (2002). Functional morphology of raptor hindlimbs: implications for resource partitioning. *The Auk: Ornithological Advances* **119**, 1052-1063.
- WARTON D. I. & HUI, F. K. C. (2011). The arcsine is asinine: the analysis of proportions in ecology. *Ecology* **92**, 3-10.
- WATSON J. (2010). *The Golden Eagle*. Bloomsbury Publishing, London, UK.
- WEI Z.-Y. & LI, L. (2011). Discovery of a new enantiornithine bird from Lower Cretaceous of western Liaoning, China. *Global Geology*, 655-662.
- WELNHOFER P. (2010). A short history of research on *Archaeopteryx* and its relationship with dinosaurs. *Geological Society, London, Special Publications* **343**, 237-250.
- WHETSTONE K. N. (1983). Braincase of Mesozoic birds: I. new preparation of the "London" *Archaeopteryx*. *Journal of Vertebrate Paleontology* **2**, 439-452.
- WILMAN H., BELMAKER, J., SIMPSON, J., DE LA ROSA, C., RIVADENEIRA, M. M. & JETZ, W. (2014). EltonTraits 1.0: Species-level foraging attributes of the world's birds and mammals. *Ecology* **95**, 2027.
- WILSON L. E., CHIN, K. & CUMBAA, S. L. (2016). A new hesperornithiform (Aves) specimen from the Late Cretaceous Canadian High Arctic with comments on high-latitude hesperornithiform diet. *Canadian Journal of Earth Sciences* **53**, 1476-1483.
- WINKLER D. E., SCHULZ-KORNAS, E., KAISER, T. M. & TÜTKEN, T. (2019). Dental microwear texture reflects dietary tendencies in extant Lepidosauria despite their limited use of oral food processing. *Proceedings of the Royal Society B: Biological Sciences* **286**, 20190544.
- WISEMAN K. D., GREENE, H. W., KOO, M. S. & LONG, D. J. (2019). Feeding ecology of a generalist predator, the California kingsnake (*Lampropeltis californiae*): why rare prey matter. *Herpetological Conservation and Biology* **14**, 1-30.
- WOOD J. R., RAWLENCE, N. J., ROGERS, G. M., AUSTIN, J. J., WORTHY, T. H. & COOPER, A. (2008). Coprolite deposits reveal the diet and ecology of the extinct New Zealand megaherbivore moa (Aves, Dinornithiformes). *Quaternary Science Reviews* **27**, 2593-2602.
- WROE S. (2008). Cranial mechanics compared in extinct marsupial and extant African lions using a finite-element approach. *Journal of Zoology* **274**, 332-339.
- WROE S., CLAUSEN, P., MCHENRY, C., MORENO, K. & CUNNINGHAM, E. (2007). Computer simulation of feeding behaviour in the thylacine and dingo as a novel test for convergence and niche overlap. *Proceedings of the Royal Society B: Biological Sciences* **274**, 2819-2828.
- WROE S., MCHENRY, C. & THOMASON, J. (2005). Bite club: comparative bite force in big biting mammals and the prediction of predatory behaviour in fossil taxa. *Proceedings of the Royal Society B: Biological Sciences* **272**, 619-625.
- XIA D., KYES, R. C., WANG, X., SUN, B., SUN, L. & LI, J.-H. (2019). Grooming networks reveal intra- and intersexual social relationships in *Macaca thibetana*. *Primates* **60**, 223-232.

- XING L., O'CONNOR, J. K., MCKELLAR, R. C., CHIAPPE, L. M., TSENG, K., LI, G. & BAI, M. (2017). A mid-Cretaceous enantiornithine (Aves) hatchling preserved in Burmese amber with unusual plumage. *Gondwana Research* **49**, 264-277.
- XING L. D., BELL, P. R., PERSONS, W. S., JI, S. A., MIYASHITA, T., BURNS, M. E., JI, Q. & CURRIE, P. J. (2012). Abdominal contents from two large Early Cretaceous compsognathids (Dinosauria: Theropoda) demonstrate feeding on confuciusornithids and dromaeosaurids. *PLOS ONE* **7**, e44012.
- XU X., YOU, H., DU, K. & HAN, F. (2011). An *Archaeopteryx*-like theropod from China and the origin of Avialae. *Nature* **475**, 465-470.
- XU X., ZHENG, X., SULLIVAN, C., WANG, X., XING, L., WANG, Y., ZHANG, X., O'CONNOR, J. K., ZHANG, F. & PAN, Y. (2015). A bizarre Jurassic maniraptoran theropod with preserved evidence of membranous wings. *Nature* **521**, 70-73.
- XU X., ZHOU, Z. & WANG, X. (2000). The smallest known non-avian theropod dinosaur. *Nature* **408**, 705.
- XU X., ZHOU, Z., WANG, Y. & WANG, M. (2020). Study on the Jehol Biota: recent advances and future prospects. *Science China Earth Sciences*.
- XU X., ZHOU, Z. H., DUDLEY, R., MACKEM, S., CHUONG, C. M., ERICKSON, G. M. & VARRICCHIO, D. J. (2014). An integrative approach to understanding bird origins. *Science* **346**, 1341-+.
- YANG T.-R. & SANDER, P. M. (2018). The origin of the bird's beak: new insights from dinosaur incubation periods. *Biology Letters* **14**, 20180090.
- YOSIBASH Z., TAL, D. & TRABELSI, N. (2010). Predicting the yield of the proximal femur using high-order finite-element analysis with inhomogeneous orthotropic material properties. *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences* **368**, 2707-2723.
- YOUNG N. M., WAGNER, G. P. & HALLGRÍMSSON, B. (2010). Development and the evolvability of human limbs. *Proceedings of the National Academy of Sciences* **107**, 3400-3405.
- YUAN C. (2008). A new genus and species of Sapeornithidae from Lower Cretaceous in western Liaoning, China. *Acta Geologica Sinica (English Edition)* **82**, 48-55.
- ZANI P. A. (2000). The comparative evolution of lizard claw and toe morphology and clinging performance. *Journal of Evolutionary Biology* **13**, 316-325.
- ZANNO L. E. & MAKOVICKY, P. J. (2011). Herbivorous ecomorphology and specialization patterns in theropod dinosaur evolution. *Proceedings of the National Academy of Sciences* **108**, 232-237.
- ZELDITCH M. L., SWIDERSKI, D. L., SHEETS, H. D. & FINK, W. L. (2004). *Geometric Morphometrics for Biologists: A Primer*, 1st edition. Academic Press, San Diego, CA, USA.
- ZELENKOV N. V. & AVERIANOV, A. O. (2016). A historical specimen of enantiornithine bird from the Early Cretaceous of Mongolia representing a new taxon with a specialized neck morphology. *Journal of Systematic Palaeontology* **14**, 319-338.
- ZHANG F., ERICSON, P. G. P. & ZHOU, Z. (2004). Description of a new enantiornithine bird from the Early Cretaceous of Hebei, northern China. *Canadian Journal of Earth Sciences* **41**, 1097-1107.
- ZHANG F., ZHOU, Z. & BENTON, M. J. (2008). A primitive confuciusornithid bird from China and its implications for early avian flight. *Science in China Series D: Earth Sciences* **51**, 625-639.
- ZHANG F., ZHOU, Z., XU, X. & WANG, X. (2002). A juvenile coelurosaurian theropod from China indicates arboreal habits. *Naturwissenschaften* **89**, 394-398.
- ZHANG F., ZHOU, Z., XU, X., WANG, X. & SULLIVAN, C. (2008). A bizarre Jurassic maniraptoran from China with elongate ribbon-like feathers. *Nature* **455**, 1105-1108.

- ZHANG J. J., RAYNER, M., VICKERS, S., LANDERS, T., SAGAR, R., STEWART, J. & DUNPHY, B. (2019). GPS telemetry for small seabirds: using hidden Markov models to infer foraging behaviour of common diving petrels (*Pelecanoides urinatrix urinatrix*). *Emu - Austral Ornithology* **119**, 126-137.
- ZHANG Y. G., O'CONNOR, J., DI, L., MENG, Q. J., SIGURDSEN, T. & CHIAPPE, L. M. (2014). New information on the anatomy of the Chinese Early Cretaceous Bohaiornithidae (Aves: Enantiornithes) from a subadult specimen of *Zhouornis hani*. *PeerJ* **2**, e407.
- ZHANG Z. H., CHIAPPE, L. M., HAN, G. & CHINSAMY-TURAN, A. (2013). A large bird from the Early Cretaceous of China: new information on the skull of enantiornithines. *Journal of Vertebrate Paleontology* **33**, 1176-1189.
- ZHANG Z. H., HOU, L. H., YOSHIKASU, H., O'CONNOR, J., MARTIN, L. D. & CHIAPPE, L. M. (2006). The first Mesozoic heterodactyl bird from China. *Acta Geologica Sinica (English Edition)* **80**, 631-635.
- ZHENG X., MARTIN, L. D., ZHOU, Z., BURNHAM, D. A., ZHANG, F. & MIAO, D. (2011). Fossil evidence of avian crops from the Early Cretaceous of China. *Proceedings of the National Academy of Sciences* **108**, 15904-15907.
- ZHENG X., O'CONNOR, J. K., HUCHZERMAYER, F., WANG, X., WANG, Y., ZHANG, X. & ZHOU, Z. (2014). New specimens of *Yanornis* indicate a piscivorous diet and modern alimentary canal. *PLOS ONE* **9**, e95036.
- ZHENG X., O'CONNOR, J. K., WANG, X., WANG, Y. & ZHOU, Z. (2018a). Reinterpretation of a previously described Jehol bird clarifies early trophic evolution in the Ornithuromorpha. *Proceedings of the Royal Society B: Biological Sciences* **285**, 20172494.
- ZHENG X., O'CONNOR, J., HUCHZERMAYER, F., WANG, X., WANG, Y., WANG, M. & ZHOU, Z. (2013). Preservation of ovarian follicles reveals early evolution of avian reproductive behaviour. *Nature* **495**, 507-511.
- ZHENG X., O'CONNOR, J. K., WANG, X., PAN, Y., WANG, Y., WANG, M. & ZHOU, Z. (2017). Exceptional preservation of soft tissue in a new specimen of *Eoconfuciusornis* and its biological implications. *National Science Review* **4**, 441-452.
- ZHENG X., WANG, X., SULLIVAN, C., ZHANG, X., ZHANG, F., WANG, Y., LI, F. & XU, X. (2018b). Exceptional dinosaur fossils reveal early origin of avian-style digestion. *Scientific Reports* **8**, 1-8.
- ZHENG X., ZHANG, Z. & HOU, L. (2007). A new enantiornithine bird with four long rectrices from the Early Cretaceous of northern Hebei, China. *Acta Geologica Sinica (English Edition)* **81**, 703-708.
- ZHOU S., ZHOU, Z. & O'CONNOR, J. (2014a). A new piscivorous ornithuromorph from the Jehol Biota. *Historical Biology* **26**, 608-618.
- ZHOU S., ZHOU, Z. & O'CONNOR, J. K. (2012). A new basal beaked ornithurine bird from the Lower Cretaceous of western Liaoning, China. *Vertebrata Palasiatica* **50**, 9-24.
- ZHOU S., ZHOU, Z. & O'CONNOR, J. K. (2013). Anatomy of the basal ornithuromorph bird *Archaeorhynchus spathula* from the Early Cretaceous of Liaoning, China. *Journal of Vertebrate Paleontology* **33**, 141-152.
- ZHOU S., ZHOU, Z. H. & O'CONNOR, J. (2014b). A new piscivorous ornithuromorph from the Jehol Biota. *Historical Biology* **26**, 608-618.
- ZHOU Z. (1999). *Early Evolution of Birds and Avian Flight: Evidence from Mesozoic Fossils and Modern Birds*, University of Kansas.

- ZHOU Z., BARRETT, P. M. & HILTON, J. (2003a). An exceptionally preserved Lower Cretaceous ecosystem. *Nature* **421**, 807.
- ZHOU Z. & HOU, L. (2002a). The Discovery and Study of Mesozoic Birds in China. In *Mesozoic Birds: Above the Heads of Dinosaurs* (ed. L. M. Chiappe and L. M. Witmer), pp. 160-183. University of California Press, Berkeley and Los Angeles, CA, USA.
- ZHOU Z. & LI, F. Z. Z. (2010). A new Lower Cretaceous bird from China and tooth reduction in early avian evolution. *Proceedings of the Royal Society B: Biological Sciences* **277**, 219-227.
- ZHOU Z., WINKLER, D. E., FORTUNY, J., KAISER, T. M. & MARCÉ-NOGUÉ, J. (2019). Why ruminating ungulates chew sloppily: Biomechanics discern a phylogenetic pattern. *PLOS ONE* **14**, e0214510.
- ZHOU Z. & ZHANG, F. (2002b). A long-tailed, seed-eating bird from the Early Cretaceous of China. *Nature* **418**, 405-409.
- ZHOU Z. & ZHANG, F. (2003b). Anatomy of the primitive bird *Sapeornis chaoyangensis* from the Early Cretaceous of Liaoning, China. *Canadian Journal of Earth Sciences* **40**, 731-747.
- ZHOU Z. & ZHANG, F. (2005). Discovery of an ornithurine bird and its implication for Early Cretaceous avian radiation. *Proceedings of the National Academy of Sciences* **102**, 18998-19002.
- ZHOU Z. & ZHANG, F. (2006). A beaked basal ornithurine bird (Aves, Ornithurae) from the Lower Cretaceous of China. *Zoologica Scripta* **35**, 363-373.
- ZHOU Z., ZHANG, F. & LI, Z. (2009). A new basal ornithurine bird (*Jianchangornis microdonta* gen. et sp. nov.) from the Lower Cretaceous of China. *Vertebrata Palasiatica* **47**, 299-310.
- ZHOU Z. H., CHIAPPE, L. M. & ZHANG, F. C. (2005). Anatomy of the early cretaceous bird *Eoenantiornis buhleri* (Aves : Enantiornithes) from China. *Canadian Journal of Earth Sciences* **42**, 1331-1338.
- ZHOU Z. H., CLARKE, J. & ZHANG, F. C. (2008). Insight into diversity, body size and morphological evolution from the largest Early Cretaceous enantiornithine bird. *Journal of Anatomy* **212**, 565-577.
- ZHOU Z. H., CLARKE, J., ZHANG, F. C. & WINGS, O. (2004). Gastroliths in *Yanornis*: an indication of the earliest radical diet-switching and gizzard plasticity in the lineage leading to living birds? *Naturwissenschaften* **91**, 571-574.
- ZINOVIEV A. V. (2009). An attempt to reconstruct the lifestyle of confuciusornithids (Aves, Confuciusornithiformes). *Paleontological Journal* **43**, 444-452.
- ZUBER M., LAAB, M., HAMANN, E., KRETSCHMER, S., HAUSCHKE, N., VAN DE KAMP, T., BAUMBACH, T. & KOENIG, T. (2017). Augmented laminography, a correlative 3D imaging method for revealing the inner structure of compressed fossils. *Scientific Reports* **7**, 41413.
- ZUM GAHR K.-H. (1998). Wear by hard particles. *Tribology International* **31**, 587-596.
- ZUSI R. L. (1984). A functional and evolutionary analysis of rhyngokinesis in birds. *Smithsonian Contributions to Zoology* **395**.

## Figures




Fig. 1.

Methods employed in measuring the angle of bird claw curvature. Methods are diagrammed onto the digit I ungual of the enantiornithine *Mirarce eatoni*, as depicted in (Atterholt *et al.*, 2018). (A–B) Inner arc angle (A) as codified by (Feduccia, 1993) and (B) (Fowler *et al.*, 2009). (C) Outer arc angle as codified by (Pike *et al.*, 2004). In all cases, an initial chord (red) is drawn between the tip of the claw and the (A) proximal walking surface of the ungual or the (B) ventroproximal or (C) dorsoproximal extent of the horny sheathe of the claw [visible as inflection points in bony cores]. A perpendicular line (yellow) is drawn from the midpoint of this chord to the ventral (A and B) or dorsal (C) arc of the ungual. Lines connecting the endpoints (green) are drawn, as well as lines perpendicular to the midpoint of these green lines (blue). The intersection of the blue lines defines the centre of a circle approximated by the arc. From this centre, lines (magenta) are drawn to the endpoints of the initial chord, and the angle between the magenta lines is the angle of curvature of the claw arc.






Fig. 2.

Length, width, and height measurements of the avian pes. Length (red), width (green) and height (blue), diagrammed onto the digit I phalanx I and digit I ungual of *Mirarce eatoni* as depicted in (Atterholt *et al.*, 2018). (A–C) Measurements of the non-ungual phalanges in (A) lateral, (B) dorsal, and (C) distal views. (D–E) Measures of the ungual phalanx in (D) lateral and (E) proximal views. Note that most measures refer to the greatest distance measurable in a given dimension, but some such as  $w_{ft}$  are shown as being less than the greatest measurable distance without explanation in the works in which they are used.

Abbreviations:  $l_p$ , proximodistal length of phalanx;  $l_c$ , chord length of ungual;  $l_a$ , arc length of ungual;  $l_{ft}$ , proximodistal length of flexor tubercle;  $w_p$ , mediolateral width of phalanx;  $w_{ft}$ , mediolateral width of flexor tubercle;  $h_p$ , dorsoventral height of phalanx;  $h_b$ , height at the base of the ungual;  $h_m$ , height at the mid-arc of the ungual;  $h_{ft}$ , dorsoventral height of the flexor tubercle.




Fig. 3.

Third-class lever nature of animal jaws. Here a reconstruction of *Shenqiornis* after (O'Connor *et al.*, 2011b) is used as an example. The load (**green**) is represented as a square of theoretical foodstuff between the teeth. The effort (**blue**) is represented as a simplification of the *m. adductor mandibulae* complex (see Figure 4A for full reconstruction of attachments). The fulcrum (**red**) is represented as the circled articulation between the upper and lower jaws.




Fig. 4.

Illustration of the functional measurements used by Ma *et al.* (2020) as well as characters 22 and 23 from Button and Zanno (2020). Example is a reconstruction of *Shenqiornis* after (O'Connor *et al.*, 2011b). C2–7 of Button and Zanno (2020) are combined into anterior and posterior jaw-closing MA as defined by Ma *et al.* (2020). Articular offset is identical between the two. (A) Reconstruction of skull muscle attachments after (Holliday, 2009). Reconstructions of *m. adductor mandibulae externus* (purple) and *m. adductor mandibulae posterior* (green) are more certain while reconstruction of *m. pterygoideus* (orange) is tentative due to the uncertain nature of the pterygoid in enantiornithines (Chiappe *et al.*, 2001; O'Connor *et al.*, 2011b). The yellow star is the centroid of the irregular shape bounding all attachments, treated as the center of force for adduction. Uncertainty of attachment area size precludes more precise weighting of

the attachment centroid. The abductor muscle for the jaw, the *m. depressor mandibulae* (cyan), attaches perpendicular to the viewing plane (Lautenschlager *et al.*, 2014a) and so is represented as a line along the back of the skull. (B–D) Diagrams illustrating inlevers (red) and outlevers (blue) for calculating (B) anterior jaw-closing mechanical advantage, (C) posterior jaw-closing mechanical advantage, and (D) jaw-opening mechanical advantage. (E–G) Illustrations of measurements to calculate. (E) relative articular offset, (F) relative maximum rostral height, and (G) relative average rostral height. See (Ma *et al.*, 2020) for an explanation of calculations. (H–I) Close-up of the premaxilla indicating measurements of (H) C22 tooth angle and (I) C23 tooth slenderness index. See (Data S4 in Button *et al.*, 2020) for explanation of calculations.




Fig. 5.

Comparison of finite element analysis (FEA) failure criteria. Comparisons are made using a reconstruction of the upper jaw of *Shenqiornis* after (O'Connor *et al.*, 2011b). All models use isotropic material properties of ostrich mandible (Rayfield, 2011a), make plane strain assumptions, constrain the articular condyle in all directions, constrain the first premaxillary tooth in dorsoventral translation, and load the skull with an equivalent amount of force. Force was applied using the macro in (Morales-García *et al.*, 2019) which replicates muscle fibres, with attachments based on those pictured in Figure 4. Legends are scaled to make the models look as similar as possible. (A) Von Mises [VM] stress. (B)

principal strain. Note that while both map very similarly onto the model, the region of high distortion at the occipital condyle is smaller in B than A. In human studies, this smaller region represents a narrower margin of error for the location of failure (Schileo *et al.*, 2008; Yosibash *et al.*, 2010). This implies greater validity for principal strain as a strength criterion.




Fig. 6.

Comparison of construction of finite element models with flexible connective tissue using a reconstruction. Comparisons are made using the upper jaw of *Shenqiornis* after (O'Connor *et al.*, 2011b). Base models were constructed as in Figure 5. Breaks were created in said model and filled with beam elements to replicate connective tissue, after the techniques of (Cost *et al.*, 2019). A bite force of 9.3N

was chosen to be similar to those recorded by (Corbin *et al.*, 2015). (A) Beams using the cross-sectional area and Young's modulus of rat cranial sutures, as detailed in (Chien *et al.*, 2008). These properties were used by (Cost *et al.*, 2019) to model the flexible components of the skull in *Gekko* and *Psittacus*, animals of similar size to *Shenqiornis*. (B) Beams using the cross-sectional area and Young's modulus of canine patellar tendon, as detailed in (Haut *et al.*, 1992). These properties were used by (Cost *et al.*, 2019) to model flexible components in the skull of *Tyrannosaurus*. Note that in A the jugal is dislocated ventrally to a biologically unreasonable degree and in B the quadrate is dislocated cranially to a lesser but still unreasonable degree. (C) Beams assigned properties extrapolated from those of a variety of connective tissues reported in the literature and normalized to body mass (Table S1). Note both the lower degree of dislocation and the lower peak principal strains experienced. However, it is unclear how valid this model is. Only three data points are available for calculating cross-section trends. For Young's Modulus, even after excluding outliers,  $R^2$  values for a trend line could not be increased above 0.25. Sensitivity analyses show cross-section of beams has a stronger control on excursion; rat suture models can only achieve similar excursion to the scaled property model if tendon cross-sections are 45 times greater or if their Young's modulus increases 60-fold. In all three models the overall strain experienced is reduced relative to those in Figure 5B, but artificially inflated at the locations of beam attachment [clearest at the jugal/cranium contact] due to singularities.




Fig. 7.

Sensitivity analysis of rhamphotheca thickness in *Buteo jamaicensis*. Comparisons are made using a radiograph of the lower jaw of *Buteo jamaicensis* after (Smith *et al.*, 1990). All models use isotropic material properties of ostrich mandible and rhamphotheca (Rayfield, 2011a), make plane strain assumptions, constrain the articular glenoid in all directions, constrain the rostral tip of the rhamphotheca in dorsoventral translation, and load the mandible with an equivalent amount of force. Force was applied using the macro in (Morales-García *et al.*, 2019) which replicates muscle fibres, with attachments based on (Lautenschlager *et al.*, 2014a). A bite force of 9.0 N was calculated with the regressions of (Sustaita & Hertel, 2010) assuming a body mass of 1kg. Rhamphotheca is highlighted with a red outline. Length of the jaw overall is kept constant with the bone underneath modelled (A) realistically [i.e. as in the radiograph], (B) with greatly thinned rhamphotheca, and (C) with greatly thickened rhamphotheca. Note that the strain magnitude and distribution in each is nearly identical. Thus, precise thickness of the rhamphotheca appears unimportant in constructing 2D FE models of avian lower jaws.






Fig. 8.

Summary chart of our recommended framework for the study of non-avian avian diet. Approaches are followed by a brief description of specific prescribed techniques, the body part it would be performed on, relevant results, and recommended dietary and/or modelling interpretations of said results.




Fig. 9

Summary workflow for using the framework described in this paper. **Identify Fossil Specimen(s):** Section VI The Framework suggests several taxa of particular interest, and Application/Discussion sections throughout list published specimens with particular promise. **Determine Tests:** See the second column of Figure 8 to find out which preserved body parts are necessary for which tests. Note that while stable isotope analysis can be performed on bones in addition to teeth, this is not recommended as the signatures they give are less reliable. **Test:** Perform a given test [one of the seven listed] on the specimen of interest. Application sections for each technique provide references for methodology. Dashed lines pointing to the test types indicate that not all tests may be possible. The solid lines pointing away from them indicate that any tests which can be performed should have their interpretations contribute to the final synthesis. **Interpret Test Results:** The final column of Figure 8 provides idealised interpretations for test results into a set of potential diets. Section V.4 Complications may also be pertinent to this workflow stage. **Synthesise into Diet Assignment:** Combine test results with those of other tests performed. In the simplest terms, this means finding the common elements between each tests' interpretations, though the possibility of feeding styles not seen in any extant group may require more creativity at this stage.






Fig. 10.

Summary of our recommended study framework, as applied to the extant golden eagle [*Aquila chrysaetos*]. Morphometrics and Lever Modelling are transposed from Figure 8 for clarity. Lever modelling shows adaptation for vulnerable and possibly mobile prey, pointing to either predation or molliphagous herbivory. Morphometrics and FEA of the pes provide evidence of raptorial use, and thus refine the lever modelling findings to predation. Given that the animal is carnivorous, then, body mass makes vertebrate consumption much more likely than invertebrate consumption. Thus, these four lines of

evidence point to *Aquila chrysaetos* being a raptorial vertebrate predator, which it indeed is (Watson, 2010).

## Tables

Table 1.

Glossary of dietary categories mentioned in this review. Note that these are general classifications that may or may not be mutually exclusive, and may be operationalized differently by different studies. A reference going into more detail about each classification [“Source”] is provided as well. The prefix hyper- is occasionally applied to diet categories, indicating a particularly high percentage of the animal’s diet consists of the relevant food source.

Term	Definition	Source
Carnivorous	Energy acquired primarily by consuming animal tissue.	(Ullrey, 2018)
Durophagous	Consuming hard parts of organisms, or otherwise breaking their hard parts before consumption.	(Crofts & Summers, 2014)
Frugivorous	Consuming the nutritive tissue ['flesh'] of fruits.	(Jordano, 2000)
Granivorous	Consuming plant seeds, before or after dispersal.	(Hulme & Benkman, 2002)
Herbivorous	Energy acquired primarily by consuming plant tissue.	(Karban & Agrawal, 2002)
Invertivorous	Consuming invertebrate animal tissue.	(Thomas, 2014)
Molliphagous	Consuming food that is soft, i.e. requiring relatively little energy to fracture.	This study
Nectarivorous	Consuming nectar, a sugary liquid exuded by flowers.	(Nicolson & Fleming, 2014)
Omnivorous	Consuming a variety of foods, with no one source providing the majority of energy.	(Thompson, Hemberg, Starzomski <i>et al.</i> , 2007)
Osteophagous	Consuming bone or bone marrow.	(Wroe <i>et al.</i> , 2005)
Piscivorous	Consuming 'fish' [non-tetrapod gnathostome] tissue.	(Eklöv & Diehl, 1994)
Predatory	Consuming tissue of animals killed by the consumer.	(Taylor, 2013)
Raptorial	Predation in which the pes plays a major role in killing and/or restraining the prey.	(Fowler <i>et al.</i> , 2009)
Scavenging	Consuming tissue of animals not killed by the consumer.	(Turner, Abernethy, Conner <i>et al.</i> , 2017)
Vertivorous	Consuming vertebrate animal tissue.	(Garrard, McCarthy, Vesk <i>et al.</i> , 2012)

Table 2.

Mass estimates of avialan individuals. Input measurements are taken from scaled images in the literature. Estimates are made using the equations of Serrano *et al.* (2015). The ORNnl equation is less precise and used in cases where a key component in the ENAN or ORPH equations is not preserved. The source text has a typographical error in four of the equations (Serrano, 2020). The corrected equations are:

$$\text{JEHO: } -1.933 + 1.486 \text{ HL} + 0.416 \text{ bcL} + 0.965 \text{ dHW} - 0.36 \text{ deHW} - 1.536 \text{ UL} + 0.635 \text{ peUW} - 0.397 \text{ FL} + 0.834 \text{ dFWml} + 0.302 \text{ TL}$$

$$\text{SAPE: } -2.876 + 0.952 \text{ HL} + 0.352 \text{ bcL} + 0.424 \text{ dHW} + 0.967 \text{ UL} + 0.615 \text{ deUW} + 0.340 \text{ dUW} - 1.891 \text{ RL} - 0.343 \text{ DCmW} + 0.589 \text{ dFWcc} + 0.878 \text{ TL} - 0.446 \text{ TmL}$$

$$\text{ENAN: } -2.626 + 1.528 \text{ HL} + 0.34 \text{ bcL} + 0.828 \text{ dHW} - 1.451 \text{ UL} + 0.811 \text{ dUW} + 0.378 \text{ TL}$$

$$\text{ORNnl: } -2.392 + 1.799 \text{ HL} + 0.355 \text{ bcL} + 1.014 \text{ dHW} + 1.003 \text{ dUW} - 1.475 \text{ RL} - 0.363 \text{ DCmW}$$

See (Serrano *et al.*, 2015) for explanation of abbreviations. Body mass correction factors were not included in the original paper and so were back calculated from the reported values; all were very close to 1.

Taxon	Specimen	Mean Mass Estimate (kg)	Lower Mass Estimate (kg)	Upper Mass Estimate (kg)	Equation Used
<b>Jeholornithiformes</b>					
Jeholornithiformes indet.	DLNM D2139	1.418	1.173	1.664	JEHO
<i>Jeholornis curvipes</i>	YFGP-yb2	1.504	1.244	1.764	JEHO
<i>Jeholornis prima</i>	STM 2-15	1.442	1.193	1.692	JEHO
<i>Kompsornis longicaudus</i>	AGB-6997	0.952	0.787	1.117	JEHO
<i>Shenzhouraptor sinensis</i>	LPM 00193	0.883	0.730	1.036	JEHO
<b>Confuciusornithidae</b>					
<i>Confuciusornis sanctus</i>	IVPP V13313	0.598	0.497	0.700	CONF
<i>Yangavis confucii</i>	IVPP V18929	0.564	0.468	0.659	CONF
<b>Sapeornithiformes</b>					

<i>Omnivoropteryx sinousaorum</i>	CAGS 02-IG-gausa-3	1.429	1.193	1.665	SAPE
<b>Enantiornithes</b>					
<i>Alethoalaornis agitornis</i>	LPM 00038	0.158	0.127	0.189	ENAN
<i>Bohaiornis guoi</i>	IVPP V17963	0.300	0.242	0.358	ENAN
<i>B. guoi</i>	LPM B00167	0.032	0.026	0.038	ENAN
<i>Cathayornis yandica</i>	IVPP V9769a/b	0.062	0.050	0.074	ENAN
<i>Chiappeavis magnapremaxillo</i>	STM 29-11	0.465	0.375	0.556	ENAN
<i>Dalingheornis liweii</i>	CNU VB2005001	0.008	0.007	0.010	ENAN
<i>Dapingfangornis sentisorhinus</i>	LPM 00039	0.204	0.164	0.243	ENAN
<i>Dunhuangia cuii</i>	GSGM-05-CM-030	0.124	0.099	0.149	ORNnl
<i>Elsornis keni</i>	MPD-b 100/201	1.512	1.206	1.817	ORNnl
<i>Eopengornis martini</i>	STM 24-1	0.193	0.155	0.230	ENAN
<i>Fortunguavis xiaotaizicus</i>	IVPP V18631	0.296	0.236	0.356	ORNnl
<i>Grabauornis lingyuanensis</i>	IVPP V14595	0.127	0.102	0.151	ENAN
<i>Gracilornis jiufotangensis</i>	PMOL-AB00170	0.027	0.021	0.032	ENAN
<i>Gretcheniao sinensis</i>	BMNHC Ph-829	0.455	0.367	0.543	ENAN
<i>Houornis caudatus</i>	IVPP V10917/1, IVPP V10917/2	0.107	0.086	0.129	ORNnl
<i>Huoshanornis huji</i>	DNM D2126	0.071	0.057	0.085	ENAN
<i>Jibeinia luanhera</i>	Drawing in (Hou, 1997), holotype lost	0.065	0.053	0.078	ENAN
<i>Junornis houi</i>	BMNHC-PH 919a/b	0.074	0.059	0.088	ENAN

<i>Liaoningornis longidigitris</i>	IVPP V11303	0.180	0.145	0.215	ENAN
<i>Linyiornis amoena</i>	STM 11-80	0.215	0.173	0.256	ENAN
<i>Longipteryx chaoyangensis</i>	DNHM D2889	0.751	0.605	0.897	ENAN
<i>Longusunguis kurochkini</i>	IVPP V17964	0.171	0.137	0.204	ENAN
<i>L. kurochkini</i>	IVPP V18693	0.237	0.191	0.283	ENAN
<i>Microenantiornis vulgaris</i>	PMOL AB00171	0.067	0.054	0.080	ENAN
<i>Monoenantiornis sihedangia</i>	IVPP V20289	0.355	0.286	0.424	ENAN
<i>Noguerornis gonzalezi</i>	LP.1702.P	0.020	0.016	0.024	ORNnl
<i>Orienantius ritteri</i>	BMNHC Ph-1154a/b	0.071	0.057	0.085	ENAN
<i>O. ritteri</i>	BMNHC Ph-1156a/b	0.083	0.067	0.100	ENAN
<i>Parabohaiornis martini</i>	IVPP V18691	0.221	0.178	0.263	ENAN
<i>Parapengornis eurycaudatus</i>	IVPP V18687	0.429	0.345	0.512	ENAN
<i>Paraprotopteryx gracilis</i>	STM V001	0.046	0.037	0.055	ENAN
<i>Parvavis chuxiongensis</i>	IVPP V18586/1, IVPP V18586/2	0.024	0.020	0.029	ENAN
<i>Piscivorenantiornis inusitatus</i>	IVPP V22582	0.281	0.227	0.336	ENAN
<i>Protopteryx fengningensis</i>	BMNHC Ph-1060a/b	0.109	0.088	0.130	ENAN
<i>P. fengningensis</i>	BMNHC Ph-1158a/b	0.088	0.071	0.105	ENAN

<i>Pterygornis dapingfangensis</i>	IVPP V20729	0.080	0.064	0.095	ENAN
<i>Shangyang graciles</i>	IVPP V25033	0.108	0.087	0.129	ENAN
<i>Shanweiniao cooperorum</i>	DNHM D1878/1, DNHM D1878/2	0.062	0.050	0.074	ENAN
<i>Shengjingornis yangi</i>	PMOL AB00179	0.340	0.274	0.406	ENAN
<i>Shenqiornis mengi</i>	DNHM D2950/1	0.340	0.274	0.406	ENAN
<i>Sulcavis geeorum</i>	BMNH Ph-000805	0.333	0.268	0.397	ENAN
<i>Yuanjiawaornis viriosus</i>	PMOL AB00032	0.418	0.337	0.499	ENAN
<i>Zhouornis hani</i>	BMNHCPH 756	0.253	0.204	0.303	ENAN
<i>Z. hani</i>	CNUVB-0903	0.758	0.611	0.905	ENAN
<b>Non-avian Ornithuromorpha</b>					
<i>Archaeorhynchus spathula</i>	IVPP V17075	0.282	0.227	0.336	ORPH
<i>A. spathula</i>	IVPP V17091	0.153	0.123	0.183	ORPH
<i>Archaeornithura meemannae</i>	STM 7-145	0.136	0.109	0.162	ORPH
<i>Bellulia rectusunguis</i>	IVPP V17970	0.778	0.627	0.928	ORPH
<i>Changzuiornis ahgm</i>	AGB5840	0.240	0.193	0.286	ORPH
<i>Dingavis longimaxilla</i>	IVPP V20284	0.526	0.424	0.629	ORPH
<i>Eogranivora edentulata</i>	STM 35-3	0.291	0.235	0.348	ORPH
<i>Gansus yumenensis</i>	GSGM-05-CM-014	0.142	0.114	0.169	ORPH
<i>Hongshanornis longicresta</i>	DNHM D2945	0.075	0.061	0.090	ORPH
<i>Patagopteryx deferrariisi</i>	MACN-N-11	1.130	0.911	1.349	ORPH
<i>Piscivoravis lii</i>	IVPP V17078	0.849	0.684	1.013	ORPH
<i>Schizoura lii</i>	IVPP V16861	0.377	0.304	0.450	ORPH

<i>Tianyuornis cheni</i>	STM 7-53	0.112	0.090	0.133	ORPH
<i>Xinghaiornis lini</i>	XHPM 1121	0.539	0.434	0.643	ORPH
<i>Yanornis martini</i> (juvenile?)	IVPP V13358	0.117	0.094	0.140	ORPH
<i>Yanornis</i> sp.	STM 9-15	0.577	0.465	0.689	ORPH
<i>Yanornis</i> sp.	STM 9-46	0.984	0.793	1.175	ORPH
<i>Yumenornis huangi</i>	GSGM-06-CM-013	0.321	0.256	0.386	ORNnl



Table 3.

**Listing of published non-avian avialan skulls.** Developed from Table 1 of O'Connor and Chiappe (2011b). Here material is described as partial when >50% complete and nearly complete when >75% complete. For preservation, IVPP V1165 (*Protopteryx*) is the standard for poor, DNHM-D2522 (*Rapaxavis*) for good, and IVPP V15336 (*Pengornis*) for excellent. If multiple publications use the same picture the Pictured In column refers to that which has the highest-resolution image. An NA for picturing publication means that a published image of the skull could not be found.

Taxon	Specimen Number	Country	Material	View	Preservation	Crushed	Disarticulated	Pictured In
<b>Scansoriopterygidae</b> [probably non-avian pennaraptorans]								
<i>Ambopteryx longibrachium</i>	IVPP V24192	China	indistinct	lateral?	poor	X		(Wang, O'Connor, Xu <i>et al.</i> , 2019c)
<i>Epidexipteryx hui</i>	IVPP V15471	China	nearly complete skull	lateral	poor; some voids	X		(Zhang, Zhou, Xu <i>et al.</i> , 2008)
<i>Scansoriopteryx heilmanni</i>	CAGS02-IG-gausa-1/DM 607	China	partial skull	dorsolateral	good	X	X	(Czerkas & Feduccia, 2014)
<i>Scansoriopteryx heilmanni</i>	IVPP V12653	China	frontal, parietal, sclerotic ring, mandible	dorsal	poor; only voids; juvenile		X	(Zhang, Zhou, Xu <i>et al.</i> , 2002)
<i>Yi qi</i>	STM 31-2	China	nearly complete skull	lateral	poor; some voids			(Xu <i>et al.</i> , 2015)
<b>Anchiornithinae</b> [possibly troodontids]								
<i>Anchiornis huxleyi</i>	HGM-41HIII 0404	China	partial skull	lateral	good; some voids	X		(Guo, Xu & Jia, 2018)
<i>A. huxleyi</i>	HGM-41HIII 0415	China	nearly complete skull	ventrolateral	good	X		(Guo <i>et al.</i> , 2018)
<i>A. huxleyi</i>	BMNHC Ph-804	China	partial skull	dorsolateral	poor	X		(Pei, Li, Meng <i>et al.</i> , 2017)
<i>A. huxleyi</i>	BMNHC Ph-822	China	partial skull	dorsal	poor	X		(Pei <i>et al.</i> , 2017)
<i>A. huxleyi</i>	BMNHC Ph-823	China	partial skull	dorsolateral	poor	X		(Pei <i>et al.</i> , 2017)
<i>A. huxleyi</i>	PKUP V1068	China	nearly complete skull	lateral	good	X		(Pei <i>et al.</i> , 2017)

<i>Anchiornis</i> sp.	STM 0-179	China	partial skull	lateral	poor; some voids	X		(Zheng <i>et al.</i> , 2018b)
<i>Anchiornis</i> sp.	STM 0-224	China	nearly complete skull	lateral	poor	X		(Zheng <i>et al.</i> , 2018b)
<i>Anchiornis</i> sp.	STM A0-4	China	nearly complete skull	lateral	good	X		(Zheng <i>et al.</i> , 2018b)
<i>Aurornis xui</i>	YFGP-T5198	China	partial skull	lateral	poor	X		(Godefroit, Cau, Dong-Yu <i>et al.</i> , 2013)
<i>Eosipnopteryx brevipenna</i>	YFGP-T5197	China	partial skull	dorsolateral	good	X		(Godefroit, Demuynck, Dyke <i>et al.</i> , 2013)
<i>Serikornis sungei</i>	PMOL-AB00200	China	partial skull	lateral	poor; some voids	X	X	(Lefèvre, Cau, Cincotta <i>et al.</i> , 2017)
<i>Xiaotingia zhengi</i>	STM 27-2	China	partial skull	dorsolateral	good; several voids	X		(Xu <i>et al.</i> , 2011)
<b>Archaeopterygidae</b>								
London <i>Archaeopteryx</i>	BMNH PV OR 37001	Germany	braincase	3D	excellent		X	(Alonso <i>et al.</i> , 2004; Whetstone, 1983)
Berlin <i>Archaeopteryx</i>	MB.Av.101	Germany	nearly complete skull	lateral	excellent			(Christiansen & Bonde, 2004; Wellnhofer, 2010)
Eichstätt <i>Archaeopteryx</i>	JM 2257	Germany	nearly complete skull	dorsolateral	good	X		(Ostrom, 1976; Whetstone, 1983)
Solnhofen <i>Archaeopteryx</i>	BMMS 500	Germany	partial skull	lateral	poor; mostly voids	X	X	(Elzanowski, 2001a)
Munich <i>Archaeopteryx</i>	BSP 1999 I 50	Germany	nearly complete skull	dorsal	good	X	X	(Rauhut, 2014)

Daiting <i>Archaeopteryx</i>	SNSB BSPG VN-2010/1	Germany	partial skull	dorsolateral	poor	X	X	(Kundrát, Nudds, Kear <i>et al.</i> , 2019)
Thermopolis <i>Archaeopteryx</i>	WDC-CSG-100	Germany	braincase	lateral	good	X		(Rauhut, 2014)
11th <i>Archaeopteryx</i>	[private]	Germany	premaxilla, dentary, surangular	dorsal	excellent	X	X	(Foth, Tischlinger & Rauhut, 2014)
12th <i>Archaeopteryx</i>	DNWK 02924 [privately held]	Germany	nearly complete skull	dorsolateral	good	X	X	(Rauhut <i>et al.</i> , 2018)
<b>Jeholornithiformes</b>								
Jeholornithiformes indet.	DLNM D2139	China	partial skull	dorsolateral	poor	X		(Wang <i>et al.</i> , 2020b)
<i>Jeholornis curvipes</i>	YFGP-yb2	China	partial skull	dorsolateral	good	X		(Lefèvre, Hu, Escuillié <i>et al.</i> , 2014)
<i>Jeholornis palmapenis</i>	SDM 2009.01	China	partial skull	dorsolateral	poor	X		(O'Connor, 2019)
<i>Jeholornis palmapenis</i>	SDM 20090109.1/2	China	partial skull	lateral	poor	X		(O'Connor, Sun, Xu <i>et al.</i> , 2012)
<i>Jeholornis prima</i>	BMNHC Ph-780	China	complete skull	lateral	excellent			(Chiappe <i>et al.</i> , 2016 pg. 37)
<i>J. prima</i>	CDL-02-04-001	China	nearly complete skull	lateral	good	X		(Chiappe <i>et al.</i> , 2016 pg. 34)
<i>J. prima</i>	IVPP V13274	China	partial skull	lateral	poor; ingested seeds	X		Zhou and Zhang 2002
<i>J. prima</i>	IVPP V13350	China	partial skull	dorsolateral	poor	X		(O'Connor, 2019)
<i>J. prima</i>	STM 2-15	China	nearly complete skull	lateral	good; ingested seeds			(O'Connor <i>et al.</i> , 2018)
<i>J. prima</i>	STM 2-31	China	partial skull	lateral	poor; some voids; gastroliths	X		(O'Connor <i>et al.</i> , 2018)

<i>J. prima</i>	STM 2-41	China	partial skull	dorsolateral	poor; ingested seeds	X	X	(O'Connor <i>et al.</i> , 2018)
<i>J. prima</i>	STM 3-28	China	partial skull	lateral	poor; gastroliths	X		(O'Connor <i>et al.</i> , 2018)
<i>Jeholornis</i> sp.	IVPP V14978	China	partial skull	lateral	good	X	X	(Wang, O'Connor & Zhou, 2019b)
<i>Jeholornis</i> sp.	STM 2-51	China	partial skull	lateral	poor	X		(Zheng <i>et al.</i> , 2013)
<i>Jixiangornis orientalis</i>	CDPC-02-04-001	China	partial skull	dorsolateral	poor	X		(Ji, Ji, Zhang <i>et al.</i> , 2002)
<i>Kompsornis longicaudus</i>	AGB-6997	China	nearly complete skull	lateral	poor	X		(Wang <i>et al.</i> , 2020b)
<i>Shenzhouraptor sinensis</i>	LPM 00193	China	partial skull	dorsolateral	poor	X	X	(Ji, Ji, You <i>et al.</i> , 2003)
<b>Early pygostylians</b>								
<i>Zhongornis haoae</i> [possibly scansoriopterygid or early-diverging avialan]	DNHM D2456	China	partial skull	dorsolateral	poor; juvenile	X		(Chiappe <i>et al.</i> , 2016 pg. 38)
<b>Confuciusornithidae</b>								
Confuciusornithidae indet.	BMNHC Ph-870	China	partial skull	dorsolateral	poor	X	X	(Navalón, Meng, Marugán-Lobón <i>et al.</i> , 2018b)
<i>Changchengornis hengdaoziensis</i>	GMV 2129	China	partial skull	lateral	poor	X	X	(Wang <i>et al.</i> , 2019b)
<i>Confuciusornis dui</i>	IVPP V11553	China	nearly complete skull	lateral	poor; rhamphotheca	X		(Falk <i>et al.</i> , 2019) [cast]
<i>Confuciusornis sanctus</i>	IVPP V10918	China	partial skull	lateral	poor	X	X	(Zhou <i>et al.</i> , 2002a)
<i>C. sanctus</i>	IVPP V13313	China	nearly complete skull	lateral	good	X		(Dalsätt <i>et al.</i> , 2006)

<i>C. sanctus</i>	IVPP V14412	China	partial skull	lateral	poor; mostly voids			(Wang <i>et al.</i> , 2019b)
<i>C. sanctus</i>	BMNHC Ph-766	China	nearly complete skull	dorsolateral	good	X		(Chiappe <i>et al.</i> , 2016 pg. 44)
<i>C. sanctus</i>	BMNHC Ph-986	China	complete skull	lateral	excellent; rhamphotheca			(Chiappe <i>et al.</i> , 2016 pg. 156)
<i>C. sanctus</i>	BMNHC Ph-987	China	nearly complete skull	lateral	good	X		(Chiappe <i>et al.</i> , 2016 pg. 48) p 48
<i>C. sanctus</i>	BSP 1999 I 15	China	nearly complete skull	ventrolateral	good			(Nudds & Dyke, 2010)
<i>C. sanctus</i>	DNHM D1874	China	nearly complete skull	lateral	good	X		(Chinsamy-Turan, Chiappe, Marugán-Lobón <i>et al.</i> , 2013)
<i>C. sanctus</i>	DNHM D2151	China	partial skull	ventral	good	X		(Chiappe <i>et al.</i> , 2016 pg. 121) p 121
<i>C. sanctus</i>	DNHM D2454	China	nearly complete skull	ventral	good			(Wang <i>et al.</i> , 2019b)
<i>C. sanctus</i>	DNHM D2859	China	partial skull	dorsal	good	X		(Chiappe <i>et al.</i> , 2016pg. 42)
<i>C. sanctus</i>	GMV 2130	China	nearly complete skull	lateral	poor	X		(Chiappe <i>et al.</i> , 1999)
<i>C. sanctus</i>	GMV 2131	China	partial skull	lateral	poor	X		(Chiappe <i>et al.</i> , 1999)
<i>C. sanctus</i>	GMV 2132	China	nearly complete skull	lateral	poor	X		(Chiappe <i>et al.</i> , 1999)
<i>C. sanctus</i>	GMV 2133	China	nearly complete skull	dorsal	poor	X		(Chiappe <i>et al.</i> , 1999)
<i>C. sanctus</i>	HGM-41HIII0400	China	partial skull	dorsal	poor	X		(Chiappe <i>et al.</i> , 2016 pg. 189)

<i>C. sanctus</i>	IVPP V12352	China	complete skull	dorsolateral	excellent; rhamphotheca	X		(Falk <i>et al.</i> , 2019)
<i>C. sanctus</i>	IVPP V13156	China	partial skull	ventral	poor; mostly voids	X		(O'Connor, 2019)
<i>C. sanctus</i>	IVPP V13168	China	nearly complete skull	dorsolateral	good	X		(Wang <i>et al.</i> , 2019b)
<i>C. sanctus</i>	IVPP V13171	China	nearly complete skull	dorsolateral	excellent	X	X	(Wang <i>et al.</i> , 2019b)
<i>C. sanctus</i>	IVPP V13313	China	nearly complete skull	lateral	good; fish pellet?			Dalsätt et al 2006
<i>C. sanctus</i>	JME 2005/1	China	partial skull	dorsolateral	good	X		(Elzanowski <i>et al.</i> , 2018)
<i>C. sanctus</i>	LPM 0233	China	partial skull	dorsolateral	poor	X		(Chiappe <i>et al.</i> , 2016 pg. 120)
<i>C. sanctus</i>	LPM 0228	China	nearly complete skull	dorsolateral	good			(Chiappe <i>et al.</i> , 2016 pg. 28)
<i>C. sanctus</i>	LPM 0229	China	nearly complete skull	ventrolateral	good	X		(Chiappe <i>et al.</i> , 2016 pg. 30)
<i>C. sanctus</i>	MB.Av.1168	China	complete skull	lateral	excellent			(Elzanowski <i>et al.</i> , 2018)
<i>C. sanctus</i>	MHNF 11186	China	partial skull	ventral	good	X		(Elzanowski <i>et al.</i> , 2018)
<i>C. sanctus</i>	NIGPAS-139379	China	nearly complete skull	ventrolateral	excellent	X		(Chiappe <i>et al.</i> , 2016 pg.125)
<i>C. sanctus</i>	NMB Ca. 258	China	partial skull	dorsolateral	good	X	X	(Elzanowski <i>et al.</i> , 2018)
<i>C. sanctus</i>	PMOL-AB00114	China	partial skull	dorsolateral	poor	X		(Wang <i>et al.</i> , 2019b)
<i>C. sanctus</i>	SMF Av 412	China	nearly complete skull	ventrolateral	good	X		(Elzanowski <i>et al.</i> , 2018)

<i>C. sanctus</i>	SMF Av 416	China	nearly complete skull	lateral	excellent	X	X	(Elzanowski <i>et al.</i> , 2018)
<i>C. sanctus</i>	SMF Av 420	China	partial skull	ventrolateral	good	X	X	(Elzanowski <i>et al.</i> , 2018)
<i>C. sanctus</i>	SMF Av 423	China	partial skull	lateral	good	X		(Elzanowski <i>et al.</i> , 2018)
<i>C. sanctus</i>	STM 13-162	China	partial skull	lateral	poor; rhamphotheca	X		(Miller <i>et al.</i> , In Press)
<i>Confuciusornis</i> sp.	CUGB P1401	China	partial skull	lateral	poor; only voids			(Li, Clarke, Gao <i>et al.</i> , 2018)
<i>Confuciusornis</i> sp.	IVPP V13156	China	nearly complete skull	dorsal	good	X		(Falk, Kaye, Zhou <i>et al.</i> , 2016)
<i>Eoconfuciusornis zhengi</i>	IVPP V11977	China	nearly complete skull	ventrolateral	excellent; rhamphotheca	X		(Falk <i>et al.</i> , 2019)
<i>Eoconfuciusornis</i> sp.	STM 7-144	China	partial skull	lateral	poor	X		(Zheng, O'Connor, Wang <i>et al.</i> , 2017)
<i>Yangavis confucii</i>	IVPP V18929	China	nearly complete skull	lateral	good	X		(Wang & Zhou, 2019d)
<b>Jinguofortisidae</b>								
<i>Jinguofortis perplexus</i>	IVPP V24194	China	partial skull	dorsolateral	good	X		(Wang, Stidham & Zhou, 2018)
<b>Sapeornithiformes</b>								
<i>Omnivoropteryx sinousaorum</i>	CAGS 02-IG-gausa-3	China	complete skull?	lateral?	unprepared		X	(Pomeroy, 2013)
<i>Sapeornis chaoyangensis</i>	BMNHC Ph-1067	China	nearly complete skull	lateral	excellent	X		(Chiappe <i>et al.</i> , 2016 pg 59)
<i>S. chaoyangensis</i>	CDL-08-02-01	China	partial skull	ventral	poor	X		(Chiappe <i>et al.</i> , 2016 pg. 60)
<i>S. chaoyangensis</i>	CDPC-02-08-001	China	nearly complete skull	ventral	poor	X		(Yuan, 2008)

<i>S. chaoyangensis</i>	DNHM D3078	China	partial skull	dorsal	poor	X		(Gao, Chiappe, Zhang <i>et al.</i> , 2012)
<i>S. chaoyangensis</i>	DNHM D2523	China	partial skull	lateral	poor	X		(Chiappe <i>et al.</i> , 2016 pg. 54)
<i>S. chaoyangensis</i>	HGM-41HIII0405	China	nearly complete skull	dorsolateral	excellent	X		(Chiappe <i>et al.</i> , 2016 pg. 56; Hu <i>et al.</i> , 2020a)
<i>S. chaoyangensis</i>	IVPP V13275	China	partial skull	dorsal	poor	X	X	(Hu <i>et al.</i> , 2020a)
<i>S. chaoyangensis</i>	IVPP V13276	China	partial skull	ventral	poor	X	X	(Zhou <i>et al.</i> , 2003b)
<i>S. chaoyangensis</i>	IVPP V13396	China	partial skull	lateral	good	X		(Chiappe <i>et al.</i> , 2016 pg. 61; Provini, Zhou & Zhang, 2009)
<i>S. chaoyangensis</i>	IVPP V19058	China	partial skull	lateral	poor	X	X	(Hu <i>et al.</i> , 2020a)
<i>S. chaoyangensis</i>	LPM B00018	China	nearly complete skull	lateral	poor	X		(Hu, Li, Hou <i>et al.</i> , 2010)
<i>S. chaoyangensis</i>	STM 15-7	China	dentary	lateral	good		X	(Wang, Hu, O'Connor <i>et al.</i> , 2017e)
<i>S. chaoyangensis</i>	STM 15-15	China	nearly complete skull	lateral	good; ingested seeds; gastroliths	X		(O'Connor, 2019)
<i>S. chaoyangensis</i>	STM 15-29	China	nearly complete skull	lateral	good; ingested seeds; coprolite	X	X	(Zheng <i>et al.</i> , 2011)
<i>S. chaoyangensis</i>	STM 16-18	China	partial skull	ventrolateral	good	X		(Wang <i>et al.</i> , 2017e)

# Enantiornithes



Enantiornithes indet.	NIGPAS-130723	China	nearly complete skull	lateral	poor; juvenile	X		(Chiappe <i>et al.</i> , 2016 pg. 115; Hou <i>et al.</i> , 1999a)
Enantiornithes indet.	PVL 4698	Argentina	right mandibular ramus	3D	excellent		X	(Chiappe & Walker, 2002)
Enantiornithes indet.	STM 29-8	China	partial skull	lateral	poor	X		(O'Connor <i>et al.</i> , 2014)
Enantiornithes indet.	??	Brazil	premaxilla	3D	excellent		X	Work in press (William Roberto Nava pers. com. 2019; described in Nava, Alvarenga, Chiappe <i>et al.</i> , 2015)
Enantiornithes indet.	??	Brazil	premaxilla	3D	excellent		X	Work in press (William Roberto Nava pers. com. 2019; described in Nava <i>et al.</i> , 2015)
Bohaiornithidae indet.	BMNHC Ph-1204	China	nearly complete skull	lateral	good	X		(Chiappe <i>et al.</i> , 2016 pg. 91)
Bohaiornithidae indet.	CUGB P1202	China	partial skull	lateral	poor	X		(Peteya <i>et al.</i> , 2017)
<i>Alethoalaornis agitornis</i>	LPM 00009	China	partial skull	ventrolateral	poor	X	X	(Li, Hu, Duan <i>et al.</i> , 2007)
<i>A. agitornis</i>	LPM 00038	China	nearly complete skull	lateral	good	X		(Li <i>et al.</i> , 2007)
<i>A. agitornis</i>	LPM 00040	China	partial skull	dorsal	poor	X	X	(Li <i>et al.</i> , 2007)
<i>Bohaiornis guoi</i>	IVPP V17963	China	nearly complete skull	lateral	good; gastroliths?	X		(Li, Zhou, Wang <i>et al.</i> , 2014)

<i>B. guoi</i>	LPM B00167	China	nearly complete skull	lateral	poor	X	X	(Hu, Li, Hou <i>et al.</i> , 2011)
<i>Boluochia zhengi</i>	IVPP V9770	China	premaxilla, maxilla, nasal, dentary	lateral?	poor; only voids			(O'Connor, Zhou & Zhang, 2010c)
<i>Cathayornis yandica</i>	IVPP V9769a/b	China	nearly complete skull	lateral	good		X	(Wang & Liu, 2016a)
<i>Chiappeavis magnapremaxillo</i>	STM 29-11	China	complete skull	lateral	good	X		(O'Connor <i>et al.</i> , 2016c)
<i>Cratoavis cearensis</i>	UFRJ-DG 031Av	Brazil	partial skull	lateral	poor; primarily voids	X		(Carvalho, Novas, Agnolín <i>et al.</i> , 2015)
<i>Cruralispennia multidonta</i>	IVPP V21711	China	premaxilla, maxilla, nasal, frontal, jugal, surangular, dentary	dorsal	poor		X	(Wang, O'Connor, Pan <i>et al.</i> , 2017b)
<i>Cuspirostrisornis houi</i>	IVPP V10897	China	premaxilla, nasal, frontal, dentary	lateral	?		X	(Hou, 1997) [drawing]
<i>Dalingheornis liweii</i>	CNU VB2005001	China	nearly complete skull	dorsal	poor	X		(Zhang, Hou, Yoshikasu <i>et al.</i> , 2006)
<i>Dapingfangornis sentisorhinus</i>	LPM 00039	China	nearly complete skull	ventrolateral	good; poorly prepared			(Li, Duan, Hu <i>et al.</i> , 2006)
<i>Eocathayornis walkeri</i>	IVPP V10916	China	partial skull	lateral	poor; primarily voids		X	(Wang <i>et al.</i> , 2016a)
<i>Eoenantiornis buhleri</i>	IVPP V11537	China	nearly complete skull	dorsal	good	X		(Zhou, Chiappe & Zhang, 2005)

<i>Eopengornis martini</i>	STM 24-1	China	nearly complete skull	ventral	good	X		(Wang <i>et al.</i> , 2014e)
<i>Fortunguavis xiaotaizicus</i>	IVPP V18631	China	nearly complete skull	ventral	poor	X		(Wang, O'Connor & Zhou, 2014b)
<i>Gobipipus reshetovi</i>	PIN 4492 3	Mongolia	partial skull	3D	?; embryo			(Kurochkin <i>et al.</i> , 2013) [drawing]
<i>Gobipteryx minuta</i>	IGM-100/1011	Mongolia	rostral half of skull	3D	excellent			(Chiappe <i>et al.</i> , 2001)
<i>G. minuta</i>	PIN 4492	Mongolia	partial skull	3D	excellent			(Chiappe <i>et al.</i> , 2001)
<i>G. minuta</i>	ZPAL MgR 1/12	Mongolia	partial skull	3D	poor			(Elzanowski, 1977)
<i>G. minuta</i>	ZPAL MgR 1/32	Mongolia	partial skull	3D	poor			(Elzanowski, 1977)
<i>G. minuta</i>	ZPAL MgR-I/33	Mongolia	partial skull	3D	poor; recrystalized; embryo			(Elzanowski, 1981)
<i>G. minuta</i>	ZPAL MgR-I/34	Mongolia	partial skull	3D	poor; embryo	X		(Elzanowski, 1981)
<i>G. minuta</i>	ZPAL MgR-I/88	Mongolia	partial skull	3D	good; embryo			(Elzanowski, 1981)
<i>Grabauornis lingyuanensis</i>	IVPP V14595	China	nearly complete skull	lateral	poor; only voids			(Dalsätt, Ericson & Zhou, 2014)
<i>Gracilornis jiufotangensis</i>	PMOL-AB00170	China	partial skull	lateral	good			(Li & Hou, 2011)
<i>Gretcheniao sinensis</i>	BMNH Ph-829	China	partial skull	lateral	good	X	X	(Chiappe <i>et al.</i> , 2019b)
<i>Holbotia ponomarenkoi</i>	PIN 3147-200	China	premaxilla, maxilla?, dentary	medial	poor; several voids		X	(Zelenkov & Averianov, 2016)
<i>Houornis caudatus</i>	IVPP V10917	China	premaxilla, frontal, parietal, dentary	dorsolateral	poor; several voids	X	X	(Wang <i>et al.</i> , 2016a)
<i>Huoshanornis huji</i>	DNHM D2126	China	partial skull	lateral	poor	X		(Wang, Zhang, Gao <i>et al.</i> , 2010a)

<i>Junornis houi</i>	BMNHC Ph-919	China	partial skull	lateral	poor	X		(Liu, Chiappe, Serrano <i>et al.</i> , 2017)
<i>Largirostrornis sexdentoris</i>	IVPP V10531	China	partial skull	ventral	?	X	X	(Hou, 1997) [drawing]
<i>Linyiornis amoena</i>	STM 11-80	China	partial skull	dorsolateral	good	X		(Wang <i>et al.</i> , 2016d)
<i>Longchengornis sanyanensis</i>	IVPP V10530	China	frontal, parietal, dentary	lateral?	poor; only voids		X	(Hou, 1997) [drawing]
<i>Longipteryx chaoyangensis</i>	BMNHC Ph-1071	China	nearly complete skull	lateral	good	X		(Chiappe <i>et al.</i> , 2016 pg. 89)
<i>L. chaoyangensis</i>	BMNHC Ph-826	China	partial skull	lateral	good	X		(Chiappe <i>et al.</i> , 2016 pg. 27)
<i>L. chaoyangensis</i>	BMNHC Ph-930B	China	nearly complete skull	lateral	good	X		(Chiappe <i>et al.</i> , 2016 pg. 176)
<i>L. chaoyangensis</i>	DNHM D2566	China	nearly complete skull	lateral	good	X		NA
<i>L. chaoyangensis</i>	DNHM D2889	China	complete skull	lateral	excellent			(Wang <i>et al.</i> , 2015c)
<i>L. chaoyangensis</i>	HGM-41HIII0319	China	nearly complete skull	lateral	good	X		(Chiappe <i>et al.</i> , 2016 pg. 88)
<i>L. chaoyangensis</i>	IVPP V11309	China	nearly complete skull	lateral	poor	X		(Chiappe <i>et al.</i> , 2016 pg. 86; Hou <i>et al.</i> , 2004)
<i>L. chaoyangensis</i>	IVPP V12325	China	nearly complete skull	lateral	good	X		(Wang <i>et al.</i> , 2015c)
<i>L. chaoyangensis</i>	IVPP V12552	China	nearly complete skull	lateral	good; juvenile	X		(O'Connor <i>et al.</i> , 2010c) [drawing]
<i>L. chaoyangensis</i>	SG2005-B1	China	nearly complete skull	lateral	good; several voids	X	X	(Li, Gong, Zhang <i>et al.</i> , 2010)

<i>Longusunguis kurochkini</i>	IVPP V17964	China	partial skull	lateral	excellent	X	X	(Wang <i>et al.</i> , 2014c)
<i>L. kurochkini</i>	IVPP V18693	China	nearly complete skull	dorsal	good	X	X	(Hu <i>et al.</i> , 2020b)
<i>Microenantiornis vulgaris</i>	PMOL-AB00171	China	nearly complete skull	lateral	poor	X		(Wei & Li, 2011)
<i>Monoenantiornis sihedangia</i>	IVPP V20289	China	partial skull	ventrolateral	good	X	X	(Hu & O'Connor, 2017)
<i>Neuquenornis volans</i>	MUCPv-142	Argentina	braincase	3D	good		X	(Chiappe <i>et al.</i> , 1994)
<i>Orienantius ritteri</i>	BMNHC Ph-1154	China	partial skull	lateral	poor	X		(Liu, Chiappe, Zhang <i>et al.</i> , 2019)
<i>O. ritteri</i>	BMNHC Ph-1156	China	partial skull	lateral	poor	X		(Liu <i>et al.</i> , 2019)
<i>Parabohaiornis martini</i>	IVPP V18690/1	China	partial skull	lateral	good	X		(Wang <i>et al.</i> , 2014c)
<i>Parapengornis eurycaudatus</i>	IVPP V18687	China	nearly complete skull	lateral	excellent	X		(Hu <i>et al.</i> , 2015)
<i>Paraprotapteryx gracilis</i>	STM V001	China	partial skull	lateral	poor	X		(Zheng, Zhang & Hou, 2007)
<i>Parvavis chuxiongensis</i>	IVPP V18586	China	exoccipital, surangular	cranial	poor; primarily voids	X	X	(Wang, Zhou & Xu, 2014d)
<i>Pengornis houi</i>	IVPP V15336	China	complete skull	lateral	excellent			(Zhou, Clarke & Zhang, 2008)
<i>Piscivorenantiornis inusitatus</i>	IVPP V22582	China	premaxilla, frontal, quadrate, braincase, dentary	various	good; fish pellet?		X	(Wang <i>et al.</i> , 2017c)
<i>Protopteryx fengningensis</i>	BMNHC Ph-1060	China	partial skull	lateral	poor	X		(Chiappe, Di, Serrano <i>et al.</i> , 2019a)

<i>P. fengningensis</i>	BMNHC Ph-1158	China	nearly complete skull	lateral	poor	X		(Chiappe <i>et al.</i> , 2019a)
<i>P. fengningensis</i>	IVPP V1165	China	nearly complete skull	lateral	poor	X		(Chiappe <i>et al.</i> , 2019a)
<i>Pterygornis dapingfangensis</i>	IVPP V16363	China	partial skull	dorsolateral	good	X	X	(Wang, Li & Zhou, 2017a)
<i>P. dapingfangensis</i>	IVPP V20729	China	maxilla, nasal, jugal, quadratojugal, quadrate, angular, surangular	medial	good		X	(Wang <i>et al.</i> , 2017a)
<i>Rapaxavis pani</i>	DNHM D2522	China	nearly complete skull	dorsolateral	good	X		(Morschhauser <i>et al.</i> , 2009)
<i>Shangyang graciles</i>	IVPP V25033	China	partial skull	dorsal	poor	X	X	(Wang & Zhou, 2019e)
<i>Shanweiniao cooperorum</i>	DNHM D1878/1	China	partial skull	lateral	poor	X		(O'Connor <i>et al.</i> , 2009)
<i>Shengjingornis yangi</i>	PMOL-AB00179	China	nearly complete skull	lateral	poor	X		(Li, Wang, Zhang <i>et al.</i> , 2012)
<i>Shenqiornis mengi</i>	DNHM D2950/1	China	nearly complete skull	lateral	good	X		(Wang <i>et al.</i> , 2010b)
<i>Sinornis santensis</i>	BPV 538a/b	China	partial skull	lateral?	poor		X	(O'Connor & Dyke, 2010a)
<i>Sulcavis geeorum</i>	BMNH Ph-805	China	partial skull	lateral	good; several voids	X		(O'Connor <i>et al.</i> , 2013)
<i>Vescornis hebeiensis</i>	NIGPAS-130722	China	nearly complete skull	ventral	good; several voids	X		(Zhang <i>et al.</i> , 2004)
<i>Zhouornis hani</i>	BMNHC Ph-756	China	partial skull	lateral	excellent; juvenile		X	(Zhang <i>et al.</i> , 2014)

<i>Z. hani</i>	CNU VB0903	China	partial skull	lateral	good	X	X	(Zhang <i>et al.</i> , 2013)
<b>Non-avian Ornithuromorpha</b>								
<i>Apsaravis ukhaana</i>	IGM 100/1017	Mongolia	orbit, dentary, jugal?	lateral	excellent	X	X	(Clarke & Norell, 2002)
<i>Archaeorhynchus spathula</i>	IVPP V14287	China	partial skull	ventral; gastroliths	poor	X	X	(Zhou & Zhang, 2006)
<i>A. spathula</i>	IVPP V17075	China	nearly complete skull	dorsal	excellent; gastroliths	X		(Zhou <i>et al.</i> , 2013)
<i>A. spathula</i>	IVPP V17091	China	partial skull	dorsolateral	good	X	X	(Zhou <i>et al.</i> , 2013)
<i>A. spathula</i>	IVPP V20312	China	frontal, dentary, surangular	dorsolateral	good; gastroliths	X	X	(Wang & Zhou, 2016b)
<i>A. spathula</i>	STM 7-11	China	partial skull	dorsolateral	good	X	X	(Wang, O'Connor, Maina <i>et al.</i> , 2018)
<i>Archaeornithura meemannae</i>	STM 7-145	China	basicranium	cranial?	poor	X	X	(Wang <i>et al.</i> , 2015b)
<i>A. meemannae</i>	STM 7-163	China	basicranium	cranial?	poor	X	X	(Wang <i>et al.</i> , 2015b)
<i>Baptornis advenus</i>	AMNH 5101	USA	frontal, quadrate	3D	excellent		X	(Martin <i>et al.</i> , 1976) (lost according to Bell <i>et al.</i> , 2020)
<i>B. advenus</i>	FMNH 395	USA	mandible	3D	excellent		X	(Martin <i>et al.</i> , 1976)
<i>Dingavis longimaxilla</i>	IVPP V20284	China	nearly complete skull	lateral	poor; gastroliths	X		(O'Connor, Wang & Hu, 2016b)
<i>Enaliornis barretti</i>	SMC B54404	UK	braincase	3D	excellent		X	(Bell <i>et al.</i> , 2020; Elzanowski <i>et al.</i> , 1991)

<i>E. barretti</i>	YORYM 585	UK	braincase	3D	poor		X	(Elzanowski <i>et al.</i> , 1991)
<i>Eogranivora edentulata</i>	STM 35-3	China	premaxilla, nasal, frontal, dentary	dorsolateral	poor; ingested seeds; gastroliths	X	X	(Zheng <i>et al.</i> , 2018a)
<i>Hesperornis regalis</i>	KUVP 71012	USA	nearly complete skull	3D	excellent		X	(Bell <i>et al.</i> , 2020)
<i>H. regalis</i>	USNM V 4978	USA	partial skull	3D	excellent		X	(Elzanowski, 1991)
<i>H. regalis</i>	USNM V 6622	USA	premaxilla	3D	excellent		X	(Elzanowski, 1991)
<i>H. regalis</i>	YPM VP 903	USA	mandible	3D	excellent		X	NA
<i>H. regalis</i>	YPM VP 1206	USA	premaxilla, maxilla, nasal, dentary	3D	excellent		X	(Dumont <i>et al.</i> , 2016; Gingerich, 1973)
<i>H. regalis</i>	YPM VP 1207	USA	skull fragments including braincase	3D	good	X	X	(Elzanowski, 1991)
<i>cf. Hesperornis</i>	NUVF 286	Canada	teeth	3D	excellent		X	(Wilson, Chin & Cumbaa, 2016)
<i>Hongshanornis longicresta</i>	DNHM D2945/6	China	partial skull	lateral	poor; gastroliths	X		(Chiappe <i>et al.</i> , 2014)
<i>H. longicresta</i>	IVPP V14533	China	partial skull	lateral	poor	X		(Zhou & Zhang, 2005)
<i>Ichthyornis dispar</i>	ALMNH 3316	USA	premaxilla, maxilla, mandible	3D	excellent		X	(Field <i>et al.</i> , 2018b)
<i>I. dispar</i>	BHI 6421	USA	quadrate, quadratojugal, mandible	3D	excellent	X	X	(Field <i>et al.</i> , 2018b)
<i>I. dispar</i>	FHSM VP-18702	USA	partial skull	3D	good	X	X	(Field <i>et al.</i> , 2018b)



<i>I. dispar</i>	KUVP 119673	USA	jugal, quadrate, mandible	3D	excellent		X	(Field <i>et al.</i> , 2018b)
<i>I. dispar</i>	YPM VP 1450	USA	cranium, nasal, lacrimal, mandible	3D	good	X		(Field <i>et al.</i> , 2018b)
<i>I. dispar</i>	YPM VP 1459	USA	premaxilla	3D	excellent		X	(Field <i>et al.</i> , 2018b)
<i>I. dispar</i>	YPM VP 1728	USA	cranium, nasal, frontal	3D	good	X	X	(Field <i>et al.</i> , 2018b)
<i>I. dispar</i>	YPM VP 1735	USA	mandible	3D	good		X	(Clarke, 2004)
<i>I. dispar</i>	YPM VP 1749	USA	quadrate	3D	excellent		X	(Field <i>et al.</i> , 2018b)
<i>I. dispar</i>	YPM VP 1761	USA	mandible	3D	good		X	(Clarke, 2004)
<i>I. dispar</i>	YPM VP 1775	USA	mandible	3D	excellent		X	(Field <i>et al.</i> , 2018b)
<i>I. dispar</i>	YPM VP 6264	USA	mandible	3D	good	X	X	(Clarke, 2004)
<i>Iteravis huchzermeyeri</i>	BMNHC Ph-1318	China	nearly complete skull	lateral	good; gastroliths			(Chiappe <i>et al.</i> , 2016 pg. 104)
<i>I. huchzermeyeri</i>	BMNHC Ph-1343	China	nearly complete skull	dorsolateral	poor	X		(Chiappe <i>et al.</i> , 2016 pg. 106)
<i>I. huchzermeyeri</i>	BMNHC Ph-1392	China	partial skull	dorsolateral	good	X		(Chiappe <i>et al.</i> , 2016 pg. 187)
<i>Jianchangornis microdonta</i>	IVPP V16708	China	nearly complete skull	lateral	poor	X		(Zhou, Zhang & Li, 2009)
<i>Longicrusavis houi</i>	PKUP V1069	China	partial skull	ventrolateral	poor	X	X	(O'Connor <i>et al.</i> , 2010b)
<i>Parahesperornis alexi</i>	KUVP 2287	USA	complete skull	3D	poor	X		(Bell <i>et al.</i> , 2020)
<i>Pasquiaornis tankei</i>	RSM P2831.18	Canada	angular	3D	good		X	(Sanchez, 2010)
<i>P. tankei</i>	RSM P2831.52	Canada	quadrate	3D	excellent		X	(Sanchez, 2010)
<i>P. tankei</i>	RSM P2831.6	Canada	dentary	3D	good		X	(Sanchez, 2010)

<i>P. tankei</i>	RSM P2957.12	Canada	frontal	3D	good		X	(Sanchez, 2010)
<i>P. tankei</i>	RSM P2985.10	Canada	dentary	3D	good		X	(Sanchez, 2010)
<i>P. tankei</i>	RSM P2985.9	Canada	splénial	3D	good		X	(Sanchez, 2010)
<i>P. tankei</i>	RSM P2986.2	Canada	angular	3D	good		X	(Sanchez, 2010)
<i>P. tankei</i>	RSM P2988.11	Canada	dentary	3D	good		X	(Sanchez, 2010)
<i>P. tankei</i>	RSM P2988.25	Canada	quadrate	3D	excellent		X	(Sanchez, 2010)
<i>P. tankei</i>	RSM P2989.19	Canada	articular, prearticular, surangular	3D	good		X	(Sanchez, 2010)
<i>P. tankei</i>	RSM P2989.21	Canada	articular, prearticular, surangular	3D	excellent		X	(Sanchez, 2010)
<i>P. tankei</i>	RSM P2995.4	Canada	frontal	3D	excellent		X	(Bell <i>et al.</i> , 2020)
<i>P. tankei</i>	RSM P2995.5	Canada	maxilla	3D	excellent		X	(Sanchez, 2010)
<i>Patagopteryx deferrariisi</i>	MACN-N-11	Argentina	braincase, mandible	3D	excellent		X	(Chiappe, 2002)
<i>Piscivoravis lii</i>	IVPP V17078	China	dentary, quadrate, surangular	lateral	excellent; ingested fish			(Zhou, Zhou & O'Connor, 2014b)
<i>Potamornis skutchi</i>	UCMP 73103	USA	quadrate	3D	excellent		X	(Bell <i>et al.</i> , 2020)
<i>Schizooura lii</i>	IVPP V16861	China	complete skull	lateral	good	X		(Zhou <i>et al.</i> , 2012)
<i>Songlingornis linghensis</i>	IVPP V10913	China	premaxilla, dentary	lateral	?	X	X	(Hou, 1997) [drawing]
<i>Tianyuornis cheni</i>	STM 7-53	China	partial skull	lateral	poor; juvenile	X		(Zheng <i>et al.</i> , 2014)
<i>Xinghaiornis lini</i>	XHPM 1121	China	complete skull	dorsolateral	excellent			(Wang, Chiappe, Teng <i>et al.</i> , 2013)
<i>Yanornis martini</i>	BMNHC Ph-1043	China	nearly complete skull	lateral	good			(Chiappe <i>et al.</i> , 2016 pg. 100)
<i>Y. martini</i>	BMNHC Ph-928	China	nearly complete skull	lateral	poor	X		(Chiappe <i>et al.</i> , 2016pg. 102)

<i>Y. martini</i>	DNHM D3069	China	partial skull	ventrolateral	poor	X		(Chiappe <i>et al.</i> , 2016 pg. 166)
<i>Y. martini</i>	IVPP V12558	China	partial skull	lateral	poor	X		(O'Connor, 2019)
<i>Y. martini</i>	IVPP V13358	China	nearly complete skull	lateral	poor; gastroliths	X		(Zhou <i>et al.</i> , 2004)
<i>Y. martini</i>	STM 9-15	China	nearly complete skull	lateral	good; ingested fish	X		(Zheng <i>et al.</i> , 2014)
<i>Y. martini</i>	XHPM 1205	China	nearly complete skull	lateral	good	X		(Chiappe <i>et al.</i> , 2016 pg. 168)
<i>Yixianornis grabaui</i>	IVPP V12631	China	partial skull	dorsal	good	X		(Chiappe <i>et al.</i> , 2016 pg. 101)
<i>Y. grabaui</i>	IVPP V13631	China	partial skull	ventral	poor	X		(Clarke <i>et al.</i> , 2006)
<i>Zhongjianornis yangi</i>	IVPP V15900	China	nearly complete skull	dorsolateral	poor	X		(Zhou <i>et al.</i> , 2010)

Table 4.

**Summary table of which physical approaches a given complication has considerable effect upon.** Complications with an asterisk (\*) are those for which methods to mathematically correct for are established.

	Phylogenetic Signal*	Allometric Signal*	Many-To-One-Mapping	Liem's Paradox	Integration	Modularity	Behavioural Signals
Traditional Morphometrics	<b>X</b>	<b>X</b>	<b>X</b>		<b>X</b>	<b>X</b>	<b>X</b>

Geometric Morphometrics	<b>X</b>	<b>X</b>	<b>X</b>		<b>X</b>	<b>X</b>	<b>X</b>
Lever Modelling		<b>X</b>		<b>X</b>		<b>X</b>	<b>X</b>

# **Supplement for: The Diet of Early Birds Based on Modern and Fossil Evidence and a New Framework for its Reconstruction**

## **Supplemental Figures**

Each of the figures herein are comparisons between key graphs from the literature [redrawn] and revised versions produced using reanalysed data. The figures mimic the aesthetics of the original graphs to make comparisons easier. Specifics are described in each figure caption.




Fig. S1.

Principal Component Analysis [PCA] of avian pedal measurements. Fowler *et al.* (2009; 2011) use Correspondence Analysis [CA], designed to analyse discrete data, to analyse the continuous length measurements of the avian pes. (A) The original CA of Fowler *et al.* (2011) is provided, excluding

samples of *Deinonychus*. It generally resembles that of Fowler et al (2009) but has several additional strigiform samples included. This is contrasted to (B) PCA of the data used in (Fowler *et al.*, 2011) excluding *Deinonychus*. Note relative occupations of the morphospace by each group remains somewhat constant, but several differences are pertinent between the CA and PCA plots. In PCA: Pandionidae inhabits the same region of the morphospace as Strigiformes; Phasianidae, Piciformes, and Cathartidae cluster more closely to each other than any other groups; and in the set of Accipitridae, Falconidae, and Passeriformes, Passeriformes replaces Falconidae as the intermediary region and all three cluster more closely together.




Fig. S2.

Principal Component Analysis of theropod skulls. The analysis of (A,B) Foth and Rauhut (2013) [convex hulls added] defines nine taxa as carnivorous, herbivorous, or omnivorous that this study considers indeterminate. In addition, their study does not scale the variance of the inputs when computing principal components. (C,D) Reanalysis with these taxa labelled as indeterminate and input variances scaled maintains most carnivores in their own region of the morphospace and lessens the overlap between carnivore and herbivore taxa (though the extreme morphologies of *Anchiornis* and *Bambiraptor* cause the convex hulls of the two to overlap broadly). No hull is generated for omnivores as the only taxon included with convincing evidence of omnivory is *Lesothosaurus* (Sciscio, Knoll, Bordy *et al.*, 2017). Note that the Y-axis of (A,C) and both axes of (B,D) are inverted relative to the original publication as the sign of principal components calculated with the `prcomp` function in R is arbitrary (R Core Team 2019).





Fig. S3.

Principal Component Analysis of theropod skulls. (A,B) The analysis of Schaeffer *et al.* (2019) defines 14 taxa as herbivorous/omnivorous or as small carnivores that this study considers indeterminate. (C,D) Reanalysis with these taxa labelled as indeterminate decreases the overlap between herbivores/omnivores



and small carnivores primarily by removing therizinosaurs and avialans, respectively, from their groups. The total morphospace occupied by herbivores/omnivores is also greatly reduced by excluding oviraptorids which occupy their own unique area of the morphospace. PC1 and PC3 of landmark results are also plotted (E) to display their similarity to (A) and (C). This similarity is because PC3 of the landmark data describes similar shape variation to PC2 of the outline data (Figure 4 in Schaeffer *et al.*, 2019).



**Fig. S4.**  
Relationship between mechanical advantage and plant consumption in Passerines. Plot based on the appendix of Corbin *et al.* (2015), (A) including and (B) excluding *Zenaida macroura*. *Z. macroura* is the

only columbiform bird included in the otherwise passerine data set. Correlation including this point included is weak to nonexistent, but of moderate strength when the data is restricted to passerine taxa.




Fig. S5.

Biomechanical morphospace of Dinosauria. (A) The analysis of Button and Zanno (2020) incorporates 34 functional metrics to examine the affinities of various dinosaur groups in order to investigate convergence in dietary adaptation. Their observed trends are maintained in (B) an analysis incorporating only nine of their measured indices which have both theoretical validity and are shown to discriminate diet among extant animals [their C2–8, 22, and 23]. The most noteworthy difference in trends is that in (B) there is more separation in the morphospace between edentulous theropods [Ornithomimosauria and Oviraptorosauria] and sauropods along the PC2 axis. Note also that (B) accounts for a much higher proportion of the variance in the first two principal components [60.4% as opposed to 32.2% in (A)].

Both axes of the graphs are inverted relative to the original publication. An interactive three-dimensional graph of the data can be generated from the supplemental R code.

## Supplemental Tables

The following are tables to assist in interpreting the details of this review. Specifics are given with each table caption.

Table S1.

Calculation of cranial connective properties of *Shenqiornis*. Connective tissue properties recorded in the literature are compiled and used to estimate connective tissue properties for *Shenqiornis*. Regressions of cross-sectional area [X-Sect] and Young's Modulus [E] were made vs. [body mass]<sup>0.33</sup> using a polynomial [Poly] and power [Pow] fit to the properties of cranial sutures. As Cost *et al.* (2019) considered dog patellar tendon as a potential model, an additional power-fit regression using the full sample [PowFS] was also made. Macaques were excluded from all regressions as outliers. The properties of the Pow regression are used in Figure 6C; Poly and PowFS properties produced dislocation similar to Figure 6A and B.

Animal	Part	Mass (g)	X-Sect (mm <sup>2</sup> )	E (MPa)	Source	Mass Source
Human Child	Cranial suture	27260.90	14.06	1100.00	(Davis, Loyd, Shen <i>et al.</i> , 2012)	(Davis <i>et al.</i> , 2012)
Human Infant	Cranial suture	10714.29		381.48	(Wang, Zou, Li <i>et al.</i> , 2014)	WHO weight-for-age 50th percentile at 18 months, weighted by gender
Human Postnatal	Cranial suture	8650.00		0.60	(Grau, Daw, Patel <i>et al.</i> , 2006)	WHO weight-for-age 50th percentile at 9 months
Macaque (male)	Cranial suture	5360.00		7700.00	(Kupczik, Dobson, Fagan <i>et al.</i> , 2007)	(Smith & Jungers, 1997)
Macaque (female)	Cranial suture	3590.00		1900.00	(Kupczik <i>et al.</i> , 2007)	(Smith <i>et al.</i> , 1997)
Mouse	Cranial suture	24.00	0.13	0.58	(Chien, Wu, Chao <i>et al.</i> , 2008)	(Chien <i>et al.</i> , 2008)
Lewis Rat	Cranial suture	250.50	0.51	2.35	(Chien <i>et al.</i> , 2008)	(Chien <i>et al.</i> , 2008)

Wistar Rat	Cranial suture	62.12		1.08	(McLaughlin, Zhang, Pashley <i>et al.</i> , 2000)	(Extrapolated from Novelli, Diniz, Galhardi <i>et al.</i> , 2007)
Rabbit	Facial suture	1590.57		1.27	(Radhakrishnan & Mao, 2004)	(Jones, Bielby, Cardillo <i>et al.</i> , 2009)
Goat	Nasal suture	46900.00		400.00	(Farke, 2008)	(Jones <i>et al.</i> , 2009)
Camel	Nuchal ligament	488000.00	850.00	0.55	(Dimery, Alexander & Deyst, 1985)	(Jones <i>et al.</i> , 2009)
Deer	Nuchal ligament	20000.00	51.60	0.61	(Dimery <i>et al.</i> , 1985)	(Dimery <i>et al.</i> , 1985)
Sheep	Nuchal ligament	21000.00	84.00	0.57	(Dimery <i>et al.</i> , 1985)	(Dimery <i>et al.</i> , 1985)
Pig	Palatal suture	84500.00	462.71	47.43	(Savoldi, Xu, Tsoi <i>et al.</i> , 2018)	(Jones <i>et al.</i> , 2009)
Dog	Patellar tendon	26700.00	0.24	30.80	(Haut, Lancaster & DeCamp, 1992)	(Haut <i>et al.</i> , 1992)
Reptile (validated in <i>Sphenodon</i> ?)	Suture	195.29		10.00	(Curtis, Jones, Evans <i>et al.</i> , 2013)	(Average of Herrel, Moore, Bredeweg <i>et al.</i> , 2010 Supplement 1)
<i>Shenqiornis</i> Poly		340.00	0.63	2.76	Regressions	Table 1, this review
<i>Shenqiornis</i> Pow		340.00	7.07	2.86	Regressions	Table 1, this review
<i>Shenqiornis</i> PowFS		340.00	0.68	2.51	Regressions	Table 1, this review

## Supplemental References

- BUTTON D. J. & ZANNO, L. E. (2020). Repeated evolution of divergent modes of herbivory in non-avian dinosaurs. *Current Biology* **30**, 158-168. e4.
- CHIEN C. H., WU, Y. D., CHAO, Y.-J., CHEN, T., CHEN, W. F., YU, J. C. & LI, X. (2008). The effects of different cranial modules on mechanical properties of cranial suture in Lewis Rats and same-aged C57BL/6 Mice. *Strain* **44**, 272-277.
- CORBIN C. E., LOWENBERGER, L. K. & GRAY, B. L. (2015). Linkage and trade-off in trophic morphology and behavioural performance of birds. *Functional Ecology* **29**, 808-815.
- COST I. N., MIDDLETON, K. M., SELLERS, K. C., ECHOLS, M. S., WITMER, L. M., DAVIS, J. L. & HOLLIDAY, C. M. (2019). Palatal biomechanics and its significance for cranial kinesis in *Tyrannosaurus rex*. *The Anatomical Record: Advances in Integrative Anatomy and Evolutionary Biology*.
- CURTIS N., JONES, M. E. H., EVANS, S. E., O'HIGGINS, P. & FAGAN, M. J. (2013). Cranial sutures work collectively to distribute strain throughout the reptile skull. *Journal of the Royal Society Interface* **10**, 20130442.
- DAVIS M. T., LOYD, A. M., SHEN, H.-Y. H., MULROY, M. H., NIGHTINGALE, R. W., MYERS, B. S. & BASS, C. D. (2012). The mechanical and morphological properties of 6 year-old cranial bone. *Journal of Biomechanics* **45**, 2493-2498.
- DIMERY N. J., ALEXANDER, R. M. & DEYST, K. A. (1985). Mechanics of the ligamentum nuchae of some artiodactyls. *Journal of Zoology* **206**, 341-351.
- FARKE A. A. (2008). Frontal sinuses and head-butting in goats: a finite element analysis. *Journal of Experimental Biology* **211**, 3085-3094.
- FOTH C. & RAUHUT, O. W. M. (2013). Macroevolutionary and morphofunctional patterns in theropod skulls: a morphometric approach. *Acta Palaeontologica Polonica* **58**, 1-16.
- FOWLER D. W., FREEDMAN, E. A. & SCANNELLA, J. B. (2009). Predatory functional morphology in raptors: interdigital variation in talon size is related to prey restraint and immobilisation technique. *PLOS ONE* **4**, e7999.
- FOWLER D. W., FREEDMAN, E. A., SCANNELLA, J. B. & KAMBIC, R. E. (2011). The predatory ecology of *Deinonychus* and the origin of flapping in birds. *PLOS ONE* **6**, e28964.
- GRAU N., DAW, J. L., PATEL, R., EVANS, C., LEWIS, N. & MAO, J. J. (2006). Nanostructural and nanomechanical properties of synostosed postnatal human cranial sutures. *Journal of Craniofacial Surgery* **17**, 91-98.
- HAUT R. C., LANCASTER, R. L. & DECAMP, C. E. (1992). Mechanical properties of the canine patellar tendon: some correlations with age and the content of collagen. *Journal of Biomechanics* **25**, 163-173.
- HERREL A., MOORE, J. A., BREDEWEG, E. M. & NELSON, N. J. (2010). Sexual dimorphism, body size, bite force and male mating success in tuatara. *Biological Journal of the Linnean Society* **100**, 287-292.
- JONES K. E., BIELBY, J., CARDILLO, M., FRITZ, S. A., O'DELL, J., ORME, C. D. L., SAFI, K., SECHREST, W., BOAKES, E. H., CARBONE, C., CONNOLLY, C., CUTTS, M. J., FOSTER, J. K., GRENYER, R., HABIB, M., PLASTER, C. A., PRICE, S. A., RIGBY, E. A., RIST, J., TEACHER, A.,

- BININDA-EMONDS, O. R. P., GITTLEMAN, J. L., MACE, G. M. & PURVIS, A. (2009). PanTHERIA: a species-level database of life history, ecology, and geography of extant and recently extinct mammals. *Ecology* **90**, 2648-2648.
- KUPCZIK K., DOBSON, C. A., FAGAN, M. J., CROMPTON, R. H., OXNARD, C. E. & O'HIGGINS, P. (2007). Assessing mechanical function of the zygomatic region in macaques: validation and sensitivity testing of finite element models. *Journal of Anatomy* **210**, 41-53.
- MCLAUGHLIN E., ZHANG, Y. G., PASHLEY, D., BORKE, J. & YU, J. C. (2000). The load-displacement characteristics of neonatal rat cranial sutures. *The Cleft Palate-Craniofacial Journal* **37**, 590-595.
- NOVELLI E. L. B., DINIZ, Y. S., GALHARDI, C. M., EBAID, G. M. X., RODRIGUES, H. G., MANI, F., FERNANDES, A. A. H., CICOGNA, A. C. & NOVELLI FILHO, J. L. V. B. (2007). Anthropometrical parameters and markers of obesity in rats. *Laboratory animals* **41**, 111-119.
- R CORE TEAM. (2019). prcomp function | R Documentation Version 3.6.2.
- RADHAKRISHNAN P. & MAO, J. J. (2004). Nanomechanical properties of facial sutures and sutural mineralization front. *Journal of dental research* **83**, 470-475.
- SAVOLDI F., XU, B., TSOI, J. K. H., PAGANELLI, C. & MATINLINNA, J. P. (2018). Anatomical and mechanical properties of swine midpalatal suture in the premaxillary, maxillary, and palatine region. *Scientific Reports* **8**, 1-12.
- SCHAEFFER J., BENTON, M. J., RAYFIELD, E. J. & STUBBS, T. L. (2019). Morphological disparity in theropod jaws: comparing discrete characters and geometric morphometrics. *Palaeontology* **63**, 283-299.
- SCISCIO L., KNOLL, F., BORDY, E. M., DE KOCK, M. O. & REDELSTORFF, R. (2017). Digital reconstruction of the mandible of an adult *Lesothosaurus diagnosticus* with insight into the tooth replacement process and diet. *PeerJ* **5**, e3054.
- SMITH R. J. & JUNGERS, W. L. (1997). Body mass in comparative primatology. *Journal of Human Evolution* **32**, 523-559.
- WANG J., ZOU, D., LI, Z., HUANG, P., LI, D., SHAO, Y., WANG, H. & CHEN, Y. (2014). Mechanical properties of cranial bones and sutures in 1–2-year-old infants. *Medical Science Monitor: International Medical Journal of Experimental and Clinical Research* **20**, 1808.