

The Unique Role of Jason-2 Geodetic Mission for High Resolution Gravity Field Modelling

Ole Baltazar Andersen¹, Shengjun Zhang², David T. Sandwell³, Gérald Dibarboure⁴, Walter H.F. Smith⁵, and Adili Abulaitijiang⁶

¹DTU Space, National Space Institute

²School of Resources and Civil Engineering Northeastern University

³UCSD

⁴CNES

⁵National Oceanic and Atmospheric Administration (NOAA)

⁶DTU space, National Space Institute

November 26, 2022

Abstract

The Geodetic Mission (GM) of Jason-2 was planned to provide ground-tracks with a systematic spacing of 4 km after 2 years and 2 km after 4 years to increase the spatial resolution of global altimetric gravity fields. Jason-2 ceased operation after 2 years of GM but provided a fantastic dataset. We highlight and evaluate the improvement to the gravity field which has been derived from the GM. The ageing Jason-2 suffered from several safe-holds and instrument outages. Here, we try to quantify the effect of safe-holds on marine gravity and discuss suitable approaches advising future GM like Jason-3. We evaluate the importance of attempting to “rewind” the mission to recover missing tracks as well as the possibility to continue an existing GM by using the same orbital plane. The latter idea would allow bisecting the already 2-years Jason-2 GM creating a 2 km grid after 2 years of Jason-3 GM.

Geophysical Research Letters

Supporting Information for

The Unique Role of Jason-2 Geodetic Mission for High Resolution Gravity Field Modelling

O. B. Andersen¹, S. Zhang², D. T. Sandwell³, G. Dibarboure⁴, W. H. F. Smith⁵, A. Abulaitijiang¹

¹ DTU Space, Technical University of Denmark, Lyngby, Denmark

² School of Resources and Civil Engineering, Northeastern University, Shenyang, China

³ Institute of Geophysics and Planetary Physics, Scripps Institute of Oceanography, University of California San Diego, La Jolla, CA92093, USA

⁴ CNES, Av. Edouard Belin, Toulouse, France,

⁵ NOAA, 5830 Univ Research Court, College Park, MD, USA

Corresponding author: O. B. Andersen (oa@space.dtu.dk)

Content of this document:

Table S1.

Figure S1.

Text S1.

Text S2.

References

Supporting material Table S1.

	Jason-1	Jason-2
Altitude	1324 km	1309 km
Period	7 th May 2012 until 21 st of June 2013,	14 th Sep 2017 until 1 st of October 2019,
LRO cycle length	406 days	371 days
GM total length	411 days	371+371 days = 742 days
Sub cycles	3.9, 10.9, 47.5, 179.5 days	4, 17, 79, 145 days

Table S1. Orbital characteristics of the LRO of Jason-1 and Jason-2 satellites.

Supporting material Figure S1.




Figure S1. Geographic distribution and values of marine gravity measurements. These marine gravity observations have previously been used in the derivation of the EGM2008 gravity model (Pavlis et al., 2013) and have been extensively edited for outliers.

Supporting information Text S1.

We conducted an investigation to establish that the range precision is comparable between the measurements from the two satellites. We apply the two-pass waveform retracker proposed by Sandwell and Smith (2005) to enhance the range precision. Figure 3 shows estimates of noise level for Jason-1 (left) and Jason-2 (right) as a function of significant wave height for global sample tracks. For both Jason-1 and Jason-2 the range precision is improved from 68 mm before retracking to 42 mm after retracking for 2 meters of significant wave height (SWH) (red and blue curves in Figure S2) for individual points (dots in upper figures) and the median over 0.5 m SWH bins (lines in lower figures).

Supporting information Figure S2




Figure S2. Standard deviation of retracked height with respect to EGM2008 for Jason-1 and Jason-2 sample cycle. Upper figures statistics for individual points. Lower figures: medians over 0.5 meter SWH intervals. (Red: height from sensor geophysical data record; Green: height from first step of two-pass retracking; Blue: height from second step of two-pass retracking).

Supporting information Text S2.

The first step is to retrack the raw SGDR waveforms from Jason-1 and Jason-2 altimeter missions using a

two-pass waveform retracker (Sandwell et al. 2014). The second step is to low-pass filter and resample the 20-Hz retracked height into ~ 5 Hz, to enhance the signal to noise. The third step is to correct the retracked height using state of the art geophysical and range corrections (Andersen and Scharroo, 2011). The fourth step is to perform outlier editing through comparing height and along-track slopes with the associated heights and along track slopes from EGM2008 as described by Zhang et al., (2020). The fifth step is to remove the slopes of the EGM2008 geoid and the slopes of the Mean Dynamic Topography (DOT_min1x1_EGM08) associated with the EGM2008 geoid) and to apply a low-pass filter in order to obtain along-track filtered sea surface height gradients.

Finally, the along-track sea surface slopes were turned into residual vertical deflections and then residual gravity anomalies. Subsequently the marine gravity anomalies at $1' \times 1'$ resolution were then computed by restoring the EGM2008 gravity field associated with the EGM2008 geoid model (Zhang et al., 2017).

References

- Pavlis, N. K., Holmes, S. A., Kenyon, S. C., & Factor, J. K. (2012), The development and evaluation of the Earth Gravitational Model 2008 (EGM2008), *J. Geophys Res*, D. 117, B4, <https://doi.org/10.1029/2011JB008916>
- Sandwell, D. T., Dietmar-Müller R., Smith W., Garcia, E. & Francis, R. (2014), New global marine gravity model from CryoSat-2 and Jason-1 reveals buried tectonic structure. *Science*. 346 no. 6205 pp. 65-67
- Sandwell, D. T., and W. H. F. Smith (2005), Retracking ERS-1 altimeter waveforms for optimal gravity field recovery, *Geophys. J. Int.*, 163, 79– 89, doi:10.1111/j.1365-246X.2005.02724.x
- Zhang, S., Andersen, O., Kong X., & Li H., (2020), Inversion and Validation of Improved Marine Gravity Field Recovery in South China Sea by Incorporating HY-2A Altimeter Waveform Data. *Remote Sensing*, 12(5), 802. <https://doi.org/10.3390/rs12050802>.
- Zhang S., Sandwell D., Jin T., & Li D., (2017), Inversion of Marine Gravity Anomalies over Southeastern China Seas from Multi-Satellite Altimeter Vertical Deflections. *J of Applied Geophysics*, 137, 128-137.

The Unique Role of Jason-2 Geodetic Mission for High Resolution Gravity Field Modelling

O. B. Andersen¹, S. Zhang², D. T. Sandwell³, G. Dibarboure⁴, W. H. F. Smith⁵, A. Abulaitijiang¹

¹ DTU Space, Technical University of Denmark, Lyngby, Denmark

² School of Resources and Civil Engineering, Northeastern University, Shenyang, China

³ Institute of Geophysics and Planetary Physics, Scripps Institute of Oceanography, University of California San Diego, La Jolla, CA92093, USA

⁴ CNES, Av. Edouard Belin, Toulouse, France,

⁵ NOAA, 5830 Univ Research Court, College Park, MD, USA

Corresponding author: O. B. Andersen (oa@space.dtu.dk)

Key Points

Jason-2 is unique in performing a controlled geodetic mission as part of the Extension-of-life phase with 4 km groundtrack spacing.

Mission-rewind to recover missing tracks following safe-holds for Jason-2 was found to be extremely important for gravity field modelling.

Our finding support interleaving a possible Jason-3 GM with Jason-2 to bisect the Jason-2 GM creating a 2 km GM after 2 years.

Abstract

The Geodetic Mission (GM) of Jason-2 was planned to provide ground-tracks with a systematic spacing of 4 km after 2 years and 2 km after 4 years to increase the spatial resolution of global altimetric gravity fields. Jason-2 ceased operation after 2 years of GM but provided a fantastic dataset. We highlight and evaluate the improvement to the gravity field which has been derived from the GM. The ageing Jason-2 suffered from several safe-holds and instrument outages. Here, we try to quantify the effect of safe-holds on marine gravity and discuss suitable approaches advising future GM like Jason-3. We evaluate the importance of attempting to “rewind” the mission to recover missing tracks as well as the possibility to continue an existing GM by using the same orbital plane. The latter idea would allow bisecting the already 2-years Jason-2 GM creating a 2 km grid after 2 years of Jason-3 GM.

Plain Language Summary.

36 In this paper we investigate the final extension-of-Life phase of Jason-2 and its importance and
37 impact on high resolution gravity field mapping. Jason-2 has been a unique satellites in the sense
38 that it performed a controlled and accurate mapping of the Ocean's surface 4 km resolution during
39 2 years of operation for the first time.

40 During the extension-of-life phase the ageing Jason-2 satellite encountered several safe-holds
41 resulting in missing tracks and no available sea surface heights observations. For the first time, a
42 mission rewind maneuver was performed by the space agencies to recover the missing tracks and
43 we demonstrate its fundamental importance to gravity field modelling and suggest this to be
44 considered for future geodetic missions.

45 For the upcoming extension-of-life phase of Jason-3 we suggest interleaving Jason-3 with Jason-2
46 to bisect the already 2-years Jason-2 GM creating a 2 km GM after only 2 years and this way
47 mapping even finer scale gravity field signal.

48

49 1 Introduction

50 A number of satellite altimeters have performed a "Geodetic Mission" (GM) during their
51 lifetime (i.e. Geosat, ERS-1, Cryosat-2, Jason1/2 and Saral/AltiKa). The GM is basically a
52 Long Repeat Orbit (LRO) where the orbital pattern is designed for mainly geodetic purposes.
53 This means that the spatial sampling is optimized to map short wavelength in the geoid or
54 gravity field at the price of no or long temporal sampling. Typically, the GM consist of one or
55 more repeated or interleaved LRO (Cryosat-2 has repeated LRO, Jason-2 interleaved LRO).
56 For geodetic purposes the smallest possible cross-track resolution is the ultimate goal in
57 order to map the finest scales in the gravity field as the cross-track distance governs the
58 gravity signal which can be resolved. As an example: 8 km resolution requires a little more
59 than 1 year GM and 4 km requires a little more than 2 years of SSH observations at Jason
60 orbital altitude.

61 Exact repeat missions (ERM) are primarily designed for oceanographic purposes to map
62 oceanographic signals optimally. This requires frequent temporal sampling at the price of
63 coarse spatial sampling. As an example, the Jason-1 ERM sampled the ocean every 9.9156
64 days at 314 km across-track spatial sampling. Subsequently, the satellite was moved into a
65 LRO mission with 7.5-km across-track spatial sampling but with a temporal sampling of 406
66 days.

67 The GM and ERM missions mutually support each other in the sense that the GM drives the
68 mapping of the fine structures in the Mean Sea Surface (MSS) model (Andersen et al.,
69 2015), which are applied to derive accurate sea level anomalies for the ERM. ERM are
70 important to derive long-term mean for the MSS. With more satellites flying in recent years,
71 ERM are also becoming increasingly important in determining ocean variability, which in-
72 turn can be used to correct the GM data (Dufau et al., 2016).

73 When the Jason satellites have served their main commitment to oceanographic science
74 and ensured the tandem mission obligations with future missions in the same orbit and the
75 satellites are getting toward the end of their lifetime an Extension of Life (EoL) phase is
76 considered. During the EoL phase the old satellite is moved away from the nominal orbit
77 located at 1336 km altitude. Moving the satellite away from the nominal orbit prevents a

78 collision of the satellite with active and future missions that must fly in a prescribed orbital
79 tube to achieve multi-decadal measurement along the tracks that were initiated by TOPEX
80 in 1992. Through orbital maneuvers, the satellite is lowered or raised a number of kilometers
81 into the EoL orbit which eventually will become the graveyard orbit for the satellite.

82 During their EoL missions, Long Repeat Orbits (LRO) were selected for both Jason-1 and
83 Jason-2, where for each the repeat was longer than 1 year. Contingent on the remaining
84 lifetime of the satellites, the LRO could be interleaved to create a GM with very high spatial
85 resolution in a systematic and controlled way.

86 Resolution of current global altimetric gravity models is around 12 km wavelength resolving
87 6km (Sandwell et al. 2014, Andersen et al., 2017) partly limited by the 8-km groundtrack
88 spacing of previous GM (Geosat, ERS-1, Cryosat-2). The Jason-2 is unique in this way, as
89 it is the first GM mission planned so that the ground-track distances could be systematically
90 bisected beyond 8 km to provide 4 km at 2 years and 2 km after 4 years. The ongoing
91 SARAL/AltiKa mission provides the most accurate sea level observations (Sandwell et al.,
92 2019), but the SARAL GM is un-controlled, making it more difficult to resolve the short
93 wavelengths in the gravity field in a systematic way.

94 During the EoL both Jason-1 and Jason-2 have suffered from a number of safe-holds
95 causing one or more of the instruments onboard the satellite to be shut-off. This is typically
96 due to ageing of the instrument onboard the satellite or due to collision avoidance. Most
97 noticeable is the last safe-hold of Jason-2 causing the instrument to be shut-off for around
98 100 days.

99 The errors in the derived altimetric marine gravity grids originate in omission and
100 commission errors (Pujol et al., 2018). The omission errors will be dominated by the spatial
101 distribution of the data. Lack of data, related to duration and safe-holds will increase this
102 error. The commission errors are largely related to measurement errors such as the range
103 precision, the retracker and oceanographic noise, but also errors due to an imperfect
104 gridding process. In this paper we mainly study the omission error on the gravity field
105 modelling due to the spatial distribution of data.

106 We have investigated altimetric marine gravity fields from the Jason-1 and Jason-2 Long
107 Repeat Orbits or GM in order to demonstrate the value of designing these EoL missions with
108 multiple LRO cycles interleaved to gather the best spatial coverage. Safe-holds will
109 ultimately lead to a degradation of the derived gravity field and we quantify the impact of
110 these using observations from sub-cycles of Jason-1 and Jason-2. As an example, we
111 derived gravity from the two 378-days LRO cycles of Jason-2 and the two 178-days sub-
112 cycle of Jason-1 as these (sub)cycles were affected differently by safe-holds. Such analysis
113 is important to guide future EoL missions (most profoundly the EoL for Jason-3) and their
114 strategies to remedy the effect of future safe-holds.

115

116 2 Geodetic Mission orbit choice for Jason-1 and 2

117

118 Jason-1 was launched in 2002. When it had served its main commitment to oceanographic
119 science and ensured the tandem mission obligations towards Jason-2 an EoL mission was
120 researched and initiated in 2012 (Bronner and Dibarboure, 2012). During the EoL for Jason-
121 1 the satellite was put into a LRO with a 406-day cycle with a ground track resolution of 7.5
122 km serving geodetic purposes (Sandwell et al. 2014). The Jason-1 GM lasted from 7 May
123 2012 until 21 June 2013 when the mission was terminated due to instrument failure.
124 Fortunately, Jason-1 collected exactly one full GM cycle of 406 days before the mission was
125 terminated shortly after, when the orbit became the graveyard orbit for Jason-1.

126
127 When designing the LRO for both Jason-1 and Jason-2 a number of simulations were
128 performed (Dibarboure, 2012; Dibarboure and Morrow, 2016) and a number of orbit choices
129 were investigated prior to the selection of the final orbit. The simulations are performed to
130 optimize the usefulness of the LRO for both geodesy and oceanography by designing the
131 orbit with a number of sub-cycles of varying length. During each sub-cycle a near-regular
132 ground track pattern is measured. This pattern automatically shifts longitudinally for each
133 following sub-cycle. The choice and duration of the sub-cycles are normally governed by
134 their utility for oceanographic purposes, but also with consideration of their value to geodesy
135 in the event that the GM terminates early due to satellite failure. These considerations are
136 important for the subsequent gravity field modelling in two ways. First, they ensure that safe-
137 holds will result in outages scattered evenly throughout the globe. Second, they enable the
138 possibility of rewinding the LRO by one or more sub-cycles in case of longer safe-holds.
139 This proved particularly important for the final cycle of Jason-2. The final choice of orbits
140 and sub-cycles for Jason-1 and Jason-2 LRO are shown in Table S1.

141
142 Upon designing the EoL of Jason-2 one could argue to inject Jason-2 in another 406 days
143 LRO interleaved with the Jason-1 GM in a similar orbit to speed up geodetic sampling and
144 get a 4 km sampling by combining the 406 days of Jason-1 and 2. Unfortunately, such
145 interleaved GM would require Jason-2 EoL to use exactly the same orbital altitude of Jason-
146 1GM, which was avoided due to collision risks. Consequently, another EoL orbit at another
147 altitude had to be selected. Such orbit provided irregular sampling with the Jason-1GM
148 where the tracks were on nearly identical locations in some regions but perfectly interleaved
149 in other regions. These so-called moiré patterns appear when two grids of different
150 resolution are superimposed (Dibarboure et al, 2012).

151 The consequence of this is, that Jason-2 was planned to perform its own dedicated multiyear
152 GM gradually filling up the globe with denser and denser ground tracks through multiple
153 cycles of interleaved LRO.

154
155 Through a number of simulations following the work by Dibarboure (2012) and Dibarboure
156 and Morrow, 2016) a LRO orbit for Jason-2 with clear advantages to both geodetic and
157 oceanographic research was selected and on July 11th 2017 Jason-2 began measuring the
158 first cycle of the LRO. This had a 371-day repeat period at an altitude of 1309km (27km
159 lower than the nominal TOPEX altitude). This resulted in an across track distance of around
160 8.5 km at the Equator.

161 The partnership between NOAA, NASA and CNES agreed to extend the Jason-2 mission
162 for an additional two years, from 1 January 2018 until the end of 2019 considering to extend
163 with EoL with two further cycles thereby lowering the groundtrack distance by a factor of two
164 to a little over 2 km after 4 years.

165 On July 18th 2018 Jason-2 successfully completed the first LRO cycle and operations started
166 to move the satellite into its new groundtrack in-between the ground tracks of the first LRO
167 cycle. This entailed a shift of the ground track of a little more than 4 km, which was completed
168 on the 25th of July, where the second LRO cycle was initiated. In theory, the second LRO
169 cycle should be completed by 31 July 2019 resulting in a systematic groundtrack distance
170 of a little more than 4 km. Unfortunately, Jason-2 only managed to perform 350 days of the
171 planned 371 days of the second LRO before the mission was terminated on October 8th,
172 2019.

174 2.1 Safe-holds

176 When the EOLs of the Jason satellites were initiated, both satellites were around 10 years
177 old and ageing, and during the LRO both satellites encountered safe-holds to safeguard
178 the instrument and to extend the mission as long as possible. These are shown in Table 1.
179

Satellite	Start date	End date	Duration
Jason-1	28/02/2013	18/03/2013	18 days
Jason-2 Cycle 1	14/09/2017	13/10/2017	30 days
	20/02/2018	02/03/2018	9 days
Jason-2 Cycle 2	19/10/2018	25/10/2018	6 days
	26/12/2018	07/01/2019	14 days
	16/02/2019	24/05/2019	100 days (21 days*)

180

181 **Table 1.** Safe-holds for Jason-1 and Jason-2 during the two LRO cycle. Courtesy of Christoph
182 Marechal, CNES. *se explanation in the text on mission “rewind” maneuver to remedy the safe-
183 hold.

184

185 During the LRO Jason-1 completed one sub-cycle of 179 days without safe-hold but suffered
186 one safe-hold of 18 days during the second sub-cycle. For Jason-2 the story is more
187 dramatic. Both LRO cycles suffered from several safe-holds lasting a total of more than 30
188 days. The last and most severe safe-hold lasted 100 days from February 16th, 2019 until
189 May 24th 2019. The second LRO cycle should, in theory have be completed by July 31st
190 2019, but the partnership between NOAA, NASA and CNES agreed to conduct an orbital
191 maneuver and “rewind“ the mission by 79 days to recover the missing observations.
192 Rewinding the mission to recover gaps is possible because the LRO orbit is designed with
193 multiple interleaved sub-cycles and a relatively cheap maneuver (in terms of fuel) can
194 “rewind” the mission by a sub-cycle (e.g., 17, 79 or 145 days). It is, in theory, possible to
195 rewind the mission by any amount of days, but at significant increased fuel cost and this is
196 normally avoided. By rewinding the mission by 79 days the resulting gap in data collection
197 due to the safe-hold was limited to 21 days. In theory, the second LRO should have been
198 completed on October, 21st 2019. Unfortunately, the instruments ceased working just 20
199 days before this date.

200

201 3 Marine gravity from Jason subsets


202

203 To aid in the design of future GM we have investigated a number of different sub-sets of
204 Jason data. First of all, we studied the importance of establishing a GM with multiple LRO
205 cycles interleaved to gather the best spatial coverage and hereby lowering the cross-track
206 distance as much as possible in a controlled way. Here we compared gravity from the first
207 178-days sub-cycle of Jason-1 (cross track distance 17 km) with gravity from the first cycle
208 of Jason-2 (cross track distance of 8.5 km) and gravity derived from the full 406 days GM
209 (cross track distance = 7.5 km) and finally gravity derived from the full 2x371 days GM of
210 Jason-2 (cross track distance of 4.25 km). Figure 1 illustrates the altimetric data in a subset
211 close to Bermuda in the Northwest Atlantic Ocean.

212 Safe-holds have shown to have a significant impact on the quality of the derived gravity field.
213 In order to quantify the impact of these we compared gravity computations using
214 observations the two 371-days cycles of Jason-2 and the two 178-days sub-cycle of Jason-
215 1 as these (sub)-cycles were affected differently by safe-holds. During the first 179-days
216 sub-cycle of Jason-1 the mission only encountered normal accidental outages of around 10
217 tracks. During the second sub-cycle the satellite encountered 18 days or 10% data-loss.

218 Jason-2 encountered 2 safe-holds during the first LRO cycle, losing data for 39 days or 11%
219 data loss, and 41 days safe-hold plus 20 days early failure resulting in 17% data loss for the
220 second LRO. The geographical distribution of the tracks is seen in Figure 1.

221



222

223

224 **Figure 1.** Geographic distribution of Jason GM altimeter measurements for a section in the NW Atlantic
225 Ocean close to Bermuda (in grey). Upper left: J1 sub-cycle 1, Upper right: Jason 1 sub-cycle 2; Center left:
226 J2 LRO Cycle 1, Center right: Jason 2 LRO cycle 2; Lower left: J1 Entire GM; Lower right: Jason 2 Entire GM
227 (Both LRO cycles)

228

229 **3.1 Marine Gravity observations**

230 A high-precision dataset with its assessed accuracy superior to ~2 mGal was obtained through a
231 cooperation with the (U.S.) National Geospatial-Intelligence Agency (NGA). Over 1.4 million high
232 quality measurements are distributed within the northwest Atlantic Ocean bounded by (20°~90°W,
233 20°~55°N) and their observed marine gravity anomalies are shown in Figure S1.

234

235 4 Geoid slope and gravity anomalies evaluation

236 The Sensor Geophysical Data Record (SGDR) altimeter data products including 20 Hz
237 waveforms are obtained from the Archiving, Validation and Interpretation of Satellite
238 Oceanographic (AVISO) data service. In order to compare results between Jason-1 and
239 Jason-2 it is important that the two datasets have the same range precision to ensure that
240 the differences we are seeing are not due to different commission errors related to the
241 instrument onboard the two satellites. This investigation is described in detail in Supporting
242 information Text S1

243 Gravity anomalies can be derived from altimetric sea surface height observations by
244 isolating the geoid height (Andersen et al., 2017) or from the geoid slopes (Sandwell et al.,
245 2014). In this investigation, we decided to derive the gravity anomalies using the geoid
246 slopes using the method detailed in Supporting information Text S2.

247 A global evaluation of the impact of the various combinations of the Jason subsets can be
248 performed by comparing with the multi-mission global slope grid SS V28.1. The median
249 absolute deviation of the along-track slopes with respect to the full model is a good indication
250 of the un-modelled signal in the Jason subsets combined with the noise in the altimeter
251 profiles (Sandwell et al., 2019). The median absolute deviation of the along-track slope data
252 with respect to the full (V28.1) slope grids is calculated and gridded within the latitudinal
253 range of 66°N and 66°S. These are shown in Figure 2 which illustrates the oceanographic
254 noise related to the major current systems and the residual geoid noise which is generally
255 significantly smaller. As the Jason measurements accumulate with time, the RMS decreases
256 from 2.8 μrad for sub-cycle 1 of Jason 1 to 2.4 μrad for Jason 2 cycle 1 to 2.25 μrad for J-2
257 full GM. (1 μrad of surface slope is 1 mm change in sea surface height per 1 km of horizontal
258 distance.)




Figure 2. Median absolute of along-track sea surface slope differences with respect to the SSV28.1 vertical deflection model derived from a multi-satellite altimeter dataset. Upper left: J1 sub-cycle 1, Upper right: Jason 1 sub-cycle 2; Center left: J2 LRO Cycle 1, Center right: Jason 2 LRO cycle 2; Lower left: J1 Entire LRO; Lower right: Jason 2 Entire GM (Both LRO cycles)

The derived marine gravity grids at 1'x1' resolution were spline interpolated to the location of the marine gravity observations and the standard deviation of the differences are shown in Table 2. This table also highlights the difference in the standard deviation for shallow water regions close to the coast and for deep water regions.

	All depth	< 50 meters	>2000 meters
No. Of Observation	1409700	122108	900969
J1 Sub-cycle 1	5.36	5.25	5.13
J1 Sub-cycle 2	5.53	5.95	5.37
J1 Full GM	4.66	5.14	4.34
J2 LRO cycle 1	4.83	5.40	4.43
J2 LRO cycle 2	4.92	5.55	4.66
J2 Full GM	4.08	4.21	3.72

270 **Table 2.** Comparison with marine gravity data shown in Figure S2. The values in mGal are
271 shown for all depth and for shallow water (less than 50 meters) and for deep water (greater
272 than 2000 meters).

273 It is clear how the length of the GM directly affects the accuracy with which gravity can be
274 derived from the Jason observations. The shortest GM corresponding to the first 179-day
275 cycle of J1 shows a STD of 5.35 mGal with marine gravity observations. Gravity from the
276 first J2 371-day LRO compares at 4.83 mGal and the 406-day GM of Jason-1 compares at
277 4.66 mGal. Gravity from the full J2 GM corresponding to 742 days compares at 4.08 mGal.
278 The latter is nearly 20% better than gravity from one 371-day LRO cycle of Jason-2. The
279 two interleaved LRO cycles of Jason-2 were only efficiently operating for 640 days, as the
280 Jason-2 GM suffered from nearly 100 days of safe-hold despite mission rewind. This
281 indicates that the comparison could have been significant better had the two LRO cycles
282 been completed.

283 Safe-holds degrade the derived marine gravity. Comparing the Jason-2 1st and 2nd LRO
284 cycles which encountered 39 and 60 days (40 days safe-hold plus 20 days early mission
285 termination) exhibit 4.83 vs 4.92 mGal respectively. The numbers are also inferior to gravity
286 derived from Jason-1 GM at 4.66 mGal. However, this GM lasted 30 days longer and only
287 had 18 days of safe-holds. The impact is even larger for particularly coastal regions as also
288 indicated in Table 2. Safe-hold degradation becomes more significant when comparing the
289 Jason-1 first and second sub-cycle where the numbers are 5.36 and 5.53 mGal, respectively.
290 The 18 days safe-hold corresponds to 10% of the time of the second cycle but resulted in a
291 degradation of roughly 5% overall, with degradation in coastal regions of more than 10%
292 (from 5.25 to 5.95 mGal).

293 When the second sub-cycle of Jason-1 was completed the satellite naturally transferred into
294 a subsequent 3rd sub-cycle repeating the same ground track pattern along shifted tracks.
295 The question arises if it would be better to design future GM to “rewind” the mission to
296 remedy any significant safe-hold or to continue with the subsequent sub-cycle.

297 This was examined by adding data from the 3rd sub-cycle to the “safe-hold” affected 2nd sub-
298 cycle of Jason-1. Adding 20 or even 50 days achieved an accuracy of 5.47 and 5.40 mGal,
299 which is still inferior to the comparison from the first sub-cycle. However, the result after 50
300 days (nearly 1/3 of a sub-cycle) approaches the same accuracy as could have been
301 achieved by a “mission rewind”. Unfortunately Jason-1 ceased operating at this stage. A
302 possible “mission rewind” is even more important in the coastal zone. Here the degradation
303 from 5.25 to 5.95 mGal for the second cycle only improves to 5.82 and 5.77 adding altimetry
304 from 20 and 50 additional days from sub-cycle 3. This is somewhat expected as shorter
305 wavelengths will dominate more in the shallow coast zone. This stresses the importance of
306 seriously considering “rewinding missions” in case of significant safe-holds for future GM.

307 In case the graveyard orbit of Jason-2 could be used for a future Jason-3 LRO in a way that
308 avoids collision risk, we explore the idea of moving Jason-3 into interleaved tracks with
309 Jason-2 and bisecting the already 2-years or 4 km Jason-2 GM creating a 2 km grid after
310 only 2 years of Jason-3 GM. This approach would re-use and build on the existing 2 years
311 of Jason-2 GM rather than starting over with a new ground track pattern for the Jason-3 GM.

312 We created a grid from the first 371-day cycle for Jason-2 (having data for 332 days) and
313 the first two 179-days sub-cycles of Jason-1 (totally 340 days) to directly compare the effect
314 of a 2-years systematically densified GM versus two separate 1-year un-coordinated GM
315 affected by the moiré patterns (Dibarboure et al, 2012). The investigation showed that the
316 standard deviation increases from 4.08 mGal for the 2-year densified mission to 4.20 mGal
317 for 2 years of un-coordinated GM. For coastal regions the numbers increase significantly
318 more from 4.21 mGal to 4.50 mGal. The difference might appear small but it is important
319 and it should be noted that Jason-2 suffered from significant safe-hold problems during the
320 second cycle. Hence the gain from densifying an existing GM will be significantly larger than
321 starting all over with a new GM in a different orbit.

322

323 5 Summary and recommendations

324 The GM carried out as the EoL mission for Jason-2 was the first systematic attempt to
325 provide satellite ground-tracks with a systematic track distance of 4 km after 2 years (and
326 planned 2 km after 4 years). The track distance is a limiting factor to the derived global
327 altimetric gravity fields.

328 Starting out with data from the first 179-day cycle of J1 (track distance of 17 km) we found
329 a standard deviation of 5.35 mGal with marine gravity observations. Gravity from the first
330 371-days LRO of Jason-2 (track distance of 8.5 km) compared at 4.83 mGal and the 406-
331 day GM of Jason-1 compares at 4.66 mGal (track distance of 7.8 km). Gravity from the full
332 Jason-2 GM corresponding to 742 days and a track density of 4.3 km clearly compared
333 favorable at 4.08 mGal demonstrating the value of gradually decreasing the track-distance
334 using multiple LRO for the GM. The result was obtained despite the fact that Jason-2 was
335 only efficiently measuring for 642 days of the planned two LRO cycles lasting 742 days.

336 During its GM Jason-2 suffered from significant safe-holds. The most noticeable was a 100-
337 day safe-hold in early 2019. The partnership between NOAA, NASA and CNES agreed to
338 conduct an orbital maneuver and “rewind” the mission by 79 days to recover the missing
339 observations and limit the safe-hold gap to 21 days. This was the first time such was
340 attempted for a GM and stresses the importance of an LRO orbit design having multiple
341 interleaved sub-cycles.

342 Investigating the two 179-day sub-cycles of Jason-1 GM (one was nearly complete and one
343 suffered 18 days or 10% data loss) showed that it very important to consider recovering data
344 from significant safe-holds for future GM mission rather than just continuing the GM into the
345 next interleaved cycle. Jason-3 will be the next satellite to be considered for a GM as its
346 successor Sentinel-6/Jason-CS is scheduled for launch on 10 November 2020.

347 Rewinding the GM to recover mission tracks is particularly important as global marine gravity
348 continues to increase in accuracy with more and more GM data becoming available and
349 integrated with the Jason altimetry (e.g. from the uncontrolled GM of SARAL/AltiKa).

350 Considering minimizing the effect of significant safe-holds is equally important for Mean Sea
351 Surface determination paramount to deriving accurate sea level anomalies. Here the GM
352 data governs the accuracy of the fine scales of the MSS. This is particularly important for

353 future high-resolution altimetric mission like the NASA/CNES Surface Water and Ocean
354 topography (SWOT) to be launched in 2022.

355 In case collision risk between Jason-2 and Jason-3 in the same graveyard orbit could be
356 assessed and found to be controlled we explored the idea of moving Jason-3 into tracks
357 interleaved with Jason-2 and bisecting the already 2-years or 4 km Jason-2 GM creating a
358 2 km grid after only 2 years of GM. This would enable a global gravity field and more
359 importantly a global MSS with unprecedented resolution in time for the SWOT mission. If
360 technically possible our findings strongly recommend to reuse the Jason-2 LRO orbit with
361 Jason-3 to bisect and densify the geodetic grid in a regular way as opposed to a new GM
362 orbit where the grids will not be aligned and therefore will have Moiré patterns.

363

364 **Acknowledgement.** The authors are thankful to the space agencies for considering the Geodetic
365 mission as part of the Extension of Life and for providing these data. We are also thankful to Jim
366 Beale at the National Geospatial-intelligence Agency for sharing the marine gravity for this
367 investigation. All altimeter Level-1 data are available from AVISO data archive
368 (aviso.altimetry.fr/en/data.html). The 1-minute derived marine gravity fields using various
369 combination of the Jason GM is available from <http://dx.doi.org/10.11583/DTU.12865505> and from
370 <http://ftp.space.dtu.dk/pub/Altimetry/JASON-GRAVITY-GRL2020> The scientific results and
371 conclusions, as well as any views or opinions expressed herein, are those of the authors and do not
372 necessarily reflect those of NOAA or the Department of Commerce.

373

374 References

375

376 Andersen, O. B., Knudsen, P., Kenyon, S., Factor, J. K., & Holmes, S. (2017) Global gravity field
377 from recent satellites (DTU15) - Arctic improvements. *First Break*, 35 (12), 37-40,
378 <http://dx.doi.org/10.3997/1365-2397.2017022>

379

380 Andersen, O. B., Knudsen, P., & Stenseng, L. (2016), The DTU13 MSS (Mean Sea Surface) and
381 MDT (Mean Dynamic Topography) from 20 Years of Satellite Altimetry. *IAG Symposia*, 1-10,
382 Springer Verlag, Heidelberg.

383

384 Bronner E., & Dibarboure G. (2012), Technical Note about the J1 Geodetic Mission, [https://podaac-](https://podaac-tools.jpl.nasa.gov/drive/files/allData/jason1/L2/docs/Technical_Note_J1_Geodetic_Mission.pdf)
385 [tools.jpl.nasa.gov/drive/files/allData/jason1/L2/docs/Technical_Note_J1_Geodetic_Mission.pdf](https://podaac-tools.jpl.nasa.gov/drive/files/allData/jason1/L2/docs/Technical_Note_J1_Geodetic_Mission.pdf)

386

387 Dibarboure, G., & R. Morrow (2016), Value of the Jason-1 geodetic phase to study rapid oceanic
388 changes and importance for defining a Jason-2 geodetic orbit. *J. Atmos. Oceanic Technol.* 33 (9):
389 1913–1930. <https://doi.org/10.1175/JTECH-D-16-0015.1>

390

391 Dibarboure, G., [P. Schaeffer](#), [P. Escudier](#), [M.-I. Pujol](#), [J. F. Legeais](#) et al. (2012), Finding
392 desirable orbit options for the “extension of life” phase of Jason-1. *Marine Geodesy*, 35(sup1), 363-
393 399.

394

395 Dufau, C., Orstynowicz, M., Dibarboure, G., Morrow, R., & Le-Traon, P.-Y. (2016), Mesoscale
396 Resolution Capability of altimetry: present & future, *J. Geophys. Res.*, 121, 4910–4927, 2016.

397

398 Pavlis, N. K., Holmes, S. A., Kenyon, S. C., & Factor, J. K. (2012), The development and
399 evaluation of the Earth Gravitational Model 2008 (EGM2008), *J. Geophys Res.*, D. 117, B4,

400 <https://doi.org/10.1029/2011JB008916>

401

402 Pujol, M.-I., Schaeffer, P., Faugère, Y., Raynal, M., Dibarboure, G., & Picot, N. (2018), Gauging the
403 improvement of recent mean sea surface models: A new approach for identifying and quantifying
404 their errors. *J. Geophys. Res.*, C, 123, 5889–5911. <https://doi.org/10.1029/2017JC013503>

405

406 Sandwell, D. T., Dietmar-Müller R., Smith W., Garcia, E. & Francis, R. (2014), New global marine
407 gravity model from CryoSat-2 and Jason-1 reveals buried tectonic structure. *Science*. 346 no. 6205
408 pp. 65-67

409

410 Sandwell D., Harper H., Tozer, B., & Smith W., (2019), Gravity field recovery from geodetic
411 altimeter missions. *Adv in Space Res*, <https://doi.org/10.1016/j.asr.2019.09.011>.