

# Spatial and temporal variability of seasonal rainfall and mean temperature over different region of Bangladesh

Mohammad Sarker<sup>1,1</sup>

<sup>1</sup>The Higher Ground Foundation

November 30, 2022

## Abstract

Bangladesh is a small country of South Asia which is considered as one of the most vulnerable countries in the world to climate change and it is affected by severe weather and climate events. In this paper, the climatic change and variability over Bangladesh has been studied. The time series analysis was applied to investigate the variability and trend over three zone [(Northern (North West & North East), Middle and Southern (Coastal, Island, Hilly)] of Bangladesh. The study was based upon the regional differences of climatic trends for the country and tested it for each region against other regions via a t-test. Result suggest that as a topical country, there is small temperature variation in Bangladesh. But there is regional variation between northwest and south zone. In winter there exists a north-south temperature gradient which reverses during summer (pre-monsoon and monsoon). There is an extreme large seasonal variation from winter- to summer- monsoon in Bangladesh. In all seasons the difference of maximum and minimum temperature is higher in the north zone than in the south zone. A remarkable correlation was discovered between this temperature range and the rainfall occurrence. All seasons the mean maximum temperature is increasing except in winter for the northwest and middle zone ( $-0.004$  °C/year and  $-0.0069$  °C/year). Hilly region is showing highest increasing rates of mean maximum Temperature in all season ( $+0.0754$  °C/year in winter,  $+0.0635$  °C/year in Pre monsoon,  $+0.0554$  °C/year in monsoon,  $+0.0679$  °C/year in Post monsoon); Where as in all season mean minimum temperature is decreasing only hilly region. Overall the temperature is increasing in whole of the country. We also find a positive trend in the rainfall, especially large during the summer monsoon and in the southern zone (in Hilly region, increasing rate of rainfall  $+17.784$  mm/year). The northwest area on the contrary is characterized by decreasing rainfall values both in winter and summer. So we can conclude that the northwest is clearly becoming more arid. May be this is enhanced by the large scale deforestation processes going on.

## **Spatial and temporal variability of seasonal rainfall and mean temperature over different region of Bangladesh**

Mohammad Shohrab Hossain Sarker

The Higher Ground Foundation, London, UK

International Center for Climate Change and Development, Dhaka, Bangladesh

Corresponding to author: msh.sarker@yahoo.com

### **Key findings:**

- (i) The seasonal variation for both maximum and minimum temperature is small. In winter there exists a north-south temperature gradient which reverses during summer (pre-monsoon and monsoon).
- (ii) The seasonal variation for the rainfall is extremely large. In winter there exists a south-north precipitation gradient which reverses during the other seasons.
- (iii) Overall we found a positive trend in the rainfall, especially large during the summer monsoon and in the southern zone.
- (iv) The northwest area is characterized by decreasing rainfall values both in winter and summer.
- (v) Trend in the south zone (Coastal) is higher than other zone of the country.
- (vi) The regional analysis identified two regions in Bangladesh, which behave differently, i.e. the Hilly region in the southeast and the northwest area.
- (vii) Difference between mean maximum temperature and mean minimum temperature ( $\Delta T$ ) is increasing in the Southern Zone.
- (viii) Overall the temperature of whole of the country is increasing (the rate of max temp is  $0.028^{\circ}/\text{year}$ ) concurrent with the global temperature increase.

## 26    **Abstract**

27    Bangladesh is a small country of South Asia which is considered as one of the most vulnerable  
28    countries in the world to climate change and it is affected by severe weather and climate events. In this  
29    paper, the climatic change and variability over Bangladesh has been studied. The time series analysis  
30    was applied to investigate the variability and trend over three zone [(Northern (North West & North  
31    East), Middle and Southern (Coastal, Island, Hilly)] of Bangladesh. The study was based upon the  
32    regional differences of climatic trends for the country and tested it for each region against other regions  
33    via a t-test. Result suggest that as a topical country, there is small temperature variation in Bangladesh.  
34    But there is regional variation between northwest and south zone. In winter there exists a north-south  
35    temperature gradient which reverses during summer (pre-monsoon and monsoon). There is an extreme  
36    large seasonal variation from winter- to summer- monsoon in Bangladesh. In all seasons the difference  
37    of maximum and minimum temperature is higher in the north zone than in the south zone. A  
38    remarkable correlation was discovered between this temperature range and the rainfall occurrence. All  
39    seasons the mean maximum temperature is increasing except in winter for the northwest and middle  
40    zone ( $-0.004$  °C/year and  $-0.0069$  °C/year). Hilly region is showing highest increasing rates of mean  
41    maximum Temperature in all season ( $+0.0754$  °C/ year in winter,  $+0.0635$  °C/year in Pre monsoon,  
42     $+0.0554$  °C/ year in monsoon,  $+0.0679$  °C/year in Post monsoon); Where as in all season mean  
43    minimum temperature is decreasing only hilly region. Overall the temperature is increasing in whole of  
44    the country. We also find a positive trend in the rainfall, especially large during the summer monsoon  
45    and in the southern zone (in Hilly region, increasing rate of rainfall  $+17.784$  mm/year). The northwest  
46    area on the contrary is characterized by decreasing rainfall values both in winter and summer. So we  
47    can conclude that the northwest is clearly becoming more arid. May be this is enhanced by the large  
48    scale deforestation processes going on.

49    [Key words: Pre monsoon, Summer monsoon, Post monsoon, Maximum temperature, Minimum  
50    temperature, Precipitation]

|    |   |    |
|----|---|----|
| 51 | <b>Table of Contents</b>  |    |
| 52 | 1. Introduction: .....  | 3  |
| 53 | 2. Location and physiographic setting of Bangladesh.....                      | 6  |
| 54 | 3. Methodology .....  | 7  |
| 55 | 3.1 Data process .....  | 7  |
| 56 | 3.2 Physio-climate of classified zone ( <i>as noted Sarker, 2009</i> ) .....  | 8  |
| 57 | 3.3 Classification of Meteorological seasons:.....                            | 10 |
| 58 | 3.4 Statistical analysis .....  | 10 |
| 59 | 4. Climate of Bangladesh .....  | 10 |
| 60 | 5. Result and Discussion .....  | 15 |
| 61 | 5.1 Seasonal variation.....   | 15 |
| 62 | 5.1.1 Seasonal variation of Temperature:.....                                 | 15 |
| 63 | 5.1.2 Seasonal variation of Rainfall.....                                     | 19 |
| 64 | 5.2 Relation between Maximum-Minimum Temperature Difference and Rainfall..... | 20 |
| 65 | 5.3 Time Series Analysis .....  | 22 |
| 66 | 5.3.1 Trend of Temperature.....   | 22 |
| 67 | 5.3.2 Trend of Precipitation.....   | 33 |
| 68 | 5.4 Regional differences .....  | 35 |
| 69 | 6. Conclusion:.....   | 36 |
| 70 | Acknowledgements.....   | 38 |
| 71 |   |    |
| 72 |   |    |

## 73      **1. Introduction:**

74      Climate change is the one of the major alarm for the earth. “While most climate scientist agree that  
75      global temperature are rising and will continue to rise with so-called greenhouse warming there is  
76      much less agreement on how such changes will affect particular region of the globe” (Brammer, 2004).  
77      IPCC AR5 report shows a linear trend of 0.85°C from 1880 to 2012 (IPCC, 2013). The total increase  
78      between the average of the 1850–1900 period and the 2003–2012 period is 0.78 [0.72 to 0.85] °C,  
79      based on the single longest dataset available (IPCC, 2013). The IPCC Fourth Assessment Report for  
80      South Asia shows that warming is likely to be above the global mean (IPCC, 2014). Rainfall in the  
81      summer season is likely to increase and it is very likely that there will be an increase in the frequency  
82      of intense precipitation events in parts of South Asia (IPCC, 2014). MMD-IPCC FAR Scenario A1B

83 models show a median increase of 3.3°C in annual mean temperature and 11% precipitation median  
84 increase by the end of the 21<sup>st</sup> century in South Asia (Christensen *et al.*, 2007). The IPCC Special  
85 Report on the Regional Impacts of Climate Change (2007) indicated that Bangladesh will experience  
86 5–6 % increase in precipitation by 2030 (Shahid, 2011).

87 The climate of Bangladesh is dominated by the Southwest monsoon. Southwest monsoon season shows  
88 considerable year-to-year rainfall variability over South Asia (Asian Development Bank, 2014).  
89 Current climatic projection suggest that temperature could increase by 1.1-1.25°C in the monsoon and  
90 by 1.25-1.5 °C in the dry season and monsoon rainfall could increase by 10-15 percent by 2030 and  
91 dry season rainfall by 5-10 percent in and around Bangladesh (Brammer, 2000, 2004). Such a changes  
92 could increase the frequent and severity of floods and keep some basin land wet longer into dry season  
93 than occur at present (Brammer, 2000, 2004). Changes to precipitation and temperature are expected to  
94 result in severe droughts and strong flooding in different parts of Bangladesh. It could be made more  
95 severe as the Himalaya's glaciers are melting; they feed seven rivers in this region, among which are  
96 the two big rivers Ganga and Brahmaputra, which lead into the Bay of Bengal through Bangladesh.  
97 Year to year variability of summer monsoon season influences agricultural activities, river drainage  
98 systems and hydrological behaviour all of which, in turn, influences the life and economy of the  
99 country (Johnson, 1982; Ahmed and Karmakar, 1993a; Ahmed and Kim, 2003). According to the IPCC  
100 fifth assessment report, the loss of life and property associated with floods is highest in India and  
101 Bangladesh (IPCC, 2014). Thus, it is important to describe and distinguish the regional and seasonal  
102 climatic variability and climate change of Bangladesh. Therefore, this research aim to assess the  
103 present status of several climate changes. At the same time comprehensive environmental management  
104 plans can be made by using the data and results (Sarker, 2009).

105 There were a high number of research had been conducted on climatic event in Bangladesh. The  
106 variability of annual rainfall (Shamsuddin and Ahmed, 1974), extreme climatic even (Ahmed, 1989),  
107 variability of arrival and withdrawal date of the summer monsoon period (Ahmed and Karmakar,

108 1993a; Ahmed and Kim, 2003), temperature and rainfall trend (Ericksen, Ahmad and Chowdhury,  
109 1997; Karmakar, 2000), intra-annual and inter-annual variation of rainfall over different regions of  
110 Bangladesh (Debsarma, 2003), were vital work. A comparison of study of three individual  
111 metrological station such as Rajshahi (Northwest), Cox.sBazar (Coastal) and Sylhet (Northeast)  
112 conducted by Ahmed and Kim, 2004). They mentioned frequency of consecutive rain day of various  
113 duration. They found, in wettest monsoon season, Northwest (Rajshahi) region have fewer consecutive  
114 rain day 8-10 days than coastal 25-30 days (Cox's Bazar). Whereas, during driest monsoon season, the  
115 consecutive rainfall day range from 3-6 days in Northwest (Rajshahi) and 5-7 in coastal (Cox's Bazar).  
116 Asian Development Bank, (1994) suggested that rainfall for the nation as whole has slightly increased  
117 over time between 1951 and 1991 (the rate of increase is 0.19% per year) (Asian Development Bank,  
118 1994). Mirza *et al.*, (1998) showed that precipitation over all the GBM basins of Bangladesh have been  
119 increasing over a period of 31 years. Lutz *et al.*, (2016) examined extreme event consider Ganges and  
120 Brahmaputra river basins, (that include Bangladesh as a study area as well), using mean air  
121 temperature, precipitation and ETCCDI indices between current scenario (1971–2000) and future  
122 (2071-2100). This indices includes R99pTOT [precipitation due to extremely wet days (>99th  
123 percentile)], Consecutive dry days (CDD), Warm spell duration index (WSDI), Cold spell duration  
124 index (CSDI). Similarly (Shahid, 2011) investigated trend in extreme rainfall using index with more  
125 indicator. He found CDD decreasing in Bangladesh. There are few studies projected temperature  
126 (Ramamasy and Baas, 2007; Immerzeel, 2008; Gosling *et al.*, 2011) and precipitation (May, 2004;  
127 Immerzeel, 2008; Gosling *et al.*, 2011) increases in future.

128 Before this study, none of the research has been considered to -combined temporal-spatial and variation  
129 of temperature and rainfall at seasonal and annual timescale over different regions or for the entire  
130 country. However, there are few studies examine spatiotemporal variation after this studies. Definitely,  
131 research to date, no studies have been investigated spatial and temporal variation of delta T (difference  
132 between maximum and minimum temperature) for at seasonal and annual timescale over different

133 regions and entire Bangladesh. In addition, very little focus given on seasonal timescale. Some studies  
134 focus on individual station rather regional basis. In addition, motive for spatial and at seasonal  
135 timescale variation poorly described. Furthermore, there is no previous study of comparison between  
136 different regions. Given the lack of work, a range of schemes such as spatial and temporal variation at  
137 seasonal and annual timescale of different region and tested the trends for each region against the  
138 trends for the other regions via a t-test has been explored in this study.

## 139 **2. Location and physiographic setting of Bangladesh**

140 Bangladesh is a small country of 145,452 km<sup>2</sup>, extending from 20°45' N to 26°40' N, and from 88°05'  
141 E to 92°40' E (Salahuddin *et al.*, 2006). About 80% of the land surface is plain and most of the country  
142 lies below 10 meters elevation (Sarker, 2015) (see fig 1). It is bordered by the Assam Hills in the east,  
143 by the Meghalaya plateau in the north, with the lofty Himalayas lying beyond (Ahmed, 2003; Ahmed  
144 and Kim, 2003; Sarker, 2015). (Figure 1)

145




Figure 1 : Elevation map of Bangladesh. (Sources author PhD Thesis)

### 3. Methodology

#### 3.1 Data source and data processing

Series of 23 years (1981-2003) monthly mean maximum temperature and minimum temperature and rainfall data for 23 stations were obtained from Bangladesh Metrological Department (BMD). There were a few missing data in the early eighties. This was overcome by filling in average values obtained by linear interpolation technique. Total missing data 1.88%. Value of each stations transformed to seasonal (e.g. winter) by calculating -average for both temperature (Tmax and Tmin) –summation for rainfall. Similarly seasonal value converted to regional value by calculating average value of selected stations for a region (as an example, average value of Rangpur, Dinajpur, Rajshahi, Bogra stations converted to Northwest region). Apart from the study of Tmax and Tmin the Temperature range  $\Delta T$ ,



159 defined as Tmax-Tmin, was also analyzed

160 **3.2 Physio-climate of classified zone (as noted Sarker, 2009):**

161  
162 The country is split into three zone and six regions determined by the physiography of the country and  
163 its response to the general climatic patterns discussed below. (See fig 2, Table1).

164 **Northern Zone:**

165 **North West Region:** It is located near the foothills of the Himalayas and The contiguous plain of west  
166 Bengal and the Gangetic plain lies to its west. Relatively dry, shortest duration of the summer  
167 monsoon, least number of rain days and least amount of seasonal rainfall Temperature is higher in pre  
168 monsoon and coolest) in winter.

169 **North East Region:** It is located near the foothills of the Meghalaya Plateau (windward side) which  
170 add an orographic effect on the monsoon rainfall.

171 **Middle Zone:** It is central part of the country and consists of flat plains. High temperature and heavy  
172 rainfall is Characteristics of the zone.

173 **Southern Zone:** it is located near Bay of Bengal. This zone (southeast) is the entry and exit point of  
174 the Bay of Bengal Branch of the South Asian Monsoon (Spate and Learmonth, 1984; Ahmed and  
175 Karmakar, 1993b; Ahmed and Kim, 2003). Relatively wet, longest duration of summer monsoon. High  
176 humidity, heavy rainfall and less temperature variation between day and night; are characteristics of  
177 Southern zone.




Figure 2: The Study area with 23 metrological station

Table 1: Distribution Meteorological stations in study area:

| 3 Zones | 6 Regions  | 23 Meteorological stations                                      | Number of Station Based on Zone |
|---------|------------|---|---------------------------------|
| North   | North West | Rangpur, Dinajpur, Rajshahi, Bogra                              | 7                               |
|         | North East | Mymensingh, Sylhet, Srimongal.                                  |                                 |
| Middle  | Middle     | Jessore, Chuadanga, Faridpur, Dhaka, Comilla, Tangail.          | 6                               |
| South   | Coastal    | Khulna, Satkhira, Patuakhali, Barisal, Chittagong, Cox's Bazar. | 10                              |
|         | Island     | Bhola, Hatiya, Sandwip  |                                 |
|         | Hilly      | Rangamati.  |                                 |

### 3.3 Classification of Meteorological seasons:

Bangladesh is characterized by three distinct meteorological seasons with a relatively cool, sunny and dry season from November through February, a pre-monsoon hot season from March through May, the rainy summer monsoon (Rainy season) from June through mid-October and a short and mild autumn (Post-monsoon) season from mid-October to mid-November (Asian Development Bank, 1994; Ahmed, 2003, 2004; Rasheed, 2008; as noted by Sarker, 2009). Nevertheless, four seasons were recognised by (Brammer, 2000, 2004).

### 3.4 Statistical analysis

Using the familiar simple formula  $Y=mx+b$ , linear regression analysis was carried out with time (year) as an independent variable. The slope of the regression line indicates a per year increase or decrease in rainfall and temperature. The regional differences of climatic trends (mean TMAX and TMIN) for the country were tested for each region against other regions conducted via a t-test. The statistical test will provide a p-value that bounded 0 and 1. The critical value ( $\alpha$ ) is tested at 95% (0.05) confidence level. Finally, in this research, the seasonal mean maximum and minimum temperature and rainfall variation and trends are cartographically represented by iso-lines on maps, created by Arc View software.

## 4. Climate of Bangladesh

The climate of Bangladesh is primarily influenced by the tropical summer and winter circulation, and partly by pre-monsoon and post-monsoon circulation. The southwest summer monsoon originates over the Indian Ocean and carries warm, moist, and unstable air. The winter monsoon originates over northwest India and carries cool and dry air. The seasonal variability of the climate of Bangladesh for the period 1981-2003 is given below.

In winter, two branches of the jet stream are anchored by the Himalayas, one to the north and the other to the south (Figure 3. (a)). The southern branch of the jet stream causes subsidence in the atmosphere in the north-western part of India and consequently, a centre of high pressure develops there, while low pressure develops over the relatively warmer Indian Ocean (Ahmed, 2004). As a result of this pressure

distribution, winds flow from the interior of India toward the Indian Ocean, in what is known as the winter monsoon circulation (Sarker, 2015). Since the wind in this season is characterized by the land-to-ocean flow, nearly dry conditions prevail during this season. January is the coldest month in Bangladesh. In this study it was found that mean maximum temperature in January varies from about 23.20 °C (Rangpur) to 26.88°C ( Cox's bazaar) where is The mean minimum temperature in from 15.23 °C (Cox's bazaar) to 9.93°C (Srimongal) (Table 2). This season is very dry due to the effect of the wind flow from north India to the ocean. The average winter rainfall in Bangladesh is 48.86 mm (Table 2) or only 2% of the yearly total. **The pre-monsoon** season is the transition period when the northerly or north-westerly winds of the winter season gradually change to the southerly or southwesterly winds of the summer monsoon (Ahmed, 2003). During the early pre-monsoon season, a narrow zone of air-mass discontinuity lies over Bangladesh, which extends from the southwest corner to the northeast corner of the country (Ahmed, 2004). This line of discontinuity separates the warm moist air coming in from the Bay of Bengal and the hot dry air flowing from the interior of northern India. (Huq, 1974; Ahmed and Karmakar, 1993; Ahmed and Kim, 2003; Ahmed, 2004), resulting in high temperatures and the occurrence of thunderstorms and tornadoes (Sarker, 2015). April is the hottest month. The mean maximum temperature in April varies from about 30.04°C (Sandwip) to 35.92°C (Rajshahi) (Table 2) and mean minimum temperature varies from about 20.82 °C (Sylhet) to 24.49°C ( Hatiyal) (Table 2). Rainfall in the pre-monsoon hot season (March-May) accounts for 21% of the total annual rainfall. The rain during this period is caused by convective storms (thunderstorms) or nor`western (locally called *Kalbaishakhi*) (Ahmed, 2003). The average total pre-monsoon rainfall in Bangladesh equals 526.35 mm. It varies from about 212.86 mm (Hatiya) to 1281.13 mm (Sandwip)

#### ***In summer monsoon***

In the summer monsoon season a centre of low pressure develops over the western part of India because of the intense heating of the landmass, while high pressure develops over the relatively cooler Indian Ocean, and the Inter-Tropical Convergence Zone (ITCZ) of the North Indian Ocean moves to

the north (Ahmed, 2004) (Fig 3b). This flow pattern is known as the summer monsoon circulation..

Also the Equatorial low Pressure or Inter Tropical Convergence Zone (ITCZ) of North Indian Ocean progresses to the North. As the trough associated with the ITCZ moves to about 25°N over India at the height of the summer, the release of latent heat in the upper troposphere as well as the heating of the Tibetan Plateau causes a north south temperature and pressure gradient resulting in the development of an easterly jet stream at a height of about 150 hPa (15km) (Robinson et al., 1999). As the trough associated with the ITCZ moves to about 25°N over India at the height of the summer, the release of latent heat in the upper troposphere, as well as the heating of the Tibetan Plateau, causes a north-south temperature and pressure gradient and the development of an easterly jet stream at about 150 hPa (Robinson and Henderson-Seller, 1999). The summer monsoon flow enters Bangladesh in late May or early June, and continues to flow toward the centre of low pressure that lies over the western-central part of India. The temperature during this season is slightly lower than during the pre-monsoon season. June is the hottest month of this season. Heavy and prolonged rainfall characterizes the season, caused by tropical depressions that enter the country from the Bay of Bengal. June is the hottest month of this season. In June the mean maximum temperature varies from about 30.50°C (Sandwip) to 33.90°C (Chuadanga) (Table 2) and mean minimum temperature varies from about 24.45 °C (Sylhet) to 26.27°C (Satkhira) (Table 2). Heavy and prolonged rainfall characterize the season, caused by tropical depressions that enter the country from the Bay of Bengal. During this season the rainfall in the northwest and middle region is less than in other areas. The average total rainfall varies from about 750.26 mm (Rajshahi) to 2593.41 mm (Coxs' Bazar) (Table 2).

The post-monsoon (autumn) represents a transitional period (Ahmed, 2004) similar to the pre-monsoon. The temperature remains mild during this season and it is more comfortable due to less moisture in the air. The mean maximum temperature varies from 30.14°C (Rangpur) to 31.78°C (Jessore) and the mean minimum temperature varies from 21.30°C (Dinajpur) to 25.46°C (Hatiya) (Table 2). This season is comfortable as there is less moisture in the air. The average total post

monsoon rainfall in Bangladesh is 538.95 mm (21.8%) and it varies from 331.38 mm (Sandwip) to 806.78 mm (Sylhet) (Table 2)




Figure 3: The airflow patterns over southern Southern Asia in (a) winter and (b) summer (Sources author PhD Thesis)

**Table 2. Mean Minimum, Maximum Temperature and Rainfall for the period 1981-2003**

| Zones      | Regions   | Stations    | Min Temp (°C) | Max Temp (°C) | Rainfall (mm) | Min Temp (°C) | Max Temp (°C) | Rainfall (mm) | Min Temp (°C) | Max Temp (°C) | Rainfall (mm) | Min Temp (°C) | Max Temp (°C) | Rainfall (mm) | Min Temp (°C) | Max Temp (°C) | Rainfall (mm) |
|------------|-----------|-------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|---------------|
| North      | Northwest | Rangpur     | 12.12         | 24.94         | 65            | 20.32         | 31.28         | 421.06        | 25.78         | 31.82         | 1346.14       | 21.62         | 30.14         | 587.27        | 19.85         | 29.54         | 2512.57       |
|            |           | Dinajpur    | 11.63         | 24.97         | 60            | 23.23         | 32.1          | 302.24        | 25.53         | 32.17         | 1176.48       | 21.3          | 30.43         | 564.75        | 20.37         | 29.92         | 2079.73       |
|            |           | Rajshahi    | 11.96         | 25.83         | 37.09         | 21.38         | 34.64         | 227.7         | 25.89         | 32.8          | 825.91        | 21.97         | 31.3          | 470.13        | 20.19         | 31.14         | 1560.83       |
|            |           | Bogra       | 13.14         | 26.18         | 44.98         | 21.48         | 32.48         | 212.86        | 26.04         | 32.28         | 1427.63       | 22.33         | 31.32         | 490.9         | 20.64         | 30.56         | 2176.37       |
|            | Northeast | Mymensing   | 13.23         | 25.88         | 38.65         | 21.33         | 31.32         | 526.57        | 25.91         | 31.33         | 1184.21       | 22.3          | 30.67         | 588.43        | 20.58         | 29.8          | 2337.86       |
|            |           | Sylhet      | 13.8          | 26.28         | 59.91         | 20.56         | 30.7          | 1091.97       | 24.84         | 31.48         | 2250.26       | 21.86         | 30.7          | 806.78        | 20.16         | 29.79         | 4208.92       |
|            |           | Srimongal   | 11.46         | 26.48         | 55.81         | 20.48         | 32.09         | 738.11        | 24.98         | 32.19         | 1099.26       | 21.13         | 30.89         | 508.08        | 19.41         | 30.41         | 2401.26       |
| Middle     | Middle    | Jessore     | 12.88         | 26.9          | 54.38         | 22.58         | 34.62         | 318.44        | 25.94         | 32.77         | 917.21        | 22.09         | 31.78         | 404.47        | 21.42         | 31.52         | 489.19        |
|            |           | Chuadanga   | 12.43         | 26.27         | 46.55         | 22.39         | 34.91         | 229.53        | 26.15         | 33.2          | 754.13        | 22.37         | 31.66         | 446.85        | 20.73         | 31.51         | 530.76        |
|            |           | Faridpur    | 13.68         | 26            | 49.95         | 22.2          | 33.32         | 451.38        | 25.84         | 31.77         | 945           | 22.99         | 30.79         | 451.04        | 21.06         | 30.47         | 533.35        |
|            |           | Dhaka       | 14.42         | 26.6          | 44.64         | 22.87         | 33.01         | 537.39        | 26.24         | 31.96         | 987.37        | 22.95         | 31.15         | 522.34        | 21.53         | 30.68         | 605.7         |
|            |           | Tangail     | 13.46         | 25.64         | 49.24         | 21.98         | 32.67         | 464.06        | 25.39         | 32.01         | 885.94        | 22.28         | 30.91         | 461.56        | 20.68         | 30.31         | 135.86        |
|            |           | Comilla     | 12.94         | 26.48         | 45.81         | 21.6          | 31.76         | 536.46        | 25.94         | 31.47         | 1085.41       | 22.61         | 30.99         | 418.12        | 20.65         | 30.17         | 499.93        |
| South      | Coastal   | Khulna      | 13.58         | 26.88         | 62.98         | 22.99         | 33.93         | 333.51        | 26.08         | 32.29         | 984.7         | 22.99         | 31.4          | 535.31        | 21.92         | 31.13         | 619.76        |
|            |           | Satkhira    | 13.56         | 27.15         | 56.6          | 23.31         | 34.37         | 301.73        | 26.18         | 32.58         | 1246.51       | 22.51         | 31.59         | 455.43        | 21.94         | 31.42         | 540.39        |
|            |           | Patuakhali  | 15.12         | 26.86         | 40.86         | 23.43         | 32.68         | 429.12        | 25.98         | 31.09         | 1636.78       | 23.37         | 30.78         | 588.7         | 22.43         | 30.35         | 672.27        |
|            |           | Barishal    | 14.17         | 26.81         | 43.82         | 23.25         | 32.89         | 403.52        | 25.92         | 31.37         | 1206.98       | 22.93         | 30.97         | 479.7         | 21.48         | 30.51         | 562.66        |
|            |           | Chittagong  | 15.19         | 27.18         | 47.37         | 22.82         | 31.64         | 526.91        | 25.25         | 31.36         | 1859.12       | 23.12         | 31.16         | 503.68        | 21.49         | 30.34         | 586.67        |
|            |           | Cox's Bazar | 16.48         | 27.89         | 45.44         | 23.35         | 32.11         | 462.21        | 25.21         | 30.61         | 2593.41       | 23.46         | 31.15         | 715.31        | 22.04         | 30.44         | 798.94        |
|            | Island    | Bhola       | 14.17         | 26.9          | 46.26         | 23.25         | 32.41         | 414.47        | 25.92         | 31.09         | 1323.32       | 22.93         | 30.95         | 490.91        | 22.06         | 30.34         | 574.25        |
|            |           | Hatiya      | 16.92         | 26.69         | 29.36         | 20.54         | 30.08         | 454.76        | 25.28         | 30.82         | 1898.48       | 25.07         | 30.65         | 684.25        | 21.75         | 29.56         | 766.2         |
|            |           | Sandwip     | 15.95         | 26.14         | 70            | 23.86         | 30.41         | 1281.13       | 25.59         | 30.23         | 1902.26       | 23.56         | 30.32         | 331.38        | 22.15         | 29.28         | 413.13        |
|            | Hilly     | Rangamati   | 14.51         | 26.82         | 46.92         | 21.85         | 32.59         | 564.84        | 24.71         | 31.34         | 1474.58       | 22.57         | 30.86         | 593.43        | 20.8          | 30.4          | 675.49        |
| Bangladesh |           |             | 13.5          | 26            | 40            | 21            | 31.5          | 500           | 24            | 31            | 1400          | 23            | 30.4          | 550           | 20.5          | 29.73         | 630.63        |

## 5. Result and Discussion

### 5.1 Seasonal variation

#### 5.1.1 Seasonal variation of Temperature:

Winter is the coldest and pre-monsoon is the hottest season in this country. All regions of the country show their highest mean maximum temperature in pre-monsoon except the northeast region (31.37°C in pre monsoon and 31.66° C in monsoon) while the highest mean minimum temperature occurs during the summer monsoon (Figure 4 and table 3). This is because the atmospheric moisture content and cloud cover are the highest during the summer monsoon season. At night they absorb the outgoing long wave radiation very efficiently, and thereby they trap the heat at night. As a result, in the summer monsoon, the minimum temperature is slightly higher than expected (Ahmed, 2005; Sarker, 2005). An analogue explanation holds to explain the high maxima in the pre-monsoon dry period.


*In winter* mean maximum temperature is slightly lower in the north zone (25.48 °C in northwest region) and higher in south zone of the country (27.13 °C mean maximum temperature in coastal region). [(Figure 4 & 5(A) Table 3)]. The northwest region is clearly the coolest region of the country. This is certainly due to the advection of cold continental air that enters the country from the northwest and being farthest north, the angle of the sun is also smaller than in any other region in winter. In addition the proximity of the Himalaya's contributed to the relatively lower maximum temperatures than in any other region (Ahmed, 2005).. The mean minimum temperature of the northwest region also shows lower values than in any other region of the country (12.21°C Table 3), because atmospheric moisture content in this region is generally low. Due to the high moisture content of the atmosphere over the Island region and the coast, these areas show highest mean minimum temperature values of 15.68°C and 14.68°C (Table 3).



290 Table 3. Seasonal Mean Minimum, Maximum, Difference of maximum & minimum ( $\Delta T$ ) Temperature and Rainfall (Average total per season)


| Zone       | Region      | Winter        |               |                 |               | Pre monsoon   |               |                 |               | Monsoon       |               |                 |               | Post monsoon  |               |                 |               | Annual        |               |                 |               |
|------------|-------------|---------------|---------------|-----------------|---------------|---------------|---------------|-----------------|---------------|---------------|---------------|-----------------|---------------|---------------|---------------|-----------------|---------------|---------------|---------------|-----------------|---------------|
|            |             | Min Temp (°C) | Max Temp (°C) | $\Delta T$ (°C) | Rainfall (mm) | Min Temp (°C) | Max Temp (°C) | $\Delta T$ (°C) | Rainfall (mm) | Min Temp (°C) | Max Temp (°C) | $\Delta T$ (°C) | Rainfall (mm) | Min Temp (°C) | Max Temp (°C) | $\Delta T$ (°C) | Rainfall (mm) | Min Temp (°C) | Max Temp (°C) | $\Delta T$ (°C) | Rainfall (mm) |
| North      | Northwest   | 12.21         | 25.48         | 13.27           | 51.77         | 21.6          | 32.63         | 11.02           | 290.97        | 25.81         | 32.27         | 6.46            | 984.7         | 21.8          | 30.8          | 8.99            | 528.26        | 20.26         | 30.29         | 10.03           | 1855.69       |
|            | Northeast   | 12.83         | 26.22         | 13.39           | 51.46         | 20.79         | 31.37         | 10.58           | 785.55        | 25.24         | 31.66         | 6.42            | 1511.24       | 21.76         | 30.75         | 8.99            | 634.43        | 20.05         | 30            | 9.95            | 2982.68       |
| Middle     | Middle zone | 13.3          | 26.31         | 13.01           | 48.43         | 22.27         | 33.38         | 11.11           | 422.88        | 25.92         | 32.2          | 6.28            | 943.7         | 22.55         | 31.21         | 8.66            | 450.73        | 21.01         | 30.78         | 9.76            | 1865.73       |
| South      | Coastal     | 14.68         | 27.13         | 12.45           | 49.51         | 23.19         | 32.94         | 9.75            | 409.5         | 25.77         | 31.55         | 5.78            | 1587.92       | 23.07         | 31.18         | 8.11            | 546.36        | 21.88         | 30.7          | 8.81            | 2593.28       |
|            | Islands     | 15.68         | 26.58         | 10.9            | 48.54         | 22.55         | 30.97         | 8.42            | 716.79        | 25.6          | 30.71         | 5.12            | 1708.02       | 23.85         | 30.64         | 6.79            | 502.18        | 21.98         | 29.72         | 7.74            | 2975.53       |
|            | Hilly       | 14.51         | 26.82         | 12.41           | 46.92         | 21.85         | 32.59         | 10.62           | 564.84        | 24.71         | 31.34         | 6.81            | 1474.58       | 22.57         | 30.86         | 7.84            | 593.43        | 20.8          | 30.4          | 9.41            | 2679.77       |
| Bangladesh |             | 13.78         | 26.35         | 12.57           | 49.44         | 21.97         | 32.3          | 10.25           | 531.75        | 25.45         | 31.59         | 6.15            | 1368.36       | 22.64         | 31            | 8.23            | 542.56        | 20.97         | 30.31         | 9.29            | 2492.11       |

291



292

293 Fig. 4. Regional variation of the mean maximum and minimum temperature and Rainfall in different season



294

295

296

297

Fig. 5. The mean maximum Temperature in (A) Winter (B) Pre monsoon; the mean minimum Temperature in (C) Winter (D) Pre monsoon; The average total rainfall in (E) Winter (F) Monsoon

298 ***In the Pre-monsoon*** and monsoon seasons The mean maximum temperature, is highest in the middle  
 299 zone (33.38 °C in figure 4, 5.(B) & Table 3)]. There are several reasons for this reversal in regional  
 300 temperature gradient as compared tot the winter season : (a) during the pre monsoon season, the sun is  
 301 nearly vertical over this zone, (b) coastal regions and islands are slightly cooler due to the maritime  
 302 effect, (c) hilly regions are slightly cooler because of the elevation, and (d) the northern regions are  
 303 slightly cooler because the sun shines at a smaller angle over the northern regions resulting in lesser  
 304 intensity of the sun over these (Ahmed, 2005).

305 Temperature during the ***summer monsoon*** season is generally lower than the temperature during the  
 306 pre-monsoon season. Widespread cloud cover during the monsoon season reduces the insulation which,  
 307 in turn, causes the temperature to be lower than expected for the season (Ahmed, 2005; Sarker, 2005).

308 In the pre-monsoon, the coastal region shows the highest mean minimum temperature (23.19°C) while  
 309 the northern regions show lower mean minimum temperature values of 20.79 °C in (Figure 4, & Table  
 310 3)] in the northeast and 21.6 in the northwest region. This indicates that the mean minimum  
 311 temperature increases from the north region to the coastal areas. In monsoon season mean maximum  
 312 temperature in the northwest region shows the highest value of 32.27°C while the islands area show  
 313 the lowest value of 30.71°C. It indicates that the mean maximum temperature decreases from the  
 314 northwest region to the south zone (Coastal, island and hilly regions). Compared to other regions, the  
 315 northwest region has the least widespread cloud cover during the summer monsoon season. So, under  
 316 less cloud cover, insulation is relatively high which causes the maximum temperature in northwest  
 317 region to be higher than any other region. In this season the middle zone shows higher mean minimum  
 318 temperatures (25.92°C) than the other regions (Figure 4 and Table 3)

319 ***In Post monsoon*** the mean maximum temperature in the middle zone and coastal area is slightly higher  
 320 (31.21°C and 31.18°C in coastal) than in the north zone (30.75 °C in northeast and 30.80°C in  
 321 northwest region) (Figure 4 and Table 3). During the post monsoon period a branch of the mid latitude  
 322 western jet stream settles south of the Himalayan's. It enters into Bangladesh through the northwest

323 corner of the country. For this reason the northern zone is cooler than the middle and coastal region  
324 these (Ahmed, 2005).

### 325 **5.1.2 Seasonal variation of Rainfall**

326

327 Rainfall is most important phenomena for the country as it is an agro based country and economy of the  
328 country depend on agriculture. There is an extreme large seasonal variation from winter- to summer-  
329 monsoon in Bangladesh as wind blows from different direction for winter and summer. In monsoon  
330 rainfall varies from 943.7 mm to 1708.02 mm. The Seasonal rainfall variation of all regions are given  
331 in the figure 4 and 6.

332 ***In winter*** rainfall varies from 46.92 mm to 51.77 mm. The northwest region is showing higher winter  
333 rainfall than any other region of the country (Figure 4, Fig 5(E) and Table 3). In this season the  
334 remnants of the Mediterranean disturbances enter the sub-continent through the north western corner of  
335 the sub-continent. These weather disturbances travel from west to east along the foothills of the  
336 Himalayas, causing precipitation along their travel path. Some of these disturbances reach the north  
337 western region of Bangladesh. That is why the northwest region of Bangladesh receives more winter  
338 rainfall than any other regions in the country.

339 ***In pre monsoon*** rainfall varies from 290.97 mm in the northwest region to 785.55 mm in the northeast  
340 region (table 3) in this season. Sandwip and Sylhet receive rainfall values of 1281.13 and 1091.97 mm  
341 (Table 2). The middle zone shows less rainfall. (Figure 4)

342 ***In monsoon seasons***, all the regions of the country receive heavy rainfall. As during monsoon the  
343 surface wind blows from the southwest direction and carries a lot of moisture from the Bay of Bengal  
344 and North Indian Ocean this causes a lot of precipitation over Bangladesh and adjoining countries.  
345 High orographic precipitation occurs over the northern hills and mountain slopes of India. The  
346 upstream regions over Meghalaya and Assam get very high amount of precipitation, Cherrapunji which  
347 received world's highest annual precipitation 11387 mm lies over the hilly region close to northeast  
348 region of Bangladesh (Quadir *et al.*, 2003) for which the Sylhet, the one of weather station of northeast

receives highest monsoon rainfall (2250.26 mm) in the country. (Table 2) and northeast receive 1511.24 mm where south zone receive (Islands 17.08.02 mm, Coastal 1587.92 mm and hilly 1474.58 mm) higher rainfall than other zone. In addition the periods of consecutive rain days range is higher in Northeast and southeast than other regions. Ahmed and Kim, (2003) has shown very long episodes of consecutive rain days in the northeast (Sylhet) are due to the additional effect of orographic uplift on the rainfall provided by the Meghalaya Plateau. Similarly, very long episodes of consecutive rain days in the coastal southeast (Cox's Bazar) are due to the additional effect of orographic uplift on the rainfall provided by the coastal hills. From the table 3 and Figure 5.(F), we can conclude that the northeastern and southeaster part of Bangladesh gets the highest precipitation and the north western and southwestern part is relatively dry.

***In post monsoon*** rainfall is less than the monsoon rainfall. In this season highest rainfall occurs in Sylhet (806.78 mm). The Northeast region receives rainfall amount of 634.43 mm. and the middle region receives 450.73 mm (Figure 4 & table 3.). The rainfall amount of the season is slightly enhanced in the northeast region than other region (Same as orographic effect).

363

## 364 **5.2 Relation between Maximum-Minimum Temperature Difference and Rainfall:**

365

366 A remarkable correlation was discovered between this temperature range and the rainfall occurrence (Fig 6). In summer-monsoon, the difference of temperature ( $\Delta T$ ) is lower than any other season while rainfall is higher than any other season, in all regions of the country. It indicates a negative correlation between  $\Delta T$  and the rainfall. Again in monsoon, island and coastal region (south zone) is showing  $\Delta T$  value (lowest  $\Delta T$  value is 5.12°C in islands). Table 3, reveals the variation of  $\Delta T$  over different region and the value is decreasing from the north to the south while at the same way the rainfall increases from northwest to the south (Figure: 7) as south zone located in coastal area and humidity of that zone is higher.









Fig. 6. Seasonal variation of the mean max, min Temperature and Rainfall in different region




Fig. 7. Seasonal variation of the difference of mean Max & Min temperature and rainfall in south zone.

### 5.3 Time Series Analysis

#### 5.3.1 Trend of Temperature

To investigate the temporal variability and trends over different region of Bangladesh, the time series plot and regression analysis has been applied (Appendix fig A-D). Table 4 & 5 and figure 8, and 9 show that in all seasons the mean maximum temperature is increasing except in winter for the northwest and middle zone and mean minimum temperature is increasing as well except hilly region for all season.

***In winter***, the mean maximum temperature is rising by  $0.0754^{\circ}\text{C}/\text{year}$  (Hilly region) to  $0.0076^{\circ}\text{C}/\text{year}$  (islands regions). The northwest and middle zone are showing decreasing rates of  $-0.0004^{\circ}\text{C}/\text{year}$  and  $-0.0069^{\circ}\text{C}/\text{year}$  [Table 4 and Figure 8 (A)]. The mean minimum temperature is increasing in northwest region with rate of  $+0.0281^{\circ}\text{C}/\text{year}$  while the middle zone and coastal region are showing very small increasing rates. The northeast, island and hilly region are showing decreasing rates [(Table 5 and figure 9(A))].

***In the pre-monsoon*** period, mean maximum temperature is rising from  $+0.0034^{\circ}\text{C}/\text{year}$  (middle zone) to  $+0.064^{\circ}\text{C}/\text{year}$  (Hilly region) [(Table 4 and figure 8 (B))]. The mean minimum temperature is increasing in most regions except island and hilly region, where the mean minimum temperature are decreasing with the rate of  $-0.006^{\circ}\text{C}/\text{year}$  and  $-0.047^{\circ}\text{C}/\text{year}$  [(Table 5 and figure 9(B))].

***In the monsoon*** period, mean maximum temperature is rising from  $+0.0167^{\circ}\text{C}/\text{year}$  (northeast region) to  $+0.056^{\circ}\text{C}/\text{year}$  (Hilly region) [(Table 4 and figure 8(C))]. Like pre monsoon, the mean minimum temperature at island and hilly region are decreasing with values of  $-0.004^{\circ}\text{C}/\text{year}$  and  $-0.003^{\circ}\text{C}/\text{year}$  [(Table 5 and figure 9(C))].

***In the post monsoon*** period, it was found that in all regions the mean maximum temperature is increasing with a rate varying from  $+0.029^{\circ}\text{C}/\text{year}$  (Island) to  $+0.068^{\circ}\text{C}/\text{year}$  (Hilly) [(Table 4 and figure 8(D))]. The mean minimum temperature is decreasing only in the hilly region with a rate of  $-0.04^{\circ}\text{C}/\text{year}$  [(Table 5 and figure 9(D))].



404

405

406


407

408

409

410

From the above calculation we can conclude that overall the temperature is increasing over the whole country (the rate of max temp is  $+0.028^{\circ}/\text{year}$ ) concurrent with the global temperature increase. From the table 3 and 4, it is clear that the increasing rate of mean maximum temperature is higher in southern zone than in the other zones. It may be caused by increasing of sea surface temperature. (Josheph, 1995) shown increase of SST (Sea Surface temperature) using data for the Bay of Bengal over the period 1951-1987, but why this should affect in the first place the maximum temperatures is not clear.



411

412

413

Fig. 8. The mean maximum Temperature trend in (A) Winter (B) Pre monsoon (C) Monsoon (D) Post monsoon



414

415 **Table 4. Mean maximum temperature trends (°C/year)**

| <b>Regions</b>     | <b>Winter</b>  | <b>Pre monsoon</b> | <b>Monsoon</b> | <b>Post monsoon</b> |
|--------------------|----------------|--------------------|----------------|---------------------|
| <b>Northwest</b>   | <i>-0.0004</i> | <i>+0.008</i>      | <i>+0.024</i>  | <i>+0.025</i>       |
| <b>Northeast</b>   | <i>+0.013</i>  | <i>+0.017</i>      | <i>+0.017</i>  | <i>+0.032</i>       |
| <b>Middle zone</b> | <i>-0.007</i>  | <i>+0.003</i>      | <i>+0.031</i>  | <i>+0.031</i>       |
| <b>Coastal</b>     | <i>+0.036</i>  | <i>+0.038</i>      | <i>+0.043</i>  | <i>+0.047</i>       |
| <b>Islands</b>     | <i>+0.008</i>  | <i>+0.049</i>      | <i>+0.026</i>  | <i>+0.018</i>       |
| <b>Hilly</b>       | <i>+0.075</i>  | <i>+0.066</i>      | <i>+0.055</i>  | <i>+0.068</i>       |

416

417 **Table 5. Mean minimum temperature trends (°C/year)**

| <b>Regions</b>     | <b>Winter</b> | <b>Pre monsoon</b> | <b>Monsoon</b> | <b>Post monsoon</b> |
|--------------------|---------------|--------------------|----------------|---------------------|
| <b>Northwest</b>   | <i>+0.028</i> | <i>+0.048</i>      | <i>+0.035</i>  | <i>+0.06</i>        |
| <b>Northeast</b>   | <i>-0.013</i> | <i>+0.023</i>      | <i>+0.014</i>  | <i>+0.047</i>       |
| <b>Middle zone</b> | <i>+0.002</i> | <i>+0.010</i>      | <i>+0.002</i>  | <i>+0.0169</i>      |
| <b>Coastal</b>     | <i>+0.001</i> | <i>+0.009</i>      | <i>+0.011</i>  | <i>+0.0315</i>      |
| <b>Islands</b>     | <i>-0.037</i> | <i>-0.006</i>      | <i>-0.004</i>  | <i>+0.0006</i>      |
| <b>Hilly</b>       | <i>-0.044</i> | <i>-0.045</i>      | <i>-0.003</i>  | <i>-0.0411</i>      |

418

419





Fig. 9. The mean minimum Temperature trend in (A) Winter (B) Pre monsoon (C) Monsoon (D) Post monsoon


### 5.3.2 Trend of Precipitation

Several studies (Quadir *et al.*, 2003) have indicated that during the recent decade the precipitation of Bangladesh has been increasing (Fig 10). Restricting ourselves to the two extreme seasons (winter and summer-monsoon) I found that the precipitation trends in the monsoon season are positive in all the regions of the country except the northwest region (-3.06 mm/ year) . The Hilly region shows the highest increase in rainfall (+17.78 mm/ year). (Table 6, and Figure 11). But in winter, precipitation shows a decreasing trend in all the regions except the islands and hilly region. The increasing rate of

precipitation for the island has been found as +0.016 mm/ year. Hilly region shows 0.93mm/ year. (Table 6, and Figure 11). So we can conclude that in the monsoon period the rainfall is increasing in all regions except in the northwest region. The northwest is clearly becoming more arid. May be this is enhanced by the large scale deforestation processes going on.



**Fig. 10. Time series of monsoon rainfall of Bangladesh over the period 1981-2003**



**Fig. 11. The Rainfall trend in (A) Winter (B) Monsoon**

441  
442  
443  
444  
445  
446  
447  
448  
449  
450  
451  
452  
453  
454  
455  
456  
457  
458  
459

**Table 6. Rainfall trends in mm/year**

| <b>Regions</b>     | <b>Winter</b>  | <b>Monsoon</b> |
|--------------------|----------------|----------------|
| <b>Northwest</b>   | <i>-1.0109</i> | <i>-3.057</i>  |
| <b>Northeast</b>   | <i>-0.833</i>  | <i>+0.096</i>  |
| <b>Middle zone</b> | <i>-0.734</i>  | <i>+0.727</i>  |
| <b>Coastal</b>     | <i>-0.131</i>  | <i>+3.827</i>  |
| <b>Islands</b>     | <i>+0.161</i>  | <i>+12.183</i> |
| <b>Hilly</b>       | <i>+0.932</i>  | <i>+17.784</i> |

**5.4 Regional differences:**

As the emphasis in this research was laid upon the regional differences of the climatic trends for Bangladesh, we present maps (Figure 8 & 9) displaying the trends and tested the trends for each region against the trends for the other regions (t-test). The t-test was only carried out for the maximum and minimum temperatures and the results shown in table 7. From this table we can see that essentially the Hilly region and the northwest region stand out as separate regions from the overall Bangladesh area: the Hilly region as a region with an exceptional large increase in maximum temperature and decrease in minimum temperature. The northwest region with a very small increase in maximum temperature and a comparative large increase in minimum temperature.

Table 7: Statistical t-test.

| Maximum temperature                  |           |           |             |         |         |       |
|--------------------------------------|-----------|-----------|-------------|---------|---------|-------|
| Region                               | Northwest | Northeast | Middle zone | Coastal | Islands | Hilly |
| Northwest                            | I         | I         | I           | S       | I       | S     |
| Northeast                            | I         | I         | I           | S       | I       | S     |
| Middle zone                          | I         | I         | I           | I       | I       | S     |
| Coastal                              | S         | S         | I           | I       | I       | S     |
| islands                              | I         | I         | I           | I       | I       | S     |
| Hilly                                | S         | S         | S           | S       | S       | I     |
| Minimum temperature                  |           |           |             |         |         |       |
| Region                               | Northwest | Northeast | Middle zone | Coastal | islands | Hilly |
| Northwest                            | I         | I         | S           | S       | S       | S     |
| Northeast                            | I         | I         | I           | I       | I       | S     |
| Middle zone                          | S         | I         | I           | I       | I       | S     |
| Coastal                              | S         | I         | I           | I       | I       | S     |
| islands                              | S         | I         | I           | I       | I       | S     |
| Hilly                                | S         | S         | S           | S       | I       | I     |
| Insignificant = I<br>Significant = S |           |           |             |         |         |       |

## 6. Conclusion:

In this research, the recent climatic variability and trend of temperature and precipitation over Bangladesh have been studied. The conclusion of this study can be summarized as follows:

- (ix) The seasonal variation for both maximum and minimum temperature is rather small. In winter there exists a north-south temperature gradient which reverses during summer (pre-monsoon and monsoon).
- (x) The seasonal variation for the rainfall is extremely large. In winter there exists a south-north precipitation gradient which reverses during the other seasons.
- (xi) In all seasons the difference of maximum and minimum temperature is higher in the north zone than in the south zone. We discovered a remarkable correlation between this temperature range and the rainfall occurrence.

475 (xii) Overall we find a positive trend in the rainfall, especially large during the summer monsoon and  
476 in the southern zone. The northwest area on the contrary is characterized by decreasing rainfall  
477 values both in winter and summer. Thus there appears to be an increasing contrast between the  
478 summer and winter monsoon.

479 (xiii) The maximum temperature shows a remarkable increasing trend especially in winter. The  
480 minimum temperature on the other hand shows an overall decreasing trend except in the north.  
481 This results in a conspicuous increasing trend for the temperature range (Maximum –  
482 minimum). Overall the temperature of whole of the country is increasing as same as global  
483 temperature is increasing. It was found in this study that trend in the south zone is higher than  
484 other zone of the country. It may be caused by increasing of sea surface temperature. Global  
485 warming would result in an increase in sea surface temperatures as a result of which changes in  
486 the frequency, intensity or tracks in cyclones hitting the coastal zones may take place. So an  
487 increasing in SST will lead to higher storm surges and higher risk of coastal disaster in low  
488 coastal areas particularly of Bangladesh.

489 (xiv) The regional analysis identified two regions in Bangladesh which behave differently, i.e. the  
490 Hilly region in the southeast and the northwest area. The northwest area is clearly under the  
491 influence of processes more typical for middle latitudes, while in the hilly region orographic  
492 and coastal processes might be at work.

493

494

495

496

497

498

499

500    **Acknowledgements**

501    This article is based on my MSc thesis, completed at Vrije Universiteit, Brussel, Belgium as part of  
502    Master of Environmental Science and Technology, programme in 2005. First, I would like to thank my  
503    thesis supervisor, Prof. Dr. Hugo Decler (Retired), Department of Geography, Faculty of Sciences,  
504    Vrije Universiteit Brussel. I would also like to express my thanks to late Professor Rafique Ahmed,  
505    Dep. Of Geography and Earth Science, University of Wisconsin-La Crosse, La Crosse, Wisconsin  
506    54601, USA for his careful guidance and cordial help right from the beginning of the research and also  
507    reviewing the paper. I would like to express my gratitude and thanks to Goutam Chandra Mridha,  
508    Senior Specialist, IWM for his careful guidance to create map and demonstration right from the initial  
509    stage of my research.

510

511    **References:**

- 512    ADB (Asian Development Bank) (2014) ‘Bangladesh’, in Bank, A. D. (ed.) *ASIAN DEVELOPMENT*  
513    *OUTLOOK 2014 FISCAL POLICY FOR INCLUSIVE GROWTH*, pp. 158–163.
- 514    Ahmed, R. (1989) ‘Probabilistic estimates of rainfall extremes in Bangladesh during the pre-monsoon  
515    season’, *Indian Geographical Journal*, 64, pp. 39–53.
- 516    Ahmed, R. (2003) ‘Climate of Bangladesh.’, in Islam, S. (ed.) *Banglapedia*. Bangladesh: Asiatic  
517    Society of Bangladesh., pp. 45–48.
- 518    Ahmed, R. (2004) ‘Climate of Bangladesh.’, in *Abhaao O Jalabayu Bijnan” (in Bangla)*. Dhaka:  
519    Gyankosh Prakashani.
- 520    Ahmed, R. (2005) *Personal communication with Professor Rafique Ahmed, Dep. of Geography and*  
521    *Earth Science, University of Wisconsin-La Crosse, La Crosse, Wisconsin 54601, USA .*
- 522    Ahmed, R. and Karmakar, S. (1993a) ‘Arrival a n d withdrawal dates of the summer monsoon in  
523    bangladesh’, *International Journal of Climatology*, 13, pp. 727–740.
- 524    Ahmed, R. and Karmakar, S. (1993b) ‘ARRIVAL A N D WITHDRAWAL DATES OF THE

525 SUMMER MONSOON IN BANGLADESH', *International Journal of Climatology*, 13, pp. 727–740.

526 Ahmed, R. and Kim, K. (2003) 'Patterns of Daily Rainfall in Bangladesh During the Summer Monsoon

527 Season: Case Studies at Three Stations', *Physical Geography*, 24(4), pp. 295–318.

528 Asian Development Bank (1994) *Climate Change in Asia: Bangladesh*. Manila: Asian Development

529 Bank.

530 Brammer, H. (2000) *Agroecological Aspect of Agricultural Research in Bangladesh*. Dhaka: The

531 university press limited.

532 Brammer, H. (2004) *Can Bangladesh be Protected from Floods?* Dhaka: The university press limited.

533 Christensen *et al.* (2007) 'The Physical Science Basis. Contribution of Working Group I to the Fourth

534 Assessment Report of the Intergovernmental Panel on Climate Change', in Solomon, S., D. et al. (eds)

535 *Regional Climate Projections*. New York: Cambridge University Press, Cambridge.

536 Debsarma, S. K. (2003) 'Intra-Annual and Inter- Annual Variation of Rainfall over different Regions of

537 Bangladesh', in *SAARC seminar on Climatic variability in the south Asian region and its impact*.

538 Ericksen, N. J., Ahmad, Q. K. and Chowdhury, A. R. (1997) *Socio-Economic Implications of Climate*

539 *Change for Bangladesh*. Dhaka: Bangladesh Unnayan Parishad (BUP). Available at:

540 [http://www.nirapad.org.bd/admin/soft\\_archive/1308128275\\_Socio Economic Implications of Climate](http://www.nirapad.org.bd/admin/soft_archive/1308128275_Socio_Economic_Implications_of_Climate_Change_for_Bangladesh.pdf)

541 [Change for Bangladesh.pdf](http://www.nirapad.org.bd/admin/soft_archive/1308128275_Socio_Economic_Implications_of_Climate_Change_for_Bangladesh.pdf).

542 Gosling, S. N. *et al.* (2011) *Climate: Observations, projections and impacts: Bangladesh*, MetOffice.

543 Exeter.

544 Huq, M. S. (1974) *Climate of Bangladesh*. In: *Studies in Bangladesh Geography Savar*: Edited by K.

545 F. M. Department of Geography, Jahangir Nagar University.

546 Immerzeel, W. (2008) 'Historical trends and future predictions of climate variability', *International*

547 *Journal of Climatology*, 254(May 2007), pp. 243–254. doi: 10.1002/joc.

548 IPCC (2013) *Summary for Policymakers*. In: *Climate Change 2013: The Physical Science Basis*.

549 *Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on*



550 *Climate Change, Energy and Environment*. Edited by V. B. and P. M. M. (eds. )]. T.F., D. Qin, G.-K.

551 Plattner, M. Tignor, S.K. Allen, J. Boschung, A. Nauels, Y. Xia. Cambridge, UK, New York USA. doi:

552 10.1260/095830507781076194.

553 IPCC (2014a) *The IPCC 's Fifth Assessment Report What 's in it for South Asia ?*

554 IPCC (2014b) *The IPCC 's Fifth Assessment Report What 's in it for South Asia ?* Available at:

555 <http://cdkn.org/wp-content/uploads/2014/04/CDKN-IPCC-Whats-in-it-for-South-Asia-AR5.pdf>.

556 Johnson, B. L. . (1982) *Bangladesh*. London Uk: Heinemann.

557 Josheph, P. . (1995) 'Changes in the frequency and tracks of tropical cyclone in the indian Ocean seas',

558 in *The workshop of Global Chhange and Tropical Cyclone, Dhaka, Bangladesh*,. Dhaka, Bangladesh.

559 Karmakar, s. and S. M. L. (2000) *Recent Climatic Change in Bangladesh, SAARC Meteorological*

560 *Research Centre (SMRC)*. Dhaka, Bangladesh.

561 Lutz, A. F. *et al.* (2016) 'Selecting representative climate models for climate change impact studies: an

562 advanced envelope-based selection approach', *International Journal of Climatology*, 36(12), pp. 3988–

563 4005. doi: 10.1002/joc.4608.

564 May, W. (2004) 'Simulation of the variability and extremes of daily rainfall during the Indian summer

565 monsoon for present and future times in a global time-slice experiment', *Climate Dynamics*, 22(2), pp.

566 183–204. doi: 10.1007/s00382-003-0373-x.

567 Mirza, M. Q. *et al.* (1998) 'Trends and persistence in precipitation in the Ganges, Brahmaputra and

568 Meghna river basins', *Hydrological Sciences Journal*, 43(6), pp. 845–858. doi:

569 10.1080/02626669809492182.

570 Quadir, A. D. *et al.* (2003) 'Climate change and its Impacts on Bangladesh Floods over the Past

571 Decades', in *Processing of SAARC (South Asian Association for Regional Co-operation) seminar on*

572 *Climate Variability in the South Asian Region and Its Impacts*. Dhaka, Bangladesh: SAARC

573 Meteorological Research Centre (SMRC),.

574 Ramamasy, S. and Baas, S. (2007) *Climate variability and change: adaptation to drought in*

575 *Bangladesh, Case Study - Institutions for Rural Development, FAO*. Rome: FAO. Available at:  
576 cabi:20083071262.

577 Rasheed, S. K. B. (2008) *Resource and Environmental Profile*. Dhaka, Bangladesh: A H Development  
578 Publishing House.

579 Robinson, P. J. and Henderson-Seller, A. (1999) *Contemporary Climatology*. 2nd edn. Harlow: Pearson  
580 Education Limited.

581 Salahuddin, A. *et al.* (2006) ‘Teleconnections between the sea surface temperature in the Bay of Bengal  
582 and monsoon rainfall in Bangladesh’, *Global and Planetary Change*, 53(3), pp. 188–197. doi:  
583 <https://doi.org/10.1016/j.gloplacha.2006.06.001>.

584 Sarker, M. S. H. (2005) *Temperature and Precipitation Changes in Bangladesh*. Vrije Universiteit  
585 Brussel: Belgium.

586 Sarker, M. S. H. (2009) “‘Climatic Variability and Its trend Over Different Region of Bangladesh’.”, in  
587 *The 21st Conference on Climate Variability and Change which was as part of the AMS (American*  
588 *Meteorological Society) 89th Annual Meeting, (11-15 January 2009) in Phoenix, AZ, USA*.

589 Sarker, M. S. H. (2015) *Environmental change and its impact on migration in Bangladesh*. University  
590 of Sheffield.

591 Shahid, S. (2011) ‘Trends in extreme rainfall events of Bangladesh’, *Theoretical and Applied*  
592 *Climatology*, 104(3–4), pp. 489–499. doi: 10.1007/s00704-010-0363-y.

593 Shamsuddin, S. D. and Ahmed, R. (1974) ‘Variability of Annual Rainfall of Bangladesh’, *Journal of*  
594 *Bangladesh National Geographical Association*, 2, pp. 13–20.


595 Spate, O. H. K. and Learmonth, A. T. . (1984) *Spate, Oskar Hermann Khristian*. Delhi, India:  
596 Munshiram Manoharlal.

600

601

602 Appendix:

603



604

605

606 Figure A: Time series of winter mean maximum temperature.

607









Figure B: Time series of pre monsoon mean maximum temperature.




Figure C: Time series of monsoon mean maximum temperature.









Figure D: Time series of post monsoon mean maximum temperature.