The Common Representative Intermediates Mechanism version 2 in the United Kingdom Chemistry and Aerosols Model

Scott Archer-Nicholls¹, Nathan Luke Abraham², Youngsub Matthew Shin³, James Weber⁴, Maria Rosa Russo⁵, Douglas Lowe⁶, Steven Utembe⁷, Fiona M. O'Connor⁸, Brian Kerridge⁹, Barry Latter¹⁰, Richard Siddans¹¹, Michael Jenkin¹², Oliver Wild¹³, and Alexander Thomas Archibald¹

¹University of Cambridge
²NCAS, University of Cambridge
³Centre for Atmospheric Science, Department of Chemistry, University of Cambridge
⁴Centre for Atmospheric, Department of Chemistry, University of Cambridge
⁵NCAS Climate, Centre for Atmospheric Science
⁶University of Manchester
⁷Environmental Protection Authority Victoria
⁸Met Office Hadley Centre
⁹Rutherford Appleton Laboratories
¹⁰RAL
¹¹Rutherford Appleton Laboratory
¹²Atmospheric Chemistry Services
¹³Lancaster University

November 24, 2022

Abstract

We document the implementation of the Common Representative Intermediates Mechanism version 2, reduction 5 (CRIv2-R5) into the United Kingdom Chemistry and Aerosol model (UKCA) version 10.9. The mechanism is merged with the stratospheric chemistry already used by the StratTrop mechanism, as used in UKCA and the UK Earth System Model (UKESM1), to create a new CRI-Strat mechanism. CRI-Strat simulates a more comprehensive treatment of non-methane volatile organic compounds (NMVOCs) and provides traceability with the Master Chemical Mechanism (MCM). In total, CRI-Strat simulates the chemistry of 233 species competing in 613 reactions (compared to 87 species and 305 reactions in the existing StratTrop mechanism). However, while more than twice as complex than StratTrop, the new mechanism is only 75% more computationally expensive. CRI-Strat is evaluated against an array of in situ and remote sensing observations and simulations using the StratTrop mechanism in the UKCA model. It is found to increase production of ozone near the surface, leading to

higher ozone concentrations compared to surface observations. However, ozone loss is also greater in CRI-Strat, leading to less ozone away from emission sources and a similar tropospheric ozone burden compared to StratTrop. CRI-Strat also produces more carbon monoxide than StratTrop, particularly downwind of biogenic VOC emission sources, but has lower burdens of nitrogen oxides as more is converted into reservoir species. The changes to tropospheric ozone and nitrogen budgets are sensitive to the treatment of NMVOC emissions, highlighting the need to reduce uncertainty in these emissions to improve representation of tropospheric chemical composition.

The Common Representative Intermediates Mechanism version 2 in the United Kingdom Chemistry and Aerosols Model

S. Archer-Nicholls¹, N. L. Abraham^{1,2}, Y. M. Shin¹, J. Weber¹, M. R. Russo^{1,2}, D. Lowe³, S. Utembe⁴, F. M. O'Connor⁵, B. Kerridge^{6,7}, B. Latter^{6,7}, R. Siddans^{6,7}, M. Jenkin⁸, O. Wild⁹, A. T. Archibald^{1,2}

4 5 6

7 8	¹ Department of Chemistry, University of Cambridge, Cambridge, UK ² National Centre for Atmospheric Science, Department of Chemistry, University of Cambridge,
9	Cambridge, UK
10	³ University of Manchester, Manchester, UK
11	⁴ University of Melbourne, Melbourne, Australia
12	⁵ Met Office Hadley Centre, Exeter, UK
13	⁶ Remote Sensing Group, STFC Rutherford Appleton Laboratory, UK
14	⁷ NERC National Centre for Earth Observation, Rutherford Appleton Laboratory, UK
15	⁸ Atmospheric Chemistry Services, Okehampton, Devon, UK
16	⁹ Lancaster University, Lancaster, UK

17	Key Points:
18	• The CRI-Strat mechanism has been integrated into the UKCA model, greatly in-
19	creasing the complexity of VOC chemistry compared to StratTrop.
20	• CRI-Strat simulates higher surface ozone compared to StratTrop due to greater
21	production, but tropospheric ozone burden is similar.
22	• The ozone and oxidised nitrogen budgets when running with the CRI-Strat mech
23	anism show high sensitivity to the input NMVOC emissions.

Corresponding author: Alex T. Archibald, ata27@cam.ac.uk

24 Abstract

We document the implementation of the Common Representative Intermediates Mech-25 anism version 2, reduction 5 (CRIv2-R5) into the United Kingdom Chemistry and Aerosol 26 model (UKCA) version 10.9. The mechanism is merged with the stratospheric chemistry 27 already used by the StratTrop mechanism, as used in UKCA and the UK Earth System 28 Model (UKESM1), to create a new CRI-Strat mechanism. CRI-Strat simulates a more 29 comprehensive treatment of non-methane volatile organic compounds (NMVOCs) and 30 provides traceability with the Master Chemical Mechanism (MCM). In total, CRI-Strat 31 simulates the chemistry of 233 species competing in 613 reactions (compared to 87 species 32 and 305 reactions in the existing StratTrop mechanism). However, while more than twice 33 as complex than StratTrop, the new mechanism is only 75% more computationally ex-34 pensive. CRI-Strat is evaluated against an array of *in situ* and remote sensing observa-35 tions and simulations using the StratTrop mechanism in the UKCA model. It is found 36 to increase production of ozone near the surface, leading to higher ozone concentrations 37 compared to surface observations. However, ozone loss is also greater in CRI-Strat, lead-38 ing to less ozone away from emission sources and a similar tropospheric ozone burden 39 compared to StratTrop. CRI-Strat also produces more carbon monoxide than StratTrop, 40 particularly downwind of biogenic VOC emission sources, but has lower burdens of ni-41 trogen oxides as more is converted into reservoir species. The changes to tropospheric 42 43 ozone and nitrogen budgets are sensitive to the treatment of NMVOC emissions, highlighting the need to reduce uncertainty in these emissions to improve representation of 44 tropospheric chemical composition. 45

46 Plain Language Summary

To understand the climate and predict how it will change in the future, we need 47 to understand its chemical composition - the trace gases and small particles that exist 48 in tiny quantities in the atmosphere. A key tool we use to do this are computer mod-49 els which simulate the atmosphere and processes within it. Key processes include the 50 formation of ozone, a harmful pollutant and greenhouse gas in the lower atmosphere. How-51 ever, the chemistry involved in forming ozone is very complicated, so computer simula-52 tions of the atmosphere must greatly simplify the chemistry. These simple schemes may 53 introduce errors in the model. We also have much more complex chemical mechanisms 54 which simulate our best understanding of all chemical reactions, but these complex schemes 55 require too much computational power to be used when simulating the whole atmosphere. 56 In this paper, we describe the implementation of a chemical mechanism that sits between 57 these levels of complexity, realistically simulating the formation and destruction of ozone 58 without being too slow to run. We compare this new mechanism against measurements 59 taken of the atmosphere and the preexisting, simpler chemical mechanism and show that 60 the new mechanism greatly enhances the amount of ozone that is produced. 61

62 1 Introduction

Understanding chemical processes in the lower atmosphere is of vital importance 63 for tackling the problems of air pollution and making accurate projections of how the 64 Earth system will change due to human activity (Sillman, 1999; Akimoto, 2003; Von Schnei-65 demesser et al., 2015; Monks et al., 2015; Boucher et al., 2013). However, the chemistry 66 of the troposphere is extremely complicated because of the wide variety of non methane 67 volatile organic compounds (NMVOCs) whose structures are diverse and whose lifetimes 68 and abundances cover many orders of magnitude (Atkinson, 1990; M. Jenkin et al., 1997; 69 Goldstein & Galbally, 2007). Two particularly challenging aspects of tropospheric chem-70 istry regard the understanding of the formation and destruction of tropospheric ozone 71 and the impacts of aerosols, with much of the difficulty in both of these research top-72 ics stemming from the importance of NMVOCs. The rate of production of ozone is non-73

linearly dependent on the combination of nitrogen oxides $(NO_x = NONO_2)$ and NMVOCs 74 levels, with high rates of ozone production occurring when levels of both are high and 75 net ozone destruction occurring when there is a large excess of NMVOCs (NO_x-limited 76 regime) or an excess of NO_x (VOC-limited regime) (Sillman, 1999; Monks et al., 2015). 77 Larger NMVOC molecules, such as monoterpenes, aromatic compounds and long chain 78 n-alkanes $(C_{>10}$ – where C indicates the number of carbon atoms in the NMVOC) play 79 an important role in the generation of secondary organic aerosol (SOA). SOA can make 80 up over 50% of submicron aerosol mass (Jimenez et al., 2009), but we still have signif-81 icant uncertainty in the exact chemical makeup, and models still fail to accurately sim-82 ulate it (Tsigaridis et al., 2014; Hodzic et al., 2020). Unfortunately, many thousands of 83 different NMVOC species have been identified in the atmosphere, and many more are 84 yet to be discovered, making a complete representation of all NMVOC species and their 85 chemistry in a model an impossible task (Goldstein & Galbally, 2007; Heald & Kroll, 2020). 86

The key mapping between the input NMVOCs and their effects on ozone and SOA 87 is their oxidation mechanism. Such mechanisms are well known from laboratory exper-88 iments for the simplest NMVOCs ($C_{<5}$). Based on the wealth of experimental data (McGillen 89 et al., 2020), structure activity relationships (SARs) have been derived to fill in the gaps 90 (M. E. Jenkin et al., 2018b, 2018a; M. E. Jenkin, Valorso, et al., 2019) and to extrap-91 olate our understanding of NMVOC oxidation to cover a wide range of structures and 92 configurations. Aumont et al. (2005) have shown that the oxidation process in the at-93 mosphere can be treated as a geometric problem, with the number of species produced 94 during the oxidation of an alkane NMVOC with n carbon atoms being given by: 95

$$\Omega \approx \sum_{i=2}^{n} \frac{1}{2} (11)^2 (7)^{i-2}.$$
 (1)

For a C₅ compound, this equation leads to $\approx 10^5$ species forming, presenting a huge amount of complexity.

96

To approach the problems of complexity relating to NMVOC chemistry in the at-99 mosphere, researchers have typically followed one of two routes for developing chemistry 100 mechanisms depending on the research questions being tackled and relevant spatial and 101 temporal scales. Detailed, explicit mechanisms (Aumont et al., 2005) and near-explicit 102 mechanisms, such as the Master Chemical Mechanism (MCM; Saunders et al. (2003); 103 M. E. Jenkin et al. (2003)), comprise of the amalgamation of known relevant chemistry 104 as measured from laboratory chemical kinetic studies (Atkinson et al., 2006; S. P. Sander 105 et al., 2011) and SARs (M. E. Jenkin et al., 2018b, 2018a; M. E. Jenkin, Valorso, et al., 106 2019) and evaluated against field and chamber experiments (M. E. Jenkin et al., 2012; 107 Novelli et al., 2018). These mechanisms represent our best understanding of chemical 108 processes in the atmosphere, often comprising of thousands of species and many times 109 more reactions, and are continuously updated and expanded as new data and understand-110 ing comes to light. However, this makes them computationally very expensive to run and 111 hence these explicit mechanisms are mostly used in box model studies (M. E. Jenkin et 112 al., 2015; Derwent, 2017). 113

An alternative design approach is to construct mechanisms that are as simple as 114 possible but as complicated as necessary, the aim being to represent the key chemical 115 processes and their interactions with as few chemical species and reactions as possible. 116 Such atmospheric chemistry schemes typically have 10s of species and dozens to hun-117 dreds of reactions (e.g., the StratTrop scheme which simulates 81 species, 291 reactions 118 (Archibald et al., 2020)). Chemical processes are still informed by the best available data, 119 but similar species are lumped together or represented by surrogate species and more 120 complex but important processes are parameterised to reduce the mechanism complex-121 ity. Mechanisms may also become quite specialist, with those designed to simulate ur-122 ban air pollution (e.g., (Stockwell et al., 1990)) differing greatly from those intended to 123 simulate the whole atmosphere at a coarse resolution (e.g., (Archibald et al., 2020)), as 124

they focus on different aspects of the chemistry system occurring in the real atmosphere.
These approximations are necessary to run interactive chemistry in 3D models that must
also simulate other key processes such as transport, deposition, clouds and radiation. Even
so, the chemistry component is often the most computationally expensive part of a 3D
model (Esenturk et al., 2018).

These different approaches leave open a gap between our most comprehensive mech-130 anisms and simpler ones. Simpler schemes can perform well for the photochemical con-131 ditions they were designed for, but may perform poorly when simulating other regions, 132 133 or fail to properly represent how chemical conditions should change in response to changes in emissions or climate. Due to their complexity, it is impractical to use near-explicit chem-134 istry schemes in the same 3D model setup as simpler mechanisms to see how they would 135 perform instead. While they can be evaluated against comprehensive schemes in box model 136 simulations, when results differ it is difficult to pinpoint which aspect of the chemistry 137 is causing the differences, due to the lack of traceability, and it can be unclear whether 138 the simpler mechanisms respond realistically in photochemical conditions outside of those 139 evaluated in box model experiments. There is therefore a clear need for intermediate com-140 plexity mechanisms - ones which are fully traceable to more comprehensive schemes and 141 are known to respond similarly to changes in forcings, but are still simple enough to be 142 used in 3D interactive models. 143

The Common Representative Intermediates (CRI) mechanism (M. E. Jenkin et al., 144 2002; M. Jenkin et al., 2008) is just such an intermediate complexity mechanism. The 145 number of species and reactions are reduced by over an order of magnitude compared 146 with the MCM. However, through a rigorous development process, in which it was fully 147 and systematically evaluated against the more complex MCM mechanism at each stage 148 of complexity reduction and no change that significantly degraded representation of ozone 149 production was allowed through (Watson et al., 2008), the scheme is fully traceable to 150 the MCM. The end result is a mechanism that is simple enough to be run in a 3D model, 151 but which we can be confident responds to changes in emissions and conditions accord-152 ing to our best understanding as represented in the MCM. Intermediate complexity mech-153 anisms offer enormous benefits when used in 3D models as a research tool to study the 154 importance of chemical processes which are ignored in simpler schemes and as a bench-155 mark against which to test, evaluate and inform development of these simpler schemes. 156 By having more confidence in the representation of gas-phase chemistry, it is possible 157 to attribute remaining model biases to other structural components of the model. 158

The CRI mechanism has been used in several models now, including the STOCHEM 159 Lagrangian global chemical transport model (S. Utembe et al., 2010; M. A. H. Khan et 160 al., 2015), the Weather Research and Forecasting model with chemistry (WRF-Chem), 161 an online regional coupled model (Archer-Nicholls et al., 2014; Lowe et al., 2015; M. A. Khan 162 et al., 2019) and in a regional nested configuration of the European Monitoring and Eval-163 uation Programme chemical transport model for the UK (EMEP4UK) (Hood et al., 2018). 164 However, to the best of our knowledge, it has never been used in a global chemistry-climate 165 model. In this paper we document the implementation of the CRIv2-R5 mechanism in 166 the United Kingdom Chemistry and Aerosol (UKCA) model (Morgenstern et al., 2009; 167 O'Connor et al., 2014), as used in the Met Office Unified Model (UM) and the UK's Earth 168 System Model (UKESM1) (Sellar et al., 2019), evaluate it against a suite of observations 169 and rigorously compare the new mechanism to the existing chemical mechanism (Strat-170 Trop; Archibald et al. (2020)). We note that there are differences in the reaction rate 171 coefficients in the two mechanisms that reflect their independent development and re-172 liance on different assessments of kinetic parameters, which has a bearing on model sim-173 ulations (Newsome & Evans, 2017). We also explore how differences in the allocation of 174 NMVOC emissions in the two mechanisms contributes to the differences between them. 175

The UKESM1 model is used for quantifying and understanding climate forcing, including as part of the Coupled Model Intercomparison Project Phase 6 (CMIP6) (Eyring et al., 2016), making projections of future air quality (including crop yields and human health impacts), and increasingly being used to quantify impacts of mitigation. The implementation of the CRI mechanism into the UKCA, UM and UKESM1 models represents a step change the potential for simulating the complex chemistry-climate interactions between ozone and NMVOCs in the coupled Earth System across many chemical environments and multi-century timescales.

¹⁸⁴ 2 Model Description

2.1 The UKCA model

The United Kingdom Chemistry and Aerosols (UKCA) model is a sub-model of 186 the Met Office Unified Model (UM) and is designed to simulate atmospheric composi-187 tion for weather and climate modelling. UKCA is a part of the UKESM1 (Sellar et al., 188 2019) Earth system model, and uses the Chemistry of Stratosphere and Troposphere (Strat-189 Trop) chemical mechanism (Archibald et al., 2020), which merges the Stratospheric and 190 Tropospheric chemical mechanisms described by (Morgenstern et al., 2009) and (O'Connor 191 et al., 2014), respectively. The UKCA model provides ozone, methane, and nitrous ox-192 ide fields to the UM radiation scheme, as well as calculating oxidant fields that are used 193 to drive the GLOMAP-mode aerosol scheme (Mann et al., 2010; Mulcahy et al., 2018, 194 2020).195

The StratTrop chemical mechanism and its implementation in UKESM1 is described 196 in detail by Archibald et al. (2020). It uses the Fast-JX photolysis scheme (Neu et al., 197 2007), which was implemented in UKCA as described by Telford et al. (2013). The ASAD 198 chemical mechanism framework (Carver et al., 1997) is used to provide a flexible and ex-199 tendable approach to mechanism development and to enable a choice of numerical in-200 tegration schemes. A sparse-matrix Newton-Raphson chemical solver (Wild & Prather, 201 2000) is used here, applying the quasi-Newton approximations recommended by Esenturk 202 et al. (2018) to reduce run time. Wet deposition of soluble chemical compounds is pa-203 rameterised following Giannakopoulos et al. (1999) and dry deposition is based on the 204 resistance type model of Wesley (1989), as described by O'Connor et al. (2014). 205

206

185

2.2 The Common Representative Intermediates Mechanism

The Common Representative Intermediates (CRI) Mechanism has been described 207 in detail in previous papers (M. E. Jenkin et al., 2002; M. Jenkin et al., 2008; Watson et al., 2008; S. R. Utembe et al., 2009), so this section only covers a brief overview of the 209 design philosophy. CRI is a fully traceable, reduced complexity representation of the Mas-210 ter Chemical Mechanism (MCM) (M. E. Jenkin et al., 2002; Saunders et al., 2003), with 211 CRI version 2 optimised against the MCM version 3.1. The chemistry of inorganic com-212 pounds and smaller initial organic molecules (such as methane, ethane and ethene) is 213 functionally identical in CRI to the full MCM. However, the CRI scheme substantially 214 reduces the total number of species and reactions by lumping the intermediate oxida-215 tion products of larger NMVOC species based on an index defined as the total number 216 of carbon-carbon (C - C) and carbon-hydrogen (C - H) bonds, counting double bonds 217 (C = C) as two. This "CRI index" can be understood as the maximum potential num-218 ber of O_x molecules ($O_x = O_3 + NO_2$) generated by the VOC in question, assuming com-219 plete oxidation to CO_2 and H_2O , with every HO_2 and RO_2 molecule created convert-220 ing one NO molecule to NO_2 , thereby generating one ozone molecule when the NO_2 is 221 photolysed (M. E. Jenkin et al., 2002). This means that every primary NMVOC species 222 in CRI will produce the same number of ozone molecules as its equivalent in the MCM, 223 even though the mechanism is greatly simplified by lumping together similar interme-224 diate species. The intermediate species are named according to their structure, CRI in-225 dex and functional group as explained in the supplementary Table S1. Using this lump-226 ing method, the number of species and reactions in the CRIv2 mechanism is reduced to 227

434 species and 1183 reactions from 4500 species and \approx 12600 reactions in MCMv3.1, covering the degradation of the same 115 emitted NMVOC species without compromising the mechanism's ability to simulate ozone production.

The CRIv2 scheme also underwent several further stages of complexity reduction 231 by lumping together emitted species, with the reduced mechanisms evaluated against the 232 MCM at each stage to preserve ozone forming potential, as described by Watson et al. 233 (2008). The version implemented in the UKCA model is the simplest of these reductions, 234 reduction number five (CRIv2-R5) with 196 species, the same version as implemented 235 into the WRF-Chem model (Archer-Nicholls et al., 2014) and STOCHEM-CRI model 236 (S. Utembe et al., 2010; M. A. H. Khan et al., 2015). This version includes the full CRIv2 237 degradation of isoprene, α -pinene, β -pinene and several aromatic species known to con-238 tribute to secondary organic aerosol (SOA) production, as described by S. R. Utembe 239 et al. (2009). 240

²⁴¹ 3 Implementation of CRIv2-R5 in UKCA

The CRIv2-R5 chemical mechanism was designed for use in boundary layer/urban air quality scenarios. Compared to the StratTrop mechanism used in UKCA, it represents the chemistry of the following primary NMVOCs and their products which either do not exist or are heavily parameterised in StratTrop:

- ²⁴⁶ 1. C4 alkane chemistry via n-butane (C4H10)
- 247
 2. C2-C4 alkene chemistry via ethene, propene and trans-2-butene (C2H4, C3H6 and TBUT2ENE).
- ²⁴⁹ 3. Alkyne chemistry via C2H2.
- 4. Expanded biogenic volatile organic compound (BVOC) chemistry with explicit isoprene (C5H8) degradation and additional monoterpene chemistry via α -pinene and β -pinene (APINENE and BPINENE).
- 5. Aromatic chemistry via benzene, toluene and o-xylene (BENZENE, TOLUENE and OXYLENE).
- 6. Higher oxidised NMVOC chemistry with the addition of Ethanol (EtOH), propanal (EtCHO) and methyl ethyl ketone (MEK).

In addition, it expands on several key chemical processes, for example all peroxy rad-257 icals (RO_2) can react with all other peroxy radicals $(RO_2 + R'O_2)$ and most can form 258 organonitrates from $RO_2 + NO$ reactions. However, CRIv2-R5 lacks key photolysis re-259 actions and chemical species which are important for the chemistry of the upper atmo-260 sphere. The UKCA chemistry-aerosol model is designed to be used in the UK Met Of-261 fice Unified Model (UM) and UK Earth System Model (UKESM1) configurations (Sellar 262 et al., 2019), which simulate the whole atmosphere up to a model top of 85 km in the 263 lower mesosphere. Hence, some changes were made to make the CRIv2-R5 scheme suitable for simulating stratospheric chemical conditions as well as the troposphere. In all, the following development tasks were made to import the CRIv2-R5 mechanism into the 266 UKCA model such that it was suitable for use in a 3D Earth System model configura-267 tion: 268

- 1. Capability for peroxy radical self- and cross-reactions with summed RO_2 pool.
 - 2. Adding stratospheric chemistry, using the same species and reactions as used in the StratTrop mechanism.
- 3. Coupling with the GLOMAP aerosol mechanism.
- 4. Linking with online Fast-JX photolysis.

270 271

5. Wet and dry deposition of species added.

Table 1. Comparison of gas-phase chemical mechanisms: the StratTrop chemical mechanism as described by (Archibald et al., 2020); the CRIv2-R5 mechanism used as the basis of development as described by M. Jenkin et al. (2008); Watson et al. (2008); S. R. Utembe et al. (2009); CRI-Strat which is CRIv2-R5 combined with species and reactions needed for simulating the stratosphere taken from the StratTrop mechanism; and CRI-Strat+GLOMAP-mode which further includes species and reactions needed for aerosol production.

	StratTrop + GLOMAP-mode	CRI-v2-R5	CRI-Strat	CRI-Strat + GLOMAP-mode
No. Species	87	198	219	233
No. Tracers	83	146	167	181
No. Non transported prognostics	4	52	52	52
No. Peroxy radicals [*]	9	47	47	47
No. Emitted species	23	27	27	38
No. Photolysis reactions	60	100	124	128
No. Bimolecular reactions	212	451	536	554
No. Termolecular reactions	25	29	36	39
No. Heterogeneous reactions	8	0	5	8
No. Wet deposited species	34	74	80	83
No. Dry deposited species	41	124	128	131

* Peroxy radicals are transported tracers in the StratTrop mechanism,

Table 1 gives a summary of the chemical mechanisms as integrated into the UKCA 275 model, with the CRI-Strat mechanism with coupling to GLOMAP-mode aerosol being 276 the full mechanism described and analysed in this paper. The CRI-Strat mechanism was 277 implemented into UKCA using the ASAD framework (Carver et al., 1997), meaning it 278 can now easily be ported to other models which share the same framework such as TOM-279 CAT/SLIMCAT chemical transport model (CTM) (Chipperfield, 2006), the offline GLObal 280 Model of Aerosol Processes (GLOMAP) (Spracklen et al., 2006) or the Frontier Research 281 System for Global Change (FRSGC) version of the University of California, Irvine (UCI) 282 global CTM(CTM) (Wild et al., 2000). The full CRI-Strat+GLOMAP-mode mechanism 283 has over twice the number of species (83 to 181) and reactions (305 to 729) as the Strat-284 Trop mechanism. However, the run time is only about 75% longer, hence achieving a greater 285 than doubling of complexity for less than double the cost (details in Section S1 of the 286 supplement). While still computationally expensive and not designed to replace Strat-287 Trop for all model studies, multi-century Earth System model simulations with the mech-288 anism are plausible. 289

The CRIv2-R5 mechanism used to build CRI-Strat was originally optimised against 290 the MCMv3.1 (M. E. Jenkin et al., 2003; Saunders et al., 2003), which drew heavily on 291 kinetic parameters evaluated by the International Union of Pure and Applied Chemistry 292 (IUPAC) Task Group on Atmospheric Chemical Kinetic Data Evaluation (e.g., Atkinson 293 et al. (1997, 2004)). In contrast, the StratTrop scheme (Archibald et al., 2020) draws on 294 a mixture of data from the MCMv3.2 website, the IUPAC Task Group web pages and 295 the NASA JPL Evaluation No. 17 (S. P. Sander et al., 2011). The reaction rate coef-296 ficients for common reactions therefore do not always agree, including for some reactions 297 which are extremely important for tropospheric composition. In some cases CRI-Strat 298 is out of date, in others the mechanisms are simply different as they have different sources. 299 Key differences in rate coefficients are documented in Section S4 of the supplement. The 300 CRI version 2.2 mechanism was recently released (M. E. Jenkin, Khan, et al., 2019) and, 301 among other developments, had some of its reaction rates coefficients revised to match 302

recent assessments to be consistent with MCMv3.3.1 (M. E. Jenkin et al., 2015). Unfortunately, this release was too late to be integrated into the development cycle for the mechanism presented in this paper. However, updating the mechanism to CRI version 2.2 is part of ongoing work.

The remainder of this section describes the developments and implementation of the CRI-Strat mechanism in detail.

3.1 Peroxy Radical Chemistry

Peroxy radicals (RO_2) , a class of short-lived compounds formed during oxidation 310 of VOCs, play a crucial role in the formation of tropospheric ozone (Lightfoot et al., 1992) 311 and secondary organic aerosols (Bianchi et al., 2019; Mcfiggans et al., 2019). While in 312 high-NOx environments they tend to react with NO, forming NO_2 and ozone (M. Jenkin 313 & Clemitshaw, 2000; Monks, 2005), in low-NO_x environments they usually react either 314 with HO2 to form hydroperoxides (ROOH), themselves (self-reactions) or other RO_2 species 315 (cross-reactions) (Tyndall et al., 2001; Orlando & Tyndall, 2012). However, because in-316 dividually simulating the reaction of each RO_2 species with every other RO_2 species for 317 the 47 RO_2 species in CRIv2-R5 would be prohibitively expensive, the approximation 318 described by M. Jenkin et al. (1997) is used, whereby each peroxy radical undergoes a 319 reaction with the summed total of all peroxy radical species (the " RO_2 pool"). The sec-320 ond order reaction rate coefficient is calculated as the geometric mean of the self-reaction 321 rate of the peroxy radical species in question and the $CH_3O_2 + CH_3O_2$ reaction rate. The 322 RO_2 pool is not consumed by these reactions, hence these RO_2 -permutation reactions 323 can be thought of as pseudo-unimolecular reactions with a first order rate coefficient pro-324 portional to the total RO_2 concentration. 325

To enable this chemistry within the framework of the UKCA model, peroxy rad-326 icals were defined as a unique type - that of a non transported prognostic field. Within 327 the ASAD framework and chemical solver, peroxy radicals are treated like other chem-328 ical tracers, but in the rest of the model they are not transported as this is unnecessary 329 for such short lived species. Avoiding the transportation of the 47 peroxy radical species 330 offers substantial savings as some of the largest computational costs of the UKCA model 331 come from tracer transport (Esenturk et al., 2018). The definition of peroxy radicals as 332 a unique type also provides a simple method of tagging the species for their concentra-333 tions to be summed when calculating the total RO_2 pool. The RO_2 pool is defined as 334 a field that can affect the rates of reactions but does not have its concentration directly 335 changed from chemical reactions. Instead, its concentration is calculated for each grid-336 cell and is updated as the concentrations of constituent RO_2 species change. Because 337 ASAD uses an implicit backward euler solver with Newton Raphson iteration (Carver 338 et al., 1997; Wild et al., 2000), the RO_2 pool is recalculated with each iteration of the 330 solver, not just at each timestep. 340

Overall, this method enables efficient representation of peroxy radical chemistry without adding an excessive number of reactions or transported tracers. This framework can be built on further, for example to parameterise the formation of RO₂ accretion products which are important for the formation of highly oxidised organic material (HOM) (Weber et al., 2020).

346

309

3.2 Merging with Stratospheric Chemistry

As both the MCM and CRI mechanisms are designed to simulate polluted boundary layer chemistry, they lack many reactions which are not relevant in the boundary layer but are important in the upper troposphere and stratosphere due to the differing photochemical conditions at different altitudes. The UKCA model is used as part of UKESM1 (Sellar et al., 2019) with a model top at 85 km in the standard configuration, therefore it is essential that the chemical mechanism can adequately represent both troposphericand stratospheric chemistry.

The CRIv2-R5 scheme was merged with the Stratospheric chemistry in UKCA, de-354 scribed by Morgenstern et al. (2009) and updated into the StratTrop mechanism by Archibald 355 et al. (2020), to produce the new CRI-Strat mechanism (see Table 1). In total, 20 species 356 and 121 reactions that are important for stratospheric chemistry were added. Some of 357 the added reactions involve chemical species already present in the CRIv2-R5 mecha-358 nism. In all cases, imported reactions used the same rate coefficients or cross sections 359 as the equivalent reactions in StratTrop. Full details of the CRI-Strat mechanism are 360 given in Tables S2-S5 of the supplement. 361

Photolysis reactions were added for methyl peroxy nitrate $(CH_3O_2NO_2, MeO2NO2$ in CRI-Strat) using the same cross sections as for HO_2NO_2 (Browne et al., 2011). The species $CH_3O_2NO_2$ is not part of the StratTrop mechanism but is an important reservoir for NO_x in cold temperatures and so photolysis of this species is needed to prevent accumulation in the upper troposphere and stratosphere. Reactions forming water vapour with a sizeable flux in the upper atmosphere were adjusted to ensure that H_2O was specified as a product; while chemical production of water vapour is typically neglected in the troposphere it forms an important part of the budget in the stratosphere.

370

394 395

396 397

398

3.3 Coupling with GLOMAP Aerosols

The UKCA model incorporates a modal representation of aerosol size distribution using the two-moment aerosol microphysics scheme from the Global Model of Aerosol Processes (GLOMAP-mode) (Mann et al., 2010) for all aerosol components (sulphate, sea-salt, black carbon and organic carbon, but not currently ammonium nitrate), with the exception of mineral dust which employs a bin scheme as described by Woodward (2001). GLOMAP-mode is fully coupled with the StratTrop mechanism in UKCA (Archibald et al., 2020) as described by (Mulcahy et al., 2020).

To couple the CRI-Strat mechanism to GLOMAP-mode aerosol, the standard Strat-378 Trop couplings have been mimicked wherever possible, except for secondary organic aerosol 379 (SOA) and dimethyl sulphide (DMS) chemistry as discussed below. Because the oxidant 380 fields are different in CRI-Strat, the production of aerosols may differ greatly between 381 CRI-Strat and StratTrop. The species and reactions in CRI-Strat needed for coupling with GLOMAP-mode are marked with an A in supplementary Tables S2-S6. As the main 383 focus of this paper is to evaluate changes in the gas phase due to implementation of the 384 CRI-Strat mechanism, here we only document how the aerosol is coupled. Evaluating 385 the changes to the aerosol fields and their impacts on atmospheric composition and cli-386 mate will be the subject of a follow up paper. 387

Formation of SOA is parameterised in GLOMAP-mode through formation of the SEC_ORG tracer, which represents the extremely low volatility products of biogenic volatile organic compound (BVOC) oxidation. SEC_ORG does not undergo any further chemistry, and is either permanently condensed to the aerosol phase as organic aerosol or is lost via deposition processes. When the StratTrop mechanism is coupled with GLOMAPmode aerosol, SEC_ORG is formed via the reactions:

$$Monoterp + OH \to F * SEC_ORG \tag{2}$$

 $Monoterp + O3 \to F * SEC_ORG \tag{3}$

$Monoterp + NO3 \to F * SEC_ORG \tag{4}$

with rate coefficients equal to the equivalent oxidation reactions of α -pinene with OH, O₃ and NO₃ respectively. The factor F represents the yield of SEC_ORG from Monoterp and is set at runtime to be equal to 26% for all of the above reactions, based on a 13% yield from α -pinene (Tunved et al., 2004; Spracklen et al., 2006) and doubled to account for a lack of SOA production from isoprene and anthropogenic species (Mulcahy et al.,
2020).

To simulate formation of SEC_ORG in the CRI-Strat mechanism, all of the AP-405 INENE and BPINENE initial oxidation reactions produce SEC_ORG at the same yield 406 as the StratTrop reactions with Monoterp, in addition to all of the gas-phase products 407 which are important for ozone chemistry (see supplementary Table S5). This method 408 is not ideal as it does not conserve carbon. However, it is designed to produce a simi-409 lar amount of SEC_ORG as the StratTrop mechanism in order to enable fair compar-410 ison of the gas-phase chemistry between the mechanisms and it serves as a placeholder 411 until a more complete coupling between gas-phase chemistry and aerosol can be devel-412 oped. Working on more explicit couplings between the organic gas-phase chemistry and 413 aerosol routines is ongoing and will build on CRI version 2.2 (M. E. Jenkin, Khan, et 414 al., 2019) and CRIv2.2 with Highly Oxygenated Organic Material (CRI_HOM) (Weber 415 et al., 2020) versions of the mechanism. In the long term, this potential for more real-416 istic chemical coupling between gas-phase organic chemistry and SOA formation is one 417 of the key advantages of using a semi-explicit mechanism such as CRI-Strat over a sim-418 pler mechanism such as StratTrop. 419

Dimethyl sulphide (DMS) is a sulphur containing compound emitted in large quan-420 tities by natural sources and is a critically important source of SO₂, particularly in the 421 preindustrial atmosphere, which can be oxidised to form sulphate aerosols (Andreae, 1990). 422 The DMS chemistry used here is the same as in the CRIv2-R5 implementation in the 423 WRF-Chem model (Archer-Nicholls et al., 2014) and uses the von Glasow and Crutzen 424 (2004) DMS chemistry scheme. This description of multi-generational DMS oxidation 425 is more comprehensive than that used in StratTrop (Archibald et al., 2020) and fits bet-426 ter with the complexity of the rest of the CRI-Strat mechanism. Evaluating and updat-427 ing the DMS chemistry in both the CRI-Strat and StratTrop mechanisms is part of on-428 going work. 429

430

3.4 Coupling with FastJ-X photolysis

The Fast-JX photolysis scheme implemented in UKCA calculates photolysis rates 431 ("j" rates) for each reaction based on experimentally determined cross sections and quan-432 tum yields at a range of wavelength bins (Telford et al., 2013). The CRI-Strat mecha-433 nism includes many more photolysis reactions than StratTrop (128 compared to 60). How-434 ever, most of the added species do not have experimentally determined photolysis cross 435 sections and quantum yields. In the original CRI scheme, used in a box model, the pho-436 tolysis rates for these species were calculated using a two stream isotropic photolysis scheme 437 described by Hayman (1997). When the mechanism was ported to WRF-Chem (Archer-438 Nicholls et al., 2014), photolysis cross section and quantum yield data were adopted from 439 those of species that were already available in WRF-Chem. However, for new species, 440 a box model was used to generate photolysis rate profiles as a function of solar zenith 441 angle. Profiles with similar shapes generated in WRF-Chem were then scaled to match 442 the profiles from the box model. These scaling factors were then applied to the corre-443 sponding cross section and quantum yield data to calculate surrogate photolysis rates for the new species. 445

For this implementation, cross sections already available in UKCA are used for added
photolysis reactions wherever possible, otherwise the surrogate cross sections and scaling factors used in the WRF-Chem implementation are applied. References for the photolysis cross-section data are given in Telford et al. (2013). Details of all photolysis reactions in the CRI-Strat mechanism are provided in supplementary Table S3.

451 **3.5** Wet and dry deposition

There are many new species in the CRIv2-R5 mechanism which are efficiently wet 452 and dry deposited but would otherwise have long chemical lifetimes in the atmosphere. 453 The deposition of new species not included in the StratTrop mechanism is based on the 454 functional group(s) of the species in question. The dry deposition rates and Henry's law 455 coefficients used are described in more detail by O'Connor et al. (2014) and Archibald 456 et al. (2020). Appropriate dry deposition velocities and Henry's Law coefficients were 457 taken from those of existing species in the StratTrop mechanism with a similar functional 458 group or structure; Table 2 shows which surrogate species are used for deposition rate 459 coefficients for each additional species in CRI-Strat. In the case of a species fitting in 460 more than one category, the class with the fastest deposition rates was used. This en-461 sures that all species undergoing deposition are removed at an appropriate rate, but data 462 for specific species can be updated in the future when experimental data becomes avail-463 able. 464

465 4 Methods

466

4.1 Description of Model Setup

The experiments conducted in this study use the UM version 10.9. The model con-467 figuration is based on UKESM1, with 85 vertical levels using terrain-following hybrid height 468 coordinates up to 85 km and a horizontal resolution of $1.25^{\circ} \times 1.875^{\circ}$ (N96) (Walters 469 et al., 2019). All of the simulations are run using nudging of wind and temperature fields 470 and prescribed sea surface temperatures (Telford et al., 2008) to the ERA-interim re-471 analysis product by ECMWF (Dee et al., 2011) to constrain all simulations to observed 472 meteorological evolution so the evaluation can focus on how changes to the chemical mech-473 anism affect atmospheric composition (Zhang et al., 2014). The model simulations run 474 from 1 September 2008 to 1 January 2019, with the analysis period from 1 January 2010 475 to the end of the runs. Well mixed greenhouse gases are not emitted, rather carbon diox-476 ide levels are set as a constant field while methane, nitrous oxide and CFCs are prescribed 477 with constant lower boundary conditions, all at 2014 levels (Archibald et al., 2020). 478

479 4.2 Emissions

The emissions used in this study are those developed for the Coupled-Model In-480 tercomparison Project 6 (CMIP6) (Collins et al., 2017). Anthropogenic and biomass burn-481 ing emissions data for CMIP6 are from the Community Emissions Data System (CEDS), 482 as described by Hoesly et al. (2018), and can be downloaded from http://www.globalchange 483 .umd.edu/ceds/ceds-cmip6-data/. All of the experiments use repeated 2014 emissions 484 as the closest to present day available in the inventories. Anthropogenic emissions are 485 based on the Emission Database for Global Atmospheric Research (EDGARv4.3.1) (http:// 486 edgar.jrc.ec.europa.eu/overview.php?v=431) across the globe, incorporating more 487 detailed regional datasets where available (Hoesly et al., 2018), while biomass burning 488 emissions for the modern period are largely based on Global Fire Emissions Database 489 version 4 with small fires (GFED4s) (Van Der Werf et al., 2017) and merged with other 490 datasets as described by Van Marle et al. (2017). Combination of these datasets onto a unified grid is described by Feng et al. (2020). Offline biogenic emissions are derived 492 from the Model of Emissions of Gases and Aerosols from Nature (MEGAN) version 2.1 493 (Guenther et al., 2012). Oceanic BVOC emissions are included from the POET inven-494 tory (Olivier et al., 2003). 495

Table 3 shows the mappings used to link the CEDS emissions to CRI-Strat. Lumping of raw NMVOC species to CRIv2-R5 speciation is based on methods described in (Watson et al., 2008). We also include equivalent mappings to the StratTrop scheme, as well as total global emissions for 2014 in TgC yr⁻¹.

Compound class	Functional group(s) / molecule	${ m Dry/wet/both}$	CRIv2-R5 species	StratTrop surrogate
Alcohols	HO-	Both	EtOH, i-PrOH, n-PrOH, AROH14, AROH17, ARNOH14, ARNOH17	MeOH
Aldehydes	-CHO	Dry	HOCH2CHO, CARB14, CARB17, CARB11A, UCARB10, UCARB12, NUCARB12, UDCARB8, UDCARB11, UDCARB14, TNCARB26, TNCARB10, TNCARB12, TNCARB11, CCARB12, TNCARB15, TXCARB24, ANHY, TXCARB22 UDCARB15,	MeCHO
Carboxylic Acids	$-CO_2H$	Both	RCOOH25	MeCO2H
Glyoxals	2x -C(O)-	Both	CARB3, CARB6, CARB9, CARB12, CARB15	MGLY
Hydroxy-ketones	-OH and $R(O)R'$	Both	CARB7, CARB10, CARB13, CARB16	HACET
Hydroxy-nitrates	-OH and -ONO2	Dry	HOC2H4NO3, RN9NO3, RN12NO3, RN15NO3, RN18NO3. RU14NO3. RTN28NO3. RTX28NO3	NOSI
Monoterpenes	$\mathrm{C}_{10}\mathrm{H}_{16}$	Dry	APINENE, BPINENE	Monoterp
Nitrate-carbonyls	$-ONO_2$ and $-C(O)$ -	Both	NOA, NUCARB12, RTX24NO3, RTN25NO3, DTV99NO9 DTN99NO9	NALD
:		Ę	KI AZZNU3, KI NZJNU3 Hoten toon dateoon dateoon dateoon	
Peroxides	H00-	Both	НОС2Н4ООН, КИ10ООН, КИ13ООН, КИ16ООН, RN19ООН, RN8ООН, RN110OH, RN14ООН,	EtOOH
			RN1700H, RU1400H, RU1200H, RU1000H, NR11400H NR11300H RN300H RN1300H	
			RN1500H, RN1800H, NRN600H, NRN900H,	
			NRN1200H,RA1300H, RA1600H, RA1900H, PENNSSOCH NERVISSOCH PENNSSOCH PENNSSOCH	
			RTN2400H, RTN2300H, RTN1400H, RTN1000H,	
Peroxv Acids	-CO3H	Both	RTX2800H, RTX2400H, RTX2200H, NRTX2800H E4C03H. H0CH2C03H	MeCO3H
Peroxyacyl Nitrates	-PAN	Dry	PHAN, RU12PAN, RTN26PAN	PAN

Table 2. CRI-Strat species which undergo wet or dry deposition but do not exist in StratTrop do so using coefficients based on surrogate StratTrop species.

aw CMIP6 NMVOC emissions to CRI-Strat and StratTrop mechanisms, with total emitted carbon mass for 2014 from Anthropogenic,	genic sources.
Mapping of raw (rning and biogenic
Table 3.	biomass bu

Raw emission classes	Anthropogenic TgC yr ⁻¹	Biomass Burning TgC yr ⁻¹	$\begin{array}{c} {\rm Biogenic} \\ {\rm TgC} \ {\rm yr}^{-1} \end{array}$	CRIv2-R5 Species	Strat Trop Species
VOC1: Alcohols	0.4 3.	3.5 0.1	$\frac{48.5}{9.5}$	MeOH EtOH	MeOH to MeOH
VOC2: Ethane VOC7: Ethene VOC9: Ethyne	5.3 4.9 3.1	2.8 4.1 1.1	$\begin{array}{c} 1.0\\ 25.8\\ \end{array}$	C2H6 C2H4 C2H2 C2H2	C2H6 to C2H6 to C2H6
VOC3: Propane VOC8: Propene	5.5 3.0	0.6 3.5	$1.0 \\ 14.$	C3H8 C3H6	C3H8 to C3H8
VOC4-6: Butanes and higher alkanes	4.8	0.4	0.1	C4H10	N/A
VOC10: Isoprene		0.6	588	C5H8	C5H8
VOC11: Monoterpenes	1	1.2	94.7	67% to APINENE* 33% to BPINENE*	Monoterp [†]
VOC12: Other Alkenes and Alkyenes	6.5	0.8	2.6	TBUT2ENE	N/A
VOC13: Benzene VOC14: Toluene VOC15-17: Xylenes and higher aromatics	6.1 7.0 3.1	2.0 3.9 1.1		BENZENE TOLUENE •XYLENE	N/A N/A N/A
VOC21: Formaldehyde VOC22: Other Aldehydes	1.0 0.5 0.6	2.1 3.4 0.8	1.8 10.0 2.0	HCHO MeCHO EtCHO	HCHO MeCHO to MeCHO
VOC23: Ketones	1.5	$1.1 \\ 0.9$	$22.9 \\ 0.5$	Me2CO MEK	Me2CO to Me2CO
VOC24: Acids	4.4	0.5 7.1	$1.4 \\ 1.9$	HCOOH MeCO2H	N/A^{\ddagger}
Total CRI: Total StratTrop:	70.5 27.9	40.6 23.9	900.6 710.6		
* Two to one split between APINENE and [†] In the StratTrop mechanism Monoterp of Monoterp are not included in the total NM [‡] Acids have historically not been mapped	[BPINENE emissi nly forms Sec_Org IVOC emissions fc to StratTrop mec	ions according to G ¹ and does not contr or the StratTrop me hanism in UKESM1	uenther et al ibute to ozoi chanism. (Archibald	. (2012) and Sindelarova ϵ ne formation. Therefore, ϵ et al., 2020), even though	et al. (2014). emissions mapped to HCOOH and MeCO2H

are existing species.

The CRI-Strat mechanism utilises a wider breadth of input NMVOC emissions, us-500 ing almost all of the available data from the CEDS emission database, therefore has a 501 greater amount of total carbon mass emitted compared to an equivalent StratTrop run. 502 For anthropogenic emissions, the most significant changes are due to the addition of aro-503 matic species and C_4 alkane/alkene emissions. For biogenic emissions, the largest change 504 is that monoterpenes are mapped to the chemically reactive APINENE and BPINENE 505 tracers in CRI-Strat, which undergo oxidation and contribute to ozone formation, whereas 506 in StratTrop they are mapped to the MONOTERP tracer which is only considered as 507 a precursor of SOA and does not contribute to ozone production). The total NMVOC 508 emissions for the year 2014 using the CEDS emissions inventory are given in Table 3. 509

510

4.3 Model Simulations

Two base simulations are conducted, labelled StratTrop and CRI-Strat (Table 4). 511 These use the emissions associated with their respective mechanism, as described in Ta-512 ble 3. The StratTrop simulation uses a slightly modified version of the mechanism, where 513 the reactions of $NO_3 + DMS$ and $NO_3 + Monoterp$ were adjusted so that they conserved 514 nitrogen, when previously they did not. These changes were made to enable a fair com-515 parison between the mechanisms for Section 5.6. These changes have a minimal impact 516 on the overall chemical composition in StratTrop and are explained in detail in Section S3 517 of the Supplement. 518

The treatment of emissions can be seen as an intrinsic part of a chemical mecha-519 nism. However, the emissions of additional NMVOC species in the CRI-Strat simula-520 tion which are not represented in StratTrop pose a dilemma when comparing the two 521 mechanisms: are the differences between the simulations due to the different approaches 522 in representing chemistry, there being more reactive carbon available in CRI-Strat, or 523 a combination of these factors? Two additional simulations were therefore designed to 524 better understand the effects of additional NMVOC emissions. A CRI-Strat run was con-525 ducted with identical emissions to the StratTrop base run (CRI_Emiss_ST) to isolate the 526 changes just due to the chemical mechanism without any changes in emissions. The fi-527 nal CRI-Strat simulation only uses emissions from NMVOC emission classes that are also 528 used by StratTrop, but these are mapped to the appropriate CRI-Strat species, hence 529 it can be used to identify the effects of changing the speciation of NMVOC emissions with-530 out increasing the total carbon mass. The key difference is that C_2 and C_3 species are 531 mapped into different species, hence this scenario is called speciated C2-C3 emissions (CRI_Emiss_C2C3). 532 For example, emissions of ethane, ethene and ethyne are mapped to C2H6, C2H4 and 533 C2H2 respectively in CRI_Emiss_C2C3, rather than being lumped to C2H6 as in the Strat-534 Trop and CRI_Emiss_ST scenarios. The MONOTERP tracer is unreactive in StratTrop 535 (it can only be oxidised to form SEC_ORG and cannot contribute to ozone production), 536 therefore all monoterpene emissions are mapped to MONOTERP in CRI_Emiss_ST and 537 CRI_Emiss_C2C3, with the respective reactions copied over from StratTrop. The sim-538 ulations are summarised in Table 4. 539

540 5 Results

In this paper, we focus on understanding how changes in chemistry affect gas-phase 541 species which are important for describing the global tropospheric composition as drivers 542 to climate: ozone, carbon monoxide, hydroxy radicals, methane lifetime and nitrogen ox-543 ides. We present a short evaluation against surface and remote sensing products, but the 544 main focus is on comparing the performance of the CRI-Strat with the StratTrop mech-545 anism, which is already well evaluated (Archibald et al., 2020), and on understanding 546 how these changes are sensitive to the treatment of NMVOC emissions. While the changes 547 in chemistry and oxidant fields are also expected to affect formation of aerosols, these 548 influences will be explored in more detail in follow up studies. 549

Scenario Name	Chemical Mechanism	Emissions	Purpose
CRI-Strat	CRI-Strat	Standard CRI	Base CRI scenario
StratTrop	StratTrop	Standard ST	Base StratTrop scenario
CRI_Emiss_ST	CRI-Strat	Standard ST	Isolating effect of chemical representation
CRI_Emiss_C2C3	CRI-Strat	Speciated C2-C3 emissions	without changes to emissions Isolating effect of NMVOC speciation without changing total emitted mass

 Table 4.
 Summary of simulations.

550 5.1 Model evaluation

551

5.1.1 Surface ozone

Lowest model level ozone concentrations from the base CRI-Strat and StratTrop model simulations are evaluated against the global dataset of rural surface ozone sensors compiled for the Tropospheric Ozone Assessment report (TOAR) (Schultz et al., 2017). The TOAR dataset compiles ozone measurements from surface station-based observations around the globe including data from regional and national air quality networks, as well as larger global collaborations.

The data used in this work are the gridded surface ozone datasets, which include 558 mean surface ozone concentrations on a $2^{\circ} \times 2^{\circ}$ global grid. Schultz et al. (2017) also 559 provide gridded station means separated by urban or rural classification, based on the 560 population density, nightlight intensity, and OMI satellite NO₂ column. The model low-561 est level ozone concentration was evaluated against the rural mean surface ozone grid-562 ded dataset, because the relatively large grid size in the UKCA model $(1.25^{\circ} \times 1.875^{\circ})$ 563 approximately 135 km horizontal grid spacing at the equator) means that urban-scale 564 chemistry is not well resolved. In all comparisons with TOAR gridded data, the model 565 output was regridded to the same $2^{\circ} \times 2^{\circ}$ grid as the TOAR data before analysis. 566

Figure 1 shows comparison of CRI-Strat and StratTrop model simulations with TOAR 567 observations across the globe, grouped by region. Further plots comparing model out-568 put with TOAR data are included in the supplement Figures S6-S9. The CRI-Strat sim-569 ulation has consistently higher surface ozone values across almost the entire world, with 570 this increase most pronounced in populated regions such as Europe and East Asia. In 571 more remote regions, the difference between the CRI-Strat and StratTrop simulations 572 is much smaller. The CRI-Strat and StratTrop simulations follow similar seasonal trends, 573 showing this variation is driven more by the parent model and shared traits, such as sea-574 sonal variation in emissions. In many regions, both simulations are low compared to ob-575 servations in winter months and high in summer months. Due to CRI-Strat having higher 576 ozone in general, the global summer high bias is greater in CRI-Strat (+12.6 ppbv) com-577 pared to StratTrop (+7.4 ppbv), but has a smaller negative bias in winter (-4.3 ppbv com-578 pared to -9.2ppby). These results indicate that the CRI-Strat mechanism has a higher 579 ozone production efficiency than StratTrop. 580

The high bias in surface summer ozone and low bias in winter are likely due to structural issues in using a coarse, global model as has been documented elsewhere (Young et al., 2013, 2018; Archibald et al., 2020). For example, the coarse horizontal resolution results in the emissions being smoothed and less heterogeneous, particularly around urban areas and large point sources, leading to greater mixing of NO_x and NMVOCs and ozone production, particularly in summer (Wild & Prather, 2006; Stock et al., 2014; Fenech et al., 2018). The low bias in winter, when local production is lower, may be due to in-





sufficient long range transport of ozone or loss due to deposition being too great. In some 588 regions, such as East Asia, the seasonal cycle in the model simulations is out of phase, 589 with the model runs showing peak ozone in the summer when the observations are at 590 their lowest. This is likely indicative of missing model processes, such as heterogeneous 591 chemistry or poor representations of local weather processes, such as monsoon cycles. 592 Any biases caused by such structural weaknesses in the parent model are common to both 593 mechanisms, but may be accentuated in CRI-Strat due to greater ozone production when 594 photochemical conditions allow. Given these issues, the higher bias in CRI-Strat com-595 pared to the TOAR surface ozone observations likely more clearly reveal other biases in 596 the model which compensate for errors associated with the less accurate description of 597 the underlying chemistry in StratTrop. 598

599

618

5.1.2 Surface carbon monoxide

Modeled carbon monoxide (CO) surface fields are evaluated against the National 600 Oceanic and Atmospheric Administration (NOAA) climate Monitoring and Diagnostics 601 Laboratory (CMDL) dataset (Pétron et al., 2002), as shown in Figure 2. The NOAA data 602 are derived from regular in situ flask samples and screened for local pollution events. The 603 observational data evaluated here are calculated as a monthly average climatology derived from observations made between 2010-15. The StratTrop simulation shows a low 605 bias at most sites in the Northern hemisphere, which are more influenced by anthropogenic 606 pollution, but is close to observed values in the tropics and over the southern hemisphere 607 which are more remote and dominated by biogenic and biomass burning emissions. These 608 effects have been documented elsewhere (Archibald et al., 2020). The CRI-Strat simu-609 lation has greater CO at all sites and over all seasons compared to StratTrop. This re-610 duces the size of the negative bias at northern hemisphere sites but creates a positive 611 bias over the remote southern hemisphere sites. The seasonal trends in both simulations 612 are similar, showing that these are more sensitive to model dynamics and seasonal vari-613 ation in emissions than chemical mechanism. The higher CO in CRI-Strat is due to sec-614 ondary production of CO from oxidation of the additional NMVOCs but this has mixed 615 effects on the comparison with observations, improving biases in the northern hemisphere 616 whilst making them worse in the southern hemisphere. 617

5.1.3 Tropospheric ozone column

Although surface ozone is a pressing concern for air quality, the influence of ozone 619 on climate is more dependent on ozone in the upper troposphere and the tropospheric 620 ozone burden (Bowman et al., 2013). We use two different monthly mean gridded satel-621 lite data products to evaluate tropospheric column ozone (TCO) in the CRI-Strat and 622 StratTrop simulations. The OMI-MLS TCO monthly gridded data, determined by sub-623 tracting the Microwave Limb Sounder (MLS) stratospheric column ozone (Waters et al., 624 2006) from the Ozone Monitoring instrument (OMI) total column ozone (Dobber et al., 625 2006), is available between 60° S- 60° N with a horizontal resolution of $1^{\circ} \times 1.25^{\circ}$, as de-626 scribed by Ziemke et al. (2006, 2019). For comparison with OMI-MLS data (2010-2018), 627 the modelled TCO is calculated by vertically integrating the model ozone between the 628 surface and the tropopause (defined as 380K + 2 PV; Hoerling et al. (1993). The OMI 629 data (Boersma et al., 2007) were produced by the Remote-Sensing Group at Rutherford 630 Appleton Laboratory using a profile retrieval scheme developed first for GOME-2 (Miles 631 et al. 2015). Individual profiles were gridded on a monthly basis with a horizontal res-632 olution of $1.5^{\circ} \times 1.5^{\circ}$ and a correction applied in each layer for bias with respect to ozoneson-633 des which had been derived as a function of month of year and latitude (R. Siddans pri-634 vate communication). For comparison with OMI data (2010-2017) in the surface-450 hPa 635 and 450-170 hPa layers, we use monthly, gridded averaging kernels and a priori infor-636 mation to minimise vertical sampling differences between OMI and UKCA. Although 637 potential issues with using monthly mean rather than individual averaging kernels can 638



Figure 2. Average surface CO concentrations from CMDL network (black) compared with CRI-Strat (red) and StratTrop (blue) model simulations, showing average seasonal variation, correlation coefficient and mean bias error over 2010-2018 evaluation period.

arise for certain species and instruments (von Clarmann & Glatthor, 2019), agreement
 between model and observation are found to be improved substantially through appli cation of monthly mean averaging kernels in this analysis.

In Figure 3, average tropospheric ozone column is compared between the OMI-MLS 642 satellite product and the CRI-Strat and StratTrop simulations. Differences in tropospheric 643 ozone column between each of the model simulations are presented in supplementary Fig-644 ure S10. The tropospheric ozone column is remarkably similar in the CRI-Strat and Strat-645 Trop simulations, both showing a similar high bias over the tropics and a low bias at high 646 latitudes (Figure 3 (c, e)). The CRI-Strat and StratTrop simulations also have similar 647 tropospheric ozone burden between 60° S and 60° N (303 Tg and 308 Tg respectively, com-648 pared to 301 Tg from OMI-MLS). 649

The seasonal zonal mean ozone in Figure 3 (b, d, f) is also more similar in the two 650 model simulations compared to OMI-MLS. Both mechanisms show a clear high bias over 651 the tropics throughout the year, with a strong peak between 0 and 30° N between March 652 and June. In contrast, the satellite product has higher ozone column values around 40° N 653 in June-July, with a slightly smaller peak around 30°S in October-November, a pattern 654 not represented in the model simulations. Considering the large differences in surface 655 ozone between the two mechanisms (Figure 1) it is perhaps surprising how similar the 656 total tropospheric ozone is between the two mechanisms, and this is probably a sign that 657 common model deficiencies (such as errors in emissions or transport) cause similar bi-658 ases in both simulations. However, these tropospheric column comparisons may hide sig-659 nificant differences in vertical distribution of ozone. 660

The high bias in tropospheric ozone column seen over equatorial regions in both 661 model simulations (Fig. 3 c-f) is absent in the lower troposphere (Fig. 4 c, e), but ap-662 pears in the upper troposphere (Fig. 4 d, f). As the mechanisms show a similar bias in this region of the atmosphere, similar structural weaknesses must be causing the bias in 664 both simulations, likely contenders being errors in lightning NOx emissions (Banerjee 665 et al., 2014) or convective transport (Hoyle et al., 2011). CRI-Strat has higher ozone columns 666 than StratTrop above polluted regions such as India and downwind of Europe, mostly 667 in the lower troposphere, but has reduced ozone compared to StratTrop in the less pol-668 luted southern hemisphere (4 g, h). 669

670 671

683

5.2 Comparison of the ozone budget and processes in CRI-Strat and Strat-Trop

In this section, we investigate why tropospheric chemical composition differs be-672 tween the CRI-Strat and StratTrop mechanisms. This is assessed using the base CRI-673 Strat and StratTrop simulations, and the two CRI-Strat simulations with modified NMVOC 674 emissions (CRI_Emiss_ST and CRI_Emiss_C2C3). Collectively, these simulations allow 675 us to explore the impact of changing chemistry without changing emissions, the differ-676 ence attributable to explicit speciation of emissions, and the impact of the additional chem-677 istry of C4 alkanes and alkenes, aromatics and monoterpene. Given that the StratTrop 678 mechanism does not use a large fraction of the reactive carbon mass included in the in-679 put inventory, this analysis may also illustrate the causes of certain biases in the Strat-680 Trop mechanism and how these might be improved if it were to be modified to use a wider 681 selection of NMVOC species, informing future development of this mechanism. 682

5.2.1 Ozone comparison

Boundary layer ozone is much higher over land in CRI-Strat compared to Strat-Trop, but is lower over remote oceans (Figure 5). Ozone levels are mostly lower in CRI_Emiss_ST and CRI_Emiss_C2C3 than StratTrop, but are higher over polluted regions. The variability in ozone is greater in CRI_Emiss_C2C3 than in CRI_Emiss_ST, and there are higher



Figure 3. Map of tropospheric ozone column (DU) from OMI-MLS (a) and annual variation in zonal means (b), averaged between 2010-2018. Difference in tropospheric ozone column between CRI-Strat and OMI-MLS (c, d), between StratTrop and OMI-MLS (e, f), and between CRI-Strat and StratTrop noting different colour scale (g, h). Burdens given above panels a, c and e are calculated by summing ozone over the troposphere between 60°S to 60°N as this is the range covered by the OMI-MLS product.



Figure 4. Ozone column between surface and 450 hPa from OMI satelite product as map (a) and from 450 to 170 hPa (b), averaged between 2010-2017. Difference in ozone column between CRI-Strat and OMI over respective pressure ranges (c, d), between StratTrop and OMI (e, f), and between CRI-Strat and StratTrop, noting different colour scale (g, h). Burdens are given over the respective pressure ranges for between 60°S to 60°N then over 90°S to 90°N.



Figure 5. Mean ozone averaged over lower 1 km of the atmosphere in StratTrop (a) and CRI-Strat (b). Absolute difference (c-e) and relative difference (f-h) in ozone over lower 1 km of atmosphere between CRI-Strat and StratTrop (c, f), CRI_Emiss_ST - StratTrop (d, g), and CRI_Emiss_C2C3 - StratTrop (e, h).

ozone levels over highly populated regions such as the Indo-Gangetic plain but lower levels in remote regions. This is due to CRI_Emiss_C2C3 having emissions of alkenes such as C_3H_6 which are more reactive and have a higher ozone production efficiency than equivalent alkanes such as C_3H_8 , but have shorter lifetimes and contribute less to ozone production downwind of sources (M. E. Jenkin et al., 2002).

Looking at the vertical distribution of ozone, high ozone levels in CRI-Strat are lo-693 calised to the lower atmosphere in the northern hemisphere (Figure 6 (a, d)). Ozone con-694 centrations are lower in the upper tropical troposphere and across the southern hemi-695 sphere. CRI-Strat has higher ozone in the lower stratosphere, but these differences are 696 relatively small (< 2%) and the largest fractional differences are in the lower atmosphere. 697 In the CRI_Emiss_ST and CRI_Emiss_C2C3 scenarios, ozone levels are lower through-698 out most of the atmosphere, and the only region that shows similar zonal average ozone is the boundary layer between 30N and 60N (Figure 6 (b,c, e, f)). This demonstrates that 700 the CRI-Strat mechanism is more efficient at producing ozone near emission sources, but 701 that losses are greater in remote regions of the atmosphere. 702

The relationship between surface ozone and emissions is shown in the ozone isopleths in Figure 7. Ozone concentrations in all of the simulations are lower in NO_x -limited and VOC-limited regimes (in the bottom-right and top-left of each panel respectively), and drop slightly faster in CRI-Strat than StratTrop as NO_x emissions increase. Peak ozone concentrations in StratTrop occur in regions with lower VOC and NO_x emissions,



Figure 6. Zonal mean absolute difference (a-c) and percentage difference (d-f) in ozone between CRI-Strat and StratTrop (c, f), CRI_Emiss_ST - StratTrop (d, g), and CRI_Emiss_C2C3 -StratTrop (e, h). Black lines show average tropopause height over simulation period.



Figure 7. Ozone isopleths showing average surface ozone levels with respect to surface emissions of VOCs and NOx in StratTrop (a), CRI-Strat (b), CRI-Emiss_ST (c) and CRI-Emiss_C2C3 (d) model simulations. Isopleths have been drawn by mapping average surface ozone concentrations from each gridcell of each of the model simulations with NO and NMVOC emissions at each gridcell, then interpolating results to give a smooth field. Stippling shows regions of the phase space which have sufficient data from the model output.

between 0.01-0.1 mg C m⁻²hr⁻¹ and 0.01-0.1 mg N m⁻²hr⁻¹ respectively), and then drop 708 off at high emission regions (Fig. 7 (a)). The CRI-Strat simulation has the highest ozone 709 levels, with the increased NMVOC emissions spurring ozone production in more regions 710 with high NO_x emissions (Fig. 7 (b)). The CRI_Emiss_ST scenario shows a similar dis-711 tribution to StratTrop but peak ozone levels occur in regions with higher emissions (Fig. 712 7 (c)). CRI_Emiss_C2C3 has peak ozone at even higher emission levels and the peak in 713 ozone covers a broader region of the phase space, demonstrating that increased speci-714 ation of NMVOC emissions allows for greater ozone production in polluted regions as 715 more is emitted as short lived, reactive compounds (Fig. 7 (d)). 716

717 5.2

5.2.2 Production and loss of ozone

To investigate these differences in ozone between the simulations, we present a full budget analysis of tropospheric ozone. Tropospheric production and loss of ozone is calculated using diagnostics which track the odd oxygen family, including NO₂ and its reservoir species, collectively known as O_x (Wang et al., 1998):

$$O_x = O_3 + O + O(^1D) + NO_2 + 2NO_3 + 3N_2O5 + HONO_2 + HO_2NO_2 + PANs.$$
(5)

This definition of O_x is based on the principle that the rate limiting step for ozone production following NO₂ photolysis is the conversion from NO to NO₂. Hence production (P_{Ox}) is primarily via the pathways HO₂ + NO, CH₃O₂ + NO and R'O₂ + NO (where R'O₂ is the sum of all peroxy radicals apart from CH₃O₂). Chemical loss of ozone (L_{Ox}) is defined as the sum of all chemical pathways which result in net loss of ozone, primar⁷²⁸ ily via reaction of $O(^{1}D)$ with water vapour and catalytic loss of O_{3} with HO_{x} but also ⁷²⁹ via minor reaction pathways involving NO₃. Net ozone production is therefore defined ⁷³⁰ as the difference between chemical production and loss:

731

737

740

$$Net_{Ox} = P_{Ox} - L_{Ox} \tag{6}$$

 732 O_x is also lost via deposition (D_{Ox}), both directly through dry deposition of O₃ and indirectly via deposition of NO₂ and its reservoir species. The final part of the budget is from transfer of ozone from the stratosphere to the troposphere (S_{Ox}), which can be inferred from the excess loss of ozone assuming no long-term changes in tropospheric burden.

$$S_{Ox} = L_{Ox} + D_{Ox} - P_{Ox} \tag{7}$$

Finally, the lifetime of O_x in the troposphere is calculated by dividing the total tropospheric burden by the total loss via chemical sinks and deposition (Young et al., 2018).

$$\tau_{Ox} = B_{Ox} / (L_{Ox} + D_{Ox}) \tag{8}$$

In the CRI-Strat simulation, ozone production is much higher than StratTrop in 741 the lower atmosphere, but also in most of the troposphere between $60^{\circ}S$ and $90^{\circ}N$ (Fig. 742 8 (a) and supplementary Figure S11). The only region where StratTrop has faster pro-743 duction is in the southern upper tropical troposphere. In CRI_Emiss_ST and CRI_Emiss_C2C3, 744 ozone production is even more concentrated in the boundary layer, with much less pro-745 duction than StratTrop in the upper troposphere (Fig. 8 (b, c)). However, all simula-746 tions using the CRI-Strat mechanism have greater ozone loss over most of the troposphere 747 (Fig. 8 (d-f)), so that the main region of greater net ozone production is in the bound-748 ary layer and there is a layer of net loss immediately above (Fig. 8 (g-i)). As shown in 749 Supplementary Figure S12, the difference in distribution of L_{Ox} is almost identical to 750 the flux through the $O(^{1}D) + H_{2}O$ reaction, one of the reactions that has different re-751 action rate coefficients between the mechanisms. This reaction drives a large component 752 of the ozone changes between the mechanisms. 753

Table 5 gives an overview of the tropospheric ozone budget, production and loss 754 terms. While CRI-Strat and StratTrop have similar total tropospheric ozone burdens 755 of 331.8 Tg and 336.8 Tg respectively, the burdens of CRI_Emiss_ST and CRI_Emiss_C2C3 756 are both around 30 Tg lower. Compared to the TOAR model intercomparison (Young 757 et al., 2018), the tropospheric ozone burdens in the StratTrop and CRI-Strat simulations 758 are within the model interquartile range (320-370 Tg), whereas CRI_Emiss_ST and CRI_Emiss_C2C3 759 are below that range. Gaudel et al. (2018) calculate total ozone burdens between 333 760 and 345 Tg for the 2010-2014 period from satellite products which can observe all lat-761 itude bands, overlapping with StratTrop and slightly above CRI-Strat but much higher 762 than CRI_Emiss_ST and CRI_Emiss_C2C3. Ozone production efficiency (OPE, defined 763 as moles of O_x produced per mole of NO_x emitted) is also higher in CRI-Strat (31.4) than 764 StratTrop (27.2). 765

The lifetime of ozone in all of the simulations using CRI-Strat chemistry is much 766 shorter (17.3-17.5 days) than with StratTrop (19.8 days), hence the lower ozone in CRI_Emiss_ST 767 and CRI_Emiss_C2C3 likely reflects the shorter lifetime of O_x . However, the CRI-Strat 768 simulation has a similar ozone burden to StratTrop because increased production com-769 pensates for the increased losses. In fact, the lifetime of ozone in all of these simulations 770 is short compared to previous analysis (e.g. 22.3 ± 2.0 days from Young et al. (2013), 771 although these were for the year 2000 rather than 2014). Note also that the lifetime cal-772 culation is dependent on the definition of O_x , for example Bates and Jacob (2020) find 773 O_3 lifetimes of approximately 75 days as they do not consider the $O(^1D) + H_2O$ reac-774 tion to cause net loss of O_x . The short lifetimes could partly explain why both CRI-Strat 775 and Strat-Trop have lower ozone column values at high latitude than OMI-MLS (Fig-776 ure 3), as insufficient ozone is transported from the tropics, where production is high-777 est. 778



Figure 8. Zonal mean difference in chemical production of Ox between CRI-Strat and Strat-Trop (a), CRI_Emiss_ST and StratTrop (b), and CRI_Emiss_C2C3 and StratTrop (c) over 2010-2018 simulation period. Zonal mean difference in chemical loss of Ox between CRI-Strat and StratTrop (d), CRI_Emiss_ST and StratTrop (e), and CRI_Emiss_C2C3 and StratTrop (f). Zonal mean difference in net chemical production of Ox between CRI-Strat and StratTrop (g), CRI_Emiss_ST and StratTrop (h), and CRI_Emiss_C2C3 and StratTrop (i). Black lines show average tropopause height over simulation period.

		CRI-Strat	$\operatorname{StratTrop}$	CRI_Emiss_ST	CRI_Emiss_C2C3
O ₃ burden (Tg)		331.9	336.8	308.0	306.8 17 4
O _x meume (days) NMVOC emissions (Tg C vear ⁻¹		1012	762	762 762	16.4 762^{\dagger}
OPE (mole _{O₃} mole _{NOx})		31.4	27.2	28.7	28.9
Chemical production (P_{Ox})	Total	6624	5725	6057	6092
$(Tg O_3 year^{-1})$	$HO_2 + NO$	4152 (62.7%)	$3853 \ (67.3\%)$	$3886\ (64.2\%)$	$4001 \ (65.7\%)$
	$CH_3O_2 + NO$	1540(23.3%)	$1285\ (22.5\%)$	$1452\ (24.0\%)$	$1419\ (23.3\%)$
	$ m R'O_2+NO$	882 (13.3%)	545 (9.5%)	$676\ (11.2\%)$	629 $(10.3%)$
	$Other^*$	64.5(0.7%)	$41.3\ (0.7\%)$	41.5(0.7%)	$43.2\ (0.7\%)$
Chemical loss (Lox)	Total	5853	5128	5355	5380
$(Tg O_3 year^{-1})$	$O(^{1}D) + H_{2}O$	3196(45.5%)	2660(42.9%)	$3022 \ (47.1\%)$	$3005 \ (46.6\%)$
	$HO_2 + O_3$	1713 $(24.4%)$	1596(25.7%)	1498(23.4%)	$1513\ (23.5\%)$
	$OH + O_3$	708 (10.1%)	$714 \ (11.5\%)$	$667\ (10.4\%)$	$680 \ (10.6\%)$
	$O_3 + Alkene$	160(2.3%)	96.5(1.6%)	115(1.79%)	129(2%)
	$Other^{\ddagger}$	76.3(1.1%)	$61.5 \ (1.0\%)$	$53.1\ (0.8\%)$	$52.6\ (0.8\%)$
Deposition (D _{Ox})	Total	1164	1081	1061	1068
	$O_3 dry dep$	974 (13.9%)	896 (14.4%)	$872\ (13.6\%)$	878~(13.6%)
	NO_{y} dep	190.3(2.7%)	$185\ (3.0\%)$	189(2.95%)	190(2.94%)
Inferred STT (Tg O ₃ year–1)		394	483	361	355

Table 5. Overview of Ox burden, lifetime, ozone production efficiency (OPE), chemical production, chemical loss, deposition and inferred stratosphere to tropo-

manuscript submitted to Journal of Advances in Modeling Earth Systems (JAMES)

The ozone production and loss terms for CRI-Strat (6624 and 5853 Tg yr^{-1} respec-779 tively) are both much higher than for StratTrop (5725 and 5128 Tg yr⁻¹), and are higher 780 than most models from the TOAR assessment (Young et al., 2018). It has been noted 781 before that models with representation of higher aromatics and monoterpenes have high 782 P_{Ox} and L_{Ox} terms (Porter et al., 2017), but the calculation of the net chemical tendency 783 $(P_{Ox} - L_{Ox})$ shows that the CRI-Strat results in a greater overall propensity to form 784 ozone in spite of the emissions of NMVOCs (c.f. StratTrop and CRI_Emiss_ST exper-785 iments Table 4). The P_{Ox} and L_{Ox} terms in CRLEmiss_ST and CRLEmiss_C2C3 are 786 very similar to each other, in both cases higher than StratTrop and lower than the base 787 CRI-Strat run. There is a general trend for "hotter" chemistry - more ozone is produced 788 and lost - in all of the simulations using the CRI-Strat chemical mechanism. We can also 789 say that the addition of higher NMVOC emissions has a larger impact on the ozone bud-790 get (CRI-Strat vs CRI_Emiss_ST and CRI_Emiss_C2C3) than the improved speciation 791 of NMVOC emissions (CRI_Emiss_ST vs CRI_Emiss_C2C3). 792

The increase in ozone production in CRI-Strat occurs over all channels, but the largest 793 relative increase is via $R'O_2 + NO$ (882 Tg yr⁻¹), which is over 50% larger in CRI-Strat 794 than in StratTrop (545 Tg yr⁻¹), partly because there is more $R'O_2$ in CRI-Strat. The 795 CRI_Emiss_ST simulation shows its largest increase in production compared to Strat-796 Trop via the $R'O_2 + NO$ channel, as well as the $CH_3O_2 + NO$ channel, whereas in CRI_Emiss_C2C3 797 it is via the $HO_2 + NO$ channel. Note though that the increased production via $HO_2 + NO$ 798 and $CH_3O_2 + NO$ is also linked to oxidation of larger NMVOCs, as many reactions pro-799 duce secondary HO₂ and CH₃O₂ as larger molecules are oxidised. 800

Chemical loss of O_x via the $O(^1D) + H_2O$ channel is greater in CRI-Strat (3196 Tg yr⁻¹) 801 than in StratTrop (2660 Tg yr⁻¹). In addition, the CRLEmiss_ST and CRLEmiss_C2C3 802 scenarios have similar losses via this channel (3022 Tg yr^{-1} and 3005 Tg yr^{-1} respec-803 tively) to the base CRI-strat simulation, scaling linearly with ozone burden. The $O(^{1}D) + H_{2}O(^{1}D)$ 804 pathway is the primary driver behind why O_x lifetime is much shorter in the CRI-Strat 805 chemical mechanism compared to StratTrop. This is down to both the reaction rate co-806 efficients for the hydroxyl radical producing $O(^{1}D) + H_{2}O \rightarrow 2OH$ reaction being greater 807 in CRI-Strat than in StratTrop, and the stabilising reaction $O(^{1}D) + M \rightarrow O(^{3}P) + M$ 808 is smaller, which collectively result in excited $O(^{1}D)$ being about 25% more likely to re-809 act with H_2O than stabilise with M for a given concentration of water vapour (see Fig-810 ure S2 and Section S4 in the supplement for details). 811

5.3 Hydroxyl and Hydroperoxyl Radicals

812

The hydroxyl radical (OH) is the major oxidant in the troposphere for almost all 813 NMVOC species. Together with the hydroperoxyl radical (HO_2) it forms the HO_x fam-814 ily $(HO_x = OH + HO_2)$. Zonal means of OH and HO₂ from the StratTrop simulation and 815 differences to other simulations are shown in Figure 9. All of the simulations with CRI-816 Strat chemistry show a bimodal pattern in OH of high levels in the lower troposphere 817 and low levels in the upper troposphere compared to StratTrop (Fig. 9 (c-e)), and a pat-818 tern of more HO_2 in both the lower and upper troposphere (Fig. 9 (f-h)). Total HO_x , 819 which is roughly equal to HO_2 , is much higher in all of the simulations using CRI-Strat 820 chemistry than StratTrop (Fig. 9 (f-h)), which is unsurprising given the main source is 821 the $O(^{1}D) + H_{2}O$ reaction which we already know has a much higher flux in CRI-Strat 822 than StratTrop (Table 5). 823

In the lower atmosphere, which has higher O_3 in the CRI-Strat simulation and is abundant with water vapour, we have greater HO_x production and more OH and HO_2 . There are also more O_3 + Alkene reactions in CRI-Strat, which produce OH, and HO_2 is produced as a byproduct of many VOC oxidation and photolysis reactions. Together, these differences lead to the "hotter" chemistry in CRI-Strat.



Figure 9. Zonal mean OH (a) and HO₂ (b) in StratTrop simulation. Differences in zonal mean OH and HO₂ between CRI-Strat and StratTrop (c, f), CRI_Emiss_ST and StratTrop (d, g), and CRI_Emiss_C2C3 and StratTrop (e, h). Black lines show average tropopause height over simulation period.

	CRI-Strat	StratTrop	CRI_Emiss_ST	CRI_Emiss_C2C3
$[OH] (10^6 \text{ molecules } \text{cm}^{-3})$	1.335	1.339	1.348	1.375
OH NH:SH ratio	1.38	1.35	1.4	1.4
$[HO_2]$ (pptv)	6.27	5.90	6.02	6.06
$OH: HO_2$ ratio (%)	1.49	1.67	1.61	1.63
CH ₄ lifetime W.R.T. OH (years)	7.77	8.13	7.71	7.60
$HO_2 + HO_2$ flux (P mole year ⁻¹)	60.5	32.2	38.8	39.9

Table 6. Overview of mass weighted annual average tropospheric HOx concentrations, OH northern hemisphere to southern hemisphere ratio, $OH : HO_2$ ratio, methane lifetime with respect to OH and $HO_2 + HO_2$ flux.

In the upper troposphere, OH concentrations are much lower in CRI-Strat com-829 pared to StratTrop for a number of reasons. It is less moist than the lower atmosphere 830 and away from emission sources of short-lived alkenes (which can make OH), hence pri-831 mary OH production is lower. However, CRI-Strat has more formaldehyde than Strat-832 Trop and other carbonyls which can be photolysed in the upper troposphere to produce 833 HO₂. CRI-Strat has more long-lived NMVOCs and CO in the upper troposphere which 834 primarily react with OH converting it into HO2. Ozone levels in the upper troposphere 835 are also lower in CRI-Strat. Hence CRI-Strat has more HO_x in the upper troposphere, 836 but a greater proportion is as HO_2 and there is less OH than in StratTrop. 837

Overall, CRI-Strat has slightly lower average OH concentration than StratTrop, 838 whereas the CRI_Emiss_ST and CRI_Emiss_C2C3 simulations have more OH overall than 839 the base CRI-Strat and StratTrop simulations (Table 6). Hence, given the same emis-840 sions the CRI-Strat mechanism produces more HO_x than StratTrop, but this is coun-841 tered by the additional NMVOCs in the full CRI-Strat mechanism which primarily re-842 act with OH, converting it into HO₂, leading to a lower OH : HO₂ ratio in CRI-Strat. 843 The total $HO_2 + HO_2$ flux, a major sink of HO_x in the atmosphere, is almost twice as 844 high in CRI-Strat than in StratTrop due to both the increase in HO_2 and a faster re-845 action rate (see supplementary Figures S2 and S4). 846

The methane (CH₄) lifetime is slightly shorter in CRI-Strat than in StratTrop, even though OH is higher in StratTrop. The reason is because the OH + CH₄ reaction rate coefficient is slightly faster in CRI-Strat (see Figure S5 in the supplement). While this difference has little impact on the concentration of CH₄, because this is set by surface boundary conditions, it influences tropospheric chemistry because CH₄ is a major source of the methyl peroxy radical (CH₃O₂), the most abundant peroxy radical and an important component of the tropospheric ozone forming process (Table 5).

5.4 Carbon Monoxide

854

Carbon monoxide (CO) is important as the main chemical sink for OH, converting it into HO_2 , and as a toxic air pollutant in its own right. It is produced from both primary emission sources, generally from incomplete combustion of biomass and fossil fuels, and secondary production from the oxidation of VOCs (Grant et al., 2010).

The total tropospheric burden of CO is over 50 Tg higher in CRI-Strat compared to StratTrop, and the CRI_Emiss_ST and CRI_Emiss_C2C3 simulations also have higher CO burdens (Table 7). The CO lifetime varies linearly with OH concentration between the simulations (Table 6), as the OH + CO reaction occurs at the same rate in both mechanisms, and so is slightly longer in CRI-Strat as it has slightly less OH overall compared



Figure 10. Average tropospheric column CO in StratTrop simulations for DJF (a) and JJA (b). Differences in tropospheric CO column in DJF and JJA between CRI-Strat and StratTrop (c, f), CRI_Emiss_ST and StratTrop (d, g), and CRI_Emiss_C2C3 and StratTrop (e, h).

to StratTrop. As all of the simulations use the same primary CO emissions, the differ-864 ences in CO burden are due to secondary production and oxidant profiles. Although most 865 of this secondary production occurs via formaldehyde (HCHO) oxidation and photoly-866 sis channels, there is a large increase in contribution from other channels in CRI-Strat 867 compared to StratTrop (these being oxidation and photolysis of larger carbonyls, O_3 +alkene 868 reactions and PAN oxidation). The increase in CO production from HCHO and other 869 channels is also apparent to a lesser extent in the CRI_Emiss_ST and CRI_Emiss_C2C3 870 scenarios. The more explicit NMVOC degradation chemistry in the CRI-Strat mecha-871 nism produces more HCHO and other oxidised NMVOCs, which go on to form CO as 872 they are further oxidised and undergo photolysis. 873

The bulk of the additional CO burden is in the southern hemisphere downwind of 874 regions with high BVOC emissions, such as the Amazon, Central Africa and Australia 875 (Fig. 10). In CRI_Emiss_ST and CRI_Emiss_C2C3, increases in CO column are localised 876 to these regions and downwind of them (Fig. 10 (d, e, g, h)), implying that the primary 877 cause of increased CO is the more explicit, multigenerational representation of isoprene 878 degradation in the CRI-Strat mechanism. The base CRI-Strat simulation has much more 879 CO over these BVOC dominated regions and a background increase in column CO of 880 around 5-10 DU across almost the entire world, including the northern hemisphere (Fig. 881 10 (c, f)). This is due to additional CO production from degradation of higher NMVOCs 882 emitted by anthropogenic sources, as well as from the explicit APINENE and BPINENE 883 chemistry in CRI-Strat. 884

Tropospheric CO in the StratTrop mechanism was evaluated against multiple datasets by Archibald et al. (2020). They found significant negative biases in column CO of 10-20 DU over much of the Northern hemisphere, but similar values across the Southern



Figure 11. Average tropospheric column HCHO in StratTrop simulations for DJF (a) and JJA (b). Differences in tropospheric HCHO column density in DJF and JJA between CRI-Strat and StratTrop (c, f), CRI_Emiss_ST and StratTrop (d, g), and CRI_Emiss_C2C3 and StratTrop (e, h).

hemisphere and a high bias over BVOC rich Central Africa and Amazonia compared to 888 the MOPITT satellite product. The CRI-Strat simulation has a lot more CO in the North-889 ern hemisphere, reducing the negative bias there, but far too much CO in the southern 890 hemisphere particularly downwind of BVOC sources. Although the CRI-Strat mecha-891 nism captures secondary production of CO better than StratTrop, it highlights that too 892 much CO production occurs in the southern hemisphere, likely due to errors in BVOC 893 emissions. In addition, the model setup either lacks some key CO or CO precursor emissions in the Northern hemisphere, CO lifetime is too short in the model and/or not enough 895 CO is being transported from low to high latitudes. 896

This importance of HCHO in CO production is emphasised in Figure 11, which shows much higher HCHO column densities above high BVOC emission regions in all of the simulations that use CRI-Strat chemistry. As HCHO is the dominant source of secondary CO, this will lead to higher CO burdens downwind. The more explicit representation of isoprene and monoterpene degradation chemistry clearly leads to greater secondary production of HCHO and as a result CO.

5.5 Peroxy Radicals

903

Peroxy radicals (RO₂) are formed from the oxidation of VOCs and are important but short-lived intermediates in tropospheric ozone formation. The largest differences between the two chemical mechanisms centre around treatment of RO₂ species, as discussed in section 3.1. The CRI-Strat has a total of 47 RO₂ species (as opposed to 9 in StratTrop), all of which undergo reactions with NO, NO₃, HO₂ and the summed RO₂

su
eaction
al re
radic
уху
perc
for
and
OSS
on/]
ucti
bord
0 p
or C
es fe
flux
key
and
len a
burd
$^{\circ}$
Õ
eric
hqsc
ropc
ge t
vera
of a
ew e
rervi
õ
2
le
ab
E

Flux		CRI-Strat	$\operatorname{StratTrop}$	CRI_Emiss_ST	CRI_Emiss_C2C3
CO burden (Tg) CO lifetime (days		354.9 39.1	300.2 38.6	317.2 38.1	315.2 37.2
CO production (Tg CO year ⁻¹)	TotalEmissionsHCHO + OHHCHO + $h\nu$ Other ChemOther Phot	$\begin{array}{c c} 3402 \\ 11111 & (32.7\%) \\ 580.3 & (17.1\%) \\ 1293 & (38\%) \\ 125 & (3.7\%) \\ 293 & (8.6\%) \end{array}$	$\begin{array}{c} \textbf{2915}\\ \textbf{1111} \ (38.1\%)\\ 492 \ (16.9\%)\\ 1076 \ (36.9\%)\\ 71.5 \ (2.5\%)\\ 164 \ (5.6\%)\\ \end{array}$	$\begin{array}{c} \textbf{3121}\\ \textbf{3121}\\ \textbf{1111} \ (\textbf{35.6\%})\\ \textbf{551} \ (\textbf{17.7\%})\\ \textbf{163} \ (\textbf{37.3\%})\\ \textbf{101} \ (\textbf{3.25\%})\\ \textbf{194} \ (\textbf{6.23\%}) \end{array}$	$\begin{array}{c} \textbf{3171}\\ \textbf{1111} \ (35\%)\\ 566 \ (17.8\%)\\ 1177 \ (37.1\%)\\ 107 \ (3.4\%)\\ 2111 \ (6.7\%)\end{array}$
CO Loss (Tg CO year ⁻¹)	Total CO + OH CO Dry Dep	$\begin{array}{c} \textbf{3311}\\ \textbf{3157} (95.4\%)\\ \textbf{153} (4.6\%) \end{array}$	$\begin{array}{c} \textbf{2841} \\ \textbf{2704} \ (95.2\%) \\ \textbf{137} \ (4.8\%) \end{array}$	$\begin{array}{c} \textbf{3041} \\ \textbf{2900} \ (95.4\%) \\ \textbf{141} \ (4.7\%) \end{array}$	$\begin{array}{c} 3090 \\ 2948 \ (95.4\%) \\ 142 \ (4.6\%) \end{array}$
$\frac{\text{RO}_2 \text{ fluxes}}{\text{(P mole year}^{-1})}$	$\begin{array}{c} \textbf{Total}\\ \text{RO}_2 + \text{NO}\\ \text{RC}(0)\text{O}_2 + \text{NO}_2 \ (\text{PAN Prod})\\ \text{RO}_2 + \text{NO}_3\\ \text{RO}_2 + \text{HO}_2\\ \text{RO}_2 + \text{HO}_2\\ \text{RO}_2 + \text{RO}_2 \end{array}$	$\begin{array}{c} \textbf{111.5}\\ \textbf{18.4} \ (16.5\%)\\ 20.6 \ (18.5\%)\\ 0.3 \ (0.3\%)\\ 60.4 \ (54.2\%)\\ 11.8 \ (10.6\%)\end{array}$	75.8 11.4 (15.0%) 19.6 (25.9%) 0.2 (0.3%) 41.2 (54.4%) 3.4 (4.5%)	$\begin{array}{c} \textbf{91.5}\\ \textbf{14.1} (15.4\%)\\ \textbf{14.7} (16.1\%)\\ \textbf{0.3} (0.3\%)\\ \textbf{52.4} (57.3\%)\\ \textbf{10.1} (11.04\%)\end{array}$	$\begin{array}{c} \textbf{87.9}\\ 13.1 \ (14.9\%)\\ 14.0 \ (16.0\%)\\ 0.2 \ (0.3\%)\\ 51.0 \ (58.0\%)\\ 9.5 \ (10.8\%)\end{array}$

radical pool in ways that conserve the number of potential ozone forming steps. Acetyl 909 peroxy radicals $(RC(O)O_2)$ are also crucial as they form thermally stable peroxyacetyl 910 nitrate (PAN) compounds, important NO_y reservoir species. CRI-Strat has multiple ad-911 dition PAN species from degradation of higher NMVOC species. Table 7 gives a sum-912 mary of the key RO_2 reaction fluxes in the troposphere. The CRI-Strat simulation has 913 a 50% higher RO_2 flux compared to StratTrop. The CRLEmiss_ST and CRLEmiss_C2C3 914 simulations have total fluxes 20% and 16% higher than StratTrop respectively, showing 915 that a large fraction of the additional flux is a result of the more explicit multigenera-916 tion chemistry, and not just from the additional NMVOC emissions in CRI-Strat. 917

There are big differences in the respective fates of RO_2 species. In the CRI-Strat 918 mechanism, RO_2 species are more than twice as likely to react with another RO_2 species 919 than in StratTrop, because the full range of these cross-reactions are parameterised for 920 all RO₂ species. The CRI-Strat mechanisms also have a greater flux through the $RO_2 + HO_2$ 921 branches, which form ROOH species which are then often lost via wet and dry deposi-922 tion, due to higher HO_2 and RO_2 concentrations in CRI-Strat (see Table 6). There is 923 a much larger flux through PAN forming reactions in StratTrop. The MPAN forming 924 reaction in StratTrop is much faster than the equivalent reaction in CRI-Strat (see sup-925 plementary Figure S5), and hence the production of MPAN and PAN-type species is greater 926 in StratTrop even though all other PAN-type species (including PAN) are formed in greater 927 abundance in CRI-Strat. However, as PAN-type species rapidly dissociate back to the 928 original reactants in high temperature conditions, the higher flux through the $RC(O)O_2 + NO_2$ 929 pathway does not necessarily lead to permanent oxidation, unlike via the other branches. 930

5.6 Nitrogen Oxides and their reservoirs

There are stark differences in nitrogen oxides between the different model simu-932 lations, as shown in Figure 12. In this analysis, we use the StratTrop simulation as a base-933 line to evaluate CRI-Strat, as StratTrop in UKESM1 has been thoroughly evaluated by 934 Archibald et al. (2020). In the StratTrop simulation, the tropospheric column density 935 of NO_2 is highest over polluted regions, such as China and Europe, and is higher in lo-936 cal winter than summer, due to higher emissions, and lower oxidising capacity and pho-937 tolysis rates in winter (Fig. 12 (a, b)). Archibald et al. (2020) found a high bias down-938 wind of these polluted regions in winter of up to 5×10^{15} molecules cm⁻² (approximately 939 50%) compared to the OMI satellite product, but a small low bias over most of the rest 940 of the world (see Archibald et al. (2020) Fig. 18). The CRI-Strat simulation has much 941 lower column density of NO_2 across the Northern Hemisphere continental regions in win-942 ter such as over China where CRI-Strat has column densities more than 5×10^{15} molecules cm⁻² 943 lower than StratTrop (Fig. 12 (c)). CRLEmiss_ST has lower NO₂ column than Strat-944 Trop directly over high emission regions but is similar downwind (Fig. 12 (d, g). CRI_Emiss_C2C3 945 is somewhere between the base CRI-Strat and CRI-Emiss_ST simulations, with a lower 946 column density of NO_2 over polluted regions and downwind of them over continents, but 947 not to the same degree as CRI-Strat (Fig. 12 (e, h). 948

Nitrogen is conserved through chemical reactions (noting that some minor changes were made to the StratTrop mechanism in order for it to conserve nitrogen, see Section S3 in the supplement for details) and all simulations have exactly the same NO_x emissions, therefore if NO_2 is lower it must have been converted into some other form and/or lost via deposition. It is conventional to use the shorthand NO_x as the sum of nitrogen oxides NO and NO_2 , NO_z is the sum of their reservoir species, and NO_y is the sum of all oxidised nitrogen compounds:

956 957

959

960

931

$$NO_x = NO + NO_2 \tag{9}$$

 $NO_{z} = HONO_{2} + NO_{3} + 2*N_{2}O_{5} + HO_{2}NO_{2} + XONO_{2} + PANs + RONO_{2} + CH_{3}O_{2}NO_{2} + Nitrophenols$ (10)

$$NO_y = NO_x + NO_z \tag{11}$$


Figure 12. Average tropospheric column NO₂ in StratTrop simulations for DJF (a) and JJA (b). Differences in tropospheric NO₂ column in DJF and JJA between CRI-Strat and StratTrop (c, f), CRI_Emiss_ST and StratTrop (d, g), and CRI_Emiss_C2C3 and StratTrop (e, h).

The conversion of NO_x to NO_z can inhibit the ozone forming process. The most component of NO_z is nitric acid (HONO₂), which is efficiently dry and wet deposited and so acts as a sink for reactive nitrogen in the troposphere. However, some reservoir species (such as PAN) can also aid overall ozone production if they are transported, releasing NO_x into other regions of the troposphere where the ozone forming potential per NO_x molecule is greater.

When analysing how oxidised nitrogen differs between these model simulations, its 967 important to understand some key differences between the two chemical mechanisms. 968 The CRI-Strat mechanism has the same inorganic nitrogen species as StratTrop, but it 969 has many more organic nitrogen containing species which contribute to NO_z . Firstly, in 970 CRI-Strat almost every $RO_2 + NO$ reaction has a minor branch which forms an organon-971 itrate $(RONO_2)$, whereas in StratTrop only CH_3ONO_2 and ISON (which represents organon-972 itrates from isoprene oxidation) exist. Peroxy radicals with nitrate groups are also formed 973 from NO₃ initiated reactions with alkenes, which can also go on to form stable organon-974 itrates. These organonitrates can act as NO_x reservoirs, transported long distances be-975 fore undergoing further oxidation or photolysis to release NO_x or getting deposited out 976 of the atmosphere. CRI-Strat also has several more PAN-type species, formed from ther-977 mal equilibrium between peroxyacyl radicals and NO_2 . Finally, there are some species 978 in CRI-Strat for which there is no equivalent in StratTrop: CH₃O₂NO₂ is formed from 979 the thermal equilibrium between CH_3O_2 and NO_2 and can be an important reservoir species 980 in the upper troposphere (Browne et al., 2011), while nitrated phenols are formed dur-981 ing the oxidation of benzene and toluene in the presence of NO_x (Harrison et al., 2005). 982

Zonal mean differences in these summed species are shown in Figure 13. There is reduced NO_x in the CRI-Strat mechanism compared to StratTrop (Fig. 13 (a)), partic-



Figure 13. Average zonal mean NO_x (a), NOy (e) and NO_z (f) in StratTrop run from 2010-2018. Zonal mean differences in NO_x , NO_y and NO_z between CRI-Strat and StratTrop (b, f and j), CRI_Emiss_ST and StratTrop (c, g and k), and CRI_Emiss_C2C3 and StratTrop (d, h, l) respectively. Black lines show average tropopause height over simulation period.

ularly in the lower atmosphere in the northern hemisphere and in the upper troposphere, 985 that is visible in the base CRI-Strat simulation (Fig. 13 (b)) and in the runs with same 986 total NMVOC mass emissions to StratTrop (Fig. 13 (c and d)). In contrast, total ox-987 idised nitrogen NO_v is generally higher in CRI-Strat compared to StratTrop (Fig. 13 (e 988 and f)), except in the lower atmosphere around 30°N (specifically over East Asia and 989 India, see supplementary Figure S13), but lower almost everywhere in CRI_Emiss_ST 990 and CRLEmiss_C2C3 (Fig. 13 (g and h)). The increase in NO_y is shown to be due to 991 there being more NO_z almost everywhere in the CRI-Strat model simulation compared 992 to StratTrop (Fig. 13 (j)). However, in the CRI-STEmiss and CRI-Emiss_C2C3 simu-993 lations higher NO_z is only found in the lower atmosphere northern hemisphere down-994 wind of high emitting regions (Fig. 13 (k and l)). Hence the increased NO_z in the CRI-995 Strat simulation is clearly tied in with the extra NMVOC emissions. Without these added 996 organic species, the CRI-Strat chemical mechanism has greater propensity to produce 997 more NO_z in polluted regions but loose oxidised nitrogen mass overall. 998

⁹⁹⁹ The base CRI-Strat simulation has more tropospheric NO_y than StratTrop (1.112 Tg N ¹⁰⁰⁰ vs. 1.018 Tg N), but has less of the total nitrogen as NO_x (10.3% vs 14.9%) and more ¹⁰⁰¹ as NO_z reservoirs (89.7% vs 85.1%), as shown in Table 8. However, CRI-Strat has a smaller ¹⁰⁰² fraction of NO_y as HONO₂ (45.8% vs 50.4%), the single largest component of NO_z . In-¹⁰⁰³ stead, a greater fraction is stored as PANs and RONO₂, as well as CH₃O₂NO₂ and ni-¹⁰⁰⁴ trophenol species which are not in the StratTrop chemical mechanism.

Table 8. Overview of tropospheric oxidised nitrogen burdensstratosphere-troposphere transfer (STT) and NO _y lifetime in the	rdens (fraction of total NO in the troposphere (fractic	y in brackets), troj on of total NO _y dej	oospheric oxidised nitro position in brackets).	ogen emission and de
	-			

	CRI-Strat	StratTrop	CRI_Emiss_ST	CRI_Emiss_C2C3
	$\begin{array}{c c} 1.112 \\ 0.115 & (10.3\%) \\ 0.998 & (89.7\%) \end{array}$	$\begin{array}{c} 1.018 \\ 0.152 \ (14.9\%) \\ 0.866 \ (85.1\%) \end{array}$	$\begin{array}{c} 0.950\\ 0.136 \ (14.3\%)\\ 0.814 \ (85.7\%)\end{array}$	$\begin{array}{c} 0.910 \\ 0.131 \ (14.4\%) \\ 0.779 \ (85.6\%) \end{array}$
HONO ₂ Burden (Tg N) Other inorganic NO _z (Tg N) PANs (Tg N) RONO ₂ (Tg N) CH ₃ O ₂ NO ₂ (Tg N) Nitrophenols (Tg N)	$ \left \begin{array}{c} 0.509 & (45.8\%) \\ 0.021 & (1.9\%) \\ 0.367 & (33.0\%) \\ 0.061 & (5.5\%) \\ 0.008 & (0.7\%) \\ 0.031 & (2.8\%) \end{array} \right $	$\begin{array}{c} 0.513 \ (50.4\%) \\ 0.018 \ (1.7\%) \\ 0.296 \ (29.1\%) \\ 0.039 \ (3.9\%) \\ N/A \\ N/A \end{array}$	$\begin{array}{c} 0.506 \ (53.2\%) \\ 0.018 \ (1.9\%) \\ 0.245 \ (25.8\%) \\ 0.038 \ (4.0\%) \\ 0.007 \ (0.7\%) \\ 0.0 \end{array}$	$\begin{array}{c} 0.512 & (56.2\%) \\ 0.019 & (2.1\%) \\ 0.206 & (22.7\%) \\ 0.035 & (3.9\%) \\ 0.007 & (0.7\%) \\ 0.0 \end{array}$
Total NO _x Emissions (Tg N year ⁻¹) Total NO _y Deposition (Tg N year ⁻¹) Inferred STT (Tg N year ⁻¹)	61.5 63.0 1.46	61.5 62.9 1.40	$\begin{array}{c} 61.5\\ 63.0\\ 1.43\end{array}$	$\begin{array}{c} 61.5\\ 63.0\\ 1.44\end{array}$
NO _x Dry deposition (Tg N year ⁻¹) HONO ₂ Wet deposition (Tg N year ⁻¹) HONO ₂ Dry deposition (Tg N year ⁻¹) Other inorganic NO _z deposition (Tg N year ⁻¹) PANs dry deposition (Tg N year ⁻¹) RONO ₂ deposition (Tg N year ⁻¹) Nitrophenol deposition (Tg N year ⁻¹)	$ \begin{array}{c} 6.83 \ (10.8\%) \\ 29.0 \ (46.0\%) \\ 22.0 \ (34.9\%) \\ 1.21 \ (1.9\%) \\ 1.70 \ (2.7\%) \\ 2.09 \ (3.3\%) \\ 0.22 \ (0.4\%) \end{array} $	$\begin{array}{c} 7.70 & (12.2\%) \\ 30.1 & (47.8\%) \\ 21.6 & (34.3\%) \\ 0.97 & (1.6\%) \\ 1.28 & (2.0\%) \\ 1.30 & (2.1\%) \\ \mathrm{N/A} \end{array}$	$\begin{array}{c} 7.36 \ (11.7\%) \\ 30.1 \ (47.8\%) \\ 22.3 \ (35.5\%) \\ 1.04 \ (1.6\%) \\ 0.894 \ (1.4\%) \\ 1.22 \ (2.0\%) \\ 0.0 \end{array}$	$\begin{array}{c} 7.25 \ (11.5\%) \\ 30.0 \ (47.7\%) \\ 22.4 \ (35.6\%) \\ 1.08 \ (1.7\%) \\ 0.918 \ (1.5\%) \\ 1.28 \ (2.0\%) \\ 0.0 \end{array}$
NO _y deposition lifetime (days) HONO ₂ deposition lifetime (days) PANs deposition lifetime (days) RONO ₂ deposition lifetime (days)	6.44 3.65 78.9 10.7	$\begin{array}{c} 5.91 \\ 3.62 \\ 84.5 \\ 11.0 \end{array}$	5.51 3.53 100.1 11.4	5.28 3.57 82.0 9.99

Many of these differences can be attributed to the added NMVOC emissions in the 1005 CRI-Strat simulations. The greater total quantity of NMVOC emissions, added larger 1006 NMVOCs and the explicit depiction of their multigenerational degradation provides more 1007 opportunities for NO_z reservoir species to form. In the CRLEmiss_ST simulation, the total tropospheric NO_v is 0.95 Tg N, showing that the CRI-Strat mechanism has lower 1009 NO_v when if run with the same NMVOC emissions as StratTrop (Table 8). The frac-1010 tion of NO_x (14.3%) is slightly lower than StratTrop, but still higher than CRI-Strat. 1011 In the CRLEmiss_C2C3 simulation, NO_v is lower still, at 0.91 Tg N. The driving fac-1012 tor behind these differences appears to be a greater fraction of total NO_{v} as $HONO_{2}$ in 1013 CRI_Emiss_ST (53.2%) and CRI_Emiss_C2C3 (56.2%) compared to StratTrop (50.4%) 1014 or the base CRI-Strat run (45.8%). However, CRI_Emiss_ST and CRI_Emiss_C2C3 both 1015 have slightly less $HONO_2$ than StratTrop in absolute terms. 1016

The importance of these differences in oxidised nitrogen species burdens can be bet-1017 ter understood by analysing the sources and sinks of oxidised nitrogen in the troposphere. 1018 In all of the simulations, total NO_x emissions are identical, and total NO_y deposition al-1019 most identical, as shown in Table 8. Total NO_v deposition is greater than total emissions because of net transfer of NO_v species (mostly $HONO_2$) from the stratosphere to 1021 the troposphere, with this additional NO_v originating from reaction of N_2O and $O(^1D)$ 1022 in the stratosphere. From the tropospheric NO_v burden and sum of all deposition sinks, 1023 we can calculate the mean tropospheric $NO_{\rm v}$ lifetime, which ranges from around 6.4 days 1024 in CRI-Strat to 5.9 days in StratTrop, 5.5 days in CRI_Emiss_ST and only 5.3 days in 1025 CRI_Emiss_C2C3. By decomposing the channels by which NO_v is deposited, we can see 1026 that less is deposited as NO_x or $HONO_2$ in CRI-Strat compared to StratTrop, and a larger 1027 fraction is deposited as organic nitrogen containing species. In general these organic nitrogen-1028 containing species deposit less efficiently than HONO₂ (R. Sander, 2015) and have longer 1029 lifetimes with respect to deposition, effectively extending the $NO_{\rm v}$ lifetime and allow-1030 ing total NO_z to accumulate (Table 8). They are also more effective at transporting NO_x 1031 from polluted regions and releasing it in cleaner regions which are more NO_x sensitive, 1032 going a long way to explaining why the CRI-Strat simulation has the highest overall rate 1033 of ozone production (Table 5). 1034

¹⁰³⁵ Compared to the StratTrop model run, the CRI_Emiss_ST and CRI_Emiss_C2C3 ¹⁰³⁶ simulations have similar wet deposition loss of HONO₂ but greater loss via dry depo-¹⁰³⁷ sition. This is because there is more HONO₂ production in the boundary layer (see sup-¹⁰³⁸ plementary Figure S14), where it can be rapidly lost via dry deposition before it has time ¹⁰³⁹ to be transported away from emission sources. Hence, total tropospheric NO_z and NO_y ¹⁰⁴⁰ levels, and NO_y lifetime, are overall lower in CRI_Emiss_ST and CRI_Emiss_C2C3 com-¹⁰⁴¹ pared to StratTrop.

1042

5.7 Summary and synthesis

 1043 Collating all of the information covered in this analysis, we can understand that the variations in tropospheric ozone are largely driven by the variations in the NO_y fields between the simulations:

- 10461. In polluted regions, ozone production is more efficient in the CRI-Strat mecha-
nism due to the more explicit representation of multigenerational NMVOC-NOx
chemistry, particularly when the amount of NMVOC available is increased. How-
ever, loss of ozone is also more rapid, particularly due to the increased proportion
of O_x lost via the $O(^1D) + H_2O$ reaction.
- The enhanced HONO₂ production in CRI-Strat leads to greater NO_y removal in the boundary layer via dry deposition, reducing the amount entering the free troposphere.

- 3. In the base CRI-Strat run, faster HONO₂ production is compensated by greater production of organic nitrogen-containing species which extend the lifetime of NO_y and enhance ozone production in remote regions.
 4. These accurations effects accurately belonge some the schole term containing the schole term containing term.
 - 4. These competing effects roughly balance over the whole troposphere, leading to similar tropospheric ozone burdens in the CRI-Strat and StratTrop simulations, albeit with more ozone near the surface in CRI-Strat.
- 5. The CRI_Emiss_ST and CRI_Emiss_C2C3 simulations are less able to form organonitrates than the CRI-Strat simulation so lose more nitrogen to HONO₂ deposition and therefore have lower NO_y burdens. Combined with the shorter ozone lifetime, these two simulations have the lowest tropospheric ozone burdens despite having higher ozone production rates than StratTrop in the polluted boundary layer.

¹⁰⁶⁵ Overall, we find a strong sensitivity between emissions of NMVOCs and the oxi-¹⁰⁶⁶ dised nitrogen budget, which leads to considerable knock on effects to the tropospheric ¹⁰⁶⁷ ozone burden and oxidising capacity of the atmosphere. This has important implications ¹⁰⁶⁸ for the need to improve emissions of NMVOCs (which are typically poorly constrained ¹⁰⁶⁹ (Huang et al., 2017)) and how these emissions are treated in simpler mechanisms such ¹⁰⁷⁰ as StratTrop. The dependence of NO_x to emissions of NMVOCs is also relevant for how ¹⁰⁷¹ we interpret comparisons of NO_x between models and observations.

The CRI mechanism has now been implemented in a number of models, including 1072 STOCHEM (S. Utembe et al., 2010) and WRF-Chem (Archer-Nicholls et al., 2014). The 1073 STOCHEM implementation also saw an increase in ozone production and loss compared 1074 to its existing mechanism, but this led to much higher ozone burdens across the tropo-1075 sphere, particularly over the oceans. In contrast, CRI-Strat has similar tropospheric ozone 1076 burdens to StratTrop and lower ozone concentrations across much of the world's oceans. 1077 We believe the higher tropospheric ozone columns did not occur in UKCA because of 1078 the shorter O_x lifetime in CRI-Strat compared to StratTrop due to the faster flux through 1079 the $O(^{1}D) + H_{2}O$ reaction in CRI-Strat. The WRF-Chem implementation also saw an 1080 increase in ozone production but little overall increase in ozone over the model domain compared to the CBM-Z mechanism. However, in a regional model a species such as ozone, 1082 with a typical lifetime of a few weeks, is very sensitive to the effects of boundary con-1083 ditions. In Archer-Nicholls et al. (2014), boundary conditions were prescribed by the MOZART 1084 model for both CBM-Z and CRIv2-R5. Combining WRF-Chem with boundary condi-1085 tions driven by UKCA simulations, with both models using the same CRI mechanism, 1086 could offer a solution to the problem of how to account for the impact of boundary con-1087 ditions. 1088

1089 6 Conclusions

1054

1055

1056

1057

1058

1059

1060

1061

1063

1064

The CRIv2-R5 chemical mechanism has been integrated into the UKCA model, merged 1090 with the existing stratospheric chemistry and coupled to GLOMAP-mode aerosol to cre-1091 ate the CRI-Strat mechanism. The mechanism is constrained to reliably reproduce the 1092 ozone forming potential from MCMv3.1, enabling traceability to our best understand-1093 ing and providing a benchmark to evaluate simpler mechanisms and test other aspects 1094 of model setup. This new mechanism marks a step change in chemical complexity and 1095 comprehensiveness, improving representation of important tropospheric processes, such 1096 as BVOC chemistry, peroxy radicals and organonitrates, while remaining sufficiently af-1097 fordable for use in an Earth System Model (approximately 75% longer runtime compared 1098 to an equivalent run with StratTrop). CRI-Strat can now be used in UKESM1, a flag-1099 ship model used for CMIP6 (Eyring et al., 2016), opening up new potential scientific en-1100 quiries. As the implementation has been done using the ASAD framework (Carver et 1101 al., 1997), it can also be ported to other models which share the same framework. 1102

In this paper we critically evaluate and compare CRI-Strat to the well-established
 StratTrop mechanism, highlighting some key differences:

 CRI-Strat has higher surface ozone and CO compared to StratTrop, improving some biases compared to observations but worsening others.
 CRI-Strat chemistry is generally "hotter" than StratTrop: it has much higher production and loss of ozone, more HO_x and more secondary production of CO.
 Total tropospheric ozone burden and ozone column density are surprisingly sim-

- ilar between the simulations, given the large differences in production rate.
- 4. CRI-Strat partitions a greater fraction of nitrogen into reservoir forms (NO_z) with a lower fraction in the reactive form (NO_x) .
- 11135. Many differences are related to the speciation of NMVOCs. Significantly more emit-
ted species are included in CRI-Strat than in StratTrop. When using exactly the
same emissions, CRI-Strat has 8.6% lower tropospheric ozone burden than Strat-
Trop.116Trop.

Some of these differences reflect differences in reaction rate coefficients for key reactions, 1117 a number of which are out of date in CRIv2-R5 and therefore CRI-Strat. The faster pro-1118 duction of ozone that occurs in CRI-Strat when photochemical conditions allow means 1119 that it is more sensitive to model structural uncertainties than StratTrop, particularly 1120 relating to emissions, model resolution and parameterisations such as for lightning- NO_x 1121 emissions. The more complex CRI-Strat mechanism is not designed to be a replacement 1122 for StratTrop, but provides a new tool that expands the possible scientific questions that 1123 can be tackled with the model and a benchmark to evaluate against. 1124

The tests and evaluation described in this paper set out to fully characterise the 1125 CRI-Strat mechanism against the StratTrop mechanism (the reference mechanism for 1126 UEKSM1) for the gas-phase composition of species relevant for the climate. This pro-1127 vides information that is essential to understand and make use of the new mechanism. 1128 However, the experiments performed do not use CRI-Strat to its full potential. We ex-1129 pect it to exceed the capabilities of StratTrop when run at higher spatial resolution and 1130 in evaluation against field campaigns with a focus on oxidants. Future work will also fo-1131 cus on highly polluted environments or those dominated by BVOCs and production of 1132 SOA, for which CRI has been shown to provide a robust framework for simulating (S. Utembe 1133 et al., 2010). We also plan to run experiments for different climate and emission regimes 1134 such as the pre-industrial atmosphere; these experiments (combined with multi model 1135 analyses) will enable us to understand if we can be confident that UKESM1 represents 1136 the changes in composition and chemistry-climate feedbacks from pre-industrial to the present day realistically. This evaluation has also neglected analysis of aerosols, whose 1138 formation rates will differ with CRI-Strat due to changes in oxidant fields, and will be 1139 properly evaluated in the future. There is also scope to improve the coupling between 1140 CRI-Strat and the GLOMAP-mode scheme, for example updating reaction rates and ex-1141 tending the isoprene chemistry with inclusion of CRIv2.2 (M. E. Jenkin, Khan, et al., 1142 2019), improving representation of BVOC environments, OH recycling and further ex-1143 pansion with HOM chemistry (Weber et al., 2020). 1144

1145 Acknowledgments

1110

1111

1112

SAN would like to acknowledge funding on the NERC PROMOTE grant, reference 1146 NE/P016383/1. NLA, ATA, and MRR are supported by NERC and NCAS through the 1147 ACSIS project. FMO'C acknowledges support from the the Met Office Hadley Centre 1148 Climate Programme funded by BEIS and Defra (GA01101) and the EU Horizon 2020 1149 Research Programme CRESCENDO project, grant agreement number 641816. UKCA 1150 development is supported by the NERC – Met Office Joint Weather and Climate Re-1151 search Programme. This work used Monsoon2, a collaborative High Performance Com-1152 puting facility funded by the Met Office and the Natural Environment Research Coun-1153 cil. Analysis used JASMIN, the UK collaborative data analysis facility. We thank the 1154 Global Emission Initiative (GEIA) for providing access to emissions data via the Emis-1155

sions of atmospheric Compounds and Compilation of Ancillary Data (ECCAD) archive.
We would also like to thank the Tropospheric Ozone Assessment Report (TOAR) initiative for providing access to the aggregated surface ozone statistics that they have calculated using data collected from multiple stations and networks globally. Development
and application of the OMI ozone profile retrieval scheme was funded through NERCNCEO and ESA Climate Change Initiative. The authors declare no conflicts of interest.

Due to intellectual property right restrictions, we cannot provide either the source 1163 code or documentation papers for the UM. The Met Office Unified Model is available 1164 for use under licence. The functionality described here is fully available in the UM trunk 1165 from version 11.8. A number of research organisations and national meteorological ser-1166 vices use the UM in collaboration with the UK Met Office to undertake basic atmospheric 1167 process research, produce forecasts, develop the UM code and build and evaluate Earth 1168 system models. For further information on how to apply for a licence; see https://www 1169 .metoffice.gov.uk/research/approach/modelling-systems/unified-model (last 1170 access: 24 November 2020). Data tables of the full CRI-Strat mechanism as described 1171 in this paper are included in the supplement. 1172

UM simulations are compiled and run in suites developed using the Rose suite en-1173 gine (http://metomi.github.io/rose/doc/html/index.html, last access: 24 Novem-1174 ber 2020) and scheduled using the Cylc workflow engine (https://cylc.github.io/, 1175 last access: 24 November 2020). Both Rose and Cylc are available under version 3 of the GNU General Public License (GPL). In this framework, the suite contains the informa-1177 tion required to extract and build the code as well as configure and run the simulations. 1178 Each suite is labelled with a unique identifier and is held in the same revision-controlled 1179 repository service in which we hold and develop the model's code. This means that these 1180 suites are available to any licensed user of the UM. The input emissions data are avail-1181 able from Input4MIPs (https://esgf-node.llnl.gov/projects/input4mips/, last ac-1182 cess: 24 November 2020). 1183

1184 **References**

- Akimoto, H.
 (2003, dec).
 Global air quality and pollution.
 Science (New York,

 N.Y.), 302(5651), 1716-9.
 Retrieved from http://www.ncbi.nlm.nih.gov/

 pubmed/14657488
 doi: 10.1126/science.1092666
- Andreae, M. O. (1990). Ocean-atmosphere interactions in the global biogeochemical sulfur cycle. *Marine Chemistry*, 30(C), 1–29. doi: 10.1016/0304-4203(90)90059 -L
- 1191Archer-Nicholls, S., Lowe, D., Utembe, S., Allan, J., Zaveri, R. A., Fast, J. D., ...1192McFiggans, G. (2014, nov). Gaseous chemistry and aerosol mechanism1193developments for version 3.5.1 of the online regional model, WRF-Chem.1194Geoscientific Model Development, 7, 2557–2579. Retrieved from http://1195www.geosci-model-dev.net/7/2557/2014/
- 1196Archibald, A., O'Connor, F., Abraham, N. L., Archer-Nicholls, S., Chipperfield,1197M., Dalvi, M., ... Zeng, G. (2020). Description and evaluation of the1198UKCA stratosphere-troposphere chemistry scheme (StratTrop vn 1.0) im-1199plemented in UKESM1. Geoscientific Model Development, 13, 1223–1266. doi:120010.5194/gmd-2019-246
- 1201Atkinson, R.(1990).Gas-phase tropospheric chemistry of organic compounds:1202a review.Atmospheric Environment, 24A(1), 1-41.Retrieved from1203http://linkinghub.elsevier.com/retrieve/pii/S13522310070099341204doi: 10.1016/j.atmosenv.2007.10.068
- 1205Atkinson, R., Baulch, D. L., Cox, R. A., Crowley, J. N., Hampson, R. F., Hynes,1206R. G., ... Rossi, M. J. (2004). Evaluated kinetic and photochemical data for1207atmospheric chemistry : Volume I gas phase reactions of Ox, HOx, NOx and

1208	SOx species. Atmospheric Chemistry and Physics, 4, 1461–1738.
1209	Atkinson, R., Baulch, D. L., Cox, R. A., Crowley, J. N., Hampson, R. F., Hynes,
1210	R. G., Troe, J. (2006, sep). Evaluated kinetic and photochemical
1211	data for atmospheric chemistry: Volume II – gas phase reactions of or-
1212	ganic species. Atmospheric Chemistry and Physics, $6(11)$, $3625-4055$.
1213	Retrieved from http://www.atmos-chem-phys.net/6/3625/2006/ doi:
1214	10.5194/acp-6-3625-2006
1215	Atkinson, R., Baulch, D. L., Cox, R. A., Hampson, R. F., Kerr, J. A., Rossi, M. J.,
1216	& Troe, J. (1997). Evaluated Kinetic and Photochemical Data for Atmospheric
1217	Chemistry Supplement VI. J. Phys. chem. Ref. Data, 26(6), 1329–1499.
1218	Aumont, B., Szopa, S., & Madronich, S. (2005). Modelling the evolution of organic
1219	carbon during its gas-phase tropospheric oxidation: Development of an ex-
1220	plicit model based on a self generating approach. Atmospheric Chemistry and
1221	<i>Physics</i> , $5(9)$, 2497–2517. doi: 10.5194/acp-5-2497-2005
1222	Banerjee, A., Archibald, A. T., Maycock, A. C., Telford, P., Abraham, N. L., Yang,
1223	X., Pyle, J. A. (2014). Lightning NOx, a key chemistry-climate interaction:
1224	Impacts of future climate change and consequences for tropospheric oxidis-
1225	ing capacity. Atmospheric Chemistry and Physics, $14(18)$, $9871-9881$. doi:
1226	10.5194/acp-14-9871-2014
1227	Bates, K. H., & Jacob, D. J. (2020). An Expanded Definition of the Odd Oxy-
1228	gen Family for Tropospheric Ozone Budgets: Implications for Ozone Lifetime
1229	and Stratospheric Influence. Geophysical Research Letters, $47(4)$, 1–9. doi:
1230	10.1029/2019GL 084486
1231	Bianchi, F., Kurtén, T., Riva, M., Mohr, C., Rissanen, M. P., Roldin, P., Ehn,
1232	M. (2019). Highly Oxygenated Organic Molecules (HOM) from Gas-Phase
1233	Autoxidation Involving Peroxy Radicals: A Key Contributor to Atmospheric
1234	Aerosol. Chemical Reviews. doi: 10.1021/acs.chemrev.8b00395
1235	Boersma, K. F., Eskes, H. J., Veefkind, J. P., Brinksma, E. J., Van Der A, R. J.,
1236	Sneep, M., Bucsela, E. J. (2007). Atmospheric Chemistry and Physics
1237	Near-real time retrieval of tropospheric $\ MO_2$ from OMI. Atmos.
1238	Chem. Phys, 7, 2103-2118. Retrieved from www.atmos-chem-phys.net/7/
1239	2103/2007/
1240	Boucher, O., Randall, D., Artaxo, P., Bretherton, C., Feingold, G., Forster, P.,
1241	Zhang, X. (2013). Clouds and Aerosols. In T. Stocker et al. (Eds.), <i>Climate</i>
1242	change 2013: The physical science basis. contribution of working group i to
1243	the fifth assessment report of the intergovernmental panel on climate change.
1244	Cambridge, United Kingdom and New York, NY, USA: Cambridge University
1245	Press.
1246	Bowman, K. W., Shindell, D. T., Worden, H. M., Lamarque, J. F., Young, P. J.,
1247	Stevenson, D. S., Worden, J. R. (2013). Evaluation of ACCMIP out-
1248	going longwave radiation from tropospheric ozone using TES satellite ob-
1249	servations. Atmospheric Chemistry and Physics, 13(8), 4057–4072. doi:
1250	10.5194/acp-13-4057-2013
1251	Browne, E. C., Perring, A. E., Wooldridge, P. J., Apel, E., Hall, S. R., Huey, L. G.,
1252	Cohen, R. C. (2011). Global and regional effects of the photochemistry of
1253	CH3O2NO2: Evidence from ARCTAS. Atmospheric Chemistry and Physics,
1254	11(9), 4209–4219. doi: 10.5194/acp-11-4209-2011
1255	Carver, G., Brown, P., & Wild, O. (1997). The ASAD atmospheric chemistry inte-
1256	gration package and chemical reaction database. Computer Physics Communi-
1257	cations, $105(2)$, $197-215$. doi: $10.1016/S0010-4655(97)00056-8$
1258	Chipperfield, M. P. (2006). New version of the TOMCAT/SLIMCAT off-line chem-
1259	ical transport model: Intercomparison of stratospheric tracer experiments.
1260	Quarterly Journal of the Royal Meteorological Society, 132(617), 1179–1203.
1261	doi: 10.1256/qj.05.51
1262	Collins, W. J., Lamarque, JF., Schulz, M., Boucher, O., Eyring, V., Hegglin, M. I.,

1263	Smith, S. J. (2017). AerChemMIP: Quantifying the effects of chemistry
1264	and aerosols in CMIP6. Geoscientific Model Development, 10, 585–607. Re-
1265	trieved from http://www.geosci-model-dev-discuss.net/gmd-2016-139/
1266	doi: 10.5194/gmd-2016-139
1267	Dee, D. P., Uppala, S. M., Simmons, A. J., Berrisford, P., Poli, P., Kobayashi, S.,
1268	Vitart, F. (2011). The ERA-Interim reanalysis: Configuration and performance
1269	of the data assimilation system. Quarterly Journal of the Royal Meteorological
1270	Society $1.37(656)$ 553–597 doi: 10.1002/di.828
1270	Derwort $B = (2017)$ Intercomparison of chamical mechanisms for air quality policy.
1271	formulation and assessment under North American conditions
1272	to the Air 6 Weste Management Association $00(00)$ 10062247 2017 1202060
1273	the Atr \mathfrak{S} waste Management Association, $U(00)$, $10902247.2017.1292909$.
1274	Retrieved from https://www.tandionline.com/doi/full/10.1080/
1275	10962247.2017.1292969 doi: 10.1080/10962247.2017.1292969
1276	Dobber, M. R., Dirksen, R. J., Levelt, P. F., Van Den Oord, G. H., Voors, R. H.,
1277	Kleipool, Q., Rozemeijer, N. C. (2006). Ozone monitoring instrument
1278	calibration. $IEEE Transactions on Geoscience and Remote Sensing, 44(5),$
1279	1209-1238. doi: $10.1109/TGRS.2006.869987$
1280	Esenturk, E., Abraham, L., Archer-Nicholls, S., Mitsakou, C., Griffiths, P., & Pyle,
1281	J. (2018). Quasi-Newton Methods for Atmospheric Chemistry Simulations :
1282	Implementation in UKCA UM Vn10.8. Geoscientific Model Development, 11,
1283	1–31. doi: 10.5194/gmd-2018-32
1284	Evring, V., Bony, S., Meehl, G. A., Senjor, C. A., Stevens, B., Stouffer, R. J., &
1285	Taylor K E (2016) Overview of the Coupled Model Intercomparison Project
1205	Phase 6 (CMIP6) experimental design and organization <i>Geoscientific Model</i>
1200	Development 0(5) 1037-1058 doi: 10.5104/gmd-0-1037-2016
1287	Earsch & Dehenty D M Harrisida C Vandaulalia & Macintuma H L fr
1288	reflection, S., Domerty, R. M., fleaviside, C., vardoulakis, S., Macintyre, H. L., $\&$
1289	O Connor, F. M. (2018). The influence of model spatial resolution on sim-
1290	ulated ozone and nne particulate matter for Europe: Implications for health
1291	impact assessments. Atmospheric Chemistry and Physics, 18(8), 5765–5784.
1292	doi: 10.5194/acp-18-5765-2018
1293	Feng, L., Smith, S. J., Braun, C., Crippa, M., Gidden, M. J., & Hoesly, R. (2020).
1294	The generation of gridded emissions data for CMIP6. Geoscientific Model De-
1295	$velopment, \ 13, \ 461-482.$
1296	Gaudel, A., Cooper, O. R., Ancellet, G., Barret, B., Boynard, A., Burrows, J. P.,
1297	Lewis, A. (2018). Tropospheric Ozone Assessment Report: Present-
1298	day distribution and trends of tropospheric ozone relevant to climate and
1299	global atmospheric chemistry model evaluation. Elem Sci Anth. 6. doi:
1300	10.1525/elementa.291
1301	Giannakopoulos, C., Chipperfield, M. P., Law, K. S., & Pyle, J. A. (1999). Vali-
1302	dation and Intercomparison of Wet and Dry Model Deposition Schemes Using
1302	210Ph in a Clobal Three-Dimensional off-line Chemical Transport Model
1303	Journal of Geophysical Research 10/(D19) 23 761–23 784
1304	Coldatein A. H. & Colholly, J. E. (2007)
1305	Goldstein, A. H., & Galoally, I. E. (2007). Known and unexplored organic con-
1306	stituents in the earth's atmosphere. Environmental Science and Technology, $(1/5)$, 1514, 1501, 1.5 , 10, 1001, (-0.75) 476
1307	41(5), $1514-1521$. doi: $10.1021/es0/2476p$
1308	Grant, A., Archibald, A. T., Cooke, M. C., & Shallcross, D. E. (2010). Mod-
1309	elling the oxidation of seventeen volatile organic compounds to track yields
1310	of CO and CO2. Atmospheric Environment, $44(31)$, 3797–3804. Re-
1311	trieved from http://dx.doi.org/10.1016/j.atmosenv.2010.06.049 doi:
1312	10.1016/j.atmosenv. $2010.06.049$
1313	Guenther, A. B., Jiang, X., Heald, C. L., Sakulyanontvittaya, T., Duhl, T., Em-
1314	mons, L. K., & Wang, X. (2012, nov). The Model of Emissions of Gases and
1315	Aerosols from Nature version 2.1 (MEGAN2.1): an extended and updated
1316	framework for modeling biogenic emissions. <i>Geoscientific Model Development</i> .
1317	5, 1471-1492. Retrieved from http://www.geosci-model-dev.net/5/1471/

1318	2012 / doi: 10.5194/gmd-5-1471-2012
1319	Harrison, M. A., Barra, S., Borghesi, D., Vione, D., Arsene, C., & Iulian Olariu, R.
1320	(2005). Nitrated phenols in the atmosphere: A review. Atmospheric Environ-
1321	ment, 39(2), 231-248. doi: 10.1016/j.atmosenv.2004.09.044
1322	Hayman, G. D. (1997). Effects of Pollution Control on UV Exposure, AEA Tech-
1323	nology Final Report. prepared for the Department of Health on Contract
1324	121/6377, AEA Technology, Reference AEA/RCEC/22522001/R/002 IS-
1325	SUE1.
1326	Heald, C. L., & Kroll, J. H. (2020). The fuel of atmospheric chemistry: Toward a
1327	complete description of reactive organic carbon. Science Advances, $6(6)$, 1–9.
1328	doi: 10.1126/sciadv.aay8967
1329	Hodzic, A., Campuzano-Jost, P., Bian, H., Chin, M., Colarco, P. R., Day, D. A.,
1330	Jimenez, J. L. (2020). Characterization of organic aerosol across the global
1331	remote troposphere: A comparison of ATom measurements and global chem-
1332	istry models. Atmospheric Chemistry and Physics, $20(8)$, $4607-4635$. doi:
1333	10.5194/acp-20-4607-2020
1334	Hoerling, M. P., Schaack, T. K., & Lenzen, A. J. (1993). A Global Analysis of
1335	Stratospheric-Tropospheric Exchange during Northern Winter. Monthly
1336	$Weather \ Review, \ 121(1), \ 162-172.$
1337	Hoesly, R. M., Smith, S. J., Feng, L., Klimont, Z., Janssens-Maenhout, G., Pitka-
1338	nen, T., Zhang, Q. (2018). Historical (1750-2014) anthropogenic emis-
1339	sions of reactive gases and aerosols from the Community Emissions Data
1340	System (CEDS). Geoscientific Model Development, 11(1), 369–408. doi:
1341	10.5194/gmd-11-369-2018
1342	Hood, C., MacKenzie, I., Stocker, J., Johnson, K., Carruthers, D., Vieno, M., &
1343	Doherty, R. (2018). Air quality simulations for London using a coupled
1344	regional-to-localmodelling system. Atmospheric Chemistry and Physics, 18,
1345	11221–11245. doi: 10.5194/acp-2017-1202
1346	Hoyle, C. R., Marécal, V., Russo, M. R., Allen, G., Arteta, J., Chemel, C., Zeng,
1347	G. (2011). Representation of tropical deep convection in atmospheric mod-
1348	els - Part 2: Iracer transport. Atmospheric Chemistry and Physics, 11(15),
1349	8103-8131. doi: 10.5194/acp-11-8103-2011
1350	Huang, G., Brook, R., Orippa, M., Janssens-Maennout, G., Schleberle, C., Dore,
1351	non methane veletile errorie compounds. A global gridded data get for
1352	$1070\ 2012$ Atmospheric Chemistry and Physics $17(12)\ 7683-7701$ doi:
1353	1010^{-2012} . $1010000000000000000000000000000000000$
1354	In $M = k$ Clemitshaw K (2000) Ozone and other secondary photochemi-
1355	cal pollutants: chemical processes governing their formation in the planetary
1257	boundary layer Atmospheric Environment 3/(16) 2499–2527 Retrieved from
1358	http://linkinghub.elsevier.com/retrieve/pii/S1352231099004781 doi:
1359	10.1016/S1352-2310(99)00478-1
1360	Jenkin M Saunders S M & Pilling M J (1997 jan) The tropospheric
1361	degradation of volatile organic compounds: a protocol for mechanism de-
1362	velopment. Atmospheric Environment, 31(1), 81–104. Retrieved from
1363	http://linkinghub.elsevier.com/retrieve/pii/S1352231096001057
1364	doi: 10.1016/S1352-2310(96)00105-7
1365	Jenkin, M., Watson, L., Utembe, S., & Shallcross, D. (2008, oct). A Common
1366	Representative Intermediates (CRI) mechanism for VOC degradation. Part
1367	1: Gas phase mechanism development. Atmospheric Environment, 42, 7185–
1368	7195. Retrieved from http://linkinghub.elsevier.com/retrieve/pii/
1369	S1352231008006742 doi: 10.1016/j.atmosenv.2008.07.028
1370	Jenkin, M. E., Khan, M. A., Shallcross, D. E., Bergström, R., Simpson, D., Mur-
1371	phy, K. L., & Rickard, A. R. (2019). The CRI v2.2 reduced degradation
1070	scheme for isoprene Atmospheric Environment 212 (May) 172–182 doi:

1373	10.1016/j.atmosenv.2019.05.055
1374	Jenkin, M. E., Saunders, S. M., Derwent, R. G., & Pilling, M. J. (2002, oct). De-
1375	velopment of a reduced speciated VOC degradation mechanism for use in
1376	ozone models. Atmospheric Environment, 36(30), 4725–4734. Retrieved from
1377	http://linkinghub.elsevier.com/retrieve/pii/S1352231002005630 doi:
1378	10.1016/S1352-2310(02)00563-0
1379	Jenkin, M. E., Saunders, S. M., Wagner, V., & Pilling, M. J. (2003, feb). Proto-
1380	col for the development of the Master Chemical Mechanism, MCM v3 (Part
1381	B): tropospheric degradation of aromatic volatile organic compounds. At-
1382	mospheric Chemistry and Physics, 3(1), 181–193. Retrieved from http://
1383	www.atmos-chem-phys.net/3/181/2003/ doi: 10.5194/acp-3-181-2003
1384	Jenkin, M. E., Valorso, R., Aumont, B., & Rickard, A. R. (2019). Estimation of
1385	rate coefficients and branching ratios for reactions of organic peroxy radicals
1386	for use in automated mechanism construction. Atmospheric Chemistry and
1387	<i>Physics</i> (19), 7691–7717. doi: 10.5194/acp-2019-44
1388	Jenkin, M. E., Valorso, R., Aumont, B., Rickard, A. R., & Wallington, T. J. (2018a).
1389	Estimation of rate coefficients and branching ratios for gas-phase reactions of
1390	OH with aliphatic organic compounds for use in automated mechanism con-
1391	struction (Vol. 18) (No. 13). doi: 10.5194/acp-18-9297-2018
1392	Jenkin, M. E., Valorso, R., Aumont, B., Rickard, A. R., & Wallington, T. J.
1393	(2018b). Estimation of rate coefficients and branching ratios for gas-phase
1394	reactions of OH with aromatic organic compounds for use in automated mech-
1395	anism construction. Atmospheric Chemistry and Physics, 18(13), 9329–9349.
1396	doi: 10.5194/acp-18-9329-2018
1397	Jenkin, M. E., Wyche, K. P., Evans, C. J., Carr, T., Monks, P. S., Alfarra, M. R.,
1398	Rickard, A. R. (2012). Development and chamber evaluation of the MCM
1399	v3.2 degradation scheme for β -caryophyllene. Atmospheric Chemistry and
1400	<i>Physics</i> , $12(11)$, 5275–5308. doi: 10.5194/acp-12-5275-2012
1401	Jenkin, M. E., Young, J. C., & Rickard, A. R. (2015). The MCM v3.3.1 degrada-
1402	tion scheme for isoprene. Atmospheric Chemistry and Physics, 15(20), 11433-
1403	11459. doi: $10.5194/acp-15-11433-2015$
1404	Jimenez, J. L., Canagaratna, M. R., Donahue, N. M., Prevot, a. S. H., Zhang,
1405	Q., Kroll, J. H., Worsnop, D. R. (2009, dec). Evolution of organic
1406	aerosols in the atmosphere. Science (New York, N.Y.), 326(5959), 1525–9.
1407	Retrieved from http://www.ncbi.nlm.nih.gov/pubmed/20007897 doi:
1408	10.1126/science.1180353
1409	Khan, M. A., Clements, J., Lowe, D., McFiggans, G., Percival, C. J., & Shall-
1410	cross, D. E. (2019). Investigating the behaviour of the CRI-MECH gas-
1411	phase chemistry scheme on a regional scale for different seasons using the
1412	WRF-Chem model. Atmospheric Research, 229(June), 145–156. Re-
1413	trieved from https://doi.org/10.1016/j.atmosres.2019.06.021 doi:
1414	10.1016/j.atmosres.2019.06.021
1415	Khan, M. A. H., Cooke, M. C., Utembe, S. R., Archibald, A. T., Derwent, R. G.,
1416	Xiao, P., Shallcross, D. E. (2015). Global modeling of the nitrate rad-
1417	ical (NO3) for present and pre-industrial scenarios. Atmospheric Research,
1418	164-165(3), 347-357. doi: $10.1016/j.atmosres.2015.06.006$
1419	Lightfoot, P. D., Cox, R. A., Crowley, J. N., Destriau, M., Hayman, G. D., Jenkin,
1420	M. E., Zabel, F. (1992). Organic peroxy radicals: kinetics, spectroscopy
1421	and tropospheric chemistry. Atmospheric Environment, $26(10)$, $1805-1961$.
1422	Lowe, D., Archer-Nicholls, S., Morgan, W., Allan, J., Utembe, S., Ouyang, B.,
1423	McFiggans, G. (2015). WRF-chem model predictions of the regional impacts
1424	of N2O5 heterogeneous processes on nighttime chemistry over north-western
1425	Europe. Atmospheric Chemistry and Physics, 15, 1385–1409. Retrieved
1426	trom http://www.atmos-chem-phys-discuss.net/14/20883/2014/ doi:
1427	10.5194/acp-15-1385-2015

Mann, G. W., Carslaw, K. S., Spracklen, D. V., Ridley, D. A., Manktelow, P. T., & 1428 (2010).Description and evaluation of GLOMAP-mode Chipperfield, M. P. 1429 : a modal global aerosol microphysics model for the UKCA composition-1430 climate model. Geoscientific Model Development, 519–551. doi: 10.5194/ 1431 gmd-3-519-2010 1432 Mcfiggans, G., Mentel, T. F., Wildt, J., Pullinen, I., Kang, S., Kleist, E., ... 1433 (2019).Kiendler-scharr, A. Secondary organic aerosol reduced by mixture 1434 of atmospheric vapours. Nature, 0–6. Retrieved from http://dx.doi.org/ 1435 10.1038/s41586-018-0871-y doi: 10.1038/s41586-018-0871-y 1436 McGillen, M. R., Carter, W. P., Mellouki, A., J. Orlando, J., Picquet-Varrault, B., 1437 & J. Wallington, T. (2020). Database for the kinetics of the gas-phase atmo-1438 spheric reactions of organic compounds. Earth System Science Data, 12(2), 1439 1203–1216. doi: 10.5194/essd-12-1203-2020 1440 Monks, P. S. (2005, may). Gas-phase radical chemistry in the troposphere. *Chemi*-1441 cal Society reviews, 34(5), 376-95. Retrieved from http://www.ncbi.nlm.nih 1442 .gov/pubmed/15852151 doi: 10.1039/b307982c 1443 Monks, P. S., Archibald, A. T., Colette, A., Cooper, O., Coyle, M., Derwent, R., ... 1444 Williams, M. L. (2015). Tropospheric ozone and its precursors from the urban 1445 to the global scale from air quality to short-lived climate forcer. Atmospheric 1446 Chemistry and Physics, 15(15), 8889–8973. doi: 10.5194/acp-15-8889-2015 1447 Morgenstern, O., Braesicke, P., O'Connor, F. M., Bushell, A. C., Johnson, C. E., 1448 Osprey, S. M., & Pyle, J. A. (2009). Evaluation of the new UKCA climatecomposition model – Part 1: The stratosphere. Geoscientific Model Develop-1450 ment, 2, 43–57. Retrieved from http://www.geosci-model-dev.net/2/43/ 1451 2009/gmd-2-43-2009.html doi: 10.5194/gmd-2-43-2009 1452 Mulcahy, J. P., Johnson, C., Jones, C., Povey, A., Sellar, A., Scott, C. E., ... Yool, 1453 A. (2020). Description and evaluation of aerosol in UKESM1 and HadGEM3-1454 GC3.1 CMIP6 historical simulations. Geoscientific Model Development. 1455 Mulcahy, J. P., Jones, C., Sellar, A., Johnson, B., Boutle, I. A., Jones, A., ... Mc-1456 Coy, D. T. (2018). Improved Aerosol Processes and Effective Radiative Forcing 1457 in HadGEM3 and UKESM1. Journal of Advances in Modeling Earth Systems, 1458 10(11), 2786-2805. doi: 10.1029/2018MS0014641459 Neu, J. L., Prather, M. J., & Penner, J. E. (2007). Global atmospheric chemistry: 1460 Integrating over fractional cloud cover. Journal of Geophysical Research Atmo-1461 spheres, 112(11), 1–12. doi: 10.1029/2006JD008007 1462 Newsome, B., & Evans, M. (2017).Impact of uncertainties in inorganic chemi-1463 cal rate constants on tropospheric composition and ozone radiative forcing. At-1464 mospheric Chemistry and Physics, 17(23), 14333–14352. doi: 10.5194/acp-17 1465 -14333-20171466 Novelli, A., Kaminski, M., Rolletter, M., Acir, I. H., Bohn, B., Dorn, H. P., ... 1467 Fuchs, H. (2018).Evaluation of OH and $\ \ HO_2\$ concentrations and 1468 their budgets during photooxidation of 2-methyl-3-butene-2-ol (MBO) in 1469 the atmospheric simulation chamber SAPHIR. Atmospheric Chemistry and 1470 *Physics*, 18(15), 11409–11422. doi: 10.5194/acp-18-11409-2018 1471 O'Connor, F. M., Johnson, C. E., Morgenstern, O., Abraham, N. L., Braesicke, P., 1472 Dalvi, M., ... Pyle, J. A. (2014).Evaluation of the new UKCA climate-1473 composition model-Part 2: The troposphere. Geoscientific Model Development, 1474 7(1), 41-91. doi: 10.5194/gmd-7-41-2014 1475 Olivier, J. G. J., Peters, J., Granier, C., Petron, G., Muller, J.-F., & Wallens, S. 1476 (2003).Present and future surface emissions of atmospheric compounds. 1477 POET Report #3, EU project EVK2-1999-00011. 1478 Orlando, J. J., & Tyndall, G. S. (2012).Laboratory studies of organic peroxy 1479 radical chemistry: An overview with emphasis on recent issues of atmo-1480 spheric significance. Chemical Society Reviews, 41(19), 6294–6317. doi: 1481 10.1039/c2cs35166h 1482

Pétron, G., Granier, C., Khattatov, B., Lamarque, J. F., Yudin, V., Müller, J. F., 1483 Inverse modeling of carbon monoxide surface emissions & Gille, J. (2002).1484 using Climate Monitoring and Diagnostics Laboratory network observa-1485 tions. Journal of Geophysical Research Atmospheres, 107(24), 1–23. doi: 10.1029/2001JD001305 1487 Porter, W. C., Safieddine, S. A., & Heald, C. L. (2017). Impact of aromatics and 1488 monoterpenes on simulated tropospheric ozone and total OH reactivity. Atmo-1489 spheric Environment, 169, 250–257. doi: 10.1016/j.atmosenv.2017.08.048 1490 Sander, R. (2015). Compilation of Henry's law constants (version 4.0) for water as 1491 solvent. Atmospheric Chemistry and Physics, 15, 4399-4981. doi: 10.5194/acp 1492 -15 - 4399 - 20151493 Sander, S. P., Friedl, R. R., Barker, J. R., Golden, D. M., Kurylo, M. J., Sciences, 1494 G. E., ... Orkin, V. L. (2011). Chemical Kinetics and Photochemical Data 1495 for Use in Atmospheric Studies Evaluation Number 17 NASA Panel for Data 1496 Evaluation :. (17). 1497 Saunders, S. M., Jenkin, M. E., Derwent, R. G., & Pilling, M. J. (2003, feb). Pro-1498 tocol for the development of the Master Chemical Mechanism, MCM v3 (Part 1499 A): tropospheric degradation of non-aromatic volatile organic compounds. At-1500 mospheric Chemistry and Physics, 3, 161–180. doi: 10.5194/acp-3-161-2003 1501 Schultz, M. G., Schröder, S., Lyapina, O., Cooper, O. R., Galbally, I., 1502 (2017).Petropavlovskikh, I., ... Zhiqiang, M. Tropospheric Ozone Assess-1503 ment Report: Database and metrics data of global surface ozone observations. 1504 Elementa, 5. doi: 10.1525/elementa.244 1505 Sellar, A. A., Jones, C. G., Mulcahy, J., Tang, Y., Yool, A., Wiltshire, A., ... 1506 Zerroukat, M. (2019).UKESM1: Description and evaluation of the UK 1507 Earth System Model. Journal of Advances in Modeling Earth Systems. doi: 1508 10.1029/2019ms001739 1509 Sillman, S. (1999, jun). The relation between ozone, NOx and hydrocarbons in 1510 urban and polluted rural environments. Atmospheric Environment, 33(12), 1511 Retrieved from http://linkinghub.elsevier.com/retrieve/ 1821 - 1845.1512 pii/S1352231098003458 doi: 10.1016/S1352-2310(98)00345-8 1513 Sindelarova, K., Granier, C., Bouarar, I., Guenther, A., Tilmes, S., Stavrakou, T., 1514 ... Knorr, W. (2014). Global data set of biogenic VOC emissions calculated 1515 by the MEGAN model over the last 30 years. Atmospheric Chemistry and 1516 *Physics*, 14(17), 9317–9341. doi: 10.5194/acp-14-9317-2014 1517 Spracklen, D. V., Carslaw, K. S., Kulmala, M., Kerminen, V. M., Mann, G. W., & 1518 Sihto, S. L. (2006).The contribution of boundary layer nucleation events 1519 to total particle concentrations on regional and global scales. Atmospheric 1520 Chemistry and Physics, 6(12), 5631–5648. doi: 10.5194/acp-6-5631-2006 1521 Stock, Z. S., Russo, M. R., & Pyle, J. A. (2014).Representing ozone extremes 1522 in European megacities: The importance of resolution in a global chemistry 1523 climate model. Atmospheric Chemistry and Physics, 14(8), 3899–3912. doi: 1524 10.5194/acp-14-3899-2014 1525 Stockwell, W. R., Middleton, P., & Chang, J. S. (1990).The Second Generation 1526 Regional Acid Deposition Model Chemical Mechanism for Regional Air Qual-1527 ity Modeling The most important in the gas Alkanes with between. Journal of 1528 Geophysical Research, 95(D10), 16,343–16,367. 1529 Telford, P. J., Abraham, N. L., Archibald, A. T., Braesicke, P., Dalvi, M., Mor-1530 genstern, O., ... Pyle, J. A. (2013).Implementation of the Fast-JX 1531 Photolysis scheme into the UKCA component of the MetUM chemistry 1532 climate model. Geoscientific Model Development, 6, 161–177. doi: 1533 10.5194/gmdd-5-3217-2012 1534 Telford, P. J., Braesicke, P., Morgenstern, O., & Pyle, J. A. (2008).Technical 1535 note: Description and assessment of a nudged version of the new dynamics 1536 Atmospheric Chemistry and Physics, 8(6), 1701–1712. Unified Model. doi: 1537

1538	10.5194/acp-8-1701-2008
1539	Tsigaridis, K., Daskalakis, N., Kanakidou, M., Adams, P. J., Artaxo, P., Bahadur,
1540	R., Zhang, X. (2014). The AeroCom evaluation and intercomparison of
1541	organic aerosol in global models. Atmospheric Chemistry and Physics, 14(19),
1542	10845–10895. doi: 10.5194/acp-14-10845-2014
1543	Tunved, P., Ström, J., & Hansson, HC. (2004). An investigation of processes con-
1644	trolling the evolution of the boundary laver aerosol size distribution properties
1544	at the Swedish background station Aspyreten Atmospheric Chemistry and
1545	Physics Discussions $\lambda(4)$ 4507–4543 doi: 10.5194/acnd-4.4507.2004
1540	Turdell C S Cov P A Cranier C Leseloux P Moortget C K Pilling
1547	M J Wallington T J (2001) Atmospheric chargington of apoll argonic
1548	M. J., Wallington, I. J. (2001). Atmospheric chemistry of small organic
1549	peroxy radicals. Journal of Geophysical Research, $100(D11)$, $12157-12182$.
1550	Utembe, S., Cooke, M., Archibald, A. T., Jenkin, M., Derwent, R., & Shallcross, D.
1551	(2010, apr). Using a reduced Common Representative Intermediates (CRIv2-
1552	R5) mechanism to simulate tropospheric ozone in a 3-D Lagrangian chemistry
1553	transport model. Atmospheric Environment, 44(13), 1609–1622. Retrieved
1554	from http://linkinghub.elsevier.com/retrieve/pii/S1352231010001056
1555	doi: $10.1016/j.atmosenv.2010.01.044$
1556	Utembe, S. R., Watson, L. A., Shallcross, D. E., & Jenkin, M. E. (2009, apr).
1557	A Common Representative Intermediates (CRI) mechanism for VOC
1558	degradation. Part 3: Development of a secondary organic aerosol mod-
1559	ule. $Atmospheric Environment, 43(12), 1982–1990.$ Retrieved from
1560	http://linkinghub.elsevier.com/retrieve/pii/S1352231009000284
1561	doi: 10.1016/j.atmosenv.2009.01.008
1562	Van Der Werf, G. R., Randerson, J. T., Giglio, L., Van Leeuwen, T. T., Chen,
1563	Y., Rogers, B. M., Kasibhatla, P. S. (2017). Global fire emissions esti-
1564	mates during 1997-2016. Earth System Science Data, 9(2), 697–720. doi:
1565	10.5194/essd-9-697-2017
1566	Van Marle, M. J., Kloster, S., Magi, B. I., Marlon, J. R., Daniau, A. L., Field,
1567	R. D., Van Der Werf, G. R. (2017). Historic global biomass burning emis-
1568	sions for CMIP6 (BB4CMIP) based on merging satellite observations with
1569	proxies and fire models $(1750-2015)$. Geoscientific Model Development, $10(9)$.
1570	3329–3357. doi: 10.5194/gmd-10-3329-2017
1671	Von Schneidemesser E. Monks P.S. Allan J.D. Bruhwiler L. Forster P
1572	Fowler, D.,, Sutton, M. A. (2015). Chemistry and the Linkages between
1573	Air Quality and Climate Change Chemical Reviews 115(10) 3856–3897 doi:
1575	10 1021/acs chemrev 5b00089
1574	von Clarmann T & Clatthor N (2010) The application of mean averaging kernels
1575	to mean trace are distributions. Atmospheric Measurement Techniques Discus
1576	sione 1 11 doi: 10 5104/amt 2010 61
1577	stons, 1–11. doi: 10.5194/ann-2019-01 von Classer P. & Creater P. J. (2004 ann) Model study of multiplese DMS
1578	von Grasow, n., & Orutzen, r. J. (2004, apr). Model study of multiphase DMS
1579	oxidation with a focus of halogens. Atmospheric Chemistry and Physics, $4(3)$,
1580	589-608. Retrieved from http://www.atmos-chem-phys.net/4/589/2004/
1581	doi: 10.5194/acp-4-589-2004
1582	Walters, D., Baran, A. J., Boutle, I., Brooks, M., Earnshaw, P., Edwards, J.,
1583	Zerroukat, M. (2019). The Met Office Unified Model Global Atmosphere
1584	7.0/7.1 and JULES Global Land 7.0 configurations. <i>Geoscientific Model Devel</i> -
1585	opment, 12(5), 1909-1963. doi: 10.5194/gmd-12-1909-2019
1586	Wang, Y., Jacob, D. J., & Logan, J. A. (1998). Global simulation of tropospheric
1587	O3-NOx-hydrocarbon chemistry - 3. Origin of tropospheric ozone and effects
1588	of nonmethane hydrocarbons. Journal of Geophysical Research: Atmospheres,
1589	103(3339), 10757-10767. doi: $10.1029/98$ jd00156
1590	Waters, J. W., Froidevaux, L., Harwood, R. S., Jarnot, R. F., Pickett, H. M., Read,
1591	W. G., Walch, M. J. (2006). The Earth Observing System Microwave Limb
1592	Sounder (EOS MLS) on the aura satellite. IEEE Transactions on Geoscience

1593	and Remote Sensing, 44(5), 1075–1092. doi: 10.1109/TGRS.2006.873771
1594	Watson, L., Shallcross, D., Utembe, S., & Jenkin, M. (2008, oct). A Common
1595	Representative Intermediates (CRI) mechanism for VOC degradation. Part 2:
1596	Gas phase mechanism reduction. Atmospheric Environment, 42(31), 7196–
1597	7204. Retrieved from http://linkinghub.elsevier.com/retrieve/pii/
1598	S1352231008006845 doi: 10.1016/j.atmosenv.2008.07.034
1599	Weber, J., Archibald, A., Griffiths, P., Archer-Nicholls, S., Berndt, T., Jenkin, M.,
1600	Knote, C. (2020). CRI-HOM: A novel chemical mechanism for simu-
1601	lating Highly Oxygenated Organic Molecules (HOMs) in global chemistry-
1602	aerosol-climate models. Atmospheric Chemistry and Physics, 20, 1–31. doi:
1603	10.5194/acp-2020-154
1604	Wesley, M. L. (1989). Parameterization of Surface Resistances to Gaseous Dry
1605	Deposition in Regional-Scale Numerical Models. Atmospheric Environment,
1606	23(6), 1293-1304. doi: 10.1016/S0950-351X(05)80241-1
1607	Wild, O., & Prather, M. J. (2000). Excitation of the primary tropospheric chem-
1608	ical mode in a global three-dimensional model. Journal of Geophysical Re-
1609	search, 105(D20), 24647. Retrieved from http://doi.wiley.com/10.1029/
1610	2000JD900399 doi: 10.1029/2000JD900399
1611	Wild, O., & Prather, M. J. (2006). Global tropospheric ozone modeling: Quantifying
1612	errors due to grid resolution. Journal of Geophysical Research Atmospheres,
1613	111(11), 1-14. doi: $10.1029/2005JD006605$
1614	Wild, O., Zhu, X. I. N., & Prather, M. J. (2000). Fast-J: Accurate Simulation of In-
1615	and Below-Cloud Photolysis in Tropospheric Chemical Models. Journal of At-
1616	mospheric Chemistry, 37, 249-282.
1617	minoral dust in the Hadley Contra climate model — <i>Journal of Combusical Re</i>
1618	search 106(D16) 18 155–18 166
1619	Young P. I. Archibald A. T. Bowman K. W. Lamaraque IF. Stevenson D. S.
1620	Tilmes S Zeng C (2013) Pre-industrial to end 21st century projections
1622	of tropospheric ozone from the Atmospheric Atmospheric Chemistry and Cli-
1623	mate Model Intercomparison Project (ACCMIP). Atmospheric Chemistry and
1624	<i>Physics</i> , 13(13), 2063–2090, doi: 10.5194/acp-13-5277-2013
1625	Young, P. J., Naik, V., Fiore, A. M., Gaudel, A., Guo, J., Lin, M. Y., Zeng,
1626	G. (2018). Tropospheric Ozone Assessment Report: Assessment of
1627	global-scale model performance for global and regional ozone distribu-
1628	tions, variability, and trends. $Elem Sci Anth, 6(1), 10.$ Retrieved from
1629	http://www.elementascience.org/article/10.1525/elementa.265/ doi:
1630	10.1525/elementa.265
1631	Zhang, K., Wan, H., Liu, X., Ghan, S. J., Kooperman, G. J., Ma, P. L.,
1632	Lohmann, U. (2014). Technical note: On the use of nudging for aerosol-
1633	climate model intercomparison studies. Atmospheric Chemistry and Physics,
1634	14(16), 8631-8645. doi: $10.5194/acp-14-8631-2014$
1635	Ziemke, J. R., Chandra, S., Duncan, B. N., Froidevaux, L., Bhartia, P. K., Levelt,
1636	P. F., & Waters, J. W. (2006). Tropospheric ozone determined from Aura OMI
1637	and MLS: Evaluation of measurements and comparison with the Global Mod-
1638	eling Initiative's Chemical Transport Model. Journal of Geophysical Research
1639	Atmospheres, 111(19), 1–18. doi: 10.1029/2006JD007089
1640	Ziemke, J. R., Oman, L. D., Strode, S. A., Douglass, A. R., Olsen, M. A., McPeters,
1641	R. D., Taylor, S. L. (2019). Trends in Global Tropospheric Ozone Inferred
1642	rrom a Composite Record of TOMS/OMI/MLS/OMPS Satellite Measurements
1643	and the MEKKA-2 GMI Simulation. Atmospheric Chemistry and Physics, 19, 2257–2260, doi: 10.5104/ger. 2019.716
1644	3237-3209. doi: 10.3194/acp-2018-710

Supporting Information for "The Common Representative Intermediates Mechanism version 2 in the United Kingdom Chemistry and Aerosols Model"

S. Archer-Nicholls¹, N. L. Abraham^{1,2}, Y. M. Shin¹, J. Weber¹, M. R.

Russo^{1,2}, D. Lowe³, S. Utembe⁴, F. M. O'Connor⁵, B. Kerridge^{6,7},

B.Latter^{6,7}, R. Siddans^{6,7}, M. Jenkin⁸, O. Wild⁹, A. T. Archibald^{1,2}

^[1]Department of Chemistry, University of Cambridge, Cambridge, UK ^[2]National Centre for Atmospheric Science, Department of Chemistry, University of Cambridge, Cambridge, UK ^[3]University of Manchester, Manchester, UK ^[4]University of Melbourne, Melbourne, Australia ^[5]Met Office Hadley Centre, Exeter, UK ^[6]Remote Sensing Group, STFC Rutherford Appleton Laboratory, UK ^[7]NERC National Centre for Earth Observation, Leicester, UK ^[8]Atmospheric Chemistry Services, Okehampton, Devon, UK ^[9]Lancaster University, Lancaster, UK

Contents of this file

- 1. S1. Comparison of mechanism runtime
- 2. S2. Details of the CRI-Strat mechanism
- 3. S3. Changes to StratTrop mechanism to conserve Nitrogen
- 4. S4. Differences in reaction rates coefficients between mechanisms

Corresponding author: A. T. Archibald, Department of Chemistry, University of Cambridge, Cambridge, UK. (ata27@cam.ac.uk) 5. S5. Supporting Analysis of ozone

6. S6. Supporting Analysis of Nitrogen Containing Species

7. Figure S1. Difference in zonal mean HONO₂, NO_y, NO_x, NO_z and O₃ between StratTrop_NCon and StratTrop_Orig.

8. Figure S2. Comparison of selected inorganic bimolecular reaction rates in CRIv2 to equivalent reactions in StratTrop over an atmospherically relevant temperature range. Blue line is CRI rate, orange line StratTrop, both using scale on left axis, and black dot-dashed line shows ratio of CRI/StratTrop.

9. Figure S3. Comparison of inorganic bimolecular reaction rates in CRIv2 to equivalent reactions in StratTrop over an atmospherically relevant temperature range. Blue line is CRI rate, orange line StratTrop, both using scale on left axis, and black dot-dashed line shows ratio of CRI/StratTrop.

10. Figure S4. Comparison of selected organic bimolecular reaction rates in CRIv2 to equivalent reactions in StratTrop over an atmospherically relevant temperature range. Blue line is CRI rate, orange line StratTrop, both using scale on left axis, and black dot-dashed line shows ratio of CRI/StratTrop.

11. Figure S5. Ratios of selected termolecular reaction rate coefficients in CRIv2 to equivalent reactions in StratTrop over atmospherically relevant temperature and pressure ranges.

12. Figure S6. Average surface ozone concentrations from rural sites on the TOAR network over the whole world and biases between StratTrop and CRI-Strat model simulations and observations from TOAR network using data from 2010 to 2014, averaged over whole year (a-c), June to August (d-f) and December to February (g-i).

13. Figure S7. Average surface ozone concentrations from rural sites on the TOAR network over North America and biases between StratTrop and CRI-Strat model simulations and observations from TOAR network using data from 2010 to 2014, averaged over whole year (a-c), June to August (d-f) and December to February (g-i).

14. Figure S8. Average surface ozone concentrations from rural sites on the TOAR network over Europe and biases between StratTrop and CRI-Strat model simulations and observations from TOAR network using data from 2010 to 2014, averaged over whole year (a-c), June to August (d-f) and December to February (g-i). Mean bias between model and observations given in titles of panels b, c, e, f, h, and i.

15. Figure S9. Average surface ozone concentrations from rural sites on the TOAR network over East Asia and biases between StratTrop and CRI-Strat model simulations and observations from TOAR network using data from 2010 to 2014, averaged over whole year (a-c), June to August (d-f) and December to February (g-i). Mean bias between model and observations given in titles of panels b, c, e, f, h, and i.

16. Figure S10. Tropospheric ozone column (DU) in StratTrop mechanism in DJF (a) and JJA (b). Difference in tropospheric ozone column between CRI-Strat and StratTrop in DJF (c) and JJA (f). Difference in tropospheric ozone column between CRI-Emiss_ST and StratTrop in DJF(d) and JJA (g). Difference in tropospheric ozone column between CRI-Emiss_C2C3 and StratTrop in DJF(e) and JJA (h).

17. Figure S11. Difference in chemical production of Ox averaged over the lower 1km of the atmosphere between CRI-Strat and StratTrop (a), CRI_Emiss_ST and StratTrop (b), and CRI_Emiss_C2C3 and StratTrop (c). Difference in chemical loss of Ox averaged over lower 1 km of the atmosphere between CRI-Strat and StratTrop (d), CRI_Emiss_ST

and StratTrop (e), and CRI_Emiss_C2C3 and StratTrop (f). Difference in deposition of Ox averaged over lower 1 km of the atmosphere between CRI-Strat and StratTrop (g), CRI_Emiss_ST and StratTrop (h), and CRI_Emiss_C2C3 and StratTrop (i). Difference in deposition of Ox averaged over lower 1 km of the atmosphere between CRI-Strat and StratTrop (j), CRI_Emiss_ST and StratTrop (k), and CRI_Emiss_C2C3 and StratTrop (l).

18. Figure S12. Zonal mean flux though the $O(^{1}D) + H_{2}O$ reaction in StratTrop (a), and difference in zonal mean flux between CRI-Strat and StratTrop (b), CRI_Emiss_ST and StratTrop (c), and CRI_Emiss_C2C3 and StratTrop (d).

19. Figure S13. Mean differences in NO_x (a-c), NO_y (d-f), and NO_z (g-i) over the lower 1 km of the atmosphere between CRI-Strat and StratTrop (a, d, g), CRI_Emiss_ST and StratTrop (b, e and h), and CRI_Emiss_C2C3 and StratTrop (c, f, i).

20. Figure S14. Zonal mean flux through the OH + NO2 + M reaction in StratTrop (a), and difference in zonal mean flux between CRI-Strat and StratTrop (b), CRI_Emiss_ST and StratTrop (c), and CRI_Emiss_C2C3 and StratTrop (d).

21. Table S1. Structural codes and functional types used to describe CRI intermediates. 22. Table S2. Species treated by the CRI-Strat chemistry mechanism. Where the name of the species has been changed from that used by the original CRIv2-R5 mechanism in order to follow UKCA standards, the original name is also given. Species with 1 in the Strat column have been added to make the mechanism suitable for running in the stratosphere, species with 1 in the Aero column are only activated if running with

GLOMAP-MODE.

23. Table S3. Photolysis reactions in the CRI-Strat chemical mechanism. Reactions with 1 in the Strat column have been added to make the mechanism suitable for running

in the stratosphere, reactions with 1 in the Aero column are only activated if running with GLOMAP-MODE. References for cross section data for reactions can be found in Telford et al. (2013). ?Reaction rates calculated online from the cross section data are multiplied by the scaling factor.

24. Table S4. Bimolecular reactions in CRI-Strat chemical mechanism. Reactions with 1 in the Strat column have been added to make the mechanism suitable for running in the stratosphere, reactions with 1 in the Aero column are only activated if running with GLOMAP-MODE. Temperature dependent reaction rate coefficients k_0 , a_0 and b_0 are given for the equation $k(T) = k_0 (T/300)^{a_0} \exp(-b_0/T)$, where T is temperature in K.

25. Table S5. Termolecular reactions in CRI-Strat chemical mechanism. Reactions with 1 in the Strat column have been added to make the mechanism suitable for running in the stratosphere, reactions with 1 in the Aero column are only activated if running with GLOMAP-MODE. Temperature and pressure dependent reaction rate coefficients k_1 , a_1 , b_1 , k_2 , a_2 , b_2 and F are given for the equation k(T) = $(\frac{k_0(T)[M]}{1+k_0(T)[M]/k_i(T)})F_c^{(1+(\log(k_0(T)/k_i))^2)^{-1}}$, where T is temperature in K; [M] is the total number density in molecules cm⁻³; $k_0 = k_1(T/300)^{a_1}\exp(-b_1/T)$ is the low pressure limit rate coefficient and $k_i = k_2(T/300)^{a_2}\exp(-b_2/T)$ is the infinite pressure limit rate coefficient. If $k_2 = 0$ then $k(T) = k_0(T)[M]$; if $k_1 = 0$ then $k(T) = k_i(T)$. The broadening term F_c is calculated from F as follows: if 0 > F < 1 then $F_c = F$; if F > 1 then $F_c = \exp(-F/T)$; if F = 0 then the F_c term in the equation is ignored.

26. Table S6. Overview of tropospheric Ox burden, lifetime, ozone production efficiency (OPE), chemical production, chemical loss, deposition and inferred stratosphere to troposphere transfer. Values in brackets give fraction of total chemical production for

the production terms and fraction of total losses $(L_{Ox} + D_{Ox})$ for the loss and deposition terms.

27. Table S7. Overview of air mass weighted OH concentration, CO burden and CO lifetime.

28. Table S8. Overview of tropospheric oxidised nitrogen burdens (fraction of total NO_y in brackets), tropospheric oxidised nitrogen emission and deposition fluxes, stratosphere-troposphere transfer (STT) and NO_y lifetime in the troposphere (fraction of total NO_y deposition in brackets).

Additional Supporting Information (Files uploaded separately)

- 1. Table S2. CRI-Strat_species.xcls
- 2. Table S3. CRI-Strat_photol.xcls
- 3. Table S4. CRI-Strat_bimol.xcls
- 4. Table S5. CRI-Strat_termol.xcls

Introduction

This supplement provides additional data, results and analysis to support the main paper. It starts in Section S1 with a comparison of computational costs for the CRI-Strat and StratTrop mechanisms in UKCA run on the same model architecture. In then documents details of the CRI-Strat mechanism in Section S2. Here, we provide supporting tables in excel files which contain the entire CRI-Strat mechanism as used in UKCA. These tables were generated directly from the source code used by the model. Section S3 shows a comparison between the standard version of StratTrop and the version used in the paper, which contains some minor changes to make it conserve nitrogen to enable a fair comparison with CRi-Strat. Section S4 documents some differences in reaction rate

Х - 7

coefficients between the CRI-Strat and StratTrop mechanisms. Finally, Sections S5 and S6 provide some supporting analysis of ozone and nitrogen containing species respectively to support the analysis conducted in the main paper.

X - 8

S1. Comparison of mechanism runtime

Both the CRI-Strat and StratTrop mechanisms were run in the UKCA model version $10.9 \text{ at } 1.25^{\circ} \times 1.875^{\circ}$ (N96) resolution with 85 vertical levels up to 85 km, as described in the main paper. Simulations were run on the the joint Met Office - NERC MONSooN2 supercomputer, which is a CRAY XC40 machine using Xeon E5-2695v4 18C 2.1GHz processors. The model simulations were carried out on 12 nodes with 36 cores per node for a total of 432 cores. Over the course of a year, the CRI-Strat run had an average runtime per month of 9056 ± 245 s, compared with 5170 ± 137 s for the StratTrop simulation. Hence the CRI-Strat simulation took 75.2% longer than StratTrop. Memory usage was 252Gb for CRI-Strat compared to 183Gb for StratTrop, an increase of 30%.

S2. Details of the CRI-Strat mechanism

Each intermediate species in CRI is named using a code signifying its general structure, CRI index and functional group. For example, "RN10O2" is an n-aklyl peroxy radical with 10 C - H and C - C bonds (equivalent to the n-propyl peroxy radical, or "NC3H7O2" in the MCM). The radicals generated after the first few steps of oxidation will typically form intermediate lumped carbonyl species (labelled with "CARB") which can undergo further oxidation or photolysis. For example "CARB3" is a dicarbonyl with 3 C-H or C-C bonds (equivalent to glyoxal or "GLYOX" in MCM). An intermediate with a CRI index 3 higher than another intermediate is usually a related molecule with an alkyl chain one C-atom longer (+1 C - C bond, +2 C - H bonds). So "CARB6" is equivalent to methylglyoxal, called "MGLYOX" in MCM. Note that while smaller intermediates (CRI index \leq 10) generally have a single clear analogue species or represent a small group of isomers, larger intermediates often represent many different molecules which may have different chemical

formulae and molecular masses. A full list of potential structure and functional codes are

provided in Table S1.

Full documentation of all species and reactions in the CRI-Strat mechanism are given in Tables S2-S5. In many case species are renamed from how they appear in the original CRI mechanism, for example "CH3" is changed to "Me" and "C2H5" is changed to "Et". This is done both for consistency with how other species are named within UKCA and because there is a hard limit of 10 characters for each species name in ASAD. Those species and reactions with a number 1 in the Strat column are copied from the StratTrop mechanism, to add stratospheric chemistry to the base CRIv2-R5 scheme.

S3. Changes to StratTrop mechanism to conserve Nitrogen

In making the comparison of nitrogen reservoir species (NO_y) between CRI-Strat and StratTrop, it became clear that there was substantially less NO_y in StratTrop than there should be and there was no way of explain where it was going via the diagnostics used to carry out the NO_y budget analysis (see Fig. 13 and Table 9 in main paper). The cause of this problem was found to be these two reactions in StratTrop did not conserve NO_y :

$$DMS + NO_3 \rightarrow SO_2$$
 (1)

$$MONOTERP + NO_3 \rightarrow 0.26 * SEC_ORG$$
 (2)

Both of these reactions are linked to the GLOMAP-mode aerosol scheme, and were copied from the offline oxidants version of the model (Mulcahy et al., 2020) in which concentrations of oxidants are prescribed and therefore unaffected by chemistry. However, when being used in a coupled model setup (such as in UKCA with the StratTrop mechanism), this means that the nitrogen in NO_3 is not being conserved, which would have knock on effects to the NO_v and Ox budgets. These reactions are simulated differently in CRI-Strat

due to its different handling of monoterpene and DMS chemistry, meaning that in CRI-Strat the nitrogen is conserved and tracked through diagnostics. Some minor changes were therefore made to the StratTrop scheme in order to enable a fair and rigorous comparison between the CRI-Strat and StratTrop mechanisms.

It was decided that the simplest solution to this problem would be for both of these reactions to make $HONO_2$ by replacing the two reactions with the following:

$$DMS + NO_3 \rightarrow SO_2 + HONO_2$$
 (3)

$$MONOTERP + NO_3 \rightarrow 0.26 * SEC_ORG + HONO_2$$
 (4)

The HONO₂ produced is most likely to be lost via wet or dry deposition and have little further effect on chemistry (unless it is photolyzed or oxidised). However, it would be tracked by the diagnostics for NO_y deposition, enabling full conservation of reactive nitrogen. For the case of DMS, HONO₂ is a product of NO₃ + DMS oxidation, as it proceeds via hydrogen abstraction (von Glasow & Crutzen, 2004), so this change to the reaction is reasonable. However, monoterpene oxidation via NO₃ typically proceeds with the NO₃ attaching to the double bond to form an organonitrate. In CRI-Strat, when NO₃ reacts with APINENE or BPINENE, the products will propagate, but the nitrogen is always conserved and its deposition is tracked via diagnostics, mostly likely as RONO₂ or as HONO₂ if when formed following further reaction. The monoterpene parameterisations are simplified in StratTrop, hence assuming HONO₂ is formed will enable tracking of NO_y that is in keeping with the rest of the mechanism, enabling a fair comparison with CRI-Strat.

In this section, we document differences between the original StratTrop mechanism (StratTrop_orig) with the modified version that conserves nitrogen (StratTrop_Ncon). In

the main paper, the StratTrop_Ncon mechanism is used for all analysis and is simply referred to as StratTrop. Differences in zonal mean concentrations of $HONO_2$, NO_y , NO_x , NO_z and O_3 are shown in Figure S1. There is a considerable increase $HONO_2$, all of which is lost via reactions 1 and 2 in StratTrop_orig. There is a corresponding increase in NO_y and NO_z in StratTrop_Ncon due to the additional $HONO_2$. However, there is only slight increase in NO_x and O_3 , showing that the additional $HONO_2$ in StratTrop_Ncon is mostly lost to deposition without having a huge influence on the overall composition of the atmosphere.

The differences between StratTrop_orig and StratTrop_Ncon are detailed further in the following tables. The conservation of nitrogen leads to an increase in tropospheric ozone burden of approximately 0.5%, as shown in Table S6. Production and loss terms for O_3 are similar but slightly higher in StratTrop_Ncon. Loss via NO_y deposition is also higher, although this is an artifact because StratTrop_orig does not have a full diagnostic of O_x loss because it is missing the fluxes via reactions 1 and 2. Due to how the stratosphere-troposphere transport (STT) of ozone is calculated from the difference in production and loss of O_x , StratTrop_orig also underestimates how much tropospheric ozone is coming from the stratosphere.

Table S7 shows that OH is slightly higher in StratTrop_Ncon, presumably from HONO₂. However, this has only a small impact on CO burden and lifetime.

Table S8 documents the changes to the NO_y in StratTrop_Ncon compared to Strat-Trop_orig. The HONO₂ burden is 4.5% larger in StratTrop_Ncon compared to Strat-Trop_orig, however this leads to a 7.9% increase in total NO_y due to the large contribution of deposition via HONO₂. In StratTrop_orig, this large fraction of NO_y loss is missing

from the diagnostics, which leads to invalid results where the calculated total emissions are greater than total deposition, implying that net flux of NO_y is into the stratosphere rather than from the stratosphere.

S4. Differences in reaction rates coefficients between mechanisms

The version of the CRI mechanism used for these developments was CRIv2-R5, as documented by M. Jenkin, Watson, Utembe, and Shallcross (2008); Watson, Shallcross, Utembe, and Jenkin (2008); S. R. Utembe, Watson, Shallcross, and Jenkin (2009); the same version used in the WRF-Chem model (Archer-Nicholls et al., 2014) and STOCHEM-CRI model (S. Utembe et al., 2010; Khan et al., 2015). CRIv2-R5 was merged with the stratospheric component of the StratTrop mechanism to make the whole-atmosphere version of the mechanism evaluated in this paper.

Chemical mechanism developers often make use of the most up-to-date kinetic information as possible. However, as improvements are made in experimental techniques and *ab initio* methods, kinetic information (rate constants, product yields etc) change with time. This can be seen through the updates in the evaluations of kinetic data for use in atmospheric chemistry modelling (Sander et al., 2011; Atkinson, 2000). The CRIv2-R5 mechanism was originally optimised against the MCMv3.1 (M. E. Jenkin et al., 2003; Saunders et al., 2003), which drew heavily on kinetic parameters evaluated by the IU-PAC Task Group on Atmospheric Chemical Kinetic Data Evaluation (e.g., Atkinson et al. (1997, 2004)). The StratTrop scheme (Archibald et al., 2020) drew on a mixture of data from the MCMv3.2 website, the IUPAC Task Group web pages and the NASA JPL Evaluation No. 17 (Sander et al., 2011).

In this section, differences in reaction rate coefficients for key bimolecular and termolecular reactions are documented and commented on where relevant. There are some reactions which are out of date in CRI-Strat, and in more up-to-date versions (e.g. version 2.2 M. E. Jenkin et al. (2019)) the updated reaction rate coefficients is either identical or closer

to StratTrop. In other cases, the reactions are simply different, with CRI-Strat drawing from IUPAC evaluations and StratTrop from JPL. In these cases, it is not straightforward to say which mechanism is right. These cumulative differences in reaction rate coefficients are a driving factor in why the CRI_Emiss_ST scenario, in which the CRI-Strat mechanism is run with StratTrop emissions, differs from the StratTrop run.

We first document differences in termolecular reactions in Figure S5. CRI-Strat forms HONO₂ faster under all atmospheric conditions, particularly in colder low pressure conditions present in the upper atmosphere. HONO is also formed more rapidly. In both cases, these increase the rate of transfer from reactive NO_x into reservoir species. HO₂NO₂ is formed faster in the lower atmosphere in CRI-Strat, and PAN is also formed faster under most atmospheric conditions, but both of these important reservoir species are more thermally unstable in CRI-Strat, resulting in more HO₂NO₂ and PAN near emission sources but less in the free troposphere. Formation of MPAN is approximately 50 times faster in StratTrop than in CRI-Strat, one of the largest differences in reaction rates between the mechanisms. This causes substantial changes to chemistry in regions dominated by BVOC emissions, as MACRO2 (or RU10O2 in CRI-Strat) is formed in the degradation of isoprene.

The set of $O(^{1}D)$ reactions with H₂O, N₂ and O₂ are extremely important for tropospheric chemistry as the $O(^{1}D) + H_{2}O$ reaction is both the major source of the OH radical and a key sink of O_x, whereas the $O(^{1}D) + N_{2}$ and $O(^{1}D) + O_{2}$ (collectively $O(^{1}D) + M$) reactions stabilise the exited odd oxygen into its ground electronic state. In CRIv2, the $O(^{1}D) + H_{2}O$ reaction rate coefficient is faster and the $O(^{1}D) + N_{2}$ coefficient slower compared to StratTrop. Collectively, these differences mean any $O(^{1}D)$ atom is between 20 to

25% more likely to react with H_2O in CRI-Strat than in StratTrop, as shown in Figure S2, leading to shorter O_x lifetime and higher HO_x production in CRI-strat, leading to net loss of ozone in Southern hemisphere in CRI-Strat, and loss of ozone across the troposphere in CRI-Emiss_ST and CRI-Emiss_C2C3, compared to StratTrop as shown in Figure S10.

Figure S3 compares reaction rates of all inorganic bimolecular reactions that differ between CRI-Strat and StratTrop, except for the $O(^{1}D)$ reactions documented in Figure S2. All other inorganic bimolecular reactions are identical in both mechanisms. The $O_x + NO_x$ reactions are subtly different between the mechanisms; it is hard to say how significant these changes would be but the faster $O_3 + NO$ reaction in CRI-Strat at boundary layer temperatures would lead to faster titration of ozone in polluted regions. $HO_2 + NO$ is approximately 9% faster in CRI-Strat across all temperatures, leading to faster O_x production The OH + HO₂NO₂ reactions faster in CRI-Strat, lowering the lifetime of this key reservoir species. $OH + H_2$ is slightly slower in CRI-Strat. The $HO_2 + HO_2$ reaction is faster in CRI-Strat, leading to a greater HO_x sink.

Figure S4 compares reaction rates of selected organic bimolecular reactions that differ between CRI-Strat and StratTrop. Due to there being many more organic reaction, this section focuses on those which have a large flux in the troposphere so are important for ozone formation, and for which both mechanisms share the same or similar reactions. The $OH + CH_4$ is slightly faster in CRI-Strat compared to StratTrop, this is not a big difference but it in an important reaction for atmospheric composition and does affect calculations of methane lifetime (Table 6 in main paper). $MeO_2 + NO$ is faster in StratTrop, could aid Ox production away from emission sources in StratTrop and more rapid production of $MeONO_2$. The $OH + MeONO_2$ is 20-100 times faster over typical temperature range in

CRI-Strat compared to StratTrop, and has the opposite temperature dependence, results in much less MeONO₂ in CRI-Strat and loss of an important reservoir species. This is one of the largest differences in reaction rate coefficients between the mechanisms. OH+PAN faster in CRI-Strat, contributing to PAN having a shorter lifetime in CRI-Strat. OH + C_5H_8 and OH + MeCHO are slightly faster in CRI-Strat, whereas OH + EtCHO is faster in StratTrop.

Collectively, all these differences help explain why CRI_Emiss_ST has less NO_x than StratTrop, as the NO_x is more rapidly taken up into reservoir species that can are deposited out. The only exception being the formation of MPAN, which occurs in much greater burdens in StratTrop than CRI-Strat.

S5. Supporting Analysis of ozone

To support Figure 1 in the main paper, Figures S7, S8 and S9 show comparisons between the CRI-Strat and StratTrop model simulations and the TOAR network with a focus on North America, European and East Asian regions respectively, with the majority of the East Asian region site located in Japan. In all three cases, both mechanisms have a low bias compared to observations in winter and a high bias in summer, but CRI-Strat has higher surface ozone throughout the year. Both mechanisms also tend to have a higher bias over low latitude regions and a lower bias over high latitude regions. In North America and Europe, StratTrop is overall biased low over the whole year (-2.2 ppbv and -2.3 ppbv respectively) while CRI-Strat has a high bias (+2.8 ppbv and +2.1 ppbv; Figures S7 and S8). In contrast, StratTrop has a small high bias over the East Asian region (2.8 ppbv) while CRI-Strat has a much higher bias (12.7 ppbv; Figure S9).

Tropospheric ozone in the StratTrop scenario and in comparison to all of the other model scenarios are shown in Figure S10. Tropospheric ozone is higher in the polluted northern hemisphere and lower in the cleaner southern hemisphere in the CRI-Strat simulation compared to StratTrop. CRI_Emiss_ST and CRI_Emiss_C2C3 both have much lower ozone compared to StratTrop across the world. These results indicate that the increased northern hemisphere ozone in CRI-Strat is due to the increase in anthropogenic NMVOC emissions, while the decreased ozone in the southern hemisphere is largely due to different kinetic parameters in CRI-Strat, as explained in Section 1.

Figure S11 shows differences in production and loss of odd oxygen (Ox) in the lowest 1km of the atmosphere between the model scenarios. Production and loss of Ox is typically higher in all scenarios that use the CRI-Strat mechanism compared to Strattrop, but

this is most apparent in the base CRI-Strat scenario. The CRI_Emiss_C2C3 scenario is similar to CRI_Emiss_ST, but has slightly faster production in polluted regions such as the Indo-Gangentic plain.

The difference in flux through the $O(^{1}D) + H_{2}O$ drives much of the difference in loss of Ox between the CRI-Strat mechanism and StratTrop, as can be seen if you compare Figure S12 with Figure 8 (d-f) in the main paper. The reaction rate coefficients for the $O(^{1}D)$ reactions in StratTrop and CRI-Strat are compared in Section 1.

S6. Supporting Analysis of Nitrogen Containing Species

Differences in NO_x , NO_y and NO_z summed species for the lowest 1 km of the atmosphere are shown in Figure S13. There is a clear trend for reduced NO_x and higher NO_z in the CRI-Strat mechanism compared to StratTrop (Fig. S13 (a, c)), with more NO_y everywhere except the highly polluted East Asia and India regions. The CRI-STEmiss and CRI_Emiss_C2C3 simulations have lower NO_x and NO_y almost everywhere compared to StratTrop (Fig. S13 (b, c, e, f)), and the only regions with higher NO_z are downwind of pollution centres such as East Asia, Western Europe and the Indo Gangentic plain (Fig. S13 (h, i)). Overall, CRI-Strat forms HONO2 faster than StratTrop in polluted regions (see Section 1), causing net loss of NO_y in CRI-STEmiss and CRI_Emiss_C2C3. However, with the extra NMVOC emissions in CRI-Strat, more of the NO_y is locked up as RONO₂ which generally have longer lifetimes than HONO₂ and so CRI-Strat has more NO_y everywhere except in very polluted regions where HONO₂ production dominates.

Differences in flux through the $OH + NO_2 + M$ reaction are shown in Figure S14. CRI-Strat has much greater flux through this reaction near the surface, due to the higher concentrations of HO_x in the CRI-Strat simulation. However, away from emission sources, the flux becomes lower in CRI-Strat due to it having NO_x concentrations compared to StratTrop as more of the NO_x is converted into reservoir species. The CRI-Emiss_ST simulation has somewhat greater flux through $OH + NO_2 + M$ than StratTrop through the boundary layer. These differences are mostly driven by the differences in OH and NO2 between the schemes, as although the reaction rate coefficient for the $OH + NO_2 + M$ reaction differs between the mechanisms, it is a relatively small change over the temperature and pressure range of the boundary layer (see Section 1).

X - 20

References

- Archer-Nicholls, S., Lowe, D., Utembe, S., Allan, J., Zaveri, R. A., Fast, J. D., ... McFiggans, G. (2014, nov). Gaseous chemistry and aerosol mechanism developments for version 3.5.1 of the online regional model, WRF-Chem. *Geoscientific Model Development*, 7, 2557–2579. Retrieved from http://www.geosci-model-dev.net/ 7/2557/2014/ doi: 10.5194/gmd-7-2557-2014
- Archibald, A., O'Connor, F., Abraham, N. L., Archer-Nicholls, S., Chipperfield, M., Dalvi, M., ... Zeng, G. (2020). Description and evaluation of the UKCA stratospheretroposphere chemistry scheme (StratTrop vn 1.0) implemented in UKESM1. *Geoscientific Model Development*, 13, 1223–1266. doi: 10.5194/gmd-2019-246
- Atkinson, R. (2000). Atmospheric chemistry of VOCs and NOx. Atmospheric Environment, 34, 2063-2101. Retrieved from http://linkinghub.elsevier.com/ retrieve/pii/S1352231099004604 doi: 10.1016/S1352-2310(99)00460-4
- Atkinson, R., Baulch, D. L., Cox, R. A., Crowley, J. N., Hampson, R. F., Hynes, R. G., ... Rossi, M. J. (2004). Evaluated kinetic and photochemical data for atmospheric chemistry : Volume I – gas phase reactions of Ox, HOx, NOx and SOx species. *Atmospheric Chemistry and Physics*, 4, 1461–1738.
- Atkinson, R., Baulch, D. L., Cox, R. A., Hampson, R. F., Kerr, J. A., Rossi, M. J., & Troe, J. (1997). Evaluated Kinetic and Photochemical Data for Atmospheric Chemistry Supplement VI. J. Phys. chem. Ref. Data, 26(6), 1329–1499.
- Jenkin, M., Watson, L., Utembe, S., & Shallcross, D. (2008, oct). A Common Representative Intermediates (CRI) mechanism for VOC degradation. Part 1: Gas phase mechanism development. Atmospheric Environment, 42, 7185–7195. Retrieved

from http://linkinghub.elsevier.com/retrieve/pii/S1352231008006742 doi: 10.1016/j.atmosenv.2008.07.028

- Jenkin, M. E., Khan, M. A., Shallcross, D. E., Bergström, R., Simpson, D., Murphy, K. L., & Rickard, A. R. (2019). The CRI v2.2 reduced degradation scheme for isoprene. Atmospheric Environment, 212(May), 172–182. doi: 10.1016/j.atmosenv .2019.05.055
- Jenkin, M. E., Saunders, S. M., Wagner, V., & Pilling, M. J. (2003, feb). Protocol for the development of the Master Chemical Mechanism, MCM v3 (Part B): tropospheric degradation of aromatic volatile organic compounds. Atmospheric Chemistry and Physics, 3(1), 181–193. Retrieved from http://www.atmos-chem-phys.net/3/181/ 2003/ doi: 10.5194/acp-3-181-2003
- Khan, M. A. H., Cooke, M. C., Utembe, S. R., Archibald, A. T., Derwent, R. G., Xiao, P., ... Shallcross, D. E. (2015). Global modeling of the nitrate radical (NO3) for present and pre-industrial scenarios. *Atmospheric Research*, 164-165(3), 347–357. doi: 10.1016/j.atmosres.2015.06.006
- Mulcahy, J. P., Johnson, C., Jones, C., Povey, A., Sellar, A., Scott, C. E., ... Yool, A. (2020). Description and evaluation of aerosol in UKESM1 and HadGEM3-GC3.1CMIP6 historical simulations. *Geoscientific Model Development*.
- Sander, S. P., Friedl, R. R., Barker, J. R., Golden, D. M., Kurylo, M. J., Sciences, G. E., ... Orkin, V. L. (2011). Chemical Kinetics and Photochemical Data for Use in Atmospheric Studies Evaluation Number 17 NASA Panel for Data Evaluation :. (17).
- Saunders, S. M., Jenkin, M. E., Derwent, R. G., & Pilling, M. J. (2003, feb). Protocol for
the development of the Master Chemical Mechanism, MCM v3 (Part A): tropospheric degradation of non-aromatic volatile organic compounds. *Atmospheric Chemistry and Physics*, *3*, 161–180. doi: 10.5194/acp-3-161-2003

- Telford, P. J., Abraham, N. L., Archibald, A. T., Braesicke, P., Dalvi, M., Morgenstern, O., ... Pyle, J. A. (2013). Implementation of the Fast-JX Photolysis scheme into the UKCA component of the MetUM chemistry climate model. *Geoscientific Model Development*, 6, 161–177. doi: 10.5194/gmdd-5-3217-2012
- Utembe, S., Cooke, M., Archibald, A. T., Jenkin, M., Derwent, R., & Shallcross, D. (2010, apr). Using a reduced Common Representative Intermediates (CRIv2-R5) mechanism to simulate tropospheric ozone in a 3-D Lagrangian chemistry transport model. *Atmospheric Environment*, 44(13), 1609–1622. Retrieved from http://linkinghub.elsevier.com/retrieve/pii/S1352231010001056 doi: 10 .1016/j.atmosenv.2010.01.044
- Utembe, S. R., Watson, L. A., Shallcross, D. E., & Jenkin, M. E. (2009, apr). A Common Representative Intermediates (CRI) mechanism for VOC degradation. Part 3: Development of a secondary organic aerosol module. *Atmospheric Environment*, 43(12), 1982–1990. Retrieved from http://linkinghub.elsevier.com/retrieve/pii/S1352231009000284 doi: 10.1016/j.atmosenv.2009.01.008
- von Glasow, R., & Crutzen, P. J. (2004, apr). Model study of multiphase DMS oxidation with a focus on halogens. Atmospheric Chemistry and Physics, 4(3), 589-608. Retrieved from http://www.atmos-chem-phys.net/4/589/2004/ doi: 10.5194/acp-4-589-2004

Watson, L., Shallcross, D., Utembe, S., & Jenkin, M. (2008, oct). A Common Repre-

sentative Intermediates (CRI) mechanism for VOC degradation. Part 2: Gas phase mechanism reduction. *Atmospheric Environment*, 42(31), 7196–7204. Retrieved from http://linkinghub.elsevier.com/retrieve/pii/S1352231008006845 doi: 10.1016/j.atmosenv.2008.07.034



Figure S1. Difference in zonal mean $HONO_2$, NO_y , NO_x , NO_z and O_3 between Strat-

 $Trop_NCon \ and \ StratTrop_Orig.$



Figure S2. Comparison of selected inorganic bimolecular reaction rates in CRIv2 to equivalent reactions in StratTrop over an atmospherically relevant temperature range. Blue line is CRI rate, orange line StratTrop, both using scale on left axis, and black dot-dashed line shows ratio of CRI/StratTrop.



Figure S3. Comparison of inorganic bimolecular reaction rates in CRIv2 to equivalent reactions in StratTrop over an atmospherically relevant temperature range. Blue line is CRI rate, orange line StratTrop, both using scale on left axis, and black dot-dashed line shows ratio of CRI/StratTrop.



Figure S4. Comparison of selected organic bimolecular reaction rates in CRIv2 to equivalent reactions in StratTrop over an atmospherically relevant temperature range. Blue line is CRI rate, orange line StratTrop, both using scale on left axis, and black dot-dashed line shows ratio of CRI/StratTrop.



Figure S5. Ratios of selected termolecular reaction rate coefficients in CRIv2 to equivalent reactions in StratTrop over atmospherically relevant temperature and pressure ranges.



Figure S6. Average surface ozone concentrations from rural sites on the TOAR network across the world and biases between StratTrop and CRI-Strat model simulations and observations from TOAR network using data from 2010 to 2014, averaged over whole year (a-c), June to August (d-f) and December to February (g-i). Mean bias between model and observations given in titles of panels b, c, e, f, h, and i.



Figure S7. Average surface ozone concentrations from rural sites on the TOAR network over North America and biases between StratTrop and CRI-Strat model simulations and observations from TOAR network using data from 2010 to 2014, averaged over whole year (a-c), June to August (d-f) and December to February (g-i). Mean bias between model and observations given in titles of panels b, c, e, f, h, and i.



Figure S8. Average surface ozone concentrations from rural sites on the TOAR network over Europe and biases between StratTrop and CRI-Strat model simulations and observations from TOAR network using data from 2010 to 2014, averaged over whole year (a-c), June to August (d-f) and December to February (g-i). Mean bias between model and observations given in titles of panels b, c, e, f, h, and i.



Figure S9. Average surface ozone concentrations from rural sites on the TOAR network over East Asia and biases between StratTrop and CRI-Strat model simulations and observations from TOAR network using data from 2010 to 2014, averaged over whole year (a-c), June to August (d-f) and December to February (g-i). Mean bias between model and observations given in titles of panels b, c, e, f, h, and i.



Figure S10. Tropospheric ozone column (DU) in StratTrop mechanism in DJF (a) and JJA (b). Difference in tropospheric ozone column between CRI-Strat and StratTrop in DJF (c) and JJA (f). Difference in tropospheric ozone column between CRI_Emiss_ST and StratTrop in DJF(d) and JJA (g). Difference in tropospheric ozone column between CRI_Emiss_C2C3 and StratTrop in DJF(e) and JJA (h).



Figure S11. Difference in chemical production of Ox averaged over the lower 1km of the atmosphere between CRI-Strat and StratTrop (a), CRI_Emiss_ST and StratTrop (b), and CRI_Emiss_C2C3 and StratTrop (c). Difference in chemical loss of Ox averaged over lower 1 km of the atmosphere between CRI-Strat and StratTrop (d), CRI_Emiss_ST and StratTrop (e), and CRI_Emiss_C2C3 and StratTrop (f). Difference in deposition of Ox averaged over lower 1 km of the atmosphere between CRI-Strat and StratTrop (g), CRI_Emiss_ST and StratTrop (h), and CRI_Emiss_C2C3 and StratTrop (i). Difference in deposition of Ox averaged over lower 1 km of the atmosphere between CRI-Strat and StratTrop (g), CRI_Emiss_ST and StratTrop (h), and CRI_Emiss_C2C3 and StratTrop (i). Difference in deposition of Ox averaged over lower 1 km of the atmosphere between CRI-Strat and StratTrop (j), CRI_Emiss_ST and StratTrop (k), and CRI_Emiss_C2C3 and StratTrop (l).



Figure S12. Zonal mean flux though the $O(^{1}D) + H_{2}O$ reaction in StratTrop (a), and difference in zonal mean flux between CRI-Strat and StratTrop (b), CRI_Emiss_ST and StratTrop (c), and CRI_Emiss_C2C3 and StratTrop (d).



Figure S13. Mean differences in NO_x (a-c), NO_y (d-f), and NO_z (g-i) over the lower 1 km of the atmosphere between CRI-Strat and StratTrop (a, d, g), CRI_Emiss_ST and StratTrop (b, e and h), and CRI_Emiss_C2C3 and StratTrop (c, f, i).



Figure S14. Zonal mean flux through the OH + NO2 + M reaction in StratTrop (a), and difference in zonal mean flux between CRI-Strat and StratTrop (b), CRI_Emiss_ST and StratTrop (c), and CRI_Emiss_C2C3 and StratTrop (d).

Structural Code Definition	
RN-	Generic N-alkyl radical
NRN-	N-alkyl radical from NO3+alkene oxidation with nitrate group
RA-	Radical from aromatic oxidation
RU-	Unsaturated radical from C5H8 oxidation
NRU-	Unsaturated radical from NO3+C5H8 oxidation with nitrate group
RTN-	Radical from APINENE oxidation
NRTN-	Radical from NO3+APINENE oxidation with nitrate group
RTX-	Radical from BPINENE oxidation
NRTX-	Radical from NO3+BPINENE oxidation with nitrate group
RCOOH-	Carboxylic acid
CARB-	Generic carbonyl
UDCARB-	Second generation dicarbonyl from aromatic oxidation
UCARB-	Unsaturated carbonyl from C5H8 oxidation
NUCARB-	Second generation carbonyl product of Isoprene+NO3 oxidation with nitrate
TNCARB-	Second generation carbonyl product of APINENE oxidation
CCARB-	Second generation carbonyl product of BPINENE oxidation
TXCARB-	Second generation carbonyl product of BPINENE+O3 oxidation
AROH-	Phenol
RAROH-	Phenol radical
ARNOH-	Nitrophenol
Functional Group Code	Definition
-02	Peroxy radical
-OOH	Peroxide
-NO3	Nitrate
-PAN	Peroxyacyl nitrate

 Table S1.
 Structural codes and functional types used to describe CRI intermediates.

Table S2. Species treated by the CRI-Strat chemistry mechanism. Where the name of the species has been changed from that used by the original CRIv2-R5 mechanism in order to follow UKCA standards, the original name is also given. Species with 1 in the Strat column have been added to make the mechanism suitable for running in the stratosphere, species with 1 in the Aero column are only activated if running with GLOMAP-MODE.

Table S3. Photolysis reactions in the CRI-Strat chemical mechanism. Reactions with 1 in the Strat column have been added to make the mechanism suitable for running in the stratosphere, reactions with 1 in the Aero column are only activated if running with GLOMAP-MODE. References for cross section data for reactions can be found in Telford et al. (2013). Reaction rates calculated online from the cross section data are multiplied by the scaling factor.

Table S4. Bimolecular reactions in CRI-Strat chemical mechanism. Reactions with 1 in the Strat column have been added to make the mechanism suitable for running in the stratosphere, reactions with 1 in the Aero column are only activated if running with GLOMAP-MODE. Temperature dependent reaction rate coefficients k_0 , a_0 and b_0 are given for the equation $k(T) = k_0(T/300)^{a_0} \exp(-b_0/T)$, where T is temperature in K.

Table S5. Termolecular reactions in CRI-Strat chemical mechanism. Reactions with 1 in the Strat column have been added to make the mechanism suitable for running in the stratosphere, reactions with 1 in the Aero column are only activated if running with GLOMAP-MODE. Temperature and pressure dependent reaction rate coefficients k_1 , a_1 , b_1 , k_2 , a_2 , b_2 and F are given for the equation $k(T) = \left(\frac{k_0(T)[M]}{1+k_0(T)[M]/k_i(T)}\right)F_c^{(1+(\log(k_0(T)/k_i))^2)^{-1}}$, where T is temperature in K; [M] is the total number density in molecules cm⁻³; $k_0 = k_1(T/300)^{a_1}\exp(-b_1/T)$ is the low pressure limit rate coefficient and $k_i = k_2(T/300)^{a_2}\exp(-b_2/T)$ is the infinite pressure limit rate coefficient. If $k_2 = 0$ then $k(T) = k_0(T)[M]$; if $k_1 = 0$ then $k(T) = k_i(T)$. The broadening term F_c is calculated from F as follows: if 0 > F < 1 then $F_c = F$; if F > 1 then $F_c = \exp(-F/T)$; if F = 0 then the F_c term in the equation is ignored.

Table S6. Overview of tropospheric Ox burden, lifetime, ozone production efficiency (OPE), chemical production, chemical loss, deposition and inferred stratosphere to troposphere transfer. Values in brackets give fraction of total chemical production for the production terms and fraction of total losses $(L_{Ox} + D_{Ox})$ for the loss and deposition terms.

		StratTrop_orig	StratTrop_Ncon
$\overline{O_3 \text{ burden (Tg)}}$		335.2	336.8
O_x lifetime (days)		19.9	19.8
OPE $(mole_{O_3}mole_{NO_x}^{-1})$		27.0	27.2
$\overline{O_x \text{ production}}$	Total	5692	5725
$(Tg O_3 year^{-1})$	$HO_2 + NO$	3832~(67.3%)	3853~(67.3%)
	$CH_3O_2 + NO$	1276~(22.4%)	1285~(22.5%)
	$R'O_2 + NO$	544 (9.6%)	545~(9.5%)
	Other ^a	40.2 (0.7%)	41.3(0.7%)
$\overline{O_x}$ chemical Loss	Total	5099	5128
$(Tg O_3 year^{-1})$	$O(^{1}D) + H_{2}O$	2649~(43.0%)	2660~(42.9%)
	$HO_2 + O_3$	1586 (25.7%)	$1596\ (25.7\%)$
	$OH + O_3$	706 (11.5%)	714~(11.5%)
	$O_3 + Alkene$	96.4~(1.6%)	96.5~(1.6%)
	Other ^b	60.8(1.0%)	61.5(1.0%)
$\overline{O_x}$ Deposition	Total	1061	1081
	O_3 dry dep	892~(14.5%)	896~(14.4%)
	NO_y dep	170 (2.8%)	185~(3.0%)
Inferred STT (Tg O_3 year -1)		468	483

^a O_x production channels are the sum of inorganic acid oxidation, RONO₂ oxidation and

RONO₂ photolysis.

^b "Other" O_x loss channels are the sum of $O(^{3}P) + O_3$, $O^{3}P + NO_2$, $N_2O_5 + H_2O$ and NO_3

chemical losses.

Table S7.	Overview	of air	mass	weighted	OH	concentration,	CO	burden	and	CO	lifetime.
-----------	----------	--------	------	----------	----	----------------	----	--------	-----	----	-----------

	StratTrop_orig	StratTrop_Ncon
$\overline{[OH]}$ (10 ⁶ molecules cm ⁻³)	1.33	1.34
CO burden (Tg)	301.2	300.2
CO lifetime (days)	38.8	38.6

Table S8. Overview of tropospheric oxidised nitrogen burdens (fraction of total NO_y in brackets), tropospheric oxidised nitrogen emission and deposition fluxes, stratosphere-troposphere transfer (STT) and NO_y lifetime in the troposphere (fraction of total NO_y deposition in brackets).

	StratTrop_orig	StratTrop_Ncon
NO _y Burden (Tg N)	0.993	1.018
NO_x Burden (Tg N)	0.151~(15.2%)	0.152~(14.9%)
NO_z Burden (Tg N)	0.842~(84.8%)	0.866~(85.1%)
$HONO_2$ Burden (Tg N)	0.491 (49.4%)	0.513(50.4%)
Other inorganic NO_z (Tg N)	0.017~(1.7%)	0.018~(1.7%)
PANs (Tg N)	0.295~(29.7%)	0.296~(29.1%)
$RONO_2 (Tg N)$	0.039~(3.9%)	0.039~(3.9%)
Total NO_x Emissions (Tg N year ⁻¹)	61.5	61.5
Total NO_y Deposition (Tg N year ⁻¹)	58.3	62.9
Inferred STT (Tg N year ^{-1})	-3.19	1.40
NO_x Dry deposition (Tg N year ⁻¹)	7.7 (13.2%)	7.70 (12.2%)
HONO ₂ Wet deposition (Tg N year ⁻¹)	27.8~(47.7%)	30.1~(47.8%)
$HONO_2$ Dry deposition (Tg N year ⁻¹)	19.3~(33.1%)	21.6~(34.3%)
Other inorganic NO_z deposition (Tg N year ⁻¹)	0.97~(1.7%)	0.97~(1.6%)
PANs dry deposition (Tg N year ^{-1})	1.28~(2.2%)	1.28~(2.0%)
$RONO_2$ deposition (Tg N year ⁻¹)	1.28~(2.2%)	1.30~(2.1%)