## The phase of water ice which forms in cold clouds in the mesospheres of Mars, Venus and Earth

Thomas Mangan<sup>1</sup>, John Plane<sup>1</sup>, and Benjamin Murray<sup>2</sup>

<sup>1</sup>School of Chemistry, University of Leeds <sup>2</sup>School of Earth and Environment, University of Leeds

November 24, 2022

#### Abstract

Water ice clouds form in the mesospheres of terrestrial planets in the solar system (and most likely elsewhere) by vapor deposition at low pressures and temperatures. Under these conditions a range of crystalline and amorphous phases of ice might form. The phase is important because it influences nucleation kinetics, density, vapor pressure over the solid, growth rates and particle shape. In the past the temperature range over which these different phases exist has been defined on the basis of depositing ice at low temperature and warming it while observing phase changes. However, the direct deposition of ice at a range of temperatures relevant for the terrestrial planets has not been systematically investigated. Here we present X-ray Diffraction (XRD) measurements of water ice deposited at temperature intervals between 88 and 145 K in a vacuum chamber. XRD patterns showed that low density amorphous ice was formed at [?] 120 K, stacking disordered ice I formed from 121 – 135 K and hexagonal ice I formed at 140 and 145 K. Direct deposition results in the stable hexagonal phase at much lower temperatures than when warming stacking disordered ice. All three phases of water ice observed here are possible in clouds in the mesospheres of Earth and Mars, while on Venus only amorphous ice is likely to form.

# The phase of water ice which forms in cold clouds in the mesospheres of Mars, Venus and Earth

T. P. Mangan<sup>1</sup>, J. M. C. Plane<sup>1</sup>, B. J. Murray<sup>2</sup>

<sup>1</sup>School of Chemistry, University of Leeds, LS2 9JT.

<sup>2</sup>School of Earth and Environment, University of Leeds, LS2 9JT.

Email: B.J.Murray@leeds.ac.uk.

## **Key Points**

- Low density amorphous ice was formed at 88-120 K, stacking disordered ice I formed from 121-135 K and hexagonal ice I formed at 140-145 K.
- Direct deposition results in the stable hexagonal phase at much lower temperatures than when warming stacking disordered ice.
- All three phases of ice are possible in clouds in the mesospheres of Earth and Mars, while on Venus only amorphous ice is likely to form.

## Abstract

Water ice clouds form in the mesospheres of terrestrial planets in the solar system (and most likely elsewhere) by vapor deposition at low pressures and temperatures. Under these conditions a range of crystalline and amorphous phases of ice might form. The phase is important because it influences nucleation kinetics, density, vapor pressure over the solid, growth rates and particle shape. In the past the temperature range over which these different phases exist has been defined on the basis of depositing ice at low temperature and warming it while observing phase changes. However, the direct deposition of ice at a range of temperatures relevant for the terrestrial planets has not been systematically investigated. Here we present X-ray Diffraction (XRD) measurements of water ice deposited at temperature intervals between 88 and 145 K in a vacuum chamber. XRD patterns showed that low density amorphous ice was formed at ≤ 120 K, stacking disordered ice I formed from 121 -135 K and hexagonal ice I formed at 140 and 145 K. Direct deposition results in the stable hexagonal phase at much lower temperatures than when warming stacking disordered ice. All three phases of water ice observed here are possible in clouds in the mesospheres of Earth and Mars, while on Venus only amorphous ice is likely to form.

## Plain Language Summary

Ice clouds made up of water ice crystals can form in the thin upper atmospheres (mesospheres) of terrestrial planets, where the temperatures are very cold (< 150 K or <  $-190^{\circ}$ F or <  $-123^{\circ}$ C). Under these conditions a range of crystalline (hexagonal, cubic, or a mixture of both known as stacking disordered ice) and amorphous (lacking a defined crystal structure) phases of ice might form. The phase is important because it influences a range of cloud properties. Experimentally, rather than warming ice formed at low temperature (as has been done in the past) we deposit the ice directly between 88 – 145 K. Measurements showed that the amorphous, stacking disordered

and hexagonal phases of water ice (ice I) can form depending on the specific temperature and direct deposition results in the hexagonal phase at much lower temperatures than when warming ice. All three phases of water ice observed here are possible in clouds in the mesospheres of Earth and Mars, while on Venus only amorphous ice is likely to form.

## Index Terms

0320 Cloud physics and chemistry, 0343 Planetary atmospheres, 5422 Ices, 3311 Clouds and aerosols, 0305 Aerosols and particles

## Keywords

Mesospheric clouds, Martian Clouds, Venusian Clouds, Amorphous ice, Stacking disordered ice, Hexagonal ice

## 1. Introduction

Water ice clouds can form in the coldest regions of rarefied planetary upper atmospheres, where particles grow through deposition from the vapor phase. Despite the very low partial pressures of water in the upper atmospheres of Venus, Earth and Mars, the temperatures can be low enough (~<150 K, depending on the planet) to lead to large supersaturations and the formation of ice particles. At these temperatures multiple metastable forms of ice can exist, but the phases that can form remain poorly defined. In this paper we focus on understanding the phase of ice that forms in this class of very cold clouds which exist in the upper atmospheres of three of the terrestrial planets in the solar system.

In Earth's upper mesosphere (80-90 km), nanoparticles composed primarily of water ice form between 100 and 150 K, producing clouds known as Polar Mesospheric Clouds (PMCs) or noctilucent clouds (NLCs) [Thomas, 1991; Hervig et al., 2001; Rapp and Thomas, 2006]. In the Martian atmosphere, water ice clouds have also been observed planet wide at altitudes (30 - 90 km) where temperatures can drop to < 120 K [Forget et al., 2009; Vincendon et al., 2011; Fedorova et al., 2020]. In addition, water ice particles may serve as seeds for CO<sub>2</sub> ice particles in the Martian mesosphere [Plane et al., 2018]. On Venus, the possibility of water ice clouds has been hypothesised by Turco et al. [1983]. At the mesopause (~120 km) temperatures can drop below 100 K, leading to supersaturated conditions conducive for ice particle formation [Mahieux et al., 2015; Chamberlain et al., 2020]. Icy grains containing H<sub>2</sub>O are also present in the Inter-Stellar Medium (ISM) and past the frost line within the solar system, where it is cold enough ( $\leq$  150 K) for  $\mu$ m-sized ice particles to form [Gillett and Forrest, 1973; D'Angelo and Podolak, 2015]. Examples of temperature profiles for Earth, Mars and Venus, with calculated saturation ratios with respect to hexagonal ice I are shown in Figure 1. Super-saturated regions with respect to water ice are apparent in the mesospheres of all three planets, with temperatures where  $S \ge 1$  ranging from ~140 K down to < 80 K.

In the case of PMCs on Earth, the prevailing view is that ice particles are predominantly crystalline ice I based on satellite observations, but the specific crystalline phase is

undetermined [*Hervig and Gordley*, 2010]. However, the transitory presence of Low-Density Amorphous (LDA) ice should not be ruled out. LDA ice may nucleate homogeneously depending on the temperature and supersaturation, but may rapidly transition to a crystalline phase [*Murray and Jensen*, 2010]. Also, it has been proposed that the initial phase that would nucleate in PMCs is always LDA ice irrespective of the nucleation mechanism [*Nachbar et al.*, 2018b]. Crystallization timescales at different temperatures for this LDA ice then become important for the evolution of cloud particles under these low temperature conditions [*Jenniskens and Blake*, 1996]. For Martian and Venusian mesospheric clouds the temperature minima are generally more extreme than the mesopause on Earth and the phase of ice which forms has not been considered in the literature.

In total, water ice is known to exist in 19 distinct polymorphs, as well as low and high density amorphous forms [Loerting et al., 2011; Salzmann et al., 2011; Falenty et al., 2014; del Rosso et al., 2016; Millot et al., 2018; Thoeny et al., 2019]. Under the low temperature and pressure conditions in planetary upper atmospheres, Ice I is the only crystalline phase expected to form. Historically, ice I was thought to exist as hexagonal ice (ice I<sub>h</sub>) or cubic ice (ice I<sub>c</sub>) polymorphs [Hobbs, 1974], however it has become apparent that most samples of ice identified as being cubic were actually made of ice consisting of interlaced layers of cubic and hexagonal sequences, known as stacking disordered ice (ice Isd) which has trigonal symmetry [Hobbs, 1974; Hansen et al., 2008; Kuhs et al., 2012; Malkin et al., 2012; Malkin et al., 2015; Salzmann and Murray, 2020]. It was only very recently that true ice I<sub>c</sub> has been made [del Rosso et al., 2020; Komatsu et al., 2020; Salzmann and Murray, 2020], but the synthesis routes were not relevant for atmospheric ice formation. Both ice Ih and Ic structures are made up of honeycomb layers of hydrogen bonded water molecules, structural differences arise due to the way these layers are arranged. For ice I<sub>h</sub>, each layer is a mirror image of the last, for ice I<sub>c</sub> each layer is shifted, while a mixture of both stacking sequences leads to ice Isd.

The specific phase formed in these clouds is important as it influences the properties of the particles that form, including the nucleation kinetics, density, vapor pressure over the solid, growth rates and particle shape. These factors then impact the growth and sedimentation of the ice particles, a process which redistributes water vapour, resulting in a layer with depleted water mixing ratio and below that a layer of enhanced water vapour where the particles sublime [*Hervig et al.*, 2015]. In Earth's mesosphere this is thought to alter odd hydrogen and oxygen chemistry [*Murray and Plane*, 2003; 2005a], and particle phase may also influence the heterogeneous chemistry on mesospheric ice particles [*Plane et al.*, 2004; *Murray and Plane*, 2005b; *Mangan et al.*, 2017a]. Similar processes and interactions may occur in the atmospheres of Venus and Mars. Both the boundary between amorphous and crystalline ice formation, and the specific crystal structure of ice formed, is of interest. All of this suggests a complex range of possibilities for the form of Ice I, and how much LDA ice is present in planetary upper atmospheres, both of which may depend on factors including deposition temperature, particle warming, saturation ratio and pressure.

Previous experimental studies of the crystal structure of ice I below 200 K have typically deposited the ice at a single base temperature and observed structural

changes via warming [*Shilling et al.*, 2006; *Kuhs et al.*, 2012; *Murray et al.*, 2015a]. Indeed, the amorphous to crystalline boundary has primarily been investigated this way, in terms of the transition, rather than determining the temperature at which the ice is no longer amorphous at deposition [*Jenniskens and Blake*, 1994; 1996; *Smith et al.*, 1996]. In contrast, in the present study we use X-ray Diffraction (XRD) to investigate the effect of changing initial deposition temperature on the crystal structure of ice I between 88 -145 K by vapor depositing ice at regular temperature intervals.



Figure 1. Example temperature profiles and saturation ratios for H<sub>2</sub>O ice in the upper atmospheres of Earth, Mars and Venus. Saturation is defined using H<sub>2</sub>O vapor mixing ratios of 5 ppm (Earth), 0.7 ppm (Mars) and 2 ppm (Venus) [*Forget et al.*, 1999; *Hervig* 

*et al.*, 2009; *Chamberlain et al.*, 2020]. Saturation ratios were calculated using the *Nachbar et al.* [2019] amorphous correction to the vapor pressure over ice  $l_h$  [*Murphy and Koop*, 2005]. The atmospheric profile for Earth was taken from the Solar Occultation for Ice Experiment (SOFIE) database (http://sofie.gats-inc.com), event profile 56463 [*Marshall et al.*, 2011]; for Mars, the Mars Pathfinder entry profile [*Magalhães et al.*, 1999]; and for Venus, an averaged profile at 70 - 80° Lat [*Mahieux et al.*, 2012].

## 2. Methods

To investigate the crystal structure of  $H_2O$  under different deposition conditions, we used a temperature-controlled stage within a vacuum chamber, probed using the powder X-ray Diffraction (XRD) technique which we have employed previously to study deposition of  $H_2O$  ice [*Shilling et al.*, 2006; *Murray et al.*, 2015a]. Shown in Figure 2, the experimental system described briefly below is in the same configuration as described in *Mangan et al.* [2017b] for CO<sub>2</sub> deposition, with some modifications to allow for  $H_2O$  vapor deposition.

The X-Ray diffractometer (Bruker D8 Advance) was configured in a standard reflectance geometry and equipped with a Cu K- $\alpha$  X-ray source ( $\lambda$ =1.540598 Å). Diffracted X-rays from a sample mounted on an Anton Paar TTK 450 temperature-controlled stage were detected by a VÅNTEC detector (shown in Fig. 1). The TTK 450 chamber was pumped by a Pfeiffer Duo 2.5A vacuum pump allowing a pressure range of 1 × 10<sup>-3</sup> – 1 × 10<sup>3</sup> mbar. The stage was cooled with liquid nitrogen to temperatures down to 88 ± 0.2 K, measured using a Pt-100 probe inserted into the sample support. The temperature was controlled using a Eurotherm PID controller and resistive heaters. The aluminium sample support was covered with a borosilicate glass slide (160 µm thickness) adhered using a thin layer of vacuum grease.

For each experiment the environmental chamber was first pumped to a pressure of <  $1 \times 10^{-2}$  mbar. The sample support was then cooled from room temperature at a rate of 20 K min<sup>-1</sup> to the specific deposition temperature (110 – 145 K at deposition temperature increments of 5 K, one experiment at 88 K, as well as 1 K increments around 120 K), at which temperature a diffraction pattern was recorded ( $2\theta = 20 - 50^{\circ}$ ) to ensure no ice frosting had occurred on the glass slide. Water vapor was produced by flowing dry N<sub>2</sub> (10 sccm) through two bubblers containing deionised water cooled to 278 K. This wet flow was combined downstream of the bubblers with a dry N<sub>2</sub> flow of 290 sccm, before entering the front of the chamber via a needle valve, with a stable total chamber pressure in each experiment of 10 mbar (20 mbar at 88 K) maintained by throttling the vacuum pump. Low resolution, 30 second scans at  $2\theta = 22 - 30^{\circ}$  were taken during 20 minutes of ice growth to monitor the signal (examples at 121 K are shown in Figure 3). The humid N<sub>2</sub> flow was then stopped and a high resolution XRD pattern (5-minute scan) with increments of  $2\theta = 0.0426$  for  $2\theta = 20 - 50^{\circ}$  measured.  $2\theta$ values were corrected against a silicon standard, accounting for peak shifting due to sample height.



Figure 2. Diagrams of the experimental set-up, including; a front on view of the vacuum chamber with temperature-controlled stage (TTK 450), detector and X-ray source (left), top down view of the experiment including the  $H_2O$  vapor flow system (right).



Figure 3. XRD patterns of ice growth at 121 K monitored over 20 minutes at  $2\theta = 22 - 30^{\circ}$ . The patterns are offset for clarity.

#### 3. Results

Figure 4 shows the background corrected diffraction patterns of water ice grown for 20 minutes and deposited at intervals from 88 - 145 K at 10 mbar (88 K measurement taken at 20 mbar) are compared with calculated patterns for ice I<sub>sd</sub>, I<sub>h</sub> and I<sub>c</sub>. The diffraction patterns at 110, 115 and 120 K show an absence of defined Bragg peaks with two broad amorphous features. The first, centred around  $24^{\circ}$  and a broader, weaker feature at >35° centred around  $40 - 45^{\circ}$  are consistent with LDA ice [*Dowell and Rinfret*, 1960; *Shilling et al.*, 2006]. The diffraction pattern taken at 88 K  $2\theta = 22 - 30^{\circ}$  also shows the broad feature centred at 24°, suggesting LDA ice is deposited from 88 - 120 K.

In order to probe the specific conditions at which the deposited ice is no longer amorphous, experiments were undertaken at 1 K intervals, over a smaller temperature range (119 – 122 K), shown in Figure 5. The presence of crystalline ice, with defined sharp Bragg peaks is seen consistently in diffraction patterns of ice deposited at 121 K and above. The experimental peak positions at 24°, 40° and 48° are consistent with the simulated patterns of I<sub>c</sub> and ice I<sub>h</sub>. The shoulder peak at 23° is consistent with ice I<sub>h</sub>, but the remaining prominent ice I<sub>h</sub> peaks are missing (26°, 33.5° and 44°). Combining these inconsistencies with the feature from 23 - 26° shows that neither pure ice I<sub>h</sub> or I<sub>c</sub> formed at 121-135 K and instead the crystalline ice formed is consistent with ice I<sub>sd</sub>. Ice I<sub>sd</sub> can have a range of cubicities and memory in the stacking sequences, but the shape of the pattern (particularly around 23 - 26°) suggests that this ice is roughly 50% cubic and 50% hexagonal sequences which are randomly arranged [*Malkin et al.*, 2015; *Salzmann and Murray*, 2020].

The experiments at 140 and 145 K show marked differences from the ice  $I_{sd}$  deposited at 120 -135 K. Firstly, the 40° and 23° peaks are no longer present and neither is the broad feature from 23 - 26° suggesting the ice formed is no longer ice  $I_{sd}$ . At 140 K, only two main peaks are present; a high intensity 24° peak (consistent with ice  $I_h$  and  $I_c$ ) and a weaker intensity 43° peak (only seen in the simulated pattern for ice  $I_h$ ). The absence of stacking disordered features and a peak in the diffraction pattern at 2 $\theta$  that is only found in ice  $I_h$  leads to the conclusion that at 140 K the ice formed is ice  $I_h$ , growing with strong preferred crystallographic orientation (we explain this below) rather than ice  $I_c$  or  $I_{sd}$ . Further evidence for pure ice  $I_h$  forming can be found in the ice deposited at 145 K, where two additional small Bragg peaks appear at 26° and 33.5° that are only found in ice  $I_h$ . These findings on the structure of ice I with deposition temperature are discussed in the context of the literature in the following section.

The relative intensities of the patterns of crystalline ice are a poor match to the ideal simulated patterns because the samples exhibit preferred crystallographic orientation. The simulated patterns are derived assuming a large population of randomly orientated crystallites are present. However, in our XRD experiments this is not the case and the vapor deposited ice shows evidence of varying degrees of preferred orientation. This occurs because of ordering of the crystallites via oriented growth on the sample surface. In the case of the diffraction patterns shown here, the crystallites are partially ordered, leading to varying intensities of the Bragg peaks for ice  $I_{sd}$  and in the case of ice  $I_h$ , missing peaks relative to the simulated pattern. While the strong preferred orientation precludes further detailed analysis of these diffraction patterns (such as stacking probability in ice  $I_{sd}$ ), we can identify the phases.

sample temperature / K	Sliq <sup>a</sup>	Sice In <sup>a</sup>	SLDAice b
88	3.09 × 10 <sup>16</sup>	1.86 × 10 <sup>17</sup>	9.57 × 10 <sup>15</sup>
110	2.83 × 10 <sup>10</sup>	1.11 × 10 <sup>11</sup>	1.08 × 10 <sup>10</sup>
115	2.80 × 10 <sup>9</sup>	1.03 × 10 <sup>10</sup>	1.11 × 10 <sup>9</sup>
120	3.35 × 10 <sup>8</sup>	1.15 × 10 <sup>9</sup>	1.38 × 10 <sup>8</sup>
125	4.74 × 10 <sup>7</sup>	1.54 × 10 <sup>8</sup>	2.01 × 10 <sup>7</sup>
130	7.77 × 10 <sup>6</sup>	2.39 × 10 <sup>7</sup>	3.41 × 10 <sup>6</sup>
135	1.45 × 10 <sup>6</sup>	4.24 × 10 <sup>6</sup>	6.55 × 10⁵
140	3.05 × 10⁵	8.51 × 10⁵	1.42 × 10⁵
145	7.14 × 10 <sup>4</sup>	1.90 × 10⁵	3.39 × 10 <sup>4</sup>

Table 1. Experimental saturation ratios for liquid ( $S_{liq}$ ), ice I<sub>h</sub> ( $S_{ice Ih}$ ) and the LDA ice correction to  $S_{ice Ih}$  ( $S_{LDA ice}$ ).

<sup>a</sup> [Murphy and Koop, 2005], <sup>b</sup> [Nachbar et al., 2019].



Figure 4. XRD patterns of  $H_2O$  ice, vapor deposited at temperatures from 88 -145 K. Included at the top of the figure are predicted patterns of ice  $I_h$ ,  $I_c$  and  $I_{sd}$  (assuming randomly arranged and equal probability of ice  $I_h$  and  $I_c$  sequences). A summary of the experimental conditions is shown in Table 1. The patterns are offset for clarity.



Figure 5. XRD patterns of vapor deposited ice close to the amorphous- crystalline boundary of ice I at 115 – 122 K. The patterns are offset for clarity.

### 4. Discussion

#### 4.1 The amorphous – stacking disordered boundary at 120 K.

In this study we observe LDA ice at deposition temperatures of  $\leq 120$  K while ice I<sub>sd</sub> is formed at 121 – 135 K. The formation of LDA ice at the coldest deposition conditions investigated here is consistent with other studies looking at very low temperature ice [*Dowell and Rinfret*, 1960; *Hobbs*, 1974; *Jenniskens and Blake*, 1994]. Also, the observation in this study that ice I<sub>sd</sub> is the coldest observed crystalline form of ice I formed by direct vapor deposition is consistent with other literature, whether the ice was produced by direct vapor deposition or via warmed LDA ice. Using a similar experimental setup, *Murray et al.* [2015a] warmed amorphous ice at 1 K min<sup>-1</sup> from 110 K and observed crystallisation to ice I<sub>sd</sub> at 120 K. However, this crystallisation behaviour in warmed ice is often seen at temperatures much higher than 120 K. For example, in *Jenniskens and Blake* [1994], the transition from amorphous to crystalline ice was gradual, where warming at 1 - 2 K min<sup>-1</sup> caused changes in diffraction patterns at 131 K. In another example, *Hallbrucker et al.* [1989] observed a structural relaxation at 136 ± 1 K prior to the onset of crystallisation in amorphous ice generated via hyperquenching water. In other studies, complete crystallization upon warming was not observed until 150 - 160 K [Johari et al., 1991; Jenniskens and Blake, 1994; Smith et al., 1996; Mitlin and Leung, 2004; Shilling et al., 2006].

The gradual transition to crystalline ice observed in the literature is in stark contrast to what is observed in Figure 5, where the diffraction patterns of ice deposition at  $\geq$ 121 K show a completely crystalline structure with defined Bragg peaks. Outside of the features associated with ice I<sub>sd</sub>, the diffraction patterns show no evidence of the broad features associated with amorphous ice. *Hobbs* [1974] provides a summary of the early experimental research into ice I at low temperatures, covering 16 studies highlighting that the amorphous-crystalline boundary was observed to start and complete over a wide range of temperatures (approximately 110 – 160 K). However, at least some of these diffraction studies may have misinterpreted the broad feature between 23 - 26° in ice I<sub>sd</sub> as evidence for the persistence of amorphous ice to relatively high temperatures [*Dowell and Rinfret*, 1960]. Hence, we suggest that the phase transition temperatures reported in some of these earlier studies may not always be reliable.

In a study of the vapor pressure of low temperature ice I, Nachbar et al. [2018a] concluded that because the same nano-crystalline ice polymorph was observed both by warming of amorphous ice and direct vapor deposition, amorphous ice is the initial phase formed by vapor deposition below 160 K before crystallization occurs above 130 K. Crystallization timescales could play a part in the temperature dependent behaviour of deposited ice and are an important variable atmospherically. The higher in temperature the amorphous ice is, the quicker crystallization occurs [Dowell and Rinfret, 1960; Jenniskens and Blake, 1996]. Isothermal experiments by Jenniskens and Blake [1996] showed amorphous ice warmed to and then held at 125 K took ~5000 seconds to show evidence of crystallinity in the diffraction patterns. In contrast, in our study there was a maximum crystallization timescale of about 1800 s and we observe crystalline ice deposited at this temperature. In the present study evidence of defined Bragg peaks associated with crystalline ice are present in the first scans taken only 300 s from the onset of deposition at 121 K (shown in Figure 3), while crystallization timescales are several days at this temperature according to Jenniskens and Blake [1996]. If amorphous ice forms first then it must have crystallized on timescales shorter than were observable in this experiment, in contrast to slower crystallization in annealed ice from other studies.

#### 4.2 The deposition of ice *I<sub>h</sub>* at relatively low temperatures

A striking observation in our study was the very low temperature (140 - 145 K) at which deposition produced ice l<sub>h</sub>, as shown in Figure 4. This result contrasts with the results of a number of previous studies: amorphous ice warmed from the deposition temperature of 110 K by *Murray et al.* [2015a] was found to be ice I<sub>sd</sub> from 120 K all the way up to the highest temperature measured (160 K). In neutron diffraction experiments of annealed ice deposited initially under liq N<sub>2</sub> temperatures, *Kuhs et al.* [2012] showed that the ice was stacking disordered up to the 175 – 190 K range. A fraction of cubicity was still apparent during isothermal experiments over the course of 12 hours, especially at the lowest (175 K) temperature studied. While LDA ice warmed from 90 K by *Shilling et al.* [2006] was still in the form ice I<sub>sd</sub> at 180 K and persisted for

at least 60 minutes. The literature review in *Hobbs* [1974] shows that ice I<sub>h</sub> is typically observed from ~170 – 200 K. Experiments by *Dowell and Rinfret* [1960] suggested conversion to ice I<sub>h</sub> occurs at  $\geq$  140 K. However, there is a caveat in that these early studies were carried out before stacking disorder in ice I was recognised. It appears from the discussion in *Dowell and Rinfret* [1960] that the peak at ~23° was interpreted as evidence for ice I<sub>h</sub>, but we now know that this peak is consistent with ice I<sub>sd</sub> [*Malkin et al.*, 2015]. Overall, these literature studies suggest that ice I<sub>sd</sub> persists to temperatures as high as 190 K, in strong contrast to our study where we deposited directly. To verify that our experimental system was not promoting ice I<sub>h</sub>, we deposited LDA ice at 110 K and warmed it to 160 K (see Figure 6). The ice at 160 K was ice I<sub>sd</sub>, similar to the findings of *Murray et al.* [2015a] and the other studies listed above. Hence there does appear to be a strong difference between the ice phase produced through direct deposition and that of ice deposited at low temperatures and warmed.



Figure 6. XRD patterns of  $H_2O$  ice, vapor deposited at 110 K and then warmed to 160 K. Included at the top of the figure is a predicted pattern for stacking disordered ice, ice  $I_{sd}$  (assuming randomly arranged and equal probability of ice  $I_h$  and  $I_c$  sequences). The patterns are offset for clarity.

The effect of saturation ratio (and therefore growth rate) on the formation of ice is an experimentally and atmospherically relevant variable that could influence which polymorph of ice I is produced. In this study the flow of water vapor and the chamber pressure was fixed, leading to higher saturation ratios with respect to the solid phase at the lower temperatures (Table 1: at 145 K the saturation ratio ( $S_{LDAice}$ ) is ~3 x 10<sup>4</sup> while at 110 K it is ~1 x 10<sup>10</sup>). *Hudait and Molinero* [2016] grew ice I between 200 to 260 K in a modelling study and concluded that the crystal structure was linked to growth rate and therefore also linked to saturation ratio. Ices formed at higher saturation ratios (> 1.6 x 10<sup>4</sup>) were found to form ice I<sub>sd</sub>, with ice I<sub>h</sub> forming at lower saturation ratios. This trend in saturation although not directly comparable is consistent with the data observed here, where the highest temperatures (140 - 145 K) and lowest saturation ratios (~(0.3 - 1) x 10<sup>5</sup>) leads to ice I<sub>h</sub> formation. This may also be the

situation in planetary atmospheres where supersaturation is driven primarily by temperature with relatively constant water mixing ratio.

We now turn to why ice Ih is favoured when deposited at and above 140 K, whereas ice Isd is favoured at these temperatures when ice is deposited at lower temperatures and warmed. This issue is related to the fact that ice Isd is favoured when ice is grown from liquid water. Hudait and Molinero [2016] concluded that there is a significant surface energy cost of ice Isd over ice Ih. In fact, the energy cost against cubic ice sequences at the ice-vapour interface is about 25 times greater than that in the iceliquid interface [Hudait and Molinero, 2016; Hudait et al., 2016]. This means that when ice grows from the vapour, there is a strong thermodynamic driver for any cubic sequences to relax to the hexagonal configuration. This is only overcome at relatively high growth rates (at high supersaturation) where the cubic sequences are buried in the ice structure before they get a chance to relax to the more stable hexagonal configuration. Similarly, when ice grows from an aqueous environment, the growth is rapid relative to reorganisation, locking the ice in the kinetic product, i.e. ice Isd [Malkin et al., 2015]. This also offers an insight into why ice Isd is produced at temperatures ≤ 135 K and persists when warmed to higher temperatures. At low temperatures ice Isd is favoured in part because the rate of reorganisation at the surface slows with decreasing temperature, but also because the supersaturation and growth rate increases for a given mixing ratio of water vapour [Hudait and Molinero, 2016]. The metastable cubic sequences are then locked within the 'bulk' ice, so when the ice is warmed, the stacking disorder persists until the vapour transport route becomes relatively rapid (on an hour-minute time scale) at temperatures above ~190 K [Murphy, 2003]. It has been suggested in the past that ice Isd crystals might grow in cirrus clouds in the Earth's Tropical Tropopause Layer (TTL) (180 - 200 K) [Murray et al., 2005; Kuhs et al., 2012], but our results suggest that even if aqueous droplets freeze to ice Isd, subsequent growth will result in ice Ih. This finding is consistent with the molecular simulations by Hudait and Molinero [2016], although some stacking faults in predominantly hexagonal structures might be possible, and would be consistent with observed trigonal symmetry in some ice crystals [Kuhs et al., 2012; Murray et al., 2015b].

## 5. Implications for clouds in planetary mesospheres

The experimental results presented in this study suggest that deposition of LDA ice, ice  $I_{sd}$  and ice  $I_h$  may be possible across just a 20 K temperature range, with implications for the crystal phase present in these clouds. It should be noted, as discussed in the previous section, that a range of temperatures have been reported for each ice I phase in the literature, and so a range of conditions for each may be possible atmospherically.

A comparison of the super-saturated regions (regions kinetically favourable for stable ice crystals with  $S \ge 1$ ) are shown in Figure 1 for the upper atmospheres of Earth, Mars and Venus at fixed water vapor mixing ratios (see the figure caption for values). It is worth noting that super-saturation is exponentially dependent on temperature and only linearly dependent on the water vapor mixing ratio, making temperature the key driver.

The implications of the experimental results presented here are discussed for water ice clouds on each planet.

#### 5.1 Earth

The temperature profile shown for Earth is a satellite measurement from June 30<sup>th</sup>, 2012 at 67°N, which represents a typical cold profile seen during the northern hemisphere PMC season occurring between late May and late August. In the mesosphere of Earth, the temperature range where  $S \ge 1$  for the profile shown is approximately 125 – 140 K at 84 – 88 km altitude, around the mesopause altitude. Generally, PMC ice particle temperatures exist in the range of 120 - 150 K [*Hervig and Gordley*, 2010]. PMCs which scatter enough light to be optically visible tend to occur several km below the cold point of the mesopause (when they can appear as NLCs to a ground-based observer), but ice particles are thought to also exist close to the mesopause as evidenced by the presence of Polar Mesospheric Summer Echoes (PMSEs) [*Rapp and Lübken*, 2004; *Russell III et al.*, 2010]. These ice particles are thought to nucleate near the cold point, and then grow and sediment to form the visible cloud [*von Zahn and Berger*, 2003; *Rapp and Thomas*, 2006].

The evidence from this study and others suggests most PMCs that form between 120 and 135 K will be ice Isd crystals. As the ice particles sediment and reach warmer temperatures, the ice I<sub>sd</sub> structure will likely persist [*Murray et al.*, 2015a]. Ice crystals that sediment into regions warmer than ~140 K may then grow through the deposition of ice Ih. At the mesopause, conditions are sometimes cold enough that LDA ice particles can form, as previously suggested [Lubken et al., 2009; Murray and Jensen, 2010]. Upon warming these LDA ice particles would likely transition to ice Isd so that LDA ice is probably a transient phase and will not be present in the bulk of PMCs, consistent with previous satellite measurement which indicate crystalline ice dominates PMCs [Hervig and Gordley, 2010]. However, LDA ice may be important in the early stages of cloud development. For example, the equilibrium vapour pressure of LDA ice is substantially larger than crystalline ices, hence this may drive a mass transfer process analogous to the Werner-Bergeron-Findeisen process in Earth's tropospheric clouds composed of mixtures of liquid water and ice [Murray et al., 2012; Nachbar et al., 2018b]. In this process, a few particles of crystalline ice grow at the expense of LDA particles, potentially leading to a reduced number of hydrometeors of larger size. Overall, it is striking that all three phases of ice may be present and important in the evolution of mesospheric clouds on Earth.

### 5.2 Mars

For Mars, as can be seen from the Mars Pathfinder entry profile shown in Figure 1, the situation is distinct to that on Earth. A colder temperature profile, with a wide altitude range of extreme cold temperatures leads to a large saturated region ( $S \ge 1$  for < 135 K at 58 – 87 km in this profile), one that is significantly more supersaturated than for Earth. Mars lacks a significant ozone layer and a clear mesopause, leading to a much more variable altitude at which the temperature minimum occurs. This temperature minimum is influenced heavily by dynamical processes such as gravity waves causing a much larger range of altitudes (30 – 90 km) in the atmosphere (especially in the mountainous equatorial regions for solar longitude Ls = ~0 - 150°)

where water ice clouds are typically observed [*Vincendon et al.*, 2011; *Fedorova et al.*, 2020; *Heavens et al.*, 2020].

Water ice clouds observed in the mesosphere (~80 km) with the strongest temperature minima (~80 K) will likely form as particles of LDA ice which could transform to ice I<sub>sd</sub> at warmer temperatures, in a similar way to PMCs on Earth. Observations during the dust storm season have highlighted the presence of water ice at altitudes of ~80 – 90 km at temperatures < 150 K, suggesting that ice I<sub>sd</sub> and ice I<sub>h</sub> may also be possible for these warmer mesospheric clouds [*Fedorova et al.*, 2020]. Similarly, clouds forming in the lower mesosphere (< 60 km) are more likely to lead to ice I<sub>sd</sub> and possibly ice I<sub>h</sub> particle formation depending on the specific conditions. Water ice clouds can also form all the way down to the Martian surface (~0 – 30 km) at temperatures from ~180 – 215 K, under similar temperature conditions to TTL cirrus on Earth, and so should also form primarily particles of ice I<sub>h</sub> [*Whiteway et al.*, 2009; *Wilson and Guzewich*, 2014].

### 5.3 Venus

The profile shown in Figure 1 for the Venusian upper atmosphere is taken from the SOIR Instrument onboard the Venus Express satellite in the 70 - 80° latitude region [Mahieux et al., 2012]. A temperature minimum (~80 K) is observed in this profile around an altitude of 125 km, although the temperature can drop as low as 60 K based on other observations [Mahieux et al., 2015]. This temperature minimum is often observed around 120 - 130 km for measurements taken at the terminators, though the extent of the temperature minima is variable. The super-saturated region shown here occurs at temperatures < 130 K, over a 119 - 129 km altitude range. The Venusian mesosphere where ice clouds may form is more rarefied (~10<sup>-5</sup> Pa) than the coldest regions observed on Earth and Mars  $(1 - 10^{-2} Pa)$ , hence much more extreme temperature minima are required for nucleation. It is suggested that if clouds form at these high altitudes they will likely produce small subvisible nanoparticles [Turco et al., 1983]. Small particles of water ice forming at ~60 - 120 K could persist as LDA ice and are unlikely to undergo significant crystallization to ice Isd before sublimation under warmer conditions. The temperatures on Venus are probably too high to form the high density counterpart of LDA ice, HDA ice, which requires vapour deposition below 38 K to form [Jenniskens and Blake, 1994]. Given the need for temperatures below 130 K to produce supersaturated air, Venus is the only atmosphere discussed here where ice  $I_h$  is unlikely to form and where the dominant phase may be LDA.

#### 5.4 Ice nucleation and crystal shape

It is thought that an amorphous phase is the initial material to nucleate when particles nucleate homogeneously from the vapour, since the nucleation barrier for the formation of an amorphous cluster is smaller than for a crystalline cluster [*Huang and Bartell*, 1995; *Murray and Jensen*, 2010]. Of particular relevance to planetary mesospheric clouds, *Nachbar et al.* [2018a] concluded that the initial phase that forms for temperatures < 160 K is always LDA ice. Furthermore, an ice activation model using the bulk properties of LDA ice was also able to satisfactorily reproduce heterogeneous ice nucleation experiments at ~140 - 150 K [*Duft et al.*, 2019]. The mesospheric clouds discussed for Earth, Mars and Venus all form below 150 K, suggesting that at least the nucleation kinetics of ice particles forming by deposition in

these atmospheres will be governed by the properties of LDA ice, including associated parameters such as vapor pressure, surface tension and density. Depending on the temperature, timescale, growth rate and size of the ice particles, the initial amorphous clusters formed below 160 K may transform to ice  $I_{sd}$ . If the temperature is low enough ( $\leq 120$  K) or either growth or particle lifetime is limited, they can instead persist as LDA ice. *Hudait and Molinero* [2016] suggest that at the high supersaturations possible in experiments and in planetary upper atmospheres (e.g.  $S = 10^6 - 10^8$ ), crystalline ice  $I_{sd}$  is more likely to form than ice  $I_h$ . This saturation range is comparable to that seen over the 120 – 135 K range in this study (1.4 x  $10^8 - 6.6 \times 10^5$ ) where ice  $I_{sd}$  is observed. Precipitation of particles from colder to warmer regions, short-lived perturbations of the temperature profile cause by gravity waves, and the effect of varying supersaturation, provides the possibility of multiple phases being present in these clouds. The phase that grows then determines the possibilities for the overall shape of the crystal.

The lack of long-range order in LDA ice suggests that once the particles grow large enough that the shape is no longer significantly influenced by the addition of a single water molecule (diameter of a H<sub>2</sub>O molecule ~0.27 nm), they will be approximately spherical. Crystals with a structure of ice I<sub>sd</sub> (the likely most abundant crystalline phase in the mesospheric clouds for Earth and Mars) are trigonal, forming columns or plates with aspect ratios larger than spherical particles [*Murray et al.*, 2015b]. Crystals of ice I<sub>h</sub> will also be anisotropic and will lead to columns and plates, albeit with six-fold symmetry. Ice I<sub>c</sub> (which is not expected to form for the clouds discussed here based on experimental evidence), would form more compact isotropic crystals. For PMCs on Earth, the expectation of primarily ice I<sub>sd</sub> and ice I<sub>h</sub> is consistent with observations of high aspect ratio, non-spherical crystals [*Rapp et al.*, 2007; *Hervig and Gordley*, 2010]. For Mars, the elongated crystal shapes of the tropospheric water ice (ice I<sub>h</sub>) clouds differentiate them from CO<sub>2</sub>-I ice clouds, which form more compact crystal shapes similar to ice I<sub>c</sub>, such as cubes and octahedra [*Mangan et al.*, 2017b].

## 6. Conclusions

Diffraction patterns of unannealed, vapor deposited water ice at 10 mbar (20 mbar at 88 K) have been presented in this study at temperatures from 88 – 145 K. LDA ice was found to form at deposition temperatures of 88 – 120 K with a boundary at ~120 K between amorphous and crystalline ice deposition. Ice  $I_{sd}$  was deposited at ~120 – 135 K and ice  $I_h$  at 140 and 145 K. LDA ice was also annealed from 110 up to 160 K producing ice  $I_{sd}$ , highlighting the difference in crystal structure between ice deposited directly at a particular temperature and ice deposited at lower temperature and then warmed to the same temperature.

The crystal structure of ice I in the mesospheric clouds of Earth, Mars and Venus was inferred from the experiments undertaken in this study. For Earth and Mars conditions should exist for all three phases of ice I to occur, with the predominant structure in mesospheric clouds likely to be ice  $I_{sd}$ , and LDA ice under extreme low temperature conditions. In contrast, lower altitude clouds on Earth and Mars are likely to be comprised of ice  $I_h$  particles. On Venus, clouds that may form around 125 km would likely be LDA ice. Particles of ice  $I_{sd}$  and  $I_h$  will form anisotropic crystals, such as

needles and plates, while LDA ice should lead to more spherical particles. Overall, the results of this study show that amorphous, metastable stacking disordered and stable hexagonal ice may form and influence the properties of clouds in the mesospheres of terrestrial planets.

## Acknowledgments

This work was supported by grant ST/T000279/1 from the UK Science and Technology Facilities Council. The data from this study can be accessed via (awaiting DOI for Research Data Leeds Repository).

## References

- Chamberlain, S., A. Mahieux, S. Robert, A. Piccialli, L. Trompet, A. C. Vandaele, and V. Wilquet (2020), SOIR/VEx observations of water vapor at the terminator in the Venus mesosphere, *Icarus*, *346*, 113819. <u>https://doi.org/10.1016/j.icarus.2020.113819</u>.
- D'Angelo, G., and M. Podolak (2015), Capture and Evolution of Planetesimals in Circumjovian Disks, *The Astrophysical Journal*, *806*(2), 203. <u>https://doi.org/10.1088/0004-637x/806/2/203</u>.
- del Rosso, L., M. Celli, F. Grazzi, M. Catti, T. C. Hansen, A. D. Fortes, and L. Ulivi (2020), Cubic ice Ic without stacking defects obtained from ice XVII, *Nature Materials*, *19*(6), 663-668. <u>https://doi.org/10.1038/s41563-020-0606-y</u>.
- del Rosso, L., F. Grazzi, M. Celli, D. Colognesi, V. Garcia-Sakai, and L. Ulivi (2016), Refined Structure of Metastable Ice XVII from Neutron Diffraction Measurements, *J. Phys. Chem. C*, 120(47), 26955-26959. https://doi.org/10.1021/acs.jpcc.6b10569.
- Dowell, L. G., and A. P. Rinfret (1960), Low-Temperature Forms of Ice as Studied by X-Ray Diffraction, *Nature*, *188*(4757), 1144-1148. <u>https://doi.org/10.1038/1881144a0</u>.
- Duft, D., M. Nachbar, and T. Leisner (2019), Unravelling the microphysics of polar mesospheric cloud formation, *Atmos. Chem. Phys.*, *19*(5), 2871-2879. https://doi.org/10.5194/acp-19-2871-2019.
- Falenty, A., T. C. Hansen, and W. F. Kuhs (2014), Formation and properties of ice XVI obtained by emptying a type sll clathrate hydrate, *Nature*, *516*(7530), 231-233. https://doi.org/10.1038/nature14014.
- Fedorova, A. A., F. Montmessin, O. Korablev, M. Luginin, A. Trokhimovskiy, D. A. Belyaev, et al. (2020), Stormy water on Mars: The distribution and saturation of atmospheric water during the dusty season, *Science*, eaay9522. https://doi.org/10.1126/science.aay9522.
- Forget, F., F. Hourdin, R. Fournier, C. Hourdin, O. Talagrand, M. Collins, et al. (1999), Improved general circulation models of the Martian atmosphere from the surface to above 80 km, *J. Geophys. Res. Planets*, *104*(E10), 24155-24175. https://doi.org/10.1029/1999je001025.
- Forget, F., F. Montmessin, J.-L. Bertaux, F. González-Galindo, S. Lebonnois, E. Quémerais, et al. (2009), Density and temperatures of the upper Martian atmosphere measured by stellar occultations with Mars Express SPICAM, *J. Geophys. Res. Planets*, 114(E1). <u>https://doi.org/10.1029/2008je003086</u>.
- Gillett, F. C., and W. J. Forrest (1973), Spectra of the Becklin-Neugebauer point source and the Kleinmann-Low nebula from 2.8 to 13.5 microns, *The Astrophysical Journal*, *179*, 483. <u>https://doi.org/10.1086/151888</u>.
- Hallbrucker, A., E. Mayer, and G. P. Johari (1989), Glass-liquid transition and the enthalpy of devitrification of annealed vapor-deposited amorphous solid water: a comparison with hyperquenched glassy water, *Journal of Physical Chemistry*, *93*(12), 4986-4990. <u>https://doi.org/10.1021/j100349a061</u>.

- Hansen, T. C., M. M. Koza, and W. F. Kuhs (2008), Formation and annealing of cubic ice: I. Modelling of stacking faults, *Journal of Physics: Condensed Matter*, *20*(28), 285104. https://doi.org/10.1088/0953-8984/20/28/285104.
- Heavens, N. G., D. M. Kass, A. Kleinböhl, and J. T. Schofield (2020), A multiannual record of gravity wave activity in Mars's lower atmosphere from on-planet observations by the Mars Climate Sounder, *Icarus*, *341*, 113630. https://doi.org/10.1016/j.icarus.2020.113630.
- Hervig, M. E., and L. L. Gordley (2010), Temperature, shape, and phase of mesospheric ice from Solar Occultation for Ice Experiment observations, *J. Geophys. Res. Atmos.*, 115(D15). <u>https://doi.org/10.1029/2010jd013918</u>.
- Hervig, M. E., D. E. Siskind, S. M. Bailey, and J. M. Russell (2015), The influence of PMCs on water vapor and drivers behind PMC variability from SOFIE observations, *Journal* of Atmospheric and Solar-Terrestrial Physics, 132, 124-134. <u>https://doi.org/10.1016/j.jastp.2015.07.010</u>.
- Hervig, M. E., M. H. Stevens, L. L. Gordley, L. E. Deaver, J. M. Russell III, and S. M. Bailey (2009), Relationships between polar mesospheric clouds, temperature, and water vapor from Solar Occultation for Ice Experiment (SOFIE) observations, *J. Geophys. Res. Atmos.*, 114(D20). <u>https://doi.org/10.1029/2009jd012302</u>.
- Hervig, M. E., R. E. Thompson, M. McHugh, L. L. Gordley, J. M. Russell III, and M. E. Summers (2001), First confirmation that water ice is the primary component of polar mesospheric clouds, *Geophys. Res. Lett.*, 28(6), 971-974. <u>https://doi.org/10.1029/2000gl012104</u>.
- Hobbs, P. V. (1974), Ice physics, Clarendon Press, Oxford.
- Huang, J., and L. S. Bartell (1995), Kinetics of Homogeneous Nucleation in the Freezing of Large Water Clusters, *Journal of Physical Chemistry*, *99*(12), 3924-3931. https://doi.org/10.1021/j100012a010.
- Hudait, A., and V. Molinero (2016), What Determines the Ice Polymorph in Clouds?, *Journal* of the American Chemical Society, 138(28), 8958-8967. https://doi.org/10.1021/jacs.6b05227.
- Hudait, A., S. Qiu, L. Lupi, and V. Molinero (2016), Free energy contributions and structural characterization of stacking disordered ices, *Physical Chemistry Chemical Physics*, *18*(14), 9544-9553. <u>https://doi.org/10.1039/C6CP00915H</u>.
- Jenniskens, P., and D. F. Blake (1994), Structural transitions in amorphous water ice and astrophysical implications, *Science*, *265*(5173), 753-756. https://doi.org/10.1126/science.11539186.
- Jenniskens, P., and D. F. Blake (1996), Crystallization of Amorphous Water Ice in the Solar System, *The Astrophysical Journal*, *473*(2), 1104-1113. <u>https://doi.org/10.1086/178220</u>.
- Johari, G. P., A. Hallbrucker, and E. Mayer (1991), The dielectric behavior of vapordeposited amorphous solid water and of its crystalline forms, *J. Chem. Phys.*, *95*(4), 2955-2964. <u>https://doi.org/10.1063/1.460897</u>.
- Komatsu, K., S. Machida, F. Noritake, T. Hattori, A. Sano-Furukawa, R. Yamane, et al. (2020), Ice Ic without stacking disorder by evacuating hydrogen from hydrogen hydrate, *Nature Communications*, *11*(1), 464. <u>https://doi.org/10.1038/s41467-020-14346-5</u>.
- Kuhs, W. F., C. Sippel, A. Falenty, and T. C. Hansen (2012), Extent and relevance of stacking disorder in "ice I&It;sub>c&It;/sub>" *Proceedings of the National Academy of Sciences*, 109(52), 21259. <u>https://doi.org/10.1073/pnas.1210331110</u>.
- Loerting, T., K. Winkel, M. Seidl, M. Bauer, C. Mitterdorfer, P. H. Handle, et al. (2011), How many amorphous ices are there?, *Physical Chemistry Chemical Physics*, *13*(19), 8783-8794. <u>https://doi.org/10.1039/C0CP02600J</u>.
- Lubken, F. J., J. Lautenbach, J. Hoffner, M. Rapp, and M. Zecha (2009), First continuous temperature measurements within polar mesosphere summer echoes, *Journal of Atmospheric and Solar-Terrestrial Physics*, *71*(3-4), 453-463. https://doi.org/10.1016/j.jastp.2008.06.001.

- Magalhães, J. A., J. T. Schofield, and A. Seiff (1999), Results of the Mars Pathfinder atmospheric structure investigation, *J. Geophys. Res. Planets*, *104*(E4), 8943-8955. <u>https://doi.org/10.1029/1998je900041</u>.
- Mahieux, A., A. C. Vandaele, S. W. Bougher, R. Drummond, S. Robert, V. Wilquet, et al. (2015), Update of the Venus density and temperature profiles at high altitude measured by SOIR on board Venus Express, *Planetary and Space Science*, 113-114, 309-320. https://doi.org/10.1016/j.pss.2015.02.002.
- Mahieux, A., A. C. Vandaele, S. Robert, V. Wilquet, R. Drummond, F. Montmessin, and J. L. Bertaux (2012), Densities and temperatures in the Venus mesosphere and lower thermosphere retrieved from SOIR on board Venus Express: Carbon dioxide measurements at the Venus terminator, *J. Geophys. Res. Planets*, *117*(E7). <u>https://doi.org/10.1029/2012je004058</u>.
- Malkin, T. L., B. J. Murray, A. V. Brukhno, J. Anwar, and C. G. Salzmann (2012), Structure of ice crystallized from supercooled water, *Proceedings of the National Academy of Sciences*, 109(4), 1041. <u>https://doi.org/10.1073/pnas.1113059109</u>.
- Malkin, T. L., B. J. Murray, C. G. Salzmann, V. Molinero, S. J. Pickering, and T. F. Whale (2015), Stacking disorder in ice I, *Physical Chemistry Chemical Physics*, 17(1), 60-76. <u>https://doi.org/10.1039/C4CP02893G</u>.
- Mangan, T. P., V. L. Frankland, B. J. Murray, and J. M. C. Plane (2017a), The fate of meteoric metals in ice particles: Effects of sublimation and energetic particle bombardment, *Journal of Atmospheric and Solar-Terrestrial Physics*, 161, 143-149. <u>https://doi.org/10.1016/j.jastp.2017.07.002</u>.
- Mangan, T. P., C. G. Salzmann, J. M. C. Plane, and B. J. Murray (2017b), CO2 ice structure and density under Martian atmospheric conditions, *Icarus*, *294*, 201-208. <u>https://doi.org/10.1016/j.icarus.2017.03.012</u>.
- Marshall, B. T., L. E. Deaver, R. E. Thompson, L. L. Gordley, M. J. McHugh, M. E. Hervig, and J. M. Russell Iii (2011), Retrieval of temperature and pressure using broadband solar occultation: SOFIE approach and results, *Atmos. Meas. Tech.*, *4*(5), 893-907. https://doi.org/10.5194/amt-4-893-2011.
- Millot, M., S. Hamel, J. R. Rygg, P. M. Celliers, G. W. Collins, F. Coppari, et al. (2018), Experimental evidence for superionic water ice using shock compression, *Nature Physics*, 14(3), 297-302. <u>https://doi.org/10.1038/s41567-017-0017-4</u>.
- Mitlin, S., and K. T. Leung (2004), Temporal evolution of an ultrathin, noncrystalline ice deposit at crystallization near 160 K studied by FT-IR reflection-absorption spectroscopy, *Canadian Journal of Chemistry*, 82(6), 978-986. https://doi.org/10.1139/v04-076.
- Murphy, D. M. (2003), Dehydration in cold clouds is enhanced by a transition from cubic to hexagonal ice, *Geophys. Res. Lett.*, *30*(23). <u>https://doi.org/10.1029/2003GL018566</u>.
- Murphy, D. M., and T. Koop (2005), Review of the vapour pressures of ice and supercooled water for atmospheric applications, *Q.J.R. Meteorol. Soc.*, *131*(608), 1539-1565. <u>https://doi.org/10.1256/qj.04.94</u>.
- Murray, B. J., and E. J. Jensen (2010), Homogeneous nucleation of amorphous solid water particles in the upper mesosphere, *Journal of Atmospheric and Solar-Terrestrial Physics*, *72*(1), 51-61. <u>https://doi.org/10.1016/j.jastp.2009.10.007</u>.
- Murray, B. J., D. A. Knopf, and A. K. Bertram (2005), The formation of cubic ice under conditions relevant to Earth's atmosphere, *Nature*, *434*(7030), 202-205. <u>https://doi.org/10.1038/nature03403</u>.
- Murray, B. J., T. L. Malkin, and C. G. Salzmann (2015a), The crystal structure of ice under mesospheric conditions, *Journal of Atmospheric and Solar-Terrestrial Physics*, 127, 78-82. <u>https://doi.org/10.1016/j.jastp.2014.12.005</u>.
- Murray, B. J., D. O'Sullivan, J. D. Atkinson, and M. E. Webb (2012), Ice nucleation by particles immersed in supercooled cloud droplets, *Chemical Society Reviews*, 41(19), 6519-6554. <u>https://doi.org/10.1039/C2CS35200A</u>.

- Murray, B. J., and J. M. C. Plane (2003), The uptake of atomic oxygen on ice films: Implications for noctilucent clouds, *Physical Chemistry Chemical Physics*, *5*(19), 4129-4138. <u>https://doi.org/10.1039/B3055555H</u>.
- Murray, B. J., and J. M. C. Plane (2005a), Modelling the impact of noctilucent cloud formation on atomic oxygen and other minor constituents of the summer mesosphere, *Atmos. Chem. Phys.*, *5*(4), 1027-1038. <u>https://doi.org/10.5194/acp-5-1027-2005</u>.
- Murray, B. J., and J. M. C. Plane (2005b), Uptake of Fe, Na and K atoms on lowtemperature ice: implications for metal atom scavenging in the vicinity of polar mesospheric clouds, *Physical Chemistry Chemical Physics*, *7*(23), 3970-3979. <u>https://doi.org/10.1039/B508846A</u>.
- Murray, B. J., C. G. Salzmann, A. J. Heymsfield, S. Dobbie, R. R. Neely, III, and C. J. Cox (2015b), Trigonal Ice Crystals in Earth's Atmosphere, *Bulletin of the American Meteorological Society*, *96*(9), 1519-1531. <u>https://doi.org/10.1175/BAMS-D-13-00128.1</u> %J Bulletin of the American Meteorological Society.
- Nachbar, M., D. Duft, and T. Leisner (2018a), The vapor pressure over nano-crystalline ice, *Atmos. Chem. Phys.*, *18*(5), 3419-3431. <u>https://doi.org/10.5194/acp-18-3419-2018</u>.
- Nachbar, M., D. Duft, and T. Leisner (2018b), Volatility of Amorphous Solid Water, *Journal of Physical Chemistry B*, 122(43), 10044-10050. https://doi.org/10.1021/acs.jpcb.8b06387.
- Nachbar, M., D. Duft, and T. Leisner (2019), The vapor pressure of liquid and solid water phases at conditions relevant to the atmosphere, *J. Chem. Phys.*, *151*(6), 064504. https://doi.org/10.1063/1.5100364.
- Plane, J. M. C., J. D. Carrillo-Sanchez, T. P. Mangan, M. M. J. Crismani, N. M. Schneider, and A. Määttänen (2018), Meteoric Metal Chemistry in the Martian Atmosphere, J. Geophys. Res. Planets, 123(3), 695-707. <u>https://doi.org/10.1002/2017je005510</u>.
- Plane, J. M. C., B. J. Murray, X. Chu, and C. S. Gardner (2004), Removal of Meteoric Iron on Polar Mesospheric Clouds, *Science*, *304*(5669), 426. https://doi.org/10.1126/science.1093236.
- Rapp, M., and F. J. Lübken (2004), Polar mesosphere summer echoes (PMSE): review of observations and current understanding, *Atmos. Chem. Phys.*, 4(4), 2601-2633. <u>https://doi.org/10.5194/acp-4-2601-2004</u>.
- Rapp, M., and G. E. Thomas (2006), Modeling the microphysics of mesospheric ice particles: Assessment of current capabilities and basic sensitivities, *Journal of Atmospheric and Solar-Terrestrial Physics*, 68(7), 715-744. https://doi.org/10.1016/j.jastp.2005.10.015.
- Rapp, M., G. E. Thomas, and G. Baumgarten (2007), Spectral properties of mesospheric ice clouds: Evidence for nonspherical particles, *J. Geophys. Res. Atmos.*, 112(D3). <u>https://doi.org/10.1029/2006jd007322</u>.
- Russell III, J. M., P. Rong, S. M. Bailey, M. E. Hervig, and S. V. Petelina (2010), Relationship between the summer mesopause and polar mesospheric cloud heights, *J. Geophys. Res. Atmos.*, *115*(D16). <u>https://doi.org/10.1029/2010JD013852</u>.
- Salzmann, C. G., and B. J. Murray (2020), Ice goes fully cubic, *Nature Materials*, *19*(6), 586-587. <u>https://doi.org/10.1038/s41563-020-0696-6</u>.
- Salzmann, C. G., P. G. Radaelli, B. Slater, and J. L. Finney (2011), The polymorphism of ice: five unresolved questions, *Physical Chemistry Chemical Physics*, *13*(41), 18468-18480. <u>https://doi.org/10.1039/C1CP21712G</u>.
- Shilling, J. E., M. A. Tolbert, O. B. Toon, E. J. Jensen, B. J. Murray, and A. K. Bertram (2006), Measurements of the vapor pressure of cubic ice and their implications for atmospheric ice clouds, *Geophysical Research Letters*, 33(17). <u>https://doi.org/10.1029/2006gl026671</u>.
- Smith, R. S., C. Huang, E. K. L. Wong, and B. D. Kay (1996), Desorption and crystallization kinetics in nanoscale thin films of amorphous water ice, *Surface Science*, 367(1), L13-L18. <u>https://doi.org/10.1016/S0039-6028(96)00943-0</u>.

- Thoeny, A. V., T. M. Gasser, and T. Loerting (2019), Distinguishing ice β-XV from deep glassy ice VI: Raman spectroscopy, *Physical Chemistry Chemical Physics*, 21(28), 15452-15462. <u>https://doi.org/10.1039/C9CP02147G</u>.
- Thomas, G. E. (1991), Mesospheric clouds and the physics of the mesopause region, *Reviews of Geophysics*, *29*(4), 553-575. <u>https://doi.org/10.1029/91RG01604</u>.
- Turco, R. P., O. B. Toon, R. C. Whitten, and R. G. Keesee (1983), Venus: Mesospheric hazes of ice, dust, and acid aerosols, *Icarus*, *53*(1), 18-25. https://doi.org/10.1016/0019-1035(83)90017-9.
- Vincendon, M., C. Pilorget, B. Gondet, S. Murchie, and J. P. Bibring (2011), New near-IR observations of mesospheric CO<sub>2</sub> and H<sub>2</sub>O clouds on Mars, *J. Geophys. Res. Planets*, *116.* <u>https://doi.org/10.1029/2011je003827</u>.
- von Zahn, U., and U. Berger (2003), Persistent ice cloud in the midsummer upper mesosphere at high latitudes: Three-dimensional modeling and cloud interactions with ambient water vapor, *J. Geophys. Res. Atmos.*, *108*(D8). https://doi.org/10.1029/2002JD002409.
- Whiteway, J. A., L. Komguem, C. Dickinson, C. Cook, M. Illnicki, J. Seabrook, et al. (2009), Mars Water-Ice Clouds and Precipitation, *Science*, 325(5936), 68. <u>https://doi.org/10.1126/science.1172344</u>.
- Wilson, R. J., and S. D. Guzewich (2014), Influence of water ice clouds on nighttime tropical temperature structure as seen by the Mars Climate Sounder, *Geophys. Res. Lett.*, 41(10), 3375-3381. <u>https://doi.org/10.1002/2014gl060086</u>.