

Applicability of the Long-chain Diol Index (LDI) as a sea surface temperature proxy in the Arabian Sea

Zeynep Erdem¹, Julie Lattaud², Marit R van Erk³, Eveline M Mezger³, Gert-Jan Reichart⁴, Andreas Lückge⁵, Jaap S Sinninghe Damste¹, and Stefan Schouten⁶

¹Royal Netherlands Institute for Sea Research

²ETHZ

³NIOZ Royal Netherlands Institute for Sea Research

⁴NIOZ

⁵Federal Institute for Geosciences and Natural Resources

⁶Royal NIOZ

November 24, 2022

Abstract

The long-chain diol index (LDI) is a relatively new proxy for sea surface temperature (SST) which has been rarely applied in upwelling regions. Here, we evaluated its application by comparison with other SST records obtained by commonly used proxies, i.e. the Mg/Ca ratio of the planktonic foraminifera species *37* and $\text{Mg}/\text{Ca}_{G.\text{ruber}}$. SST records revealed an increase of 0.6-3.4°C from the Last Glacial Maximum to the late Holocene with somewhat higher amplitude in the northern part of the Arabian Sea than compared to the western part. In contrast, the LDI SSTs did not reveal major changes during the last glacial-interglacial transition which was followed by a decreasing trend during the Holocene. The LGM vs the Holocene LDI SSTs ranged between -0.2 and -2.7°C. Particularly at one record, offshore Oman, the SST decrease from during the Holocene was high in amplitude, suggesting a potential cold bias, possibly related to changes in upwelling intensity during the Holocene. This indicates that care has to be taken when applying the LDI for annual mean SST reconstruction in upwelling regions.

Hosted file

erdem et al_supp.info_arabian sea ldi.docx available at <https://authorea.com/users/549601/articles/603642-applicability-of-the-long-chain-diol-index-ldi-as-a-sea-surface-temperature-proxy-in-the-arabian-sea>

Applicability of the Long chain Diol Index (LDI) as a sea surface temperature proxy in the Arabian Sea

Z. Erdem¹, J. Lattaud^{1,*}, M.R. van Erk^{1,}, E. Mezger¹, G.-J. Reichart^{1,2}, A. Lückge³, J.S. Sinninghe Damsté^{1,2}, S. Schouten^{1,2}**

¹ NIOZ Royal Netherlands Institute for Sea Research, P.O. Box 59, 1790 AB Den Burg, Texel, the Netherlands

² Utrecht University, Faculty of Geosciences, P.O. Box 80115, 3508 TC Utrecht, the Netherlands

³ Bundesanstalt für Geowissenschaften und Rohstoffe, Stilleweg 2 30655 Hannover, Germany

Corresponding author: Zeynep Erdem (zeynep.erdem@nioz.nl)

* current address: Biogeoscience group, ETH Zurich, Sonneggstrasse 5, Zurich, Switzerland.

** current address: Max Planck Institute for Marine Microbiology, Celsiusstrasse 1, 28359 Bremen, Germany

Key Points:

- We applied the LDI as a paleothermometer to sedimentary records from the Arabian Sea focusing on the last glacial – interglacial transition.
- The sea surface temperature records showed decreasing values with the onset of the deglaciation, different from other proxy records.
- Strong upwelling and high nutrient input may cause this cold bias, which was observed particularly at one record, offshore Oman.

Abstract

The long-chain diol index (LDI) is a relatively new proxy for sea surface temperature (SST) which has been rarely applied in upwelling regions. Here, we evaluated its application by comparison with other SST records obtained by commonly used proxies, i.e. the Mg/Ca ratio of the planktonic foraminifera species *Globigerinoides ruber* and the alkenone paleothermometer $U^{K'}_{37}$. We focused on the last glacial–interglacial transition of four different sedimentary archives from the western and northern Arabian Sea, which are currently under the influence of monsoon-induced upwelling and the associated development of an oxygen minimum zone. The $U^{K'}_{37}$ and $Mg/Ca_{G.ruber}$ SST records revealed an increase of 0.6–3.4°C from the Last Glacial Maximum to the late Holocene with somewhat higher amplitude in the northern part of the Arabian Sea than compared to the western part. In contrast, the LDI SSTs did not reveal major changes during the last glacial–interglacial transition which was followed by a decreasing trend during the Holocene. The LGM vs the Holocene LDI SSTs ranged between -0.2 and -2.7°C. Particularly at one record, offshore Oman, the SST decrease from during the Holocene was high in amplitude, suggesting a potential cold bias, possibly related to changes in upwelling intensity during the Holocene. This indicates that care has to be taken when applying the LDI for annual mean SST reconstruction in upwelling regions.

1 Introduction

Reconstructing past ocean temperature is essential for understanding past climate change, its drivers and responsible mechanisms. Several proxies were established over the past decades in order to reconstruct sea surface temperature (SST) and changes therein. Planktonic foraminiferal assemblages (by transfer function; (Imbrie and Kipp, 1971; Pflaumann et al., 1996), stable oxygen isotopes ($\delta^{18}\text{O}$) (Erez and Luz, 1983; Urey et al., 1954) and Mg/Ca ratios of planktonic foraminifera species (Elderfield and Ganssen, 2000; Nürnberg et al., 1996) and organic proxies, such as the U^{K}_{37} index (Brassell et al., 1986; Prahl and Wakeham, 1987) based on the long chain alkenones synthesized by haptophyte algae, have been widely applied as paleothermometers. However, discrepancies between various SST proxy records as well as with SST reconstructions based on climate model simulations have been observed. This is particularly the case concerning the reconstructed Last Glacial Maximum (LGM) SSTs reported in the tropical oceans and the upwelling regions where proxy records and models show disagreement with each other as well as with global estimates (e.g., Kucera et al., 2005; Mix et al., 2001; Shakun et al., 2012; Waelbroeck et al., 2009). Accordingly, there is a need for an increased fundamental understanding of the processes that control the imprint of temperature on these proxies as well as the processes underlying the observed offsets.

In the last decade, the LDI (Long-chain Diol Index; Rampen et al., 2012), which is based on the distribution of certain long-chain diols (LCDs), has been developed. The LDI is based on the relative abundances of 1,13- and 1,15-diols of which culture studies showed that they are produced by freshwater and marine eustigmatophyte algae. However, in the marine realm, it is still uncertain what the source organisms are (e.g., Balzano et al., 2018; Rampen et al., 2014; Volkman et al., 1999; Volkman et al., 1992). The number of studies of late Quaternary LDI applications has been rapidly increasing recently (Dauner et al., 2019; de Bar et al., 2018; Häggi et al., 2019; Jonas et al., 2017; Lattaud et al., 2018; Lopes dos Santos et al., 2013; Rodrigo-Gámiz et al., 2015; Warnock et al., 2018; Zhu et al., 2018) giving broader information on global SST variations in the past. Recently, a more extended calibration for the LDI became available, providing further insight into the main constraints on its applicability (de Bar et al., 2020). Overall, the new calibration dataset was in line with downcore applications but also showed that i) the LDI has limitations in certain regions, such as in the front of river mouths, in low salinity regions, and at high latitudes; ii) the abundance of LCDs other (e.g., 1,12 and 1,14-diols) than those used for the LDI may indicate a cold bias of LDI-derived SST estimations. It was, therefore, suggested to use other LCD-based indices (e.g., diol upwelling index (DI) and the relative abundance of C_{28} 1,12-diol (FC_{28} 1,12-diol)) prior to LDI applications to constrain potential biases (de Bar et al., 2020). However, many uncertainties with respect to the application of the LDI still remain, such as those related to changing nutrient regimes, seasonality, and other potential changes in the water column (e.g., mixed layer depth, oxygenation). Moreover, SST reconstructions from (sub)tropical upwelling regions indicated discrepancies and uncertainties between proxies as well as modelling approaches (e.g., Otto-Bliesner et al., 2009). Multi-proxy

approaches can, therefore, be helpful to constrain these uncertainties (Kucera et al., 2005; Mix et al., 2001; Waelbroeck et al., 2009) as well as to better our understanding of past climate in tropical upwelling regions.

Here, for the first time, we applied the LDI to generate four sedimentary records from different locations in the Arabian Sea focusing on the last glacial-interglacial period. The Arabian Sea is characterised by strong upwelling conditions dependent on monsoon activity, particularly in the west, with relatively warm surface waters during the year (24-27°C Arabian Sea (Locarnini et al., 2013)). The Arabian Sea has been the focus of studies on past SST reconstructions for more than two decades (e.g., Böll et al., 2015; Doose-Rolinski et al., 2001; Emeis et al., 1995; Gaye et al., 2018; Huguët et al., 2006; Ivanochko et al., 2005; Rostek et al., 1997; Schulz et al., 1998; Sonzogni et al., 1998). These studies have revealed contrasting trends and magnitudes of SST changes as reconstructed by various proxies as well as simulated using state-of-the-art climate models, particularly concerning the last glacial-interglacial transition. Application of the foraminifera transfer function suggested that the western Arabian Sea experienced almost no cooling during the LGM (CLIMAP Project Members, 1988), whereas the alkenone paleothermometer indicated 1-4°C warming between LGM and the late Holocene (Sonzogni et al., 1998). Moreover, modeling approaches reported a variable LGM SST range (1-3°C cooler compared to modern-day) for the Arabian Sea (Mix et al., 2001; Otto-Bliesner et al., 2009; Waelbroeck et al., 2009). The disagreement between proxy-based estimations have been related to organisms' habitats as well as to regional environmental changes (e.g., Dahl and Opp, 2006; Rosell-Melé et al., 2004; Sonzogni et al., 1998) and has been recognized as a common issue in the SST reconstruction of the LGM-Holocene warming in tropical regions (e.g., Waelbroeck et al., 2009). In this study, our aim is to test the applicability of the LDI using four sedimentary records recovered from the western and northern Arabian Sea by comparing the obtained reconstructed SST records with other SST proxy estimates from the same records and evaluate the potential application of the LDI in a tropical upwelling region.

2 Research Setting


In the Arabian Sea, the seasonal cycle is regulated by the Indian monsoon, which is driven by the pressure gradient between the Tibet Plateau and the South Indian Ocean (Rixen et al., 2000; Webster et al., 1998; Wyrtki, 1973). Low temperatures over the Tibetan Plateau and the Himalayas in boreal winter (January – March) causes a high-pressure system over Central Asia and hence a strong pressure gradient towards the low-pressure cell located in the Indian Ocean. During this period, dry and cold northeasterly winds prevail over the Arabian Sea, resulting in cooler surface waters (Webster et al., 1998), particularly in the north of the Arabian Sea (Figure 1a). During boreal summer (July – September), a strong pressure gradient between the Indian-Tibetan low-pressure cell and a high-pressure belt over the Southern Ocean results in powerful southwesterly winds. During this southwest (SW) monsoon period, the Intertropical Convergence Zone (ITCZ) is positioned in the north (Figure 1b). As a result of these strong, warm and humid SW winds, upwelling of nutrient-rich waters occurs along the margin of Oman

to Somalia, leading to lower SSTs (24-25°C) in the western Arabian Sea in comparison with the rest of the region and annual mean SST (e.g., (Haake et al., 1993; Locarnini et al., 2013; Wakeham et al., 2002). During the NE monsoon the upwelling is surprised and waters above the thermocline are convectively mixed until the top of the oxygen minimum zone (Wakeham et al., 2002), as northernmost Arabian Sea exhibits 3-4°C cooling in the surface waters (Dahl and Oppo, 2006).

The Arabian Sea is one of the most productive open marine systems in the world today (Qasim, 1982; Smith et al., 1998). Whilst primary productivity has its annual maximum during the SW monsoon, particularly in the west due to enhanced upwelling, it is substantially less during the NE monsoon (Haake et al., 1993; Honjo et al., 1999; Wakeham et al., 2002). The combination of high productivity, subsequent mineralization of organic matter as it settles through the water column, and consumed dissolved oxygen together with sluggish ventilation of intermediate waters in the Arabian Sea results in one of the strongest OMZs in the world (e.g., (Helly and Levin, 2004; Paulmier and Ruiz-Pino, 2009; Wyrski, 1971). The OMZ, defined as waters with oxygen concentrations <0.5 ml/l (~ 22 μ M), is located in the Arabian Sea at water depths of ca. 150-1200 m (Figure 1c) (Acharya and Panigrahi, 2016; Helly and Levin, 2004; Olson et al., 1993; Sarma, 2002; Wyrski, 1973). Despite the higher primary productivity and organic matter export to the deep sea in the western part of the region, the OMZ is much more pronounced in the northeast east (Acharya and Panigrahi, 2016; Naqvi, 1991; Rixen et al., 2014). This is mainly caused by the prevailing subsurface currents and intermediate water masses in the region, which is strongly influenced by the lack of an opening to the north. Ventilation of OMZ waters is facilitated by three water masses. Firstly, high salinity Persian Gulf Water (PGW) in the north flowing south and eastwards (Prasad et al., 2001), secondly, less saline but denser Red Sea Water (RSW) flowing along the Oman coast (Shankar et al., 2005) and, thirdly, oxygenated intermediate depth Indian Central Water (ICW) entering the Arabian Sea from the south. The ICW is initially rich in oxygen but by the time it reaches the Arabian Sea, it is oxygen depleted and nutrient-rich (Olson et al., 1993). Recently it was shown that the influence of ICW in the region is observed predominantly during the SW Monsoon (Schmidt et al., 2020). The intense OMZ causes denitrification within the water column (Wyrski, 1973) with strongest rates observed in the northeast - east Arabian Sea (Gaye-Haake et al., 2005), where the maximum expansion of the OMZ is observed (Figure 1c).

Sedimentary archives from the Arabian Sea suggest a teleconnection between the Indian monsoon and Northern Hemisphere (NH) climate variability (e.g., Böning and Bard, 2009; Clemens and Prell, 2003; Deplazes et al., 2013; Deplazes et al., 2014; Reichert et al., 1998; Reichert et al., 2002; Schulte and Müller, 2001; Schulz et al., 1998; Sirocko et al., 1996). Interstadials are generally characterized by an intensified Arabian Sea OMZ, while during stadials the strength of the OMZ is substantially reduced (Altabet et al., 2002; Gaye et al., 2018; Pichevin et al., 2007; Reichert et al., 1998; Schulz et al., 1998). Meanwhile, proxy evidence has suggested contrasting differences in the west and east of Arabian Sea in terms of upwelling intensity and surface water productivity (e.g., Emeis et al., 1995; Schulte et al., 1999). Moreover,

with the onset of the last deglaciation with interruptions during the cold periods, stronger SW Monsoon, intensified upwelling and enhanced productivity were reported in the records from the western Arabian Sea (Gaye et al., 2018; Marcantonio et al., 2001; Rampen et al., 2008; Singh et al., 2011; Sirocko et al., 1996).



3 Materials and Methods

3.1 Sediment cores and age models

Four sediment cores were studied. They were collected from different areas during different expeditions; SO42-74KL (in 1986; Kasten core) and NIOP905 (in 1993; piston core) from western Arabian Sea, Oman - Somali upwelling zone; SO130-289KL (in 1998; Kasten core) and PASOM3 (in 2009; gravity core) from northern Arabian Sea within the OMZ from offshore Pakistan and the Murray Ridge, respectively (Figure 1, Table 1). The age models of cores NIOP905 and 74KL were published by Huguët et al. (2006) and of core 289KL by

Deplazes et al. (2013). We tuned the age model of core 74KL by fine-tuning of total organic carbon (TOC) to reflectance (L*) profile of core 289KL (Supp. Info. Figure S1). The age model of core PASOM3 is primarily based on fine-tuning of L* data to the NGRIP (Andersen et al., 2004) and to L* profile of core 289KL (Supp. Info. Text S1). In addition, three AMS radiocarbon ages, based on mixed planktonic foraminifera species (predominantly that belong to genus *Orbulina*) were determined (Beta Analytic Inc., USA). Calibrated ages, (cal kyr BP; 0 BP equivalent to 1950 AD), were calculated using the conventional radiocarbon ages and the Marine13 calibration curve (Reimer et al., 2013). A regional reservoir age of 232 ± 26 years were applied according to the CALIB 7.1 database (Stuiver et al., 2017). The final correlation was based on a total of 41 tie points and the age model was constructed by linear interpolation between these control points. The abundance of the deep-dwelling planktonic foraminifera *Globorotalia truncatulinoides* and *G. crassaformis* were used as an independent verification of the age model, and was based on the existence of so-called *Globorotalia* events (Reichart et al., 1998).

Table 1. Metadata of the sediment cores and references. * age model is tuned (see Supp. Info. Figure S1). n.a.: not available

Core Name	Core location		Water Depth (m)	Metadata				
	Lat (N)	Long (E)		Age Model	U ^{K'} ₃₇	$\delta^{18}\text{O}_{G.ruber}$	Mg/Ca _{G.ruber}	LDI
NIOP905	10°47.0'	51°56.0'	1586	Huguet et al. (2006)	Huguet et al. (2006)	Sirocko et al. (1993)	Anand et al. (2008)	this study
SO42-74KL	14°19.2'	57°20.8'	3212	Huguet et al. (2006)*	Huguet et al. (2006)	Jung et al. (2001)	Dahl and Oppo (2006)	this study
PASOM3	22°19.9'	63°36.0'	1172	this study	this study	this study	this study	this study
SO130-289KL	23°07.3'	66°29.8'	571	Deplazes et al. (2013)	n.a.	n.a.	n.a.	this study

3.2 *Globigerinoides ruber* $\delta^{18}\text{O}$ and Mg/Ca analysis of core PASOM3

For elemental and isotopic analysis, 50 specimens of *G. ruber* were hand-picked per sample from the 250-355 μm fraction. Picked material was later crushed and divided into two parts; one third was saved for stable isotope analysis and the remaining part was used for analyses of the elemental composition. The crushed foraminiferal specimens were cleaned following the Barker protocol (Barker et al., 2003) without the reductive cleaning step. This cleaning protocol includes removing clay particles using ultrapure water, ultrasonication and methanol and removal of organic particles using an alkali buffered 1% H_2O_2 solution with NH_4OH . Stable oxygen ($\delta^{18}\text{O}$) isotope analyses were performed using an automated carbonate device (Kiel IV, Thermo) connected to a Thermo Finnigan MAT 253 Dual Inlet IRMS. Values are reported in permille relative to the Peedee belemnite (PDB) standard in δ notation. For calibration we used NBS19 (limestone) and in-house standard for drift detection (NFHS-1; (Mezger et al., 2016), showing a precision better than 0.1‰ for $\delta^{18}\text{O}$. Measurements for

foraminiferal Mg/Ca composition were performed on an Element 2-ICPMS using low mass resolution. Samples were measured against four ratio calibration standards with a similar matrix and with a drift standard (NFHS-1) measured on every third sample. The monitor standards JCT-1 and NFHS-1 were measured for quality control (Mezger et al., 2016; Okai et al., 2002).

3.3 Long Chain Diol and Alkenone analysis

For LCD analysis, we extracted sediments obtained from cores PASOM3 and 289KL. For cores NIOP905 and 74KL we analyzed the polar fractions previously obtained by Huguët et al. (2006). For PASOM3 and 289KL freeze-dried and homogenized sediments (9 to 18 g from PASOM3 and 0.75 to 11 g from 289KL) were extracted using an Accelerated Solvent Extractor 350 (ASE 350, DIONEX; 100°C and 7.6×10^6 Pa) and subsequently separated over an activated Al_2O_3 column into apolar, ketone and polar fractions. The solvents hexane:DCM 9:1 (v:v), hexane:DCM 1:1 (v:v) and DCM:MeOH 1:1 (v:v) were used to elute the apolar, ketone and polar fractions, respectively. Ketone fractions were used for alkenone analysis of PASOM3 that were conducted on a Hewlett Packard 6890 gas chromatograph (GC) equipped with a flame ionization detector (FID) and helium as the carrier gas, with an autosampler and an injection volume of 1 μL after re-dissolving the ketone fractions in 100 to 800 μL ethyl acetate. Separation was achieved on a fused silica column CP Sil-5 (50 m x 0.32 mm; film thickness 0.12 μm). Initial oven temperature of 70°C increased with 20°C per min to 200°C and subsequently with 3°C per min to 320°C, a temperature which was held for 25 min. Peak areas of $\text{C}_{37:2}$ and $\text{C}_{37:3}$ alkenones were determined by integration. The $U^{K'}_{37}$ values were calculated using the calibration of Müller et al. (1998) and after the equation described by Prahl and Wakeham (1987):

$$\text{with } U^{K'}_{37} = [\text{C}_{37:2}] / ([\text{C}_{37:2}] + [\text{C}_{37:3}]) \quad (1)$$

For LCD analysis polar fractions were silylated by the addition of 15 μL pyridine and 15 μL BSTFA, followed by heating at 60 °C for 30 min. The analysis were carried out using an Agilent 7990B GC coupled to an Agilent 5977A MS following de Bar et al. (2017). The GC was equipped with an on-column injector and fused silica column (25 m x 0.32 mm) coated with CP Sil-5 (film thickness 0.12 μm). Samples were re-dissolved in 300 - 600 μL ethyl acetate and the injection volume was 1 μL . The oven temperature was 70°C at injection and was programmed to increase to 130°C by 20°C/min, and subsequently to 320°C by 4°C /min, which was held for 25 min. Helium flow was held constant at 2 ml/min. The MS operated with an ionization energy of 70 eV and the temperature of the ion source was 250°C and of the interface was 320°C. The total runtime was 75.5 min. Peak areas were defined using SIM of characteristic m/z ratios of LCD ions: 299.3 (C_{26} 1,12-diol and C_{28} 1,14-diol), 313.3 (C_{28} 1,13-diol and C_{30} 1,15-diol), 327.3 (C_{28} 1,12-diol and C_{30} 1,14-diol) and 341.3 (C_{30} 1,13-diol) (Rampen et al., 2012; Versteegh et al., 1997). Full-scan mode allowed for identity confirmation, since the different LCDs show characteristic spectra of fragmentation (Versteegh et al., 1997). The LDI SST were calculated following the calibration of de Bar et al. (2020), based on globally distributed marine surface sediments and covers a temperature range from -1.8 to 30.3°C with a calibration error of 3°C

with $LDI = [C_{30} \text{ 1,15-diol}] / ([C_{28} \text{ 1,13-diol}] + [C_{30} \text{ 1,13-diol}] + [C_{30} \text{ 1,15-diol}])$ (2)

After calculation of the fractional abundances of the C_{28} and C_{30} 1,14-diols, C_{30} 1,15-diol and C_{28} and C_{30} 1,13-diols, the Diol Index (DI) was calculated following Rampen et al. (2008)

with $DI = ([C_{28} \text{ 1,14-diol}] + [C_{30} \text{ 1,14-diol}]) / [C_{30} \text{ 1,15-diol}]$ (3)

The distribution of LCDs in core NIOP905 has been reported before (Rampen et al., 2008) but LCDs were re-analyzed here using an improved GC-MS method (de Bar et al., 2017). Comparisons showed that our results were similar with the previously published values (Supp. Info. Figure S2).

4 Results

We compiled and generated SST records of cores 74KL and NIOP905 for 0-25 kyr and of cores PASOM3 and 289KL 0-35 kyr (Table 1). Except for core PASOM3, we compiled data from previous studies concerning $U^{K'}_{37}$, and the $\delta^{18}O$ and Mg/Ca ratio of the planktonic foraminiferal species *Globigerinoides ruber*. We applied the same calibrations for all proxies and, therefore, re-calculated the $U^{K'}_{37}$ - and $Mg/Ca_{G.ruber}$ -derived SSTs (after Müller et al. (1998) and Anand et al. (2003), respectively) for all records. We generated LDI-derived SST records for all cores as well those obtained by $U^{K'}_{37}$, and $\delta^{18}O$ and Mg/Ca ratio data of *G. ruber* from the PASOM3 core. Based on these records, we estimated average SSTs for the Last Glacial Maximum (LGM; 19 – 22 kyr) and the late Holocene (0 – 5 kyr for NIOP905 and 74KL; 3 – 5 kyr for PASOM3 and 2 – 5 kyr for 289KL) and calculated an approximate SST change ΔSST (LGM-late Holocene) (Table 2). The downcore Mg/Ca ratio based SST estimations of core 74KL is limited to four discrete specific time intervals (Dahl and Oppo, 2006).

The PASOM3 *G. ruber* oxygen isotopic record ($\delta^{18}O_{G.ruber}$) showed changes coinciding with the glacial-interglacial transition with positive values during the glacial and negative values during the Holocene (Figure 2a). The $Mg/Ca_{G.ruber}$ SST in PASOM3 showed fluctuations between 22°C and 27°C, showing substantial differences within relatively short time intervals, particularly during the Glacial, while the Holocene SSTs were relatively stable at this core location. The LGM – late Holocene difference (ΔSST) was 1.2°C at PASOM3 core (Table 2).




Figure 2. SST proxy records (b-d) for four Arabian Sea sediment cores together with oxygen isotope records of *G. ruber* ($\delta^{18}\text{O}_{G. ruber}$) and, for reference, the Greenland ice core (GRIP; $\delta^{18}\text{O}$) (a); see Table 1). Thick lines show the 5-point moving average. The late Holocene (0-5 kyr BP) of core PASOM3 is not well represented due to low sedimentation rates during this period and loss of the core top (first available data point is, therefore, from 3 kyr BP). The present-day mean annual SST in the Arabian Sea is indicated on each scale bar. Specific periods are indicated by grey shaded bars: i.e. Younger Dryas (YD), Heinrich Stadial 1 (H1) and LGM: Last Glacial Maximum

The $U^{K'}_{37}$ SST estimations of PASOM3 showed substantial fluctuations throughout the record. The SSTs during 29-35 kyr BP period were slightly lower than present day whereas during the following period SSTs were variable with minimum temperatures of 20.4°C during the LGM. The maximum estimated SST was 26.8°C during the Holocene (Figure 2c). The LGM-late Holocene $U^{K'}_{37}$ SST difference was 3.4°C (Table 2). This SST record showed continuous warming during the deglaciation and Holocene with interruptions during the Heinrich Stadial 1 (H1) and the Younger Dryas (YD; Figure 2c). The $U^{K'}_{37}$ -derived SST for the core top (ca. 3.1 kyr BP) was 25.8°C.

The LDI SST records of the northern cores, PASOM3 and 289KL, did not reveal major fluctuations during the last 35 kyr, with minimum temperatures of 24.4°C and 24.8°C and maximum temperatures of 26.3°C and 26.6°C, respectively (Figure 2d). The LDI SST records from the cores in the western Arabian Sea, NIOP905 and 74KL, covering the last 25 kyr indicated fluctuating SSTs from the LGM (19 – 22 kyr BP) and onwards with minimum SSTs of 25°C and 21°C, and maximum SSTs of 27°C and 26°C, respectively. Overall, all LDI-derived SST records of NIOP905, 289KL and PASOM3 showed a small decreasing trend from H1 to the late Holocene, with a major change observed at core 74KL. The difference in LDI SSTs from the LGM to the late Holocene (0-5 cal kyr BP) was observed to be -1.6°C at PASOM3, -0.2°C at 289KL, -0.8°C at NIOP905 and -2.7°C at 74KL, respectively (Table 2). All cores revealed higher reconstructed SSTs during the H1 and YD with the largest changes observed at 74KL. Core top (0 kyr BP for NIOP905 and 74KL) LDI-derived SSTs in the western cores were 25.4°C and 22.1°C, respectively, whereas those in the northern cores (ca. 3.1 kyr BP for PASOM3 and ca. 2.2 kyr BP for 289KL) were 24.7°C and 26.1°C, respectively.

Table 2. Modern annual mean SST observed at the site location (Locarnini et al., 2013), SST ranges and SST changes (Δ SST) between the Last Glacial Maximum (19 – 22 kyr) and late Holocene (0 – 5 kyr; ca. 3 – 5 kyr for PASOM3, and 2 – 5 kyr for 289KL) calculated for each core for each proxy. *The values of Δ SST from the Mg/Ca ratio for core 74KL was calculated for time intervals of 24.3-24.5 kyr and 0.5-1.5 kyr after Dahl and Oppo (2006) considering the core depths reported in their study and the new age model reported in this study. Continuous downcore record of 74KL is not available. n.a.: not available.

	Annual mean SST	$U^{K'}_{37}$ (°C)		LDI (°C)		Mg/Ca _{G.ruber} (°C)	
Core Name	modern	Range	Δ SST	Range	Δ SST	Range	Δ SST
NIOP905	26.2	24.8 - 26.3	0.6	25.0 - 26.5	-0.5	22.7 – 25.6	1.7
SO42-74KL	26.5	24.8 – 27.4	2.0	21.1 - 26.0	-2.7	21.4 – 24.4	2.5*
PASOM3	26.7	20.4 - 26.8	3.4	24.4 - 26.3	-1.6	22.3 – 26.8	1.2
SO130-289KL	26.9	n.a.	n.a.	24.8 - 26.6	-0.2	n.a.	n.a.

5 Discussion

5.1 Arabian Sea temperature records using established proxies.

Our newly generated PASOM3 $\delta^{18}\text{O}_{G.ruber}$ record is similar to other $\delta^{18}\text{O}_{G.ruber}$ Arabian Sea records and they all show agreement with the Greenland ice core $\delta^{18}\text{O}$ trend (Johnsen et al., 2001), indicating that these records reflect global climate dynamics (Figure 2). The Mg/Ca ratio-derived SST record from the PASOM3 core shows a larger variability than that of NIOP905 but shows a similar trend from LGM to late Holocene with an approximately 1-2°C warming (Table 2). This is somewhat smaller than the 2.5°C warming reported for the 74KL record (Dahl and Oppo, 2006; Table 2) but it should be noted that that was only determined by a single data point in the LGM. During the Holocene Mg/Ca ratio-derived SSTs of NIOP905 and PASOM3 follow a relatively stable trend with only slight cooling during the late Holocene. Considering the present day mean annual SST of the Arabian Sea (26-27°C; Locarnini et al., 2013) and the calibration error of $\text{Mg}/\text{Ca}_{G.ruber}$ (ca. 1.5°C; Anand et al., 2003) Mg/Ca ratio-derived SST estimates of 24-25 °C from core tops of the western cores (NIOP905 and 74KL) are somewhat low. Based on observations on the modern specimens of *G. ruber* and the Mg/Ca based - SST calibrations in the Arabian Sea (Conan and Brummer, 2000; Curry et al., 1992) SST estimates were observed to be $1.7 \pm 0.8^\circ\text{C}$ lower than annual mean SST (Peeters et al., 2002). This is potentially due to the present-day habitat of *G. ruber* in the Arabian Sea, where this species was observed living year around, calcifying in the uppermost part of the water column and with blooms during both monsoon seasons in relatively cold surface waters (Curry et al., 1992; Dahl and Oppo, 2006).

The newly obtained $\text{U}^{\text{K}'}_{37}$ -derived SST record of PASOM3 shows a high variability but the 5-points running mean shows a similar trend as other previously published $\text{U}^{\text{K}'}_{37}$ -derived SST records and follows the $\delta^{18}\text{O}_{G.ruber}$ as well as the global climate trend with lower SST values during the LGM and increasing towards the Holocene (Figure 2). However, $\text{U}^{\text{K}'}_{37}$ -derived SSTs for the LGM were substantially lower in our PASOM3 record from the Northern Arabian Sea compared to other two $\text{U}^{\text{K}'}_{37}$ -derived SST records from the western part and showed a much larger LGM to Holocene change (i.e. 3.4°C; Table 2). This is in accordance with previous observations based on $\text{U}^{\text{K}'}_{37}$ SSTs from the region, suggesting such a north – west difference in deglacial warming (e.g., Böll et al., 2015). This regional difference was interpreted as a result of reduced impact of NE monsoon related cooling and/or strengthening SW monsoon induced upwelling in the region after the Glacial (Dahl and Oppo, 2006; Emeis et al., 1995; Gaye et al., 2018; Rostek et al., 1997). With the onset of the deglaciation, the SW monsoon and upwelling in the western Arabian Sea intensified and resulted in cooler SSTs in comparison with the northern Arabian Sea (e.g., Anand et al., 2008; Böll et al., 2015; Gaye et al., 2018). However, the $\text{U}^{\text{K}'}_{37}$ -derived SST record of NIOP905 shows even cooler surface water conditions during the Holocene in comparison with those of the other western Arabian Sea core 74KL (Figure 2c). This is potentially due to a local feature and circulation pattern observed in the region (Figure 1b;

e.g., Sonzogni et al., 1998). Right above the core site, the so-called *Great Whirl* develops and remains stationary during the SW Monsoon, which has a major effect on the local SST and surface productivity (Fischer et al., 1996; Koning et al., 2001; Peeters et al., 1999). This local feature is potentially reflected in $U^{K'}_{37}$ SSTs from core NIOP905, particularly during the Holocene part of the record.

Overall, the SST records based on these proxies reveal relatively small SST change (1-2°C) from LGM to the Holocene, with the exception of PASOM3 (3.4°C). Besides the above-discussed reasons for specific proxy and core site, this is a well-known observation for SST reconstructions in the tropical regions (e.g., Mix et al., 2001; CLIMAP Project Members, 1981; Rosell-Melé et al., 2004; Waelbroeck et al., 2009). Basin-wide proxy applications (Dahl and Oppo, 2006; Rosell-Melé et al., 2004; Sonzogni et al., 1998) as well as model simulations (Mix et al., 2001; Otto-Bliesner et al., 2009; Waelbroeck et al., 2009) showed that the Arabian Sea SST warming from the LGM to the late Holocene ranged between 1-3.5°C, with higher amplitude in the north. The new Mg/Ca ratio and $U^{K'}_{37}$ results from core PASOM3 are also in line with previous observations and model simulations.

5.2 Arabian Sea temperature records using the LDI.

The overall trend observed for the LDI-derived SST records are consistent in three of the four records (i.e. excepting the 75KL record) but surprisingly only reveal minor changes from glacial to the Holocene (Figure 2d; Table 2) and a slight cooling trend during the Holocene (except for the 289KL record). The small Holocene to LGM SST change and cooling during the Holocene in the LDI records show similarities to the CLIMAP Project data based on planktonic foraminifera SST approaches (modern analogue technique and transfer function), which revealed relatively low LGM-Holocene warming or even cooling at sites in the Arabian Sea between 10-20°N interval (CLIMAP Project, 1981; Sonzogni et al., 1998). The LDI record of core 74KL is clearly an exception to the other three (Figure 2d) in that it shows 1.5°C lower glacial SSTs prior to H1 as well as a strong continuous decline in SST during the Holocene, arriving at SSTs of ca. 22°C in the upper Holocene. This clearly contrasts most other proxy records discussed above, which show some degree of LGM to Holocene warming and is incompatible with the global climate trend (Figure 2; Table 2). Interestingly, SSTs estimated from planktonic foraminifera-based transfer functions for core 74KL (Schulz, 1995a, 1995b) also show decreasing values for the Holocene and suggest cooler summer (SW monsoon) SSTs after the Heinrich Stadial 1 (H1) and, in this way, are partly in agreement with the LDI-derived SST trend in the Holocene (Figure 3a). The decreasing trend of LDI SST observed in most of our records during the Holocene may be a direct effect on SST due to intensified upwelling (e.g., Gaye et al., 2018; Marcantonio et al., 2001; Rampen et al., 2008; Sirocko et al., 1996) leading to a summer monsoon (i.e. cold) bias in LDI SSTs. Therefore, the LDI is potentially reflecting SW summer monsoon SSTs rather than annual mean temperatures in the Arabian Sea, particularly after the H1 event. Indeed, the deglacial decrease in LDI-based SSTs is lowest at core site 289KL (-0.2°C; Table 2), which is

also the furthest from the modern upwelling cell (Figure 1). Nevertheless, we cannot exclude that other factors not related to SST have impacted the LDI.




Figure 3. Compilation of various records from core 74KL; a) SST reconstructions based on LDI and planktonic foraminifera transfer functions for different seasons (light blue: winter, NE monsoon; brown: summer, SW monsoon; (Schulz, 1995b), b) Diol upwelling index (DI) and $C_{28,1,12}$ -diol factor showing potential influence of diatom productivity on the LDI. Largest convergence between the LDI and the planktonic foraminifera transfer function SSTs coincide with periods characterized by reduced upwelling. LDI SSTs show continuous decrease where DI and 1,12-diols increase.

Sedimentary oxic degradation has been shown to impact LDI values in Arabian Sea surface sediments due to different degradation rates of 1,13- and 1,15-diols leading to ca. 1.5-2.5°C cooler LDI-reconstructed SSTs in sediments deposited below the OMZ (Rodrigo-Gámiz et al., 2016). Enhanced bottom water ventilation and a weaker OMZ has been inferred for cold periods (e.g., Böning and Bard, 2009; Reichert et al., 1997; Reichert et al., 1998; Schulte et al., 1999), presumably leading to enhanced oxygen exposure times (OETs). This may have led to lower LDI SST estimates during these colder times for the cores obtained from stations located within the present-day OMZ (NIOP905, PASOM3 and 289KL; Figure 1). In contrast, our results actually indicate relatively warmer conditions, particularly during YD and H1, and cooler SSTs

during the Holocene (Figure 2). This suggests that the oxic degradation of the LCDs likely did not significantly affect these LDI records. Being by far the deepest core site, 74KL has always been below the OMZ (Figure 1) and the sediments deposited probably have been impacted by much longer oxygen exposure times (OET) than the other sedimentary records studied. These enhanced OET may have led to an overall cold bias compared to other cores, especially during the Holocene when other sites may have been within the OMZ and therefore experienced lower OET (Figure 2d).

A cold bias in SSTs reconstructed with the LDI could also occur when groups of organisms, such as *Proboscia* diatoms, produce, in addition to their characteristic 1,14-diols, also 1,13- and 1,15-diols as has been observed around Iceland (Rodrigo-Gámiz et al., 2015). de Bar et al. (2020) reported that such a bias in LDI could be assessed by the relative abundance of 1,12-diols, produced by certain *Proboscia* diatoms, expressed in the FC_{28} 1,12. However, in all of our records, FC_{28} 1,12 remained below the threshold (<0.1), although Holocene values at core 74KL were getting close to this value (Figure 3). Nevertheless, the Diol Index (DI), a proxy for upwelling intensity and enhanced nutrient concentrations, does show increasing values in all sites after the last deglaciation (Figures 3 and 4), as previously reported for core NIOP905 (Rampen et al., 2008), suggesting an overall increase in *Proboscia* diatom contributions. However, except for core 74KL, the LDI SST estimates during the Holocene fit well with other temperature proxies as well as modern day SST (Figure 2 and Figure 3), despite having relatively high DI values. Hence, these changes in upwelling and concomitantly increasing nutrient availability cannot explain the relatively high LDI-reconstructed SSTs during the glacial.

As discussed, LDI-reconstructed SSTs for core 74KL remain very unusual compared to the other cores as they show the highest SSTs during cold intervals, the YD and H1, and a continuously decreasing trend during the Holocene with the largest offset from the LGM to the Holocene ($\Delta SST = -2.7^{\circ}C$; Table 2). Core 74KL was collected in an area where the SW monsoon and related upwelling are much more pronounced today in comparison to the western part of the Arabian Sea. Indeed, we observe strong correlation between the DI and LDI ($r^2=0.72$) at core 74KL. Moreover, the OMZ is much stronger within the overlying waters of the 74KL core site, in comparison to the other western Arabian Sea core NIOP905 (Figure 1; Garcia et al., 2013). These local differences suggest that the extent of the OMZ in combination with enhanced upwelling might have a pronounced impact on the LDI at the 74KL location.




Figure 4: Records for the Diol upwelling index (DI) for the four stations considered in this study.

Finally, the strong enhanced upwelling and expanded OMZ at 74KL may also impact the diol producers, which likely live in the photic zone (Rampen et al., 2012). Studies show that with enhanced upwelling, surface productivity and an extended OMZ, there is enhanced nitrogen loss (denitrification) in subsurface waters (Galloway et al., 2004). As a result of intensified upwelling, oxygen and nitrate limited (denitrified) waters are brought to surface waters resulting in enhanced N_2 -fixation (Bange et al., 2005; Deutsch et al., 2007; White et al., 2013). This will cause changes within the plankton community with a relative increase being expected in the abundance of N_2 fixing organisms such as cyanobacteria or cyanobacterial symbionts of diatoms (Zehr and Capone, 2020; and the references therein), due to changes in nutrient dynamics in the upwelled waters (such as N:P ratio) (Bange et al., 2005; White et al., 2013). It is however unclear how and what extent this will affect the LCD producers and their diol composition. In an early study focusing on Baltic Sea, detection of LCDs during cyanobacterial bloom were reported suggesting some link with enhanced N_2 fixation (Morris and Brassell, 1988), although it is clear that they are not derived from the cyanobacteria as these compounds are not detected in cyanobacterial cultures (e.g., de Leeuw et al., 1992). Further research investigating the link between nutrient dynamics and diol production as well as composition of diol producers is needed to clarify this.

5 Conclusions

A multi proxy application in series of cores from the western and northeastern part of the Arabian Sea emphasize the importance of application of different proxies and on several records to reconstruct regional climate dynamics. Our results show that LDI is likely to reflect in part summer monsoonal temperatures showing little change in SST from glacial to the Holocene and is not an ideally suited proxy to reconstruct annual mean SST in the Arabian Sea. The index may

also be potentially biased by upwelling intensity and related nutrient input in regions of strong upwelling and extended OMZs. Results from LDI in upwelling regions should therefore be interpreted carefully. However, other SST proxies such as $U^{K'}_{37}$ and $Mg/Ca_{G.ruber}$ show also variable trends though they mostly document a relatively consistent overall SST increase from the Last Glacial Maximum (LGM) to the late Holocene of 1-2°C, with a larger amplitude in the northern Arabian Sea in comparison with the west.

Acknowledgments, Samples, and Data

We would like to thank the crew and scientists aboard *R/V Pelagia* during PASOM expedition 64PE301 and Rick Hennekam for his support with the XRF data analysis. This research has been funded by the European Research Council (ERC) under the European Union's Seventh Framework Program (FP7/2007-2013) ERC grant agreement [339206] to S.S. S.S and by financial support from the Netherlands Earth System Science Centre (NESSC) through a gravitation grant from the Dutch Ministry for Education, Culture and Science to GJR, JSSD and SS.

Datasets for this research are available in these in-text data citation references: Schulz, H.(1995): <https://doi.pangaea.de/10.1594/PANGAEA.134165>, Hugué et al., (2006): <https://doi.org/10.1594/PANGAEA.861247> and <https://doi.org/10.1594/PANGAEA.861245>, Dahl and Oppo (2006): <https://doi.org/10.1594/PANGAEA.834987>. New datasets are included in this paper's supplementary information files. They were also recently submitted to the same platform (www.pangaea.de) and will be linked to existing datasets once they are publicly available.

References

- Acharya, S. S., & Panigrahi, M. K. (2016). Eastward shift and maintenance of Arabian Sea oxygen minimum zone: Understanding the paradox. *Deep Sea Research Part I: Oceanographic Research Papers*, 115, 240-252.
- Altabet, M. A., Higginson, M. J., & Murray, D. W. (2002). The effect of millennial-scale changes in Arabian Sea denitrification on atmospheric CO₂. *Nature*, 415, 159-162.
- Anand, P., Elderfield, H., & Conte, M. H. (2003). Calibration of Mg/Ca thermometry in planktonic foraminifera from a sediment trap time series. *Paleoceanography*, 18(2), doi:10.1029/2002pa000846
- Anand, P., Kroon, D., Singh, A. D., Ganeshram, R. S., Ganssen, G., & Elderfield, H. (2008). Coupled sea surface temperature-seawater $\delta^{18}\text{O}$ reconstructions in the Arabian Sea at the millennial scale for the last 35 ka. *Paleoceanography*, 23(4), n/a-n/a. doi:10.1029/2007pa001564
- Andersen, K. K., Azuma, N., Barnola, J.-M., Bigler, M., Biscaye, P., Caillon, N., . . . White, J. W. C. (2004). High-resolution record of Northern Hemisphere climate extending into the last interglacial period. *Nature*, 431(7005), 147-151.
- Balzano, S., Lattaud, J., Villanueva, L., Rampen, S. W., Brussaard, C. P., Van Bleijswijk, J., . . . Schouten, S. (2018). A quest for the biological sources of long chain alkyl diols in the western tropical North Atlantic Ocean. *Biogeosciences*, 15(19), 5951-5968.
- Bange, H. W., Naqvi, S. W. A., & Codispoti, L. A. (2005). The nitrogen cycle in the Arabian Sea. *Progress in Oceanography*, 65(2-4), 145-158.
- Barker, S., Greaves, M., & Elderfield, H. (2003). A study of cleaning procedures used for foraminiferal Mg/Ca paleothermometry. *Geochemistry, Geophysics, Geosystems*, 4(9).
- Böll, A., Schulz, H., Munz, P., Rixen, T., Gaye, B., & Emeis, K.-C. (2015). Contrasting sea surface temperature of summer and winter monsoon variability in the northern Arabian Sea over the last 25 ka. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 426, 10-21.
- Böning, P., & Bard, E. (2009). Millennial/centennial-scale thermocline ventilation changes in the Indian Ocean as reflected by aragonite preservation and geochemical variations in Arabian Sea sediments. *Geochimica Et Cosmochimica Acta*, 73(22), 6771-6788.
- Brassell, S. C., Eglinton, G., Marlowe, I. T., Pflaumann, U., & Sarnthein, M. (1986). Molecular stratigraphy: A new tool for climatic assessment. *Nature*, 320(6058), 129-133.
- Clemens, S. C., & Prell, W. L. (2003). A 350,000 year summer-monsoon multi-proxy stack from the Owen ridge, Northern Arabian sea. *Marine Geology*, 201(1-3), 35-51.
- Conan, S. M.-H., & Brummer, G.-J. A. (2000). Fluxes of planktic foraminifera in response to monsoonal upwelling on the Somalia Basin margin. *Deep-Sea Research Part II-Topical Studies in Oceanography*, 47(9-11), 2207-2227. Retrieved from <Go to ISI>://000087495300023
- Curry, W. B., Ostermann, D. R., Guptha, M. V. S., & Ittekkot, V. (1992). Foraminiferal production and monsoonal upwelling in the Arabian Sea: evidence from sediment traps. In C. P. Summerhayes, W. L. Prell, & K. C. Emeis (Eds.), *Upwelling systems: evolution since the early Miocene* (pp. 93-106). London: The Geological Society.
- Dahl, K. A., & Oppo, D. W. (2006). Sea surface temperature pattern reconstructions in the Arabian Sea. *Paleoceanography*, 21(1), PA1014-1011-PA1014-1016.
- Dauner, A. L. L., Mollenhauer, G., Bicego, M. C., de Souza, M. M., Nagai, R. H., Figueira, R. C. L., . . . Martins, C. C. (2019). Multi-proxy reconstruction of sea surface and subsurface temperatures in the western South Atlantic over the last ~75 kyr. *Quaternary Science Reviews*, 215, 22-34. doi:10.1016/j.quascirev.2019.04.020
- de Bar, M. W., Hopmans, E. C., Verweij, M., Dorhout, D. J. C., Damste, J. S. S., & Schouten, S. (2017). Development and comparison of chromatographic methods for the analysis of long chain diols and alkenones in biological materials and sediment. *J Chromatogr A*, 1521, 150-160. doi:10.1016/j.chroma.2017.09.037
- de Bar, M. W., Stolwijk, D. J., McManus, J. F., Sinninghe Damste, J. S., & Schouten, S. (2018). A Late Quaternary climate record based on long-chain diol proxies from the Chilean margin. *Climate of the Past*, 14, 1783-1803. doi:10.5194/cp-14-1783-2018
- de Bar, M. W., Weiss, G., Rampen, S., Lattaud, J., Bale, N. J., Brummer, G.-J. A., . . . Donner, B. (2020). Global temperature calibration of the Long chain Diol Index in marine surface sediments. *Organic Geochemistry*, 103983.
- de Leeuw, J. W., Rijpstra, W. I. C., & Mur, L. R. (1992). The absence of long-chain alkyl diols and alkyl keto-1-ols in cultures of the cyanobacterium *Aphanizomenon flos-aquae*. *Org. Geochem*, 18(4), 575-578.

- Deplazes, G., Lückge, A., Peterson, L. C., Timmermann, A., Hamann, Y., Hughen, K. A., . . . Haug, G. H. (2013). Links between tropical rainfall and North Atlantic climate during the last glacial period. *Nature Geoscience*, 6(3), 213-217. doi:10.1038/ngeo1712
- Deplazes, G., Lückge, A., Stuut, J.-B. W., Pätzold, J., Kuhlmann, H., Husson, D., . . . Haug, G. H. (2014). Weakening and strengthening of the Indian monsoon during Heinrich events and Dansgaard-Oeschger oscillations. *Paleoceanography*, 29(2), 99-114. doi:10.1002/2013pa002509
- Deutsch, C., Sarmiento, J. L., Sigman, D. M., Gruber, N., & Dunne, J. P. (2007). Spatial coupling of nitrogen inputs and losses in the ocean. *Nature*, 445(7124), 163-167.
- Doose-Rolinski, H., Rogalla, U., Scheeder, G., Lückge, A., & von Rad, U. (2001). High-resolution temperature and evaporation changes during the late Holocene in the northeastern Arabian Sea. *Paleoceanography*, 16(4), 358-367.
- Elderfield, H., & Ganssen, G. (2000). Past temperature and $\delta^{18}\text{O}$ of surface ocean waters inferred from foraminiferal Mg/Ca ratios. *Nature*, 405(6785), 442-445.
- Emeis, K.-C., Anderson, D. M., Doose, H., Kroon, D., & Schulz, D. (1995). Sea-Surface temperatures and the history of monsoon upwelling in the Northwest Arabian Sea during the last 500,000 years. *Quaternary Research*, 43(3), 355-361.
- Erez, J., & Luz, B. (1983). Experimental paleotemperature equation for planktonic-foraminifera. *Geochimica Et Cosmochimica Acta*, 47(6), 1025-1031.
- Fischer, J., Schott, F., & Stramma, L. (1996). Currents and transports of the Great Whirl-Socotra Gyre system during the summer monsoon, August 1993. *Journal of Geophysical Research: Oceans*, 101(C2), 3573-3587.
- Garcia, H. E., Boyer, T. P., Locarnini, R. A., Antonov, J. I., Mishonov, A. V., Baranova, O. K., . . . Levitus, S. (2013). World ocean atlas 2013. Volume 3, Dissolved oxygen, apparent oxygen utilization, and oxygen saturation.
- Gaye-Haake, B., Lahajnar, N., Emeis, K.-C., Unger, D., Rixen, T., Suthhof, A., . . . Guptha, M. (2005). Stable nitrogen isotopic ratios of sinking particles and sediments from the northern Indian Ocean. *Marine Chemistry*, 96(3-4), 243-255.
- Gaye, B., Böll, A., Segschneider, J., Burdanowitz, N., Emeis, K.-C., Ramaswamy, V., . . . Rixen, T. (2018). Glacial–interglacial changes and Holocene variations in Arabian Sea denitrification. *Biogeosciences*, 15(2), 507-527. doi:10.5194/bg-15-507-2018
- Haake, B., Ittekkot, V., Rixen, T., Ramaswamy, V., Nair, R. R., & Curry, W. B. (1993). Seasonality and interannual variability of particle fluxes to the deep Arabian Sea. *Deep-Sea Res. I*, 40(7), 1323-1344.
- Häggi, C., Schefuß, E., Sawakuchi, A. O., Chiessi, C. M., Mulitza, S., Bertassoli, D. J., . . . Schouten, S. (2019). Modern and late Pleistocene particulate organic carbon transport by the Amazon River: Insights from long-chain alkyl diols. *Geochimica Et Cosmochimica Acta*, 262, 1-19. doi:10.1016/j.gca.2019.07.018
- Helly, J. J., & Levin, L. A. (2004). Global distribution of naturally occurring marine hypoxia on continental margins. *Deep Sea Research Part I: Oceanographic Research Papers*, 51(9), 1159-1168.
- Honjo, S., Dymond, J., Prell, W., & Ittekkot, V. (1999). Monsoon-controlled export fluxes to the interior of the Arabian Sea. *Deep-Sea Research Part II*, 46(8-9), 1859-1902.
- Huguet, C., Kim, J. H., Sinninghe Damste, J. S., & Schouten, S. (2006). Reconstruction of sea surface temperature variations in the Arabian Sea over the last 23 kyr using organic proxies (TEX86 and U37K'). *Paleoceanography*, 21(3).
- Imbrie, J., & Kipp, N. G. (1971). A new micropaleontological method for quantitative paleoclimatology: application to a late Pleistocene Caribbean core In K. K. Turekian (Ed.), *The Late Cenozoic Glacial Ages* (pp. 71-181). New Haven: Yale University Press.
- Ivanochko, T. S., Ganeshram, R. S., Brummer, G.-J. A., Ganssen, G., Jung, S. J. A., Moreton, S. G., & Kroon, D. (2005). Variations in tropical convection as an amplifier of global climate change at the millennial scale. *Earth and Planetary Science Letters*, 235(1-2), 302-314.
- Ivanova, E. M., Conan, S. M.-H., Peeters, F. J. C., & Troelstra, S. R. (1999). Living *Neogloboquadrina pachyderma* sin and its distribution in the sediments from Oman and Somalia upwelling areas. *Marine Micropaleontology*, 36(2-3), 91-107.
- Johnsen, S. J., Dahl-Jensen, D., Gundestrup, N., Steffensen, J. P., Clausen, H. B., Miller, H., . . . White, J. (2001). Oxygen isotope and palaeotemperature records from six Greenland ice-core stations: Camp Century, Dye-3, GRIP, GISP2, Renland and NorthGRIP. *Journal of Quaternary Science: Published for the Quaternary Research Association*, 16(4), 299-307.

- Jonas, A. S., Schwark, L., & Bauersachs, T. (2017). Late Quaternary water temperature variations of the Northwest Pacific based on the lipid paleothermometers TEX_{H86}, U_{K'37} and LDI. *Deep Sea Research Part I: Oceanographic Research Papers*, 125, 81-93. doi:10.1016/j.dsr.2017.04.018
- Jung, S. J. A., Ganssen, G. M., & Davies, G. R. (2001). Multidecadal variation in the early Holocene outflow of Red Sea Water into the Arabian Sea. *Paleoceanography*, 16(6), 658-668.
- Jung, S. J. A., Davies, G. R., Ganssen, G., & Kroon, D. (2002). Decadal-centennial scale monsoon variations in the Arabian Sea during the Early Holocene. *Geochemistry Geophysics Geosystems*, 3.
- Kim, J.-H., Rimbu, N., Lorenz, S. J., Lohmann, G., Nam, S. I., Schouten, S., . . . Schneider, R. R. (2004). North Pacific and North Atlantic sea-surface temperature variability during the Holocene. *Quaternary Science Reviews*, 23(20-22), 2141-2154.
- Koning, E., van Iperen, J. M., van Raaphorst, W., Helder, W., Brummer, G.-J. A., & van Weering, T. C. E. (2001). Selective preservation of upwelling-indicating diatoms in sediments off Somalia, NW Indian Ocean. *Deep-Sea Research I*, 48, 2473-2495.
- Kucera, M., Rosell-Melé, A., Schneider, R., Waelbroeck, C., & Weinelt, M. (2005). Multiproxy approach for the reconstruction of the glacial ocean surface (MARGO). *Quaternary Science Reviews*, 24(7-9), 813-819.
- Lattaud, J., Lo, L., Huang, J. J., Chou, Y. M., Gorbarenko, S. A., Sinninghe Damsté, J. S., & Schouten, S. (2018). A Comparison of Late Quaternary Organic Proxy-Based Paleotemperature Records of the Central Sea of Okhotsk. *Paleoceanography and Paleoclimatology*.
- Locarnini, R. A., Mishonov, A. V., Antonov, J. I., Boyer, T. P., Garcia, H. E., Baranova, O. K., . . . Seidov, D. (2013). World Ocean Atlas 2013 Volume 1: Temperature. In S. Levitus & A. Mishonov (Eds.), *NOAA Atlas NESDIS 73* (Vol. 73, pp. 40).
- Lopes dos Santos, R. A., Spooner, M. I., Barrows, T. T., De Deckker, P., Sinninghe Damsté, J. S., & Schouten, S. (2013). Comparison of organic (U^{K'37}, TEX^{H86}, LDI) and faunal proxies (foraminiferal assemblages) for reconstruction of late Quaternary sea surface temperature variability from offshore southeastern Australia. *Paleoceanography*, n/a-n/a. Retrieved from <http://dx.doi.org/10.1002/palo.20035>
- Marcantonio, F., Anderson, R. F., Higgins, S., Fleisher, M. Q., Stute, M., & Schlosser, P. (2001). Abrupt intensification of the SW Indian Ocean monsoon during the last deglaciation: constraints from Th, Pa, and He isotopes. *Earth and Planetary Science Letters*, 184(2), 505-514.
- Mezger, E. M. (2019). Controls on sodium incorporation in foraminiferal calcite. Utrecht Studies in Earth Sciences, 182. PhD Thesis. Utrecht University: Utrecht. ISBN 9789062665365. 241 pp. <https://hdl.handle.net/1874/380001>
- Mezger, E., Nooijer, L., Boer, W., Brummer, G., & Reichert, G. (2016). Salinity controls on Na incorporation in Red Sea planktonic foraminifera. *Paleoceanography and Paleoclimatology*, 31(12), 1562-1582.
- Mix, A. C., Bard, E., & Schneider, R. (2001). Environmental processes of the ice age: land, oceans, glaciers (EPILOG). *Quaternary Science Reviews*, 20(4), 627-657.
- Morris, R. J., & Brassell, S. C. (1988). Long-chain alkanediols: biological markers for cyanobacterial contributions to sediments. *Lipids*, 23(3), 256-258.
- Müller, P. J., Kirst, G., Ruhland, G., von Storch, I., & Rosell-Melé, A. (1998). Calibration of the alkenone paleotemperature index U^{K'37} based on core-tops from the eastern South Atlantic and the global ocean (60°N-60°S). *Geochimica Et Cosmochimica Acta*, 62(10), 1757-1772.
- Naqvi, W. A. (1991). Geographical extent of denitrification in the Arabian Sea in relation to some physical processes. *Oceanologica Acta*, 14(3), 281-290. Retrieved from <Go to ISI>://A1991FW22400006
- Nürnberg, D., Bijma, J., & Hemleben, C. (1996). Assessing the reliability of magnesium in foraminiferal calcite as a proxy for water mass temperatures. *Geochimica Et Cosmochimica Acta*, 60(5), 803-814. Retrieved from <Go to ISI>://A1996TY43300005
- Okai, T., Suzuki, A., Kawahata, H., Terashima, S., & Imai, N. (2002). Preparation of a new Geological Survey of Japan geochemical reference material: Coral JcP-1. *Geostandards newsletter*, 26(1), 95-99.
- Olson, D. B., Hitchcock, G. L., Fine, R. A., & Warren, B. A. (1993). Maintenance of the low-oxygen layer in the central Arabian Sea. *Deep-Sea Research Part II-Topical Studies in Oceanography*, 40(3), 673-685. Retrieved from <Go to ISI>://A1993LF20700003
- Otto-Bliesner, B. L., Schneider, R., Brady, E., Kucera, M., Abe-Ouchi, A., Bard, E., . . . Kageyama, M. (2009). A comparison of PMIP2 model simulations and the MARGO proxy reconstruction for tropical sea surface temperatures at last glacial maximum. *Climate Dynamics*, 32(6), 799-815.
- Paulmier, A., & Ruiz-Pino, D. (2009). Oxygen minimum zones (OMZs) in the modern ocean. *Progress in Oceanography*, 80(3-4), 113-128.

- Peeters, F., Ivanova, E., Conan, S., Brummer, G.-J., Ganssen, G., Troelstra, S., & van Hinte, J. (1999). A size analysis of planktic foraminifera from the Arabian Sea. *Marine Micropaleontology*, 36(1), 31-63.
- Pflaumann, U., Duprat, J., Pujol, C., & Labeyrie, L. D. (1996). SIMMAX: A modern analog technique to deduce Atlantic sea surface temperatures from planktonic foraminifera in deep-sea sediments. *Paleoceanography*, 11(1), 15-35.
- Pichevin, L., Bard, E., Martinez, P., & Billy, I. (2007). Evidence of ventilation changes in the Arabian Sea during the late Quaternary: Implication for denitrification and nitrous oxide emission. *Global Biogeochemical Cycles*, 21(4).
- Prahl, F. G., & Wakeham, S. G. (1987). Calibration of unsaturation patterns in long-chain ketone compositions for palaeotemperature assessment. *Nature*, 330, 367-369.
- Prasad, T., Ikeda, M., & Kumar, S. P. (2001). Seasonal spreading of the Persian Gulf Water mass in the Arabian Sea. *Journal of Geophysical Research: Oceans*, 106(C8), 17059-17071.
- Project, C. (1981). *Seasonal reconstructions of the Earth's surface at the last glacial maximum*: Geological Society of America.
- Qasim, S. Z. (1982). Oceanography of the Northern Arabian Sea. *Deep-Sea Research Part A-Oceanographic Research Papers*, 29(9A), 1041-1068.
- Rampen, S. W., Schouten, S., Koning, E., Brummer, G.-J. A., & Damsté, J. S. S. (2008). A 90 kyr upwelling record from the northwestern Indian Ocean using a novel long-chain diol index. *Earth and Planetary Science Letters*, 276(1), 207-213.
- Rampen, S. W., Willmott, V., Kim, J.-H., Rodrigo-Gámiz, M., Uliana, E., Mollenhauer, G., . . . Schouten, S. (2014). Evaluation of long chain 1, 14-alkyl diols in marine sediments as indicators for upwelling and temperature. *Organic Geochemistry*, 76, 39-47.
- Rampen, S. W., Willmott, V., Kim, J.-H., Uliana, E., Mollenhauer, G., Schefuß, E., . . . Schouten, S. (2012). Long chain 1, 13-and 1, 15-diols as a potential proxy for palaeotemperature reconstruction. *Geochimica Et Cosmochimica Acta*, 84, 204-216.
- Reichart, G. J., den Dulk, M., Visser, H. J., van der Weijden, C. H., & Zachariasse, W. J. (1997). A 225 kyr record of dust supply, paleoproductivity and the oxygen minimum zone from the Murray ridge (northern Arabian sea). *Palaeogeography Palaeoclimatology Palaeoecology*, 134(1-4), 149-169.
- Reichart, G. J., Lourens, L. J., & Zachariasse, W. J. (1998). Temporal variability in the northern Arabian Sea Oxygen Minimum Zone (OMZ) during the last 225,000 years. *Paleoceanography*, 13(6), 607-621.
- Reichart, G. J., Nortier, J., Versteegh, G., & Zachariasse, W. J. (2002). Periodical breakdown of the Arabian Sea oxygen minimum zone caused by deep convective mixing. In P. D. Clift, D. Kroon, C. Gaedicke, & J. Craig (Eds.), *The tectonic and climatic evolution of the Arabian Sea region* (pp. 407-419). London: The Geological Society.
- Reichart, G. J., Brinkhuis, H., Huiskamp, F., & Zachariasse, W. J. (2004). Hyperstratification following glacial overturning events in the northern Arabian Sea. *Paleoceanography*, 19(2).
- Reimer, P. J., Bard, E., Bayliss, A., Beck, J. W., Blackwell, P. G., Ramsey, C. B., . . . Friedrich, M. (2013). IntCal13 and Marine13 radiocarbon age calibration curves 0–50,000 years cal BP. *Radiocarbon*, 55(4), 1869-1887.
- Rixen, T., Baum, A., Gaye, B., & Nagel, B. (2014). Seasonal and interannual variations in the nitrogen cycle in the Arabian Sea. *Biogeosciences*, 11(20), 5733.
- Rixen, T., Ittekkot, V., Haake-Gaye, B., & Schäfer, P. (2000). The influence of the SW monsoon on the deep-sea organic carbon cycle in the Holocene. *Deep-Sea Research Part II-Topical Studies in Oceanography*, 47(14), 2629-2651.
- Rodrigo-Gámiz, M., Rampen, S. W., de Haas, H., Baas, M., Schouten, S., & Sinninghe Damsté, J. S. (2015). Constraints on the applicability of the organic temperature proxies UK'37, TEX86 and LDI in the subpolar region around Iceland. *Biogeosciences*, 12(22), 6573-6590. doi:10.5194/bg-12-6573-2015
- Rodrigo-Gámiz, M., Rampen, S. W., Schouten, S., & Sinninghe Damsté, J. S. (2016). The impact of oxic degradation on long chain alkyl diol distributions in Arabian Sea surface sediments. *Organic Geochemistry*, 100, 1-9. doi:<http://dx.doi.org/10.1016/j.orggeochem.2016.07.003>
- Rosell-Melé, A., Bard, E., Emeis, K.-C., Grieger, B., Hewitt, C., Müller, P. J., & Schneider, R. R. (2004). Sea surface temperature anomalies in the oceans at the LGM estimated from the alkenone- $U^{K'}_{37}$ index: comparison with GCMs. *Geophysical Research Letters*, 31(3), L03208-03201-L03208-03204.
- Rostek, F., Bard, E., Beaufort, L., Sonzogni, C., & Ganssen, G. (1997). Sea surface temperature and productivity records for the past 240 kyr in the Arabian Sea. *Deep-Sea Research II*, 44(6-7), 1461-1480.
- Sarma, V. (2002). An evaluation of physical and biogeochemical processes regulating perennial suboxic conditions in the water column of the Arabian Sea. *Global Biogeochemical Cycles*, 16(4).

- Schlitzer, R. (2015). Ocean data view, 513-546.
- Schmidt, H., Czeschel, R., & Visbeck, M. (2020). Seasonal variability of the Arabian Sea intermediate circulation and its impact on seasonal changes of the upper oxygen minimum zone. *Ocean Science*, 16(6), 1459-1474.
- Schulte, S., & Müller, P. J. (2001). Variations of sea surface temperature and primary productivity during Heinrich and Dansgaard-Oeschger events in the northeastern Arabian Sea. *Geo-Marine Letters*, 21, 168-175.
- Schulte, S., Rostek, F., Bard, E., Rullkötter, J., & Marchal, O. (1999). Variations of oxygen-minimum and primary productivity recorded in sediments of the Arabian Sea. *Earth and Planetary Science Letters*, 173(3), 205-221.
- Schulz, H. (1995a). Meeresoberflächentemperaturen vor 10.000 Jahren-Auswirkungen des frühholozänen Insolationsmaximums. *Berichte-Reports, Geologisch-Paläontologisches Institut und Museum, Christian-Albrechts-Universität, Kiel*, 73, 119.
- Schulz, H. (1995b). *Sea surface temperatures calculated for sediment core SO42-74KL*. <https://doi.org/10.1594/PANGAEA.51953>
- Schulz, H., von Rad, U., & Erlenkeuser, H. (1998). Correlation between Arabian Sea and Greenland climate oscillations of the past 110,000 years. *Nature*, 393(6680), 54-57.
- Shakun, J. D., Clark, P. U., He, F., Marcott, S. A., Mix, A. C., Liu, Z., . . . Bard, E. (2012). Global warming preceded by increasing carbon dioxide concentrations during the last deglaciation. *Nature*, 484, 49-55.
- Shankar, D., Shenoi, S., Nayak, R., Vinayachandran, P., Nampoothiri, G., Almeida, A., . . . Sreejith, O. (2005). Hydrography of the eastern Arabian Sea during summer monsoon 2002. *Journal of earth system science*, 114(5), 459-474.
- Singh, A. D., Jung, S. J., Darling, K., Ganeshram, R., Ivanochko, T., & Kroon, D. (2011). Productivity collapses in the Arabian Sea during glacial cold phases. *Paleoceanography*, 26(3).
- Sirocko, F., Garbe Schönberg, D., McIntyre, A., & Molfino, B. (1996). Teleconnections between the subtropical monsoons and high-latitude climates during the last deglaciation. *Science*, 272(5261), 526-529.
- Sirocko, F., Sarnthein, M., Erlenkeuser, H., Lange, H., Arnold, M., & Duplessy, J. C. (1993). Century-scale events in monsoonal climate over the past 24,000 years. *Nature*, 364(6435), 322-324.
- Smith, S. L., Codispoti, L. A., Morrison, J. M., & Barber, R. T. (1998). The 1994-1996 Arabian Sea Expedition: An integrated, interdisciplinary investigation of the response of the northwestern Indian Ocean to monsoonal forcing. *Deep-Sea Research II*, 45, 1905-1915.
- Sonzogni, C., Bard, E., & Rostek, F. (1998). Tropical sea-surface temperature during the last glacial period: a view based on alkenones in Indian Ocean sediments. *Quaternary Science Reviews*, 17, 1185-1201.
- Southon, J., Kashgarian, M., Fontugne, M., Metivier, B., & Yim, W. W.-S. (2002). Marine reservoir corrections for the Indian Ocean and southeast Asia. *Radiocarbon*, 44(1), 167-180.
- Stuiver, M., Reimer, P., & Reimer, R. (2017). CALIB 7.1 [WWW program].
- Urey, H. C., Epstein, S., Lowenstam, H., & Emiliani, C. (1954). Climatic history of the Earth by measuring the oxygen isotopic composition of the calcium carbonate in well-preserved marine fossils. Chicago Univ. Il Enrico Fermi Inst. 23p.
- Versteegh, G. J. M., Bosch, H.-J., & de Leeuw, J. W. (1997). Potential palaeoenvironmental information of C₂₄ to C₃₆ mid-chain diols, keto-ols and mid-chain hydroxy fatty acids; a critical review. *Organic Geochemistry*, 27(1-2), 1-13.
- Volkman, J. K., Barrett, S. M., & Blackburn, S. I. (1999). Eustigmatophyte microalgae are potential sources of C₂₉ sterols, C₂₂ - C₂₈ n-alcohols and C₂₈ - C₃₂ n-alkyl diols in freshwater environments. *Organic Geochemistry*, 30(5), 307-318.
- Volkman, J. K., Barrett, S. M., Dunstan, G. A., & Jeffrey, S. W. (1992). C₃₀-C₃₂ alkyl diols and unsaturated alcohols in microalgae of the class *Eustigmatophyceae*. *Org. Geochem*, 18(1), 131-138.
- von Rad, U., Schulz, H., Riech, V., den Dulk, M., Berner, U., & Sirocko, F. (1999). Multiple monsoon-controlled breakdown of oxygen-minimum conditions during the past 30,000 years documented in laminated sediments off Pakistan. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 152(1-2), 129-161.
- Waelbroeck, C., Paul, A., Kucera, M., Rosell-Melee, A., Weinelt, M., Schneider, R., . . . Members, M. P. (2009). Constraints on the magnitude and patterns of ocean cooling at the Last Glacial Maximum. *Nature Geoscience*, 2(2), 127-132. doi:10.1038/Ngeo411
- Wakeham, S. G., Peterson, M. L., Hedges, J. I., & Lee, C. (2002). Lipid biomarker fluxes in the Arabian Sea, with a comparison to the equatorial Pacific Ocean. *Deep-Sea Research Part II*, 49(12), 2265-2301.
- Warnock, J. P., Bauersachs, T., Kotthoff, U., Brandt, H.-T., & Andrén, E. (2018). Holocene environmental history of the Ångermanälven Estuary, northern Baltic Sea. *Boreas*, 47(2), 593-608. doi:10.1111/bor.12281

- Webster, P. J., Magana, V. O., Palmer, T., Shukla, J., Tomas, R., Yanai, M., & Yasunari, T. (1998). Monsoons: Processes, predictability, and the prospects for prediction. *Journal of Geophysical Research: Oceans*, 103(C7), 14451-14510.
- White, A. E., Foster, R. A., Benitez-Nelson, C. R., Masqué, P., Verdeny, E., Popp, B. N., . . . Prahl, F. G. (2013). Nitrogen fixation in the Gulf of California and the Eastern Tropical North Pacific. *Progress in Oceanography*, 109, 1-17.
- Wyrski, K. (1971). *Oceanographic atlas of the international Indian Ocean expedition*: National Science Foundation.
- Wyrski, K. (1973). Physical oceanography of the Indian Ocean. In B. Zeitschel (Ed.), *The Biology of the Indian Ocean* (pp. 18-36). Berlin: Springer.
- Zehr, J. P., & Capone, D. G. (2020). Changing perspectives in marine nitrogen fixation. *Science*, 368(6492).
- Zhu, X., Jia, G., Mao, S., & Yan, W. (2018). Sediment records of long chain alkyl diols in an upwelling area of the coastal northern South China Sea. *Organic Geochemistry*.