# Interannual variation of settling particles that reflect upper-ocean circulation in the southern Chukchi Borderland, 2010-2014

Jonaotaro ONODERA<sup>1</sup>, Eiji Watanabe<sup>2</sup>, Motoyo Itoh<sup>3</sup>, Naomi Harada<sup>2</sup>, Makio C Honda<sup>2</sup>, Anders Tengberg<sup>4</sup>, Yuichiro Tanaka<sup>5</sup>, and Takashi Kikuchi<sup>2</sup>

<sup>1</sup>Research Institute for Global Change(RIGC) JAMSTEC <sup>2</sup>Japan Agency for Marine-Earth Science and Technology <sup>3</sup>JAMSTEC (Japan Agency for Marine-Earth Science and Technology) <sup>4</sup>Aanderaa Data Instruments AS <sup>5</sup>Geological Survey of Japan, AIST

November 26, 2022

#### Abstract

Changes in the lower-trophic marine ecosystem and biogeochemical cycles were considered with respect to the warming Arctic Ocean. The time series monitoring of the hydrography and marine settling particles supplies a large amount of information regarding the marine lower-trophic ecosystem in with respect to the changing Arctic Ocean. To investigate the interannual relationship between the settling particles and hydrographic conditions of the western Arctic Ocean, bottom-tethered sediment trap moorings were deployed at the Northwind Abyssal Plain (NAP) and at the Chukchi Abyssal Plain from October 2010 to September 2014 to the east and west of the Chukchi Borderland. The settling particles at both stations contained a significant amount of lithogenic matter, with biogenic particles being a lateral advection of shelf materials from shelf to basin. The many peaks of settling particle fluxes at Station NAP in 2013–2014 corresponded to the hydrographic events of passing oceanic eddies over the station, in addition to seasonal biological production. Additionally, abundant gelatinous matter was found in some summer samples. The settling flux data of particulate organic matter and the mole ratio and stable isotope ratios of particulate organic carbon and nitrogen in the settling particles reflected the changes in the composition of trapped plankton assemblages under various hydrographic conditions over the course of the four years of the study.

Interannual variation of settling particles that reflect upper-ocean circulation in
 the southern Chukchi Borderland, 2010-2014

- 3
- 4 Jonaotaro Onodera<sup>1</sup>\*, Eiji Watanabe<sup>1</sup>, Motoyo Itoh<sup>1</sup>, Naomi Harada<sup>1</sup>, Makio C. Honda<sup>1</sup>, Anders
- 5 Tengberg<sup>2</sup>, Yuichiro Tanaka<sup>3</sup>, Takashi Kikuchi<sup>1</sup>
- 6
- 7 <sup>1</sup>: Research Institute for Global Change, Japan Agency for Marine-Earth Sciences and Technology, 2-

8 15 Natsushima-cho Yokosuka, 237-0061, Japan

- 9 <sup>2</sup>: University of Gothenburg / Aanderaa Xylem, Inc.
- 10 <sup>3</sup>: Geological Survey of Japan, National Institute of Advanced Industrial Science and Technology,
- 11 Tsukuba Central 7, 1-1-1 Higashi, Tsukuba, 305-8567, Japan
- 12

# 13 Abstract

14 Changes in the lower-trophic marine ecosystem and biogeochemical cycles were considered with 15 respect to the warming Arctic Ocean. The time series monitoring of the hydrography and marine 16 settling particles supplies a large amount of information regarding the marine lower-trophic 17 ecosystem in with respect to the changing Arctic Ocean. To investigate the interannual relationship 18 between the settling particles and hydrographic conditions of the western Arctic Ocean, bottom-19 tethered sediment trap moorings were deployed at the Northwind Abyssal Plain (NAP) and at the 20 Chukchi Abyssal Plain from October 2010 to September 2014 to the east and west of the Chukchi 21 Borderland. The settling particles at both stations contained a significant amount of lithogenic 22 matter, with biogenic particles being a lateral advection of shelf materials from shelf to basin. The 23 many peaks of settling particle fluxes at Station NAP in 2013-2014 corresponded to the 24 hydrographic events of passing oceanic eddies over the station, in addition to seasonal biological 25 production. Additionally, abundant gelatinous matter was found in some summer samples. The 26 settling flux data of particulate organic matter and the mole ratio and stable isotope ratios of 27 particulate organic carbon and nitrogen in the settling particles reflected the changes in the 28 composition of trapped plankton assemblages under various hydrographic conditions over the course 29 of the four years of the study.

30

# 31 1. Introduction

32 Changes in physical oceanographic characteristics and atmospheric conditions related to recent 33 retreats of sea ice in the Arctic Ocean influence characteristic marine ecosystems and 34 biogeochemical cycles (e.g., Wassmann et al., 2011; Grebmeier, 2012; Ardyna et al., 2014; Berge et 35 al., 2015; Hunt et al., 2015; Harada 2016). On the Pacific side of the Arctic Ocean, the annual and 36 interannual variation of the lateral transportation of Pacific water to Arctic seas and the distribution

of summer sea ice are important for changes in marine ecosystems and biogeochemical cycles. The
Pacific water transports heat, freshwater, and shelf matter, such as nutrients and suspended particles,
including plankton, to the basin. The Chukchi Borderland is one of the key areas where the spread of
Pacific-origin waters to the basin can be monitored.

41 Interannual hydrographic variations, such as changes in surface ocean circulation, oceanic 42 eddies, and sea ice condition should be reflected in the interannual variation of plankton assemblages 43 and settling particle fluxes. Such information is important both for understanding changing Arctic 44 marine ecosystems and for use as validation data for the development of numerical models for 45 marine ecosystems in the Arctic Ocean. A collaborative study of the monitoring of settling particle 46 fluxes using sediment traps and the physical oceanographic-marine ecosystem model that was 47 conducted by Watanabe et al. (2014) has suggested that the westward advection of oceanic eddies 48 containing shelf materials is key for explaining the temporal high flux of settling particles in the 49 southern Northwind Abyssal Plain (NAP) Station of the Chukchi Borderland from fall 2010 to 2011. 50 Using sediment trap samples from the NAP Station, the time series changes in the occurrence of 51 some plankton groups and their relationships with hydrographic variations have been discussed 52 (Watanabe et al., 2014; Ikenoue et al., 2015; Matsuno et al., 2014, 2015, 2016; Onodera et al., 2015, 53 2016; Tokuhiro et al., 2018). This discussion and the understanding of the settling flux and the 54 composition of particulate biogenic and lithogenic matters, along with their relationship to the 55 hydrographic background it produced are more significant in the context of data regarding 56 previously studied plankton assemblages in sediment trap samples. This paper describes the 57 interannual variation of settling particle flux and the composition of bulk particle components 58 (particulate organic matter [POM], calcium carbonate, biogenic opal, and lithogenic materials) and 59 examines the relationships between the obtained results and plankton assemblages in the studied 60 samples as well as the hydrographic conditions around mooring sites in the Chukchi Borderland.

61

# 62 2. Material and methods

## 63 2.1. Mooring operations

64 The samples and hydrographic data were obtained using bottom-tethered moorings with a 65 sediment trap in the southern Chukchi Borderland (Fig. 1). The deployment and recovery of 66 sediment trap mooring was performed by R/V Mirai of the Japan Agency for Marine-Earth Science 67 and Technology and Canadian ice breaker CCGS Sir Wilfrid Laurier, CCGS Louis S. St-Laurent, and 68 CCGS Amundsen. The first annual deployment in the southern NAP occurred at the mooring ID of 69 Station NAP10t from October 04, 2010, to September 28, 2011 (Table 1). At the turnaround of the 70 mooring system, which occurs every September or October, the mooring deployment at the position 71 NAP10t was continued until the end of September 18, 2013 (Table 1). Following this, the mooring 72 NAP13t, positioned approximately 60-km south from NAP10t-12t, was adopted owing to

unfavorable conditions of sea ice for deployment at NAP10t that began in September 2013. To compare the differences between the settling particle fluxes at NAP Station and the western area, where nitrate concentrations are higher than those at NAP Station (Nishino et al. 2011), the annual deployment of the sediment trap mooring CAP12t was conducted on the shelf slope on the southwestern side of the Chukchi Spur (Fig. 1). The following mooring deployment at the CAP12t position was cancelled owing to the conditions of the sea ice.

79

## 80 2.2. Sediment trap samples

81 The time series sediment trap SMD26S-6000 (Nichiyu Giken Kogyo, Co. Ltd.) was used for all 82 deployed moorings. The sediment trap was deployed at a depth of approximately 180–260-m depth 83 for all moorings, including CAP12t, and at a deeper layer, approximately 1300-m deep, for NAP10t-84 13t (Table 1). The water depths of NAP10t-12t, NAP13t, and CAP12t are ~1975 m, 1973 m, and 447 85 m, respectively (Table 1). The sampled period of each cup is 10-15 days (Supplementary Table 1). 86 The sample bottle attached to the sediment trap was filled with filtered sea water obtained at 1000 m 87 depth of NAP station and in the southwestern Canada Basin. The antiseptic agent for trapped organic 88 matter in the trap bottle was pH-neutralized formalin (4-5 v/v %) with sodium tetraborate. The 89 recovered sediment trap samples were sieved using a 1-mm mesh to remove swimmers. The filtrated 90 sample was evenly split into 10 subsamples using the wet sample divider WSD-10 (McLane 91 Research Laboratories, Inc.). Then, 1-3 of the 10 subsamples were filtered using a weighted 92 polycarbonate filter (pore size 0.45µm) and were desalted with Milli-Q water. This filter was dried in 93 the desiccator with diphosphorus pentoxide for a few days. The dried filter was weighed to calculate 94 total mass flux (TMF), and then the particles of the weighed sample detached from the filter were 95 lightly milled to homogenize. The powder samples were sued to analyze the bulk components 96 (carbon, nitrogen, biogenic opal, and others) and stable isotope analyses of the particulate organic 97 carbon (POC) and particulate nitrogen (PN). The contents of particulate carbon and nitrogen were 98 measured using the elemental analyzer FlashEA from Thermo Fisher Scientific. The POC was 99 analyzed after hydrochloric acid gas treatment in the desiccator to remove carbonate. The particulate 100 inorganic carbon was estimated by the difference of total particulate carbon and POC content. The 101 POM contents were estimated using Redfield ratio and the POC contents. The stable isotope 102 analyses of POC and PN were done using the mass spectrometer Delta Plux XP IRMS by Thermo 103 Fisher Scientific. This paper simply describes the stable isotope ratios of POC and PN as d<sup>13</sup>C and 104  $d^{15}N$ , respectively. The mole ratio POC/PN is given as C/N in this paper. Biogenic opal analyses 105 were conducted through alkali leaching (Mortlock and Froelich, 1989) and molybdate yellow 106 methodology. In bulk component analyses, the remainder of the POM, CaCO<sub>3</sub>, and biogenic opal in 107 the total mass was treated as other, which largely includes lithogenic matters with possible errors in 108 the bulk component analysis. In addition, elementary analysis using ICP-AES was conducted for the

109 first deployment samples of NAP10t.

110

## 111 2.3. Hydrographic data and deployed sensors

The daily concentrations of sea ice and downward short-wave radiation around the NAP and
CAP Stations were obtained from re-analyzed data from the National Centers for Environmental
Prediction/Climate Forecast System Reanalysis (Saha et al., 2010).

115 Hydrographic sensors were attached to the moorings CAP12t and NAP13t (Table 2). The 116 applied sensors are CT for CAP12t and NAP13t, current (ADCP and electromagnetic sensors), 117 turbidity, chlorophyll-a, and ice profiling sonar for NAP13t (Table 2). Two different units of 118 turbidity sensors (FTU: Formazin Turbidity Unit, or FNU: Formazin Nephelometric Unit) for 119 NAP13t were not unified at one unit. Those sensors were mainly moored in upper 300 m where there 120 are pacific-origin and melt waters. The backscattered data obtained by ADCP from NAP13t was 121 used to estimate the qualitative change in particle concentrations in water using the method of 122 Deines (1999). In the estimation process of particle concentration by ADCP, the method of removing 123 the influence of the attenuation of acoustic signals for a depth of 140 m is described in Itoh et al. 124 (submitted).

125

### 126 3. Hydrographic backgrounds

127 The Chukchi Borderland is located in a frontal area between the oceanic Beaufort Gyre, with its 128 oligotrophic surface water, in the east and a relatively eutrophic water mass in west (Nishino et al., 129 2011). The east-west gradient of the nitrate concentrations in the Chukchi Borderland suggests a 130 higher settling flux of biogenic particles in the west, where CAP12t was deployed, than in the east, 131 where the NAP Station series was deployed (Watanabe et al., 2014).

Previous research has suggested that the development of oceanic eddies off the Barrow Canyon and their westward advection to the NAP region along the outside of the Beaufort Gyre are an important temporal transport mechanism of shelf material inputs to the sediment trap samples obtained at NAP10t (Watanabe et al., 2014; Harada, 2016). The size and center of the Beaufort Gyre changes year to year (Mizobata et al., 2016). Following the physical oceanographic model for the Arctic Ocean, Onodera et al. (2015) and Watanabe et al. (2015) suggested that the NAP12t was covered by the oceanic Beaufort Gyre.

Condition of downward shortwave radiation was similar between NAP and CAP stations (Fig. 2a). The data shows that there is polar night period in November-January. Sea ice covered the NAP region from October to August (Fig. 2b). The ice-cover period at CAP12t began about 1 week earlier than it did at NAP12t. The sonar profiling record for the sea ice that was taken from the top of NAP13t mooring (~23 m depth) showed that the daily mean thickness of the sea ice grew from October 2013 to May 2014 and shrank beginning in July 2014 (Fig. 2b). The maximum daily mean

- 7 8
  - 3

thickness at the end of April 2014 was 19.6 m. Thicker sea ice (mean thickness of 5.4 m) temporarilypassed over NAP13t in mid-August 2014.

147 The current direction for the upper water column at NAP13t was largely northwest to westward 148 (Fig. 3). The mean current speed for the particle sampling period was  $6 \pm 5$  cm/s at depths of ~170 149 m, which was 20 m above the shallower sediment trap of NAP13t (Fig. 4). The increase in the speed 150 of the current and the change in its direction, along with the change in the water mass, were observed 151 at the deployed sensor depths (Fig. 3). Some of these current change events probably reflected the 152 passing of the oceanic eddy at NAP13t. The eddy event for NAP13t was recognized at least four 153 times, according to ADCP data (Table 3). The first eddy event was observed during the latter half of 154 September 2013. In this event, Bering summer water and cold, newly-ventilated Pacific winter water 155 was observed at depths of 25-45 and 170-215 m, respectively (Fig. 4a, b, c). The direction of the 156 current quickly changed from northeast to southwest. As the eddy moved westward, this eddy 157 appeared to be of an anticyclonic type, with a cold core at depths of 100–150 m. The second eddy 158 event was observed in November 2013. The rapid increase of salinity at depths of  $\sim$ 170 and  $\sim$ 200 m 159 and suggests upwelling of deeper waters for the second eddy event (Fig. 4a, b, c). If the temporal 160 upwelling for the second eddy event reflected a passing cyclonic eddy, the cyclonic eddy probably 161 moved around NAP13t from the west to the east, judging by the direction change of the current, 162 from northward to southeast. The third event, occurring in January 2014, and the fourth event, 163 extending from the end of May to early June 2014 are judged to be westward motions of 164 anticyclonic oceanic eddies. The fifth eddy event for July 2014 was minor compared to previous 165 eddy events. However, there was clock-wise current change in the period (Fig. 3). In addition, 166 Kawaguchi et al. (2016) reported the occurrences of anticyclonic oceanic eddies around the position 167 of NAP10t-12t in September 2014. The temporal upwelling of Atlantic water observed at 196 m in 168 late August-early September 2014 might be related to the eddy event around NAP10t-12t.

169

170 4. Results

#### 171 4.1. Station NAP

The TMF and the bulk component of the settling particles obtained by the shallower traps of NAP10t and NAP11t has been published by Watanabe et al (2014) and Onodera et al. (2015). Using previously published data, we compiled our results with other flux data from the deeper trap of NAP10t-11t and the entire data sets of NAP12t and NAP13t (Fig. 5).

176

#### 177 *4.1.1 TMF*

For the entire sampled duration at NAP10t-12t, from October 4, 2010, to September 17, 2012, the TMF ranged from 5.0 to 263.3 mg m<sup>-2</sup> d<sup>-1</sup> (median: 31.8 mg m<sup>-2</sup> day<sup>-1</sup>) for shallow trap and from <0.1 to 259.3 mg m<sup>-2</sup> day<sup>-1</sup> (median: 19.4 mg m<sup>-2</sup> day<sup>-1</sup>) for deep traps (Fig. 5c, d). The shallow and

- 9
- 10

deep sediment traps of NAP13t showed the similar and relatively high TMF ranged from 3.1 to
342.3 mg m<sup>-2</sup> day<sup>-1</sup> (median: 31.0 mg m<sup>-2</sup> day<sup>-1</sup>), and from 2.4 to 180.2 mg m<sup>-2</sup> day<sup>-1</sup> (median: 47.0 mg m<sup>-2</sup> day<sup>-1</sup>), respectively. The maximum TMF was clearly observed in early winter and summer,
with the exception of a low flux period of 2012 (Fig. 5).

185

# 186 *4.1.2 Bulk components of trapped settling particles*

187 The most abundant composition of the bulk components of trapped particles was "others" (total 188 mass - biogenic materials), which is almost entirely composed of lithogenic matter in many studied 189 samples (median: ~ 69% for NAP Shallow and Deep trap sample. see supplementary Table 1). The 190 elemental composition analyzed for NAP10t shows high aluminum content, ranging from 5.8 to 7.6 191 wt% and from 6.1 to 8.0 wt% for shallow and deep trap samples, respectively. The ranges of POC 192 content in the settling particles were 1.6-21.7 wt% and 1.1-15.5 wt% for shallow and deep trap 193 samples, respectively. The estimated %POM content based on %POC and the Redfield ratio was 194 4.6-62.0 wt% and 3.2-44.4 wt% for shallow and deep trap samples. Biogenic component 195 (POM+CaCO3+Biogenic Opal) ranged from 14.0 to 68.0, and from 12.0 to 74.4 wt% for shallow 196 and deep trap samples, respectively.

- 197 With the exception of 2012, the POC and PN fluxes had remarkable increase in January-198 February, July-September, and November-December (Fig. 6a, b). The mole ratio for C/N ranged 199 from 6.2 to 12.7 for all samples of the NAP Station (Fig. 6c). The average mole ratio for C/N at the 200 deep trap for the entire studied period is 1.2 higher than that for the shallower trap. The large 201 discrepancy of the C/N relationship between shallower and deeper traps was usually observed in low 202 POC flux periods at deep trap (Fig. 6c). Stable isotopes of  $d^{13}C$  and  $d^{15}N$  ranged from -29.3 to -20.9, and from 3.7 to 12.0 in all NAP Station samples (Fig. 6d, e). Although the C/N ratio, d<sup>13</sup>C, 203 204 and  $d^{15}N$  showed no clear seasonality, the changes in those data suggested temporal changes of 205 trapped particle contents. In April 2011, the mole ratio for C/N decreased and that for d<sup>15</sup>N increased 206 (Fig. 6c, e). The shallower trap sample for this event contained many specimens of tiny juvenile 207 amphipods. The deeper trap sample for this event contained fragments of biogenic matter, which 208 might have originated from higher nutrient level organisms. The lowest values for the C/N ratio, 209 d<sup>13</sup>C, and d<sup>15</sup>N were observed in shallower sediment trap samples in late July–August 2012 (Fig. 6c, 210 e). Low d<sup>15</sup>N values of less than 5 were observed only for this event. Beginning in October 2012, 211  $d^{15}N$  and  $d^{15}C$  quickly increased to standard or higher levels relative to the earlier period of the study. 212 In contrast with the result for summer 2012, the d<sup>13</sup>C and mole ratios of C/N increased in August 213 2013. At NAP13t, beginning with fall 2013, the time series pattern of mole ratio of C/N and d<sup>15</sup>N for 214 shallower trap had an opposite correlation (Fig. 6c, e). 215 The PIC flux was lower than the biogenic opal and POC (Fig. 6f, g). The %PIC was lower than
- 216 5.3 wt% and 5.7 wt% for shallow and deep trap samples. Clear seasonality was not observed in the
- 11

PIC flux (Fig. 6f), and the temporal increase of PIC was due to trapped microzooplankton bearingcalcareous shells (Watanabe et al., 2014). The relative contents of biogenic opal over the entire

219 sampled duration were 0.9–71.7 wt% (median: 14.2 wt%) and 7.4–20.2 wt% (median: 12.0 wt%) for

shallow and deep trap samples (Fig. 6g). With seasonal decreases of sea ice concentration and sea ice
thickness, settling fluxes of biogenic opal, POC and PN increased in early summer except for 2012
(Fig. 6)

**222** (Figs. 2b, 6g).

Gelatinous matter such as Appendicularian houses and small jellyfish were abundant in July– early September 2011 and 2014 (Fig. 6a, 7). Because much of this gelatinous material, including some biogenic matter, such as fecal pellets, remained in the 1 mm mesh sieve intended to separate swimmers from settling particles (<1 mm), the TMF and %POC figures for the jelly event are probably underestimates.

228

# 229 4.1.3 Turbidity and ADCP backscattered data of NAP13t

230 The daily mean for turbidity at the top buoy of NAP13t (nominal depth: 25m) was usually lower 231 than 0.3 FTU. The maximum daily mean for turbidity at the top buoy temporarily reached 13 FTU in 232 December 2013 (Fig. 8a). The daily mean turbidity at depths of  $\sim$ 170 m, which is 20 m above the 233 shallower sediment trap, ranged from 0.9 to 1.1 FNU throughout the sampled duration. The acoustic 234 strength of the ADCP data for the depth interval of 12-116 m shows relatively greater particle 235 concentrations in the upper water, at depths shallower than 50 m, in late September 2013, late 236 January 2014, and after May 2014 (Fig. 8b). The acoustic strength at a depth of 170 m was relatively 237 high in September 2013, late December 2013-early March 2014, and late May-September 2014 238 (Fig. 8c). These high signal periods at  $\sim$ 170 m basically correspond to a mid-high signal for depths 239 of 12-50 m (Fig. 8b).

240

241 4.2. CAP12t

#### 242 4.2.1 TMF

The range of the TMF at CAP12t was from 2.8 to 255.1 mg m<sup>-2</sup> day<sup>-1</sup> (Fig. 9a), which was similar to the TMF range at NAP12t. The high TMF for CAP12t was observed from March to early May 2013 (Fig. 9a), by contrast with the high TMF periods for NAP12t (Fig. 5). The particle flux in summer 2013 was lower than in high flux event of March–May 2013. However, note that the gelatinous matter that remained on the 1-mm mesh during sample treatment, was observed in the trap sample bottles from July 20 to August 08, 2013.

249

# 250 4.2.2 Bulk components of trapped settling particles

The content rate of lithogenic matters at CAP12t was 3-63% (median: 46%), which was lower
than that at NAP stations. The contribution of biogenic opal to TMF at this station (11-53 wt%) was

253 relatively higher than that at NAP stations. At the period of maximum TMF observation, in March, 254 biogenic particle fluxes such as POC, PN and biogenic opal also increased (Fig. 9). In the periods of 255 lower TMF, in winter and summer (from May 16 to July 29, 2014), the lithogenic particle flux based 256 on "others" composition decreased whereas POC and PN fluxes did not decrease or increased. The 257 POC% in the low TMF periods from December 10, 2013, to January 16, 2014, and from May 16 to 258 July 29, 2014, became higher because of relative decrease of lithogenic material. The increase of 259 POC flux (Fig. 9b) in summer reflected biological production, including gelatinous materials. The 260 increase of PN flux in summer was not so high compared to the POC flux (Fig. 9c). The mole ratio 261 of C/N ranged from 6.9 to 18.3 (Fig. 9d). The highest C/N ratio was observed in the presence of 262 abundant gelatinous matter in the trap bottles. The stable isotope for d<sup>13</sup>C and d<sup>15</sup>N ranged from 263 -29.1 to -21.9, and from 7.5 to 14.0, respectively (Figs. 8e, 8f). PIC contents were usually low, and 264 the flux was less than 0.3 mg m<sup>-2</sup> day<sup>-1</sup> (Fig. 9g), lower than that found at NAP Station (Fig. 6f). 265 Biogenic opal flux showed the maximum in March 2013 (Fig. 9h), which is comparable to the 266 summer flux maximum of shallower sediment trap at NAP13t (Fig. 6g). However, as mentioned 267 below, the microscopic observation suggested that fragment of diatom frustules is the major 268 component of trapped biogenic opal in March 2013.

269

#### 270 5. Discussion

# 271 5.1. Time series fluctuation of settling particle flux reflecting hydrography

# 272 5.1.1 Change in TMF dominated by hydrographic condition at NAP13t

273 In the intensified current condition reaching to  $\sim 0.2 \text{ m s}^{-1}$  just below the shallower trap for the 274 first oceanic eddy event in the early period (late half of September 2013, Table 3)) of NAP13t 275 observations (Fig. 4d, e), the settling particle flux at shallower trap depths may not have reflected the 276 true amount of settling particle flux (Fig. 8d). The turbidity and back-scatter strength of the single 277 point ADCP deployed at depths that are  $\sim 20$  m above the shallower trap shows that particle 278 concentrations in water were commonly abundant but lower than those of later periods (Fig. 8c). The 279 increases in TMF at deeper trap depths during the same period can be interpreted as a reflection of 280 higher particle concentrations in the upper water column. According to the vertical distribution of 281 particle concentrations and TMF data, many particles in the water during the first eddy event were in 282 the Bering summer water at a depth of around 25 m, and the cold winter water in the deeper layer 283 appeared during the eddy event (Fig. 4). When the second eddy, which was considered cyclonic, 284 passed NAP13t in late November 2013, TMF and the backscattered strength of the ADCP did not 285 increase, likely due to the upwelling of deeper waters with relatively low particle concentration. It 286 appears that the high TMF period, beginning in late December 2013, did not correspond to an eddy 287 event. However, the particle concentration in the upper water with a depth of around 20 m began to 288 increase in December 2013, and a mass of water with abundant particles temporally expanded in

289 depth when the third eddy core passed NAP13t in early January 2014 (Figs. 4 and 7). The core depth 290 of the third oceanic eddy was  $\sim$ 50 m, and mean current speeds around the shallower trap was 7.8±1.8 291 m s<sup>-1</sup> and  $4.5\pm1.9$  cm s<sup>-1</sup> at 170 m and 209 m for the event #3. Therefore, the condition of the current 292 that influences trapping efficiency for the shallower trap was stable for the eddy event #3 (Fig. 4). 293 These observations suggest that the appearance of water with large amounts of particles and an 294 increase in TMF from December 2013 can be related to the third oceanic eddy event. When the 295 fourth eddy event occurred in the end of May 2014, the acoustic signal for particle concentration 296 increased, although TMF did not increase, likely because of the high current speed during the eddy 297 event. When the settling particle fluxes increased in July 2014, no rapid increase of speed or change 298 in the direction of the current suggesting passing oceanic eddy were clearly observed. The increases 299 of particle concentration for upper 50m (Fig. 8b) and settling particle flux in July 2014 may reflect 300 an influence of seasonal sea ice retreat (Fig. 2b). However, a circular change in the direction of the 301 current was observed at depths of around 50-170 m in July (Fig. 4). Kawaguchi et al. (2016) 302 detected westward anticyclonic cold eddies with core depths of 100-150 m around the position of 303 NAP10-12t in early September 2014. If this change in the direction of the current reflected passing 304 oceanic eddies, the high TMF in August-September 2014 can be (at least partially) explained by 305 advection of shelf materials with a passing oceanic eddy in addition to an increase of vertical particle 306 transport by seasonal retreat of sea ice and increases of biological production in summer.

307

308

#### 5.1.2 Interannual variation of trapped biogenic particles reflecting hydrography at NAP Station

309 The relationships among the time series fluctuation of particle concentration in upper 50m 310 waters, settling particle fluxes and changes in hydrographic conditions observed at NAP13t suggest 311 that oceanic eddies play a partially significant role as a transporter of shelf materials advected to the 312 mooring position. This relationship was estimated through the collaborative study of settling particle 313 flux at NAP10t and the application of a physical oceanographic model developed exclusively for the 314 Arctic Ocean (Watanabe et al., 2014). Because settling particles are dominated by lithogenic 315 materials, the interannual variation of TMF reflects the status change of ocean currents for the upper 316 water column, such as the oceanic eddy passing, as mentioned above.

317 The interannual variation of  $d^{13}C$ ,  $d^{15}N$ , the ratio of C/N, and the flux of trapped plankton groups 318 for every spring and summer from 2011 to 2014 suggest a difference in biogeochemical conditions 319 for lower trophic production, in addition to the physical hydrographic conditions. When the biogenic 320 opal and POC fluxes at NAP10t increased slightly in April-May 2011 (Fig. 6a, g), large diatom 321 species such as Coscinodiscus spp. contributed to the flux of POC (Onodera et al., 2015). Although 322 the settling flux of other siliceous microplankton radiolarians slightly increased during this period 323 (Ikenoue et al., 2015), the contribution of the shell to the biogenic opal flux was low (Ikenoue et al., 324 in prep.). The abundance of the herbivore copepoda *Calanus hyperboreus* increased in March-May

325 2011 (Matsuno et al., 2013, 2015). In addition to seasonal primary production under sea ice and the 326 lateral advection of re-suspended matter, the contribution of fecal pellets resulting from the grazing 327 pressure effect suggests settling POM and biogenic opal fluxes. The sudden increase of d<sup>15</sup>N and 328 corresponding decrease of C/N in April 2011 can be primarily explained by the abundant trapped 329 small zooplankton that are less than 1 mm, which is largely a number of juvenile amphipods 330 detached from the abdominal part of one parent specimen during sample treatment. This is supported by the stable isotope values for amphipods, which show a  $d^{15}N$  of ~11‰ and a  $d^{13}C$  of ~ -25.4‰ 331 around the Beaufort Gyre (Pomerleau et al., 2014). The quite low values of d<sup>13</sup>C, reaching to 332 333 -28.6% at the shallow trap in July 2011, were lower than the  $\sim 27\%$  found for POC in the Chukchi 334 Sea (Zhang et al., 2012). The phytoplankton growing under high  $pCO_2$  in cold water are able to 335 tolerate such low d<sup>13</sup>C values (Tesi et al., 2018). The decreases of d<sup>13</sup>C in late June–July 2011 also 336 reflect zooplankton ecology. For this timing, rapid increases in numbers of silicoflagellate skeletons 337 were observed with many fecal pellets in the trapped particles in shallow traps but not in the deep 338 traps (Onodera et al., 2016). From late June to July 2011, the composition of three fecal-pellet types 339 suggested intensified grazing pressure by copepoda and raptorial amphipod (Matsuno et al., 2016). 340 The hydrographic conditions of active primary production may have a primary influence on the 341 stable isotope condition of carbon and nitrogen in zooplankton for herbivorous feeding, as reported 342 for the Beaufort Sea (Forest et al., 2011).

343 It is unclear why decreases in d<sup>13</sup>C in June–July 2011 began at the deep trap 2 weeks earlier than 344 at the shallow trap. Takeuchi et al. (unpublished) showed a difference in time series trend for the 345 composition of lithogenic matter that originates in deep and shallow trap samples, using rare earth 346 element analysis. In general, the origin area of trapped particles for the deeper sediment trap may be 347 different, and broader, than that at the shallower trap, although it can also depend on the particle 348 sinking rate (Siegel et al., 2008). Horizontal diffusion of settling particles with low sinking rate is 349 estimated under mesoscale oceanic eddy (Siegel et al., 1990). If we follow the numerical model 350 (Siegel et al., 1990), the particles transported by eddy approaching to mooring position can be 351 reached earlier at the deeper trap than at the shallower trap, for some values of distance between the 352 oceanic eddy and the mooring position.

353 Among the four annual deployment records of the settling particle fluxes at NAP Station in this 354 study, particle flux data from early 2012 to middle 2013 (Fig. 5) showed an unusual annual pattern. 355 As far as suspected from mooring depth change for NAP11t-NAP12t (Fig. 5a, b), subsurface current 356 condition around the shallower trap was stable except for July 2013 (Onodera et al. 2015). The low 357 particle fluxes at both shallow and deep traps might mainly reflect low settling particle condition in 358 addition to particle trapping efficiency mainly estimated for July 2013. Onodera et al. (2015) 359 suggested that the continuation of low particle fluxes at NAP12t and diatom assemblage in trap 360 samples similar to those in the Canada Basin reflected a westward shift of oligotrophic water masses

in the Beaufort Gyre. Matsuno et al. (2016) suggested a similar change in hydrographic condition, 361 362 based on the absence of barnacle and bivalve larvae in sediment trap samples for summer 2012. 363 Nitrogen in the southwestern Canada Basin during the summer is usually depleted in surface waters, 364 by contrast with its phosphate concentrations (Shiozaki et al., 2018). Blais et al. (2012) and Shiozaki 365 et al. (2018) reported findings of nitrogen fixation and its role in the nitrogen cycle in the Arctic 366 Ocean. In relation to those hydrographic conditions for summer 2012, the low d<sup>15</sup>N values for POM 367 may suggest that there was partial contribution of nitrogen fixation for POM formation. In the 368 central Canada Basin in September 2012, with its low sea ice concentrations, the  $pCO_2$  in surface 369 water was ~30 µatm greater than that in the northern Canada Basin, which had a great deal of sea ice 370 (Islam et al., 2017). The fact that the lowest level of  $d^{13}C$  appeared in summer 2012 seems to suggest 371 that the POM settled and trapped in the deployed sediment trap was formed owing to the condition 372 of the lower growth rate of phytoplankton and the condition of high  $pCO_2$  in surface water with 373 reduced sea ice. The temporarily intensified current, as suspected by changes in the mooring depth of 374 the sediment trap for July 2012 (Fig. 5a, b), might take trapped particles far from the vicinity of the 375 mooring station. However, the possibility for the specific influence of the intensified current such as 376 change of trapping efficiency (Matsuno et al. 2014) to lower levels of d<sup>13</sup>C and d<sup>15</sup>N remains unclear, 377 as the d<sup>13</sup>C and d<sup>15</sup>N values remained low until September 2012. The increase of stable isotope 378 values in October 2012 (Fig. 6d, e) suggests that the clear influence of the oligotrophic water ceased 379 around the end of September. The diatom assemblages in the trap samples also changed from 380 oligotrophic planktic flora to ice-related flora (Onodera, unpublished) even in the ice-free condition 381 at Station NAP for early October 2012 (Fig. 2b). The change in biogeochemical data that occurred 382 from September to October 2012 suggests a change in the major direction of the current, probably 383 associated with a shift of the distribution of the Beaufort Gyre.

384 Higher sea ice concentrations in summer 2013 led to the interannual hydrographic characteristics 385 observed at NAP Station for the four summer seasons from 2011 to 2014 (Fig. 2b). Continuous ice-386 free condition at Station NAP was observed for September (or late August - September) in 2011, 387 2012, and 2014 whereas the ice-free ocean condition was limited in summer 2013 (Fig. 2b). The 388 highest d<sup>13</sup>C, which appeared in late July–early August 2013 (Fig. 6d), suggests that ice algae 389 particles contributed to the trapped settling particles. According to microscopic observations of the 390 samples, the diatom flora in the trap samples for this period was dominated by ice-related species 391 (Onodera, unpublished). For the Chukchi and Beaufort Seas in spring-summer 2002, the d<sup>13</sup>C of the 392 POC in sea ice ranged from -25.1 to -14.2%, and it was linearly correlated with POC 393 concentrations (Gradinger, 2009). Tesi et al. (2018) showed that the POC produced by ice algae in 394 the East Siberian Sea has a heavier of  $d^{13}C$ , reaching about -21%. In general, the ice algae under sea 395 ice on the Pacific side of the Arctic Ocean are more abundant on the shelf side than in the 396 oligotrophic Canada Basin (Nelson et al., 2014). In addition to the input of particles from ice algae to

397 the deployed sediment trap, the flux of lithogenic material increased in summer 2013.

The temporary decrease in C/N and increase in d<sup>15</sup>N preceding the high biogenic particle fluxes of June–early September 2014 are essentially caused by trapped small zooplankton in the studied samples from the shallow trap, as indicated for the case of summer 2011. The highest POC and PN fluxes for the shallower trap in summer 2014 were observed in lithogenic matter (Figs. 5 and 6), which may reflect the cold anticyclonic eddies with higher suspended particle concentrations in the eddy core located ~27 km north of NAP13t in early September 2014 (Kawaguchi et al., 2016).

404 It is unclear that oligotrophic condition and low flux of settling particles at Station NAP in 2012 405 can be treated as one of future condition at Station NAP in progressed global warming. Throughout 406 four-year monitoring on settling particles, the low particle-flux trend in 2012 with minimum summer 407 ice extent was different from time-series settling particle flux in other years. The position and area of 408 oceanic Beaufort Gyre, which influenced to low flux of settling particles in 2012, reflects dynamics 409 of atmospheric circulation over the Arctic Ocean (Proshutinsky et al. 2009, Regan et al. 2019). 410 Decadal time-scale researches on Arctic climatology and physical oceanography suggests that recent 411 physical condition over the Arctic tends to bring westward expansion of oceanic Beaufort Gyre. 412 However, there is no clear research results to explain that those recent climatological physical 413 dynamics can be explained as the reflection of global warming.

414

## 415 *5.1.3 CAP Station*

416 The ranges of  $d^{13}C$  found at CAP Station (-29.1 ~ -21.9) are higher than those at shallower trap of 417 NAP12t (-28.2 ~-21.2), which suggests that much of the influence of POM originated from the 418 shallower Chukchi Sea shelf, based on the d<sup>13</sup>C distribution over the Chukchi Sea (Zhang et al., 419 2012). In contrast to what was expected before the mooring deployment was begun, based on recent 420 nitrate distribution patterns in the Chukchi Borderland (Nishino et al., 2011), the settling particle flux 421 at CAP12t was not higher than that found at NAP12t. This is probably due to the interannual 422 variations of water mass distribution in 2012-2013. Because the data obtained at CAP Station only 423 covered 1 year, additional study with more data and samples is required to for the relationship 424 between settling particles and hydrographic condition around CAP Station to be properly 425 characterized.

The high particle flux found in March 2013 can be explained as the lateral advection of resuspended sediment on the shelf. The high particle flux in March 2013 may have reflected the input of warmer Pacific water, which moved in a current westward along the shelf edge of the Chukchi Sea. The increase in water temperatures in March 2013, from about -1.2 to  $-0.8^{\circ}$ C at a depth of ~90 m (Fig. 4f), suggests an influence of advected warmer Pacific water to the sub sea-surface of the mooring position of CAP12t during the event (Watanabe et al., 2017). Microscopic observation suggests that the high biogenic opal in this period can be explained by the presence of many

fragments of diatom frustules, which are treated as re-suspended particles. As has been already noted regarding on this hydrographic event, in Watanabe et al. (2017), the westward advection of Pacific water along the northern shelf of Chukchi Sea was the main contributor to lateral material transportation. Throughout the entire observation period, the contributions of oceanic eddy advection at CAP12t was unclear or negligible because of their location.

438

## 439 5.2. Gelatinous matter

At the outset of this study, the settling particles obtained were defined as being smaller than 1 440 441 mm, following precedent in sediment trap studies (e.g. Honjo et al., 2008). The discarded 442 Appedicularian houses gravitationally sink to ocean interior, and it has a characteristic of settling 443 particles.although their sizes are larger than 1 mm (Fig. 7). Because POC flux in this study is the 444 results for particles smaller than 1mm, the contribution of gelatinous matters such as large 445 Appendicularia house to the trapped settling samples in this study is not reflected in the obtained 446 analysis data because of the methodology used. Some larger planktons such as Phaeodaria larger 447 than 1mm may partially contribute to settling POC flux (e.g., Ikenoue et al. 2019). However, the 448 particles remained on 1mm sieve in this study is almost swimmers such as copepoda and amphipoda. 449 The occurrence pattern of abundant Appendicularian houses in this study may suggest that the 450 contribution of Appendicularian houses to POC flux is patchy but locally and temporally important. 451 Regarding the temporary occurrence of abundant gelatinous matter in this study, mainly composed 452 of discarded Appendicularian houses, their likely importance for Arctic marine ecosystems should be 453 noted. Appendicularia around the study area are mainly distributed along the shelf break of the 454 Canada Basin and on the Chukchi Sea shelf (Lane et al., 2008; Hopcroft et al., 2010), and their 455 abundance is markedly increased compared to the levels noted in the 1950s (Lane et al., 2008), as an 456 effect of grazing on micro and meso plankton, including Copepoda (Lalande et al., 2011), the 457 vertical transport of POC and other suspended particles, including microplastics, to the deep ocean 458 (Deibel et al., 2005; Lalande et al., 2011; Katija et al., 2017), and the food source for fish in the 459 Chukchi Sea (Nakano et al., 2016). The Appendicularia Oikopleura vanhoeffeni were commonly 460 observed in the midst of phytoplankton blooms at a depth of around 50 m near the ice edge of the 461 shelf slope in the southwestern Canada Basin in late July 2005 (Raskoff et al., 2010). The gelatinous 462 event at NAP Station in summer 2011 was complete when the settling diatom flora changed from 463 planktic to ice-related taxa just before the retreat of the sea ice retreat (Onodera et al., 2015). 464 Oikopleura vanhoeffeni in Newfoundland has a habitat range of -1.3 to 4.6 °C for water temperature 465 and 32.3 to 32.7 for water salinity (Choe and Deibel, 2008). Because Appendicularia are filter 466 feeders, the phytoplankton bloom under sea ice around the study region in early summer (Arrigo et 467 al., 2012; Lowry et al., 2014) could be more favorable than feeding the largely ice algae attached to 468 sea ice from below in low saline water. Because detailed analysis on gelatinous matters were not

performed for the trap samples, additional study on this topic is necessary, using new sediment trapsamples around the study region.

471

## 472 5.3. Contribution of resuspended biogenic matters

473 As estimated from the high abundance of lithogenic matter in the samples, trapped biogenic 474 particle matter is expected to contain re-suspended particles originating in the sediment of the shelf 475 region. The increase of biogenic matter flux in the non-productive season can be basically explained 476 as laterally transported matter. The distinction between biogenic particle flux in the resuspended 477 component and upper water origin is important for the understanding of the potential of primary 478 productivity and the biological carbon pump around the mooring station. Magen et al. (2010) 479 estimated different origins for POM in the southern Beaufort Sea using end members for the ratio of 480 POC and total particulate nitrogen, d<sup>13</sup>C for POC and d<sup>15</sup>N for total particulate nitrogen. However, 481 the end member for  $d^{13}C$  of POC, showing the coastal origin of POC, is similar to the  $d^{13}C$  value for phytoplankton in d<sup>13</sup>C-depleted water (Tesi et al., 2017). Thus, the estimation of the quantitative 482 483 contribution of resuspended biogenic matter to total biogenic matter using analyzed items was not 484 successfully performed in this study. According to Takeuchi et al. (unpublished), who analyzed rare 485 earth elements in settling particles obtained from NAP10t, trapped lithogenic particles had a mixture 486 of three different sources: the Mackenzie Delta, the Yukon River, and East Siberian regions. They 487 suggested that sources of lithogenic matter for shallow and deep traps changed with time. 488 Resuspended biogenic matter could be an additional source of food for benthos and nekton 489 assemblages in the study region. Relative to the relationship between changing physical 490 oceanographic conditions and marine ecosystems, it is likely that monitoring the behavior of 491 advected particles will be important for assessing future Arctic marine ecosystems.

492

# 493 6. Summary

494 To understand the relationships between hydrographic conditions and settling particle fluxes, 495 time series sediment traps were deployed annually and interannually at NAP Station and at CAP12t 496 on the shelf slope of the western Chukchi Sea, from October 2010 to September 2014. Although the 497 resulting data were limited to 4 years, the influence of hydrographic conditions on settling particle 498 flux in the Chukchi Borderland was obvious. The trapped settling particles contained a great deal of 499 shelf-origin materials. At NAP Station, many maxima of settling particle fluxes corresponded to 500 passing events of oceanic eddy advections along the oceanic Beaufort Gyre. When the oligotrophic 501 surface water of the Canada Basin covered the Chukchi Borderland in 2012, the settling particle flux 502 did not increase at NAP Station, although it did increase at CAP12t in March 2013, as a reflection of 503 the westward transportation of warm Pacific water on the shelf, rather than its vertical transport of in 504 situ biological production. These lateral advections of reworked organic matter are able to support

- 505 benthos and nekton in downstream areas, and meaning that further monitoring and model prediction
- 506 will be important to come to an understanding of alterations in Arctic marine ecosystems.
- 507

#### 508 Author's contribution

- 509 JO: analysis of total mass, C, N, opal, field work for mooring deployment and recovery
- 510 EW: input as physical oceanographer
- 511 MI: supply of CT and WH sensors and their data, and Fig. 8b
- 512 NH: PI, comments to this ms
- 513 MCH: ICP element analysis of NAP10t samples
- 514 AT: supply of Aanderaa & Xylem sensors and their data for NAP13t, and Fig. 8c
- 515 YT: supply of Nichiyu equipment, comment on particle transport
- 516 TK: supply of CT data (CAP12t), and cruise logistics, field work for mooring deployment and
- 517 recovery
- 518

## 519 Acknowledgments

- 520 We sincerely thank the captains, crews, chief scientists, and technicians of the R/V Mirai 521 belonging to Japan Agency for Marine-earth Science and Technology (JAMSTEC) and the three ice 522 breakers CCGS Sir Wilfrid Laurier, CCGS Louis S. St. Laurent, and CCGS Amundsen. belonging to 523 the Canadian Coast Guard. We appreciate support by Drs. Hatta, M., Sugie, K. and Ms. Ishiwata, S. 524 on biogenic opal analysis. We thank to Dr. Chiba, S. on observation of gelatinous matters. This work 525 was funded by a Grant-in-Aid for Scientific Research (S) of the Japan Society for the Promotion of 526 Science (JSPS) JFY2010-2014, no. 22221003 to NH, YT, a Grant-in-Aid for Scientific Research (A) 527 of JSPS, no. 15H01736 to JO, EW, YT, JSPS Research Fellowships for Young Scientists no. 22-5808 528 to JO, GRENE, and Arctic Challenge for Sustainability (ArCS, ArCS II). The data used to prepare 529 this paper were tabulated in the Supplementary Table 1 of this paper, and also it will be archived in 530 the Arctic Data archive System (ADS, https://ads.nipr.ac.jp/) by June 2020.
- 531

532

#### 533 References

- Arrigo, K.R., Perovich, D., Pickart, R.S., Brown, Z.W., van Dijken, G.L., & et al. (2012). Massive
  phytoplankton blooms under Arctic sea ice. Science, 336, 1408.
- Ardyna, M., Babin, M., Gosselin, M., Devred, E., Rainville, L., & Tremblay, J.-É. (2014). Recent
  Arctic Ocean sea ice loss triggers novel fall phytoplankton blooms. Geophysical Research
  Letters, 41, 6207–6212, doi:10.1002/2014GL061047
- Berge, J., Renaud, P.E., Darnis, G., Cottier, F., Last, K., Gabrielsen, T.M., & et al. (2015). In the
  dark: A review of ecosystem processes during the Arctic polar night. Progress in Oceanography,
- 29

541 139, 258-271.

- 542 Choe, N., & Deibel, D. (2008). Temporal and vertical distributions of three appendicularian species
  543 (Tunicata) in Conception Bay, Newfoundland. Journal of Plankton Research, 30, 969–979.
- 544 Deibel, D., Saunders. P.A., Acuña, J.-L., Bochdansky, A.B., Shiga, N., & Rivkin, R.B. (2005). The
  545 role of appendicularian tunicates in the biogenic carbon cycle of three Arctic polynyas. In:
  546 Gorsky, G., Youngbluth, M.J., & Deibel, D. (Eds) Response of Marine Ecosystems to Global
- 547 Change, Ecological Impact of Appendicularians, Éditions Scientifiques GB, Paris, pp. 327–356.
- 548 Deines, K. (1999). Backscatter estimation using broadband acoustic Doppler current profilers.
  549 Proceeding of the IEEE 6th Working Conference on Current Measurements, 249–253, IEEE,
  550 Stroughton, Wis.
- Forest, A., Galindo, V., Darnis, G., Pineault, S., Lalande, C., Trambaly, J-É., & et al. (2011). Carbon
  biomass, elemental rations (C:N) and stable isotopic composition (δ<sup>13</sup>C, δ<sup>15</sup>N) of dominant
  calanoid copepods during the winter-to-summer transition in the Amundsen Gulf (Arctic
  Ocean). Journal of Plankton Research, 33, 161–178.
- Gradinger, R. (2009). Sea-ice algae: Major contributors to primary production and algal biomass in
  the Chukchi and Beaufort Seas during May/June 2002. Deep-Sea Research II, 56, 1201–1212.
- Grebmeier, J.M. (2012). Shifting patterns of life in the Pacific Arctic and sub-Arctic Seas. Annual
  Review of Marine Science, 4, 63–78. https://doi.org/10.1146/annurev-marine-120710-100926
- Harada, N. (2016). Review: Potential catastrophic reduction of sea ice in the western Arctic Ocean:
  Its impact on biogeochemical cycles and marine ecosystems. Global and Planetary Change,
  136, 1–17.
- Honjo, S., Manganini, S.J., Krishfield, R.A., & Francois, R. (2008) Particulate organic carbon fluxes
  to the ocean interior and factors controlling the biological pump: A synthesis of global sediment
  trap programs since 1983. Progress in Oceanography, 76, 217-285.
- Hopcroft, R.R., Kosobokova, K.N., & Pinchuk, A.I. (2010). Zooplankton community patterns in the
  Chukchi Sea during summer 2004. Deep-Sea Research II, 57, 27–39.
- Hunt Jr., G.L., Blanchard, A.L., Boveng, P., Dalpadado, P., Drinkwater, K.F., & Eisner, L., et al.
  (2013). The Barents and Chukchi Seas: Comparison of two Arctic shelf ecosystems. Journal of
  Marine Ecosystems, 109–110, 43–68. https://doi.org/10.1016/j.jmarsys.2012.08.003
- 570 Ikenoue, T., Bjørklund, K.R., Kruglikova, S.B., Onodera, J., Kimoto, K., & Harada, N. (2015). Flux
  571 variations and vertical distribution of siliceous Rhizaria (Radiolaria and Phaeodaria) in the
  572 western Arctic Ocean: indices of environmental changes. Biogeosciences, 12, 2019–2046.
- 573 Ikenoue, T., Kimoto, K., Okakzaki, Y., Honda, M.C., Takahashi, K., Harada, N., & Fujiki, T.
  574 (2019). Phaeodaria: An important carrier of particulate organic carbon in the mesopelagic
  575 twilight zone of the North Pacific Ocean. Global Biogeochemical Cycles, 33, 1146-1160.
  576 https://doi.org/10.1029/2019GB006258.
- 31 32

- 577 Ikenoue, T., Kimoto, K., Nakamura, Y., Bjørklund, K.R., Kuramoto, N., Ueki, M., et al. (in prep.)
  578 Quantification of species-specific biogenic silica production of Arctic radiolarians (Rhizaria)
  579 based on Microfocus X-ray computed tomography. To be submitted to Limonology and
  580 Oceanography in this April.
- Islam, F., DeGrandpre, M.D., Beatty, C.M., Timmermans, M.-L., Krishfield, R.A., Toole, J.M., & et
  al. (2016). Sea surface pCO2 and O2 dynamics in the partially ice-covered Arctic Ocean.
  Journal Geophysical Research: Oceans, 122, 1425–1438.
  https://doi.org/10.1002/2016JC012162
- Itoh, M., Kitamura, M., Fujiwara, A., Carmack, E.C., Amakasu, A., Ara, K., & et al. (submitted).
  Increasing zooplankton biomass associated with sea ice loss in the western Arctic Ocean and its
  ecological impacts. Submitted to Polar Science on November 19, 2020.
- Jakobsson, M., Mayer, L., Coakley, B., Dowdeswell, J.A., Forbes, S., & et al. (2012). The
  International Bathymetric Chart of the Arctic Ocean (IBCAO) Version 3.0. Geophysical
  Research Letters, 39, L12609. https://doi.org/10.1029/2012GL052219.
- 591 Katija, K., Choy, C.A., Sherlock, R.E., Sherman, A.D., & Robinson, B.H. (2017). From the surface
  592 to the seafloor: How giant larvaceans transport microplasctics into the deep sea. Science
  593 Advances, 3, e1700715.
- Kawaguchi, Y., Nishino, S., Inoue, J., Maeno, K., Takeda, H., & Oshima, K. (2016). Enhanced
  diapycnal mixing due to near-inertial internal waves propagating through an anticyclonic eddy
  in the ice-free Chukchi Plateau. Journal of Physical Oceanography, 46, 2457–2481.
- Lane, V.Z.P., Llinás, L., Smith, S.L., & Pilz, D. (2008). Zooplankton distribution in the western
  Arctic during summer 2002: Hydrographic habitats and implications for food chain dynamics.
  Journal of Marine Systems, 70, 97–133.
- Lowry, K.E., van Dijken, G.L., & Arrigo, K.R. (2014). Evidence of under-ice phytoplankton blooms
  in the Chukchi Sea from 1998 to 2012. Deep-Sea Research II, 105, 105–117.
- Magen, C., Chaillow, G., Crowe, S.A., Mucci, A., Sundby, B., Gao, A., & et al. (2010). Origin and
  fate of particulate organic matter in the south Beaufort Sea- Amundsen Gulf region, Canadian
  Arctic. Estuarine. Coastal and Shelf Science, 86, 31–41.
- Matsuno, K., Yamaguchi, A., Fujiwara, A., Onodera, J., Watanabe, E., Imai, I., & et al. (2014).
  Seasonal changes in mesozooplankton swimmers collected by sediment trap moored at a single
  station on the Northwind Abyssal Plain in the western Arctic Ocean. Journal of Plankton
  Research, 36, 490–502.
- Matsuno, K., Yamaguchi, A., Fujiwara, A., Onodera, J., Watanabe, E., Harada, N., & et al. (2015).
  Seasonal changes in the population structure of dominant plaknktonic copepods collected using
  a sediment trap moored in the western Arctic Ocean. Journal of Natural History, 49 (45-48),
  https://doi.org/10.1080/00222933.2015.1022613
- 33 34

- Matsuno, K., Yamaguchi, A., Fujiwara, A., Onodera, J., Watanabe, E., Harada, N., & et al. (2016).
  Seasonal changes in mesozooplankton swimmer community and fecal pellets collected by
  sediment trap moored at the Northwind Abyssal Plain in the western Arctic Ocean. Bulletin of
- 616 Fisheries Sciences, Hokkaido University, 66, 77–85.
- Mizobata, K., Watanabe, E., & Kimura, N. (2016). Wintertime variability of the Beaufort Gyre in the
  Arctic Ocean derived from CryoSat-2/SIRAL observations. Journal of Geophysical Research,
  121, 1685–1699. https://doi.org/10.1002/2015JC011218.
- Mortlock, R.A., & Froelich, P.N. (1989). A simple method for the rapid determination of biogenic
  opal in pelagic marine sediments. Deep-Sea Research, 36, 1415–1426.
- Nakano, T., Matsuno, K., Nishizawa, B., Iwahara, Y., Mitani, Y., Yamamoto, J., & et al. (2016). Diets
  and body condition of polar cod (*Boreogadus salida*) in the northern Bering Sea and Chukchi
  Sea. Polar Biology, 39, 1081–1086.
- Nelson, R.J., Ashjian, C.J., Bluhm, B.A., Conlan, K.E., Grandinger, R.R., Gradinger, R.R., & et al.
  (2014). Biodiversity and biogeography of the lower trophic taxa of the Pacific Arctic Region:
  Sensitivities to climate change. In: Grebmeier, J.M., & Maslowski, W. (Eds.) The Pacific Arctic
  Region: Ecosystem Status and Trends in a Rapidly Changing Environment. Springer, pp. 269–336.
- Nishino, S., Kikuchi, T., Yamamoto-Kawai, M, Kawaguchi, Y., Hirawake, T., & Itoh, M. (2011).
  Enhancement/reduction of biological pump depends on ocean circulation in the sea-ice
  reduction regions of the Arctic Ocean. Journal of Oceanography 67, 305–314.
- Onodera, J., Watanabe, E., Harada, N., & Honda, M.C. (2015). Diatom flux reflects water-mass
  conditions on the southern Northwind Abyssal Plain, Arctic Ocean. Biogeosciences, 12, 1373–
  1385.
- 636 Onodera, J., Watanabe, E., Nishino, S., & Harada, N. (2016). Distribution and vertical fluxes of
  637 silicoflagellates, ebridians, and the endoskeletal dinoflagellate *Actiniscus* in the western Arctic
  638 Ocean. Polar Biology, 39, 327–341.
- Pomerleau, C., Nelson, R.J., Hunt, B.P.V., Sastri, A.R., & Williams, W.J. (2014). Spatial patterns in
  zooplankton communities and stable isotope ratios (δ<sup>13</sup>C and δ<sup>15</sup>N) in relation to oceanographic
  conditions in the sub-Arctic Pacific and western Arctic regions during the summer of 2008.
  Journal of Plankton Research, 36, 757–775.
- Raskoff, K.A., Hopcroft, R.R., Kosobokova, K.N., Purcell, J.E., & Youngbluth, M. (2010). Jellies
  under ice: ROV observations from the Arctic 2005 hidden ocean expedition. Deep-Sea
  Research II, 57, 111–126.
- Saha, S., Moorthi, S., Pan, H-L., Wu, X., Wang, J., Nadiga, S., & et al. (2010). The NCEP Climate
  Forecast System Reanalysis. Bull. Amer. Meteor. Soc., 91, 1015–1057.
- 648 Shiozaki, T., Fujiwara, A., Ijichi, M., Harada, N., Nishino, S., Nishi, S., & et al. (2018). Diazotroph
- 35 36

- 649 community structure and the role of nitrogen fixation in the nitrogen cycle in the Chukchi Sea
  650 (western Arctic Ocean). Limnology and Oceanography, 63, 2191-2205.
  651 https://doi.org/10.1002/lno.10933
- Siegel, D.A., Granata, T.C., Michaels, A.F., & Dickey, T.D. (1990). Mesoscale eddy diffusion,
  particle sinking, and the interpretation of sediment trap data. Journal of Geophysical Research,
  95 (C4), 5305–5311.
- Tesi, T., Geibel, M.C., Pearce, C., Panova, E., Vonk, J.E., Karlsson, E., & et al. (2017). Carbon
  geochemistry of plankton-dominated samples in the Laptev and East Siberian shelves: contrasts
  in suspended particle composition. Ocean Science, 13, 735–748. https://doi.org/10.5194/os-13735-2017
- 659 Tokuhiro, K., Abe, Y., Matsuno, K., Onodera, J., Fujiwara, A., Harada, N., & et al. (2018). Seasonal 660 phenology of four dominant copepods in the Pacific sector of the Arctic Ocean: Insights from 661 statistical analysis of sediment trap data. Polar Science, 19. 94-111. 662 https://doi.org/10.1016/j.polar.2018.08.006
- Wassmann, P., Duarte, C.M., Agustí, S., & Sejr, M.K. (2011). Footprints of climate change in the
  Arctic marine ecosystem. Global Change Biology, 17, 1235–1249.
  https://doi.org/10.1111/j.1365-2486.2010.02311.x
- Watanabe, E., Onodera, J., Harada, N., Honda, M.C., Kimoto, K., Kikuchi, T., & et al. (2014).
  Enhanced role of eddies in the Arctic marine biological pump. Nature Communications, 5,
  3950. https://doi.org/10.1038/ncomms4950
- Watanabe, E., Onodera, J., Harada, N., Aita, M. N., Ishida, A., & Kishi, M. J. (2015). Wind-driven
  interannual variability of sea ice algal production in the western Arctic Chukchi Borderland.
- 671 Biogeosciences, 12, 6147–6168, doi:10.5194/bg-12-6147-2015.
- Watanabe, E., Onodera, J., Itoh, M, Nishino, S., & Kikuchi, T. (2017). Winter transport of subsurface
  warm water toward the Arctic Chukchi Borderland. Deep-Sea Research I, 128, 115–130.
- Zhang, R., Chen, M., Guo, L., Gao, Z., Ma, Q., & et al. (2012). Variations in the isotopic
  composition of particulate organic carbon and their relation with carbon dynamics in the
  western Arctic Ocean. Deep-Sea Research II, 81–84, 72–78.
- 677

## 678 Table titles and figure captions

- Table 1. Deployment summary for sediment trap moorings. Time of sampled duration is 00:00 AM
- 680 (UTC). Trap depths are means of moored depths during deployment.
- Table 2. Moored equipment at Stations NAP10t-13t and CAP12t.
- Table 3. Major events in passing oceanic eddies estimated for current data of NAP13t (Kawaguchi,
  2017, Pers. Comm. for events #1–4).
- 684

Figure 1. Map of research area with schematic of major surface ocean currents based on Corlett and
Pickart (2017). Abbreviations for geographic features follow. CAP: Chukchi Abyssal Plain, CS:
Chukchi Spur, NAP: Northwind Abyssal Plain, and NWR: Northwind Ridge. Bathymetry data

are from the International Bathymetric Chart of the Arctic Ocean (Jakobsson et al., 2012).

Figure 2. Climate Forecast System Reanalysis (CFSR) reanalysis data (Saha et al., 2010) for
downward shortwave radiation at sea surface (or on snow/sea ice) and sea ice concentration at
NAP and CAP Stations for the entire deployment period, with measured ice thickness for the
deployment period for NAP13t. (a) Shortwave radiation, (b) sea ice concentration (line), and
sea ice thickness (shade) for NAP13t.

Figure 3. Progressive daily current vector at depths of 50 m, 100 m, ~170 m, and ~210 m at NAP13t
from September 10, 2013 (starting point), to September 16, 2014 (respective ending points).
The dots on each line represent the first day of each month, from October 2013 to September
2014.

Figure 4. Daily means for hydrographic data from NAP13t and CAP12t. (a)–(c) Temperature and
salinity with moored depth of CT sensors of NAP13t. Color of depth curve shows water mass
group in the Chukchi Sea (Corlett and Pickart, 2017). (d) East–west current speed and (e)
north–south current speed. Arrows represent the period of major event on oceanic eddy
detection. (f) Temperature and salinity at top buoy (~90 m depth) of CAP12t. Color bar shows
water mass group (Corlett and Pickart, 2017).

Figure 5. Time series fluctuation of moored sediment trap depths and settling particle fluxes at
 NAP10t-13t. (a) and (b) daily means of moored sediment trap depths and (c) TMF for shallower
 and deeper sediment traps.

- Figure 6. Time series fluctuation of particulate organic carbon (POC) and particulate nitrogen (PN)
  at NAP10t-13t. (a) Daily settling flux for POC, (b) daily settling flux for PN, (c) mole ratio for
  C/N, (d) stable carbon isotope ratio for POC, and (e) stable nitrogen isotope ratio for PN. The
  shallow trap data are shown in gray, and those of deep trap are in black. The short bars under
  (a) show periods of temporal mass occurrence of gelatinous matters (Fig. 7).
- Figure 7. The photograph of recovered sediment trap samples containing abundant gelatinous
  matters (mainly Appendicularian house) from NAP13t in July 30 August 12, 2014. (a) the
  sample of shallow trap, and (b) the sample of deeper trap. The scale bar = 1cm.
- Figure 8. Daily means of turbidity, signal strengths of ADCP, and TMF at NAP13t. (a) Turbidity at
  depths of 25 m and 170 m: note unit differences. (b) Acoustic signal strength of ADCP WH300,
  (c) acoustic signal strength of ADCP deployed at depths of ~170 m, and (d) TMF in logarithm.
- 718 Black bar at top of panel indicates that current speed exceeded 20 cm/s at 170 m depth. The
- 719 larger dB of acoustic backscattered intensity shows relatively abundant particle concentration.
- Figure 9. Time series fluctuations of settling particle flux at ~265 m depth for CAP12t. (a) TMF, (b)
- 39
- 40

721	POC flux, (c) PN flux, (d) C/N ratio, (e) stable carbon isotope ratio for POC, (f) stable nitrogen
722	isotope ratio for PN, (g) PIC flux, and (h) biogenic opal flux. The short bar under (b) shows
723	period of temporal mass occurrence of gelatinous matters (Fig. 7).

725	Supplementary Table 1. The dataset used in this study. There are six spread sheets including used
726	data. It will be archived in the Arctic Data archive System (ADS, https://ads.nipr.ac.jp/).



Figure 1.



Figure 2.



Figure 3.



Figure 4.



Figure 5.



Figure 6.



Figure 7.



Figure 8.



Figure 9.