

# Swarm-derived indices of geomagnetic activity

Constantinos Papadimitriou<sup>1</sup>, Georgios Balasis<sup>1</sup>, Adamantia Zoe Boutsis<sup>1</sup>, Alexandra Antonopoulou<sup>1</sup>, Georgia Moutsiana<sup>2</sup>, Ioannis A. Daglis<sup>2</sup>, Omiros Giannakis<sup>1</sup>, Paola De Michelis<sup>3</sup>, Giuseppe Consolini<sup>4</sup>, Jesper W. Gjerloev<sup>5</sup>, and Lorenzo Trenchi<sup>6</sup>

<sup>1</sup>National Observatory of Athens

<sup>2</sup>National and Kapodistrian University of Athens

<sup>3</sup>Istituto Nazionale di Geofisica e Vulcanologia

<sup>4</sup>Istituto Nazionale di Astrofisica

<sup>5</sup>APL-JHU

<sup>6</sup>Serco for ESA - ESRIN, Directorate of Earth Observation Programmes

November 22, 2022

## Abstract

Ground-based indices, such as the Dst, ap and AE, have been used for decades to describe the interplay of the terrestrial magnetosphere with the solar wind and provide quantifiable indications of the state of geomagnetic activity in general. These indices have been traditionally derived from ground-based observations from magnetometer stations all around the Earth. In the last 7 years though, the highly successful satellite mission Swarm has provided the scientific community with an abundance of high quality magnetic measurements at Low Earth Orbit (LEO), which can be used to produce the space-based counterparts of these indices, such the Swarm-Dst, Swarm-ap and Swarm-AE indices. In this work, we present the first results from this endeavour, with comparisons against traditionally used parameters. We postulate on the possible usefulness of these Swarm-based products for a more accurate monitoring of the dynamics of the magnetosphere and thus, for providing a better diagnosis of space weather conditions.

# Swarm-derived indices of geomagnetic activity

Constantinos Papadimitriou<sup>1,2,3</sup>, Georgios Balasis<sup>1</sup>, Adamantia Zoe Boutsis<sup>1,2</sup>,  
Alexandra Antonopoulou<sup>1</sup>, Georgia Moutsiana<sup>2</sup>, Ioannis A. Daglis<sup>2,4,1</sup>, Omiros  
Giannakis<sup>1</sup>, Paola de Michelis<sup>5</sup>, Giuseppe Consolini<sup>6</sup>, Jesper Gjerloev<sup>7</sup>,  
Lorenzo Trenchi<sup>8</sup>

<sup>1</sup>IAASARS National Observatory of Athens, Athens, Greece

<sup>2</sup>Department of Physics, National & Kapodistrian University of Athens, Athens, Greece

<sup>3</sup>Space Applications & Research Consultancy, SPARC G.P., Athens, Greece

<sup>4</sup>Hellenic Space Center, Athens, Greece

<sup>5</sup>Istituto Nazionale di Geofisica e Vulcanologia, Rome, Italy

<sup>6</sup>INAF Istituto di Astrofisica e Planetologia Spaziali, Rome, Italy

<sup>7</sup>Johns Hopkins University Applied Physics Laboratory, USA

<sup>8</sup>ESRIN, European Space Agency, Italy

## Key Points:

- New geomagnetic activity indices based on Swarm magnetic field data are computed similar to standard ground-based indices of Dst, ap and AE
- Swarm-derived indices show excellent correlations with both the traditional and SuperMAG-derived indices
- Swarm-based AE index enable us to monitor substorm activity also at the southern hemisphere

## Abstract

Ground-based indices, such as the Dst, ap and AE, have been used for decades to describe the interplay of the terrestrial magnetosphere with the solar wind and provide quantifiable indications of the state of geomagnetic activity in general. These indices have been traditionally derived from ground-based observations from magnetometer stations all around the Earth. In the last 7 years though, the highly successful satellite mission Swarm has provided the scientific community with an abundance of high quality magnetic measurements at Low Earth Orbit (LEO), which can be used to produce the space-based counterparts of these indices, such the Swarm-Dst, Swarm-ap and Swarm-AE indices. In this work, we present the first results from this endeavour, with comparisons against traditionally used parameters. We postulate on the possible usefulness of these Swarm-based products for a more accurate monitoring of the dynamics of the magnetosphere and thus, for providing a better diagnosis of space weather conditions.

## Plain Language Summary

Ground-based geomagnetic activity indices, like Dst, ap and AE, have been used for decades to monitor the dynamics of the Earth's magnetosphere, and provide information on two major types of space weather phenomena, i.e., magnetic storm and magnetospheric substorm occurrence and intensity. This study demonstrates how magnetic field data from a Low Earth Orbit (LEO) satellite mission, like ESA's Swarm constellation, can be used to derive corresponding space-based geomagnetic activity indices. The comparison of Swarm-based with ground-based indices shows a very good agreement, indicating that Swarm magnetic field data can be used to provide new satellite-based global indices to monitor the level of geomagnetic activity. Given the fact that the official AE is constructed by data from 12 ground stations solely in the northern hemisphere and the official ap from only 2 stations in the southern hemisphere (and 11 in the northern one), it can be said that both of these indices are predominantly northern, while the Swarm-derived AE and ap indices may be more representative of a global state, since they are based on measurements from both hemispheres.

## 1 Introduction

The magnetosphere is a highly complex system of fields and currents that envelop the Earth and interact with each other producing a wide range of phenomena. One particular current system that holds special importance is the aptly named ring current. The ring current is a toroidal electric current flowing around the Earth, formed by the azimuthal motion of electrons and ions, extending from 3 to 8 Earth radii on the magnetospheric equatorial plane (Daglis et al., 1999). Because of its shape and direction, it forms its own magnetic field component, with an axis almost parallel to that of the Earth's dipole and the same polarity (southward). A direct consequence of this is that on the surface of our planet the ring currents induced magnetic field is opposite to the Earth's magnetic field. Thus, in cases where incoming current systems of the solar wind have the appropriate properties to cause particle injection in the inner magnetosphere and enhance the ring current, the terrestrial field on the surface of the planet will exhibit a decrease due to the increase of the counteracting ring current field. This is the phenomenon known as geomagnetic (or simply magnetic) storm, which seriously affects the electromagnetic environment of the Earth and can cause a wide range of significant effects ranging from telecommunication issues and satellite failures to the induction of high voltage electrical currents on electrical wires and conducting materials on the surface of the planet (Bothmer and Daglis, 2007). The monitoring of geomagnetic storms has been traditionally accomplished by means of the Disturbance storm-time (Dst) index, which represents the axially symmetric disturbance of the horizontal component of the magnetic field at the magnetic equator on the Earth's surface (Sugiura and Kamei, 1981). As such, it oper-

71 ates as a proxy for the enhancement and subsequent weakening of the ring current and  
 72 hence the onset and evolution of magnetic storms.


73 Another group of current systems that cause an array of spectacular phenomena  
 74 are the auroral electrojets, which are currents that flow in concentrated channels of high  
 75 conductivity in the Earth's ionosphere and are carried by particles that generate the au-  
 76 roral light, moving both eastward (therefore forming the East ElectroJet EEJ) and west-  
 77 ward (therefore forming the West ElectroJet WEJ). Related to these systems are the  
 78 disturbances known as magnetospheric substorms (Akasofu, 1964; McPherron, 1979),  
 79 which are collective phenomena, considered as one of the major ways for the discharge  
 80 of accumulated energy in the terrestrial magnetosphere (Chian and Kamide, 2007). Dur-  
 81 ing the onset of substorm expansion phase, a dynamical process in the near Earth mag-  
 82 netosphere causes cross tail current to be diverted into the ionosphere, forming a sub-  
 83 storm current wedge consisting of downward (upward) field aligned currents on the dawn-  
 84 side (duskside) of the wedge and a westward auroral electrojet in the ionosphere (Kepko  
 85 at al., 2015). The Auroral Electrojet index AE (Davis and Sugiura, 1966) actually mea-  
 86 sures the intensity of this substorm enhanced westward ionospheric electrojet via its dia-  
 87 magnetic result on the horizontal component of the terrestrial magnetic field.

88 Balasis et al. (2019) recently derived a 1 Hz Swarm Dst-like index based solely on  
 89 magnetic data from the Swarm mission. The scientific merit of such a geomagnetic ac-  
 90 tivity index relies on its global character (since the three satellites provide a global Earth  
 91 coverage both latitudinally and longitudinally), as well as their proximity to the region  
 92 of emergence and activity of ionospheric currents. The present work further expands upon  
 93 that first effort, by simplifying and at the same time generalizing the approach so that  
 94 all three most commonly used indices of geomagnetic activity can be derived, namely  
 95 Dst, ap (or Kp) and AE. This article is divided upon 6 sections. Following the Introduc-  
 96 tion, Section 2 presents the main steps of analysis and pre-processing of Swarm data that  
 97 must be taken before the method is applied, as well as the argumentation on the selec-  
 98 tion of an appropriate time interval for the demonstration of the effectiveness of the meth-  
 99 ods. Section 3 outlines the new process for the derivation of the Swarm-Dst index and  
 100 is complemented by a comparison against the standard Dst index. The same line of think-  
 101 ing is followed in Sections 4 and 5, correspondingly for the ap and AE indices, while Sec-  
 102 tion 6 summarizes the findings and offers some general remarks on the usefulness of the  
 103 method.

## 104 **2 Data Selection and Pre-processing**

105 Our explorers in this quest for an ever more detailed description of the Earth's mag-  
 106 netic field are the satellites of the Swarm constellation (Friis-Christensen et al., 2006).  
 107 Launched on 23 November 2013, the mission is composed of three satellites in polar Low-  
 108 Earth Orbit (LEO), with two of them (Swarm-A and Swarm-C) flying side by side at  
 109 an initial altitude of 460 km and the third (Swarm-B) flying slightly higher at 510 km.  
 110 Carrying highly precise instruments, the mission offers the most up-to-date survey of the  
 111 terrestrial magnetic field (De Michelis et al., 2015; Hulot et al., 2015; Olsen et al., 2015),  
 112 but also of the general near-Earth electromagnetic environment and its interaction with  
 113 the solar wind (Balasis et al., 2015; Olsen et al., 2016; Papadimitriou et al., 2018).

114 The derivation of the Swarm indices is based on the dataset prepared within the  
 115 framework of the INTENS (Characterization of IoNospheric TurbulENCE level by Swarm  
 116 constellation) project. The dataset was constructed from the Level 1b, MAG\_LR prod-  
 117 uct of the Swarm mission (<https://swarm-diss.eo.esa.int/>), which contains among oth-  
 118 ers, magnetic field measurements from the Vector Field Magnetometer (VFM) instru-  
 119 ment (Tffner-Clausen et al., 2016) on board the three satellites at a time sampling of 1  
 120 second. The magnetic field vector was mapped to Mean Field Aligned (MFA) coordi-  
 121 nates, i.e. one component parallel to the mean field  $B_{par}$  and two perpendicular com-



**Figure 1.** Number of measurements performed by the Swarm satellites in each MLT bin. Top panel shows Swarm-A (Swarm-C is identical) and bottom panel Swarm-B.

122 ponents  $B_{per1}$  and  $B_{per2}$  and the static, background field was subtracted by removing  
 123 the internal mode of the CHAOS-6 model (Finlay et al., 2016) from the data, which con-  
 124 sists of the core and crustal magnetic field contributions of the Earth. Obvious outliers,  
 125 i.e. isolated spikes, were removed by applying a simple threshold and as a final step, the  
 126 magnitude of the residual vector was computed and saved for further processing.

127 The time period that we selected for the illustration of the method was the year  
 128 of 2015, which is appropriate to test the Swarm indices during both quiet and disturbed  
 129 geomagnetic conditions. Indeed, 2015 was characterized by two months of intense ge-  
 130 omagnetic activity due to the storms of March 17<sup>th</sup> and June 23<sup>rd</sup>, two events that have  
 131 been extensively discussed by the scientific community (Liu et al., 2015; Kataoka et al.,  
 132 2015; Wang et al., 2016; Wu et al., 2016; Marubashi et al., 2016; Balasis et al., 2018; De  
 133 Michelis et al., 2020), as well as a third event of less intensity, yet still an important one,  
 134 the storm of the 22<sup>nd</sup> of December. Additionally, several substorms erupted between and  
 135 following the storm events, which had shorter duration and thus were interspersed by  
 136 several quiet intervals as well. As such, this period provides a blend of both storm and  
 137 substorm activity, while also exhibiting intervals of quiescence and is perfectly suited for  
 138 modelling all levels of geomagnetic activity.

139 A duration that is a multiple of a four-month interval is also important for reasons  
 140 of geographical coverage, as four months is the time it takes for the orbital plane of the  
 141 Swarm satellites to complete a  $180^\circ$  turn, with respect to the Sun-Earth line. Since the  
 142 satellites fly one half of their orbit on the day-side and the other half on the night-side,  
 143 a  $180^\circ$  rotation provides full local time (LT) coverage. The time spent by the Swarm satel-  
 144 lites in each bin of Magnetic Local Time (MLT) is shown in Figure 1 (only one of the  
 145 satellites of the lower pair is shown, Swarm-A and not Swarm-C, since their orbits are  
 146 almost identical).

### 147 3 The Swarm-derived Dst Index

148 Magnetic storms produce global magnetic disturbances on the Earth’s surface, which  
 149 serve as the basis for storm monitoring via the hourly Dst index. It is derived from the  
 150 variations of the horizontal component of the terrestrial magnetic field, using data from

151 four observatories, positioned at magnetic latitudes ranging from approximately  $-30^\circ$  to  
 152  $+30^\circ$ .

153 In order to mirror the behaviour of the ground station data, it is imperative to re-  
 154 move measurements that were performed when the satellites were at high latitudes. Thus,  
 155 the first step in the process is to discard times where the satellites were above  $+30^\circ$  or  
 156 below  $-30^\circ$  in magnetic latitude. At those near-equatorial latitudes, a good proxy for the  
 157 horizontal component of the terrestrial magnetic field is the  $B_{par}$ , so the process con-  
 158 tinues keeping only this component and ignoring the others. Then, a non-overlapping,  
 159 moving average scheme is applied on the time series, with a window of 60 seconds, so  
 160 that the series are now set to a 1-minute time resolution. Up to this point, the analy-  
 161 sis was performed separately for each of the satellites Swarm-A and Swarm-B (Swarm-  
 162 C yields the same results as A so it was not used). In the next step the two series are  
 163 merged, in a joint 1-min resolution dataset, so that if both satellites are concurrently within  
 164 the latitude limits their values are averaged, otherwise we keep only the ones from the  
 165 satellite that was within the prescribed latitudes. Even so, there are still many cases for  
 166 which all satellites are at high latitudes, so these gaps are interpolated by a simple linear  
 167 scheme, to produce a complete time series. Then, a low-pass, Chebyshev Type I fil-  
 168 ter is used, with a cutoff period of 4 hours, to filter out some of the small perturbations  
 169 in the signal that arise from the fast motion of the satellites. Finally, we apply a linear  
 170 transform of the form:


$$171 \quad \mathbf{S}_{Dst} = 1.53\mathbf{B}_f + 12.85$$

172 where  $B_f$  is the filtered series acquired in the last step, to get the final Swarm-Dst  
 173 index  $S_{Dst}$ . A flowchart of the method, showing the form of the time series at each in-  
 174 termediate step described here is shown in Figure 2. A comparison of the final Swarm  
 175 index with the traditionally used one is shown in Figure 3, where it is evident how this  
 176 method can produce an index that is strikingly similar to the traditionally used one, a  
 177 similarity that is justified by a Pearson correlation coefficient (Pearson, 1895) of 0.94 be-  
 178 tween the two. It should be noted here that to facilitate comparisons, the SYM-H in-  
 179 dex was used instead of the Dst, since it is provided with a 1-minute sampling time, in-  
 180 stead of the hourly series of the official Dst index, but the good agreement holds also with  
 181 the official Dst, since the two indices are practically identical (Iyemori et al., 2010).


182 The four parameters of the method are the latitudinal limit, the filtering cutoff and  
 183 the two parameters of the linear transform. Even though their choice is intuitive in this  
 184 case, a more formal derivation is achieved by following two simple steps. First, many dif-  
 185 ferent combinations for the latitudes and filtering cutoff are tested, trying to find the one  
 186 that maximizes the correlation with the ground-based index series and secondly, keep-  
 187 ing the first two constant, we find the linear transform parameters that minimize the root  
 188 mean square error between the filtered series and the standard index.

## 189 4 The Swarm-derived ap Index


190 The geomagnetic three-hourly Kp index was introduced by J. Bartels in 1949 and  
 191 is derived from 13 magnetic observatories, 11 of which in the Northern Hemisphere, be-  
 192 tween latitudes from  $+38^\circ$  up to  $+60^\circ$  and 2 in the Southern Hemisphere at latitudes  
 193 of  $-43^\circ$  and  $-35^\circ$  [<https://www.gfz-potsdam.de/en/kp-index/>]. It is designed to measure  
 194 solar particle radiation by its magnetic effects and today it is considered a proxy for the  
 195 energy input from the solar wind to Earth. The name Kp was derived from the German  
 196 words “planetarische Kennziffer meaning “planetary index. The index is quasi-logarithmic  
 197 in nature and since this is more difficult to assign to actual magnetic field measurements,  
 198 Bartels also proposed a correspondence between Kp values and the more linear ap in-  
 199 dex values. Due to this, we will also use the ap index for this study, keeping in mind that  
 200 a simple one-to-one relation can map the ap values to Kp and vice-versa.



**Figure 2.** Flowchart for the derivation of the Swarm-Dst index. Each of the 3 upper rows includes distinct panels for Swarm-A (blue) and Swarm-B (red) time series, while each of the next 2 rows shows a single panel of combined Swarm-A and Swarm-B time series.




**Figure 3.** Swarm-derived Dst index (red) compared to the ground-based index (black).



**Figure 4.** Swarm-derived ap index (red) compared to the standard index (black).

201 We repeat the exact same process as above, tweaking the parameters until the best  
 202 possible outcome emerges. For the Swarm-ap index we use measurements of the mag-  
 203 nitude of the residual field vector, located in a narrow band between  $+55^\circ$  and  $+60^\circ$   
 204 in the Northern Hemisphere and correspondingly  $-60^\circ$  to  $-55^\circ$  in the Southern one. The cut-  
 205 off period is set to 9 hours and the linear transform to  $S_{ap} = 0.39B_f - 4.29$  in order  
 206 to achieve a maximum correlation coefficient of 0.86. The result is shown in Figure 4.  
 207 As can be seen the Swarm index closely follows the standard ap index, missing only in  
 208 absolute magnitude a few of the highest peaks. This hints at the possibility of a non-  
 209 linear transform which would promote high values more significantly than the lower ones,  
 210 something which makes sense given the peculiar way with which these indices are pro-  
 211 duced.



**Figure 5.** Swarm-derived AE index (red) compared to the standard AE index (black). Top panel shows the entire 12 month period from Jan 1<sup>st</sup> to Dec 31<sup>st</sup> 2015, while the two bottom panels show zoomed pictures from the second half of April (bottom left) and first half of May (bottom right).

212

## 5 The Swarm-derived AE Index

213

214

215

216

217

218

219

220

221

222

223

224

The AE index represents the overall activity of the electrojets. The AE index is derived from geomagnetic variations in the horizontal component observed at selected observatories along the auroral zone, solely in the Northern Hemisphere, at magnetic latitudes between  $+60^\circ$  and  $+70^\circ$  [<http://wdc.kugi.kyoto-u.ac.jp/aedir/ae2/onAEindex.html>]. To mirror this we follow the same process outlined above, again for the magnitude of the residual field, keeping only measurements between  $+65^\circ$  and  $+75^\circ$  (and correspondingly  $-75^\circ$  to  $-65^\circ$ ) in magnetic latitude, filtering with a cutoff period of 2.6 hours and applying the transform  $S_{AE} = 2.2B_f - 8.9$  to achieve a maximum correlation coefficient of 0.85. As can be seen in Figure 5, while the Swarm index doesn't match the peaks during the intense storm events (again hinting at a possible non-linear relation), it captures with high accuracy the substorm activity during April and May, highlighted with the two bottom panels which zoom in two characteristic intervals.

225

226

227

228

229

230

231

232

233

234

235


236

237

238

239

One of the most important benefits of using a satellite observatory is that by slightly changing the formulation of the method, one can easily produce more localized versions of these indices. As an example, by imposing the latitudinal limits to only maintain measurements from the north or the south hemisphere, while keeping the rest of the method unchanged, we can construct the Swarm-AE-North and Swarm-AE-South indices, to complement the full version of Swarm-AE. This is shown in Figure 6, where the full Swarm-AE index is overplotted by its localized North and South hemisphere counterparts. Even though the two localized indices agree, in general, with each other, if we draw our attention to specific, small intervals, like the ones depicted on the two bottom subplots of Figure 6, one can see small, but important differences between the two. This comes as a verification of recent literature (Liou et al., 2018) in which they report that substorm onset is far from north-south symmetric, as was previously considered, and that it is more likely to be initiated in a dark than a sunlit oval. They additionally showed that the preferred locations of substorm onsets coincide with the local peak of the Earth's magnetic field (or a minimum in the ionospheric conductivity), a finding which is consistent with



**Figure 6.** Swarm-derived AE index (grey) compared to its regional counterparts the Swarm-AE-North (blue) and Swarm-AE-South (red) indices. Top panel shows the entire 12 month period from Jan 1<sup>st</sup> to Dec 31<sup>st</sup> 2015, while the two bottom panels show zoomed pictures from the second half of April (bottom left) and first half of May (bottom right).

240 an ionospheric feedback mechanism. Similar asymmetries were also reported by Wey-  
 241 gand et al. (2014) based on magnetic field data from ground based observatories.


## 242 6 Comparisons against SuperMAG Indices

243 There has been a huge effort in recent years to complement the traditional indices  
 244 of geomagnetic activity with new versions, utilizing data from the immense ground mag-  
 245 netometer network SuperMAG. SuperMAG is a worldwide collaboration of organizations  
 246 and national agencies which currently operate more than 300 ground-based magnetome-  
 247 ters spread across the globe (Gjerloev, 2009). Data from each of these stations are pro-  
 248 cessed according to the same guidelines (Gjerloev, 2012) and utilized, according to their  
 249 latitudinal location to produce the SuperMAG geomagnetic activity indices: the SMR  
 250 index, which is a ring current index, similar to Dst or SYM-H (Newell et al., 2012) and  
 251 the SME index, which attempts to capture the auroral electrojet behavior and thus op-  
 252 erate in a manner analogous to the AE index (Newell et al., 2011). As a first step to in-  
 253 corporate these new indices in our work, we performed two simple comparisons, calcu-  
 254 lating the correlation coefficient between the Swarm-Dst and the SuperMAG SMR in-  
 255 dex, which yielded a value of 0.95 and also compared Swarm-AE (the full version) against  
 256 SuperMAG SME, which gave the result 0.86. Both of these values are slightly higher than  
 257 the ones procured by comparisons against the traditional SYM-H and AE indices, which  
 258 indicates that the satellite indices are closer to the new indices, which are more detailed  
 259 and are being produced by many more observatories than the old ones, although further  
 260 work is needed to address these differences properly. That being said, this is a very promis-  
 261 ing result that we believe further justifies our approach and opens new avenues for ex-  
 262 ploration.

263 These results, along with all the previous ones, are presented in Table 1, while a  
 264 pictorial representation of this comparison is shown in Figure 7.

**Table 1.** Summary of results

Correlations with ring current indices	
Swarm-Dst vs SYM-H	0.94
Swarm-Dst vs SMR	0.95
Correlations with ap index	
Swarm-ap vs ap	0.86
Correlations with auroral electrojet indices	
Swarm-AE vs AE	0.85
Swarm-AE vs SME	0.86
Swarm-AE-North vs AE	0.81
Swarm-AE-South vs AE	0.77



**Figure 7.** Comparisons of SuperMAG indices (with black) SMR (top panel) and SME (bottom panel) against their Swarm-derived counterparts (with red) for the year 2015.

## 7 Discussion and Conclusions

This work shows how the magnetic field data from the Swarm mission can be utilized, by means of a simple and intuitive method to reproduce, with high accuracy, the three major indices of geomagnetic activity, namely the Dst, ap (or Kp) and AE indices. The global coverage provided by a constellation of low-Earth orbiting satellites makes them ideal for encapsulating the entirety of the magnetic field, discerning changes at larger spatial scales, while their altitude positions them right in the place of the ionospheric currents which are responsible for many of the effects that comprise our notion of space weather. We note that the Swarm AE-like index could, in principle, be more representative of a global state, since it is based on measurements from both hemispheres, while the ground-based one is computed from measurements in the Northern Hemisphere only.

Additionally, since the satellites remain at fairly constant LTs for several weeks, their data can further promote recent research on regional indices of electrojet or ring current activity, such as the regional versions of SME and SMR indices (Bergin et al., 2020). As such, satellite magnetic observatories can complement their ground-based counterparts, providing new insights into the state of the magnetosphere and new promise for more accurate diagnosis of space weather conditions.

### Acknowledgments

The results presented rely on data collected by the three satellites of the Swarm constellation. We thank the European Space Agency (ESA) that supports the Swarm mission. Swarm data can be accessed at <http://earth.esa.int/swarm>. We acknowledge the Kyoto World Data Center (WDC) for Geomagnetism and the observatories that produce and make Dst, ap and AE indices available at <http://wdc.kugi.kyoto-u.ac.jp/>. We gratefully acknowledge the SuperMAG collaborators for the provision of ground-based index data (<https://supermag.jhuapl.edu/info/?page=acknowledgement>). The authors acknowledge financial support from European Space Agency (ESA contract N. 4000125663/18/I-NB “EO 486 Science for Society Permanently Open Call for Proposals EOEP-5 BLOCK4” (INTENS)). This work was benefited from discussions within the International Space Science Institute (ISSI) Team # 455 “Complex Systems Perspectives Pertaining to the Research of the Near-Earth Electromagnetic Environment”.

### References

- Akasofu, S.-I. (1964). The development of the auroral substorms. *Planet. Space Sci.*, 12, 273.
- Balasis, G., Papadimitriou, C., Daglis, I. A., and Pilipenko, V. (2015). ULF wave power features in the topside ionosphere revealed by Swarm observations. *Geophys. Res. Lett.*, 42, 6922-6930, doi:10.1002/2015GL065424.
- Balasis, G., et al. (2018). Observation of intermittency-induced critical dynamics in geomagnetic field time series prior to the intense magnetic storms of March, June, and December 2015. *J. Geophys. Res.: Space Phys.*, 123, 4594-4613, doi:10.1002/2017JA025131.
- Balasis, G., Papadimitriou, C., and Boutsis, A. Z. (2019). Ionospheric response to solar and interplanetary disturbances: a Swarm perspective. *Philosophical Transactions A*, doi:10.1098/rsta.2018.0098.
- Bartels, J. (1949). The standardized index Ks and the planetary index Kp. *IATME Bull.* 12 (b), 97, IUGG Publ.
- Bothmer, V., and Daglis, I. A. (2007). In V. Bothmer and I. A. Daglis (Eds.), *Space Weather: Physics and Effects*, Praxis/Springer, New York, (438 pp.), ISBN 978-3-540-23907-9.
- Bergin, A., Chapman, S., and Gjerloev, J. (2020). AE, Dst and their SuperMAG Counterparts: The Effect of Improved Spatial Resolution in Geomagnetic

- 315 Indices. *J. Geophys. Res.: Space Physics*, 125, doi:10.1029/2020JA027828.
- 316 Chian, A. C.-L., and Kamide, Y. (2007). An Overview of the Solar-Terrestrial En-  
317 vironment. In Y. Kamide and A. C.-L. Chian (Eds.), *Handbook of the Solar-  
318 Terrestrial Environment*, Springer-Verlag, Berlin, Heidelberg, doi:10.1007/978-  
319 3-540-46315-3.
- 320 Daglis, I. A., Thorne, R. M., Baumjohann, W., and Orsini, S. (1999). The terres-  
321 trial ring current: Origin, formation, and decay. *Rev. Geophys.*, 37(4), 407-438,  
322 doi:10.1029/1999RG900009.
- 323 Davis, T. N., and Sugiura, M. (1966). Auroral electrojet activity index AE  
324 and its universal time variations. *J. Geophys. Res.*, 71(3), 785801,  
325 doi:10.1029/JZ071i003p00785.
- 326 De Michelis, P., Consolini, G., and Tozzi, R. (2015). Magnetic field fluctuation fea-  
327 tures at Swarms altitude: A fractal approach. *J. Geophys. Res.*, 42, 31003105,  
328 doi:10.1002/2015GL063603.
- 329 De Michelis, P., Pignalberi, A., Consolini, G., Coco, I., Tozzi, R., and Pezzopane,  
330 M., et al. (2020). On the 2015 St. Patrick's storm turbulent state of the iono-  
331 sphere: Hints from the Swarm mission. *Journal of Geophysical Research: Space  
332 Physics*, 125, e2020JA027934. <https://doi.org/10.1029/2020JA027934>
- 333 Finlay, C., et al. (2016). Recent geomagnetic secular variation from Swarm and  
334 ground observatories as estimated in the CHAOS-6 geomagnetic field model.  
335 *Earth Planets and Space*, 68, 112, doi:10.1186/s40623-016-0486-1.
- 336 Friis-Christensen, E., Lhr, H., and Hulot, G. (2006). Swarm: A constellation  
337 to study the Earth's magnetic field. *Earth Planets Space*, 58, 351358,  
338 doi:10.1186/BF03351933.
- 339 Gjerloev, J. W. (2009). A global ground-based magnetometer initiative. *Eos Trans.  
340 AGU*, 90 (27), 230231, doi:10.1029/2009EO270002.
- 341 Gjerloev, J. W. (2012). The SuperMAG data processing technique. *J. Geophys. Res.*,  
342 117, A09213, doi:10.1029/2012JA017683.
- 343 Hulot, G., Vigneron, P., Lger, J.-M., Fratter, I., Olsen, N., Jager, T., Bertrand,  
344 F., Brocco, L., Sirol, O., Lalanne, X., Boness, A., and Cattin, V. (2015).  
345 Swarms absolute magnetometer experimental vector mode, an innovative  
346 capability for space magnetometry. *Geophys. Res. Lett.*, 42, 1352-1359,  
347 doi:10.1002/2014GL062700.
- 348 Iyemori, T., Takeda, M., Nose, M., Odagi, Y., and Toh, H. (2010). Mid-latitude  
349 Geomagnetic Indices ASY and SYM for 2009 (Provisional). In *Internal Re-  
350 port of Data Analysis Center for Geomagnetism and Space Magnetism*, Kyoto  
351 University, Japan.
- 352 Kataoka, R., et al. (2015). Pileup accident hypothesis of magnetic storm on 17  
353 March 2015. *Geophys. Res. Lett.*, 42, 5155-5161, doi:10.1002/2015GL064816.
- 354 Kepko, L., McPherron, R. L., Amm, O., et al. (2015). Substorm Current Wedge  
355 Revisited. *Space Sci Rev*, 190, 1-46, doi:10.1007/s11214-014-0124-9.
- 356 Liou, K., Sotirelis, T., and Mitchell, E. J. (2018). North-South Asymmetry in the  
357 Geographic Location of Auroral Substorms correlated with Ionospheric Effects.  
358 *Sci Rep* 8, 17230, doi:10.1038/s41598-018-35091-2.
- 359 Liu, Y. D., et al. (2015). Plasma and magnetic field characteristics of solar cor-  
360 onal mass ejections in relation to geomagnetic storm intensity and variability.  
361 *Astrophys. J. Lett.*, 809, L34, doi:10.1088/2041-8205/809/2/L34.
- 362 Marubashi, K., et al. (2016). The 17 March 2015 storm: the associated magnetic  
363 flux rope structure and the storm development. *Earth Planets Space*, 68, 173,  
364 doi:10.1186/s40623-016-0551-9.
- 365 McPherron, R. I. (1979). Magnetospheric substorms. *Rev. Geophys.*, 17, 657.
- 366 Newell, P. T., and Gjerloev, J. W. (2011). Evaluation of SuperMAG auroral electro-  
367 jet indices as indicators of substorms and auroral power. *J. Geophys. Res.*, 116,  
368 A12211, doi:10.1029/2011JA016779.

- 369 Newell, P. T., and Gjerloev, J. W. (2012). SuperMAG-Based Partial Ring Current  
370 Indices. *J. Geophys. Res.*, 117, doi:10.1029/2012JA017586.
- 371 Olsen, N., Hulot, G., Lesur, V., Finlay, C. C., Beggan, C., Chulliat, A., Sabaka,  
372 T. J., Floberghagen, R., Friis-Christensen, E., Haagmans, R., Kotsiaros, S.,  
373 Lhr, H., Tffner-Clausen, L., and Vigneron, P. (2015). The Swarm Initial Field  
374 Model for the 2014 geomagnetic field. *Geophys. Res. Lett.*, 42, 1092-1098,  
375 doi:10.1002/2014GL062659.
- 376 Olsen, N., Stolle, C., Floberghagen, R., Hulot, G., and Kuvshinov, A. (2016). Special  
377 issue Swarm science results after 2 years in space. *Earth Planets Space*, 68, 13,  
378 doi:10.1186/s40623-016-0546-6.
- 379 Pearson, K. (1895). Notes on regression and inheritance in the case of two parents.  
380 *P. R. Soc. London*, 58, 240-242.
- 381 Papadimitriou, C., Balasis, G., Daglis, I. A., and Giannakis, O. (2018). An initial  
382 ULF wave index derived from two years of Swarm observations. *Ann. Geo-*  
383 *phys.*, 36, 287-299, doi:10.5194/angeo-36-287-2018.
- 384 Sugiura, M., and Kamei, T. (1981). IAGA Bulletin N°40, [http://wdc.kugi.kyoto-](http://wdc.kugi.kyoto-u.ac.jp/dstdir/dst2/onDstindex.html)  
385 [u.ac.jp/dstdir/dst2/onDstindex.html](http://wdc.kugi.kyoto-u.ac.jp/dstdir/dst2/onDstindex.html).
- 386 Tffner-Clausen, L., Lesur, V., Olsen, N., and Finlay, C. (2016). In-flight scalar cal-  
387 ibration and characterisation of the Swarm magnetometry package. *Earth,*  
388 *Planets and Space*, 68(1), 129, doi:10.1186/s40623-016-0501-6.
- 389 Wang, Y., et al. (2016). On the propagation of a geoeffective coronal mass  
390 ejection during 15–17 March 2015. *J. Geophys. Res.*, 121, 7423-7434,  
391 doi:10.1002/2016JA022924.
- 392 Weygand, J. M., Zesta, E., and Troshichev, O. (2014), Auroral electrojet indices in  
393 the Northern and Southern Hemispheres: A statistical comparison, *J. Geophys.*  
394 *Res. Space Physics*, 119, 4819-4840, doi:10.1002/2013JA019377.
- 395 Wu, C. C., et al. (2016). The first super geomagnetic storm of solar cycle 24:  
396 The St. Patrick's day event (17 March 2015). *Earth Planets Space*, 68, 151,  
397 doi:10.1186/s40623-016-0525-y.

**Figure1.**





Figure2.











Figure3.




Figure4.




Figure 5.










Figure6.



**Figure7.**

### Ring Current Indices



### Auroral Electrojet Indices

