Interpretable Machine Learning to Forecast SEP Events for Solar Cycle 23

Spiridon Kasapis¹, Lulu Zhao², Yang Chen¹, Xiantong Wang¹, Monica Bobra³, and Tamas I. I. Gombosi⁴

¹University of Michigan ²University of Alabama in Huntsville ³Stanford University ⁴University of Michigan-Ann Arbor

November 23, 2022

Abstract

We use machine learning methods to predict whether an active region (AR) which produces flares will lead to a solar energetic particle (SEP) event using Space-Weather Michelson Doppler Imager (MDI) Active Region Patches (SMARPs). This new data product is derived from maps of the solar surface magnetic field taken by the Michelson Doppler Imager (MDI) aboard the Solar and Heliospheric Observatory (SOHO). We survey the SMARP active regions associated with flares that appear on the solar disk between June 5, 1996 and August 14, 2010, label those that produced SEPs as positive and the rest as negative. The AR SMARP features that correspond to each flare are used to train two different types of machine learning methods, the support vector machines (SVMs) and the regression models. The results show that the SMARP data can predict whether a flare will lead to an SEP with accuracy (ACC) {less than or equal to}0.72{plus minus}0.12 while allowing for a competitive leading time of 55.3{plus minus}28.6 minutes for forecasting the SEP events.

Interpretable Machine Learning to Forecast SEP Events for Solar Cycle 23

Spiridon Kasapis¹, Lulu Zhao², Yang Chen³, Xiantong Wang², Monica Bobra⁴, Tamas Gombosi²

¹ Department of Naval Architecture and Marine Engineering, University of Michigan								
² Department of Climate and Space Sciences and Engineering, University of Michigan								
³ Department of Statistics, University of Michigan								
⁴ Hansen Experimental Physics Laboratory, Stanford University								

Key Points:

1

2

3

4

9

10	•	SMARP data can correctly predict whether a solar flare will lead to a solar en-
11		ergetic particle (SEP) event 72% of the times.
12	•	Flare peak intensity is the strongest SEP predictor and can be coupled with SMARP
13		data to achieve accuracy $\leq 0.92 \pm 0.07$.
14	•	The SMARP dataset provides a leading time of 55.3 ± 28.6 minutes for forecast-
15		ing the SEP events.

 $Corresponding \ author: \ Spiridon \ Kasapis, \verb+skasapis@umich.edu$

16 Abstract

We use machine learning methods to predict whether an active region (AR) which 17 produces flares will lead to a solar energetic particle (SEP) event using Space-Weather 18 Michelson Doppler Imager (MDI) Active Region Patches (SMARPs). This new data prod-19 uct is derived from maps of the solar surface magnetic field taken by the Michelson Doppler 20 Imager (MDI) aboard the Solar and Heliospheric Observatory (SOHO). We survey the 21 SMARP active regions associated with flares that appear on the solar disk between June 22 5, 1996 and August 14, 2010, label those that produced SEPs as positive and the rest 23 as negative. The AR SMARP features that correspond to each flare are used to train two different types of machine learning methods, the support vector machines (SVMs) 25 and the regression models. The results show that the SMARP data can predict whether 26 a flare will lead to an SEP with accuracy (ACC) $\leq 0.72 \pm 0.12$ while allowing for a com-27 petitive leading time of 55.3 ± 28.6 minutes for forecasting the SEP events. 28

²⁹ 1 Introduction

Large solar eruptions can potentially harm modern civilization in several different ways. Events such as large solar flares that lead to solar particle emissions, can adversely affect the near-earth environment and damage power grids, jam radio communications, incapacitate satellites, expose airline passengers to dangerous levels of radiation and even endanger life in outer space. Therefore, predicting and monitoring such events is an important task for the community.

Solar Energetic Particles are rare events that involve protons, electrons and heavy ions accelerated to high energies (up to tens of GeV while the fastest ones can accelerate to speeds of up to 80% of the speed of light) by two solar processes (Reames, 2013), the energization at a solar flare site or the shock waves associated with Coronal Mass Ejections (CMEs). Solar charged particles are accelerated in flares or CME shock waves (Wild et al., 1963) and travel preferentially along the interplanetary magnetic field to their detection point in space (McCracken & Ness, 1966).

The study of solar energetic particle (SEP) events is a relatively recent science as 43 the identification of the first solar proton event took place on 28 February, 1942 (For-44 bush, 1946). Observations of solar proton events (alternative name for SEP) were made 45 using ground-based instruments that detected ionization, neutrons, or radio disturbances 46 caused by them. The largest solar proton event recorded using these modern techniques 47 (particles exceeded 15 GeV at the top of the atmosphere) was on the 23rd of February, 48 1956. In the mid-1960s spacecraft was deployed that begun directly measuring solar pro-49 ton events. This was also the time when the first flare was associated with an SEP event 50 (Shea & Smart, 1995). 51

During the so-called Halloween storms in late October 2003, SEP events caused a number of power grid failures, 47 satellites reported malfunctions, more than 10 satellites were out of action for days, the Mars Odyssey spacecraft went into deep safe mode (Lopez et al., 2004), a Japanese satellite costing 640m USD was completely lost, the US FAA issued their first-ever high radiation dosage alert for high-altitude aircraft, and astronauts in the ISS had to seek safety into their heavily shielded service module (Webb & Allen, 2004; Horne et al., 2013).

One of the sources of solar activity phenomena that cause SEPs are the magnetically strong regions on the solar sphere that we refer to as active regions (van Driel-Gesztelyi & Green, 2015). The most flare productive active regions (ARs) are the ones that undergo large changes in sunspot area and show magnetic flux imbalance (Choudhary et al., 2013). Large active regions are also generally strong, flary, evolve rapidly and their lifetime spans from days to months (Choudhary et al., 2013). Using instruments carried onboard satellites such as the Michelson Doppler Imager (MDI) on the Solar and He-

liospheric Observatory (SOHO) or the Helioseismic and Magnetic Imager (HMI) on the

⁶⁷ Solar Dynamics Observatory (SDO), we are able to retrieve components of the magnetic

field at the solar surface, allowing us to calculate physical characteristics of the ARs (Scher-

⁶⁹ rer et al., 1995; Schou et al., 2012).

Solar particle prediction studies mainly use the flare and near-Earth space envi-70 ronment data to forecast SEP events given the knowledge that large SEPs are almost 71 always accompanied by a flare (Schrijver et al., 2012). Laurenza et al. (2009) used data 72 73 such as flare location, flare size, and evidence of particle acceleration/escape to provide short-term warnings for SEP events. Similarly, Núñez (2011) used the soft X-ray, dif-74 ferential and integral proton fluxes data to forecast the SEP events of Solar Cycle 23 recorded 75 on the NOAA/SWPC list. Although both flare and CME data are found to be useful 76 inputs to predictive models, García-Rigo et al. (2016) deemed sufficient to only use flare 77 properties as they noticed that the CME information offers insignificant increase in SEP 78 prediction accuracy. 79

Recently, machine learning (ML) methods like neural networks (in the multi-layer 80 perceptron implementation), random forests, decision trees, extremely randomized trees 81 and other, have been used in predicting SEP events. The preliminary results obtained 82 by Bain et al. (2018) show that machine learning classification techniques such as the 83 logistic regression (LR), decision trees (DTs) and support vector machine (SVM) algo-84 rithms give an improved forecasting skill over the current SWPC Proton Prediction Model 85 (Balch, 2008) based on physical parameters associated with solar flares and coronal mass 86 ejections. An even more comprehensive study that assesses the predictability of Solar 87 Energetic Particles using ML techniques was recently published by Lavasa et al. (2021). 88

Different studies have used a variety of sources to obtain the data necessary for so-89 lar particle event prediction. Richardson et al. (2018) predict the SEP events peak pro-90 ton intensity using the CME data in the Space Weather Database of Notifications, Knowl-91 edge, Information (DONKI). Papaioannou et al. (2016) have presented a catalogue which 92 includes proton flux data for 314 SEP events obtained from the Energetic Particle Sen-93 sor (EPS) aboard the Geostationary Operational Environmental Satellites (GOES; Ro-94 driguez et al., 2014) and CME data obtained by the Large Angle and Spectrometric Coro-95 nagraph (LASCO; Brueckner et al., 1995) carried onboard the SOHO spacecraft. Us-96 ing this information, Papaioannou et al. (2018) classify the solar energetic particle (SEP) 97 event radiation impact with respect to the characteristics of their parent solar events while 98 attempting to infer the possible prediction of SEP events. 99

Similarly, Anastasiadis et al. (2017) provide full-disk Helioseismic and Magnetic
 Imager (HMI) magnetograms to their novel integrated prediction system which nowcasts
 SEP events. The HMI instrument aboard the Solar Dynamics Laboratory (SDO) mea sures the solar surface magnetic field from which the Space-Weather HMI Active Region
 Patches (SHARPs) are derived. SHARPs have been used to identify flares or SEPs in
 Chen et al. (2019) and Inceoglu et al. (2018) respectively.

In this work, a new data product recently published by Bobra et al. (2021) called 106 Space-Weather MDI Active Region Patches (SMARPs) will be used to predict SEPs. SMARPs 107 are derived from the solar surface magnetic field taken by the Michelson Doppler Imager 108 (MDI) on the SOHO spacecraft and provide a continuous and seamless set of keywords 109 that describe every active region observed during Solar Cycle 23. The big difference be-110 tween the HMI (Schou et al., 2012) and the MDI (Scherrer et al., 1995) is that the first 111 measures the vector magnetic field at the solar surface whereas the later only measures 112 the line-of-sight component of the solar magnetic field. The main aim of this study is 113 to evaluate the predictive power of MDI Active Region Patches (SMARPs) on SEP events 114 as it is desirable for the space weather community to explore new datasets that, when 115

used on machine learning algorithms in the future, will be able to predict when solar pro-ton events will occur, how large they will be, and how long they will last.

118 **2** Database

In this work, we will evaluate the prediction power of the sun's magnetogram on 119 SEP events. In particular, we focus on whether an active region which is associated with 120 a solar flare will lead to an SEP event. To achieve this, five different predictors obtained 121 from the SMARP dataset (SMARP Predictors) are used, while two more predictors from 122 the NOAA solar X-ray flare dataset (Flare Predictors) are used for comparison. While 123 we are specifically interested in the responses of the ML models when only SMARP Pre-124 dictors are used, the ability to forecast SEPs by using flare data will serve as a baseline 125 capability. 126

127

2.1 SMARP Predictors

The magnetogram is measured by the Michelson Doppler Imager (MDI Scherrer 128 et al., 1995) onboard SOHO between June 5, 1996 and August 14, 2010. Based on the 129 magnetogram, Bobra et al. (2021) derived a new database called Space-Weather MDI 130 Active Region Patches (SMARPs), which contains characteristics of the active regions 131 on the solar surface. A Tracked Active Region Patch (TARP) Number is assigned to each 132 active region as its identification number and a NOAA active region number, if avail-133 able, is assigned to each active region patch. Three physical keywords, total unsigned 134 flux (USFLUXL), mean gradient of the vertical field (MEANGBL), and the logarithm 135 of the total unsigned flux near polarity inversion line (RVALUE) are calculated using the 136 pixels in the active region and stored in the SMARP header file. In addition, the SMARP 137 header file also contains four spacial features specifying the location of the correspond-138 ing AR on the solar surface: the minimum and maximum latitude (LATDMIN, LAT-139 DMAX) and the minimum and maximum longitude (LONDTMIN, LONDTMAX). The 140 SMARP data is available on the Joint Science Operations Center database (Mumford 141 et al., 2015; Barnes et al., 2020). 142

Besides the three physical keywords stored in the SMARP header file, we calcu-143 late the angular distance between the AR and the magnetic foot-point of the earth. The 144 longitude and latitude location of the active region on the sun is approximated by the 145 geometric center of the active region using the latitude and longitude keywords. The mag-146 netic foot-point of the earth on the sun is assumed to be at $W45^{\circ}$. Note that the mag-147 netic foot-point varies from event to event. One way of characterizing this variability is 148 to calculate the magnetic foot-point location using the solar wind speed measured at 1 149 AU assuming an ideal Parker spiral up to the solar source surface and reconstruct the 150 coronal magnetic fields using potential field source surface model. However, interplan-151 etary magnetic field can also be disturbed by corotating interaction regions (CIRs), in-152 terplanetary coronal mass ejections (ICMEs) and other solar transient events, especially 153 in solar maximum. In this work, for simplicity, we use $W45^{\circ}$ as an approximation. We 154 also calculate the size of the active region by multiplying the difference of longitude by 155 the difference of latitude. 156

157

2.2 Construction of SEP Event List

The SEP event list we use in this work is documented in the NOAA Space Environment Service Center website. For each SEP event, a solar flare and the corresponding NOAA active region number is assigned if exists. The solar flare list is obtained from the NOAA Solar Flare Data website. For each solar flare, the list contains the start, peak and end times, the peak intensity of the flare, the active region location and the corresponding NOAA active region number. We match the solar flare list with the SMARP database using the AR numbers. If a flare does not have a registered AR number, matching based on their occurrence time and spatial coordinates is performed. We also discard those solar flares whose AR numbers are undefined or missing. Out of the ~ 25,000 flares (A, B, C, M, X) recorded during the 14 year span between 1996 and 2010, 6,510 flares are matched with SMARP files.

During this 14 year span, 93 SEP events are detected by the GOES spacecraft. Miss-169 ing information about the SEP's associated flare or AR such as the Location and its Im-170 portance (Xray/Opt), leave only 70 SEPs with information adequate to label the 6,510 171 172 flares. We assign a label to each flare: Positive if it led to an SEP and Negative if it did not. An additional 5 SEP-flare couples were discarded due to missing physical feature 173 data about their corresponding SMARP Active Region. Therefore, the dataset used for 174 training has a Positive and a Negative component comprised of 65 and 6,510 flares re-175 spectively, making it vastly unbalanced. 176



Figure 1. Histograms for the time difference between the flare peak time and the selected SMARP data (selected row in the TARP file). The distributions range between 10 and 100 minutes. The mean time differences shown in the error bars above the graphs are 55.3 and 53.6 minutes with a standard deviation of 28.6 and 24.8 minutes for the Positive (green) and Negative (red) datasets respectively.

The SMARP header files contains rows with the physical and spatial features of 177 each active region at a 96-minute cadence throughout its entire lifetime, starting two days 178 before it emerges or rotates onto the solar disk until two days after it submerges or dis-179 appears from view behind the limb (Bobra et al., 2021). We select the SMARP header 180 file row at the time right before the flare peak time. In Figure 1, a histogram of the time 181 difference between the selected SMARP file row and the flare peak time is plotted with 182 the left panel corresponding to the positive dataset and right panel corresponding to the 183 negative dataset. The distributions range between 10 and 100 minutes. The mean time 184 difference is 55.3 and 53.6 minutes with a standard deviation of 28.6 and 24.8 minutes 185 for the Positive and Negative dataset respectively. 186

2.3 Flare Predictors

187

We will evaluate the prediction power of SMARP dataset on SEP events by comparing the prediction results with those obtained by only using the flare information, i.e. flare intensity and flare location. We use the solar X-ray flare data that NOAA's Geostationary Operational Environmental Satellites (GOES) continuously provides since 1975. Similarly to the SMARP Predictors, we calculate the flare angular distance from the earth's magnetic foot-point location, W45°, on the sun.

¹⁹⁴ **3** Preliminary Data Analysis

We conduct preliminary analysis/assessment of different predictors, i.e. the SMARP and the Flare Predictors, via comparing the histogram of each predictor for the positive with that of negative samples. Figure 2 shows the density histograms of each predictor from the SMARP dataset on the top and from the GOES flare information on the bottom. The positive data is shown in green and negative data in red.



Figure 2. The probability density values are given in the histograms for the SMARP and Flare Predictors such that the area under each histogram integrates to 1. Both the SEP (green, positive samples) and flare-only (red, negative samples) data are split into 25 bins

As shown in Figure 2, the flare peak intensity is a powerful discriminator between 200 the positive and negative dataset. The flare intensity has been used as a feature to pre-201 dict the occurrence and properties (peak proton intensity, event duration, and etc.) of 202 SEP events (Laurenza et al., 2009; Balch, 2008). This predictive power difference between 203 the flare peak intensity and the SMARP Predictors on the left of Figure 2 has a big im-204 pact when comparing the SEP prediction capability with and without SMARP data. Mod-205 erate distinction between the events that led to an SEP and those that did not can be 206 identified in the predictors acquired using the SMARP active region coordinates (Ac-207

tive Region Angular Distance and Area). Large active regions increase the likelihood of an SEP event occurrence. The total unsigned flux is related to the particle acceleration therefore the SEP events are connected to higher flux values. The flux and intensity distributions show similar trends but with the former having less predictive power. The Vertical Field Gradient distribution of the Positive dataset aligns well with that of the Negative dataset, making it the least powerful predictor along with the R Value which shows the same trend.

²¹⁵ 4 Machine Learning Methods

To investigate whether the SOHO (SMARP AR data) or the GOES (flare eruption information) dataset can predict better the response variable of the two classes defined above, we use two popular groups of machine learning algorithms provided by the scikit-learn software package v0.24.2 for Python: different variations of the Support Vector Machine (SVM; Cortes & Vapnik, 1995) and two Regression Models.

4.1 Support Vector Machine

221

246

SVMs were initially designed and have been used to solve binary classification prob-222 lems (Shao et al., 2014). In the most general case, the SVM is fitted to the data using 223 a set of vector-target pairs (x_i, y_i) where i = 1, 2, ..., n. The target for positive and neg-224 ative observations respectively is $y_i \in \{1, 0\}$ and the corresponding physical character-225 istics feature vector is $x_i = (f_{i1}, f_{i2}, ..., f_{ip})$. For all tests performed, our training data 226 length is n = 116 and the maximum feature vector length is p = 6, where all calcu-227 lated predictors are used. Each different SVM method maps the input feature vector x_i 228 to a higher dimension space using an unknown function ϕ dependent on the user-defined 229 kernel K. Given a regularization parameter C > 0 it solves an optimization problem 230 to obtain the SVM trained weight vector w (Hsu & Lin, 2002; Inceoglu et al., 2018). Dur-231 ing testing, prediction is done by multiplying the trained vector w to the projected in-232 put feature vector $\phi(x_i)$ with an addition of a bias term. A more detailed study on how 233 to solve the SVM optimization equations is out of the scope of this research and can be 234 found elsewhere (Cortes & Vapnik, 1995; Vapnik, 1998). 235

The kernel function K is defined as the inner product of data pairs that correspond 236 to different observations i and j, $K(x_i, x_j) = \phi(x_i)^T \phi(x_j)$. In this study we train four 237 different variations of the SVM (Amari & Wu, 1999). One uses the Linear kernel K =238 $\langle x_i, x_j \rangle$, two use Polynomial kernels $K = \gamma \langle x_i, x_j \rangle^d$ where $d \in \{2, 3\}$ (second and third 239 degree) and the last one uses the Gaussian Radial Basis Function (RBF) kernel K =240 $exp(-\gamma ||x_i = x_i||^2)$ which has been used in similar studies (Inceoglu et al., 2018). The 241 weighting factor γ is user-defined and controls the influence a single training example 242 has on the classification task. The different kernels help the prediction model deal with 243 complex datasets such as the physical features of solar active regions by transforming 244 the input into any desired form. 245

4.2 Linear Models

The observed physical properties of a SMARP AR can be also processed for the purpose of prediction by linear models: regression methods in which the target value is expected to be a linear combination of the input features. Assuming a model function $f(x) = w^T x + b$ where w is a set of coefficients acquired during fitting, every feature's (x_i) predicted target y_i is 1 if $f(x_i) \ge 0$ and 0 if otherwise. In this case study linear models such as Ridge and Logistic regression are being used.

The ridge regression is one of the simplest machine learning algorithms and works well for small datasets while being computationally inexpensive. To fit the coefficients w to the training data, the ridge regression minimizes an ordinary Least Squares loss function with an additional term that penalizes the size of the coefficients, as given in (1).

$$J_{Ridge} = \|w^{\top}x - y\|_{2}^{2} + \alpha \|w\|_{2}^{2}$$
(1)

Between different training runs we vary the complexity parameter α in order to con-257 trol the amount of shrinkage and find the value that produces the most robust predic-258 tions. We do not adopt a cross validation procedure for selecting the tuning parameter 259 α due to considerations of sample sparsity and because the randomized picking process 260 of the training data leads to non-significant selection bias. Although it is a model often 261 adopted when the response y takes real numbers, we chose ridge regression because it 262 reduces overfitting, guarantees that we can find a solution and offers a different approach 263 for binary classification compared to other competing models. 264

The dichotomous nature of Logistic Regression makes it a great candidate for the binary SEP prediction task. We use the default Logistic Regression module provided by the Scikit-Learn library in Python (Pedregosa et al., 2011) which includes the l_2 regularization as a penalty and the Limited-memory Broyden–Fletcher–Goldfarb–Shanno (L-BFGS) optimization algorithm (Saputro & Widyaningsih, 2017) as a solver. The L-BFGS solver fits our application as it is robust and recommended for small dataset prediction tasks. To calculate the optimal w coefficients, Logistic Regression minimizes the cost function J for w and c.

$$J_{LR} = \frac{1}{2} w^{\top} w + C \sum_{i=1}^{n} \log(e^{-y_i(x_i^{\top} w + c)} + 1)$$
(2)

The constant C controls the regularization strength of the model. Although normalization is applied to the flare data before the fitting process, the C constant is also varied throughout different training runs in order to find the value that produces the most numerically stable prediction.

4.3 Training and Tuning the ML Models

The scarcity of the SEP events along with the mission duration of the MDI/SOHO 278 limits the size of the Positive dataset and leads to difficulties in separating the data into 279 training and testing subsets in a reasonable way. To overcome this problem, every model 280 is trained on 90% of the Positive events (58) and an equal number of Negative events. 281 The training of each algorithm is followed by a similarly balanced testing on the remain-282 ing 10% Positive (7) and an equal number of Negative events. This balanced training 283 and testing procedure is repeated k number of times to provide uncertainty assessment 284 285 of the random selection of events. In our work, k is chosen to be equal to 100. Each time, a different batch of Negative events is randomly selected from the pool of 7,626 flare erup-286 tion events that did not lead to an SEP. Similarly, in every run a different split between 287 training and testing occurs for the Positive dataset. 288

289 5 Results

277

For each one of the SVMs and Linear Models, we follow the same training procedure, aiming to predict whether an AR that produces a flare will lead to an SEP event. The goal is to illustrate how useful the SMARP dataset is for this particular task, we therefore train the ML models using two separate sets of features, one that uses SMARP information and one that uses flare information (see Section 2 for the detailed descriptions of the two sets of features). The number of features vary from 2 to 4 and the machine learning algorithms are tested on a number of different predictor combinations.

The comparison between the different types of predictors and algorithms is done 297 using three metrics that characterize and quantify the predictive power of classifiers: the 298 Accuracy (ACC), the True Skill Statistics (TSS; Hanssen & Kuipers, 1965) and the Hei-200 dke Skill Score (HSS; Heidke, 1926). For every set of k different runs a cumulative con-300 fusion matrix (Figure 4) is obtained based on the results from the raw SVM and Lin-301 ear Model outputs. Each row and column in the confusion matrix represents the num-302 ber of instances in an actual class and in a predicted class respectively. More informa-303 tion about the metrics, their equations and statistical meaning can be found in the works 304 of Inceoglu et al. (2018) and Florios et al. (2018). 305

The ACC can be artificially high in the rare event where a model always predicts the majority class. In flare and SEP prediction such naive cases are common due to the data imbalance, but in this paper all models are trained on a one-to-one positive-negative ratio, so these rare cases are not a concern. Therefore, this study's basic prediction quality metric is the ACC, with the TSS and HSS being presented too as auxiliary metrics.

311

5.1 SEP Prediction with SMARP Predictors

The cumulative confusion matrix in Figure 3 shows that out of 1400 validation in-312 stances, Ridge Regression correctly classifies 552 as being Positive and 449 as being Neg-313 ative when using the SMARP Flux and the AR Distance. This is the cumulative infor-314 mation obtained from 100 different runs, each of which has 14 validation points. The mean 315 accuracy suggests that 72% of the times ($\pm 12\%$ for a single run) the Third Degree Poly-316 nomial SVM algorithm can predict whether a flare will lead or not to an SEP using the 317 its AR SMARP features. The comparison between the probability of detection and the 318 probability of false detection lead to an average TSS level of 0.47 ± 0.24 for the same 319 100 runs. Similarly, the HSS measures a fractional forecast improvement over a random 320 forecast of 0.44 ± 0.25 . 321

The results show that the Linear Models can predict whether a flare will be accom-322 panied by SEPs with ACC values $\geq 0.70 \pm 0.12$ for a number of SMARP Predictors 323 combinations (Table 2 in Appendix). The maximum corresponding TSS and HSS val-324 ues for these combinations are above the 0.40 levels, while they are generally below the 325 0.35 levels for the rest of cases. Similar to row 3 of Table 1, the Polynomial models that 326 use the AR Distance and Area in Table 2 fail to produce a meaningful decision bound-327 ary yielding ACC values $\leq 0.52 \pm 0.04$, TSS values $\leq 0.15 \pm 0.28$ and HSS values \leq 328 0.05 ± 0.09 . Note here that a zero TSS or HSS value means that the method has no skill 329 over the random forecast, therefore the these specific Polynomial examples do not show 330 any predictive power at all. 331

Although the quality of the results cannot be judged based on the variance, the scores 332 indicate that the better a model's predictive power is, the lower the variance between 333 the different runs is. Thus, the Intensity based models have an ACC standard deviation 334 of ≤ 0.09 while in SMARP examples, where prediction quality is inferior, the ACC stan-335 dard deviation is ≥ 0.10 . This pattern is even more evident when considering the TSS 336 (or the HSS) for which the standard deviation can be as high as 0.28 at the SMARP Pre-337 dictors exclusive runs. Potential reasons about this behavior of variance is the small num-338 ber of Positive data which allows for low quality runs to not converge at all. 339

Both SVM and Linear models are affected by user-defined constants such as the α and C in Equations 1 and 2. An embedded grid search is employed for each experiment, where we vary each hyper-parameter on a range between 0.05 to 20. The parameter that produced the highest-quality and most consistent results was used for the examples presented in this study.



Figure 3. The distribution of 100 different ACC, TSS and HSS values are shown in the box plots (left). The values were obtained using the SMARP Flux & AR Distance on a Third Degree Polynomial SVM and constitute the best SEP prediction the SMARP data can achieve. Adding all the individual TP, TN, FP and FN values respectively we produce a cumulative confusion matrix for the 100 different runs (right).

5.2 SEP Prediction with Flare Predictors

345

Similar to the results presented before, the prediction quality metrics for the flare-346 only cases are calculated based on the confusion matrices obtained from each different 347 run. The cumulative confusion matrix in Figure 4 shows that out of 1400 validation in-348 stances, Ridge Regression correctly classifies 626 as being Positive and 651 as being Neg-349 ative when trained on flare intensity and distance. This is the cumulative information 350 obtained from k = 100 different runs, each of which was tested on 14 data points. The 351 mean accuracy suggests that 91% of the times ($\pm 8\%$ for a single run) the Ridge algo-352 rithm can predict whether a flare will lead or not to an SEP using its physical charac-353 teristics. The comparison between the probability of detection and the probability of false 354 detection lead to an average TSS level of 0.84 ± 0.12 for the 100 runs. Similarly, the HSS 355 measures a fractional forecast improvement over a random forecast of 0.82 ± 0.14 . 356

Using the Flare Predictors (first row of Table 1) all the different SVM and Linear models produce similar results. The TSS and HSS show higher standard deviation values (varying from 0.13 to 0.17) compared to the ACC. The predictive power of flare Intensity is demonstrated when comparing the first two box plots in Figure 5 with the rest, where different predictors other than Intensity are being used instead.

The ACC, TSS and HSS values range from 0.88 ± 0.09 to 0.92 ± 0.07 (values marked 362 red and green in Table 1 of the Appendix), 0.78 ± 0.17 to 0.86 ± 0.13 and 0.76 ± 0.18 to 363 0.84 ± 0.15 respectively for the runs that include Intensity accompanied with a SMARP 364 Predictor. These results show that all models, when using the Flare Intensity, can suc-365 cessfully predict $\leq 92\%$ of the times if a flare will be accompanied with an SEP. When 366 using the SMARP Predictors along with the Flare Distance instead, the ACC values range 367 from 0.60 ± 0.09 to a maximum of 0.71 ± 0.10 , the TSS from 0.36 ± 0.3 to 0.846 ± 0.2 368 and the HSS from 0.19 ± 0.18 to 0.42 ± 0.2 . This proves that when Intensity is not in-369 volved in the prediction process, all models yield inferior results, losing at the best case 370 0.17 ± 0.09 from the accuracy metric. We only test our models on a mix of SMARP and 371 Flare Predictors to verify the prediction power of the Intensity, as in real-life applica-372 tions the two groups of predictors cannot be used together due to the leading time dif-373 ference. 374



Figure 4. The distribution of k=100 different ACC, TSS and HSS values are shown in the box plots (left). The values were obtained using the Flare Intensity & Flare Distance on a Ridge Regression model and constitute the best SEP prediction the flare data can achieve. The box range shows the interquartile range, the red line inside it the median value, the whiskers show the results range and the two red dots show two outlier values. The range of the y-axis is kept the same with Figure 3 for comparison. Adding all the individual TP, TN, FP and FN values respectively we produce a cumulative confusion matrix for the 100 different runs (right).

Although each SVM or Linear model performs differently when trained on the same predictors, the variance between the models is of high significance only for some cases where the second and third degree Polynomial SVMs encounter convergence difficulties. For the extreme case of Flare Distance & ARAREA in Table 1 the accuracy difference between the best and the worst performing models is 0.1 ± 0.12 . For the rest of the predictors combinations the difference is $\leq 0.03 \pm 0.13$.

The maximum accuracy achieved on each one of the four main categories of predictor combinations is presented in Figure 5. The resulting ACC, TSS and HSS values show that regardless the machine learning model, the Flare Predictors generally perform better than the SMARP data because of the better predictive power of the flare intensity. Although the SMARP data cannot provide SEP forecast of quality similar to the flare intensity, it provides us with a better leading time compared to the Flare Predictors.

388

5.3 Comparison with Results in Literature

Inceoglu et al. (2018) used data provided by the SHARPs, GOES, and DONKI databases 389 to train SVMs that forecast both CME and SEP events with maximum TSS and HSS 390 of 0.92 ± 0.09 and 0.92 ± 0.08 . Anastasiadis et al. (2017) use the SDO/Helioseismic and 391 Magnetic Imager (HMI) full-disk magnetograms and the flare information from the SOHO/MDI 392 database on the prediction tool they call Forecasting Solar Particle Events and Flares 393 (FORSPEF). They achieve Heidke Skill Scores (HSS) of 0.37 ± 0.011 and 0.67 ± 0.007 394 when using solar flare data and CME data respectively. While we only use GOES data 395 to forecast exclusively SEP events (not CMEs), the best TSS and HSS our SVM imple-396 mentations achieve are 0.84 ± 0.12 and 0.82 ± 0.14 , results that are comparable to both 397 aforementioned studies. 398

On the other hand, Papaioannou et al. (2018) perform a principal component analysis (PCA) on a set of six solar variables obtained from GOES and LASCO in order to calculate a decision boundary for their logistic regression. They classify events as SEP versus non-SEP and achieve a maximum POD (TSS + POFD) of 77.78%. Based on flare prediction, the warning tool García-Rigo et al. (2016) present provides long-term warn-



Maximum Accuracy Across Models

Figure 5. A cumulative box plot for the four main categories of predictor combinations outlined in the Appendix Tables. More specifically, the first plot (pink) corresponds to row 1 in Table 1, the second (blue) to rows 2-5, the third (green) to 6-10 and the fourth (yellow) corresponds to Table 2. The plot makes evident the superiority of the flare intensity over the SMARP data.

ings of possible SEP event occurrence with POD scores of up to 58.3%. Núñez (2011) presents a dual-model system called UMASEP that has a POD of all (well- and poorly connected with flares) SEP events of 80.72%. The SMARP data in Figure 3 achieves a POD 78.8%, similar to the works of Papaioannou and Nunez. If intensity gets involved in our logistic regression model, we can achieve POD scores of up to 90%. Lastly, it is important to note that all the results we report are using a probability threshold $p_t =$ 0.5. It is possible to slightly increase the prediction statistics of our models as Anastasiadis et al. (2017) show in their work, but this is out of the scope of our study.

412 6 Conclusions

To predict SEP events we use the newly published Space-Weather MDI Active Re-413 gion Patches (SMARPs) dataset which includes observations of the solar magnetogram 414 that were made during the active Solar Cycle 23. Point data selected from the SMARP 415 time series is used on a variety of machine learning algorithms such as a different Sup-416 port Vector Machines and Linear Regression models. The purpose of this study is to eval-417 uate the power of this new data product for SEP forecast. Our results (Table 1 & 2) show 418 that SMARP can accomplish this task as it can identify correctly 72% of the times whether 419 an Active Region that produces a flare will lead to an SEP or not. Although the pre-420 diction results for the SMARP dataset are worse than the ones produced using the flare 421 peak intensity and location, we demonstrate that not only SMARP data produces bet-422 ter results compared to earlier SEP prediction works, but it also provides a better lead-423 ing time than other datasets. 424

⁴²⁵ In conclusion, although the SMARP dataset is constructed from the MDI data set, ⁴²⁶ which includes only the line-of-sight component of the surface magnetic field at a rel-

- ⁴²⁷ atively long 96-minute cadence, it can produce competitive prediction results for SEPs
- ⁴²⁸ while providing a longer leading time than using Flare Predictors.

429 Acknowledgments

- ⁴³⁰ This work was supported by NASA DRIVE Science Center grant 80NSSC20K0600. The
- 431 SEP event list we use in this work is documented in the NOAA Space Environment Ser-
- vice Center https://umbra.nascom.nasa.gov/SEP/ website, the SMARP data along with
- 433 full-disk Helioseismic and Magnetic Imager (HMI) magnetograms is available on the Joint
- 434 Science Operations Center database at http://jsoc.stanford.edu/ and the NOAA solar
- 435 X-ray flare dataset can be found at https://www.ngdc.noaa.gov/stp/solar/solarflares.html.
- 436 All codes and data are included in our Github repository at
- 437 https://github.com/skasapis/SEP_Prediction_Using_SMARP.

438 Appendix

Table 1.

Maximum ACC, TSS and HSS Values for the SVM and Linear Models using Different Predictors

	$_{ m SVMs}$				Linear Models		
Flare Predictors	Linear	RBF	Polynomial 2	Polynomial 3	Logistic Reg.	Ridge	
	0.90 ± 0.08	0.91 ± 0.07	0.90 ± 0.08	0.90 ± 0.08	0.90 ± 0.08	0.91 ± 0.07	ACC
1. Intensity & Flare Distance	0.82 ± 0.16	0.84 ± 0.13	0.82 ± 0.15	0.80 ± 0.16	0.83 ± 0.16	0.84 ± 0.12	TSS
v	0.80 ± 0.17	0.82 ± 0.14	0.80 ± 0.16	0.78 ± 0.17	0.80 ± 0.17	0.82 ± 0.14	HSS
SMARP & Flare Predictors							
	0.71 ± 0.10	0.68 ± 0.13	0.70 ± 0.11	0.70 ± 0.11	0.71 ± 0.11	0.71 ± 0.12	ACC
2. Flare Distance & USFLUXL	0.46 ± 0.20	0.39 ± 0.28	0.44 ± 0.24	0.42 ± 0.24	0.45 ± 0.22	0.46 ± 0.25	TSS
	0.42 ± 0.20	0.36 ± 0.26	0.40 ± 0.23	0.39 ± 0.23	0.42 ± 0.21	0.41 ± 0.23	HSS
	0.67 ± 0.14	0.69 ± 0.13	0.61 ± 0.14	0.61 ± 0.12	0.69 ± 0.12	0.70 ± 0.10	ACC
3. Flare Distance & RVALUE	0.36 ± 0.29	0.41 ± 0.27	0.25 ± 0.32	0.30 ± 0.32	0.41 ± 0.25	0.43 ± 0.22	TSS
	0.34 ± 0.27	0.38 ± 0.26	0.22 ± 0.27	0.23 ± 0.23	0.38 ± 0.24	0.40 ± 0.21	HSS
	0.66 ± 0.13	0.68 ± 0.11	0.60 ± 0.09	0.62 ± 0.10	0.70 ± 0.12	0.69 ± 0.11	ACC
4. Flare Distance & ABABEA	0.35 ± 0.27	0.40 ± 0.25	0.36 ± 0.30	0.39 ± 0.29	0.43 ± 0.24	0.00 ± 0.11 0.42 ± 0.24	TSS
	0.32 ± 0.26	0.36 ± 0.22	0.19 ± 0.18	0.24 ± 0.21	0.41 ± 0.23	0.39 ± 0.22	HSS
	0.69 ± 0.13	0.69 ± 0.13	0.67 ± 0.12	0.66 ± 0.11	0.69 ± 0.13	0.69 ± 0.11	ACC
5. Flare Distance USFLUXL	0.00 ± 0.10 0.41 ± 0.27	0.42 ± 0.27	0.35 ± 0.30	0.36 ± 0.25	0.35 ± 0.26	0.00 ± 0.11 0.41 ± 0.24	TSS
& ARAREA	0.38 ± 0.26	0.37 ± 0.26	0.32 ± 0.28	0.32 ± 0.23	0.39 ± 0.25	0.38 ± 0.22	HSS
	0.88 ± 0.09	0.89 ± 0.07	0.90 ± 0.08	0.90 ± 0.08	0.88 ± 0.09	0.89 ± 0.09	ACC
6. Intensity & USFLUAL	0.80 ± 0.16	0.80 ± 0.13	0.82 ± 0.14	0.80 ± 0.17	0.78 ± 0.17	0.79 ± 0.17	155
	0.77 ± 0.18	0.77 ± 0.15	0.80 ± 0.15	0.79 ± 0.18	0.76 ± 0.18	0.77 ± 0.17	Н55
	0.89 ± 0.08	$\textbf{0.91}\pm\textbf{0.07}$	0.91 ± 0.08	0.89 ± 0.09	$\textbf{0.91}\pm\textbf{0.07}$	0.90 ± 0.08	ACC
7. Intensity & RVALUE	0.79 ± 0.16	0.83 ± 0.14	0.83 ± 0.15	0.80 ± 0.16	0.84 ± 0.13	0.83 ± 0.15	TSS
	0.77 ± 0.17	0.82 ± 0.15	0.81 ± 0.16	0.78 ± 0.17	0.81 ± 0.14	0.80 ± 0.16	HSS
	$\textbf{0.91} \pm \textbf{0.07}$	$\textbf{0.91} \pm \textbf{0.07}$	$\textbf{0.91} \pm \textbf{0.07}$	0.90 ± 0.07	0.92 ± 0.07	0.91 ± 0.07	ACC
8. Intensity & ARDIST	0.84 ± 0.13	0.84 ± 0.13	0.83 ± 0.14	0.83 ± 0.13	0.86 ± 0.13	0.84 ± 0.14	TSS
	0.82 ± 0.14	0.81 ± 0.14	0.81 ± 0.15	0.81 ± 0.14	0.84 ± 0.15	0.82 ± 0.15	HSS
	$\textbf{0.91} \pm \textbf{0.07}$	0.91 ± 0.08	0.90 ± 0.08	$\textbf{0.91} \pm \textbf{0.07}$	0.92 ± 0.08	0.90 ± 0.08	ACC
9. Intensity, USFLUXL	0.83 ± 0.14	0.84 ± 0.15	0.82 ± 0.15	0.83 ± 0.13	0.85 ± 0.15	0.81 ± 0.16	TSS
& ARDIST	0.82 ± 0.15	0.82 ± 0.16	0.80 ± 0.16	0.82 ± 0.14	0.83 ± 0.16	0.80 ± 0.17	HSS
	0.90 ± 0.08	0.91 ± 0.07	0.91 ± 0.08	0.90 - 0.08	0.91 ± 0.08	0.90 ± 0.08	ACC
10. Intensity & MEANGBL	0.82 ± 0.15	0.85 ± 0.13	0.84 ± 0.15	0.82 ± 0.15	0.84 ± 0.14	0.82 ± 0.16	TSS
	0.80 ± 0.17	0.83 ± 0.14	0.83 ± 0.15	0.80 ± 0.16	0.82 ± 0.15	0.80 ± 0.16	HSS

Note. The ACC values ≥ 0.91 with standard deviation ≤ 0.07 are marked in bold. In green and red are marked the higher and lower accuracy values respectively for each one of the three predictor groups.

Table 2.

Maximum ACC, TSS and HSS Values for the SVM and Linear Models using SMARP Predictors

	SVMs				Linear Models			
SMARP Predictors	Linear	RBF	Polynomial 2	Polynomial 3	Logistic Reg.	Ridge	Score	
	0.67 ± 0.12	0.67 ± 0.13	0.70 ± 0.12	0.72 ± 0.12	0.70 ± 0.12	0.71 ± 0.12	ACC	
1. USFLUXL & ARDIST	0.39 ± 0.28	0.38 ± 0.28	0.43 ± 0.25	0.47 ± 0.24	0.43 ± 0.23	0.47 ± 0.25	TSS	
	0.34 ± 0.24	0.34 ± 0.25	0.39 ± 0.23	0.44 ± 0.25	0.40 ± 0.24	0.42 ± 0.24	HSS	
	0.65 ± 0.11	0.67 ± 0.12	0.65 ± 0.12	0.67 ± 0.12	0.69 ± 0.11	0.65 ± 0.12	ACC	
2. USFLUXL & ARAREA	0.35 ± 0.27	0.38 ± 0.28	0.34 ± 0.27	0.36 ± 0.26	0.37 ± 0.23	0.30 ± 0.27	TSS	
	0.30 ± 0.23	0.35 ± 0.24	0.30 ± 0.24	0.33 ± 0.24	0.34 ± 0.22	0.27 ± 0.23	HSS	
	0.69 ± 0.11	0.69 ± 0.11	0.52 ± 0.04	0.51 ± 0.03	0.67 ± 0.12	0.70 ± 0.12	ACC	
3. ARDIST & ARAREA	0.42 ± 0.25	0.42 ± 0.23	0.15 ± 0.28	0.10 ± 0.22	0.36 ± 0.25	0.42 ± 0.26	TSS	
	0.37 ± 0.23	0.38 ± 0.22	0.05 ± 0.09	0.03 ± 0.06	0.34 ± 0.24	0.40 ± 0.25	HSS	
	0.65 ± 0.13	0.68 ± 0.10	0.58 ± 0.11	0.60 ± 0.11	0.67 ± 0.11	0.66 ± 0.12	ACC	
4. ARDIST & RVALUE	0.33 ± 0.28	0.38 ± 0.22	0.18 ± 0.29	0.26 ± 0.29	0.36 ± 0.23	0.35 ± 0.25	TSS	
	0.31 ± 0.26	0.35 ± 0.20	0.15 ± 0.22	0.21 ± 0.23	0.34 ± 0.21	0.33 ± 0.23	HSS	
	0.67 ± 0.13	0.68 ± 0.11	0.70 ± 0.11	0.67 ± 0.10	0.70 ± 0.13	0.69 ± 0.10	ACC	
5. USFLUXL, ARDIST	0.36 ± 0.28	0.38 ± 0.24	0.42 ± 0.23	0.37 ± 0.21	0.42 ± 0.27	0.41 ± 0.22	TSS	
& ARAREA	0.34 ± 0.26	0.35 ± 0.22	0.39 ± 0.22	0.34 ± 0.19	0.39 ± 0.26	0.38 ± 0.21	HSS	
	0.68 ± 0.12	0.68 ± 0.13	0.66 ± 0.13	0.69 ± 0.12	0.67 ± 0.11	0.69 ± 0.13	ACC	
6. All SMARP Predictors	0.40 ± 0.24	0.32 ± 0.27	0.35 ± 0.28	0.42 ± 0.25	0.36 ± 0.24	0.40 ± 0.27	TSS	
	0.37 ± 0.23	0.35 ± 0.25	0.33 ± 0.27	0.39 ± 0.24	0.33 ± 0.23	0.38 ± 0.26	HSS	

Note. The ACC values ≥ 0.70 are marked in bolt. In green and red are marked the higher and lower accuracy values respectively.

439 **References**

- Amari, S.-i., & Wu, S. (1999). Improving support vector machine classifiers by mod ifying kernel functions. *Neural Networks*, 12(6), 783–789.
- Anastasiadis, A., Papaioannou, A., Sandberg, I., Georgoulis, M., Tziotziou, K.,
 Kouloumvakos, A., & Jiggens, P. (2017). Predicting flares and solar energetic

Kouloumvakos, A., & Jiggens, P. (2017). Predicting flares and solar energetic
 particle events: The forspef tool. *Solar Physics*, 292(9), 1–21.

- Bain, H., Brea, P., & Adamson, E. (2018). Using machine learning techniques to
 forecast solar energetic particles. In Agu fall meeting abstracts (Vol. 2018, pp.
 SM31D-3530).
- Balch, C. C. (2008). Updated verification of the space weather prediction center's solar energetic particle prediction model. *Space Weather*, 6(1).
- Barnes, W. T., Bobra, M. G., Christe, S. D., Freij, N., Hayes, L. A., Ireland, J., ...
 others (2020). The sunpy project: Open source development and status of the
 version 1.0 core package. *The Astrophysical Journal*, 890(1), 68.
- Bobra, M. G., Wright, P. J., Sun, X., & Turmon, M. J. (2021). Smarps and sharps:
 Two solar cycles of active region data.
- Brueckner, G., Howard, R., Koomen, M., Korendyke, C., Michels, D., Moses, J., ...
 others (1995). The large angle spectroscopic coronagraph (lasco). In *The soho mission* (pp. 357–402). Springer.
- Chen, Y., Manchester, W. B., Hero, A. O., Toth, G., DuFumier, B., Zhou, T., ...
 Gombosi, T. I. (2019). Identifying solar flare precursors using time series of sdo/hmi images and sharp parameters. *Space Weather*, 17(10), 1404–1426.
- Choudhary, D. P., Gosain, S., Gopalswamy, N., Manoharan, P., Chandra, R., Uddin,
 W., ... others (2013). Flux emergence, flux imbalance, magnetic free energy and
 solar flares. Advances in Space Research, 52(8), 1561–1566.
- 464 Cortes, C., & Vapnik, V. (1995). Support-vector networks. *Machine learning*, 20(3),
 465 273–297.
- ⁴⁶⁶ Florios, K., Kontogiannis, I., Park, S.-H., Guerra, J. A., Benvenuto, F., Bloomfield,
 ⁴⁶⁷ D. S., & Georgoulis, M. K. (2018). Forecasting solar flares using magnetogram⁴⁶⁸ based predictors and machine learning. *Solar Physics*, 293(2), 1–42.
- Forbush, S. E. (1946). Three unusual cosmic-ray increases possibly due to charged particles from the sun. *Physical Review*, 70(9-10), 771.
- 471 García-Rigo, A., Núñez, M., Qahwaji, R., Ashamari, O., Jiggens, P., Pérez, G., ...
- Hilgers, A. (2016). Prediction and warning system of sep events and solar flares
 for risk estimation in space launch operations. EDP Sciences.
- Hanssen, A., & Kuipers, W. (1965). On the relationship between the frequency of
 rain and various meteorological parameters. Koninklijk Nederlands Meteorologisch
 Instituut.
- Heidke, P. (1926). Berechnung des erfolges und der güte der windstärkevorhersagen im sturmwarnungsdienst. *Geografiska Annaler*, 8(4), 301–349.
- Horne, R., Glauert, S., Meredith, N., Boscher, D., Maget, V., Heynderickx, D., &
 Pitchford, D. (2013). Space weather impacts on satellites and forecasting the
 earth's electron radiation belts with spacecast. *Space Weather*, 11(4), 169–186.
- Hsu, C.-W., & Lin, C.-J. (2002). A comparison of methods for multiclass support
 vector machines. *IEEE transactions on Neural Networks*, 13(2), 415–425.
- Inceoglu, F., Jeppesen, J. H., Kongstad, P., Marcano, N. J. H., Jacobsen, R. H., &
 Karoff, C. (2018). Using machine learning methods to forecast if solar flares will
 be associated with cmes and seps. *The Astrophysical Journal*, 861(2), 128.
- Laurenza, M., Cliver, E., Hewitt, J., Storini, M., Ling, A., Balch, C., & Kaiser, M.
- (2009). A technique for short-term warning of solar energetic particle events based
- on flare location, flare size, and evidence of particle escape. Space Weather, 7(4).
- 490 Lavasa, E., Giannopoulos, G., Papaioannou, A., Anastasiadis, A., Daglis, I., Aran,
- ⁴⁹¹ A., & Pacheco, D. (2021). Assessing the predictability of solar energetic particles

with the use of machine learning techniques. 492 Lopez, R. E., Baker, D. N., & Allen, J. (2004). Sun unleashes halloween storm. Eos, 493 Transactions American Geophysical Union, 85(11), 105–108. McCracken, K., & Ness, N. (1966). The collimation of cosmic rays by the interplane-495 tary magnetic field. Journal of Geophysical Research, 71(13), 3315–3318. 496 Mumford, S. J., Christe, S., Pérez-Suárez, D., Ireland, J., Shih, A. Y., Inglis, A. R., 497 ... others (2015). Sunpy—python for solar physics. Computational Science \mathscr{C} 498 Discovery, 8(1), 014009.499 Núñez, M. (2011). Predicting solar energetic proton events (e_i, 10 mev). Space 500 Weather, 9(7). 501 Papaioannou, A., Anastasiadis, A., Kouloumvakos, A., Paassilta, M., Vainio, R., 502 Valtonen, E., ... Abunin, A. (2018). Nowcasting solar energetic particle events 503 using principal component analysis. Solar Physics, 293(7), 1–23. 504 Papaioannou, A., Sandberg, I., Anastasiadis, A., Kouloumvakos, A., Georgoulis, 505 M. K., Tziotziou, K., ... Hilgers, A. (2016). Solar flares, coronal mass ejections 506 and solar energetic particle event characteristics. Journal of Space Weather and 507 Space Climate, 6, A42. 508 Pedregosa, F., Varoquaux, G., Gramfort, A., Michel, V., Thirion, B., Grisel, O., ... 509 others (2011). Scikit-learn: Machine learning in python. the Journal of machine 510 Learning research, 12, 2825–2830. 511 Reames, D. V. (2013). The two sources of solar energetic particles. Space Science 512 Reviews, 175(1-4), 53-92. 513 Richardson, I., Mays, M., & Thompson, B. (2018).Prediction of solar energetic 514 particle event peak proton intensity using a simple algorithm based on cme speed 515 and direction and observations of associated solar phenomena. Space Weather. 516 16(11), 1862-1881.517 Rodriguez, J., Krosschell, J., & Green, J. (2014). Intercalibration of goes 8–15 solar 518 proton detectors. Space Weather, 12(1), 92-109. 519 Saputro, D. R. S., & Widyaningsih, P. (2017). Limited memory broyden-fletcher-520 goldfarb-shanno (l-bfgs) method for the parameter estimation on geographically 521 weighted ordinal logistic regression model (gwolr). In Aip conference proceedings 522 (Vol. 1868, p. 040009). 523 Scherrer, P. H., Bogart, R. S., Bush, R., Hoeksema, J.-A., Kosovichev, A., Schou, J., 524 ... others (1995). The solar oscillations investigation—michelson doppler imager. 525 In The soho mission (pp. 129–188). Springer. 526 Schou, J., Scherrer, P. H., Bush, R. I., Wachter, R., Couvidat, S., Rabello-Soares, 527 M. C., ... others (2012). Design and ground calibration of the helioseismic and 528 magnetic imager (hmi) instrument on the solar dynamics observatory (sdo). Solar 529 Physics, 275(1), 229–259. 530 Schrijver, C., Beer, J., Baltensperger, U., Cliver, E., Güdel, M., Hudson, H., ... oth-531 ers (2012). Estimating the frequency of extremely energetic solar events, based 532 on solar, stellar, lunar, and terrestrial records. Journal of Geophysical Research: 533 Space Physics, 117(A8). 534 Shao, Y.-H., Chen, W.-J., & Deng, N.-Y. (2014). Nonparallel hyperplane support 535 vector machine for binary classification problems. Information Sciences, 263, 22-536 35.537 Shea, M., & Smart, D. (1995). History of solar proton event observations. Nuclear 538 Physics B-Proceedings Supplements, 39(1), 16–25. 539 van Driel-Gesztelvi, L., & Green, L. M. (2015). Evolution of active regions. Living 540 Reviews in Solar Physics, 12(1), 1–98. 541 Vapnik, V. (1998). The support vector method of function estimation. In Nonlinear 542 modeling (pp. 55–85). Springer. 543 Webb, D. F., & Allen, J. H. (2004). Spacecraft and ground anomalies related to the 544 october-november 2003 solar activity. Space Weather, 2(3). 545

- ⁵⁴⁶ Wild, J., Smerd, S., & Weiss, A. (1963). Solar bursts. Annual Review of Astronomy
- ⁵⁴⁷ and Astrophysics, 1, 291.