# Understanding Controlling Factors of Extratropical Humidity and Clouds with an Idealized General Circulation Model

Michelle  $Frazer^{1,1,1}$  and Yi  $Ming^{2,2,2}$ 

<sup>1</sup>Princeton University <sup>2</sup>NOAA/Geophysical Fluid Dynamics Laboratory

November 30, 2022

#### Abstract

This paper examines the physical controls of extratropical humidity and clouds by isolating the effects of cloud physics factors in an idealized model. The Held-Suarez dynamical core is used with the addition of passive water vapor and cloud tracers, allowing cloud processes to be explored cleanly. Separate saturation adjustment and full cloud scheme controls are used to consider the strength of advection-condensation theory. Three sets of perturbations to the cloud scheme are designed to test the model's sensitivity to the physics of condensation, sedimentation, and precipitation formation. The condensation and sedimentation perturbations isolate two key differences between the control cases. First, the sub-grid-scale relative humidity distribution assumed for the cloud macrophysics influences the location and magnitude of the extratropical cloud maxima, limiting isentropic transport of tropical moisture to the polar troposphere. Second, within the model's explicit treatment of cloud microphysics, re-evaporation of hydrometeors moistens and increases clouds in the lower troposphere. In contrast, microphysical processes of precipitation formation (specifically, the ratio of accretion to autoconversion) have negligible effects on humidity, cloudiness, and precipitation apart from the strength of the large-scale condensation and formation cycle. Additionally, counterintuitive relationships—such as cloud condensate and cloud fraction responding in opposing directions—emphasize the need for careful dissection of physical mechanisms. In keeping with advection-condensation theory, circulation sets the patterns of humidity, clouds, and precipitation to first order, with factors explored herein providing secondary controls. The results substantiate the utility of such idealized modeling and highlight key cloud processes to constrain.

| 1 | Understanding Controlling Factors of Extratropical Humidity                               |
|---|---|
| 2 | and Clouds with an Idealized General Circulation Model                                    |
| 3 | Michelle E. Frazer*   |
| 4 | Program in Atmospheric and Oceanic Sciences, Princeton University, Princeton, New Jersey; |
| 5 | Department of Geosciences, Pennsylvania State University, State College, Pennsylvania     |
| 6 | Yi Ming   |
| 7 | NOAA/Geophysical Fluid Dynamics Laboratory, Princeton, New Jersey                         |

<sup>8</sup> \*Corresponding author: Michelle E. Frazer, mef257@psu.edu

# ABSTRACT

This paper examines the physical controls of extratropical humidity and clouds by isolating the 9 effects of cloud physics factors in an idealized model. The Held-Suarez dynamical core is used 10 with the addition of passive water vapor and cloud tracers, allowing cloud processes to be explored 11 cleanly. Separate saturation adjustment and full cloud scheme controls are used to consider the 12 strength of advection-condensation theory. Three sets of perturbations to the cloud scheme are 13 designed to test the model's sensitivity to the physics of condensation, sedimentation, and precip-14 itation formation. The condensation and sedimentation perturbations isolate two key differences 15 between the control cases. First, the sub-grid-scale relative humidity distribution assumed for the 16 cloud macrophysics influences the location and magnitude of the extratropical cloud maxima which 17 interrupt the isentropic transport of moisture to the polar troposphere. Second, within the model's 18 explicit treatment of cloud microphysics, re-evaporation of hydrometeors moistens and increases 19 clouds in the lower troposphere. In contrast, microphysical processes of precipitation formation 20 (specifically, the ratio of accretion to autoconversion) have negligible effects on humidity, cloudi-21 ness, and precipitation apart from the strength of the large-scale condensation and formation cycle. 22 Additionally, counterintuitive relationships—such as cloud condensate and cloud fraction respond-23 ing in opposing directions—emphasize the need for careful dissection of physical mechanisms. In 24 keeping with advection-condensation theory, circulation sets the patterns of humidity, clouds, and 25 precipitation to first order, with factors explored herein providing secondary controls. The results 26 substantiate the utility of such idealized modeling and highlight key cloud processes to constrain. 27

# 28 1. Introduction

Cloud feedback is widely considered to be the largest contributor to the intermodel spread in 29 climate sensitivity among comprehensive General Circulation Models (GCMs) (e.g., Ceppi et al. 30 2017; Sherwood et al. 2020). Bony et al. (2015) argued that consensus among most comprehensive 31 GCMs does not, on its own, yield robust conclusions on cloud feedback. Rather, theories which 32 underpin physical arguments and improve understanding in a way that allows for expanded use 33 and interpretation of comprehensive GCMs are an additional requirement. Thus, simple models 34 whose workings can be clearly grasped play a key role in the midst of a complex scientific problem 35 (Pierrehumbert et al. 2007; Held 2005, 2014). If a GCM produces both observationally-constrained 36 cloud fields and multi-model consistent cloud feedbacks, but without the physical mechanisms 37 necessarily being represented appropriately, its prediction of the climatic response to a radiative 38 forcing may be significantly flawed. With the potential for unrealistic interactions between different 39 parameterized processes (Ceppi et al. 2017), decomposition of the effects of individual processes 40 could lead to improved parameterizations. 41

Here, we study under-constrained cloud macrophysical and microphysical processes by exploring 42 the underlying physical mechanisms. Since changing a stratiform cloud scheme can have significant 43 ramifications, even reversing a model's feedback with warming (Geoffroy et al. 2017), we use an 44 idealized setup to break down a cloud scheme and understand the effects of individual cloud 45 processes on atmospheric humidity and cloudiness. The processes studied herein are motivated 46 by three factors: understanding the differences between the advection-condensation theory of 47 humidity and a cloud scheme, the controls of large-scale precipitation efficiency, and the direct 48 effect of stratiform-cloud related GCM parameters on free tropospheric humidity and clouds. 49

# <sup>50</sup> a. Advection-Condensation Theory

Free tropospheric humidity is important to the distribution of clouds and precipitation. The so-called advection-condensation theory suggests that water vapor (WV) in the atmosphere is most simply reflective of the lowest temperature (lowest saturation specific humidity) experienced by the parcel since leaving the nearly saturated surface layer. This theory alone can describe WV distribution to first order (Sherwood et al. 2010). Advection-condensation theory helps explain two key features of free tropospheric humidity: dry subtropical zones and moist polar regions connected by dry isentropes.

Pierrehumbert (1998) laid out three factors which contribute to the dry subtropics. First, sub-58 sidence brings down dry air, and would keep the region at the mixing ratio of the tropopause if 59 not for other mechanisms. Second, lateral mixing brings in moist air from the tropical convective 60 region. Third, processing of air through cold extratropics dries the region. Thus, the dry subtropics 61 and moist poles are connected through nearly isentropic large-scale advection, and cycling through 62 cold polar upper tropospheric air is a key means of dehydrating air in the extratropics (Kelly et al. 63 1991). Finally, Pierrehumbert (1998) also noted the role of re-evaporation of hydrometeors as 64 a subtropical moisture source as emphasized by Sun and Lindzen (1993), but suggested this is 65 limited by weak rainfall. Also suggesting the importance of in situ moistening processes in the 66 midlatitudes, Yang and Pierrehumbert (1994) showed that in the advection-condensation model, 67 the tropical moisture source is too inefficient (that is, too weak of mixing between tropics and 68 extratropics). These factors have been expounded in further work. 69

<sup>70</sup> Using a simple saturation adjustment scheme as a representation of advection-condensation the-<sup>71</sup> ory, Galewsky et al. (2005) found that the primary dynamical control of the dry subtropics was <sup>72</sup> isentropic dehydration by mid-latitude eddies (with diabatic descent through Hadley circulation

playing a secondary role). WV is transported from the lower deep tropics to the upper polar extrat-73 ropics by baroclinic eddies along isentropes, with the moist air rising and cooling adiabatically. The 74 storm tracks interrupt the transport such that significant moisture is released through precipitation 75 before reaching the poles. Thus, the return flow supplies dehydrated air to the subtropics, and is 76 confined to isentropic layers (Held and Schneider 1999). The poleward eddy WV transport follows 77 dry isentropes but different values of equivalent potential temperature, with this moist recirculation 78 peaking on the equatorward side of the storm tracks (Laliberté et al. 2012). In this study, we 79 consider how a cloud scheme distributes moisture differently than simple saturation adjustment (as 80 in Galewsky et al. 2005), and we highlight the processes—cloud macrophysics and microphysics 81 alike-that affect extratropical humidity strongly. The physical mechanisms of these controls are 82 delineated to highlight those processes that need to be represented accurately in cloud schemes. 83

# <sup>84</sup> b. Precipitation Efficiency Controls

Differences between saturation adjustment and a cloud scheme are closely related to the controls 85 of precipitation efficiency. The residence time of water in the atmosphere is, in a full cloud scheme, 86 affected by three efficiencies: the efficiency with which WV may become cloud condensate (con-87 densation), become part of a falling hydrometeor (formation), and reach the surface as precipitation 88 (sedimentation) (Langhans et al. 2015). Advection-condensation theory reduces this complexity to 89 one efficiency since WV in excess of saturation immediately becomes surface precipitation. Thus, 90 condensation and sedimentation efficiencies highlight two of the key differences between a satura-91 tion adjustment scheme (based on advection-condensation theory) and a full cloud scheme (closer 92 to reality): condensation efficiency is affected by assumptions of small (sub-grid) scale relative 93 humidity (RH) distribution, and sedimentation efficiency by re-evaporation of precipitation. The 94 third efficiency—formation efficiency—can be affected by internal cloud scheme parameters such 95

as the assumed cloud condensation nuclei (which affects warm rain processes) or the fall speed 96 of ice. But each of the three efficiencies have the potential to significantly affect WV and cloud 97 condensate (CC) fields, the distribution of precipitation, and the overall residence time of atmo-98 spheric water. For example, precipitation efficiency (the multiplicative product of formation and 99 sedimentation efficiencies; see section 2b) is frequently highlighted as being potentially affected by 100 warmer temperatures resulting in more liquid at the expense of ice in mixed-phase clouds (Klein 101 et al. 2009; McCoy et al. 2015; Ceppi et al. 2016; McCoy et al. 2018). Here we explore the direct 102 effect of changing these efficiencies on steady-state fields which are relevant to radiative feedbacks. 103

# <sup>104</sup> c. GCM Stratiform Tuning Parameters

Thus the first two motivations are connected to the third of the direct effect of stratiform-cloud 105 related GCM tuning parameters on free tropospheric humidity and clouds. Critical RH (the 106 minimum GCM grid-box-mean RH needed for cloud condensate formation) is a useful tuning 107 parameter for radiative balance (through shortwave cloud radiative effects), but may be tuned 108 artificially high in order to compensate for too-bright clouds (McCoy et al. 2016). Critical RH is 109 important because it controls large-scale condensation, a sink of WV and source of CC. WV can 110 be altered without directly affecting CC by tuning the re-evaporation of precipitation. Another key 111 parameter is N, the assumed cloud drop number concentration: aerosols affect microphysics and 112 thus precipitation and radiation through aerosol-cloud interactions. The observed precipitation rate 113 can be expressed as a power-law function of LWP and N, with a strong correlation between liquid 114 water path (LWP) and the ratio of accretion to autoconversion processes (hereafter *accr/auto*; 115 Jiang et al. 2010). At low LWP, *accr/auto* is small because of few generated rain drops. Some 116 GCMs directly model aerosol indirect effects, but even in simpler cloud microphysics schemes 117 which lack an explicit representation of aerosol indirect effects, the autoconversion process is a 118

direct function of *N* and thus a major control of accr/auto, which is a key parameter for examining the balance of microphysical conversion processes from cloud water to rainwater (e.g., Gettelman et al. 2013).

In a GCM study implementing five different autoconversion schemes, Michibata and Takemura 122 (2015) found significant variance in *accr/auto*. But, these schemes showed a commonality of 123 the relative role of the accretion process being one or more orders of magnitude underestimated 124 compared to observations (as estimated by Gettelman et al. 2013). This incorrect ratio comes 125 from both too high simulated autoconversion rates (Gettelman et al. 2013, 2014) and in some 126 schemes, too low of an accretion enhancement factor for correct precipitation intensity (Wu et al. 127 2018). The high simulated autoconversion rates come from diagnostic precipitation which forms 128 warm rain too easily (Jing et al. 2017). Cloud condensation nuclei and *accr/auto* affect not 129 only precipitation rates but also radiative forcing. Increased *accr/auto* in GCM simulations 130 is correlated with increasing LWP (Gettelman et al. 2013), and cloud optical depth and thus 131 shortwave radiative effect is significantly controlled by LWP (e.g., Stephens 1978). As past studies 132 have likely underestimated the true sensitivity of clouds and radiation to aerosols, the negative 133 forcing of the Twomey effect (altered cloud albedo from increased anthropogenic aerosols) may be 134 underestimated (Quaas et al. 2020); though, the aerosol-cloud lifetime effect may be overestimated 135 (e.g., Quaas et al. 2009). Yet, Gettelman et al. (2013) suggested that the autoconversion rate bias 136 can be corrected by altering the relative balance of the autoconversion and accretion rates, which 137 lowers the radiative effect of aerosol cloud interactions. Thus, understanding the interplay and 138 impacts of altered N and *accr/auto* is critical. 139

#### <sup>140</sup> *d. Purpose and Organization*

The overarching purpose of this paper is to employ an idealized model setup to shed light on what 141 controls free tropospheric humidity and cloudiness. Using perturbation experiments which isolate 142 key processes, we aim at elucidating the complex connections among WV, clouds, precipitation, 143 and circulation. In analyzing the control and perturbation experiments in this study, the budgetary 144 terms of the cloud scheme which represent the conversions among WV, CC, and precipitating 145 water (P) are particularly emphasized. This method is motivated by a need for a robust physical 146 understanding to ground model representations of cloud processes in order to lend confidence to 147 model-inferred relationships (Shepherd 2014; Stevens and Bony 2013). 148

A process-based analysis is related to the secondary purpose of this work: to clearly demonstrate 149 the value of this modeling tool (a dry GCM with passive water and cloud tracers) for developing a 150 systematic understanding of physical controls on humidity and clouds and diagnosing their repre-151 sentations in models. This approach is in the same spirit as "mechanism-affirmation experiments" 152 described in Jeevanjee et al. (2017) as being the provision of a model hierarchy framework. In 153 terms of the model hierarchy, the setup used in this paper (Ming and Held 2018) is derived from the 154 Held-Suarez (HS) dry GCM, but in a different direction than the Frierson moist aquaplanet GCM 155 (Frierson et al. 2006) which extended the HS dry GCM by adding a gray radiation scheme and moist 156 physics such that latent heating affects the model's dynamics. Our model is in many aspects more 157 idealized than the Frierson model with dry dynamics and no radiation scheme, but more complex 158 in its addition of a full cloud microphysics scheme. It can be thought of as one rung higher on the 159 model hierarchy ladder than the HS dry GCM, but one rung lower than the Frierson model. This 160 setup is therefore uniquely suitable for answering specific questions about extratropical humidity 161 and cloudiness—namely the direct effects of cloud macrophysics and microphysics—as well as 162

the physical mechanisms behind these effects. With passive humidity and cloud tracers, isolated experiments are able to be performed such that the direct effect of a cloud physics process can be clearly diagnosed without the convoluting circular effects of dynamical processes.

This paper is organized as follows. Section 2 lays out the methodology of this study, describing the idealized model, experiments, and analysis framework. Section 3 describes the results from the control saturation adjustment and cloud physics experiments and the condensation, sedimentation, and formation perturbations. Section 4 discusses the implications of these results for the value of the advection-condensation paradigm, key stratiform cloud physics processes to constrain, and the utility of this idealized model.

# 172 2. Methodology

# 173 a. Control Models

The idealized model used here is based on the HS dry GCM (Held and Suarez 1994) with the 174 addition of four passive water and cloud tracers-specific humidity, cloud liquid, cloud ice, and 175 cloud fraction (CF)—as described in Ming and Held (2018). The dry GCM uses a hydrostatic 176 spectral dynamical core for an ideal gas atmosphere with no topography. For this work, a resolution 177 of T42 (referring to the maximum number of zonal waves present in the triangular truncation) is 178 used, resulting in a horizontal grid of 128 by 64 cells (about 2.8° spacing) with 20 vertical layers 179 equally spaced in the sigma coordinate. The forcing consists of Newtonian relaxation of temperature 180 toward a prescribed zonally symmetric equilibrium temperature and planetary boundary layer drag 181 represented by Rayleigh damping. This idealized setup enables the isolation of the roles of various 182 cloud processes. It assumes that latent heating or cooling from conversions among WV, CC, and 183 precipitation do not feed back on the dynamics. Also, with no explicit radiation scheme in the 184

<sup>185</sup> model, clouds do not affect circulation through cloud radiative effects. Thus, WV and clouds are <sup>186</sup> passive in that they do not affect circulation or temperature patterns.

Two control simulations are created with results explored in section 3a. The first, referred to as 187 the *Base* case, uses only the specific humidity tracer in a saturation adjustment scheme modeled 188 after Galewsky et al. (2005) as a direct representation of advection-condensation theory. Any 189 water in excess of saturation (grid-box mean) is assumed to fall out immediately as precipitation. 190 Thus, no clouds are present. The second control simulation is referred to as the *Cloud* case. It 191 carries specific humidity, cloud liquid, cloud ice, and CF tracers through the same large-scale cloud 192 macrophysics scheme as implemented in the GFDL HiRAM model (Zhao et al. 2009). The cloud 193 scheme assumes a beta distribution for sub-grid-scale total water (which includes both WV and 194 CC). CF is diagnosed from this total water-based RH, which varies only slightly from traditional 195 RH (which is based on WV only and is the RH reported in the results). The default beta distribution 196 is such that a grid-mean total water-based RH value exceeding 83.3% (the critical RH: RH<sub>c</sub>) allows 197 for sub-grid values greater than 100% and thus a non-zero CF for the grid box. 198

The pathways for conversion between WV, cloud liquid, cloud ice, and hydrometeors follow a 199 Rotstayn-Klein single-moment microphysics scheme (after Rotstayn 1997; Rotstayn et al. 2000). 200 Additionally, as the principal source of WV, surface evaporation is represented by adjusting the 201 specific humidity of grid boxes below  $\sim$ 850 hPa towards saturation with an e-folding timescale of 202 30 minutes. Microphysical sources of WV are large-scale (LS) evaporation of cloud liquid, LS 203 sublimation of cloud ice, rain evaporation, and snow sublimation. The only sinks of WV, namely 204 LS condensation and LS deposition, are also the only sources of CC. CC is lost to WV through LS 205 evaporation and LS sublimation, to rain through autoconversion, accretion, and melting of cloud 206 ice, and to snow through gravitational settling. Additionally, cloud liquid is converted to cloud ice 207 through riming, the Bergeron-Findeisen process, and homogeneous freezing, and both cloud ice 208

and snow can be converted to rain through melting. (Cloud ice and snow have identical properties such as fall speed and are simply distinguished by their location in or out of a cloud.) See Fig. 1 in Frazer and Ming (2022) and the descriptive text for more details of these conversions.

# <sup>212</sup> b. Perturbation Experiments

On the surface, there are three chief distinctions between saturation adjustment (Base control) and a full cloud scheme (Cloud control). First, clouds can form (and thus precipitation is possible) before the grid box is fully saturated through  $RH_c$  and an assumed sub-grid-scale RH distribution. Second, the cloud scheme allows precipitation to evaporate before reaching the surface through rain evaporation and snow sublimation (hereafter RESS). Third, cloud condensate may be advected before precipitating out or evaporating. The effects of the first two distinctions can be easily explored by being simply "turned-off" in the cloud scheme. The third is inferred as a residual effect.

Each of the three distinctions correspond to the three efficiencies which effect the residence time of 220 water in the atmosphere and form a key part of the analysis. We make use of the explicit/large-scale 221 precipitation efficiency (PE) as defined in Zhao (2014) to represent the total PE, since only stratiform 222 (not convective) precipitation is represented in this model. PE is the ratio of surface precipitation 223 to vertically integrated CC sources, and thus represents the fraction of condensed particles which 224 subsequently rain out. Following Langhans et al. (2015), PE can be thought of as the product of 225 formation efficiency (FE) and sedimentation efficiency (SE): PE = FE \* SE. FE represents the 226 probability of formation given condensation, and SE represents the probability of sedimentation 227 given formation. Finally, the condensation efficiency (CE) is used herein to simply represent the 228 fraction of atmospheric WV that subsequently condenses (as there is no explicit treat of entrainment 229 in this stratiform scheme). Thus, CE is the ratio of CC sources (condensation and deposition) to 230 WV sources (surface evaporation, CC evaporation and sublimation, and RESS), FE is the ratio of 231

precipitation formation (autoconversion, accretion, melting of cloud ice, and gravitational settling) 232 to CC sources, and SE is the ratio of surface precipitation to precipitation formation. Additionally, 233 the residence (or recycling) time for WV in the atmosphere is defined after Trenberth (1998) as the 234 *e*-folding time constant for the depletion of precipitable water by precipitation, that is, the global 235 ratio of column-integrated WV to the precipitation rate. These indicators of features of the water 236 cycle are used to quantify changes in the WV, CC, and precipitation budgetary terms to supplement 237 the analysis of steady-state fields. But also, as these efficiencies correspond to distinctions between 238 saturation adjustment and a cloud scheme, we intentionally alter the efficiencies to understand the 239 effects on steady-state fields. CE is affected by  $RH_c$ , SE is 100% without RESS, and FE cannot be 240 defined without CC. 241

Thus, three principal perturbation experiments are designed, testing sensitivity to condensation, 242 sedimentation, and formation cloud processes. The condensation perturbation focuses on the con-243 version between WV and CC through cloud macrophysics, specifically sub-grid-scale cloudiness. 244 The first key distinction between saturation adjustment and a cloud scheme can be eliminated 245 by removing sub-grid-scale cloudiness and requiring 100% grid-mean RH for cloud formation. 246 Accordingly, an intermediate setup between the Base and Cloud controls is created by reducing the 247 width parameter of the beta distribution defining sub-grid-scale RH from 0.2 to 0.01, effectively 248 requiring 100% grid-box-mean RH for cloud formation. This perturbation run is referred to as 249 *RHc100* (since effectively  $RH_c = 100\%$ ) with results in Section 3b. 250

The sedimentation perturbation focuses on the role of re-evaporation of hydrometeors. While saturation adjustment oversimplifies the variety of conversions in this Rotstayn-Klein microphysics scheme, it is analogous to the LS phase changes and precipitation processes. The chief remaining processes are the recycling of hydrometeors back to WV through RESS. Thus, another intermediate setup between the controls is created to illuminate the significance of RESS. For this experiment<sup>256</sup> *noRESS* which is presented in Section 3c—the rates of RESS are arbitrarily set to zero. Additionally, <sup>257</sup> to examine the combined effect of the key microphysical and macrophysical differences between <sup>258</sup> the Base and Cloud cases, a final intermediate case is considered. The *RHc100\_noRESS* case <sup>259</sup> includes the  $RH_c = 100\%$  and omission of RESS effects to examine residual differences between <sup>260</sup> the control cases, which is assumed to correspond to the third key difference between saturation <sup>261</sup> adjustment and full cloud physics—advection of CC—as explored in Section 4.

The formation perturbation is not focused directly on a difference between the Base and Cloud 262 cases. In the Base case saturation adjustment, precipitation is formed directly from WV in a manner 263 more similar to condensation than formation. Rather, formation is explored so that sensitivity to 264 all key conversions of the cloud scheme are considered. Formation consists of three major process: 265 autoconversion, accretion, and ice settling. Ice settling is a net term-the difference between ice 266 falling into and out of grid boxes. Accordingly, autoconversion and accretion were isolated as the 267 best processes to perturb in order to explore formation sensitivities. From a general perspective, 268 if autoconversion or accretion is arbitrarily reduced in this model, the other process strengthens to 269 keep formation close to constant, but somewhat reduced. Conversely, if one process is amplified, 270 the other weakens. An analogous effect results from altering the prescribed cloud drop number 271 concentration, N, the default value being 50 cm<sup>-3</sup>. For autoconversion to occur, the radius of 272 the cloud droplets—a function of N—must be greater than the critical particle radius threshold at 273 which autoconversion occurs, and autoconversion increases directly with increasing N. Increased 274 autoconversion should have two effects on accretion: increasing the flux of rain (to scavenge cloud 275 liquid) and decreasing the pool of cloud liquid available to be scavenged. Here, the second effect 276 wins out such if N is decreased, autoconversion increases and accretion decreases with a net 277 amplification of formation. An increase of N produces an opposite effect. Thus, the strength of 278

<sup>279</sup> formation and the balance between autoconversion and accretion have broader significance because <sup>280</sup> of their connection to drop number concentration parameterizations.

Here, alterations to autoconversion are used to adjust *accr/auto* (and indirectly explore a key 281 affect of altered N). The principal formation perturbation explored in Section 3d, halvAUTO, 282 consists of halving the computed value for autoconversion for each grid box at each time-step. 283 For robustness, a corresponding doubling of autoconversion, *doubAUTO* is also examined. Note 284 that the halving or doubling of autoconversion is performed in the microphysical code before the 285 enforcement of a limiter which ensures that autoconversion is limited to the amount that reduces 286 local liquid cloud condensate to the critical value at which autoconversion begins (after Rotstayn 287 1997). 288

For all control and perturbation experiments, the atmospheric state of the model (winds, temperature, etc.) is identical at every time-step. The various experiments performed are summarized in Table 1. All model runs in this study include a 300-day spin-up of the dry GCM before the next 1000 days are averaged. For figures and analysis, data is averaged between the two hemispheres because of the hemispheric symmetry of the simulated climate. 15° to 90° is considered the suband extra-tropics (STET) and is the focus of the analysis due to the lack of a convection scheme making the tropics nearly saturated (see Ming and Held 2018).

# 296 **3. Results**

# *a. Controls: Base and Cloud*

A budgetary comparison of the control cases is shown in Fig. 1a, which depicts the principal WV tendency terms for the Base and Cloud cases from a column-integrated, zonally-averaged perspective. For the Base case, the WV balance is simply between precipitation from saturation

adjustment and surface evaporation. Outside of the tropics (which are not shown), the immediate 301 precipitation dominates in the mid-latitude storm tracks while evaporation occurs mostly in the 302 subtropics, implying significant horizontal advection of water from the subtropics (including 303 that facilitated by mid-latitude baroclinic eddies). For the Cloud case, the dominant balance 304 between net LS condensation (condensation and deposition minus evaporation and sublimation 305 with condensation dominating) as the main WV sink and surface evaporation as the main WV 306 source is similar to the Base case, though RESS do make a non-negligible contribution. Cloud 307 case LS condensation is everywhere stronger than Base case saturation adjustment, while the 308 surface evaporation is nearly indistinguishable except in the high latitudes where Base surface 309 evaporation is negligible. (Surface evaporation is a direct function of low level RH, which is 310 similar between the Base and Cloud cases other than in the high latitudes, as discussed below. In 311 the high latitudes, the Base case has higher RH (near saturation) and therefore minimal surface 312 evaporation.) Thus, RESS together provide an additional source of WV, strengthening the WV 313 cycle as opposed to replacing surface evaporation as a source. Fig. 1b shows the CC budget 314 applicable only to the Cloud case. Net LS condensation as the source of CC is balanced nearly 315 perfectly latitudinally, implying minimal advection of CC. In the subtropics, autoconversion is 316 the strongest sink of CC, but ice settling (snow) dominates poleward of 40° with rain processes 317 becoming negligible poleward of  $60^{\circ}$ . 318

<sup>319</sup> While precipitation is simply saturation adjustment in the Base case but formation processes <sup>320</sup> minus RESS in the Cloud case, both precipitation and precipitation minus evaporation (P-E) have <sup>321</sup> similar latitudinal distributions in the two cases (Fig. 1c). The principal latitudinal difference is <sup>322</sup> a slight increase in precipitation (and thus P-E) in the extratropics in the Cloud case, where ice <sup>323</sup> settling (a process vastly different than saturation adjustment) dominates as the principal source of <sup>324</sup> precipitation, and surface evaporation decreases in the Base case as discussed previously. Thus, the strength of the hydrological cycle in terms of surface precipitation is largely indistinguishable with a STET average of 1.84 mm/day in the Base case and 1.91 mm/day in the Cloud case (see Table 2 which also shows a similarity in surface evaporation). This correspondence between these idealized saturation adjustment and full cloud microphysics models without any control by radiative balance suggests a significant control of the hydrological cycle by large-scale circulation perhaps mediated through RH (as discussed below).

In contrast, the strength of the WV cycle differs greatly between the two control cases. This can 331 be seen in Fig. 2a and b which depict the STET-averaged, column-integrated values and fractions 332 of the sources and sinks in the Base and Cloud cases. The total STET WV sources and sinks in the 333 Cloud case are  $3.36 \times 10^{-5}$  and  $2.82 \times 10^{-5}$  kg m<sup>-2</sup> s<sup>-1</sup>, respectively, with the regional imbalance 334 implying advection of WV into the tropics (since evaporation is strongest in the subtropics). For 335 comparison, the Base case analogs of surface evaporation (the only WV source) and condensation 336 (the only WV sink) are  $2.70 \times 10^{-5}$  and  $2.11 \times 10^{-5}$  kg m<sup>-2</sup> s<sup>-1</sup>, respectively. Thus, the strength of 337 the cycling of WV is significantly enhanced in the Cloud model by ~30%. Adding more sources 338 and sinks of WV, in particular introducing sources above the boundary layer through RESS, allows 339 for a strengthening of the WV cycle and a slight shortening of the residence time (from 13.1 to 340 12.7 days). In the Cloud case, CC is also cycled where all the WV sinks are CC sources, and 341 precipitation processes are the main CC sinks (see Fig. 2b) with CC sources and sinks balanced in 342 the STET region. 343

This overall picture of water cycling between WV, CC, precipitation, and an assumed surface reservoir can be seen in Fig. 3 and described in terms of efficiencies. For the STET WV produced through surface evaporation, RESS, and evaporation (LS evaporation and sublimation), 83.9% is condensed (through LS condensation and deposition). Of the water condensed, most forms precipitation, while some is evaporated (a very small effect in this model with only a stratiform <sup>349</sup> cloud scheme) resulting in a FE of 98.2%. (Some also persists as condensate but this effect is lost <sup>340</sup> with time-averaging). Of the precipitation formed, ~20% is returned to WV through RESS before <sup>351</sup> reaching the surface resulting in a SE of 79.7% and a PE of 78.3%. These efficiencies, along <sup>352</sup> with precipitation and residence times, are summarized in Table 2. The positive WV reservoir and <sup>353</sup> negative surface reservoir values are again indicative of moisture export (negative P-E) from the <sup>354</sup> STET region.

Fig. 3 also shows how a cloud scheme builds on saturation adjustment. In Base case, only two 355 reservoirs—WV and surface—would exist with two arrows between them representing surface 356 evaporation and saturation adjusting. Yet, qualitative similarity exists in the RH distribution of the 357 Base and Cloud cases as shown in Fig. 4a. Both cases have qualitatively realistic free tropospheric 358 RH features: the subtropics and upper troposphere are relatively dry, while the extratropics are 359 moist (Fig. 4a). As noted in Ming and Held (2018), the high RH values in the deep tropics (not 360 shown) and boundary layer (below 850 hPa) are due to the lack of a moist convection scheme and 361 the way in which surface evaporation is modeled, respectively. Fig. 4a suggests that the addition 362 of a cloud scheme has two main effects on the RH distribution, while keeping the main features 363 present. The subtropical dry zones and nearby mid-latitudes are substantially moistened with a 364 peak increase of up to around 5% RH, while much of the polar upper troposphere becomes drier 365 by a similar magnitude. The mechanisms for these changes are investigated in the condensation 366 and sedimentation perturbations. Fig. 4b shows the model isentropes, significant because of the 367 established isentropic transport of moisture from the subtropics as discussed in the introduction. 368 Here, it is clear that the polar upper troposphere (drier in the Cloud case) is connected to the 369 subtropical boundary layer via isentropes. Yet, the overall similarity between the control cases in 370 the free troposphere implies that RH is controlled to first order by general circulation, as opposed to 371

cloud processes. Thus, in keeping with advection-condensation theory, one does not need detailed
 cloud information for understanding large-scale (first-order) RH patterns.

The cloud fields generated in the Cloud case are shown in Fig. 4c-d. Free tropospheric CF values 374 peak at near 30% in the extratropical storm track region, co-incident with the 75% average RH 375 contour. Liquid cloud condensate (LCC) is concentrated in the boundary layer (unrealistically high 376 because of high RH from artificial surface evaporation as discussed above) with a secondary peak 377 near the storm tracks. Ice cloud condensate is concentrated in a broad region near the storm tracks 378 restricted to freezing temperatures (see Fig. 4b). LCC, with its higher magnitude, dominates the 379 spatial pattern of total CC, which is the sum of ice and liquid water mixing ratios. Since the focus 380 of this study is on total clouds, not on the distribution of ice versus liquid, the remainder of this 381 work will consider only total CC, which is concentrated in the tropics with a secondary peak in the 382 storm tracks. 383

#### <sup>384</sup> b. Condensation Perturbation: RHc100

As discussed in the introduction, since isentropic transport is the key source of WV for the 385 polar regions, cloud formation (and precipitation) in the extratropical storm tracks interrupt WV 386 reaching the polar regions. In the Cloud case, cloud formation (required for precipitation) takes 387 place when grid-mean RH (as defined by total water) exceeds 83.3%. Therefore one might expect a 388 correlation between the model's extratropical cloud maxima (storm tracks) in the model and 83.3% 389 RH contours. But cloud formation is based on instantaneous RH, not the long-term averages shown 390 in Fig. 4c where the storm tracks are roughly co-located with the 75% RH contours. Higher RH 391 values may occur equatorward of a given RH contour. Allowing for time variability in RH renews 392 the possibility of a connection between the location of the storm tracks and RH distribution because 393

of RH<sub>c</sub>. This possible connection is explored with the RHc100 run, where the cloud scheme is adjusted to require essentially 100% grid-mean RH for cloud formation.

In the RHc100 case, the entire WV/CC cycle slows down significantly compared to the Cloud 396 case (see Fig. 2b and c). Since clouds are now unlikely to form and remove moisture from the 397 atmosphere below 850 hPa (where the air is generally nearly, but not quite, saturated), surface 398 evaporation decreases (Fig. 5a). RESS play less of a role as WV sources, approximately half of 399 both the magnitude and percentage as in the Cloud case, and become nearly non-existent in the 400 extratropics. LS condensation decreases as a WV sink and CC source; the slowdown increases the 401 WV residence time by 1.6 days or 13% (Table 2). This slowdown ultimately leads to a general 402 increase in steady-state RH (Fig. 5d) for reasons discussed at length with the formation perturbation 403 in section 3d. 404

CE decreases only slightly (3%) despite the intense perturbation in condensation. CE is not a 405 measure of how fast WV condenses, but simply whether it eventually does (in the given region 406 which here is the STET region). Similarly, FE decreases by 3% with a greater weakening of 407 formation processes than condensation (see Fig. 5b). FE represents the likelihood that a water 408 molecule, once it condenses, forms precipitation. Here, FE decreases since LS evaporation and 409 sublimation have increased both in value and as a percentage of LS condensation/deposition. In the 410 RHc100 setup, once a cloud is formed, if it persists to another time-step where RH has decreased 411 (as from precipitation), the remaining cloud condensate must entirely re-evaporate/sublimate. In 412 contrast, in the Cloud case, only enough cloud condensate to match the RH-based PDF must 413 evaporate, as long as grid-box-mean RH is above 83.3%. 414

The most significant change in efficiencies is SE which increases from 79.7% to 89.4% resulting in an amplification in PE (= FE \* SE) from 78.3% to 85.1%. SE increases because of the drastic decrease in RESS from both decreased precipitation formation (Fig. 5c) as well as increased

steady-state RH (Fig. 5d). While RH increases everywhere, RH is most significantly increased 418 in regions where cloud formation at less than 100% RH had reduced the amount of WV from 419 being transported. Once a moist parcel (traveling largely poleward/upward) reaches a cold enough 420 temperature such that the required RH is reached, excess water vapor is condensed. Thus, 100% 421 RH required for condensation more WV is isentropically transported to the polar upper troposphere 422 (and other cold regions of high RH) before clouds are formed. Weakened RESS results from less 423 precipitation falling through moister air, especially in the extratropics where the increase in RH 424 is most significant. Ultimately, despite increased PE, there is a 10% reduction in STET surface 425 precipitation (Table 2) potentially driven by decreased CC in the boundary layer (discussed below). 426 In addition to an increase in RH, with the RHc100 setup, CF is significantly amplified in the 427 polar extratropics (Fig. 5e). With seemingly more difficult conditions for cloud formation, CF 428 increases everywhere (above 850 hPa). This can be understood by considering what triggers cloud 429 formation in the cloud scheme: high values of RH. The increase in average RH noted previously 430 does in fact correspond to a rise in occurrences of high RH as shown through a histogram of daily 431 RH values (Fig. 5g) where values in the [100%, 105%] bin increase drastically, but all other values 432 decrease slightly. A histogram of daily CF values (Fig. 5h) shows a decrease in CF values below 433 65% and a drastic rise in occurrences of the highest values with the final bin being the highest 434 populated. (Note that while RH values greater than 100% are possible, by definition, 100% is the 435 maximum possible CF value such that the final CF histogram bin represents values of exactly 100% 436 CF.) With 100% grid-mean RH required for cloud formation, when cloud formation is triggered it 437 must be 100% CF at the time-step of the model. These histograms were further broken down by 438 meridional and vertical flow directions (not shown). Poleward and upward flows accounted for the 439 highest RH values and thus the higher CF values, but overall the stratified histograms painted the 440 same picture. For every direction of flow, the RHc100 perturbation requires greater RH for cloud 441

formation, increasing high RH values and thus CF. Accordingly, the location of maximum storm 442 track cloudiness shifts poleward (to areas of greater RH) from ~  $50^{\circ}$  (Fig. 4c) to ~  $60^{\circ}$  (not shown). 443 While CF increases significantly, the change in CC in the free troposphere is small, and in most 444 places is a decrease as seen in Fig. 5f. (A significant loss of CC below 850 hPa not shown is a 445 result of the region being generally unsaturated, since surface evaporation is associated with a time 446 scale.) While changes in CF and CC need not totally align, such drastic differences are surprising 447 and are, in fact, largely an artifact of altering the macrophysics in a way that is unexpected by 448 the microphysics scheme. With the RHc100 condition, if clouds form in a grid cell, the grid cell 449 CF is 100%. Yet with higher CF, autoconversion decreases. In the microphysics scheme, the rate 450 of change of cloud liquid due to autoconversion is proportional to  $CF * (LCC/CF)^{(7/3)}$  or, in a 451 frequently-invoked limiter,  $\ln(LCC/CF) * LCC$  (see Rotstayn 1997). In other words, if CC is more 452 widely distributed over a higher CF, it triggers less autoconversion. So a rise in CF, unmatched by 453 an increase in CC (since CC is in fact more difficult to form with the RHc100 condition), causes a 454 decrease in autoconversion leading to a cycle slowdown as expected. This result highlights both 455 the non-interchangeability of CC and CF as cloud tracers and the importance of considering the 456 details of a microphysics scheme when evaluating the usefulness of performing drastic alterations. 457 The bigger picture highlighted by the RHc100 case is the significance of the storm tracks 458 interrupting isentropic flow and the way in which details of the macrophysics scheme can thus have 459 such significant effects. (Accounting for such phenomena is lacking in advection-condensation 460 theory.) Here, sub-grid-scale RH has a significant effect on extratropical clouds by affecting the 461 storm track locations and altering the frequency of high-RH values. Re-located storm tracks could 462 also have significant effects on shortwave radiation not explored here, contributing to the usefulness 463 of  $RH_c$  as a tuning parameter for radiative balance. A potential emergent constraint on storm track 464 response (which varies significantly in GCMs as noted in Bender et al. 2012) could inform  $RH_c$ 465

choice. Thus, the RHc100 case also emphasizes the additional, non-radiative, impacts of tuning through  $RH_c$ , particularly on redistributing WV and precipitation.

# 468 c. Sedimentation Perturbation: noRESS

As described previously, one of the most noteworthy differences between saturation adjustment 469 and a full cloud scheme is the addition of two significant sources of WV: RESS. As seen in Fig. 1, 470 column-integrated RESS have a significant presence at all latitudes, providing an even stronger 471 source of WV than surface evaporation poleward of approximately  $50^{\circ}$ . Fig. 2b shows that together 472 they contribute approximately 17% to STET WV sources. RESS define SE as shown in Fig. 3 473 with one-fifth of formed precipitation lost to RESS. Fig. 6a depicts the changes in WV tendencies 474 when RESS are no longer present in the Cloud scheme. While surface evaporation increases, the 475 elimination of RESS yields a net decrease in WV sources (Fig. 2d). Matching this decrease, a 476 reduction in LS condensation/deposition (WV sinks) is spatially correlated both latitudinally and 477 vertically with the eliminated RESS (Fig. 6b). Thus, as in the RHc100 case, WV and CC cycling 478 is weakened: the total WV/CC sources or sinks in noRESS are 13-16% less than in the Cloud case, 479 while still greater than in the Base case (see Fig. 2). However, at the same time, the residence time 480 of a water molecule in the atmosphere is decreased by 7% due to the elimination of RESS as WV 481 sources which come from recycled hydrometeors. 482

<sup>483</sup> Without RESS as sinks of precipitation, STET precipitation increases by ~ 5% (8% globally) <sup>484</sup> as seen in Fig. 6c and Table 2. By definition, without RESS, SE is 100%. As FE is nearly <sup>485</sup> unchanged, PE increases drastically from 78.3% to 97.9% with a moderate increase in precipitation. <sup>486</sup> The elimination of snow sublimation corresponds strongly with the pattern and magnitude of a <sup>487</sup> decrease in ice settling yielding only a slight change in precipitation poleward of 45°. However, in <sup>488</sup> the subtropics, the elimination of rain evaporation is unmatched by decreases in autoconversion and accretion, so the precipitation increase is mostly subtropical, while the storm tracks are virtually
 unaffected.

This feature can be rationalized by considering the location of WV sources and sinks and 491 the connection between these budgetary terms and the steady-state fields. From a steady-state 492 perspective, the role of RESS in redistributing WV and moistening the atmosphere can be seen 493 in Fig. 6d. Turning off RESS results in a significant decrease in RH (up to 6%), especially in the 494 subtropics and the polar lower troposphere. Additional experiments were performed with RESS 495 turned off locally, including only between 15° and 45° or elsewhere (not shown). These runs 496 resulted in RH being only reduced (with any significance) in the regions where RESS is turned 497 off, demonstrating the local nature of the contribution of RESS to moisture. In redistributing WV, 498 RESS also play a significant role in the cloud distribution. Without RESS, both CF and CC decrease 499 globally as shown in Fig. 6e-f. The change in CF is of a similar pattern to the change in RH in 500 the polar extratropics, while the change in CC is more concentrated in the storm tracks (where CC 501 is larger to begin with). RH and CF changes are directly connected, as confirmed by considering 502 histograms of extratropical RH and CF (Fig. 6g-h). The noRESS case shifts occurrences of 503 RH away from higher values (>95%) in the extratropical free troposphere corresponding with a 504 decrease in CF concentrated where RH values are highest to begin with. 505

The connection between budgetary and steady-state changes is nuanced. Globally, the general reduction in RH is to be expected since the lack of RESS results in a drying of the boundary layer. This drying triggers more surface evaporation, but no other sources of WV. Decreased higher values of RH leads to decreased clouds. But, spatially, the areas of largest RH change (free troposphere, especially the polar extratropics) do not coincide with the locations of largest RESS tendency. RESS together provide a significant source of WV throughout the boundary layer and free troposphere, especially in the tropics (not shown). However, while RESS are smallest in the extratropics, its relative importance as a source of WV is greatest there (see Fig. 1a). While surface evaporation can easily increase below 850 hPa to replace RESS as a source of WV in the boundary layer (which is always nearly saturated), its ability to replenish moisture above 850 hPa depends on circulation. The rising motions induced by the Hadley circulation in the tropics allow humidity (and thus clouds) to be less affected by the loss of RESS. In contrast, in the polar regions where less vertical motion takes place and horizontal transport is more important for WV, the lower troposphere above 850 hPa experiences significant drying.

Thus, in the storm tracks and high latitudes, the increase in precipitation is small since the 520 elimination of RESS dries the region creating two opposing effects. Precipitation is increased 521 since SE is now 100%, but this increase is nearly balanced by a reduction in precipitation due to 522 less moisture and thus fewer precipitating clouds in the region. However, in the subtropics and mid 523 latitudes, the direct increase in precipitation is largely unbalanced since clouds are less affected 524 (as clouds are few to begin with so humidity decreases have little effect). This local role of RESS 525 is further seen in the fact that P-E (Fig. 6c) remains largely unchanged. Ultimately, the role of 526 RESS in the free troposphere is to increase RH (and ultimately clouds) by providing an additional 527 source of WV, while decreasing precipitation and—to a much greater extent—the PE through the 528 introduction of an atmospheric sink for hydrometeors. 529

# <sup>530</sup> *d. Formation Perturbation: halvAUTO*

In the halvAUTO case, autoconversion decreases in the STET region by 29%. Accretion and ice settling increase by 19% and 4%, respectively, to keep total STET CC sinks only 3% less than in the Cloud case. This re-balancing can be be conceptualized as weakened autoconversion causing more cloud liquid to be present to be scavenged by ice through accretion and subsequently settling. Similarly, in the doubAUTO case, STET autoconversion increases by 34%, accretion decreases by <sup>536</sup> 22%, and ice settling increases by 6%, such that total CC sinks are only 3% more than in the Cloud <sup>537</sup> case. These changes can be seen in Fig. 2e and f. In both cases the relative balance of the WV <sup>538</sup> sources and sinks is roughly unchanged. Noting the parallel opposing changes in halvAUTO and <sup>539</sup> doubAUTO, we focus primarily on halvAUTO.

Fig. 7a shows that latitudinally the WV balance is unchanged with decreases in LS condensation, surface evaporation, and rain evaporation balancing each other. Similarly, the CC balance (Fig. 7b) stays latitudinally unchanged with a decrease in LS condensation balanced by the net decreases in CC sinks. The opposing changes in autoconversion and accretion are similar in their spatial pattern, but the decrease in autoconversion is stronger, resulting in less precipitation as shown in Fig. 7c. These changes are principally equatorward of 60° since that is where autoconversion is most significant in the first place (Fig. 1b).

Across the STET region, precipitation decreases in the halvAUTO case by 3% and increases in 547 the doubAUTO case by 4%, similar to how the strength of the WV/CC cycle changes. From an 548 efficiency perspective (see Table 2), CE and FE change slightly in the same direction as changes 549 in precipitation, decreasing in halvAUTO in line with a cycle slowdown. SE also changes slightly 550 but in the opposite way: with decreased net formation but a proportionally larger decrease in 551 RESS in the halvAUTO case, SE increases slightly. The FE and SE effects balance such that PE 552 is minimally affected. This finding holds true for smaller and larger alterations to autoconversion, 553 accretion, and N except when an artificial decrease in a process is so large that the other processes 554 cannot keep the WV/CC cycle roughly constant. For example, when autoconversion is completely 555 eliminated, total STET CC sinks decrease by 6% as accretion cannot come close to making up for 556 the difference reducing FE to 90.4% and PE to 72.1%. However, apart from such limiting cases, 557 changes in budgetary terms and efficiencies are roughly linear. The residence time increases with 558 halvAUTO with weakened precipitation since a water molecule now spends a longer time in the 559

atmosphere as CC before precipitating, while the doubAUTO case shows a corresponding decrease
 in residence time.

From a steady-state perspective, in the halvAUTO case, RH, CF, and CC all increase as shown in 562 Fig. 7d-f. The significant changes are spatially similar, concentrated equatorward of 60° (where the 563 net decrease in CC sinks was strongest) and below  $\sim$ 500 hPa, peaking in the storm tracks. These 564 steady-state changes described are qualitatively opposite in the doubAUTO case (not shown). 565 Of note, the steady-state RH and cloud fields change not in response to a shift in the balance 566 between autoconversion and accretion, but in response to changes in total sources/sinks. When 567 WV/CC cycling strengthened due to increased autoconversion, increased accretion, or decreased 568 N, a reduction of RH, CF, and CC resulted. Opposite changes are associated with WV/CC cycling 569 weakening. Re-balancing autoconversion and accretion must have a relatively innocuous effect on 570 RH and clouds in and of itself. 571

Why does a weakened (strengthened) cycle increase (decrease) RH and clouds? It is important 572 to note that this generalization does not extend past these perturbations. (The pattern is followed 573 in the RHc100 case discussed previously but not in the noRESS case, possibly because of the 574 significant spatial and physical differences resulting from replacing RESS as WV sources with 575 enhanced surface evaporation.) However, in the absence of other changes (such as adding sources 576 and sinks from the Base to the Cloud case), a longer (shorter) residence time for a water molecule in 577 the atmosphere could be expected to correspond to an increase (decrease) in the steady-state fields 578 which represent the forms that a water molecule takes as it resides in the atmosphere. Additionally, 579 steady-state RH is directly connected to the WV cycle through surface evaporation since it is 580 formulated as a function of subsaturation. RH is connected to CF as demonstrated by considering 581 histograms of RH and CF (Fig. 7g and h): the halvAUTO case slightly shifts occurrences of RH 582

toward the highest values (>100%). Without any significant changes to the cloud physics beyond 583 a re-balancing of autoconversion and accretion, CC can logically be expected to follow CF. 584 Thus, the formation perturbations demonstrate the resilience of this cloud microphysics scheme 585 to changes in the balance of formation tendencies in terms of PE. Additionally, the general patterns 586 for steady-state consequences of the WV/CC cycle weakening (strengthening) emerge showing how 587 steady-state fields are affected by changes in residence time. A weakened (strengthened) cycle, 588 apart from other changes in cloud physics, leads to an increased (decreased) residence time and 589 increased (decreased) steady-state RH, CC, and CF. 590

# **4. Discussion and Conclusions**

# 592 a. Summary

The general picture that emerges from this idealized modeling study is that circulation sets the 593 basic pattern of moisture and precipitation, as seen through the first order similarity between the 594 two control cases. In the perturbation runs, details of the physics of condensation and sedimenta-595 tion also have substantial effects on humidity, clouds, and precipitation. However, it is noteworthy 596 that while RH does differ substantially (in certain extratropical regions) between the control cases, 597 precipitation does not, as the precipitation changes in the condensation and sedimentation per-598 turbations (RHc100 and noRESS) are of opposing sign. A secondary picture is the utility of 599 this idealized GCM for understanding physical controls of free tropospheric clouds and responses 600 to perturbations since key processes can be cleanly isolated. The saturation adjustment scheme 601 (Base case) shows gross RH features, as expected from advection-condensation theory, but cloud 602 processes refine the features. In particular, cloud macrophysics are important since thresholds 603 for cloud formation change cloud distribution (including the CF/CC ratio) and hence high RH 604

and storm track location due to isentropic transport of moisture as shown in the RHc100 run. Cloud microphysics are equally important, adding a key component through the re-evaporation of hydrometeors (RESS) changing RH values by a similar magnitude, as much as 5-6%. However, the formation perturbations demonstrate that the balance of precipitation-forming processes (here autoconversion and accretion) have little significance for RH, cloudiness, precipitation, and especially PE.

# 611 b. Advection-Condensation Theory

As was discussed previously, there are, on the surface, three differences between a saturation 612 adjustment scheme (or advection-condensation theory) and a full cloud scheme: RH<sub>c</sub>, RESS, and 613 the presence of CC which can be advected and/or subject to LS evaporation/sublimation. The first 614 two differences are here individually directly removed, but the third must be explored as a residual 615 in the *RHc100\_noRESS* experiment where we remove the RH<sub>c</sub> and RESS effects together from 616 the Cloud case. If these three identified differences are exhaustive, RHc100\_noRESS represents 617 the effect of adding CC to the Base case. Additionally, if the  $RH_c$  and RESS effects are linearly 618 additive, we can mathematically manipulate the various experiments to isolate the separate effects 619 of RH<sub>c</sub> and RESS added to the Base case (as opposed to removing these effects from the Cloud case 620 as was described in the Results section). To this end, Fig. 8 explores to what extent the  $RH_c$  and 621 RESS effects are linearly additive, to what extent they can explain the full difference between the 622 Base and Cloud controls, and the characteristics of the residual differences which can be attributed 623 to CC advection. 624

The RHc100 run includes RESS and advection effects, the noRESS run includes RHc and advection effects, and the RHc100\_noRESS run is just the advection effect. So we can test for linearity of the RH<sub>c</sub> and RESS effects by comparing RHc100 plus noRESS minus RHc100\_noRESS

(Fig. 8a). The combination appears to be mostly linear except in the free tropospheric high latitudes 628 where both RHc100 and noRESS runs had significant, but opposing, effects. RHc100 leads to 629 moistening and noRESS to drying; linear addition over-emphasizes drying or under-emphasizes 630 moistening. A possible mechanism is that when both are implemented, there is less moisture (from 631 noRESS) to be exported to the high latitudes (in RHc100), but this effect should be minimal as 632 noRESS minimally dries the boundary layer. A more like explanation is that since in RHc100, 633 RESS together decrease by over 50%, the noRESS drying effect is dampened when combined. But 634 since they combine nearly linearly, we can separately analyze the three effects of adding a cloud 635 scheme to a saturation adjustment scheme. 636

When adding a cloud scheme to a saturation adjustment scheme, advection and LS evapora-637 tion/sublimation (and any other residual effects, for example, nucleation barrier and incomplete 638 fallout in cirrus as noted by Liu et al. (2010)) moistens the free tropospheric subtropics and mid-639 latitudes (Fig. 8b) as well as the polar stratosphere. Implementing a  $RH_c$  of 83.3% dries the high 640 latitudes (Fig. 8c) by allowing for more condensation and precipitation of moisture before it is 641 isentropically transported to the poles. Finally RESS moisten the free troposphere, most strongly 642 in the storm tracks and lower polar regions (Fig. 8d), by adding an additional source of WV above 643 the boundary layer. 644

Thus, this work highlights the key deficiencies with an advection-condensation paradigm. The relatively small residual effects seen when comparing RHc100\_noRESS minus Base to Cloud minus base (Fig. 8b) suggest that RH<sub>c</sub> and RESS are the key ways in which a cloud scheme alters the RH distribution from advection-condensation theory alone, in the absence of cloud processes altering the circulation through latent heat release or cloud radiative effects. The RESS effect is a cloud microphysical effect already noted as missing from the advection-condensation paradigm and important to moistening the subtropics. But here we also highlight its importance for

moistening the polar regions where less vertical motion makes surface evaporation less effective 652 at moistening the free troposphere. In contrast,  $RH_c$  is a macrophysical effect, an artifact of 653 parameterizations attempting to represent the RH variability present in the real world. Here we 654 emphasize the importance of considering sub-grid-scale humidity distribution to allow clouds to 655 form in appropriate latitudinal locations (a problem that increased resolution alone may not fix). 656 As Sherwood et al. (2010) noted, these components of why the advection-condensation paradigm 657 is inadequate are critical to understand in order to accurately model not just climatological values, 658 but importantly changes in RH (and hence clouds and precipitation) with warming. 659

# 660 c. Outlook

The picture presented here is likely to change significantly with warming. While the advection-661 condensation paradigm suggests that free tropospheric RH is unlikely to change significantly with 662 uniform warming (Sherwood et al. 2010), the specific deficiencies of advection-condensation 663 theory explored here confound predicting changes in RH with warming, already complicated by 664 non-uniform warming. Any changes in RH could also have implications for P-E changes, as the 665 wet-get-wetter paradigm (Held and Soden 2006) is predicated on unchanged lower-tropospheric RH 666 and flow. Sherwood et al. (2014) identified a mixing-induced low cloud feedback where enhanced 667 mixing with warming dehydrates the boundary layer. Here, as in advection-condensation theory, 668 we highlighted the connection between subtropical boundary layer humidity and polar upper 669 tropospheric humidity because of eddy isentropic transport. In addition to the complications of 670 dynamical effects, because of the Clausius-Clapeyron relation, WV transport is expected to increase 671 with warming for thermodynamic reasons (Lavers et al. 2015). And as noted in the introduction, 672 replacement of ice with liquid in mixed-phase clouds with warming may also effect moisture and 673 cloud distribution through changes in precipitation efficiency. Thus, modeling the mechanisms 674

<sup>675</sup> controlling extratropical humidity and clouds accurately is critical for confidently forecasting future
 <sup>676</sup> change.

Our perturbation results demonstrate the significance of key processes for defining steady-state 677 patterns of humidity and cloudiness, implying a strong need to constrain processes such as RESS and 678 sub-grid-scale RH in order to ensure the physical grounding of parameterizations so that responses 679 to altered forcings will also be physical. For example, in using RHcrit as a GCM tuning parameter, 680 the multiple ways in which it effects radiative balance which—such as through shifted storm track 681 cloud maxima and opposing changes in CF and CC—should be carefully considered, especially 682 as they may be obscured or amplified by dynamical effects. Additionally, while *accr/auto* (or 683 N) was not important here in terms of affecting steady-state fields or average precipitation, it is 684 likely to have other effects as discussed in the introduction, including modulating the intensity 685 of precipitation events. Our results suggest that the strength of warm rain processes as a whole 686 (accretion+autoconversion) plays a role in defining RH, clouds, and precipitation distribution and 687 thus is an important parameter to constrain, not just *accr/auto*. By separately analyzing the effects 688 on CF and CC and their connection to changes in RH and various components of the water cycle, 689 this study highlighted the need to carefully dissect the physical mechanisms for change instead 690 of relying on generalizations. For example, as demonstrated in the RHc100 perturbation, cloud 691 response cannot be directly predicted from changes in average RH. Relationships among RH, CF, 692 and CC in a cloud scheme may be nonintuitive and are certainly nontrivial. CC and CF have 693 varying levels of importance for cloud radiative effects depending on regime and saturation, so 694 individual, local effects are consequential. 695

<sup>696</sup> Comparing the significance of various controls of clouds cannot be precise in this idealized, <sup>697</sup> decoupled framework. Nor does this study explore the relative significance of various cloud <sup>698</sup> feedbacks to anthropogenic forcings. Yet, by allowing for a detailed exploration of cloud physics

decoupled from circulation, this type of idealized model could play a key role in the model 699 hierarchy for reducing uncertainty surrounding cloud feedback. In comprehensive GCMs with 700 coupled feedbacks, circulation feedbacks (particularly shifts in the extratropical jets) have been 701 demonstrated to be less significant than thermodynamic mechanisms of mixed-phase clouds in 702 creating the shortwave extratropical cloud feedback (Wall and Hartmann 2015; Ceppi and Hartmann 703 2016). This finding suggests that cloud parameterization mechanisms relating to mixed-phase 704 clouds may play a significant role in constraining extratropical cloudiness, an area explored in 705 related work with the idealized setup used in this paper (Frazer and Ming 2022). 706

In summary, this study takes a step forward in elucidating physical mechanisms controlling extratropical clouds, while highlighting the importance of identifying and adequately representing these mechanisms in order to accurately simulate the cloud feedbacks associated with climate change.

Acknowledgments. The authors acknowledge Nadir Jeevanjee, David Paynter, and Daniel McCoy
 for helpful feedback; three anonymous reviews were also beneficial. M.E.F. was supported by award
 NA18OAR4320123 from the National Oceanic and Atmospheric Administration, U.S. Department
 of Commerce, and award AWD1005319 from the National Science Foundation.

<sup>715</sup> *Data availability statement*. The output from the simulations described in this manuscript is
 <sup>716</sup> archived at the Geophysical Fluid Dynamics Laboratory and is available upon request.

#### 717 **References**

Bender, F. A. M., V. Ramanathan, and G. Tselioudis, 2012: Changes in extratropical storm track
 cloudiness 1983–2008: observational support for a poleward shift. *Climate Dyn.*, 38, 2037–2053,
 doi:10.1007/s00382-011-1065-6.

Bony, S., and Coauthors, 2015: Clouds, circulation and climate sensitivity. *Nat. Geosci.*, 8, 261–
 268, doi:10.1038/ngeo2398.

Ceppi, P., F. Brient, M. D. Zelinka, and D. L. Hartmann, 2017: Cloud feedback mechanisms and

- their representation in global climate models. WIREs Clim. Change, 8, doi:10.1002/wcc.465. 724 Ceppi, P., and D. L. Hartmann, 2016: Clouds and the atmospheric circulation response to warming. 725 J. Climate, 29, 783–799, doi:10.1175/JCLI-D-15-0394.1. 726 Ceppi, P., D. L. Hartmann, and M. J. Webb, 2016: Mechanisms of the negative shortwave cloud 727 feedback in middle to high latitudes. J. Climate, 29, 139–157, doi:10.1175/JCLI-D-15-0327.1. 728 Frazer, M. E., and Y. Ming, 2022: Understanding the extratropical liquid water path feedback 729 in mixed-phase clouds with an idealized global climate model. J. Climate, 2391–2406, doi: 730 10.1175/JCLI-D-21-0334.1. 731
  - Frierson, D. M. W., I. M. Held, and P. Zurita-Gotor, 2006: A gray-radiation aquaplanet moist GCM.
     Part I: Static stability and eddy scale. *J. Atmos. Sci.*, 63, 2548–2566, doi:10.1175/JAS3753.1.
  - <sup>734</sup> Galewsky, J., A. Sobel, and I. M. Held, 2005: Diagnosis of subtropical humidity dynamics using
     <sup>735</sup> tracers of last saturation. *J. Atmos. Sci.*, **62**, 3353–3367, doi:10.1175/JAS3533.1.
  - <sup>736</sup> Geoffroy, O., S. C. Sherwood, and D. Fuchs, 2017: On the role of the stratiform cloud scheme
     <sup>737</sup> in the inter-model spread of cloud feedback. *J. Adv. Model. Earth Syst.*, **9**, 423–437, doi:
     <sup>738</sup> 10.1002/2016MS000846.
  - Gettelman, A., H. Morrison, C. R. Terai, and R. Wood, 2013: Microphysical process rates and global aerosol-cloud interactions. *Atmos. Chem. Phys.*, **13**, 9855–9867, doi:10.5194/ acp-13-9855-2013.

- Gettelman, A., H. Morrison, C. R. Terai, and R. Wood, 2014: Corrigendum to "Microphysical
   process rates and global aerosol-cloud interactions". *Atmos. Chem. Phys.*, 14, 9099–9103, doi:
   10.5194/acp-14-9099-2014.
- Held, I. M., 2005: The gap between simulation and understanding in climate modeling. *Bull. Amer. Meteor. Soc.*, 86, 1609–1614, doi:10.1175/BAMS-86-11-1609.
- Held, I. M., 2014: Simplicity amid complexity. *Science*, **343**, 1206–1207, doi:10.1126/science.
   1248447.
- Held, I. M., and T. Schneider, 1999: The surface branch of the zonally averaged mass transport
   circulation in the troposphere. *J. Atmos. Sci.*, 56, 1688–1697, doi:10.1175/1520-0469(1999)
   056<1688:TSBOTZ>2.0.CO;2.
- Held, I. M., and B. J. Soden, 2006: Robust responses of the hydrological cycle to global warming.
   *J. Climate*, **19**, 5686–5699, doi:10.1175/JCLI3990.1.
- <sup>754</sup> Held, I. M., and M. J. Suarez, 1994: A proposal for the intercomparison of the dynamical
   <sup>755</sup> cores of atmospheric general circulation models. *Bull. Amer. Meteor. Soc.*, **75**, 1825–1830,
   <sup>756</sup> doi:10.1175/1520-0477(1994)075<1825:APFTIO>2.0.CO;2.
- Jeevanjee, N., P. Hassanzadeh, S. Hill, and A. Sheshadri, 2017: A perspective on climate model hierarchies. *J. Adv. Model. Earth Syst.*, **9**, 1760–1771, doi:10.1002/2017MS001038.
- Jiang, H., G. Feingold, and A. Sorooshian, 2010: Effect of aerosol on the susceptibility and efficiency of precipitation in warm trade cumulus clouds. *J. Atmos. Sci.*, **67**, 3525–3540, doi: 10.1175/2010JAS3484.1.

| 762 | Jing, X., K. Suzuki, H. Guo, D. Goto, T. Ogura, T. Koshiro, and J. Mülmenstädt, 2017: A multi- |
|-----|--|
| 763 | model study on warm precipitation biases in global models compared to satellite observations.  |
| 764 | J. Geophys. Res. Atmos., 122, 11 806–11 824, doi:10.1002/2017JD027310.                         |

- Kelly, K., A. Tuck, and T. Davies, 1991: Wintertime asymmetry of upper tropospheric water 765 between the Northern and Southern Hemispheres. Nature, 353, 244-247, doi:10.1038/353244a0. 766
- Klein, S. A., and Coauthors, 2009: Intercomparison of model simulations of mixed-phase clouds 767 observed during the ARM Mixed-Phase Arctic Cloud Experiment. I: Single-layer cloud. Quart. 768 J. Roy. Meteor. Soc., 135, 979–1002, doi:10.1002/qj.416.
- Laliberté, F., T. Shaw, and O. Pauluis, 2012: Moist recirculation and water vapor transport on dry 770 isentropes. J. Atmos. Sci., 69, 875-890, doi:10.1175/JAS-D-11-0124.1. 771
- Langhans, W., K. Yeo, and D. M. Romps, 2015: Lagrangian investigation of the precipitation 772 efficiency of convective clouds. J. Atmos. Sci., 72, 1045–1062, doi:10.1175/JAS-D-14-0159.1. 773
- Lavers, D. A., F. M. Ralph, D. E. Waliser, A. Gershunov, and M. D. Dettinger, 2015: Climate 774 change intensification of horizontal water vapor transport in CMIP5. Geophys. Res. Lett., 42, 775 5617-5625, doi:10.1002/2015GL064672. 776
- Liu, Y. S., S. Fueglistaler, and P. H. Haynes, 2010: Advection-condensation paradigm for strato-777 spheric water vapor. J. Geophys. Res. Atmos., 115, D24 307, doi:10.1029/2010JD014352. 778
- McCoy, D. T., D. L. Hartmann, and M. D. Zelinka, 2018: Mixed-phase cloud feedbacks. Mixed-779
- *Phase Clouds: Observations and Modeling*, C. Andronache, Ed., Elsevier, 215–236, doi:10. 780
- 1016/B978-0-12-810549-8.00009-X. 781

McCoy, D. T., D. L. Hartmann, M. D. Zelinka, P. Ceppi, and D. P. Grosvenor, 2015: Mixed-phase
 cloud physics and Southern Ocean cloud feedback in climate models. *J. Geophys. Res. Atmos.*,
 120, 9539–9554, doi:10.1002/2015JD023603.

McCoy, D. T., I. Tan, D. L. Hartmann, M. D. Zelinka, and T. Storelvmo, 2016: On the relationships
 among cloud cover, mixed-phase partitioning, and planetary albedo in GCMs. *J. Adv. Model. Earth Syst.*, 8, 650–668, doi:10.1002/2015MS000589.

<sup>788</sup> Michibata, T., and T. Takemura, 2015: Evaluation of autoconversion schemes in a single model
 <sup>789</sup> framework with satellite observations. *J. Geophys. Res. Atmos.*, **120**, 9570–9590, doi:10.1002/
 <sup>790</sup> 2015JD023818.

<sup>791</sup> Ming, Y., and I. M. Held, 2018: Modeling water vapor and clouds as passive tracers in an idealized <sup>792</sup> GCM. *J. Climate*, **31**, 775–786, doi:10.1175/JCLI-D-16-0812.1.

Pierrehumbert, R. T., 1998: Lateral mixing as a source of subtropical water vapor. *Geophys. Res. Lett.*, 25, 151–154, doi:10.1029/97GL03563.

Pierrehumbert, R. T., H. Brogniez, and R. Roca, 2007: On the relative humidity of the atmosphere.
 *The Global Circulation of the Atmosphere*, T. Schneider, and A. H. Sobel, Eds., Princeton
 University Press, 143–185.

<sup>798</sup> Quaas, J., and Coauthors, 2009: Aerosol indirect effects - general circulation model inter-<sup>799</sup> comparison and evaluation with satellite data. *Atmos. Chem. Phys.*, **9**, 8697–8717, doi: <sup>800</sup> 10.5194/acp-9-8697-2009.

Quaas, J., and Coauthors, 2020: Constraining the Twomey effect from satellite observations: issues and perspectives. *Atmos. Chem. Phys.*, **20**, 15 079–15 099, doi:10.5194/acp-20-15079-2020.

- Rotstayn, L. D., 1997: A physically based scheme for the treatment of stratiform clouds and
   precipitation in large-scale models. I: Description and evaluation of the microphysical processes.
   *Q. J. R. Meteorol. Soc.*, **123**, 1227–1282.
- Rotstayn, L. D., B. F. Ryan, and J. J. Katzfey, 2000: A scheme for calculation of the liquid fraction
  in mixed-phase clouds in large-scale models. *Mon. Wea. Rev.*, **128**, 1070–1088.
- <sup>808</sup> Shepherd, T. G., 2014: Atmospheric circulation as a source of uncertainty in climate change <sup>809</sup> projections. *Nat. Geosci.*, **7**, 703–708, doi:10.1038/ngeo2253.
- <sup>810</sup> Sherwood, S., S. Bony, and J.-L. Dufresne, 2014: Spread in model climate sensitivity traced to <sup>811</sup> atmospheric convective mixing. *Nature*, **505**, 37–42, doi:10.1038/nature12829.
- <sup>812</sup> Sherwood, S. C., R. Roca, T. M. Weckwerth, and N. G. Andronova, 2010: Tropospheric water <sup>813</sup> vapor, convection, and climate. *Rev. Geophys.*, **48**, doi:10.1029/2009RG000301.
- <sup>814</sup> Sherwood, S. C., and Coauthors, 2020: An assessment of Earth's climate sensitivity using multiple

lines of evidence. *Reviews of Geophysics*, **58**, e2019RG000678, doi:10.1029/2019RG000678.

Stephens, G. L., 1978: Radiation profiles in extended water clouds. II. Parameterization schemes.

J. Atmos. Sci., **35**, 2123–2132, doi:10.1175/1520-0469(1978)035<2123:RPIEWC>2.0.CO;2.

- <sup>818</sup> Stevens, B., and S. Bony, 2013: What are climate models missing? *Science*, **340**, 1053–1054, doi:10.1126/science.1237554.
- Sun, D.-Z., and R. S. Lindzen, 1993: Distribution of tropical tropospheric water vapors. J. Atmos.
   Sci., 50, 1643–1660, doi:10.1175/1520-0469(1993)050<1643:DOTTWV>2.0.CO;2.
- Trenberth, K. E., 1998: Atmospheric moisture residence times and cycling: Implications for rainfall rates and climate change. *Climatic Change*, **39**, 667–694.

| 824 | Wall, C. J., and D. L. Hartmann, 2015: On the influence of poleward jet shift on shortwave cloud |
|-----|--|
| 825 | feedback in global climate models. J. Adv. Model. Earth Syst., 7, doi:10.1002/2015MS000520.      |
| 826 | Wu, P., B. Xi, X. Dong, and Z. Zhang, 2018: Evaluation of autoconversion and accre-              |
| 827 | tion enhancement factors in general circulation model warm-rain parameterizations using          |
| 828 | ground-based measurements over the Azores. Atmos. Chem. Phys., 18, 17405-17420, doi:             |
| 829 | 10.5194/acp-18-17405-2018.   |
| 830 | Yang, H., and R. T. Pierrehumbert, 1994: Production of dry air by isentropic mixing. J. Atmos.   |
| 831 | <i>Sci.</i> , <b>51</b> , 3437–3454, doi:0.1175/1520-0469(1994)051<3437:PODABI>2.0.CO;2.         |
| 832 | Zhao, M., 2014: An investigation of the connections among convection, clouds, and climate        |
| 833 | sensitivity in a global climate model. J. Climate, 27, 1845–1862, doi:10.1175/JCLI-D-13-00145.   |
| 834 | 1.   |
| 835 | Zhao, M., I. M. Held, SJ. Lin, and G. A. Vecchi, 2009: Simulations of global hurricane climatol- |

- ogy, interannual variability, and response to global warming using a 50-km resolution GCM. J. 836
- Climate, 22, 6653-6678, doi:10.1175/2009JCLI3049.1. 837

# **LIST OF TABLES**

| 839 | Table 1. | Description of the experiments  |
|-----|----------|---|
| 840 | Table 2. | Summary of STET (15°-90°) precipitation variables: average surface precip-    |
| 841 |          | itation (P) and surface evaporation (E); condensation (CE), formation (FE),   |
| 842 |          | sedimentation (SE), and precipitation (PE) efficiencies; residence time (RT). |
| 843 |          | See text for definition of these variables                                    |

| Name          | Description  |
|---------------|--|
| Base          | control simulation with specific humidity tracer and saturation adjustment                               |
| Cloud         | control simulation with specific humidity and cloud tracers (liquid, ice, and fraction) and microphysics |
| RHc100        | variant of Cloud simulation requiring 100% grid-box-mean RH for cloud formation $(RH_c)$                 |
| noRESS        | variant of Cloud simulation without rain evaporation or snow sublimation                                 |
| halvAUTO      | variant of Cloud simulation halving the raw computed value for autoconversion at each time-step          |
| doubAUTO      | as halvAUTO, but doubling autoconversion   |
| RHc100_noRESS | variant of Cloud simulation combining both RHc100 and noRESS variations                                  |

# TABLE 1. Description of the experiments.

TABLE 2. Summary of STET (15°-90°) precipitation variables: average surface precipitation (P) and surface evaporation (E); condensation (CE), formation (FE), sedimentation (SE), and precipitation (PE) efficiencies; residence time (RT). See text for definition of these variables.

| run      | Р                      | Е    | CE   | FE   | SE   | PE   | RT   |
|----------|------------------------|------|------|------|------|------|------|
|          | mm day <sup>-1</sup> % |      |      |      |      | days |      |
| Base     | 1.84                   | 2.34 | 78.5 | -    | -    | -    | 13.1 |
| Cloud    | 1.91                   | 2.37 | 83.9 | 98.2 | 79.7 | 78.3 | 12.7 |
| RHc100   | 1.71                   | 2.17 | 81.2 | 95.2 | 89.4 | 85.1 | 14.3 |
| noRESS   | 2.00                   | 2.47 | 81.3 | 97.9 | 100. | 97.9 | 11.8 |
| halvAUTO | 1.84                   | 2.31 | 83.6 | 97.6 | 79.8 | 77.9 | 13.1 |
| doubAUTO | 1.98                   | 2.44 | 84.3 | 98.6 | 79.6 | 78.5 | 12.2 |

# **LIST OF FIGURES**

| <ul> <li>848</li> <li>849</li> <li>850</li> <li>851</li> <li>852</li> <li>853</li> <li>854</li> <li>855</li> <li>856</li> <li>857</li> <li>858</li> <li>859</li> </ul> | Fig. 1. | Comparison of zonally-averaged, column-integrated WV, CC, and precipitation (P) ten-<br>dency terms in control cases (black totals, blue sources, and red sinks). Cloud case terms<br>(depicted as indicated by the legends) shown are (a) total (WV), surface evaporation (Ev),<br>rain evaporation (RE), snow sublimation (SS), and net condensation (Co); (b) total (CC), net<br>condensation (Co), autoconversion (Au), accretion (Ac), and ice settling (IS); (c) total (P), net<br>formation (Form), net sinks (RESS), and moisture convergence (P-E, surface precipitation<br>minus evaporation). Base case terms (depicted as half-width lines and sometimes obscured<br>beneath their Cloud case counterparts)) shown are total WV, surface evaporation, saturation<br>adjustment as net condensation in (a) and precipitation in (c), and saturation adjustment<br>minus evaporation as P-E in (c). Units are $10^{-6}$ kg m <sup>-2</sup> s <sup>-1</sup> . A positive tendency value<br>denotes (a) WV, (b) CC, or (c) precipitation increasing. Totals include the less significant<br>tendency terms not shown individually.   | 44   |
|--|---------|--|------|
| 860<br>861<br>862<br>863<br>864<br>865<br>866<br>867   | Fig. 2. | Principal WV and CC sources and sinks for various model runs (see Table 1) represented as column-integrated average STET $(15^{\circ}-90^{\circ})$ tendency values. For clarity, the smallest terms are conglomerated in an other (O) category. Processes shown are WV sources: surface evaporation (Ev), rain evaporation (RE), and snow sublimation (SS); WV sink (CC source): LS condensation (Co); CC sinks: autoconversion (Au), accretion (Ac), and ice settling (IS). Base case saturation adjustment is labeled LS condensation. Tendency units (vertical axis) are $10^{-6}$ kg m <sup>-2</sup> s <sup>-1</sup> . Percentages are given with respect to total source or sink category and may not add to 100% due to rounding.  | . 45 |
| 868<br>869<br>870<br>871<br>872<br>873<br>874<br>875   | Fig. 3. | Diagram of the water cycle in the control cloud microphysics scheme (Cloud experiment). Water is cycled between four species (reservoirs): WV, CC, precipitation, and an assumed surface reservoir. The quantities shown are average STET $(15^{\circ}-90^{\circ})$ tendency values with units of $10^{-6}$ kg m <sup>-2</sup> s <sup>-1</sup> . Each reservoir shows either a balance (0.0) or an imbalance. Here, condensation comprises both LS condensation and deposition; evaporation comprises both LS evaporation and sublimation; formation includes autoconversion, accretion, ice settling, and melting of cloud ice to rain; and sedimentation represents formation processes minus RESS.  | . 46 |
| 876<br>877<br>878<br>879<br>880<br>881   | Fig. 4. | Key variables in control runs: (a) RH difference (Cloud minus Base, %) as shading and Base RH as contours (5% spacing), (b) temperature (K) as shading and potential temperature as contours (5K spacing), (c) Cloud CF (%) as shading and Cloud RH as contours (5% spacing), (d) Cloud total CC ( $10^{-6}$ kg kg <sup>-1</sup> ) as shading and liquid (solid) and ice (dashed) CC as contours ( $5 \times 10^{-6}$ kg kg <sup>-1</sup> spacing). Variables have been zonally averaged, and the x- and y-axes are latitude and pressure (hPa), respectively.   | 47   |
| 882<br>883<br>884<br>885<br>886<br>887<br>888<br>889<br>890<br>891<br>892<br>893   | Fig. 5. | Key variable changes in RHc100 perturbation from Cloud control: absolute differences in zonally averaged (a) WV, (b) CC, and (c) precipitation (P) tendency terms (y-axis units of $10^{-6}$ kg m <sup>-2</sup> s <sup>-1</sup> ); absolute differences in (d) RH, (e) CF, and (f) CC as shading with Cloud case values as contours (5%, 5%, and $5 \times 10^{-6}$ kg kg <sup>-1</sup> spacing, respectively); comparison of normalized histograms of (g) RH and (h) CF in Cloud (black) and RHc100 (grey) cases from daily data (x-axis units of %) between 15° and 90° and 850 and 250 hPa with the y-axis cut off at 0.15. For (a)-(c), WV, CC, and precipitation (P) tendency difference terms shown are as defined in Fig. 1, with units of $10^{-6}$ kg m <sup>-2</sup> s <sup>-1</sup> where a positive tendency difference denotes an increase in a WV/CC/P-increasing process or a decrease in a WV/CC/P-decreasing process. For (a)-(f) variables have been zonally averaged and the x-axis is latitude; for (d-f) the y-axis is pressure (hPa). For (g)-(h), histogram bins have widths of 5% and are all half-open except for the last bin: [0, 5), [5, 10),, [100, 105]. | 48   |

| • | 49 |
|---|----|
|   | 50 |
|   |    |
|   |    |
|   |    |
|   |    |
|   |    |
|   | 51 |
|   |    |



FIG. 1. Comparison of zonally-averaged, column-integrated WV, CC, and precipitation (P) tendency terms in 903 control cases (black totals, blue sources, and red sinks). Cloud case terms (depicted as indicated by the legends) 904 shown are (a) total (WV), surface evaporation (Ev), rain evaporation (RE), snow sublimation (SS), and net 905 condensation (Co); (b) total (CC), net condensation (Co), autoconversion (Au), accretion (Ac), and ice settling 906 (IS); (c) total (P), net formation (Form), net sinks (RESS), and moisture convergence (P-E, surface precipitation 907 minus evaporation). Base case terms (depicted as half-width lines and sometimes obscured beneath their Cloud 908 case counterparts)) shown are total WV, surface evaporation, saturation adjustment as net condensation in (a) 909 and precipitation in (c), and saturation adjustment minus evaporation as P-E in (c). Units are  $10^{-6}$  kg m<sup>-2</sup> s<sup>-1</sup>. A 910 positive tendency value denotes (a) WV, (b) CC, or (c) precipitation increasing. Totals include the less significant 911 tendency terms not shown individually. 912



FIG. 2. Principal WV and CC sources and sinks for various model runs (see Table 1) represented as columnintegrated average STET  $(15^{\circ}-90^{\circ})$  tendency values. For clarity, the smallest terms are conglomerated in an other (O) category. Processes shown are WV sources: surface evaporation (Ev), rain evaporation (RE), and snow sublimation (SS); WV sink (CC source): LS condensation (Co); CC sinks: autoconversion (Au), accretion (Ac), and ice settling (IS). Base case saturation adjustment is labeled LS condensation. Tendency units (vertical axis) are  $10^{-6}$  kg m<sup>-2</sup> s<sup>-1</sup>. Percentages are given with respect to total source or sink category and may not add to 100% due to rounding.



FIG. 3. Diagram of the water cycle in the control cloud microphysics scheme (Cloud experiment). Water is cycled between four species (reservoirs): WV, CC, precipitation, and an assumed surface reservoir. The quantities shown are average STET ( $15^{\circ}-90^{\circ}$ ) tendency values with units of  $10^{-6}$  kg m<sup>-2</sup> s<sup>-1</sup>. Each reservoir shows either a balance (0.0) or an imbalance. Here, condensation comprises both LS condensation and deposition; evaporation comprises both LS evaporation and sublimation; formation includes autoconversion, accretion, ice settling, and melting of cloud ice to rain; and sedimentation represents formation processes minus RESS.



FIG. 4. Key variables in control runs: (a) RH difference (Cloud minus Base, %) as shading and Base RH as contours (5% spacing), (b) temperature (K) as shading and potential temperature as contours (5K spacing), (c) Cloud CF (%) as shading and Cloud RH as contours (5% spacing), (d) Cloud total CC ( $10^{-6}$  kg kg<sup>-1</sup>) as shading and liquid (solid) and ice (dashed) CC as contours ( $5 \times 10^{-6}$  kg kg<sup>-1</sup> spacing). Variables have been zonally averaged, and the x- and y-axes are latitude and pressure (hPa), respectively.



FIG. 5. Key variable changes in RHc100 perturbation from Cloud control: absolute differences in zonally 931 averaged (a) WV, (b) CC, and (c) precipitation (P) tendency terms (y-axis units of 10<sup>-6</sup> kg m<sup>-2</sup> s<sup>-1</sup>); absolute 932 differences in (d) RH, (e) CF, and (f) CC as shading with Cloud case values as contours (5%, 5%, and  $5 \times 10^{-6}$ 933 kg kg<sup>-1</sup> spacing, respectively); comparison of normalized histograms of (g) RH and (h) CF in Cloud (black) 934 and RHc100 (grey) cases from daily data (x-axis units of %) between 15° and 90° and 850 and 250 hPa with 935 the y-axis cut off at 0.15. For (a)-(c), WV, CC, and precipitation (P) tendency difference terms shown are as 936 defined in Fig. 1, with units of  $10^{-6}$  kg m<sup>-2</sup> s<sup>-1</sup> where a positive tendency difference denotes an increase in a 937 WV/CC/P-increasing process or a decrease in a WV/CC/P-decreasing process. For (a)-(f) variables have been 938 zonally averaged and the x-axis is latitude; for (d-f) the y-axis is pressure (hPa). For (g)-(h), histogram bins have 939 widths of 5% and are all half-open except for the last bin: [0, 5), [5, 10), ..., [100, 105].



FIG. 6. As Fig. 5, but for noRESS perturbation.



FIG. 7. As Figs. 5 and 6, but for halvAUTO perturbation, except that the colorbar scale is reduced by a factor of 10 for (d) and (e).



FIG. 8. Comparison of absolute RH differences (%) between control cases and intermediate setups: (a) RHc100 plus noRESS minus RHc100\_noRESS minus Cloud [linearity check: should be 0 if  $RH_c = 83.3\%$  and RESS effects sum linearly], (b) RHc100\_noRESS minus Base [CC advection effect] as shading, (c) noRESS minus RHc100\_noRESS [ $RH_c = 83.3\%$  effect] as shading, (d) RHc100 minus RHc100\_noRESS [RESS effect] as color shading. All contours are Cloud minus Base difference with a spacing of 1%.