Conductivities of Titan's dusty ionosphere

Oleg Shebanits¹, Jan-Erik Wahlund², Jack Hunter Waite³, and Michele K. Dougherty⁴

¹Swedish Institute for Space Physics (IRF) ²Swedish Institute of Space Physics (IRF) ³Southwest Research Institute ⁴Imperial College

November 24, 2022

Abstract

Titan's ionosphere hosts a globally distributed non-trivial dusty ion-ion plasma, providing an environment for studies of dusty ionospheres that is in many aspects unique in our solar system. Thanks to the Cassini mission, Titan's ionosphere also features one of the largest dusty plasma datasets from 126 flybys of the moon over 13 years, from 2004 to 2017. Recent studies have shown that negatively charged dust dramatically alters the electric properties of plasmas, in particular planetary ionospheres. Utilizing the full plasma content of the moon's ionosphere (electrons, positive ions and negative ions/dust grains), we derive the electric conductivities and define the conductive dynamo region. Our results show that using the full plasma content increases the Pedersen conductivities at ~1300 km altitude by 20% compared to the earlier estimates without charged dust, while the Hall conductivities indicate a reverse Hall effect at ~900 km altitude (closest approach) and below. The dayside conductivities are shown to be factor ~7-9 larger than on the nightside, owing to higher dayside plasma densities.

Conductivities of Titan's dusty ionosphere

2 O. Shebanits¹, J-E. Wahlund¹, J. H. Waite Jr². and M. K. Dougherty³

- ³ ¹Swedish Institute of Space Physics, Uppsala, Sweden
- 4 ²Southwest Research Institute, San Antonio, USA
- ⁵ ³Imperial College London, London, UK
- 6 Corresponding author: Oleg Shebanits (<u>oleg.shebanits@irfu.se</u>)

7 Key Points:

1

- 8 Second peak of Pedersen conductivity in dusty plasma region of Titan's ionosphere
- Indications of reverse Hall effect near ~900 km altitude
- Up to x10 difference between dayside and nightside conductivities

11 Abstract

- 12 Titan's ionosphere hosts a globally distributed non-trivial dusty ion-ion plasma, providing an
- 13 environment for studies of dusty ionospheres that is in many aspects unique in our solar system.
- 14 Thanks to the Cassini mission, Titan's ionosphere also features one of the largest dusty plasma
- 15 datasets from 126 flybys of the moon over 13 years, from 2004 to 2017.
- 16 Recent studies have shown that negatively charged dust dramatically alters the electric properties
- 17 of plasmas, in particular planetary ionospheres. Utilizing the full plasma content of the moon's
- 18 ionosphere (electrons, positive ions and negative ions/dust grains), we derive the electric
- 19 conductivities and define the conductive dynamo region.
- 20 Our results show that using the full plasma content increases the Pedersen conductivities at
- \sim 21 \sim 1300 km altitude by 20% compared to the earlier estimates without charged dust, while the Hall
- 22 conductivities indicate a reverse Hall effect at ~900 km altitude (closest approach) and below.
- 23 The dayside conductivities are shown to be factor ~7-9 larger than on the nightside, owing to
- 24 higher dayside plasma densities.

25 Plain Language Summary

- 26 The largest moon of Saturn, Titan, is famous for its signature orange haze that is formed in the
- 27 top layer of its atmosphere ionosphere. The complex organic chemistry initiated mainly by
- sunlight forms grains of dust that at ~1000 km altitude reach a few nanometers in size
- 29 (comparable to finely ground flour). These grains of dust become charged in the ionosphere by
- 30 absorbing the free electrons (depleting them). Below ~1000 km altitude there are almost no
- 31 electrons left. In the absence of light electrons (negative charge), the positively charged ions
- 32 instead become the dominant mobile charge carrier, as the negatively charged dust is much
- heavier. Such a reversal of charge mobility has a large impact on the electric properties of an
- ionosphere, increasing its electric conductivity and changing the direction of its electric currents.
- 35 We use a dataset from the Cassini mission spanning an entire solar cycle and nearly half a Titan
- 36 year (≈ 15 Earth years) to calculate the electric conductivities of Titan's ionosphere and show that
- the dusty plasma increases them by 20% on average. Below 1000 km altitudes, we find

- 38 indications of a secondary conductive layer, a direct consequence of the dust grains absorbing
- 39 the electrons.

40 **1 Introduction**

41 Titan is immersed in Saturn's magnetosphere (Bertucci et al., 2008; Garnier et al., 2010) and

- 42 does not have its own magnetic field. Instead, an induced magnetic field is formed around Titan
- 43 from the interactions with Saturn's magnetosphere, similarly to interactions of Venus and Mars
- 44 with the solar wind and interplanetary magnetic field (e.g., Bertucci et al., 2011 and references
- therein). The interaction between Titan and the ambient magnetic field depends to a large degreeon Titan's conductive ionosphere. Previously, the electrical properties of Titan's ionosphere
- 47 were derived using only the ion and electron content (Rosenqvist et al., 2009). However, in the
- 48 later years, significant amounts of charged dust were detected in Titan's ionosphere (Ågren et al.,
- 49 2012; Coates et al., 2007; Shebanits et al., 2013) and a globally present dusty ion-ion plasma is
- 50 expected (Shebanits et al., 2016). The charged dust grains in the ionosphere-like plasma of
- 51 Enceladus plume (Morooka et al., 2011) and in the near-equatorial dusty ionosphere of Saturn
- 52 (Morooka et al., 2019) were shown to have a profound effect on its electric conductivities by
- 53 Simon et al. (2011) and Yaroshenko & Lühr (2016), and Shebanits et al. (2020), respectively.
- 54 In this work we derive the electrical properties of Titan's ionosphere using the full plasma
- 55 content: electrons, positive ions and negative ions/dust grains. The relevant in-situ measurements
- 56 used here are from the Radio and Plasma Wave Science Langmuir Probe (RPWS/LP, Gurnett et
- al., 2004), the Ion and Neutral Mass Spectrometer (INMS, Teolis et al., 2015; Waite et al., 2004)
- and the Cassini Fluxgate Magnetometer (MAG, Dougherty et al., 2004). The dataset spans the
- 59 entire Cassini mission and is representative of a full solar cycle and nearly half Titan year.

60 **2 Observations and Method**

61 2.1 Datasets and flyby coverage

- 62 We use data from 58 flybys where measurements of the Cassini INMS, RPWS/LP and the
- 63 Cassini MAG are simultaneously available (T#: A, B, 5, 16-21, 23, 25-30, 32, 34, 36, 38-40, 42-
- 64 44, 46, 48-51, 55-59, 61, 65, 71, 77, 83, 84, 86, 87, 91, 92, 95, 98, 100, 104, 107, 108, 113, 116-
- 65 121). The Cassini RPWS/LP dataset provides electron temperatures (Ågren et al., 2009) and
- 66 electron densities, positive ion densities, negative ion densities, derived as described in Shebanits
- 67 et al. (2016). The primary data from the INMS are the neutral densities (CH_4 and N_2). The INMS
- 68 ion data is used as supplementary for the ion mass information in the RPWS/LP analysis
- 69 (Shebanits et al., 2016).
- 70
- 71 Magnetic field data from Cassini MAG: magnetic field strength.
- 72
- 73 2.2 Method

74 Conductivities

- 75 Using the conductivity tensor representation (Schunk & Nagy, 2009) with dust term added
- 76 (Shebanits et al., 2020; Simon et al., 2011; Yaroshenko & Lühr, 2016), the Pedersen (σ_P), Hall
- 77 (σ_H) and magneticl field parallel (σ_{\parallel}) conductivities are defined as:

$$\sigma_{P} = \sum_{s=e,i,d} n_{s} \frac{q_{s}}{B} \frac{\nu_{s} \Omega_{s}}{\nu_{s}^{2} + \Omega_{s}^{2}}$$

$$\sigma_{H} = -\sum_{s=e,i,d} n_{s} \frac{q_{s}}{B} \frac{\Omega_{s}^{2}}{\nu_{s}^{2} + \Omega_{s}^{2}}$$

$$\sigma_{\parallel} = \sum_{s=e,i,d} \frac{n_{s} q_{s}^{2}}{m_{s} \nu_{s}}$$
(1)

78

- 79 Here the subscripts (s) denotes electrons (e), positive ions (i) and negative ions/dust grains (d)
- 80 and the input plasma parameters are number density (n), mass (m), charge (q), momentum
- 81 transfer collision frequency (v), gyrofrequency (Ω) and magnetic field strength (B).
- 82 Strictly speaking, the equations for conductivities must be summed over all the plasma species.
- 83 However, mass spectra information is of very limited availability for Titan's ionosphere. We 84 therefore adopt the average masses for the positive ions and for the negative ions/dust grains
- therefore adopt the average masses for the positive ions and for the negative ions/dust grains as described in Shebanits et al. (2016). Additionally, singly-charged negative ions/dust grains are
- assumed. This approach is validated using a test case of T40 flyby where the mass spectra for the
- positive ions and the negative ions and dust grains are available simultaneously (Coates et al.,
- 2009; Shebanits et al., 2016), see Appendix A.
- 89

90 Momentum Transfer Collision frequencies

- 91 In the following equations, the number density is always in cm⁻³, the rest of the units are SI. It
- 92 should be noted that since the RPWS/LP measures the total charged particle flux at a given bias
- voltage and the availability of the mass spectra is limited, we use the total charge densities for

94 the positive ions and negative ions/dust grains together with their respective mean mass as

95 derived and validated in Shebanits et al. (2016). The collision frequencies (and in turn,

- 96 conductivities) therefore represent mass-averages as well.
- 97 The momentum transfer collision frequencies (MTCFs) are largely dominated by collisions with
- 98 neutrals. However, in a dusty plasma a significant contribution from collisions with the negative
- 99 ions/dust grains is present and cannot be discarded (Shebanits et al., 2020; Yaroshenko & Lühr,
- 100 2016). We therefore use the total MTCF for each species as a sum of collision frequencies with
- 101 the other species:

102
$$v_{s,tot} = \sum_{t \neq s} v_{st}$$

- 103
- 104 The main neutral species in Titan's atmosphere are N_2 and CH_4 (Niemann et al., 2005; Waite et
- 105 al., 2005). Collisions of electrons with N_2 and CH_4 are implemented as hard-sphere collisions
- 106 (e.g., Schunk & Nagy, 2009) with the appropriate electron-dependent cross-sections $\chi(T_e)$ for N_2 107 (Itikawa 2006) and CH (Song et al. 2015):
- 107 (Itikawa, 2006) and CH_4 (Song et al., 2015):
- 108 $\nu_{e-neutral} = n_{neutral} \frac{8}{3} \sqrt{\frac{2k_b T_e}{\pi m_e}} \chi_{neutral}(T_e)$ (2)
- 109
- 110 For the negative ion/dust collisions with neutrals, both elastic (Coulomb) and hard-sphere
- 111 collisions are used (see also Shebanits et al., 2020). For the hard-sphere collisions, the cross-
- sections are replaced by the average dust grain cross-section πR_d^2 , where the average dust grain

113 radius R_d is estimated from the average grain mass by assuming a spherical grain and a mass

114 density of kerogen ($1280 \pm 300 \text{ kgm}^{-3}$, Stankiewicz et al., 2015):

115
$$v_{d-neutral} = \frac{8}{3\sqrt{\pi}} \frac{n_n m_n}{m_d + m_n} \sqrt{2k_b \frac{m_n T_d + m_d T_n}{m_n m_d} \pi R_d^2}$$
(3)

116 It should be noted that the actual dust grain geometry in Titan's ionosphere is likely to be of

117 fractal nature (e.g. Chatain et al., 2020; Michael et al., 2011; Shebanits et al., 2016; Sittler et al.,

118 2009; Waite et al., 2009). However, extensive lab experiments and sophisticated modelling are

119 required to produce a database of dust grain geometries (and their cross-sections) at Titan. We

120 therefore adapt spherical grains as a first order approximation.

121

Elastic collisions of electrons and neutrals with the heavy charged species are calculated according to Schunk & Nagy (2009), their eq. 4.144 & eq. 4.88, respectively:

124
$$v_{e-i,d} = 54.5 \frac{n_{i,d} Z_{i,d}^2}{-100}$$

$$\nu_{e-i,d} = 54.5 \frac{\pi_{i,d} - \pi_{i,d}}{T_e^{1.5}} , \qquad (4)$$

125
$$v_{i,d-neutral} = 2.5879 \times 10^{-9} n_n \sqrt{\frac{Y_n}{m_{i,d}} \frac{m_n}{m_n + m_{i,d}}},$$
 (5)

126 where the charge number is defined as $Z_{i,d} = q_{i,d}/|q_e|$.

127 MTCF equation for elastic collisions between the positive ions and negative ions/dust grains

128 (ion-dust drag) is the same as ion-ion collision (eq. 4.142 in Schunk & Nagy, 2009):

129

130
$$\nu_{id} = 1.27 Z_d^2 Z_i^2 \frac{n_d}{m_i} (m_i + m_d) \sqrt{\frac{m_d m_i}{(m_i T_d + m_d T_i)^3}}$$
(6)

131 **3 Results and Discussion**

132 3.1 Dynamo region

133 The MTCFs and gyrofrequencies defining the conductive dynamo region of Titan's ionosphere

are shown in Figure 1, plotted in altitude with the solar zenith angle (SZA) colorcoded. The error

135 bars are symmetric and represent the combined measurement uncertainty (2σ level). The

136 statistical means of the dynamo region boundaries from all included flybys are given in solid

137 black lines, with standard deviations as dashed lines.

138 The top boundary of the positive ions is at 1390 ± 90 km, in agreement with previous estimates 139 (Rosenquist et al., 2009). The negative ions and dust grains begin to conduct at ~100 km higher

140 altitudes due to their higher mass and subsequently lower gyrofrequency (see also Shebanits et

141 al., 2020). However, since the negative ions and dust grains are nearly absent above 1200 km

- 142 altitude in Titan's ionosphere, their impact on the top boundary of the dynamo region is
- 143 negligible.



144 145

Figure 1. Momentum transfer collision frequencies (color-coded by SZA) and gyrofrequencies (orange) for electrons (panel a),
positive ions (panel b) and negative ions/dust grains (panel c). The dynamo region boundary statistics from all included flybys:
the mean values are shown as solid black lines and the corresponding standard deviations as dashed lines. For the lower
boundary, the electron frequencies are extrapolated as defined in the text.

149 It should be noted that the lower boundary defined by electrons (panel a) is a rough estimate

derived from an exponential extrapolation of the electron MTCFs and a linear extrapolation of

151 their gyrofrequencies. Given the absence of magnetic field measurements below the closest

approach, a more sophisticated extrapolation is not deemed necessary for this work. In light of

this, the lower boundary is constrained to be between 700 and 1000 km altitude, with lower

154 altitudes more likely in the nightside ionosphere as indicated by the SZA trend (blue end of the

155 colorbar = nightside). Incidentally, the dusty plasma in Titan's ionosphere is expected to peak 156 below 1000 km (Shahanita et al. 2010). More sen this hal

- below 1000 km (Shebanits et al., 2016). More on this below.
- 157

158 The rather large variability in the top dynamo region boundary (Figure 1) is a combination of the

- variabilities in the background neutral densities, magnetic field, SZA and arguably also the local time. This is illustrated in Figure 2, where the top boundaries for positive ions are plotted in the
- 161 leftside panels a1 & a2 and for negative ions and dust grains in the rightside panels b1 & b2
- 161 (throughout the text, the panels in figures are labeled with letters for columns and numbers for
- (throughout the text, the panels in figures are labeled with letters for columns and numbers for
- 163 rows, for easier referencing). The magnetic field strength |B| is proportional to marker size and
- 164 the local time is colorcoded. Solid black lines are moving medians (50-points window).



Figure 2. Dynamo region top boundaries statistics for positive ions (left side panels a1 & a2) and for negative ions and dust
 grains (right side panels b1 & b2), plotted in altitude (top panels a1 & b1) and in the measured N₂ densites (bottom panels a2 & b2) versus SZA. Size of markers is proportional to the magnetic field strength |B|. Titan Local Time is color-coded.

169 The most obvious factor is |B|, with stronger field (larger markers) corresponding to lower

altitudes (panels a1, b1) and higher neutral densities (panels a2, b2). The SZA dependency is

171 noticeable for the negative ions and dust grains but is much weaker for the positive ions,

172 propagated from their respective masses. Although for the negative ions the median is roughly

173 linearly decreasing in altitude from dayside to nightside (panel b1), when plotted in the

background neutral densities instead (panel b2) the decrease is only clear on the nightside, likely

175 due to cooling and contraction of the neutral atmosphere. There seems to be a slight difference

between dawn (blue) and dusk (orange) regions for the negative ions and dust grains (right side,

177 panels b1 & b2), but with the current dataset it is statistically not conclusive.

178 3.2 Conductivities

- 179 Titan's ionospheric conductivities are shown in Figure 3 versus altitude (left side, panels a1-3)
- 180 and measured N_2 densities (right side, panels b1-3), color-coded with SZA. The error bars are
- 181 symmetric and represent the combined uncertainties of measurements, collision cross-sections
- and masses of heavy plasma species (2σ level). The error bars are one-sided for points where the
- 183 uncertainties reach zero (omitted on log scale to reduce clutter). The square markers in panels a2
- and b2 mark negative values of σ_H . The empty markers in all panels show the data points with low signal to noise ratio (positive ion densities < 10 cm⁻³). The field parallel conductivities σ_{\parallel}
- 186 separating into two profiles above ~1400 km altitude clearly demonstrate this, with the left
- profile (lower values) essentially showing the measurement limit of σ_{\parallel} for that region.
- 188 The SZA trends for the ionospheric conductivities are inherited from the ionospheric plasma
- number densities, with σ_P and σ_H being roughly linearly proportional to the positive ion densities
- 190 and σ_{\parallel} to the electron densities.
- 191 Pedersen conductivities σ_P have a peak between 1200 and 1400 km altitude and increase
- 192 sharply below 1100 km. At altitudes of ~1400 km, $\sigma_H > \sigma_P$ and the Hall currents are the
- 193 dominant horizontal currents (Ågren et al., 2011). However, at the CA the σ_H drops due to the
- 194 increasing presence of the dusty plasma. In a dusty plasma, σ_H is expected to reverse (reverse
- 195 Hall effect) due to the electron depletion onto the dusty grains (Muralikrishna & Kulkarni, 2006;
- 196 Shebanits et al., 2020; Simon et al., 2011; Yaroshenko & Lühr, 2016). The small cluster of data
- 197 points marked with a circle in panels a2 & b2 is indicative of this reversal (square markers are
- 198 $\sigma_H < 0$). With σ_H quenched or reversed and σ_P increasing below the CA, it is therefore possible
- 199 that a secondary layer of Pedersen current exists when the bottom boundary of the dynamo
- region extends enough to include the dusty plasma region below 900 km altitude.
- 201 Comparing the altitude profiles (panels a1-3) to the profiles versus measured neutral densities
- 202 (panels b1-3), the latter are more structured. Evidently, some of the variability below 1100 km 203 altitudes can be attributed to the variability of the neutral atmosphere. In particular, the increase
- of the Pedersen conductivity below 1100 km altitude is clearer, with the nightside values at 1000
- 205 km altitude corresponding to 5×10^9 cm⁻³, again consistent with Titan's atmosphere contracting
- 206 on the nightside.



Figure 3. Titan's ionospheric conductivities: Pedersen (panels a1, b1), Hall (a2, b2) and Parallel (a3, b3), plotted in altitude (left-side panels a1-3) and the measured N₂ densities (right-side panels b1-3). The SZA is color-coded. Red circle marks negative Hall conudctivities (reverse Hall effect).

- 211 3.3 Dusty plasma influence
- 212 The sharp increase in σ_P near closest approach (CA) was also evident in the earlier study and
- attributed to the decrease of the magnetic field strength (Rosenqvist et al., 2009), pre-dating the

- discovery of the negative ions/dust grains in the RPWS data (Ågren et al., 2012; Shebanits et al.,
- 215 2013).
- 216 Interestingly, although the increase of σ_P does coincide with an increase in positive ion and
- 217 negative ion/dust densities, the main influence is indeed the decrease in the magnetic field. This
- 218 is evident in Figure 4 which shows σ_P (left panels a1 & a2) and σ_H (right panels b1 & b2) below
- 219 1200 km altitude plotted versus magnetic field strength |B|. On the nightside the impact of |B| is
- 220 much weaker for both σ_P and σ_H . It is worth mentioning that the trend in |B| persists regardless
- 221 of the nominal corotational plasma ram direction.
- As mentioned above, the presence of dusty plasma strongly diminishes or even reverses σ_H . In
- 223 Titan's ionosphere, a low electron depletion ratio n_e/n_i (colorcoded) has been demonstrated to
- be a sufficient indicator for a dusty plasma (Shebanits et al., 2016). In Figure 4, the electron
- depletion ratio is colorcoded (dark = dusty) and the n_e/n_i trend in σ_H is clearly visible for the
- nightside ionosphere (panel b2) and to some extent also on the dayside (panel b1). This is in
- agreement with a globally present dusty plasma in Titan's ionosphere (Shebanits et al., 2016) and
- introduces a layer of complexity to the ionospheric current system that is unique to Titan so far.



Figure 4. Pedersen (left panels **a1** & **a2**) and Hall (right panels **b1** & **b2**) conductivities **below 1200 km** plotted versus magnetic field strength |B|. The electron depletion ratio n_e/n_i (color-coded) indicates the regions of dusty plasma. Top panels **a1** & **b1** show dayside ionosphere (SZA < 90°), bottom panels **a2** & **b2** show nightside (SZA > 90°).

233 3.4 General picture

234 Median profiles of the conductivities from Figure 3 are shown in Figure 5 versus the measured

- 235 N_2 densities (panels a1-3) and versus altitude (panels b1-3), colorcoded into dayside (SZA < 70,
- orange), terminator ($70 \le SZA < 100$, green) and nightside ($SZA \ge 100$, blue) regions.
- 237 Notably, σ_P peaks plotted in altitude (panel b1) appear broader on the dayside and terminator
- than on the nightside. This is an effect of the neutral atmosphere shifting in altitude between day and night, as evident when plotting versus the measured neutral densities instead (panel a1). As

- 240 mentioned above, the differences in the dayside, terminator and nightside profiles follow the
- 241 respective positive ion density profiles (electrons for field-parallel conductivity), with the upper
- 242 peak altitude increasing towards the terminator and roughly stabilizing towards the nightside
- 243 (Ågren et al., 2009; Shebanits et al., 2017).
- 244 The total median profiles are shown in black. Panels c1-3 show the total median profiles without
- charged dust (Rosenqvist et al., 2009) in red (marked R2009), to be compared with the medians
- from current work: solid black line for all flybys and dashed red line for the 17 flybys included in
- R2009. Comparing the total median profile to the R2009, it is evident that including the dusty
- 248 plasma shifts the first σ_P peak down in altitude by ~100 km. Expanding the dataset with the later
- 249 flybys increases this peak by \sim 20%. This increase is consistent with the enhancement of plasma
- densities during the solar maximum between 2011 and 2015 (Edberg et al., 2013; Shebanits et
- al., 2017). The profiles in panels a1-3 & b1-3 are therefore medians over a full solar cycle and
- 252 nearly half a Titan year.
- 253 The deviations between the R2009 profile and this work are attributed to two major differences.
- 254 Firstly, in the current work we treat the full plasma content including the charged dust grains –
- 255 whereas Rosenqvist et al. (2009) predated the dust detection in the RPWS/LP data. Secondly, we
- use the neutral densities measured in-situ by the INMS, while Rosenqvist et al. (2009) relied on
- an atmospheric model by Müller-Wodarg et al. (2008) this is why the σ_{\parallel} medians differ so
- much above 1200 km in panel c3. However, despite this, the total median profiles for the 17
- 259 flybys from R2009 are in a remarkable agreement down to ~1000 km with deviations only
- appearing near the CA. This again demonstrates that in Titan's ionosphere at altitudes above
- 261 ~1000 the σ_H and σ_P are mainly governed by the positive ions.



262 263 Figure 5. General conductivity profiles for all included flybys: Pedersen (panels a1, b1 & c1), Hall (panels a2, b2 & c2) and 264 Parallel (panels a3, b3 & c3). Panels a1-3 show sliding quadratic regressions with 400 point window (over N_2 densities). The 265 rest of the panels show sliding medians with a 200-point window (over altitudes). The total medians are in solid black. The 266 267 panels c1-3 also show the medians of the 17 flybys included in the Rosenqvist et al. (2009), with the R2009 profile as a solid red line and the same flybys from current work as a dashed red line. The colored lines are medians for the dayside (SZA < 70, 268 yellow), terminator ($70 \le SZA < 100$, green) and nightside ($SZA \ge 100$, blue).

269 Comparing Titan's dusty ionosphere with that of Saturn, the impact of charged dust on the 270 conductivities is very different. On Saturn, the Pedersen conductivities were increased by up to

- two orders of magnitude and the Hall conductivities reduced or reversed already above the upper 271
- 272 peak of ionospheric plasma densities (Shebanits et al., 2020). On Titan, although a 20% increase
- 273 in corresponding upper peak of Pedersen conductivities is evident, the real impact becomes
- 274 prominent near the CA. This is because the sources of the dust grains are different. On Saturn,
- the dust grains fall in from the D-ring (Hamil et al., 2018; e.g., Hsu et al., 2018; Mitchell et al., 275 276 2018) in addition to possibly forming in the top ionosphere (Waite et al., 2018), and are already
- present in significant quantities in the top ionosphere (Morooka et al., 2019), well above the 277
- 278 main photoionization peak (Hadid et al., 2018). On Titan, the dust grains are chemically formed
- in the ionosphere itself (Desai et al., 2017; Lavvas et al., 2013; Vuitton et al., 2009; Waite et al., 279

- 280 2007) and reach densities comparable to electrons and positive ions well below the
- 281 photoionization peak.
- 282

283 4 Conclusions

- The Pedersen and Hall conductivities in Titan's ionosphere down to ~1000 km altitudes
 are governed mainly by the positive ions.
- Dusty plasma influence on the Pedersen conductivity is measurable at ~1300 km
 altitude but most impact is below ~1000 km altitude (also for the Hall conductivity).
- The magnetic field strength remains the main factor for the increase in Pedersen conductivity at ~900-1100 km altitudes.
- The bulk of Titan's ionosphere in the 1000-1400 km altitude range is dominated by the
 Hall conductivity. Subsequently, Hall currents are expected to be the dominant
 horizontal currents.
- The sharp increase of the Pedersen conductivity below 1100 km altitude coincides with
 the dusty region of Titan's ionosphere, suggesting a dusty plasma layer dominated by
 the Pedersen conductivity (and subsequently, Pedersen currents).
- The Hall conductivity shows indications of a reverse Hall effect at ~950 km altitude.

297 Acknowledgments

- 298 O.S. acknowledges funding by SNSA Dnr 195/20. J.-E.W. acknowledges the support of the
- 299 Cassini RPWS/LP operations by SNSA. J.H.W. worked under internal funding from the Division
- 300 of Space Science and Engineering, Southwest Research Institute. M.K.D. acknowledges funding
- 301 by Royal Society Research Professorship RP140004.
- **Datasets availability:** the dataset used in the current study is available in a Mendeley Data
- 303 repository, doi: 10.17632/hwc5wfbvj9.1.

304 Appendix A

- 305 A validation of using the total densities and average masses of the heavy plasma species is
- 306 performed for the case of T40 (Figure 6a), where the mass spectra are available simultaneously
- 307 for the positive ions (Shebanits et al., 2016) and negative ions/dust grains (Coates et al., 2009).
- 308 The solid lines represent the average Pedersen (blue), Hall (red) and magnetic field parallel
- 309 (yellow) conductivity profiles, all in a very good agreement with the respective conductivities
- 310 derived using the full mass spectra (square markers). The corresponding plasma charge densities
- 311 are shown for reference (panel b).



Figure 6. Panel **a**: Comparison of the T40 conductivities derived using the average plasma species masses (solid lines with circle markers) and using the full mass spectra (squares and crosses) of the positive ions and negative ions/dust. Squares and solid lines are derived assuming singly charged particles, fainter crosses and dashed lines are derived using the empirical negative ion and dust grain charge estimate (Shebanits et al., 2016). Panel **b**: the plasma densities are shown for reference.

317 One of the parameters associated with the size and shape of the grains is the charge. Empirical 318 estimate of the average dust grain charge suggests values between -1.5 and -2.5 for flybys T16,

T29, T40 and T56 (Shebanits et al., 2016), while theoretical predictions (for spherical grains)

320 suggest most likely charge to be $Z_d = -1$ (Draine & Sutin, 1987; Meyer-Vernet, 2013). The

321 impact of the grain charge can be seen in Figure 6a, where the singly charged grains are used for

322 the solid lines and the empirically estimated Z_d is used for the dashed lines and. The difference

between these cases is <5% and is approximately a third of the combined measurement

uncertainties. For this reason, the conductivities for all flybys are derived assuming $Z_d = -1$. 325

326 **References**

- Ågren, K., Wahlund, J.-E., Garnier, P., Modolo, R., Cui, J., Galand, M., & Müller-Wodarg, I.
 (2009). On the ionospheric structure of Titan. *Planetary and Space Science*, 57(14–15),
 1821 1827 https://doi.org/10.1016/j.mcg.2000.04.012
- 329 1821–1827. https://doi.org/10.1016/j.pss.2009.04.012

Ågren, K., Andrews, D. J., Buchert, S. C., Coates, A. J., Cowley, S. W. H., Dougherty, M. K., et
 al. (2011). Detection of currents and associated electric fields in Titan's ionosphere from

332 Cassini data. *Journal of Geophysical Research: Space Physics*, *116*(4), A04313.

- 333 https://doi.org/10.1029/2010JA016100
- Ågren, K., Edberg, N. J. T., & Wahlund, J. E. (2012). Detection of negative ions in the deep
 ionosphere of Titan during the Cassini T70 flyby. *Geophysical Research Letters*, 39(10),
 n/a-n/a. https://doi.org/10.1029/2012GL051714
- Bertucci, C., Achilleos, N., Dougherty, M. K., Modolo, R., Coates, A. J., Szego, K., et al. (2008).
 The magnetic memory of Titan's ionized atmosphere. *Science*, *321*(5895), 1475–1478.
 https://doi.org/10.1126/science.1159780
- Bertucci, C., Duru, F., Edberg, N., Fraenz, M., Martinecz, C., Szego, K., & Vaisberg, O. (2011).
 The Induced Magnetospheres of Mars, Venus, and Titan. *Space Science Reviews*, *162*(1–4),
 113–171. https://doi.org/10.1007/s11214-011-9845-1
- Chatain, A., Carrasco, N., Ruscassier, N., Gautier, T., Vettier, L., & Guaitella, O. (2020).
 Interaction dust plasma in Titan's ionosphere: An experimental simulation of aerosols
 erosion. *Icarus*, 345(March), 113741. https://doi.org/10.1016/j.icarus.2020.113741
- Coates, A. J., Crary, F. J., Lewis, G. R., Young, D. T., Waite, J. H., Sittler, E. C., & Sittler, J. C.
 (2007). Discovery of heavy negative ions in Titan's ionosphere. *Geophysical Research Letters*, *34*(22), L22103. https://doi.org/10.1029/2007GL030978
- Coates, A. J., Wellbrock, A., Lewis, G. R., Jones, G. H., Young, D. T., Crary, F. J., & Waite, J.
 H. (2009). Heavy negative ions in Titan's ionosphere: Altitude and latitude dependence. *Planetary and Space Science*, 57(14–15), 1866–1871. https://doi.org/DOI
 10.1016/j.pss.2009.05.009
- Desai, R. T., Coates, A. J., Wellbrock, A., Vuitton, V., Crary, F. J., González-Caniulef, D., et al.
 (2017). Carbon Chain Anions and the Growth of Complex Organic Molecules in Titan's
 Ionosphere. *The Astrophysical Journal*, 844(2), L18. https://doi.org/10.3847/20418213/aa7851
- Dougherty, M. K., Kellock, S., Southwood, D. J., Balogh, A., Smith, E. J., Tsurutani, B. T., et al.
 (2004). The Cassini Magnetic Field Investigation. *Space Science Reviews*, *114*(1–4), 331–
 383. https://doi.org/10.1007/s11214-004-1432-2
- Draine, B. T., & Sutin, B. (1987). Collisional charging of interstellar grains. *The Astrophysical Journal*, 320, 803. https://doi.org/10.1086/165596
- Edberg, N. J. T., Andrews, D. J., Shebanits, O., Ågren, K., Wahlund, J. E., Opgenoorth, H. J., et
 al. (2013). Solar cycle modulation of Titan's ionosphere. *Journal of Geophysical Research: Space Physics*, *118*(8), 5255–5264. https://doi.org/10.1002/jgra.50463
- Garnier, P., Dandouras, I., Toublanc, D., Roelof, E. C., Brandt, P. C., Mitchell, D. G., et al.
 (2010). Statistical analysis of the energetic ion and ENA data for the Titan environment. *Planetary and Space Science*, 58(14–15), 1811–1822.
- 368 https://doi.org/10.1016/j.pss.2010.08.009

- Gurnett, D. A., Kurth, W. S., Kirchner, D. L., Hospodarsky, G. B., Averkamp, T. F., Zarka, P., et
 al. (2004). The Cassini radio and plasma wave investigation. *Space Science Reviews*,
- 371 *114*(1–4), 395–463. https://doi.org/DOI 10.1007/s11214-004-1434-0
- Hadid, L. Z., Morooka, M. W., Wahlund, J. -E., Persoon, A. M., Andrews, D. J., Shebanits, O.,
 et al. (2018). Saturn's Ionosphere: Electron Density Altitude Profiles and D-Ring
 Interaction From The Cassini Grand Finale. *Geophysical Research Letters*, 46(16), 9362–
- 375 9369. https://doi.org/10.1029/2018GL078004
- Hamil, O., Cravens, T. E., Reedy, N. L., & Sakai, S. (2018). Fate of Ice Grains in Saturn's
 Ionosphere. *Journal of Geophysical Research: Space Physics*, *123*(2), 1429–1440.
 https://doi.org/10.1002/2017JA024616
- Hsu, H.-W., Schmidt, J., Kempf, S., Postberg, F., Moragas-Klostermeyer, G., Seiß, M., et al.
 (2018). In situ collection of dust grains falling from Saturn's rings into its atmosphere. *Science*, 362(6410), eaat3185. https://doi.org/10.1126/science.aat3185
- Itikawa, Y. (2006). Cross Sections for Electron Collisions with Nitrogen Molecules. *Journal of Physical and Chemical Reference Data*, 35(1), 31–53. https://doi.org/10.1063/1.1937426
- Lavvas, P., Yelle, R. V., Koskinen, T., Bazin, A., Vuitton, V., Vigren, E., et al. (2013). Aerosol
 growth in Titan's ionosphere. *Proceedings of the National Academy of Sciences*, *110*(8),
 2729–2734. https://doi.org/10.1073/pnas.1217059110
- Meyer-Vernet, N. (2013). On the charge of nanograins in cold environments and Enceladus dust.
 Icarus, 226(1), 583–590. https://doi.org/10.1016/j.icarus.2013.06.014
- Michael, M., Tripathi, S. N., Arya, P., Coates, A., Wellbrock, A., & Young, D. T. (2011). Highaltitude charged aerosols in the atmosphere of Titan. *Planetary and Space Science*, 59(9),
 880–885. https://doi.org/10.1016/j.pss.2011.03.010
- Mitchell, D. G., Perry, M. E., Hamilton, D. C., Westlake, J. H., Kollmann, P., Smith, H. T., et al.
 (2018). Dust grains fall from Saturn's D-ring into its equatorial upper atmosphere. *Science*, *362*(6410), eaat2236. https://doi.org/10.1126/science.aat2236
- Morooka, M. W., Wahlund, J. E., Eriksson, A. I., Farrell, W. M., Gurnett, D. A., Kurth, W. S., et
 al. (2011). Dusty plasma in the vicinity of Enceladus. *Journal of Geophysical Research- Space Physics*, *116*(A12). https://doi.org/Doi 10.1029/2011ja017038
- Morooka, M. W., Wahlund, J.-E. J. -E. E., Hadid, L. Z., Eriksson, A. I., Edberg, N. J. T. T.,
 Vigren, E., et al. (2019). Saturn's Dusty Ionosphere. *Journal of Geophysical Research: Space Physics*, *124*(3), 1679–1697. https://doi.org/10.1029/2018JA026154
- 401 Müller-Wodarg, I. C. F., Yelle, R. V., Cui, J., & Waite, J. H. (2008). Horizontal structures and
 402 dynamics of Titan's thermosphere. *Journal of Geophysical Research*, *113*(E10), E10005.
 403 https://doi.org/10.1029/2007JE003033
- 404 Muralikrishna, P., & Kulkarni, V. H. (2006). On the height variation of the E-region cowling

405 conductivity & amp; amp; ndash; effect of charged dust particles. Annales Geophysicae, 406 24(11), 2949–2957. https://doi.org/10.5194/angeo-24-2949-2006 407 Niemann, H. B., Atreya, S. K., Bauer, S. J., Carignan, G. R., Demick, J. E., Frost, R. L., et al. 408 (2005). The abundances of constituents of Titan's atmosphere from the GCMS instrument 409 on the Huygens probe. Nature, 438(7069), 779–784. https://doi.org/10.1038/nature04122 410 Rosenqvist, L., Wahlund, J. E., Ågren, K., Modolo, R., Opgenoorth, H. J., Strobel, D., et al. 411 (2009). Titan ionospheric conductivities from Cassini measurements. Planetary and Space 412 Science, 57(14-15), 1828-1833. https://doi.org/10.1016/j.pss.2009.01.007 413 Schunk, R., & Nagy, A. (2009). Ionospheres (2nd ed.). Cambridge: Cambridge University Press. 414 https://doi.org/10.1017/CBO9780511635342 415 Shebanits, O., Wahlund, J. E., Mandt, K., Ågren, K., Edberg, N. J. T., & Waite, J. H. (2013). 416 Negative ion densities in the ionosphere of Titan-Cassini RPWS/LP results. Planetary and 417 Space Science, 84, 153–162. https://doi.org/10.1016/j.pss.2013.05.021 418 Shebanits, O., Wahlund, J.-E., Edberg, N. J. T., Crary, F. J., Wellbrock, A., Andrews, D. J., et al. 419 (2016). Ion and aerosol precursor densities in Titan's ionosphere: A multi-instrument case 420 study. Journal of Geophysical Research: Space Physics, 121(10), 10,075-10,090. 421 https://doi.org/10.1002/2016JA022980 422 Shebanits, O., Vigren, E., Wahlund, J. -E., Holmberg, M. K. G., Morooka, M., Edberg, N. J. T., 423 et al. (2017). Titan's ionosphere: A survey of solar EUV influences. Journal of Geophysical 424 Research: Space Physics, 122(7), 7491–7503. https://doi.org/10.1002/2017JA023987 425 Shebanits, O., Hadid, L. Z., Cao, H., Morooka, M. W., Hunt, G. J., Dougherty, M. K., et al. 426 (2020). Saturn's near-equatorial ionospheric conductivities from in situ measurements. 427 Scientific Reports 2020 10:1, 10(1), 1-13. https://doi.org/10.1038/s41598-020-64787-7 428 Simon, S., Saur, J., Kriegel, H., Neubauer, F. M., Motschmann, U., & Dougherty, M. K. (2011). 429 Influence of negatively charged plume grains and hemisphere coupling currents on the 430 structure of Enceladus' Alfvén wings: Analytical modeling of Cassini magnetometer 431 observations. Journal of Geophysical Research: Space Physics, 116(A4), n/a-n/a. 432 https://doi.org/10.1029/2010JA016338 433 Sittler, E. C., Ali, A., Cooper, J. F., Hartle, R. E., Johnson, R. E., Coates, A. J., & Young, D. T. 434 (2009). Heavy ion formation in Titan's ionosphere: Magnetospheric introduction of free 435 oxygen and a source of Titan's aerosols? Planetary and Space Science, 57(13), 1547–1557. 436 https://doi.org/10.1016/j.pss.2009.07.017 437 Song, M.-Y., Yoon, J.-S., Cho, H., Itikawa, Y., Karwasz, G. P., Kokoouline, V., et al. (2015). Cross Sections for Electron Collisions with Methane. Journal of Physical and Chemical 438 439 Reference Data, 44(2), 023101. https://doi.org/10.1063/1.4918630 440 Stankiewicz, A., Ionkina, N., Motherwell, B., Bennett, B., Wint, O., & Mastalerz, M. (2015). 441 Kerogen Density Revisited—Lessons From the Duvernay Shale. In Proceedings of the 3rd

442 443	Unconventional Resources Technology Conference. Tulsa, OK, USA: American Association of Petroleum Geologists. https://doi.org/10.15530/urtec-2015-2157904
444 445 446	Teolis, B. D., Niemann, H. B., Waite, J. H., Gell, D. A., Perryman, R. S., Kasprzak, W. T., et al. (2015). A Revised Sensitivity Model for Cassini INMS: Results at Titan. Space Science Reviews, 190(1–4), 47–84. https://doi.org/10.1007/s11214-014-0133-8
447 448 449	Vuitton, V., Lavvas, P., Yelle, R. V., Galand, M., Wellbrock, A., Lewis, G. R., et al. (2009). Negative ion chemistry in Titan's upper atmosphere. <i>Planetary and Space Science</i> , 57(13), 1558–1572. https://doi.org/10.1016/j.pss.2009.04.004
450 451 452	Waite, J. H., Lewis, W. S., Kasprzak, W. T., Anicich, V. G., Block, B. P., Cravens, T. E., et al. (2004). The Cassini Ion and Neutral Mass Spectrometer (INMS) investigation. <i>Space Science Reviews</i> , 114(1–4), 113–231. https://doi.org/10.1007/s11214-004-1408-2
453 454 455	Waite, J. H., Niemann, H., Yelle, R. V., Kasprzak, W. T., Cravens, T. E., Luhmann, J. G., et al. (2005). Ion Neutral Mass Spectrometer results from the first flyby of Titan. <i>Science</i> , 308(5724), 982–986. https://doi.org/10.1126/science.1110652
456 457 458	Waite, J. H., Young, D. T., Cravens, T. E., Coates, A. J., Crary, F. J., Magee, B., & Westlake, J. (2007). The Process of Tholin Formation in Titan's Upper Atmosphere. <i>Science</i> , 316(5826), 870–875. https://doi.org/10.1126/science.1139727
459 460 461	 Waite, J. H., Young, D. T., Westlake, J. H., Lunine, J. I., McKay, C. P., & Lewis, W. S. (2009). High-Altitude Production of Titan's Aerosols. In <i>Titan from Cassini-Huygens</i> (pp. 201–214). Dordrecht: Springer Netherlands. https://doi.org/10.1007/978-1-4020-9215-2_8
462 463 464	Waite, J. H., Perryman, R. S., Perry, M. E., Miller, K. E., Bell, J., Cravens, T. E., et al. (2018). Chemical interactions between Saturn's atmosphere and its rings. <i>Science</i> , 362(6410), eaat2382. https://doi.org/10.1126/science.aat2382
465 466	Yaroshenko, V. V., & Lühr, H. (2016). Electrical conductivity of the dusty plasma in the Enceladus plume. <i>Icarus</i> , 278, 79–87. https://doi.org/10.1016/j.icarus.2016.05.033
467	
468 469 470 471 472	References in supporting information must also be included in the reference list of the main paper, or in a designated section in the main paper, so that they will be discovered, linked, and indexed. A separate list in the supporting information is not necessary. References are not included in word counts for excess length fees. Data sets that are not newly reported as part of this research should also be cited in the references. New data sets or software that are deposited

- 472 this research should also be cited in the references. New data sets or software that are deposited
- 473 elsewhere with a permanent identifier should be cited. AGU follows the Joint Declaration of
 474 Data Citations Principles.
- 475 All references must be available to readers at the time of publication; there should be no
- 476 "unpublished", in preparation, under review, or "in press" references. Please write the respective
- 477 journal with any questions.
- 478 An example of a reference:

- 479 Deng, A., & Stauffer, D. R. (2006), On improving 4-km mesoscale model simulations. *Journal*
- 480 of Applied Meteorology and Climatology, 45(3), 361–381. doi:10.1175/JAM2341.1
- 481 More information on reference formatting with examples can be found in our <u>Brief Guide to</u>
- 482 <u>AGU Style</u>.
- 483

Figure 1. The figure caption should begin with an overall descriptive statement of the figure 485 followed by additional text. They should be immediately after each figure. Figure parts are 486 indicated with lower-case letters (**a**, **b**, **c**...). For initial submission, please place both the figures 487 and captions in the text near where they are cited rather than at the end of the file (not both). At 488 revision, captions can be placed in-text or at the end of the file, and figures should be uploaded 489 separately. Each figure should be one complete, cohesive file (please do not upload sub-figures 490 or figure parts in separate files).

- 491 **Table 1.** Start this caption with a short description of your table. Format tables using the Word
- 492 Table commands and structures. Additional information on table formatting can be found in our
- 493 Style Guide, <u>Table Formatting</u>. Do not create tables using spaces or tabs characters. Large
- 494 tables presenting rich data should be presented as separate excel or .csv files, not as part of the
- 495 main text.
- 496