

An Analysis of Magnetosphere-Ionosphere Coupling That Is Independent of Inertial Reference Frame

Anthony James Mannucci^{1,1}, Ryan McGranaghan^{2,2}, Xing Meng^{1,1}, and Olga P. Verkhoglyadova^{1,1}

¹Jet Propulsion Laboratory, California Institute of Technology

²Jet Propulsion Laboratory

November 30, 2022

Abstract

This paper analyses magnetosphere-ionosphere (MI) coupling from a perspective that is independent of inertial reference frame, explicitly acknowledging the role of the principle of relativity in MI coupling. For the first time in the context of MI coupling, we discuss the literature on the low-velocity limit of the theory of special relativity applied to electrodynamics. In many MI coupling theories, a particular low-velocity limit applies, known as the “magnetic limit”. Two important consequences of the magnetic limit are: 1) Maxwell’s equations cannot contain a displacement current and be consistent with the magnetic limit and 2) the magnetic field is not modified by currents created by charge densities in motion, thus charge density is approximately zero. We show how reference frame-independent descriptions of MI coupling require that ion-neutral relative velocities and ion-neutral collisions are key drivers of the physics. Electric fields, on the other hand, depend on reference frame, and can be zero in an appropriate frame. Currents are independent of reference frame and will flow when the electric field is close to zero. Starting with the same momentum equations that are typically used to derive Ohm’s law, we derive an equation that relates the perpendicular current to collisions between ions and neutrals, and electrons and neutrals, without reference to electric fields. Ignoring the relative motion between ions and neutrals will result in errors exceeding 100% for estimates of high latitude Joule heating during significant geomagnetic storms when ion-neutral velocity differences are largest near the initiation of large-scale ion convection.

An Analysis of Magnetosphere-Ionosphere Coupling That Is Independent of Inertial Reference Frame

Anthony J. Mannucci¹, Ryan McGranaghan², Xing Meng¹ and Olga P. Verkhoglyadova¹

¹Jet Propulsion Laboratory, California Institute of Technology, Pasadena, CA, USA.

²Atmospheric and Space Technology Research Associates (ASTRA), Boulder CO, USA.

Corresponding author: Anthony Mannucci (anthony.j.mannucci@jpl.nasa.gov)

Key Points:

- Relativistic transformations applied to electrodynamics are analyzed in the context of MI-coupling.
- An alternative equation to “Ohm’s law” for horizontal ionospheric currents does not require the existence of a large-scale electric field.
- Electrodynamic theories of MI coupling that do not account for the relative motion of ions and neutrals are not quantitatively accurate.

Abstract

This paper analyses magnetosphere-ionosphere (MI) coupling from a perspective that is independent of inertial reference frame, explicitly acknowledging the role of the principle of relativity in MI coupling. For the first time in the context of MI coupling, we discuss the literature on the low-velocity limit of the theory of special relativity applied to electrodynamics. In many MI coupling theories, a particular low-velocity limit applies, known as the “magnetic limit”. Two important consequences of the magnetic limit are: 1) Maxwell’s equations cannot contain a displacement current and be consistent with the magnetic limit and 2) the magnetic field is not modified by currents created by charge densities in motion, thus charge density is approximately zero. We show how reference frame-independent descriptions of MI coupling require that ion-neutral *relative velocities* and ion-neutral collisions are key drivers of the physics. Electric fields, on the other hand, depend on reference frame, and can be zero in an appropriate frame. Currents are independent of reference frame and will flow when the electric field is close to zero. Starting with the same momentum equations that are typically used to derive Ohm’s law, we derive an equation that relates the perpendicular current to collisions between ions and neutrals, and electrons and neutrals, without reference to electric fields. Ignoring the relative motion between ions and neutrals will result in errors exceeding 100% for estimates of high latitude Joule heating during significant geomagnetic storms when ion-neutral velocity differences are largest near the initiation of large-scale ion convection.

Plain Language Summary

Interactions between the magnetized and ionized solar wind, the magnetospheric cavity surrounding Earth, and Earth’s ionized upper atmosphere (ionosphere) can create rapid (~ 1 km/sec) large-scale motions of the ionosphere during periods known as geomagnetic storms. We use the principle of relativity to gain insight into the complex physics of these interactions. Relativity states that the physics governing geospace must be independent of the velocity of an observer making measurements of the system. We write key equations governing interactions of the magnetosphere-ionosphere system in terms of quantities that do not depend on the observer’s motion. In doing so, we find that previous theories had over-emphasized the importance of a large-scale electric field that grows during storms. Instead, we show that the important physics is related to collisions between the ionized portion of the atmosphere and the un-ionized “neutral” component that contains much more mass. These collisional interactions create upper

atmospheric heating and expansion, and cause large-scale currents to flow between the ionosphere and magnetosphere, resulting in a multitude of impacts to our technological society. Using the principle of relativity, and isolating the physics that is independent of observer motion, led us to a deeper understanding of key physical processes during storms.

1 Introduction

Electrodynamics as it pertains to magnetosphere-ionosphere (MI) coupling is a critical aspect of the ionospheric response during geomagnetic storms. Large-scale convection of the ionospheric plasma during disturbed geomagnetic conditions at auroral latitudes and higher, and heating of the atmosphere due to collisions between charged and neutral species, originates with electric and magnetic forces acting on the plasma that change dramatically when solar wind conditions and the state of the MI system lead to geomagnetic storms. Scientific consensus on fundamental aspects of the physical processes that occur at high latitude is not yet achieved, including the definition of Joule heating (Vasyliunas and Song, 2005; Verkhoglyadova et al. 2017). In this paper, we provide insight into the physical processes occurring during geomagnetic storms by emphasizing an approach that is independent of the inertial reference frame used to describe the phenomena.

The principle of relativity (PR) states that the laws of physics are independent of inertial frame (Pal, 2013). The PR has important implications for high latitude electrodynamics due to the electric field in the high-latitude ionosphere being dependent on inertial reference frame. The large-scale electric field at high latitudes is directed predominantly in the horizontal direction and the magnetic field tends to be predominantly in the vertical direction. The PR requires that the electric field component perpendicular to the magnetic field vary according to inertial reference frame. There are several examples in the literature where electric fields are used to describe MI coupling physics. For example, the concept that horizontal currents originating in the magnetosphere and closing in the ionosphere results in electric fields that cause high velocity **$\mathbf{E} \times \mathbf{B}$** plasma flow has been invoked for phenomena such as sub-auroral ion drifts (Anderson et al., 1993). Another example is the relationship between the ionospheric electric field and field-aligned Region 1 currents closing in the ionosphere (Fedder and Lyon, 1987; Siscoe et al., 2002; Wiltberger et al., 2003; Rothwell and Jasperse, 2006). Since the electric field depends on reference frame, such examples would seem to obscure essential physics because the same phenomena can be viewed in a reference frame for which the electric field is of very different magnitude. The notion of a “preferred inertial reference frame” for high latitude electrodynamics has been discussed (Vasyliūnas and Song, 2005; Leake et al., 2014; Strangeway, 2012) and is useful when considering that high latitude phenomena occur within the physical media of plasma

and neutral gases. A reference frame where the electric field vanishes locally in the ionospheric E-region where collisions dominate is also useful to consider in understanding the physical basis of MI coupling.

In the remainder of this paper, we apply the PR in the context of high latitude electrodynamics. We first review the PR and how it is typically expressed in the low velocity limit. For the first time in the context of MI-coupling, we introduce the literature on the low-velocity limit of special relativity as it applies to electrodynamics and discuss several implications of this limit. We next discuss an equation for the high latitude horizontal current density as an alternative to Ohm's law that contains only terms that do not depend on reference frame. In doing so, we clarify that the high-latitude current is primarily the result of the relative bulk motion between ions and neutrals resulting from large-scale convection of the ions initiated during a geomagnetic storm by the solar wind. Horizontal ionospheric currents are the by-products of ion-neutral velocity differences, rather than drivers of an electric field. We discuss the implications of ignoring the role of neutral dynamics in magnetosphere-ionosphere coupling.

2 The Low Velocity Limit of the Principle of Relativity

In this section, we review the literature of how electric and magnetic fields transform in the low velocity limit of the principle of relativity. The invariance of physical laws in this limit is known as the “Principle of Galilean Relativity” (PGR). While the PGR might appear to be a well-settled topic, there are subtleties in its application to electrodynamic phenomena, of which MI coupling is a prime example. The challenges of applying Galilean relativity to electrodynamics was first discussed by Le Bellac and Lévy-Leblond (1973) (hereafter referred to as LL73). To our knowledge, references to this paper and the subsequent literature on Galilean electrodynamics have not appeared within the literature of MI coupling, so we devote some discussion to this topic.

Throughout the text, we use the symbol \mathbf{v}_r to indicate the relative velocity between two inertial reference frames, for example frames \mathcal{A} and \mathcal{B} . We use primed variables to refer to physical quantities measured in the inertial reference frame \mathcal{B} moving with velocity \mathbf{v}_r relative to reference frame \mathcal{A} .

The theory or principle of special relativity (PSR), first deduced by Einstein in 1905, requires that electric and magnetic fields transform between inertial reference frames according to the following relationships (Rousseaux, 2014; Heras, 2010):

$$\mathbf{E}' = \gamma \left(\mathbf{E} - \frac{\gamma - 1}{\gamma} \frac{\mathbf{v}_r (\mathbf{v}_r \cdot \mathbf{E})}{v_r^2} + \mathbf{v}_r \times \mathbf{B} \right) \quad (1)$$

$$\mathbf{B}' = \gamma \left(\mathbf{B} - \frac{\gamma - 1}{\gamma} \frac{\mathbf{v}_r (\mathbf{v}_r \cdot \mathbf{B})}{v_r^2} - \frac{1}{c^2} \mathbf{v}_r \times \mathbf{E} \right) \quad (2)$$

where $(\mathbf{E}', \mathbf{B}')$ are the electric and magnetic fields in the new inertial reference frame, (\mathbf{E}, \mathbf{B}) are the fields in the original frame, and $\gamma = 1/\sqrt{1 - v_r^2/c^2}$ with c being the speed of light.

It is clear that these equations do not possess a unique low-velocity limit because the field transformations do not depend on velocity exclusively, but also on the electric and magnetic fields themselves. The “simple” approach to the low velocity limit, by setting $\gamma \approx 1$ in Equations (1) and (2), yields transformation equations that do not obey the composition law (see Equation 2 in de Montigny and Rousseaux, 2006). Applying two such transformations in succession, first with a relative frame velocity \mathbf{v}_1 followed by a transformation with relative frame velocity \mathbf{v}_2 , yields a different transformation of fields than if a single transformation were applied with velocity $\mathbf{v}_r = \mathbf{v}_1 + \mathbf{v}_2$. Such a non-conforming transformation does not obey the principle of relativity, and should not be adopted, but there are consequences, as described below.

LL73 identified two low-velocity limiting cases of the relativistic transformation equations that do obey the composition law: the “electric” and “magnetic” limits of the PGR. LL73 spawned a literature on the topic of “Galilean electromagnetism”. The electric and magnetic limits of LL73 correspond to which field magnitude is dominant. The electric limit applies when $E \gg cB$, and the magnetic limit applies when $cB \gg E$. High latitude electrodynamics and MI coupling usually encompasses only one of these limits: the magnetic. The magnetic limit of the PGR is applicable

to MI coupling when inductive electric fields are neglected, such that electric field magnitudes are given approximately by $E \approx vB$ and we assume that $v \ll c$.

For these reasons, we focus here on the magnetic limit of the PGR ($cB \gg E$) and assume that terms of order v_r/c are small. The Lorentz transformation laws (Equations (1) and (2)) become (LL73):

$$\mathbf{E}' = \mathbf{E} + \mathbf{v}_r \times \mathbf{B} \quad (3)$$

$$\mathbf{B}' = \mathbf{B} \quad (4)$$

which are familiar transformation rules in the context of space physics (e.g. Strangeway and Raeder, 2001; Parks, 2007; Vasyliūnas and Song, 2005). We refer to these low-velocity limit equations as comprising a Galilean transformation, by analogy to the more general Lorentz transformation of special relativity.

An alternative derivation of the transformation laws in the magnetic limit is possible by considering the Lorentz force law (Preti et al., 2009; Heras, 2010). The Lorentz force \mathbf{F} is given by:

$$\mathbf{F} = q(\mathbf{E} + \mathbf{v} \times \mathbf{B}) \quad (5)$$

where \mathbf{F} is the force on a charge q moving with velocity \mathbf{v} in frame \mathcal{A} , and where the electric field is \mathbf{E} and the magnetic flux density is \mathbf{B} . This force is invariant to reference frame if transformation equations (3) and (4) are used. However, we note for completeness that the Lorentz force is not invariant in the electric limit of Galilean electromagnetism (see Heras, 2010).

Equation 3 shows that the component of the electric field parallel to \mathbf{B} is unchanged under a Galilean transformation, whereas the electric field component perpendicular to both \mathbf{v}_r and \mathbf{B} changes depending on the relative velocity of frame \mathcal{B} . Equation (4) states that \mathbf{B} is a Galilean

invariant (GI) and the same in both inertial frames, but of course the invariance of \mathbf{B} only holds if there is a very small net charge density (see below), a condition generally consistent with MI-coupling theories. The frame-variant nature of the electric field has significant implications within the context of high-latitude electrodynamics, as we discuss below. Table 1 summarizes how different physical quantities relevant to high latitude electrodynamics vary under a Galilean transformation in the magnetic limit.

The source terms of the fields, net charge density ρ and current density \mathbf{J} , also transform according to the PSR. As shown by Heras (2010), they transform as a four vector according to:

$$\mathbf{J}' = \mathbf{J} - \gamma \mathbf{v}_r \rho + (\gamma - 1) \frac{\mathbf{v}_r (\mathbf{v}_r \cdot \mathbf{J})}{v_r^2} \quad (6)$$

$$\rho' = \gamma \left(\rho - \frac{(\mathbf{v}_r \cdot \mathbf{J})}{c^2} \right) \quad (7)$$

In the magnetic limit, the transformation equations become (LL73):

$$\rho' = \rho - \frac{(\mathbf{v}_r \cdot \mathbf{J})}{c^2} \quad (8)$$

$$\mathbf{J}' = \mathbf{J} \quad (9)$$

For the magnetic limit to apply, the condition $c\rho \ll J$ must hold also. The Galilean invariance of current expressed by Equation (9) is familiar in the context of high-latitude ionospheric electrodynamics (e.g. Thayer and Semeter, 2004) and is intuitive when net charge density is zero or very small ($\rho \approx 0$). It is intuitively obvious that invariance of the magnetic field (Equation (4)) *requires* very small charge densities, because to first order in \mathbf{v}_r , a charge density becomes a current in the moving reference frame, and such a current will modify the magnetic field according to Ampère's law. We note that adding a term such as $\rho \mathbf{v}_r$ to the transformation Equation (9) will cause the source transformation rules to violate the composition law.

Therefore, MI coupling theories that implicitly rely on magnetic field invariance must assume negligible charge densities. We note also (Table 1) that zero charge density in the original reference frame leads to a small charge density in the moving frame according to Equation (8), but this is a second-order effect in v_r/c .

The literature of ionospheric electrodynamics refers to non-zero charge densities or “charge accumulation” (e.g. Figure 3 of Vasyliunas, 2012), leading to “polarization electric fields” in the context of the disturbance dynamo (Richmond and Thayer, 2000). To our knowledge, there is no discussion in the ionospheric literature of the inconsistency between finite charge densities and frame-invariant currents, as required by the Galilean transformation in the magnetic limit. This inconsistency does not cause a problem since the charge densities are small ($\rho \approx 0$) and quasi-neutrality applies. It is worth noting the relativistic considerations are very relevant to the discussion in Vasyliunas (2012) regarding the neutral wind dynamo. In the “conventional approach” (Figure 3 of Vasyliunas, 2012), an electric field created in the dynamo region maps along magnetic field lines to higher altitudes where $\mathbf{E} \times \mathbf{B}$ drift causes plasma to flow. In a frame where the electric field in the dynamo region is close to zero (e.g. see Figure 1 of this paper), such a mapped electric field would not account for plasma flow at higher altitudes, but plasma would still be flowing since the plasma flow velocity will not transform away at the higher altitudes. In the “complete” physical description of the neutral wind dynamo (Figure 4 of Vasyliunas (2012)), a non-zero curl of the electric field causes a magnetic perturbation that results in plasma flow. Neither the curl of the electric field nor the magnetic perturbation is affected by a Galilean transformation, so the non-conventional physical basis offered by Vasyliunas (2012) is consistent with relativistic considerations.

It is important to understand the magnitude of electric field changes that can occur when considering relative velocities that are typical at high latitudes where MI coupling occurs. In Figure 1 we depict an idealized situation where electric and magnetic fields are at perpendicular angles to each other, to represent approximately the high latitude ionosphere where the Earth’s magnetic field is close to vertical and large-scale convection electric fields are predominantly horizontal. We will refer to this geometry elsewhere in the text. For typical high latitude electric fields of magnitude ~ 25 mV/m as viewed in an Earth-fixed frame, and typical magnetic field

247 magnitudes of $\sim 50,000$ nT (near 110 km altitude), the electric field will be reduced significantly
248 in a reference frame moving with a speed of ~ 0.5 km/s relative to the Earth. The direction of
249 such a moving frame is shown in Figure 1. A speed of ~ 0.5 km/s relative to the Earth is
250 consistent with the low velocity limit of Galilean electrodynamics and consistent with velocities
251 of the plasma or neutral species relative to the Earth. An electric field that is non-negligible
252 observed from an Earth-fixed frame can be much smaller in magnitude viewed from an inertial
253 reference frame moving at speeds consistent with the low velocity limit of the PSR, and with the
254 Galilean transformation rules of Table 1. Although an Earth fixed frame is not inertial, the small
255 acceleration in this frame is typically ignored for the purposes of these estimates. We note that
256 electric field spatial gradients will be maintained independent of reference frame, which arise
257 due to the height variation of collision frequencies (see Section 3.2 where Ohm's law is
258 discussed). However, the magnitude of the field could be rendered quite small at a particular
259 altitude and in a particular region.

260

261




Figure 1. Schematic representation of electric and magnetic fields at northern high latitudes. Magnetic fields are directed vertically (B). Electric fields are directed horizontally (e.g. in an Earth fixed frame, resulting from magnetospheric convection). In a reference frame moving with velocity \mathbf{v}_r in the direction shown (into the page), the electric field is zero if the reference frame moves with speed $\|\mathbf{E}\|/\|\mathbf{B}\|$. For “typical” disturbed conditions, non-relativistic speeds of ~ 1.6 km/s are sufficient.

Table 1. Transformation of physical quantities under change of inertial reference frame in the Galilean magnetic limit, which applies when $cB \gg E$ and $c\rho \ll J$. The primed quantities are in the new reference frame moving at velocity \mathbf{v}_r relative to the original frame. All quantities are assumed to be quasi-static or slowly varying. Galilean invariance refers to quantities that are the same in all inertial reference frames.

Physical Quantity	Transformation	Comment
Electric field \mathbf{E}	$\mathbf{E}' = \mathbf{E} + \mathbf{v}_r \times \mathbf{B}$	The electric field in a direction parallel to \mathbf{B} is invariant, as is the electric field in the absence of a magnetic field
Magnetic field \mathbf{B}	$\mathbf{B}' = \mathbf{B}$	
Current density \mathbf{J}	$\mathbf{J}' = \mathbf{J}$	
Charge density ρ	$\rho' = \rho - \frac{1}{c^2} \mathbf{v}_r \cdot \mathbf{J}$	Charge per unit volume.
Heat energy Q	$Q' = Q$	Heat energy and temperature are Galilean invariant (GI)
Velocity \mathbf{v}	$\mathbf{v}' = \mathbf{v} - \mathbf{v}_r$	
Relative velocity between two species	Invariant	
Collisional forces	Invariant	Depends on relative velocities, which are invariants

There are several implications of the electric and magnetic field transformation rules in Table 1. First, Maxwell's equations are not invariant under these transformations, unless the displacement current term is neglected in Ampere's law (LL73; Preti et al., 2009; Heras, 2010). Second, currents in a moving reference frame that arise from accumulated charges in the original frame do not generate magnetic fields (LL73). The theoretical implications of these inconsistencies are not discussed further here. The PGR is mentioned in textbooks on electrodynamics such as Jackson (1975) and Pollock and Stump (2001), primarily to show how the PGR fails in the context of electromagnetism. A point is made, however, that the physical principle of relativity demonstrates that “ \mathbf{E} and \mathbf{B} have no independent existence” (Jackson, 1975), which is true for Galilean as well as special relativity.

3 The PGR in Geospace

Discussions of the PGR in the context of geospace are diverse. On the one hand, widely-used textbooks that describe ionospheric electrodynamics (Kivelson and Russell, 1995; Kelley, 2009; Brekke, 2013) do not refer to the PSR or its Galilean limit. In the journal-based literature, several authors discuss inertial reference frames in the context of high latitude electrodynamics (examples are provided in the text below). The reference frame-dependent property of the electric field is mentioned on occasion, but not emphasized or exploited in many cases.

We note that a “preferred reference frame” is a useful construct in plasma physics because of the importance of material media that obey the laws of classical mechanics, such as the plasma and the neutral atmosphere (Vasyliunas and Song, 2005; Leake et al., 2014). Song et al. (2001) derive different versions of Ohm’s law appropriate to reference frames that move with either the plasma bulk flow or the neutrals, including different expressions for the conductivities in these two reference frames. Strangeway and Raeder (2001) consider Galilean invariance extensively in their discussion of how a single set of equations can be used to describe the transition from collisionless magnetohydrodynamic (MHD) in the magnetosphere to collisional MHD in the ionosphere.

An important construct to examine from the perspective of the PGR, and widely seen in the literature, is based on the following quantity: $\mathbf{E} + \mathbf{V} \times \mathbf{B}$, where \mathbf{V} is the velocity of a constituent of the material medium (ions, electrons, neutrals, etc.). In several publications the quantity $\mathbf{E} + \mathbf{V} \times \mathbf{B}$ is referred to as “the electric field in the reference frame of species X ” where \mathbf{V} is the bulk velocity of that species (Vasyliūnas, 2012; Vasyliūnas and Song, 2005; Thayer and Semeter, 2004; Richmond, 1995; Leake et al., 2014; Hidekatsu, 2009; Strangeway, 2012). The symbol \mathbf{E}' is often used to denote this quantity, but in this work, we will use the convention $\mathbf{E}^* = \mathbf{E} + \mathbf{V} \times \mathbf{B}$ (following Vasyliūnas and Song, 2005) to maintain use of the prime symbol to refer to transformations between inertial reference frames. The second term in the expression for \mathbf{E}^* has been referred to as a dynamo electric field when $\mathbf{V} = \mathbf{V}_n$, the velocity of the neutral species (Richmond, 1995). However, $\mathbf{V}_n \times \mathbf{B}$ has the units of electric field, but not the transformation properties of an electric field, so this second term in the expression for \mathbf{E}^* does not refer to a physical electric field.

The GI property of \mathbf{E}^* holds no matter what the velocity \mathbf{V} refers to, whether that of the neutral species, or the ions, etc. The use of \mathbf{E}^* has encouraged the exploration of reference frames tied to material media (e.g. Leake et al., 2014), whereas for electrodynamic quantities there is no need for a preferred inertial reference frame. In many situations applicable to the high latitude ionosphere, *there exists a reference frame for which the electric field is close to zero* (Figure 1). This frame is not tied to any particular medium, but is instructive to consider. As we show in the discussion of Ohm’s law, currents can arise in the absence of electric fields for precisely the reason that currents depend on \mathbf{E}^* rather than \mathbf{E} . Considering this special inertial reference frame – where the electric field is close to zero – provides insight into the physical basis for momentum and energy changes at high latitudes.

3.1 The PGR and Magnetosphere-Ionosphere Coupling

We focus in this paper on the quasi-static electrodynamic perspective where inductive electric fields and inertial effects are not important and where force balance applies between the different species: ions, electrons and neutrals. Even within this restricted domain, where the MI coupling theories describe large-scale plasma convection and slowly-varying currents, the MI coupling literature contains several examples of theoretical formulations that do not adhere to the PSR or PGR. Formulations that do not follow the PGR tend to de-emphasize the important role of the neutral wind – specifically the relative velocity of the neutrals and ions – in MI coupling.

A foundational MI coupling publication is Fedder and Lyon (1987) that couples an ideal MHD model for the magnetosphere with the ionosphere, via the equation (7) in that paper:

$$\nabla \cdot \bar{\bar{\Sigma}} \cdot \mathbf{E} = J_{\parallel} \quad (10)$$

where \mathbf{E} is the electric field in the ionosphere, $\bar{\bar{\Sigma}}$ is the ionospheric conductivity tensor, and J_{\parallel} is the field aligned current entering the ionosphere from the magnetosphere. This equation expresses the fact that the currents flowing between ionosphere and magnetosphere are divergence-free and thus there is no net charge accumulation. The divergence of the horizontally flowing current in the ionosphere given by $\bar{\bar{\Sigma}} \cdot \mathbf{E}$ must be balanced by the field aligned current J_{\parallel} .

Fedder and Lyon (1987) addresses fundamental aspects of MI coupling and is an important scientific contribution. However, Equation (10) does not adhere to the PGR and this could lead to large numerical errors in models that use this equation (see discussion at the end of Section 4). By contrast, in an earlier paper by Vasyliunas (1970), a key MI coupling equation is expressed as follows (equations 7 and 8 in that paper):

$$\nabla \cdot \bar{\Sigma} \cdot (\mathbf{E} + \mathbf{V}_n \times \mathbf{B}) = J_{\parallel} \quad (11)$$

where we have assumed that the magnetic field inclination is vertical to match the approximation in Fedder and Lyon (1987). Equation (11), unlike Equation (10), is consistent with the PGR.

3.2 The PGR and Ohm's Law

Ohm's law is derived from the plasma force balance equations taking collisions into account (Song et al., 2001; Richmond, 1995). These equations can be written, for ions and electrons, as:

$$qN_e(\mathbf{E} + \mathbf{u}_i \times \mathbf{B}) = N_e m_i v_{in}(\mathbf{u}_i - \mathbf{u}_n) + N_e m_i v_{ie}(\mathbf{u}_i - \mathbf{u}_e) \quad (12)$$

$$-qN_e(\mathbf{E} + \mathbf{u}_e \times \mathbf{B}) = N_e m_e v_{en}(\mathbf{u}_e - \mathbf{u}_n) - N_e m_e v_{ei}(\mathbf{u}_i - \mathbf{u}_e) \quad (13)$$

Charge neutrality is assumed such that N_e is the charge density of either electrons or ions. q is the elementary charge, \mathbf{u}_e , \mathbf{u}_n and \mathbf{u}_i are the electron, neutral and ion velocities, respectively, and m_i and m_e are the ion and electron masses, respectively. We assume a single ion species for simplicity. v_{in} , v_{ie} , v_{en} , and v_{ei} are the ion-neutral, ion-electron, electron-neutral and electron-ion momentum transfer rates, respectively. These rates are sometimes referred to as “collision rates”. The following reciprocity relation applies to these rates: $m_k v_{kl} = m_l v_{lk}$, where k, l represent one of the species: electron, ion, or neutral (see Gombosi, 2004), and $k \neq l$. We are ignoring pressure forces and other forces such as gravity, centrifugal, etc. Since these are force balance equations, the acceleration of each species is zero. It is clear these equations are consistent with the PGR and are valid in any inertial reference frame, as can be readily seen by applying the transformations from Table 1. A simplification we assume is that the momentum transfer rates do not depend on the relative velocities; see Richmond (1995) for a statement

regarding this limitation. Equations (12) and (13) apply under quasi steady-state conditions and do not apply during rapid changes or in the presence of dynamical processes associated with reconnection-driven substorms.

A form of Ohm's law that is derived from these force balance equations is as follows (see derivation provided by Song et al., 2001):

$$\mathbf{J} = \sigma_{\parallel} \mathbf{E}_{\parallel} + \sigma_P (\mathbf{E}_{\perp} + \mathbf{V} \times \mathbf{B}) + \sigma_H \mathbf{B} \times (\mathbf{E} + \mathbf{V} \times \mathbf{B})/B \quad (14)$$

where the current density \mathbf{J} is given by $qN_e(\mathbf{u}_i - \mathbf{u}_e)$ (charge neutrality is assumed). σ_{\parallel} , σ_P and σ_H are the parallel, Pedersen and Hall conductivities, respectively. \mathbf{E}_{\perp} is the component of the electric field perpendicular to \mathbf{B} . \mathbf{V} might be the bulk plasma velocity or the neutral wind velocity (see Song et al., 2001, Equations (17) and (22). In our Equation (14), we have corrected typographical errors of Song et al.'s (2001) equation (17)). Song et al. (2001) discuss how the expressions for the conductivities in Equation (14) depend on whether \mathbf{V} represents plasma or neutral wind velocity. The conductivities typically involve terms that are GI, such as collision frequencies and gyrofrequencies that depend on the invariant magnetic field. Since conductivities are GI, this form of Ohm's law is explicitly GI, since both the left-hand side and right-hand side do not change under the Galilean transformation.

Ohm's law Equation (14) could be interpreted to imply that currents must accompany 1) electric fields alone, 2) neutral winds alone, or 3) a combination of the two. The choice of inertial reference frame influences these three different possibilities. If an inertial reference frame is chosen such that the perpendicular electric field is zero, then in that reference frame the currents are not associated with electric fields. Strangeway (2012) also questions whether electric fields cause the currents, instead suggesting that both the electric field and the currents are a consequence of the plasma flow. From the perspective of the PGR, it must be concluded that it is inconsistent to assert that electric fields cause plasma flow, at least in certain reference frames. As we discuss below, it is consistent with the PGR to suggest that plasma motion relative to neutrals caused by magnetospheric dynamics leads to ionospheric currents at altitudes where the conductivities are sufficiently large.

Further clarification is found in Vasyliunas' (2012) work on the “physical basis for ionospheric electrodynamics”. Vasyliunas (2012) suggests that the ionospheric current is primarily a stress-balance current ultimately due to the relative motion between plasma and neutrals. That paper thus makes the direct link between two GI quantities, currents and relative velocities, without the problematic intermediary of the reference frame-dependent electric field. In the discussion section, we remark on the implications of a direct relationship between currents and electric fields, as has been proposed in the context of Subauroral Polarization Stream (SAPS) electric fields.

We are motivated to seek an alternative to the traditional Ohm's law, starting with the same force balance equations (12) and (13), but with all quantities being explicitly invariant to reference frame (see also Strangeway and Raeder, 2001, who discuss the frame-invariance of Ohm's law). Equations (12) and (13) can be written as:

$$qN_e\mathbf{F}_i = N_em_iv_{in}\boldsymbol{\delta v}_{in} + \left(\frac{m_iv_{ie}}{q}\right)\mathbf{J} \quad (15)$$

$$-qN_e\mathbf{F}_e = N_em_ev_{en}\boldsymbol{\delta v}_{en} - \left(\frac{m_iv_{ie}}{q}\right)\mathbf{J} \quad (16)$$

where now all terms in these expressions are explicitly GI. \mathbf{F}_i and \mathbf{F}_e are the Lorentz force per unit volume and per unit charge on the ions and electrons, respectively, and $\boldsymbol{\delta v}_{in}$ and $\boldsymbol{\delta v}_{en}$ are the relative velocities between ions and neutrals and electrons and neutrals, respectively.

Recognizing that the electron and ion Lorentz forces are related as follows:

$$\mathbf{F}_e = \mathbf{F}_i - \left(\frac{1}{qN_e}\right)\mathbf{J} \times \mathbf{B} \quad (17)$$

and substituting \mathbf{F}_e into Equation (16) we find that:

$$\mathbf{J} \times \mathbf{B} = N_em_iv_{in}\boldsymbol{\delta v}_{in} + N_em_ev_{en}\boldsymbol{\delta v}_{en} \quad (18)$$

which shows directly the dependence of the current on the relative velocities of ions and electrons to neutrals, independent of reference frame. (This is essentially the same as Equation 10 of Strangeway and Raeder (2001) without the force term \mathbf{F}). Multiplying Equation (18) on the left by \mathbf{B} and using the appropriate vector identity, we find that:

$$\mathbf{J} = \mathbf{B} \times (N_e m_i v_{in} \delta \mathbf{v}_{in} + N_e m_e v_{en} \delta \mathbf{v}_{en}) / B^2 + \mathbf{B}(\mathbf{B} \cdot \mathbf{J}) / B^2 \quad (19)$$

It is clear from Equation (18) that \mathbf{J} is defined up to an arbitrary vector parallel to \mathbf{B} , so we can write:

$$\mathbf{J} = \mathbf{B} \times (N_e m_i v_{in} \delta \mathbf{v}_{in} + N_e m_e v_{en} \delta \mathbf{v}_{en}) / B^2 + k \mathbf{B} \quad (20)$$

where k is an arbitrary constant. The component of \mathbf{J} perpendicular to \mathbf{B} is of primary interest in the high latitude E-region ionosphere where collisions predominate. Equation (20) shows explicitly that electric fields are not required for horizontal currents to flow at high latitude. What is required is that ionosphere-magnetospheric interaction results in relative motion between ions, electrons and neutrals. We note that Equation (20) is consistent with the physical interpretation of Vasyliūnas (2012) that the “ionospheric current is ... not an Ohmic current in the usual sense.”

For comparable relative velocities with the neutrals, the ion term in Equation (20) dominates by three orders of magnitude over the electron term, for typical conditions at local noon near ~110 km altitude where horizontal currents typically flow (see Figure 1 of Song et al., 2001 for quantitative estimates. At 110 km altitude, the constant preceding the ion-neutral relative velocity is $\sim 2.1 \times 10^{-12}$ versus $\sim 1.0 \times 10^{-15}$ for the electron-neutral term (SI units)). Although FACs originating in the magnetosphere that close horizontally in the ionosphere are often carried by electrons due to their high mobility (Carlson et al., 1998; Sugino et al., 2002), these currents likely do not play an important role in the force balance leading to Ohm’s law. The primary factor is ion-neutral relative velocity.

3.3 The PGR and High Latitude Heating

Energy input from the magnetosphere to the ionosphere increases significantly during geomagnetic disturbances. Poynting's theorem (PT) and Poynting flux are often used in the context of high latitude electrodynamics to understand energy deposition (Kelley, 1989; Thayer and Semeter, 2004; Richmond and Matsuo, 2008).

Poynting's theorem is:

$$\frac{\partial W}{\partial t} = -\nabla \cdot \mathbf{S} - \mathbf{J} \cdot \mathbf{E} \quad (21)$$

where W is the energy density of the electromagnetic (EM) field:

$$W = \frac{1}{2} \left(\epsilon_0 \mathbf{E} \cdot \mathbf{E} + \frac{1}{\mu_0} \mathbf{B} \cdot \mathbf{B} \right) \quad (22)$$

and \mathbf{S} is the Poynting vector:

$$\mathbf{S} = \mathbf{E} \times \frac{1}{\mu_0} \mathbf{B} \quad (23)$$

Physically, Equation (21) represents energy conservation. It states that the rate of change of EM energy density W within a volume equals the energy leaving that volume, via divergence of Poynting flux ($\nabla \cdot \mathbf{S}$), plus the rate of work done by the EM field within the volume ($\mathbf{J} \cdot \mathbf{E}$).

Matsuo and Richmond (2008) divide the electromagnetic work done into two terms as follows:

$$\mathbf{J} \cdot \mathbf{E} = \mathbf{J} \cdot (\mathbf{E} + \mathbf{V}_n \times \mathbf{B}) + \mathbf{V}_n \cdot (\mathbf{J} \times \mathbf{B}) \quad (24)$$

where the first term on the RHS represents the Joule heating or thermal energy contribution (e.g. Cole, 1962; Thayer and Vickrey, 1992) and the second term on the RHS represents the rate at which mechanical energy is transferred to the neutrals. The following standard expression for Joule heating (JH)

$$\mathbf{JH} = \mathbf{J} \cdot (\mathbf{E} + \mathbf{V}_n \times \mathbf{B}) \quad (25)$$

is composed of two terms, each of which depends on reference frame. Following in the same spirit as Equation (18), we wish to write \mathbf{JH} in terms of quantities that are independent of reference frame. Another expression for \mathbf{JH} is given by ηJ_\perp^2 (Cole, 1962) where J_\perp is the current perpendicular to \mathbf{B} and η is the inverse of the Cowling conductivity (see eqn. 11 in Vasyliūnas and Song, 2005 for an expression for η). Therefore, using the component of the current perpendicular to \mathbf{B} from Equation (20), \mathbf{JH} can be written as:

$$\mathbf{JH} = \eta |\mathbf{B} \times (N_e m_i v_{in} \delta \mathbf{v}_{in} + N_e m_e v_{en} \delta \mathbf{v}_{en})|^2 / B^4 \quad (26)$$

where all terms are now invariant to Galilean reference frame.

Equation (26) explicitly relates heating to the relative motions of the different plasma species. Such collisional heating has also been termed “frictional heating” in the literature. The discussion in the appendix of Thayer and Semeter (2004) starts with the more conventional Equation (25) and concludes that heating at high latitudes is due to friction between plasma and neutrals. Vasyliūnas and Song (2005) reach the same conclusion and state that the heating is not “Joule heating in the physical sense” (see also Brekke and Rino, 1978). Strangeway (2012) considers additional forces than are implicit in the momentum balance Equations (12) and (13), but concludes that ionospheric heating predominantly increases plasma temperature due to plasma motion relative to the neutrals.

4 Discussion

A standard interpretation of magnetosphere-ionosphere coupling is that magnetospheric currents lead to electric fields that in turn lead to plasma motion, ion-neutral velocity differences and finally heating (Milan et al., 2017; Cowley, 2000; Kan, 1997). The electric field is not an invariant, so explanations linking currents to electric fields are problematic from the perspective of the PGR. In addition, Vasyliūnas (2001) has shown that electric fields do not cause bulk plasma motion (see also the introduction of Strangeway, 2012). However, the paper by Cole

(1962) states that electric fields induce plasma motion, in contradiction to Vasyliunas (2001) and Strangeway (2012).

Consistency with the PGR is achieved if ion-neutral velocity differences are viewed as the primary causative factor of heating (as per the appendix in Thayer and Semeter (2004) or Vasyliūnas and Song (2005)) and currents (Vasyliūnas (2012)). In the introduction section of Vasyliūnas and Song (2005) it is stated that “by virtue of the plasma momentum equation” the ion-neutral velocity difference is proportional to the current density J . Cause and effect is explicitly considered in the dynamic treatment by Tu et al. (2011) who conclude that heating can be understood from ion-neutral velocity differences without invoking field-aligned currents, ionospheric conductance or the electric field. From a cause-and-effect perspective, ion-neutral velocity difference *causes* the existence of J .

The PGR is relevant to explanations for the high velocity ion flows known as Subauroral ion drifts (SAID), which are often considered as a consequence of Sub-auroral Polarization Stream (SAPS) electric fields. SAPS are postulated to be the result of magnetospheric currents closing in a low-conductivity region of the ionosphere, thus leading to large electric fields (Anderson et al., 1993; Clausen et al., 2012). Currents as the primary driver for high velocity plasma flows in the lower ionosphere is problematic from the perspective of the PGR, because the plasma flow velocity depends on inertial reference frame, but the currents do not. Asserting that currents lead to large SAPS electric fields is similar to reasoning based on Equation (10), which is not consistent with the PGR. Considering large-scale MI coupling (not SAPS), the dynamic simulations of Tu and Song (2016) conclude that the coupling is not established by FACs closing in the ionosphere. From the perspective of the PGR, it is more satisfactory to suggest that large-scale ionospheric convection is driven by the imposed velocity differences between the solar wind and magnetospheric plasmas without requiring an electric field as intermediary in the causal chain.

One might well ask how the velocity of the neutral species (Equation (25)) is relevant when calculating the heating that results from work done by the electromagnetic field, since the neutral species do not experience the electromagnetic force. The answer is that this equation is based on

the single-fluid magnetohydrodynamic (MHD) energy equation (Richmond, 1983), for which the velocity of the fluid is the mass-weighted velocity of the individual species. In the ionosphere, the neutral species dominate in the collisional E-region, so the mass weighted fluid velocity is nearly equal to the velocity of the neutral species. Considering only the electrodynamic contributions to energy, we can equate the energy lost by the electromagnetic field, $\mathbf{J} \cdot \mathbf{E}$, to the work done by the $\mathbf{J} \times \mathbf{B}$ force on the fluid, plus the heating of the fluid:

$$\mathbf{J} \cdot \mathbf{E} = \mathbf{V}_n \cdot (\mathbf{J} \times \mathbf{B}) + JH \quad (27)$$

which immediately implies Equation (25). Thus, the standard expression for Joule heating is closely tied to single fluid MHD theory, whereas the expression for JH embodied in Equation (26) admits of multiple species explicitly, and is simpler to reconcile with the concept of “frictional heating” between species. The appendix of Thayer and Semeter (2004) goes to considerable lengths to reconcile the single fluid and multi-species treatments as they pertain to high latitude heating.

The chain of interactions that lead to large-scale MI coupling would then appear to be:

- Coupling between the solar wind and magnetosphere imparts momentum to the magnetospheric plasma at altitudes of a few Earth radii. This coupling is primarily electrodynamic in character, since collisions between solar wind and magnetospheric ions are infrequent at such altitudes. This is referred to as “mechanical coupling” in Strangeway and Raeder (2001).
- This collisionless interaction, which also leads to magnetic flux being frozen into the magnetospheric plasma, causes magnetospheric plasma motion at progressively lower altitudes, eventually driving ion motion at low altitudes in the ionosphere.
- The momentum imparted to the ionospheric plasma by the magnetospheric plasma motion creates a velocity difference between the ionospheric plasma and the neutral species at ionospheric altitudes, leading to transfer of momentum to the neutral species via collisional processes that become important in the lower ionosphere (for time scales associated with ion-neutral coupling, see Song et al., 2009). The ion-

neutral velocity differences lead to the presence of horizontal currents and heating in the ionosphere (Tu et al., 2011; Vasyliūnas, 2012; Mannucci et al., 2018).

In this picture, partly alluded to in Vasyliunas and Song (2005), the currents are by-products of momentum transfer (as the plasma equations suggest) and not the primary drivers in the causal chain that couples the ionosphere to the magnetosphere (see also Vasyliunas, 2012).

Momentum transfer between ions and neutral species appears central to understanding cause and effect relationships in magnetosphere-ionosphere coupling (e.g. using the multispecies plasma equations in Vasyliunas and Song, 2005). Momentum transfer is independent of inertial reference frame, and so can be considered a primary focus for understanding high latitude processes without difficulty. Conversely, in the high latitude ionosphere, the electric field depends on the inertial reference frame, so electric fields as the primary cause of high latitude electrodynamics can be problematic.

Physical insight from the PGR is particularly relevant to models that relate region 1 currents directly to the convection electric field. For example, Siscoe et al. (2002) writes in the introduction that “ Φ_m (magnetospheric convection potential) is then impressed via equipotential magnetic field lines onto the ionosphere, where it becomes the Φ_{pc} (transpolar ionospheric potential) that generates region 1 currents.” This statement suggests a causative relation between the high latitude convection electric field and region 1 currents. We have discussed above how Ohm’s law (Equation (14)) is not in all cases a relationship between currents and electric fields. Consistency with the PGR does not permit a direct relationship between currents (Galilean invariant) and electric fields (not invariant), without the involvement of other terms. *Differences* between the ion and neutral velocities are Galilean invariant (although the velocities themselves are not). Such velocity differences can play an important role in generating currents.

The role of the neutrals seems to have become de-emphasized in several studies of magnetosphere-ionosphere coupling, where the following equation (or close versions) is often used to relate field aligned currents $J_{||}$ to electric fields:

$$\nabla \cdot (\bar{\bar{\Sigma}} \cdot \mathbf{E}) = J_{\parallel} \quad (28)$$

where $\bar{\bar{\Sigma}}$ is the conductance tensor (Fedder and Lyon, 1987). Siscoe et al. (2002) use a simplified version of this equation. Raeder (2003), and Rothwell and Jasperse (2006) use similar equations to (28), citing Vasyliunas (1970) in that context. However, the relevant equation (8) in Vasyliunas (1970) has the term $\mathbf{E} + \mathbf{V}_n \times \mathbf{B}$ (SI units) in the above equation, instead of \mathbf{E} alone. In the derivation in Wiltberger et al. (2003), the starting point is $\mathbf{J} = \bar{\bar{\Sigma}} \cdot \mathbf{E}$, (their Equation 1), which differs from Ohm's law (our Equation (14)). In these works, the neutral wind term, needed to maintain consistency with the PSR, has been dropped despite its being in the original reference from Vasyliunas (1970).

In a review of large-scale high-latitude ionospheric electrodynamic fields and currents focused on the Assimilative Mapping of Ionospheric Electrodynamics (AMIE) procedure, Lu (2017) discusses the neglect of neutral wind effects that can have a significant impact on JH, citing examples where the neutral winds can enhance or decrease JH by 400% or 40% respectively. In an Earth fixed frame, high latitude convection velocities can reach 1 km/s, which implies that velocity differences between the ions and neutrals can reach similar magnitudes, particularly in the early phases of storms. The $\delta \mathbf{v} \times \mathbf{B}$ magnitudes can reach values of 50 mV/m, comparable to the largest ionospheric electric fields during intense storms. Therefore, MI coupling theories based on formulations that use Equation (28) will be quantitatively challenged by neglecting the relativistic implications of (28) where a velocity term is missing (see Equation (11)). Errors associated with neglecting neutral winds are discussed from a theoretical perspective by Song et al. (2009).

5 Conclusions

We have reviewed the principle of relativity in the context of high latitude electrodynamics, where large-scale ionospheric convection is occurring. For the first time, we have discussed the literature of Galilean electromagnetism (specifically, the treatment of LL73) in this context. The “magnetic limit” of Galilean electromagnetism, which applies when considering MI-coupling, implies that Maxwell's equations are invariant under the Galilean transformation only if the

displacement current term is removed. The magnetic limit also requires that the plasma be quasi-neutral.

The PGR provides insight into the physical basis for magnetosphere-ionosphere coupling. We have considered Ohm's law from the perspective of the PGR and noted that Ohm's law need not be considered as a relationship between currents and electric fields. We suggest that physical insight is gained by considering an inertial reference frame that is not tied to a particular species of the material medium: the reference frame in which the large-scale electric field is close to zero. This provided motivation to develop a version of the equation for the current that is an alternative to Ohm's law, for which the current does not depend on the electric field and for which all terms in the equation are Galilean invariants (Equation (20)). We suggest that more accurate modelling of the coupled MI-system will occur when divergence of horizontal ionospheric currents is based on Equation (11), that includes the neutral winds, versus equations such as (28) that are not compliant with the PGR. Relativistic invariance requires that MI-coupling is strongly dependent on neutral winds. Undersampling of neutral winds, as noted by Lu (2017), is a significant scientific issue that needs to be addressed to improve understanding of MI coupling and to predict space weather. A planned NASA mission, the Geospace Dynamics Constellation (GDC STDT, 2019) will provide valuable new scientific information. A scientific objective of GDC is: "Determine how high-latitude plasma convection and auroral precipitation drive thermospheric neutral winds." The scientific emphasis of GDC on the interactions between charged and neutral particle populations is a needed scientific basis from the perspective of the relativity principle.

Joule heating at high latitudes can be understood without reference to an electric field, as being due to friction between plasma and neutrals, as shown by detailed multi-species plasma calculations (Brekke and Rino, 1978, Vasyliunas and Song, 2005; Thayer and Semeter, 2004 and Strangeway, 2012) and by calculations that account for dynamical effects (Tu et al., 2011). We express JH in a form for which all terms are Galilean invariants, by combining the MHD-based formulation that JH is proportional to J_{\perp}^2 and using Equation (20) to compute J_{\perp} (set $k = 0$). Despite the term "Joule heating" that is associated with experiments where heating is caused by

currents generated by electric fields, high latitude heating cannot depend exclusively on electric fields, the latter being reference frame-dependent.

The Galilean transformation rules for currents and electric fields are very different from each other. It is problematic to suggest that field-aligned currents of magnetospheric origin closing in the ionosphere *cause* horizontal electric fields to appear in the ionosphere (Tu et al., 2011). Such considerations suggest that horizontal currents in the ionosphere flowing through low-conductivity regions are not the primary driver of high-velocity plasma flows known as SAID. Such a causal relation would seem to be reference frame-dependent. At E-layer altitudes in the ionosphere, currents are independent of inertial reference frame, whereas perpendicular electric fields are not. Similarly, equations that relate field aligned currents to the divergence of high latitude horizontal currents must invoke the full Ohm's law, that shows the currents are not dependent solely on high latitude electric fields, but also on the neutral wind.

Acknowledgments

This research was carried out at the Jet Propulsion Laboratory, California Institute of Technology, under a contract with the National Aeronautics and Space Administration. Government sponsorship acknowledged. One of the authors (AJM) wishes to acknowledge constructive discussions with Professor Paul Song, University of Massachusetts at Lowell. We acknowledge sponsorship of the NASA and National Science Foundation Partnership for Collaborative Space Weather Modeling and to the NASA Living With a Star Science program. No data were used in this paper.

References

- Anderson, P. C., W. B. Hanson, and R. A. Heelis (1993), A proposed production model of rapid subauroral ion drifts and their relationship to substorm evolution, *Journal of Geophysical Research*, 98(A4), 6069, doi:10.1029/92ja01975.
- Brekke, A. (2013), *Physics of the Upper Polar Atmosphere*, Springer Berlin Heidelberg, Berlin, Heidelberg, doi:10.1007/978-3-642-27401-5.
- Brekke, A., and C. L. Rino (1978), High-resolution altitude profiles of the auroral zone energy dissipation due to ionospheric currents, *Journal of Geophysical Research*, 83(A6), 2517-2524, doi:10.1029/JA083iA06p02517.
- Clausen, L. B. N., et al. (2012), Large-scale observations of a subauroral polarization stream by midlatitude SuperDARN radars: Instantaneous longitudinal velocity variations, *Journal of Geophysical Research*, 117(A), A05306, doi:10.1029/2011JA017232.
- Cole, K. D. (1962), Joule Heating of the Upper Atmosphere, *Australian Journal of Physics*, 15(2), 223-, doi:10.1071/PH620223.
- Cowley, S. W. H. (2000), Magnetosphere-Ionosphere Interactions: A Tutorial Review, *Magnetospheric Current Systems. Geophysical Monograph 118*. Edited by Shin-ichi Ohtani, 118, 91-, doi:10.1029/GM118p0091.
- De Montigny, M., and G. Rousseaux (2006), On the electrodynamics of moving bodies at low velocities, *European Journal of Physics*, 27(4), 755-768, doi:10.1088/0143-0807/27/4/007.
- Fedder, J. A., and J. G. Lyon (1987), The solar wind-magnetosphere-ionosphere current-voltage relationship, *Geophysical Research Letters* (ISSN 0094-8276), 14(8), 880-883, doi:10.1029/GL014i008p00880.
- Gombosi, T. I. (2004), *Physics of the Space Environment*, Revised ed., 339 pp., Cambridge University Press.
- Heras, J. A. (2010), The Galilean limits of Maxwell's equations, *American Journal of Physics*, 78(10), 1048-1055, doi:10.1119/1.3442798.
- Hidekatsu, J. (2009), Ionospheric Dynamo Process, *Journal of the NICT*, 56, 1-13.
- Jackson, J. D. (1975), *Classical Electrodynamics*, 2 ed., John Wiley and Sons, New York.
- Kan, J. R. (1987), Generation of field-aligned currents in magnetosphere-ionosphere coupling in a MHD plasma, *Planetary and Space Science* (ISSN 0032-0633), 35(7), 903-912, doi:10.1016/0032-0633(87)90068-7.

- Kelley, M. C. (1989), *The Earth's Ionosphere: Plasma Physics and Electrodynamics*, 1st ed., Academic Press, San Diego, CA.
- Kivelson, M. G., and C. T. Russell (Eds.) (1995), *Introduction to Space Physics*, 1 ed., Cambridge University Press, New York.
- Le Bellac, M., and J. M. Lévy-Leblond (1973), Galilean electromagnetism, *Il Nuovo Cimento*, 14(2), 217-234, doi:10.1007/BF02895715.
- Leake, J. E., et al. (2014), Ionized Plasma and Neutral Gas Coupling in the Sun's Chromosphere and Earth's Ionosphere/Thermosphere, *Space Science Reviews*, 184(1-4), 107-172, doi:10.1007/s11214-014-0103-1.
- Lu, G. (2017), Large Scale High-Latitude Ionospheric Electrodynamical Fields and Currents, *Space Science Reviews*, 1-20, doi:10.1007/s11214-016-0269-9.
- Mannucci, A. J., O. P. Verkhoglyadova, X. Meng, and R. McGranaghan (2018), On the role of neutral flow in field-aligned currents, *Annales Geophysicae*, 36(1), 53-57, doi:10.5194/angeo-36-53-2018.
- Matsuo, T., and A. D. Richmond (2008), Effects of high-latitude ionospheric electric field variability on global thermospheric Joule heating and mechanical energy transfer rate, *Journal of Geophysical Research*, 113(A7), n/a-n/a, doi:10.1029/2007JA012993.
- Milan, S. E., et al. (2017), Overview of Solar Wind–Magnetosphere–Ionosphere–Atmosphere Coupling and the Generation of Magnetospheric Currents, *Space Science Reviews*, 1-27, doi:10.1007/s11214-017-0333-0.
- Pal, P. B. (2003), Nothing but relativity, *European Journal of Physics*, 24(3), 315-319, doi:10.1088/0143-0807/24/3/312.
- Parks, G. K. (2007), Importance of electric fields in modeling space plasmas, *Journal of Atmospheric and Solar-Terrestrial Physics*, 69(1-2), 18-23, doi:10.1016/j.jastp.2006.07.009.
- Pollack, G. L., and D. R. Stump (2001), *Electromagnetism*, 1 ed., Addison-Wesley, New York.
- Preti, G., F. de Felice, and L. Masiero (2009), On the Galilean non-invariance of classical electromagnetism, *European Journal of Physics*, 30(2), 381-391, doi:10.1088/0143-0807/30/2/017.
- Raeder, J. (2003), Global Magnetohydrodynamics - A Tutorial, in *Space Plasma Simulation*. Edited by J. Büchner, C. T. Dum and M. Scholer, Lecture Notes in Physics, 615, 212-246, Springer Verlag, Berlin, ISBN 3-540-00698-2.

- Richmond, A. D. (1983), Thermospheric dynamics and electrodynamics, in *Solar-terrestrial Physics*, edited by R. L. Carovillano, Forbes, J.M. , pp. 523-607, D. Reidel, Norwell, MA, USA.
- Richmond, A. D. (1995), Ionospheric electrodynamics, in *Handbook of Atmospheric Electrodynamics*, vol. 2, edited by H. Volland, 249–290, CRC Press, ISBN 978-0849325205.
- Richmond, A. D., and J. P. Thayer (2000), Ionospheric Electrodynamics: A Tutorial, in *Magnetospheric Current Systems*, American Geophysical Union Monograph volume 118, Edited by Shin-ichi Ohtani, pp. 118:131, doi:10.1029/GM118p0131.
- Rothwell, P. L., and J. R. Jasperse (2006), Modeling the connection of the global ionospheric electric fields to the solar wind, *Journal of Geophysical Research*, 111(A3), A03211-03216, doi:10.1029/2004JA010992.
- Rothwell, P. L., and J. R. Jasperse (2007), A coupled solar wind-magnetosphere-ionosphere model for determining the ionospheric penetration electric field, *Journal of Atmospheric and Solar-Terrestrial Physics*, 69(10-11), 1127-1134, doi:10.1016/j.jastp.2006.08.013.
- Rousseaux, G. (2013), Forty years of Galilean Electromagnetism (1973–2013), *The European Physical Journal Plus*, 128(8), 105207-105215, doi:10.1140/epjp/i2013-13081-5.
- Siscoe, G. L., G. M. Erickson, B. U. Ö. Sonnerup, N. C. Maynard, J. A. Schoendorf, K. D. Siebert, D. R. Weimer, W. W. White, and G. R. Wilson (2002), Hill model of transpolar potential saturation: Comparisons with MHD simulations, *Journal of Geophysical Research*, 107(A), 1075, doi:10.1029/2001JA000109.
- Song, P., T. I. Gombosi, and A. J. Ridley (2001), Three-fluid Ohm's law, *Journal of Geophysical Research*, 106(A5), 8149-8156, doi:10.1029/2000JA000423.
- Song, P., V. M. Vasyliūnas, and X. Z. Zhou (2009), Magnetosphere-ionosphere/thermosphere coupling: Self-consistent solutions for a one-dimensional stratified ionosphere in three-fluid theory, *Journal of Geophysical Research*, 114(A8), A08213-08211, doi:10.1029/2008JA013629.
- Strangeway, R. J. (2012), The equivalence of Joule dissipation and frictional heating in the collisional ionosphere, *Journal of Geophysical Research*, 117(A), A02310, doi:10.1029/2011JA017302.
- Strangeway, R. J., and J. Raeder (2001), On the transition from collisionless to collisional magnetohydrodynamics, *Journal of Geophysical Research*, 106(A), 1955-1960, doi:10.1029/2000JA900116.
- Sugino, M. (2002), Relative contribution of ionospheric conductivity and electric field to ionospheric current, *Journal of Geophysical Research*, 107(A10), 10,031-015, doi:10.1029/2001JA007545.

- Thayer, J. P., and J. F. Vickrey (1992), On the contribution of the thermospheric neutral wind to high-latitude energetics, *Geophysical Research Letters*, 19(3), 265-268, doi:10.1029/91GL02868.
- Thayer, J. P. (2000), High-latitude currents and their energy exchange with the ionosphere-thermosphere system, *Journal of Geophysical Research*, 105(A), 23015-23024, doi:10.1029/1999JA000409.
- Thayer, J. P., and J. Semeter (2004), The convergence of magnetospheric energy flux in the polar atmosphere, *Journal of Atmospheric and Solar-Terrestrial Physics*, 66(1), 807-824, doi:10.1016/j.jastp.2004.01.035.
- Tu, J., P. Song, and V. M. Vasyliūnas (2011), Ionosphere/thermosphere heating determined from dynamic magnetosphere-ionosphere/thermosphere coupling, *Journal of Geophysical Research*, 116(A9), A09311-09319, doi:10.1029/2011JA016620.
- Tu, J., and P. Song (2016), A two-dimensional global simulation study of inductive-dynamic magnetosphere-ionosphere coupling, *Journal of geophysical research*, 121(1), 11-, doi:10.1002/2016JA023393.
- Vasyliūnas, V. M. (1970), Mathematical Models of Magnetospheric Convection and Its Coupling to the Ionosphere, in *Particles and Field in the Magnetosphere*, 17 (Chapter 6), 60-71, doi:10.1007/978-94-010-3284-1_6, 1970.
- Vasyliūnas, V. M. (2001), Electric field and plasma flow: What drives what?, *Geophysical Research Letters*, 28(1), 2177-2180, doi:10.1029/2001GL013014.
- Vasyliūnas, V. M., and P. Song (2005), Meaning of ionospheric Joule heating, *Journal of Geophysical Research*, 110(A), A02301, doi:10.1029/2004JA010615.
- Vasyliūnas, V. M. (2012), The physical basis of ionospheric electrodynamics, *Annales Geophysicae*, 30(2), 357-369, doi:10.5194/angeo-30-357-2012.
- Verkhoglyadova, O. P., X. Meng, A. J. Mannucci, M. G. Mlyneczek, L. A. Hunt, and G. Lu (2017), Ionosphere-thermosphere energy budgets for the ICME storms of March 2013 and 2015 estimated with GITM and observational proxies, *Space Weather*, 27(14), 425-423, doi:10.1002/2017SW001650.
- Zhang, S. P., and G. G. Shepherd (2002), Neutral winds and emission rates in the lower thermosphere as measured with WINDII/UARS during the April 4–5th 1993 and February 1994 geomagnetic storms, *Journal of Atmospheric and Solar-Terrestrial Physics*, 64(8-11), 1201-1214, doi:10.1016/s1364-6826(02)00069-x.

843

844