

A Review of the Dynamics of Subduction Zone Initiation in the Aegean Region

Elizabeth Catlos¹ and Ibrahim Cemen²

¹The University of Texas at Austin, Jackson School of Geosciences, Department of Geological Sciences

²The University of Alabama

November 22, 2022

Abstract

The Hellenic arc, where the African (Nubian) slab subducts beneath the Aegean and Anatolian microplates, has emerged as a type-locality for understanding subduction dynamics, including slab tear, slab fragments, drips, and transfer zones. Based on field evidence and geophysical, tectonics, and geochemical studies, it has been recognized that the subducting African slab is a primary driver for extension in the Aegean and Anatolian microplates and plays a significant role in accommodating present-day westward extrusion of the Anatolian microplate. Thus, understanding the Hellenic arc subduction zone initiation (SZI) age is critical in deciphering ancient mantle flow, how plate tectonics is maintained, and the mechanisms involved in triggering the onset of subduction. The SZI for the Hellenic arc has two disparate ages based on different lines of evidence. A Late Cenozoic (Eocene-Pliocene) SZI is proposed using the analysis of topography combined with estimates of slab age and depth, paleomagnetism, the timing of metamorphism, and volcanic activity, and timing of sedimentation within its accretionary wedge, the Mediterranean Ridge. This age follows an induced-transference SZI model, where a new subduction zone initiates following the jamming of an older subduction zone by buoyant crust due to regional compression, uplift, and underthrusting. A Late Cretaceous-Jurassic SZI age has also been proposed using reconstructions of images of subducted slabs seen using tomography and timing of obducted ophiolite fragments thought to be related to the system. In this case, the induced-transference SZI model fails, and a single subduction zone persists. As a result, continental lithospheric fragments and the ancient oceans between them become incorporated into the overall system without creating a new subduction zone. The presence of a long-lived subduction zone has implications for understanding Earth's mantle dynamics and how plate tectonics operates. This paper describes and summarizes the evidence for both models in the Aegean-Western Anatolia region.

A Review of the Dynamics of Subduction Zone Initiation in the Aegean Region

E.J. Catlos¹ and I. Cemen²

¹ The University of Texas at Austin, Jackson School of Geosciences, Department of Geological Sciences, 2275 Speedway Stop C9000, Austin, TX 78712-1722, USA

² Department of Geological Sciences, The University of Alabama, Box 870338, Tuscaloosa, AL 35487-0338, USA

Corresponding author: Elizabeth Catlos (ejcatlos@jsg.utexas.edu)

Index Terms

8104 Continental margins: convergent

Keywords

Subduction zones, Hellenic arc, Aegean region, Western Anatolia, Tectonics

Abstract

The Hellenic arc, where the African (Nubian) slab subducts beneath the Aegean and Anatolian microplates, has emerged as a type-locality for understanding subduction dynamics, including slab tear, slab fragments, drips, and transfer zones. Based on field evidence and geophysical, tectonics, and geochemical studies, it has been recognized that the subducting African slab is a primary driver for extension in the Aegean and Anatolian microplates and plays a significant role in accommodating present-day westward extrusion of the Anatolian microplate. Thus, understanding the Hellenic arc subduction zone initiation (SZI) age is critical in deciphering ancient mantle flow, how plate tectonics is maintained, and the mechanisms involved in triggering the onset of subduction. The SZI for the Hellenic arc has two disparate ages based on different lines of evidence. A Late Cenozoic (Eocene-Pliocene) SZI is proposed using the analysis of topography combined with estimates of slab age and depth, paleomagnetism, the timing of metamorphism, and volcanic activity, and timing of sedimentation within its accretionary wedge, the Mediterranean Ridge. This age follows an induced-transference SZI model, where a new subduction zone initiates following the jamming of an older subduction zone by buoyant crust due to regional compression, uplift, and underthrusting. A Late Cretaceous-Jurassic SZI age has also been proposed using reconstructions of images of subducted slabs seen using tomography and timing of obducted ophiolite fragments thought to be related to the system. In this case, the induced-transference SZI model fails, and a single subduction zone persists. As a result, continental lithospheric fragments and the ancient oceans between them become incorporated into the overall system without creating a new subduction zone. The presence of a long-lived subduction zone has implications for understanding Earth's mantle dynamics and how plate tectonics operates. This paper describes and summarizes the evidence for both models in the Aegean-Western Anatolia region.

1 Introduction

Subduction zones form when two lithospheric plates converge, and one plate abruptly descends beneath the other (Figures 1 and 2) (e.g., White et al., 1970; Hayes, 2018; Stern & Gerya, 2018; Crameri et al., 2020). Large magnitude earthquakes, tsunamis, volcanic eruptions, and landslides occur near and are caused by this specific plate boundary. They are considered exceptional geological environments for recording significant ground-level changes that can trigger tsunamis, impact ground motion, and climate change. Earthquakes that occur in such zones and those triggered by the subduction process far-afield have global consequences. Understanding the dynamics of subduction zones involves diverse and multidisciplinary studies, critical for understanding their associated hazards and how they have influenced the dynamics of plate tectonics over Earth's history (e.g., Stern, 2004; Gerya, 2011; Le Pichon et al., 2019; Crameri et al., 2020).

The Aegean and Anatolian microplates (Figure 1) are significantly impacted by the dynamics of the subducting northern portion of the African (Nubian) plate, which has emerged as the primary driver for extension and the development of metamorphic core complexes in the Aegean region (e.g., Jolivet & Faccenna, 2000; Çemen et al., 2006; Dilek & Sandvol, 2009; van Hinsbergen et al., 2010; van Hinsbergen). The Hellenic and Cyprus arcs are the surface expression of the subducting Nubian Plate and eastern Mediterranean lithosphere beneath the Aegean and Anatolian microplates, respectively (e.g., Le Pichon & Angelier, 1979; Angelier et al., 1982; Anastasakis & Kelling, 1991; Papazachos et al., 2000; Ergün et al., 2005; Ganas & Parsons, 2009; Hall et al., 2009; Biryol et al., 2011; Royden & Papanikolaou, 2011; Hall et al., 2014; Symeou et al., 2018; Ventouzi et al., 2018).

Constraints regarding the subduction zone initiation (SZI) age of the present-day expression of the Hellenic arc developed from several independent approaches include timing of sedimentation within the intensely folded and faulted rocks of Mediterranean Ridge accretionary prism (Figures 1 and 2), paleomagnetism, the analysis of topography combined with estimates of slab age and depth, reconstructions of subducted slabs using tomography, and the timing of metamorphism and volcanic activity. SZI is defined as the onset of downward plate motion forming a new slab, which later evolves into a self-sustaining subduction zone (Crameri et al., 2020). Some studies suggest a Cenozoic SZI age for Hellenic arc, although estimates vary significantly, from the Eocene-Pliocene (e.g., Meulenkamp et al., 1988; Spakman et al., 1988; Papadopoulos, 1997; Brun & Sokoutis, 2010; Le Pichon et al., 2019) to Mesozoic (Late Cretaceous-Jurassic) (Faccenna et al., 2003; van Hinsbergen et al., 2005; Royden & Papanikolaou, 2011; Jolivet et al., 2013; Malandri et al., 2017; Crameri et al., 2020, van Hinsbergen et al., 2021).

The disparity in the SZI age of onset of Nubian slab subduction along the Hellenic arc is significant as it impacts the tectonic history of the entire Aegean-Anatolian region, one of the most rapidly deforming regions across the Alpine-Himalayan chain. The region has emerged as the type-locality for understanding subduction zone dynamics, including slab tear, slab fragments, drips, and the role of transfer zones triggered by subduction. Understanding its SZI is also critical in deciphering ancient mantle flow, how plate tectonics is maintained, and the mechanisms involved in triggering the onset of subduction, among other factors (e.g., Crameri et al., 2020; van Hinsbergen et al., 2021). This paper aims to summarize the approaches and results of studies that strive to constrain the SZI age of the African (Nubian) slab beneath the Aegean microplate that led to the formation of the Hellenic arc.

2 Geometry of the Hellenic arc (Greece to Western Turkey)

2.1 Definitions

The **Hellenic subduction system** extends ~1200 km from approximately 37.5°N, 20.0°E offshore the island of Zakynthos to 36.0°N, 29.0°E offshore of the island of Rhodes (Ganas & Parsons, 2009; Le Pichon et al., 2019; Papanikolaou, 2021) (Figures 2 and 3). The system defines the boundary between the northern portion of the **Nubian Plate** and the southern extent of the **Aegean microplate** within the central Mediterranean region (Pearce et al., 2012) and is sometimes referred to as the **Aegean subduction zone** (Wortel et al., 1990; Biryol et al., 2011; Bleier et al., 2007; Polat & Ozel, 2007; Taymaz et al., 2007; Crameri et al., 2020). This boundary between the **Aegean microplate** portion of Eurasia (Nyst & Thatcher, 2004) and the subducting Nubian slab is presently characterized by a strong curvature and fast slab rollback (e.g., Faccena et al., 2013; Evangelidis, 2017). Presently, the African plate advances towards Eurasia NNW at a rate of 5 mm/yr (Fernandes et al., 2003; Ganas & Parsons, 2009), but it subducts northward beneath Crete at a significantly faster rate of 35 mm/yr (McKenzie, 1972; Reilinger et al., 2006). The Aegean area also records the highest deformation rate along the entire Africa/Eurasia convergence zone (McClusky et al., 2000; Kassaras et al., 2005).

The Aegean and Anatolia microplates are sometimes classified as the single **Aegean-Anatolian microplate** (e.g., Jackson, 1994; Oral, 1995; Doutsos & Kokkalas, 2001; Le Pichon et al., 1995) with an Euler pole located north of the Sinai Peninsula (Cianetti et al., 2001). The **Anatolian microplate** itself is a distinct entity that includes over two-thirds of the country of Turkey (Figure 1) (Le Pichon et al., 1995; Oral et al., 1995; Reilinger et al., 1997; Papazachos, 1999). Şengör & Zabcı (2019) consider the whole of Turkey and the Balkan Peninsula within a plate boundary zone.

The Nubian Plate includes the African continent. When Somalia is part of the definition, it is referred to as the **African Plate** or the **Nubia-Somalia Plate** (e.g., McClusky et al., 2003). The African plate itself is defined by Nubia to the west and Somalia to the east, both of which are separated by a diffuse plate boundary (Chase, 1978, Gordon & Stein 1992, Lemaux et al., 2002; McClusky et al., 2003; DeMets & Merkouriev, 2016). The African plate is geographically significant and is, at present, the third-largest major tectonic plate (Gaina et al., 2013). A wide range of tectonic interactions characterizes its boundaries and internal dynamics, which were influenced by the collision between the plate and Eurasia (e.g., Meijer & Wortel, 1999; McClusky et al., 2003; Fernandes et al., 2006; Catalano et al., 2008; DeMets & Merkouriev, 2016). The plate boundary between the Indian and African plates is over 12,000-km-long and may have initiated around 105 Ma in the Early Cretaceous (e.g., van Hinsbergen et al., 2021). Rifting in the Gulf of Aden and the Red Sea signaled the break-off of the Arabian peninsula from Africa at 29-24 Ma (e.g., Bosworth et al., 2005; Wolfenden et al., 2005) fragmentation of the Africa plate and the onset of motion between Somalia and Nubia. Alternatively, the separation may have been more recent (11-10 Ma, Wolfenden et al., 2004; Keranen & Klemperer, 2008; Corti, 2009; see discussion in DeMets & Merkouriev, 2016).

The overriding Aegean microplate (Nyst & Thatcher, 2004) consists mainly of continental lithosphere with a similar thickness as the Anatolian plate (e.g., Zhu et al., 2006; Sodoudi et al., 2006). Large crustal thickness variations exist from western Greece to eastern Anatolia (20–47 km), but within specific regions of the Anatolian plate, thickness appears fairly uniform (Özacar et al., 2010; Karabulut et al., 2019). Moho depth across the Anatolian plate varies between 24

and 48 km, with the thinnest crustal thickness located on the coast of western Turkey and the thickest in Eastern Turkey (Tezel et al., 2013). The Aegean Sea crustal thickness averages ~25 km (Zhu et al., 2006; Tirel et al., 2004; Kind et al., 2015), similar to thickness beneath the central Menderes Massif at 25-30 km (Zhu et al., 2006; Karabulut et al., 2019). Estimates in the eastern Mediterranean overall show the Moho at an average 30 km depth (Marone et al., 2003). The Aegean microplate crustal thickness has thinner southern and central parts (20–22 km) compared to its northern portions (25–28 km) (Karagianni et al., 2005; Sodoudi et al., 2006). An exception to this observation is the crustal thickness beneath western Crete, 32.5-35 km or up to 45-50 km (Bohnhoff et al., 2001; Meier et al., 2004; Snopek et al., 2007). The eastern part of Crete appears thinner (24-26 km) compared to its western portion (32.5 km) (Bohnhoff et al., 2001). Crustal thickness also decreases to 15 km below the central Cretan Sea and to 17 km, approximately 100 km off the southern coast of Crete (Bohnhoff et al., 2001). A crustal thickness of ~40 km is reported for western Greece (Karagianni et al., 2005), north of Antalya Bay (Karabulut et al., 2019), Eastern Anatolia (Kind et al., 2015), and the Anatolian plateau (e.g., Saunders et al., 1998). In Eastern Anatolia, the crustal thickness increases towards the north, from 35 km at the Arabian foreland to 45 km further north (Özacar et al., 2010).

The subducting portion of the African plate has multiple names based on the time frame that was active and differentiate it from other nearby subduction zone systems. In the Aegean and Western Anatolia region, the subducting plate is sometimes referred to as the **Aegean slab** (Wortel & Spakman, 2000; Widiyantoro et al., 2004; Vanderhaeghe et al., 2007; Biryol, 2009; van der Meer et al., 2018; El-Sharkawy et al., 2020), the **African-Mediterranean slab** (Sachpazi et al., 2016), the **Hellenic plate** or **Hellenic slab** (Chang et al., 2010; Vernant et al., 2014; Sachpazi et al., 2016; Confal et al., 2018; Hayes, 2018; Wei et al., 2019; Blom et al., 2020). The Aegean slab usually describes the system during the Late Cretaceous, whereas the other specific names distinguish the Aegean Sea and Western Anatolia subduction system from that beneath Antalya and Cyprus. The subducting slab near Cyprus is referred to as the **Cyprus slab** (Confal et al., 2018), **Cyprus/Tethys slab** (Taylor et al., 2019), **Cyprus–Bitlis slab** (Govers and Fichtner, 2016). The **Antalya slab** is recently defined between the Hellenic and Cyprus arcs (Güvercin et al., 2021).

The western edge of the Hellenic arc is bounded by the **Kephalonia (also Kefalonia, Cephalonia) Transform Zone (KTZ)** (Figure 1) (Bocchini et al., 2018; Hansen et al., 2019). This feature is a dextral strike-slip system that delineates the boundary between the Apulian platform (Adria microplate) and the Eurasian plate or Aegean microplate (Kokinou et al., 2006; Pearce et al., 2012). The KTZ has structural trends and seismicity that extend onshore, and it is the most active zone of shallow seismicity in the broader Aegean region (Kokinou et al., 2006). Over the last 1–4 Myr, the KTZ has linked to the **North Anatolian Fault (NAF)** system via the **Central Hellenic Shear Zone** (Papanikolaou & Royden, 2007; Reilinger et al., 2010; Vassilakis et al., 2011; Royden & Papanikolaou, 2011; Halpaap et al., 2018). The KTZ is considered one of the most seismically active zones in Europe (Pearce et al., 2012; Halpaap et al., 2018) and has been represented as a vertical tear between oceanic and continental lithosphere (Suckale et al., 2009), forming a Subduction-Transform-Edge-Propagator (STEP) fault (Govers & Wortel, 2005). The STEP fault may be in its initial stages of forming (Evangelidis, 2017; Özbakır et al., 2020), or the slab may have entirely detached (Wortel & Spakman, 2000). A smooth transition is also proposed between two segments without a tear between, at least at depths shallower than 100 km (Pearce et al., 2012; Halpaap et al., 2018). Nine STEP structures may exist beneath southern Greece, segmenting the subducting African slab and contributing to seismicity and

deformation (Sachpazi et al., 2016). Overall, the southwestward expansion and stretching of the Aegean microplate during Plio-Quaternary is accommodated by a northern right lateral tectonic boundary defined by the KTZ and NASZ and a southern left-lateral tectonic boundary, marked by the Pliny and Strabo trenches (Sakellariou et al., 2013).

The active right-lateral NAF and **North Anatolian Shear Zone (NASZ)** extends for ~1200 km from the Karliova triple junction through the Sea of Marmara and Biga Peninsula (Figure 1) (Ketin, 1948; Barka, 1992; Armijo et al., 1999; Şengör & Zabcı, 2019). The NASZ contains the NAF, which accommodated from 25 to 110 km of displacement, depending on location since the Late Miocene (Westaway 1994; Yoshioka 1996; Armijo et al., 1999; Hubert-Ferrari et al. 2002; Şengör & Zabcı, 2019). The fault system disrupts the Aegean extensional domain in its northern section and may have been activated due to subduction rollback processes (e.g., Flerit et al., 2004; Facenna et al., 2006). Current timing estimates suggest the NAF initiated in east Anatolia at 10-12 Ma, and propagated westward through central Anatolia from 7-6 Ma, and was activated in the Aegean region by the Pliocene (Şengör et al., 2005; Facenna et al., 2006). Its western and eastern terminations of the NASZ are poorly defined (Barbot & Weiss, 2021), but the system itself presently accommodates ~24 mm/year of slip along northern Turkey (McClusky et al., 2000; Bulut et al., 2018).

Westward extrusion of the Anatolian plate accommodated by the NASZ and NAF dominates the present-day geodynamics of the Aegean microplate such that active extensional strain has been stated to be absent, except near the Corinth rift, south Viotia, south of Evia, and across the Sperchios-Kammena Vourla rift (Brooks & Ferentinis 1984; Papanikolaou and Royden, 2007; Chousianitis et al., 2013, 2015). However, seismogenic faults in the internal Aegean domain associated with the Hellenic subduction arc are characterized by pure normal and strike-slip kinematics or by a combination, and active thrusting is limited to the central and western sectors of the Hellenic subduction zone and the offshore regions external with respect to it (Maggini & Caputo, 2020). The nature of earthquakes varies with Hellenic arc depth, with normal senses of motion at depths up to 40 km and strike-slip faulting present parallel to strike for events deeper than 40 km (Benetatos et al., 2004). One of the strongest earthquakes of the 20th century, with an Mw 7.7 (or 7.8), occurred in the area of the South Aegean off the coast of the island of Amorgos on 09 July 1956 03:11 UTC (Figure 1) (Okal et al., 2009; Alatza et al., 2020). This event has debated focal mechanisms, as either strike-slip or normal faulting geometries (Okal et al., 2009). An Mw 4.4 earthquake with normal motion recently occurred near its epicenter on 11/27/2018 (23:16 UTC, 36.7565°N, 25.877 °E). A normal sense of motion also is found with some recent earthquakes near the NASZ, including 6/12/2017 Mw 6.2 (12:28 UTC, 38.8486°N, 26.313°E) and 2017 Mw 5.3 (10:58 UTC, 39.5275 °N, 26.1373 °E), likely associated with transtensional motion.

The Hellenic arc itself is separated into a western portion based on well-defined bathymetry. This portion is sometimes termed the **Western Hellenic Subduction Zone or Western Hellenic Arc and Trench (WHA-T) system** and extends ~400 km NW from the central Adriatic Sea to the west coast of Crete (Figures 1 and 3) (Papadopoulos et al., 2010; Pearce et al., 2012; Hansen et al., 2019). This part of the Hellenic Arc is divided into northern and southern sections termed the **Northern Hellenic arc** and **Southern Hellenic arc** (Royden & Papanikolaou, 2011). The Southern Hellenic arc exhibits classical features, but in the north, thick continental crust is subducted beneath northern Greece (Pearce et al., 2012; Halpaap et al., 2018). As a result, this area has significant variations in subduction rate, trench retreat, the occurrence of deep

seismicity, and overriding plate extension, likely related to a change in convergence regime from the subduction of oceanic crust in the south to continental crust in the north (Papanikolaou & Royden, 2007; Halpaap et al., 2018). The southern portion of the Hellenic arc near the island of Crete is also affected by the subduction of continental crust (e.g., Meier et al., 2007).

Overall, the Hellenic arc has three distinct regions: an outer compressional non-volcanic arc, a volcanic arc, and an extensional back-arc region covering the broader Aegean Sea region (Figures 1 and 2) (McKenzie 1972; Papazachos, 2019). The Pliocene-Quaternary volcanic arc is located in the Methana Peninsula, along the islands of Milos and Santorini in the Cycladic archipelago, and the island of Nisryos in the Dodecanese (Figure 1) (Le Pichon and Angelier, 1979; McKenzie, 1972; Papazachos and Nolet, 1997, see review in Scoon, 2021). The majority of the subducting Nubian Plate is oceanic, except along the central sector of its southern margin, where the Mediterranean Ridge accretionary complex collides with the African continental margin (Chaumillon & Mascle, 1997; Westbrook & Reston, 2002). High-resolution scattered wave images of the northern portion of the Western Hellenic arc also show the subduction of thick continental crust beneath northern Greece (Pearce et al., 2012; Halpaap et al., 2018) and Crete (e.g., Meier et al., 2007).

The connection of the western portion of the Hellenic arc to its central and eastern sections is diffuse (Beißer et al., 1990; Shaw & Jackson, 2010; Biryol et al., 2011; Özbakır et al., 2013). The eastern portions of the Hellenic arc are sometimes referred to as the **East Hellenic Arc and Trench system (HA-T)** (Papadopoulos et al., 2007). The boundary between the Hellenic and Cyprus arcs in this area is obstructed by up to 300-km wide, 6-10 km-thick section of sediments that comprise the **Mediterranean Ridge** (Figures 1-3; Heezen & Ewing, 1963; Emery et al., 1966; Le Pichon et al., 1982; Kenyon et al., 1982; Kastens et al., 1992; Foucher et al., 1993; Westbrook & Reston, 2002; Kopf et al., 2003). This accretionary complex is the largest structural unit of the Eastern Mediterranean Sea, extending ~2000 km from the Calabrian Rise east of Greece to the Florence Rise (Liminov et al., 1996; Cita et al., 1996). The front of subduction of the Hellenic arc is located south of the Mediterranean Ridge (Figures 1-3) (e.g., Le Pichon et al., 1995; Mascle & Chaumillon, 1997; Mascle et al., 1999; Jost et al., 2002; Westbrook & Reston, 2002; Meier et al., 2007; Jolivet et al., 2013).

The significant size of the Mediterranean Ridge indicates that the Hellenic arc is an accretionary as opposed to an erosive margin (e.g., Von Huene and Scholl, 1991, Von Huene and Scholl, 1993, Cloos and Shreve, 1996; Clift and Vannucchi, 2004). In this case, material is transferred from the subducting plate to the fore-arc wedge and plate boundary zone (e.g., DeFranco et al., 2008; Scholl and Von Huene, 2009). This type of margin is favored when subduction rates are low and seismic moment release rates are high (Le Pichon et al., 1993; Clift and Vannucchi, 2004; DeFranco et al., 2008). However, the Mediterranean Ridge accretionary complex is unusual compared to others worldwide. It formed in a continent-continent collisional setting and has shallow, Messinian-age evaporites (e.g., Cita et al., 1996; Chaumillon & Mascle, 1997). These evaporites influence its deformation and growth rates due to their mechanical properties and effect on fluid flow and pressure (Kastens, 1991; Westbrook & Reston, 2002; Kopf et al., 2003). Extensive mud volcanism associated with the Mediterranean Ridge began in the early Pleistocene 1.75-1.25 Ma (Nikitas et al., 2021).

Understanding the development of the Mediterranean Ridge is critical to determining the Hellenic arc SZI because it is thought to grow via off scraping against a backstop formed by the Alpine nappes of the Hellenic Arc (Kastens, 1991) at a very fast rate (10 km/Myr; Kastens, 1991;

Kopf et al., 2003; Papanikolaou, 2021). The ridge itself varies in geometry along strike (Cita et al., 1996; Chaumillon & Mascle, 1997; Westbrook & Reston, 2002; Kopf et al., 2003; Andronikidis et al., 2017; Papanikolaou, 2021). The wedge accumulates sediments in its western and eastern portions, but the ridge behaves unlike a typical accretionary complex in the central portion between Libya and Crete, where it experienced maximum deformation, and incipient collision had already caused uplift of the area (Chaumillon and Mascle, 1997; Kopf, 2002). Crete is situated on the central forearc of the Hellenic subduction zone and is underlain by material from two microcontinents, leading to larger observed crustal thicknesses (~50 km, e.g., Thomson et al. 1998; Stöckhert 1999; Meier et al., 2004; 2007).

South of Crete, the **Hellenic trenches**, Ptolemy, Pliny, and Strabo (Figure 1) developed between the Mediterranean Ridge and volcanic arc. These trenches are not classical ocean trenches, as earthquakes beneath them originate along low-angle thrusts at 20–40 km (Taymaz et al., 1990; Shaw & Jackson, 2010). Instead, they develop due to back-thrusting beneath the northern edge of the accretionary complex (Galindo-Zaldivar et al., 1996; Westbrook & Reston, 2002) or due to the tearing of the Nubian slab (Özbakır et al., 2013). The Hellenic Trench has been described as the surface expression of a steep (~30°) reverse fault splaying off the deeper underlying thrust-fault interface of the subduction zone (Shaw et al., 2008; Shaw & Jackson, 2010).

Low-angle thrust faults along the Aegean coast associated with subduction zone tectonics pose significant tsunami hazards (e.g., Tinti et al., 2005; Basili et al., 2013; Howell et al., 2015; Bocchini et al., 2020). Offshore Crete Island is considered one of the most tsunamigenic areas in the entire Mediterranean Sea region (Papadopoulos et al., 2010; Triantafyllou et al., 2019). However, the complexity of the overall Hellenic arc plate boundary combined with its aseismic nature makes earthquake data alone a misleading guide for identifying the likely sources of tsunamigenic earthquakes (Yolsal et al., 2007; England et al., 2015; Howell et al., 2015). Tsunamigenic earthquakes infrequently occur in the eastern Mediterranean (Papazachos and Dimitriu, 1991; Papadopoulos et al., 2007). An evaluation of historical data, including the 1956 Amorgos event that generated the largest of the most recent tsunamis, indicate that a likely trigger of some past tsunamis in the region were submarine landslides generated by earthquakes (e.g., Dominey-Howes, 2002; Okal et al., 2009; Ebeling et al., 2012). Factors that contribute to slope instability across portions of the Hellenic Arc include its sloping bottom, thick accumulations and high rates of recent sedimentation, the presence of closely spaced active faults, active earthquakes, and diapirism (e.g., Ferentinos, 1990; Hooft et al., 2017). The eruption of Santorini (Figure 1) in 1610 BCE generated a tsunami that affected civilizations throughout the eastern Mediterranean (Dominey-Howes, 2004; Friedrich, 2006; Marinatos, 1939, Papadopoulos, 2015; Hooft et al., 2017). Detailed bathymetry across the Mediterranean is critical in understanding tsunami propagation and mitigating its impacts (Figure 1) (e.g., CIESM, 2011).

The **Rhodes Basin** bounds the eastern portion of the Hellenic arc (Figure 1) (Ganas & Parsons, 2009) and is a deep depression lying in continuity with the Pliny and Strabo trenches. The depth of the basin is >4000 m and is one of the deepest portions of the Mediterranean Sea (Woodside et al., 2000). The Rhodes basin may represent an un-subducted portion of the deep Mesozoic Levantine basin (Rotsein & Ben-Avraham, 1985) or a remnant of a former upper-Miocene subduction trench that remained after a shift in the primary convergence zone (Mascle et al., 1986). Deep faults buried beneath the zone may mark the onset of extension (Woodside et al., 2000). Hall et al. (2009) suggest a two-part history of the basin. Following Miocene

convergence, the basin experienced middle Pliocene-Quaternary sinistral transpression due to actively curving Hellenic arc and change in the convergence vector of the African plate. Slab tear has been proposed to interpret the presence and structures within the deep Rhodes Basin (Woodside et al., 2000; Faccenna et al., 2014).

A STEP is also suggested for the transition between the Cyprus and Hellenic arcs (Elitez et al., 2016). Trench-parallel tear affects the subducting African lithosphere between northern Greece and the Gulf of Corinth along the Western Hellenic Arc (Hansen et al., 2019). Trench-perpendicular tear may accommodate the region between the Hellenic and Cyprian arcs, which differ in subduction steepness and material subducted (Dilek & Sandvol, 2009). The Cyprian arc involves shallower subduction dynamics with the Eratosthenes seamount and Anixamander Mountains (mud volcanoes; Lykousis et al., 2009) impinging on the trench (Kempler & Ben-Avraham 1987; Zitter et al., 2003; Biryol et al., 2011). This arc became effectively inactive during the onset of westward extrusion of the Anatolian plate (Papanikolaou, 2021).

2.2 Geometry of the Hellenic arc subduction zone

To understand when and why the Hellenic subduction zone was established, we must consider its present-day structure. The Hayes (2018) Slab2 model uses active-source seismic data interpretations, receiver functions, local and regional seismicity catalogs, and seismic tomography and models the subducting Nubian slab as uniformly northward dipping to >440 km depths in its northern portion (Figure 3). The Hellenic arc has a well-developed Wadati-Benioff zone at shallower depths but a debated slab geometry at intermediate depths (150–250 km, Suckale et al., 2009; Agostini et al., 2010; see review in Hansen et al., 2019). Figure 4A shows the slab clearly defined by earthquake depth vs. latitude across the Hellenic subduction zone. Detailed analysis of the distribution of earthquakes indicate that the western part of the subduction zone dips under 20–30° to the NE and reaches the maximum depth of 180 km, and its eastern section dips under 40° to the NW and reaches a maximum depth of 170 km (Papazachos & Comninakis, 1971; Vaněk et al., 1987; Papazachos et al., 2000; Suckale et al., 2009; Papazachos, 2019). At deeper levels (100–180 km), the Wadati–Benioff zone dips freely (without coupling) at a high angle (~45°) beneath the south Aegean trough and the volcanic arc (Mahatsente et al., 2017; Papazachos, 2019).

Seismic coupling is defined as the ratio between the observed seismic moment release and the rate calculated from plate tectonic velocities (e.g., Ruff and Kanamori, 1983; Scholz and Campos, 2012). The plate interface coupling between the Hellenic trench fault and the Nubia–Aegean is low (<20%) (Papadimitriou and Karakostas, 2008; Vernant et al., 2014) and only 10% in the southwestern portion of the arc (Papadimitriou and Karakostas, 2005). A low seismic coupling scenario tends to produce earthquakes with small rupture zones and high moment release that occur as small isolated patches on the subduction interface (Scholz 1990; Messina et al., 2007). However, in the region near the Ionian islands (Figure 1), complete seismic coupling of the subduction is most consistent with the observed moderate seismic moment release (Laigle et al., 2002; 2004). Higher degrees of seismic coupling are also found in specific regions in the Western Hellenic arc (Jackson and McKenzie, 1988; Messina et al., 2007; Ganas et al., 2020).

Classical ray tomography using body and surface waves is often applied to the Mediterranean region to image the Hellenic arc subduction system (e.g., Piromallo and Morelli, 1997; Spakman et al., 1988; Snieder, 1988; Zielhuis and Nolet, 1994; Piromallo and Morelli, 2003; Marone et al., 2004; Amaru, 2007; Schivardi and Morelli, 2009; Chang et al., 2010; Biryol et al., 2011; Salaün

et al., 2012; Legendre et al., 2012; Portner et al., 2018; Blom et al., 2019; Wei et al., 2019; El-Sharkawy et al., 2020). We present a P-wave tomographic image of the Aegean anomaly as an example in Figure 4B. Although surface wave data improves ray coverage and allows the visualization of regions with low seismicity and few stations, body wave tomography affords better resolution yet suffers from vertical smearing and an unknown background velocity model compared to surface waves (Piromallo & Morelli 1997; Kassaras et al., 2005; Schivardi & Morelli, 2009; Salaün et al., 2012). Body wave tomographic images across the Hellenic arc shows the geometry of the subduction zone clearly (e.g., Figure 4B), with the slab sometimes exhibiting a strong velocity contrast (up to 6%) compared to the background model (e.g., Papazachos & Nolet, 1997; Kassaras et al., 2005). Some regional studies show the slab forms an asymmetrical shape, shallowly dipping and larger along the western side and a steeper and smaller in the eastern section (Papazachos & Nolet, 1997; Blom et al., 2019).

Segments of the African slab have long been imaged as subducting into the lower mantle (Figure 4B) (1400 ± 100 km depth; Spakman et al., 1988; Spakman, 1990; 1991; Spakman et al., 1993; Papadopoulos, 1997; Bijwaard et al., 1998; Kárason & van der Hilst, 2000; Wortel & Spakman, 2000; Romanowicz, 2003; Kassaras et al., 2005; Chang et al., 2010; Biryol et al., 2011; Legendre et al., 2012; Govers & Fichtner, 2016; van der Meer et al., 2018; Bocchini et al., 2018; Wei et al., 2019; El-Sharkawy et al., 2020). The width of the subducting slab differs depending on the approach applied to image it. For example, Chang et al. (2010) show the slab widening beneath 660 km in the lower mantle, whereas Piromallo & Morelli (2003) has widening within the transition zone. Widening can be associated with a slab avalanche where high viscosity occurs in the lower mantle (Capitanio et al., 2009; Chang et al., 2010). Slab avalanches form when large-volume subducted slabs that temporarily stagnate within the transition zone periodically penetrate the lower mantle (e.g., Solheim and Peltier, 1994; Deschamps and Tackley, 2009; Yang et al., 2018). Slab avalanches are unstable, controlled by mantle thermal instabilities, and will accelerate as slab sinking rates increase with time (e.g., Solheim and Peltier, 1994; Yang et al., 2018). More recent tomographic images show the Hellenic slab extends from the surface to the transition zone in a bent, arcuate shape with a high-velocity structure that flattens from the 410 km discontinuity and is not seen at deeper levels (Blom et al., 2019).

Mantle tomography has also shown that not all slabs in the Mediterranean region are connected to the lithosphere at the surface due to delamination (e.g., Spakman et al., 1988; Dilek & Sandvol, 2009; Wortel & Spakman, 2000). For example, gaps are visible in the slab at depths of 60-100 km just west of the south Hellenides, and slab tear may be visible at the 660 km discontinuity (Wei et al., 2020). Multiple remnant slabs are imaged beneath the North Hellenides (e.g., Wei et al., 2020) and most of Anatolia (e.g., Salaün et al., 2012). Vote Maps, generated from stacking tomography models and identifying where the models agree based on an increasing vote count at a specified depth (Shephard et al. 2017;), show a slab interpreted to be the African plate between 250-1550 km depths (Cramer et al., 2020).

Although tomographic images are central in some studies of Hellenic arc SZI, their results are challenging to interpret, especially in locating regions of slab tear (e.g., Piromallo & Morelli 2003). The subduction zone's small size, its spatially highly variable nature, and the uneven distribution of its seismic stations make imaging the area more challenging (El-Sharkawy et al., 2020). The presence of slow velocity anomalies throughout the uppermost mantle beneath Anatolia and very slow velocities beneath eastern Anatolia have been explained by innovative

ideas regarding how the mantle and lithosphere behave in terms of mantle upwelling, drip-style lithospheric delamination (Göğüş et al., 2017), or rollback (see discussion in Portner et al., 2018). Tomographic images of both the Hellenic and Cyprus arc have transformed ideas regarding how subducting slabs operate as they enter the deep mantle and their impacts on seismicity and the creation of transfer zones (e.g., Meighan et al., 2013), large volume magmatism (e.g., Cocchi et al., 2017), and controlling ore-forming process and mineral deposits (e.g., de Boorder et al., 1998; Rabayrol et al., 2019; Rabayrol & Hart, 2021). Changes in the subducting slab dynamics significantly affect plate dynamics, including plate motion and mantle dynamics (e.g., Gianni et al., 2019).

3 Geological background of Aegean-Anatolian Suture Zones

Determining the geological relationship between the present-day Hellenic arc and ancient subduction zones associated with the Aegean and Anatolian microplates is complicated due to the amalgamation of terranes that occurred over its tectonic history. The western portion of the Anatolian microplate exposes ophiolite remnants and other evidence of ancient subduction zones that may have influenced the onset of the present-day Hellenic arc. The Anatolian microplate's development is characterized by continental fragments separated by branches of the Paleo- and Neo-Tethyan oceans that collided and ultimately combined by the Late Cretaceous-Eocene. These exposures of ophiolitic and high-pressure/low-temperature (HP/LT) rock assemblages identify suture zones (Figure 5) (e.g., Şengör & Yılmaz, 1981; Okay, 2008; Moix et al., 2008; Okay & Tuysuz, 1999; Pourteau et al., 2016; Okay et al., 2020). This section describes the evidence of ancient subduction zones that may be linked to or influenced the development of the Hellenic arc.

3.1 Intra-Pontide Suture Zone

Western Anatolia contains the Pontide Mountains to the north and the Taurides Mountains to the south (Figure 5) (e.g., Şengör & Yılmaz, 1981; Yılmaz et al., 1997; Okay & Tuysuz, 1999; Pourteau et al., 2016). The Pontide Mountains include a southern Eurasia margin affinity and are characterized by a Pan-African basement with Phanerozoic sedimentary cover units (Yılmaz et al., 1997; Moix et al., 2008; Pourteau et al., 2010; Okay et al., 2013). The Intra-Pontide suture zone (IPS) is mapped within the Pontide zone between the Sakarya continental zones and Istanbul-Zonguldak Unit (Istanbul–Zonguldak Zone, Istanbul Nappe, or Istanbul Zone, Yiğitbaş et al., 2004; Yılmaz et al., 2021). The IPS has been interpreted as an accretionary complex, a suprasubduction zone, and a remnant of a former ocean basin that may have extended into eastern Europe (e.g., Okay et al., 1996; Robertson & Ustaömer, 2004; Göncüoğlu et al., 2012; 2014; Marroni et al., 2014; Akbayram et al., 2016; Sayit et al., 2016; Frassi et al., 2018). Its complex geology, comprised of components from the Istanbul-Zonguldak and/or Sakarya zones, has led to a debate about its presence and utility in paleogeographic reconstructions (Moix et al., 2008). Geochemical data from mafic rocks collected along the IPS show signatures related to a supra-subduction zone that opened in the Middle Jurassic-Early Cretaceous, obducted during the Early-Late Cretaceous boundary, and had closed by the Late Paleocene (Göncüoğlu et al., 2014; Sayit et al., 2016).

In Greece, the Vardar suture connects to the IPS (Şengör & Yılmaz, 1981; Okay & Satir, 2000; Okay et al., 2001; Beccalotto & Jenny, 2004; Okay et al., 2010; d'Atri et al., 2012; Di Rosa et al., 2019) or Meliata-Balkan suture (Stampfli, 2000). In Turkey's Biga Peninsula, the IPS and Vardar connection may be recorded by an isolated ophiolite-bearing accretionary complex active

until the Late Cretaceous (Figure 5) (e.g., Okay et al., 1991). However, this connection is debated by several researchers (Altunkaynak & Genc, 2008; Burchfiel et al., 2008; Şengün et al., 2011). Furthermore, due to the uncertain link between IPS and Izmir-Ankara-Erzincan Suture Zone (IAESZ), the relationship of the Biga Peninsula basement assemblages to those in the Rhodope-Thrace Massif is also debated (e.g., Bonev & Beccaletto, 2007; Elmas, 2012). Both the IPS and IAESZ are thought to mark Late Cretaceous–earliest Tertiary closure of Neo-Tethyan ocean basins (e.g., Pourteau et al., 2010; Akbayram et al., 2016).

3.2 Izmir-Ankara-Erzincan Suture Zone (IAESZ)

The IAESZ separates the Pontide's Sakarya Composite Terrane in the north from the Anatolide-Tauride block to the south (Figure 5) (Şengör & Yılmaz, 1981; Okay & Tüysüz, 1999; Tekin et al., 2002; Göncüoğlu, 2010; Yılmaz et al., 2021). In the Aegean microplate, the IAESZ is thought to record the closure of the Vardar ocean and link with the Vardar ophiolite (or Axios-Vardar suture zone) (Figure 1) (Channell & Kozur, 1997; Okay & Tuysuz, 1999; Tekin et al., 2002; Moix et al., 2008). However, the feature is not exposed due to the presence of the Aegean Sea (e.g., Burtman, 1994; Stampfli, 2000; Yılmaz et al., 2001; Burchfiel et al., 2008).

In Western Anatolia, blueschist assemblages exposed along the IAESZ are intruded by Suture Zone Granitoids (SZGs) [Topuk, Orhaneli, Tepedag (Gürgenyayla and Gürgenyayla)] with Paleocene (63.5 ± 2.8 Ma) to Oligocene (31.4 ± 0.6 Ma) ages but primarily characterized by early Eocene emplacement (~ 45 – 47 Ma, Okay & Satir, 2006; Altunkaynak, 2007). The SZGs intrude the western portion of the Tavşanlı Zone, a blueschist sequence overlain by a Cretaceous accretionary complex and ophiolitic sheet. The Tavşanlı Zone is narrow (~ 50 km) and trends E–W for approximately 250–350 km (Okay & Whitney, 2010; Plunder et al., 2013). Its western and central portions contain blueschist facies metavolcanic and metasedimentary rocks with rare metabasalts (Okay, 1980a, 1980b, 1982; Okay & Kelley, 1994, see Seaton et al., 2009). The zone is thought to have formed as a result of northward-dipping subduction and represents the Mesozoic–Eocene closing of the northern branch of the Neo-Tethyan Ocean (Okay, 1986; 2008; Okay & Kelley, 1994; Sherlock et al., 1999; Moix et al., 2008; Shin et al., 2013; Plunder et al., 2013; Fornash & Whitney, 2020). The Sivrihisar Massif in the eastern portion of the Tavşanlı has eclogite, blueschist, and Barrovian sequences (Gautier, 1984; Seaton et al., 2009). Paleocene–Eocene ages from the Tavşanlı Zone granites mark the timing of the closure of the IAESZ (e.g., Okay et al., 2020).

The Afyon zone (Afyon–Bolkardag Zone of Okay, 1986; Özdamar et al., 2013 or Ören–Afyon Zone of Pourteau et al., 2013) is considered the southward palaeogeographic extension of the Tavşanlı zone (Candan et al., 2005; Pourteau et al., 2010; Akal, 2013; Özdamar et al., 2013). Although its southern extent is unclear, the Afyon zone is mapped parallel to the Tavşanlı Zone. A portion may also be exposed between the southern Menderes Massif and Lycian Nappes (Okay, 1986; Candan et al., 2005; Pourteau et al., 2013; Ustaömer et al., 2020). Afyon Zone stratigraphy resembles the Tavşanlı Zone and consists of Palaeozoic–Mesozoic Tauride shelf sequences of metasedimentary and metavolcanic rocks that experienced regional greenschist to blueschist facies (Fe–Mg carpholite and glaucophane) metamorphism overlying Pan-African-related basement (Okay, 1984; Candan et al., 2005; Pourteau et al., 2010; Özdamar et al., 2013). HP/LT metamorphism in the Afyon Zone is thought to have occurred coincident with the closure of the Neo-Tethyan Ocean at 70–65 Ma (Pourteau et al., 2010, 2013; Özdamar et al., 2013; Plunder et al., 2013). Portions of the Afyon Zone may have subducted beneath the Tavşanlı Zone

during the Late Cretaceous based on zircon ages of granites that intrude Tavşanlı Zone blueschist and altered ophiolitic assemblages (Speciale et al., 2012; Shin et al., 2013).

4. Age constraints on the initiation of subduction

4.1 Cenozoic estimates

Early estimates for the initiation age of Hellenic arc subduction were Late Miocene to Pliocene (13 ± 3 -5 Ma, Le Pichon & Angelier, 1979 and 5-10 Ma, McKenzie, 1978; Mercier, 1981) based on interpretations of seismic activity coupled with assumptions regarding the age of subducted lithosphere and subduction depths. These ages are similar to the earliest volcanic activity in the South Aegean arc (Pliocene, Pe-Piper & Piper, 2005) and the onset of the KTZ based on geodynamic modeling and GPS data (6-8 Ma, Royden & Papanikolaou, 2011). The Late Miocene time frame is also cited as when the Western Hellenic arc divided into its northern and southern portions along the KTZ (Papanikolaou, 2010). Reconstructions of fault systems in the northern margin of the eastern Mediterranean Sea are also consistent with 15 Ma (Le Pichon et al., 2019).

Interpretations of the Aegean seismic velocity structure, tomography, and seismicity data in the Aegean area suggest Eocene to Late Oligocene onset of the Hellenic arc subduction zone (40-26 Ma; Meulenkamp et al., 1988; Spakman et al., 1988; Papadopoulos, 1997; Brun & Sokoutis, 2010). These ages are consistent with the timing of granite crystallization throughout Western Anatolia and Mediterranean Ridge sedimentation (33-23.6 Ma, Fytikas et al., 1984; Kastens, 1991). Younger estimates from the ridge are also reported (~19 Ma, Kopf et al., 2003). Plate reconstructions suggest that the Northern Hellenic trench from the Western portion of the Hellenic arc experienced the onset of subduction from 34-27 Ma. In contrast, the Southern Hellenic arc was active at 34 Ma (Royden & Papanikolaou, 2011).

The progressive deceleration in motion of Africa with respect to Europe in the Mediterranean region has been observed to occur since 35 Ma, and in the eastern Mediterranean from 35 Ma to 10 Ma to a convergence rate of a few mm/yr (Savostin et al., 1986; Marsellos et al., 2010). The rate of trench retreat may have accelerated from ~0.6 cm/yr during the first 30 m.y. of subduction to 3.2 cm/yr during the past 15 m.y., perhaps due to slab tear during the Middle Miocene-Pliocene (Brun et al., 2017). Timing constraints on Aegean forearc curvature, due to opposite rotations, clockwise in the west and counterclockwise in the east, are Eocene and Middle Miocene (Morris & Robertson 1993; Cornée et al., 2018). Trench bending and rollback increased obliquity of subduction over time, which was accommodated by strain partitioning within the upper Eurasian plate (Faccenna et al., 2013; Philippon et al., 2014; Brun et al., 2016; Cornée et al., 2018). By the Early Oligocene-Eocene, back-arc basin formation was ongoing in the Aegean region as the African plate retreated as it subducted beneath Eurasia (Le Pichon & Angelier, 1981; Royden, 1993; Jolivet & Faccenna, 2000; Agostini et al., 2010; Carminati et al., 2012; Jolivet et al., 2018).

Figures 6 and 7 are N-S temporal cross-sections across Western Anatolia that describe the region's tectonic evolution from the Late Cretaceous to the present. Figure 6A (from Dilek & Altunkaynak, 2009) illustrates the closure of a branch of the Neo-Tethyan Ocean (Izmir-Ankara Ocean), which occurred during the Late Cretaceous when the IPS already marked the division between the Sakarya and Rhodope continents. Break off of the subducting Neo-Tethyan slab occurs ca. 54 Ma along the IAESZ (Figure 6B), concurrent with the development and emplacement of SZG and ophiolite obduction. The Hellenic arc forms due to subduction step-

back processes that are completed by the Miocene (~25 Ma) (Figure 6C). Subduction rollback was most likely initiated soon after the complete formation of the Hellenic Arc to start Aegean back-arc extension (Figure 6D). Some variations of this model exist. For example, Shin et al. (2013) proposed an additional subduction system concurrent with the IAESZ but located further south between the IAESZ and the Hellenic arc (Figure 7).

These models are the outcome of large data sets comprised of primarily field observations, geophysical, geochemical, petrological, geochronological, and structural data from multiple rock types, but primarily those affiliated with the upper mantle and continental crust (lithosphere). They provide testable hypotheses for the timing and generation mechanisms of particular magmatic and structural features in the Aegean and western Anatolia to predict the presence of multiple suture zones and subducted slabs. They also delineate the affinity of specific crustal fragments north of the Hellenic arc to be distinct from the African plate in terms of their origin and geological history.

4.2 Mesozoic estimates

Significantly older estimates for the onset of the Hellenic subduction exist and suggest the arc has been long-lived since the Jurassic-Late Cretaceous, with stipulations that Cenozoic age estimates are considering only the latest episode of its evolutionary history and disregard its earlier history (Figure 8) (e.g., Jolivet et al., 2018). In this model, the African slab has been part of a single, evolving subduction zone system that was initiated sometime in the Mesozoic during the initiation of the closure of the northern branch of the Neo-Tethys ocean (Faccenna et al., 2003; van Hinsbergen et al., 2005; Royden & Papanikolaou, 2011; Jolivet et al., 2013; Malandri et al., 2017; Jolivet et al., 2018; Cramer et al., 2020; van Hinsbergen et al., 2021). The model has been primarily applied to explain the evolution of the Western Hellenic arc. The Vardar suture in Greece, equivalent to the IAESZ (Channell & Kozur, 1997; Okay & Tuysuz, 1999; Moix et al., 2008), and Pindos suture zone, equivalent to units within the Antalya domain and Dilek peninsula (Stampfli & Kozur, 2006) had buoyant microcontinents that entered and locked subduction. This process triggered southward slab rollback and migration of the volcanic arc (Figure 8) (van Hinsbergen et al., 2005; Brun & Faccenna 2008; Jolivet & Brun 2010; Jolivet et al., 2013; Cornée et al., 2018). In this model, the subduction zone accumulates tectonic fragments, terrains, and oceanic lithosphere, instead of following the induced transference model of subduction (Figures 6 and 7; e.g., Stern, 2004). In this model, a suture is created, and a new subduction zone develops. Instead, the Hellenic subduction zone front has behaved like an accretionary wedge during its entire development since the Jurassic-Late Cretaceous.

The model also requires a stationary trench from 100-45 Ma to allow for the penetration of the African slab into mantle depths (Bonneau, 1982; Stampfli & Borel, 2002; Jolivet et al., 2003; Capitanio et al., 2010). Penetration to 1200-1400 km depths is estimated to have occurred by ~50 Ma, which triggers the onset of extension in the region by 45 Ma (Jolivet & Faccenna, 2000; Brun & Faccenna, 2008; Capitanio et al., 2009; 2010). The present-day curvature of the Hellenic forearc represents oblique subduction that grew systematically (Huchon et al., 1982; Le Pichon et al., 1995; ten Veen & Kleinspehn, 2003; Gautier et al., 1999; Le Pichon et al., 2002; Wallace et al., 2005, 2008; van Hinsbergen & Schmid, 2012; Philippon et al., 2014; Cornée et al., 2018).

The single subduction model was developed primarily using interpretations of mantle tomography coupled with estimates of subduction rates over long periods of geological time. Thus, ages found in suture zones that would be entirely unrelated to the Hellenic arc in early SZI

models now time the onset of subduction of the African slab. For example, the Late Cretaceous (104 Ma) estimate of Hellenic arc SZI is based on the ages of ophiolites exposed in the Tavsanli Zone of Western Anatolia, part of the IAESZ (Peters et al., 2017; Pourteau et al., 2019; Crameri et al., 2020). The single subduction system also has implications for the affinity of Aegean crustal fragments and terranes in paleogeographical reconstructions. The model requires all the lower plate continental crust to be accreted into the upper plate while subducting continental lithosphere. It also requires the entire Aegean crust from the Vardar (Izmir-Ankara) suture to the Mediterranean ridge to be derived from the lower plate (Figure 6-8). Oceans between the accreted domains were of significant size (500 km in some cases). The process would lead to significant elevation changes and crustal thicknesses that should be recorded in the geological and tomographic record. Critical changes in the zone of subduction transitions also occurred as shear zones developed to accommodate oceanic and continental lithosphere (see discussion in Le Pichon et al., 2019). The model also eliminates the need for multiple sutures and subducted slabs to be present beneath western Turkey and the Aegean and simplifies the evolution of the Aegean microplate to a single evolving, long-lived subduction system (van Hinsbergen et al., 2005).

5 Discussion

Plate interactions are thought to be most clear at convergent plate boundaries (Silver et al., 1998). However, several outstanding issues remain regarding timing the onset of convergence between the Nubian Plate and Aegean microplate. Several observations indicate that the Aegean microplate does not behave like a typical back-arc basin (Agostini et al., 2010; Doglioni et al., 2002). It lacks an ocean floor and associated ocean crust and is underlain by a thick layer of continental crust (e.g., Makris, 1978). In fact, the Aegean crust is basically comprised of stacked carbonate and oceanic nappes, metamorphic massifs, and core complexes (Jolivet et al. 2013; Papazachos, 2019). Models of deformation along the Hellenic arc intended to capture the large-scale 3-D structure of the Nubian plate suggest the region more closely resembles a continental thrust than a typical oceanic subduction zone (Ganas and Parsons, 2009).

The Aegean microplate is also bound to the north by the active strike-slip systems (NAF, NASZ, KTZ, and Central Hellenic Shear Zone; Brooks & Ferentinos, 1984; Lyberis, 1984; Nyst & Thatcher, 2004; Gürer et al., 2006; Kreemer et al., 2004; Kokkalas et al., 2006; Papanikolaou & Royden, 2007; Reilinger et al., 2010; Royden & Papanikolaou, 2011; Vassilakis et al., 2011; Halpaap et al., 2018). This results in a complex tensional regime where crustal stretching is inconsistent with the geometry and direction of the subducting Hellenic slab (e.g., Mantovani et al., 1997; Agostini et al., 2010). Differences in the timing of initiation and rate of subduction exist between segments along the Western Hellenic Arc and should also be expected to occur along other portions of the Hellenic and Cyprus arcs (Royden & Papanikolaou, 2011). Subduction zones behave chaotically and may retreat advance or remain stationary at different stages, especially if the incoming lithosphere is heterogeneous (e.g., Royden & Husson 2009; Husson et al., 2009). The extent and role of several ancient fault systems, including older Neo-Tethyan suture zones that likely played a significant role in its development, are thus debated.

The Mesozoic SZI age for the Hellenic arc requires all of the lower plate continental crust to be accreted into the upper plate while subducting continental lithosphere. The entire Aegean Crust from the Vardar suture to the Mediterranean ridge would be derived from the lower plate (Figure 8). This process requires a redefinition of components associated with ancient supercontinents and would impact paleogeographic reconstructions of the region. Oceans between the accreted domains were of significant size (500 km in some cases). The process

would lead not only to changes in elevation and crustal thicknesses, but also critical changes in the zone of subduction transitions occurring from oceanic to continental shear zones (see discussion in Le Pichon et al., 2019) (see discussion in Le Pichon et al., 2019). Not all units record blueschist facies conditions, and some experienced Barrovian prograde (burial) P-T paths, such as on the island of Naxos (e.g., Lamont et al., 2019).

Subduction zones with limited trench-parallel lengths on the order of the Hellenic arc (600-800 km) and narrow slabs (<1,500 km) typically have rapid retreat rates (Schellart et al., 2007; Bolhar et al., 2010). The requirement of a stationary trench from 100-45 Ma to allow for the African slab to enter mantle depths (Bonneau, 1982; Stampfli & Borel, 2002; Jolivet et al., 2003; Capitanio et al., 2010) seems unusual in that most of the slab today is characterized by significant slab tear. Sites of slab tear vary in scale from regional to local and include the boundary between the Hellenic and Cyprus arcs (Wortel & Spakman, 1992; Biryol et al., 2011), south of Crete at the Pliny–Strabo Shear Zone (Özbakır et al., 2013), the İzmir-Balıkesir Transfer Zone (e.g., Kaya, 1981; Gessner et al., 2013; Cemen et al., 2014; Ersoy et al., 2014), and beneath the Menderes Massif (Biryol et al., 2011; Rabayrol & Hart, 2021), and the eastern Aegean (Jolivet et al., 2015).

At present, it is unclear how the subducting African (Nubian) slab could survive through the Earth's transition zone, suggesting that its composition and convergent rate, and the material properties of the mantle were critical constraints to allow this process to occur (e.g., Káráson & van der Hilst, 2000). Interpretations of the deeper images are debated: the subducted slab may be a single folded body that overturned in the lower mantle (Faccenna et al., 2003) or two different slabs between 2000-1500 km and from 1500 km to the surface (van Hinsbergen et al., 2005; van der Meer et al., 2018). The unhindered slab penetration into lower mantle depths is unexpected due to the relationship between slab morphology and trench migration (e.g., Widiyantoro et al., 2004). However, the process of slabs reaching depths of 1700 km is not uncommon (e.g., Káráson & van der Hilst, 2000).

Numerous examples also suggest significant slab tear across the Aegean and Western Anatolia (e.g., Wortel & Spakman, 1992; Woodside et al., 2000; Biryol et al., 2011; Gessner et al., 2013; Faccenna et al., 2014; Roche et al., 2019; Rabayrol & Hart, 2021). Legrande et al. (2012) note that body wave studies show slab segments are trapped within the mantle transition zone in the western Mediterranean and Alpine region and do not dive deeper as is observed in the Aegean region. Overall, the Mediterranean Sea exhibits lateral variations in crustal and upper mantle structures (Marone et al., 2004a).

Long-lived Aegean subduction would have affected the mantle's thermal and chemical convection and impacted surface conditions, including climate over time (e.g., Sigl et al., 2005). The region contains active and quiescent volcanoes located in the Methana Peninsula (Peloponnese) and islands of the Cycladic archipelago (Figure 1) (see review in Scoon, 2021). Holocene volcanism along the Hellenic arc allows for identifying and characterizing high-frequency climatic changes during the Holocene in particular (e.g., Zanchetta et al., 2011). Fluids released from past and present subducting lithosphere have affected not only volcanism, but also grain mobility and seismic anisotropy throughout the Mediterranean and elsewhere (e.g., Karato, 1995; Van der Meijde et al., 2003; Marone et al., 2004b; Faccenda et al., 2008). This impact over the length of planetary history may have been significant in terms of triggering other subduction zones, controlling sites of seismicity and magmatism, as well as the generation of ore deposits. A Jurassic-age subduction zone that is presently active and capable of generating significant

hazards even today has important implications for our understanding of lithospheric strength, the driving force of plate tectonics over geological time, and the tectonic history of the Aegean region.

The single subduction system has its roots in tomography and the interpretation of large low shear-wave velocity anomalies in the lowermost mantle. These compositional anomalies thus appear as crucial as continents are for the surface, yet they are inaccessible and have limited data regarding their thermal histories and chemical compositions. Their role as reservoirs and impact on the evolution of the Hellenic arc over time is unclear. The interpretation that the seismic tomography images reveal a single, long-lived subduction system may be incorrect. The Aegean-Anatolian microplates have long been known to have an extensive and complex orogenic history based on surface geology. With the presence and data from mapped suture zones, it is expected that the region would record multiple subducted slabs from earlier collisional events.

Determining the Hellenic arc SZI is also important because the timing constraints shed light on the mode of initiation (i.e., induced or spontaneous; from compression or extension; Gurnis et al., 2004; Stern, 2004; Gerya, 2011; Stern & Gerya, 2018) based on our understanding of the geology at that time. Subduction initiation requires breaking a zone of weakness in the lithosphere (e.g., Stern & Gerya, 2018; Baes et al., 2020a; Cloetingh et al., 2021). The process may be accommodated along passive margins (e.g., Cloetingh et al., 1984), although the conversion of a passive margin into an active one has not been observed to have occurred during the Cenozoic (Gurnis et al., 2004; Cloetingh et al., 2021). Subduction initiation could also occur due to motion along transform or fracture zones associated with mid-ocean ridges due to large-scale changes in plate motion (e.g., Hall et al., 2003). This process could occur across extinct mid-ocean ridges or within back-arc regions within mature subduction zones, as well as mantle plume interactions with old oceanic lithosphere (Baes et al., 2020a,b; Cloetingh et al., 2021). No evidence or hypothesis exists for a mantle plume as triggering the development of the Hellenic arc. A young SZI initiation age suggests the Hellenic arc developed due to the Late Cenozoic collision of the African plate with Eurasia (e.g., Meulenkaamp et al., 1988). An older SZI links its origination to near and parallel to a spreading ridge associated with the western portion of the NeoTethyan ocean (e.g., Maffione and van Hinsbergen, 2018).

Younger estimates for the Hellenic arc are consistent with the observation that nearly half of Earth's active subduction zones were initiated during the Cenozoic (Gurnis et al., 2004). Crameri et al. (2020) indicate that SZI has happened at least thirteen times during the last hundred million years, with global occurrences of SZI occurring within 55-40 Ma and 16-6 Ma. These times mark higher SZI activity on Earth and are remarkably consistent with Cenozoic estimates of the onset of the Hellenic arc (McKenzie, 1978; Le Pichon & Angelier, 1979; Mercier, 1981; Meulenkaamp et al., 1988; Spakman et al., 1988; Papadopoulos, 1997; Brun & Sokoutis, 2010; Royden & Papanikolaou, 2011; Le Pichon et al., 2019). A significant slowdown between Africa's northward motion relative to surrounding plates has also been attributed to the Eocene-Oligocene collision of Africa and Eurasia, which altered the kinematics and tectonic evolution of not only the Aegean region but the restructuring, reorganization, and fracturing of the African plate elsewhere (e.g., DeMets & Merkouriev, 2016).

6 Conclusions

The Hellenic arc results from the convergence of the northern margin of the African (Nubian) plate with the southern portion of the Aegean microplate. The region has emerged as the type

locality for subduction zone tectonics, and many novel ideas regarding the development of this particular plate boundary have emerged from the area. Two end-member models for the SZI age of the Hellenic arc evolved using disparate datasets. Seismic tomography images help determine the present-day subduction-related structure beneath the Aegean microplate, and some interpretations suggest a long-lived subduction zone since the Jurassic to Late Cretaceous. However, tomographic images can be limited in their interpretations of deeper depths and our ability to observe slab tears or structures that disrupt the subducting slab. Surface geological data sets suggest a complex, long-lived history of the Aegean region with multiple suture zones. These observations are more consistent with multiple detached slabs, even some unrelated to the African plate. Geological, geochemical, and geochronological constraints from the continental lithosphere affected by subduction help constrain the pathways of rocks affected by subduction dynamics. However, these are limited by the currently available datasets and interpretation within an existing framework with limited exposure. The Hellenic arc is a significant plate boundary that would likely be influenced by differential retreat, inherited lower plate lithospheric heterogeneities, and mantle upwelling. Multidisciplinary studies that couple detailed field mapping of available outcrops in the Western Anatolian and Aegean microplates with geophysical observations, geochemistry, and geochronology are needed to determine its SZI age and which model, a long-lived single subduction zone or multiple subduction zones with the latest SZI in Cenozoic, applies to the region.

Acknowledgments

No real or perceived financial conflicts of interests exist for the authors. We appreciate the comments from two anonymous reviewers. We thank Jeffrey S. Horowitz for assistance with drafting figures.

References

- Agostini, S., Doglioni, C., Innocenti, F., Manetti, P., & Tonarini, S. (2010). On the geodynamics of the Aegean Rift. In I. Çemen (Ed.), *Extensional tectonics in the Basin and Range, the Aegean, and western Anatolia*. *Tectonophysics*, 488, 7-21. <https://doi.org/10.1016/j.tecto.2009.07.025>
- Akal, C. (2013). Coeval shoshonitic-ultrapotassic dyke emplacements within the Kestanol Pluton, Ezine-Biga Peninsula (NW Anatolia). *Turkish Journal of Earth Sciences*, 21, 1–20.
- Akbayram, K., Şengör, A. M. C., & Özcan, E. (2016). The evolution of the intra-Pontide suture; implications of the discovery of late Cretaceous-early Tertiary melanges. In R. Sorkhabi (Ed.), *Tectonic evolution, collision, and seismicity of southwest Asia; in honor of Manuel Berberian's forty-five years of research contributions*. Special Paper - Geological Society of America, 525. [https://doi.org/10.1130/2016.2525\(18\)](https://doi.org/10.1130/2016.2525(18))
- Alatza, S., Papoutsis, I., Paradissis, D., Kontoes, C., & Papadopoulos, G.A. (2020). Multi-Temporal InSAR analysis for monitoring ground deformation in Amorgos Island, Greece. *Sensors*, 20(2), 338. <https://doi.org/10.3390/s20020338>
- Altunkaynak, Ş. (2007). Collision-driven slab breakoff magmatism in northwestern Anatolia, Turkey. *Journal of Geology*, 115, 63–82
- Altunkaynak, Ş., & Genc, S. (2008). Petrogenesis and time-progressive evolution of the Cenozoic continental volcanism in the Biga Peninsula, NW Anatolia (Turkey). In: Xu,

- Y., Farmer, L., Menzies, M., Rudnick, R., Zhou Meifu, Z. Continental volcanism and chemistry of the Earth's interior. *Lithos*, 102, 316-340.
- Amaru, M.L. (2007). Global travel time tomography with 3-D reference models. PhD Thesis Utrecht University.
<http://dspace.library.uu.nl/bitstream/handle/1874/19338/index.htm;jsessionid=88B6AA4941C5E76FD65034D89D4065E0?sequence=17>
- Anastasakis, G., & Keling, G. (1991). Tectonic connection of the Hellenic & Cyprus arcs & related geotectonic elements. *Marine Geology*, 97(3-4), 261-277.
[https://doi.org/10.1016/0025-3227\(91\).90120-S](https://doi.org/10.1016/0025-3227(91).90120-S)
- Andronikidis, N., Kokinou, E., Vafidis, A., Kamberis, E., & Manoutsoglou, E. (2017). Deformation patterns in the southwestern part of the Mediterranean Ridge (South Matapan Trench, Western Greece). *Marine Geophysical Research*.
<https://doi.org/10.1007/s11001-017-9337-0>
- Angelier, J., Lyb  ris, N., Le Pichon, X., Barrier, E., & Huchon, P. (1982). The tectonic development of the hellenic arc & the sea of crete: A synthesis. *Tectonophysics*, 86(1-3), 159-196. [https://doi.org/10.1016/0040-1951\(82\).90066-X](https://doi.org/10.1016/0040-1951(82).90066-X)
- Armijo, R., Meyer, B., Hubert, A., & Barka, A. (1999). Westward propagation of the North Anatolian fault into the northern Aegean: Timing and kinematics. *Geology*, 27, 267-270.
[https://doi.org/10.1130/0091-7613\(1999\)027<0267:WPOTNA>2.3.CO;2](https://doi.org/10.1130/0091-7613(1999)027<0267:WPOTNA>2.3.CO;2)
- Baes, M., Sobolev, S. V., Gerya, T., & Brune, S. (2020a). Subduction initiation by plume-plateau interaction: Insights from numerical models. *Geochemistry, Geophysics, Geosystems*, 21, e2020GC009119. <https://doi.org/10.1029/2020GC009119>
- Baes, M., Sobolev, S., Gerya, T., & Brune, S. (2020b). Plume-induced subduction initiation: Single-slab or multi-slab subduction?. *Geochemistry, Geophysics, Geosystems*, 21, e2019GC008663. <https://doi.org/10.1029/2019GC008663>
- Barbot, S., & Weiss, J.R. (2021). Connecting subduction, extension and shear localization across the Aegean Sea and Anatolia. *Geophysical Journal International*, 226(1), 422–445.
<https://doi.org/10.1093/gji/ggab078>
- Barka, A.A. (1992). The north Anatolian fault zone. *Annales Tectonicae*, 6, 64-195
- Basili, R., Tiberti, M. M., Kastelic, V., Romano, F., Piatanesi, A., Selva, J., & Lorito, S. (2013). Integrating geologic fault data into tsunami hazard studies, *Natural Hazards Earth Systems Science*, 13, 1025–1050. <https://doi.org/10.5194/nhess-13-1025-2013>
- Beccaleto, L., & Jenny, C. (2004). Geology and Correlation of the Ezine Zone: A Rhodope Fragment in NW Turkey? *Turkish Journal of Earth Sciences*, 13, 45-176.
- Be   er, M., Wyss, M., & R. Kind, R. (1990). Inversion of source parameters for subcrustal earthquakes in the Hellenic Arc. *Geophysics Journal International*, 103, 439-450.
- Benetatos, C., Kiratzi, A., Papazachos, C., & Karakaisis, G. (2004). Focal mechanisms of shallow and intermediate depth earthquakes along the Hellenic Arc. *Journal of Geodynamics*, 37(2), 253-296. <https://doi.org/10.1016/j.jog.2004.02.002>
- Bijwaard, H., W. Spakman, & E. R. Engdahl (1998). Closing the gap between regional and global travel time tomography, *Journal of Geophysical Research*, 103, 30,055 – 30,078.
- Biryol, C. (2009). Complex rupture processes of the Solomon Islands subduction zone earthquake and subduction controlled upper mantle structure beneath Anatolia. PhD Thesis, University of Arizona, 150p.

- Biryol, C.B., Beck, S.L., Zandt, G., & Ozacar, A.A. (2011). Segmented African lithosphere beneath the Anatolian region inferred from teleseismic P-wave tomography. *Geophysics Journal International*, 184, 1037–1057. <https://doi.org/0.1111/j.1365-246X.2010.04910.x>
- Bleier, N., Bruhns, O., Friederich, W., Stoeckhert, B. (2007). Time variation of the stress field in the Aegean subduction zone. In Friederich, W. (Ed.), *Subduction dynamics, bridging the scales*. Bochumer Geowissenschaftliche Arbeiten, 10, 98.
- Blom, N., Gokhberg, A., Fichtner, A. (2019). Seismic waveform tomography of the central and eastern Mediterranean upper mantle. *Solid Earth*, 11, 669–690. <https://doi.org/10.5194/se-11-669-2020>.
- Bocchini, G.M., Brüstle, A., Becker, D., Meier, T., van Keken, P.E., Ruscic, M., Papadopoulos, G.A., Rische, M., & Friederich, W. (2018). Tearing, segmentation, and backstepping of subduction in the Aegean: New insights from seismicity. *Tectonophysics*, 734–735, 96–118. <https://doi.org/10.1016/j.tecto.2018.04.002>.
- Bocchini, G.M., Novikova, T., Papadopoulos, G.A. et al. (2020). Tsunami potential of moderate earthquakes: The July 1, 2009 Earthquake (Mw 6.45) and its associated local tsunami in the Hellenic Arc. *Pure Applied Geophysics*, 177, 1315–1333. <https://doi.org/10.1007/s00024-019-02246-9>
- Bohnhoff, M., Makris, J., Papanikolaou, D., & Stavrakakis, G. (2001). Crustal investigation of the Hellenic subduction zone using wide aperture seismic data. *Tectonophysics*, 343(3–4), 239–262. [https://doi.org/10.1016/S0040-1951\(01\)00264-5](https://doi.org/10.1016/S0040-1951(01)00264-5)
- Bolhar, R., Ring, U., & Allen, C.M. (2010). An integrated zircon geochronological and geochemical investigation into the Miocene plutonic evolution of the Cyclades, Aegean Sea, Greece: Part 1: Geochronology. *Contributions to Mineralogy and Petrology*, 160, 719–742. <https://doi.org/10.1007/s00410-010-0504-4>
- Bonev, N., & Beccaletto, L. (2007). From syn- to post-orogenic Tertiary extension in the north Aegean region: constraints on the kinematics in the eastern Rhodope–Thrace, Bulgaria–Greece and the Biga Peninsula, NW Turkey. *Geological Society, London, Special Publications*, 291, 113–142. <https://doi.org/10.1144/SP291.6>
- Bonneau, M. (1982). Tectonic evolution of central Aegean Hellenides from Upper Jurassic to Miocene times. *Bulletin Society Geology France*, 24, 229–242.
- Bosworth, W., Huchon, P., & McClay, K. (2006). The Red Sea and Gulf of Aden Basins. *Journal of African Earth Sciences*, 43(1–3), 334–378. <https://doi.org/10.1016/j.jafrearsci.2005.07.020>
- Brooks, M., & Ferentinos, G. (1984). Tectonics and sedimentation in the Gulf of Corinth and the Zakynthos and Kefallinia channels, Western Greece. *Tectonophysics*, 101(1–2), 25–54. [https://doi.org/10.1016/0040-1951\(84\)90040-4](https://doi.org/10.1016/0040-1951(84)90040-4)
- Brun, J.-P., & Faccenna, C. (2008). Exhumation of high-pressure rocks driven by slab rollback, *Earth and Planetary Science Letters*, 272, 1–7. <https://doi.org/10.1016/j.epsl.2008.02.038>
- Brun, J.-P., & Sokoutis, D. (2010). 45 m.y. of Aegean crust and mantle flow driven by trench retreat. *Geology*, 38(9), 815–818. <https://doi.org/10.1130/G30950.1>
- Brun, J.-P., Faccenna, C., Gueydan, F., Sokoutis, D., Philippon, M., Kydonakis, K., & Gorini, C. (2017). Effects of slab rollback acceleration on Aegean extension. *Bulletin of the Geological Society of Greece*, 50(1), 5–23. <https://doi.org/10.12681/bgsg.11697>
- Brun, J.-P., Faccenna, C., Gueydan, F., Sokoutis, D., Philippon, M., Kydonakis, K., Gorini, C. (2016). The two-stage aegean extension, from localized to distributed, a result of slab rollback acceleration. *Canadian Journal of Earth Sciences*, National Research Council Canada, 53(11), 1142–1157. <https://doi.org/10.1139/cjes-2015-0203ff>. [ffinsu-01271296f](https://doi.org/10.1139/cjes-2015-0203ff)

- Bulut, F., Özener, H., Doğru, A., Aktuğ, B., & Yaltırak, C. (2018). Structural setting along the Western North Anatolian Fault and its influence on the 2014 North Aegean Earthquake (Mw 6.9). *Tectonophysics*, 745, 382–394. <https://doi.org/10.1016/j.tec.2018.07.006>
- Burchfiel, B.C., Nakov, R., Dumurdzanov, N., Papanikolaou, D., Tzankov, T., Serafimovski, T., King, R.W., Kotzev, V., Todosov, A., & Nurce, B. (2008). Evolution and dynamics of the Cenozoic tectonics of the South Balkan extensional system. *Geosphere*, 4, 918–938.
- Burdick, S., van der Hilst, R.D., Vernon, F.L., Martynov, V., Cox, T., Eakins, J., Karasu, G.H., Tylell, J., Astiz, L., & Pavlis, G.L. (2012). Model Update March 2011: Upper Mantle Heterogeneity beneath North America from Traveltime Tomography with Global and USArray Transportable Array Data. *Seismological Research Letters*, 83 (1), 23–28. <https://doi.org/10.1785/gssrl.83.1.23>
- Burtman, V.S. (1994). Meso-Tethyan oceanic sutures and their deformation. *Tectonophysics*, 234, 305–327.
- Candan, O., Çetinkaplan, M., Oberhänsli, R., Rimmelé, G., & Akal, C. (2005). Alpine high-P/low-T metamorphism of the Afyon zone and implications for the metamorphic evolution of Western Anatolia, Turkey. *Lithos*, 84, 102–124.
- Capitanio, F. A., Faccenna, C., & Funicello, R. (2009). Opening of Sirte Basin: Result of slab avalanche? *Earth and Planetary Science Letters*, 285, 210–216. <https://doi.org/10.1016/j.epsl.2009.06.019>
- Capitanio, F. A., Stegman, D.R., Moresi, L., & Sharples W. (2010). Upper plate controls on deep subduction, trench migrations and deformations at convergent margins. *Tectonophysics*, 483, 80–92. <https://doi.org/10.1016/j.tecto.2009.08.020>
- Carminati, E., Lustrino, M., & Doglioni, C. (2012). Geodynamic evolution of the central and western Mediterranean: Tectonics vs. igneous petrology constraints. *Tectonophysics*, 579, 173–192. <https://doi.org/10.1016/j.tecto.2012.01.026>
- Catalano, S., De Guidi, G., Romagnoli, G., Torrisi, S., Tortorici, G., & Tortorici, L. (2008). The migration of plate boundaries in SE Sicily: Influence on the large-scale kinematic model of the African promontory in southern Italy. *Tectonophysics*, 449(1–4), 41–62. <https://doi.org/10.1016/j.tecto.2007.12.003>
- Çemen, I., Catlos, E. J., Gogus, O., & Ozerdem, C. (2006). Postcollisional extensional tectonics and exhumation of the Menderes Massif in the western Anatolia extended terrane, Turkey. *Special Paper - Geological Society of America*, 409, 353–379.
- Channell, J.E.T., & Kozur H.W. (1997). How many oceans? Meliata, Vardar and Pindos oceans in Mesozoic Alpine paleogeography. *Geology* 25(2), 183–186. [https://doi.org/10.1130/0091-7613\(1997\)025<0183:HMOMVA>2.3.CO;2](https://doi.org/10.1130/0091-7613(1997)025<0183:HMOMVA>2.3.CO;2)
- Chase, C.G. (1978). Plate kinematics: The Americas, East Africa, and the rest of the world. *Earth and Planetary Science Letters*, 37(3), 355–368. [https://doi.org/10.1016/0012-821X\(78\)90051-1](https://doi.org/10.1016/0012-821X(78)90051-1)
- Chaumillon, E., & Mascle, J. (1997). From foreland to forearc domains: New multichannel seismic reflection survey of the Mediterranean ridge accretionary complex (Eastern Mediterranean). *Marine Geology*, 138, 237–259. [https://doi.org/10.1016/S0025-3227\(97\)00002-9](https://doi.org/10.1016/S0025-3227(97)00002-9)
- Chousianitis, K., Ganas, A., & Evangelidis, C. P. (2015). Strain and rotation rate patterns of mainland Greece from continuous GPS data and comparison between seismic and geodetic moment release. *Journal of Geophysical Research Solid Earth*, 120, 3909–3931. <https://doi.org/10.1002/2014JB011762>

- Chousianitis, K., Ganas, A., & Gianniou, M. (2013). Kinematic interpretation of present-day crustal deformation in central Greece from continuous GPS measurements. *Journal of Geodynamics*, 71, 1-13. <https://doi.org/10.1016/j.jog.2013.06.004>
- Chousianitis, K., Ganas, A., & Evangelidis, C. P. (2015). Strain and rotation rate patterns of mainland Greece from continuous GPS data and comparison between seismic and geodetic moment release. *Journal of Geophysical Research Solid Earth*, 120, 3909–3931. <https://doi.org/10.1002/2014JB011762>
- Cianetti, S., Gasperini, P., Giunchi, C., & Boschi, E. (2001). Numerical modeling of the Aegean-Anatolian region: geodynamical constraints from observed rheological heterogeneities. *Geophysics Journal International*, 146, 760-780.
- CIESM, 2011. Marine geohazards in the Mediterranean. N°42. In: F.Briand, (Ed). CIESM Workshop Monographs, Marine Geohazards in the Mediterranean, The Mediterranean Science Commission Workshop in Nicosia, 2-5 February 2011, 192pp.
- Cita, M.B., Erba, E., Lucchi, R., Pott, M., van der Meer, R., & Nieto, L. (1996). Stratigraphy and sedimentation in the Mediterranean Ridge diapiric belt, *Marine Geology*, 132(1–4), 131-150. [https://doi.org/10.1016/0025-3227\(96\)00157-0](https://doi.org/10.1016/0025-3227(96)00157-0)
- Clift, P., & Vannucchi, P. (2004). Controls on tectonic accretion versus erosion in subduction zones: Implications for the origin and recycling of the continental crust. *Reviews in Geophysics*, 42, RG2001, <https://doi.org/10.1029/2003RG000127>
- Cloetingh, S. A. P. L., Wortel, M. J. R., & Vlaar, N. J. (1984). Passive margin evolution, initiation of subduction and the Wilson cycle. *Tectonophysics*, 109(1–2), 147–163.
- Cloetingh, S., Koptev, A., Kovács, I., Gerya, T., Beniest, A., Willingshofer, E., et al. (2021). Plume-induced sinking of intra-continental lithospheric mantle: An overlooked mechanism of subduction initiation?. *Geochemistry, Geophysics, Geosystems*, 22, e2020GC009482. <https://doi.org/10.1029/2020GC009482>
- Cloos, M., & Shreve, R.L. (1996). Shear-zone thickness and the seismicity of Chilean- and Marianas-type subduction zones. *Geology*, 24 (2), 107–110. [https://doi.org/10.1130/0091-7613\(1996\)024<0107:SZTATS>2.3.CO;2](https://doi.org/10.1130/0091-7613(1996)024<0107:SZTATS>2.3.CO;2)
- Cocchi, L., Passaro, S., Tontini, F.C., & Ventura, G. (2017). Volcanism in slab tear faults is larger than in island-arcs and back-arcs. *Nature Communications*, 8, 1451. <https://doi.org/10.1038/s41467-017-01626-w>
- Confal, J.M., Faccenda, M., Eken, T., & Taymaz, T. (2018). Numerical simulation of 3-D mantle flow evolution in subduction zone environments in relation to seismic anisotropy beneath the eastern Mediterranean region. *Earth and Planetary Science Letters*, 497, 50-61. <https://doi.org/10.1016/j.epsl.2018.06.005>
- Cornée, J-J., Quillévéré, F., Moissette, P., Fietzke, J., López-Otálvaro, G.E., Melinte-Dobrinescu, M., Philippon, M., van Hinsbergen, D.J.J., Agiadi, K., Koskeridou, E., & Münch, P. (2018). Tectonic motion in oblique subduction forearcs: insights from the revisited Middle and Upper Pleistocene deposits of Rhodes, Greece. *Journal of the Geological Society*, 176, 78-96. <https://doi.org/10.1144/jgs2018-090>
- Corti, G. (2009). Continental rift evolution: From rift initiation to incipient break-up in the Main Ethiopian Rift, East Africa. *Earth-Science Reviews*, 96(1-2), 1-53. <https://doi.org/10.1016/j.earscirev.2009.06.005>
- Crameri, F., Magni, V., Domeier, M. et al. (2020). A transdisciplinary and community-driven database to unravel subduction zone initiation. *Nature Communications*, 11, 3750. <https://doi.org/10.1038/s41467-020-17522-9>

- d'Atri, A., Zuffa, G.G., Cavazza, W., Okay, A.I., & Di Vincenzo, G. (2012). Detrital supply from subduction/accretion complexes to the Eocene–Oligocene post-collisional southern Thrace Basin (NW Turkey and NE Greece). *Sedimentary Geology*, 243–244, 117–129. <https://doi.org/10.1016/j.sedgeo.2011.10.008>
- de Boorder, H., Spakman, W., White, S.H., & Wortel, M.J.R. (1998). Late Cenozoic mineralization, orogenic collapse and slab detachment in the European Alpine Belt. *Earth and Planetary Science Letters*, 164(3–4), 569–575. [https://doi.org/10.1016/S0012-821X\(98\)00247-7](https://doi.org/10.1016/S0012-821X(98)00247-7)
- De Franco, R., Govers, R., & Wortel, R. (2008). Nature of the plate contact and subduction zones diversity. *Earth and Planetary Science Letters*, 271(1–4), 245–253. <https://doi.org/10.1016/j.epsl.2008.04.019>
- DeMets, C., & Merkouriev, S. (2016). High-resolution reconstructions of Pacific–North America plate motion: 20 Ma to present. *Geophysical Journal International*, 207(2), 741–773. <https://doi.org/10.1093/gji/ggw305>
- Di Rosa, M., Farina, F., Marroni, M., Pandolfi, L., Göncüoğlu, M.C., Ellero, A., & Ottria, G. (2019). U-Pb zircon geochronology of intrusive rocks from an exotic block in the Late Cretaceous - Paleocene Taraklı Flysch (northern Turkey): constraints on the tectonics of the Intrapontide suture zone. *Journal of Asian Earth Sciences*, 171, 277–288.
- Deschamps, F., & Tackley, P.J. (2009). Searching for models of thermo-chemical convection that explain probabilistic tomography. II—influence of physical and compositional parameters. *Physics of Earth and Planetary Interiors*, 176, 1–18.
- Dilek, Y., & Altunkaynak, Ş. (2007). Cenozoic crustal evolution and mantle dynamics of post-collisional magmatism in Western Anatolia. *International Geology Review*, 49(5), 431–453. <https://doi.org/10.2747/0020-6814.49.5.431>
- Dilek, Y., & Sandvol, E. (2009). Seismic structure, crustal architecture and tectonic evolution of the Anatolian-African Plate Boundary and the Cenozoic Orogenic Belts in the Eastern Mediterranean Region. *Geological Society, London, Special Publications*, 327, 127–160. <https://doi.org/10.1144/SP327.8>
- Doglioni, C., Agostini, S., Crespi, M., Innocenti, F., Manetti, P., Riguzzi, F., & Savaşçın, Y. (2002). On the extension in western Anatolia and the Aegean sea. In G. Rosenbaum, G.S. Lister (Eds.), *Reconstruction of the evolution of the Alpine-Himalayan Orogen*. *Journal of the Virtual Explorer*, 8, 161–176.
- Dominey-Howes, D. (2002). Documentary and geological records of tsunamis in the Aegean Sea region of Greece and their potential value to risk assessment and disaster management. *Natural Hazards*, 25, 195–224. <https://doi.org/10.1023/A:1014808804611>
- Doutsos, T., & Kokkalas, S. (2001). Stress and deformation patterns in the Aegean region. *Journal of Structural Geology*, 23, 455–472.
- Ebeling, C.W., Okal, E.A., Kalligeris, N., & Synolakis, C.E. (2012). Modern seismological reassessment and tsunami simulation of historical Hellenic Arc earthquakes. *Tectonophysics*, 530–531, 225–239. <https://doi.org/10.1016/j.tecto.2011.12.036>
- Elitez, I., Yaltırak, C., & Aktuğ, B. (2016). Extensional and compressional regime driven left-lateral shear in southwestern Anatolia (eastern Mediterranean): The Burdur-Fethiye Shear Zone. *Tectonophysics*, 688, 26–35. <https://doi.org/10.1016/j.tecto.2016.09.024>
- Elmas, A. (2012). Basement types of the Thrace Basin and a new approach to the pre-Eocene tectonic evolution of the northeastern Aegean and northwestern Anatolia: a review of

- data and concepts. *Geologische Rundschau = International Journal of Earth Sciences*, 101, 1895–1911 (2012). <https://doi.org/10.1007/s00531-012-0756-5>
- El-Sharkawy, A., Meier, T., Lebedev, S., Behrmann, J. H., Hamada, M., Cristiano, L., et al. (2020). The slab puzzle of the Alpine-Mediterranean region: Insights from a new, high-resolution, shear wave velocity model of the upper mantle. *Geochemistry, Geophysics, Geosystems*, 21, e2020GC008993. <https://doi.org/10.1029/2020GC008993>
- Emery, K.O., Heezen, B., & Allan, T.D. (1966). Bathymetry of the Eastern Mediterranean sea. *Deep Sea Research*, 13, 173-192.
- England, P., Howell, A., Jackson, J., & Synolakis, C. (2015). Palaeotsunamis and tsunami hazards in the Eastern Mediterranean *Philosophical Transactions of the Royal Society A: Mathematical, Physical and Engineering Sciences*, 373, 2053. <https://doi.org/10.1098/rsta.2014.0374>
- Ergün, M., Okay, S., Sari, C., Oral, E.Z., Ash, M., Hall, J., & Miller, H. (2005). Gravity anomalies of the Cyprus Arc and their tectonic implications. *Marine Geology*, 221(1–4), 349-358. <https://doi.org/10.1016/j.margeo.2005.03.004>
- Evangelidis, C.P. (2017). Seismic anisotropy in the Hellenic subduction zone: Effects of slab segmentation and subslab mantle flow. *Earth and Planetary Science Letters*, 480, 97-106. <https://doi.org/10.1016/j.epsl.2017.10.003>
- Faccenda, M., Burlini, L., Gerya, T. V., & Mainprice, D. (2008). Fault- induced seismic anisotropy by hydration in subducting oceanic plates. *Nature*, 455(7216), 1097–1100. <https://doi.org/10.1038/nature07376>
- Faccenna, C., Becker, T.W., Auer, L., Billi, A., Boschi, L., Brun, J.-P., Capitanio, F.A., Funicello, F., Horvath, F., Jolivet, L., Piromallo, C., Royden, L., Rossetti, F., & Serpelloni, E. (2014). Mantle dynamics in the Mediterranean, *Review of Geophysics*, 52, 283– 332. <https://doi.org/10.1002/2013RG000444>
- Faccenna, C., Becker, T.W., Jolivet, L., & Keskin, M. (2013). Mantle convection in the Middle East: Reconciling Afar upwelling, Arabia indentation and Aegean trench rollback. *Earth and Planetary Science Letters*, 375, 254-269. <https://doi.org/10.1016/j.epsl.2013.05.043>
- Faccenna, C., Bellier, O., Martinod, J., Piromallo, C., & Regard, V. (2006). Slab detachment beneath eastern Anatolia: A possible cause for the formation of the North Anatolian fault. *Earth and Planetary Science Letters*, 242(1–2), 85-97. <https://doi.org/10.1016/j.epsl.2005.11.046>
- Faccenna, C., Jolivet, L., Piromallo, C., & Morelli A. (2003). Subduction and the depth of convection in the Mediterranean mantle. *Journal of Geophysical Research*, 108(B2), <https://doi.org/10.1029/2001JB001690>
- Ferentinos G. (1990). Offshore geological hazards in the Hellenic arc. *Marine Geotechnology*, 9(4), 261-277. <https://doi.org/10.1080/10641199009388244>
- Fernandes, R.M.S., Bastos, L., Miranda, J.M., Lourenço, N., Ambrosius, B.A.C., Noomen, R., & Simons, W. (2006). Defining the plate boundaries in the Azores region. *Journal of Volcanology and Geothermal Research*, 156(1-2), 1-9. <https://doi.org/10.1016/j.jvolgeores.2006.03.019>
- Flerit, F., Armijo, R., King, G., & Meyer, B. (2004). The mechanical interaction between the propagating North Anatolian Fault and the back-arc extension in the Aegean. *Earth and Planetary Science Letters*, 224(3–4), 347-362. <https://doi.org/10.1016/j.epsl.2004.05.028>
- Fornash, K.F., & Whitney, D.L. (2020). Lawsonite-rich layers as records of fluid and element mobility in subducted crust (Sivrihisar Massif, Turkey). *Chemical Geology*, 533, 119356, <https://doi.org/10.1016/j.chemgeo.2019.119356>

- Foucher, J.P., Chamot-Rooke, N., Alexandry, S., Augustin, J.M., Monti, S., Pavlakis, P., & Voisset, M. (1993). Multibeam bathymetry and seabed reflectivity maps of the MEDRIFT corridor across the eastern Mediterranean Ridge. In: UEG VII., Strasbourg, France. *Terra Cognita, Abstracts*, pp. 278-279.
- Frassi, C., Marroni, M., Pandolfi, L., Göncüoğlu, M.C., Ellero, A., Ottria, G., Sayit, K., McDonald, C.S., Balestrieri, M.L., & Malasoma, A. (2018). Burial and exhumation history of the Daday Unit (central Pontides, Turkey); implications for the closure of the Intra-Pontide oceanic basin. In G. Capponi, A. Festa, G. Rebay (Eds.), *Birth and death of oceanic basins; geodynamic processes from rifting to continental collision in Mediterranean and circum-Mediterranean orogeny*. *Geological Magazine*, 155, 356-376.
- Fytikas, M., Innocenti, F., Manetti, P., Mazzuoli, R., Peccerillo, A., & Villari, L. (1984). Tertiary to quaternary evolution of volcanism in the Aegean region. In JE Dixon, AHF Robertson (Eds.), *The Geological Evolution of the Eastern Mediterranean*, Geological Society of London, Special Publications (Vol. 17, pp. 687–699).
- Gaina, C., Torsvik, T.H., van Hinsbergen, D.J.J., Medvedev, S., Werner, S.C., Labails, C. (2013). The African Plate: A history of oceanic crust accretion and subduction since the Jurassic. *Tectonophysics*, 604, 4-25. <https://doi.org/10.1016/j.tecto.2013.05.037>.
- Galindo-Zaldívar, J., Nieto, L., & Woodside, J. (1996). Structural features of mud volcanoes and the fold system of the Mediterranean Ridge, south of Crete. *Marine Geology*, 132(1-4), 95-112. [https://doi.org/10.1016/0025-3227\(96\)00155-7](https://doi.org/10.1016/0025-3227(96)00155-7)
- Ganas A., & Parsons T. (2009). Three-dimensional model of Hellenic Arc deformation and origin of the Cretan uplift. *Journal of Geophysical Research*, 114(B06404), 1–14. <https://doi.org/10.1029/2008JB005599>
- Ganas, A., Briole, P., Bozionelos, G., Barberopoulou, A., Elias, P., Tsironi, V., Valkaniotis, S., Moshou, A., & Mintourakis, I. (2020). The 25 October 2018 Mw = 6.7 Zakynthos earthquake (Ionian Sea, Greece): A low-angle fault model based on GNSS data, relocated seismicity, small tsunami and implications for the seismic hazard in the west Hellenic Arc. *Journal of Geodynamics*, 137, <https://doi.org/10.1016/j.jog.2020.101731>
- Ganas A., Marinou, A., Anastasiou, D., Paradissis, D., Papazissi, K., Tzavaras, P., & Drakatos, G. (2013). GPS-derived estimates of crustal deformation in the central and north Ionian Sea, Greece: 3-yr results from NOANET continuous network data. *Journal of Geodynamics*, 67, 62-71.
- Gautier, P., Brun, J-P., Moriceau, R., Sokoutis, D., Martinod, J., & Jolivet, L. (1999). Timing, kinematics and cause of Aegean extension: a scenario based on a comparison with simple analogue experiments. *Tectonophysics*, 315(1–4), 31-72. [https://doi.org/10.1016/S0040-1951\(99\)00281-4](https://doi.org/10.1016/S0040-1951(99)00281-4)
- Gautier, Y. (1984). Déformations et métamorphismes associés à la fermeture téthysienne en Anatolie centrale (région de Sivrihisar, Turquie). PhD thesis, University Paris-Sud, France. 236.
- Gerya, T. (2011). Future directions in subduction modeling. *Journal of Geodynamics*, 52, 344-378.
- Gessner, K., Gallardo, L. A., Markwitz, V., Ring, U., & Thomson, S. N. (2013). What caused the denudation of the Menderes Massif; review of crustal evolution, lithosphere structure, and dynamic topography in southwest Turkey. *Gondwana Research*, 24(1), 243-274.
- Gessner, K., Markwitz, V., & Güngör, T. (2017). Crustal fluid flow in hot continental extension: tectonic framework of geothermal areas and mineral deposits in western Anatolia.

- Geological Society, London, Special Publications, 453, 289-311.
<https://doi.org/10.1144/SP453.7>
- Gianni, G.M., Navarrete, C., & Spagnotto, S. (2019). Surface and mantle records reveal an ancient slab tear beneath Gondwana. *Scientific Reports*, 9, 19774.
<https://doi.org/10.1038/s41598-019-56335-9>
- Glodny, J., & Hetzel, R. (2007). Precise U-Pb ages of syn-extensional Miocene intrusions in the central Menderes Massif, western Turkey. *Geological Magazine*, 144, 1–12.
- Göncüoğlu, M.C. (2010). Introduction to the Geology of Turkey: Geodynamic Evolution of the Pre-Alpine and Alpine Terranes. General Directorate of Mineral Research and Exploration Monography Series, 5, 1-66.
- Göncüoğlu, M.C., Marroni, M., Pandolfi, L., Ellero, A., Ottria, G., Catanzariti, R., Tekin, U.K., & Sayit, K. (2014). The Arkot Dag Melange in Arac area, central Turkey; evidence of its origin within the geodynamic evolution of the Intra-Pontide suture zone. *Journal of Asian Earth Sciences*, 85, 117-139. <https://doi.org/10.1016/j.jseaes.2014.01.013>
- Göncüoğlu, M.C., Marroni, M., Sayit, K., Tekin, U.K., Ottria, G., Pandolfi, L., & Ellero, A. (2012). The Ayli Dag ophiolite sequence (central-northern Turkey); a fragment of Middle Jurassic oceanic lithosphere within the Intra-Pontide suture zone. *Ophioliti*, 37, 77-92.
- Gordon, R.G., & Stein, S. (1992). Global Tectonics and Space Geodesy. *Science*, 17, 333-342.
- Govers, R., & Fichtner, A. (2016). Signature of slab fragmentation beneath Anatolia from full-waveform tomography. *Earth and Planetary Science Letters*, 450, 10-19.
<https://doi.org/10.1016/j.epsl.2016.06.014>
- Govers, R., & Wortel, M.J.R. (2005). Lithosphere tearing at STEP faults: response to edges of subduction zones. *Earth and Planetary Science Letters*, 236(1–2), 505-523.
<https://doi.org/10.1016/j.epsl.2005.03.022>
- Gürer, Ö., Sangu, E., & Özburan, M. (2006). Neotectonics of the SW Marmara region, NW Anatolia, Turkey. *Geological Magazine*, 143 (2): 229–241.
<https://doi.org/10.1017/S0016756805001469>
- Gurnis, M., Hall, C., & Lavier, L. (2004). Evolving force balance during incipient subduction. *Geochemistry, Geophysics, Geosystems*, 5, Q07001.
<https://doi.org/10.1029/2003GC000681>
- Güvercin, S.E., Konca, A.Ö., Özbakır, A.D., Ergintav, S., & Karabulut, H. (2021). New focal mechanisms reveal fragmentation and active subduction of the Antalya slab in the Eastern Mediterranean. *Tectonophysics*, 805, 228792.
<https://doi.org/10.1016/j.tecto.2021.228792>
- Hall, C.E., Gurnis, M., Sdralias, M., Lavier, L.L., & Müller, R.D. (2003). Catastrophic initiation of subduction following forced convergence across fracture zones. *Earth and Planetary Science Letters*, 212(1–2), 15– 30.
- Hall, J., Aksu, A.E., Elitez, I., Yaltrak, C., & Çifçi, G. (2014). The Fethiye–Burdur Fault Zone: A component of upper plate extension of the subduction transform edge propagator fault linking Hellenic and Cyprus Arcs, Eastern Mediterranean. *Tectonophysics*, 635, 80-99.
<https://doi.org/10.1016/j.tecto.2014.05.002>
- Hall, J., Aksu, A.E., Yaltrak, C., & Winsor, J.D. (2009). Structural architecture of the Rhodes Basin: A deep depocentre that evolved since the Pliocene at the junction of Hellenic and Cyprus Arcs, eastern Mediterranean. *Marine Geology*, 258(1-4), 1-23.
<https://doi.org/10.1016/j.margeo.2008.02.007>

- Hall, R., Audley-Charles, M.G., & Carter, D.J. (1984). The significance of Crete for the evolution of the Eastern Mediterranean. Geological Society, London, Special Publications, 17, 499-516. <https://doi.org/10.1144/GSL.SP.1984.017.01.37>
- Halpaap, F., Rondenay, S., & Ottemöller, L. (2018). Seismicity, deformation, and metamorphism in the Western Hellenic Subduction Zone: New constraints from tomography. *Journal of Geophysical Research: Solid Earth*, 123, 3000–3026. <https://doi.org/10.1002/2017JB015154>
- Hansen, S.E., Evangelidis, C.P., & Papadopoulos, G.A. (2019). Imaging slab detachment within the Western Hellenic Subduction Zone. *Geochemistry, Geophysics, Geosystems*, 20, 895–912. <https://doi.org/10.1029/2018GC007810>
- Hayes, G. (2018). Slab2 - A Comprehensive Subduction Zone Geometry Model: U.S. Geological Survey data release, <https://doi.org/10.5066/F7PV6JNV>
- Heezen, B.C., & Ewing, M., 1963. The Mid Oceanic Ridge. In M.N. Hill (Ed.), *The Seas* (Vol. 3, pp. 388-410), Interscience, New York.
- Hooft, E.E.E., Nomikou, P., Toomey, D.R., Lampridou, D., Getz, C., Christopoulou, M-E., O'Hara, D., Arnoux, G.M., Bodmer, M., Gray, M., Heath, B.A., VanderBeek, B.P. (2017). Backarc tectonism, volcanism, and mass wasting shape seafloor morphology in the Santorini-Christiana-Amorgos region of the Hellenic Volcanic Arc. *Tectonophysics*, 712–713, 396-414. <https://doi.org/10.1016/j.tecto.2017.06.005>
- Hosseini, K. (2016). Global multiple-frequency seismic tomography using teleseismic and core-diffracted body waves. Dissertation, LMU München: Faculty of Geosciences. <https://doi.org/10.5282/edoc.19597>
- Hosseini, K., Matthews, K.J., Sigloch, K., Shephard, G.E., Domeier, M., & Tsekhmistrenko, M. (2018). SubMachine: Web-Based tools for exploring seismic tomography and other models of Earth's deep interior. *Geochemistry, Geophysics, Geosystems*, 19. <https://doi.org/10.1029/2018GC007431>
- Hosseini, K., Sigloch, K., Tsekhmistrenko, M., Zaheri, A., Nissen-Meyer, T., & Igel, H. (2020). Global mantle structure from multifrequency tomography using P, PP and P-diffracted waves. *Geophysical Journal International*, 220(1), 96–141. <https://doi.org/10.1093/gji/ggz394>
- Howell, A., Jackson, J., England, P., Higham, T., & Synolakis, C. (2015). Late Holocene uplift of Rhodes, Greece: evidence for a large tsunamigenic earthquake and the implications for the tectonics of the eastern Hellenic Trench System. *Geophysical Journal International*, 203(1), 459–474.
- Hubert-Ferrari, A., Armijo, R., King, G., Meyer, B., & Barka, A. (2002). Morphology, displacement, and slip rates along the North Anatolian Fault, Turkey, *Journal of Geophysical Research*, 107(B10), 2235. <https://doi.org/10.1029/2001JB000393>
- Huchon, P., Lybérís, N., Angelier, J., Le Pichon, X., & Renard, V. (1982). Tectonics of the Hellenic trench: A synthesis of sea-beam and submersible observations, *Tectonophysics*, 86(1-3), 69-112. [https://doi.org/10.1016/0040-1951\(82\)90062-2](https://doi.org/10.1016/0040-1951(82)90062-2)
- Husson, L., Brun, J-P., Yamato, P., & Faccenna, C. (2009). Episodic slab rollback fosters exhumation of HP–UHP rocks. *Geophysics Journal International*, 179, 1292–1300
- Jackson, J. (1994). Active tectonics of the Aegean region. *Annual Review of Earth and Planetary Sciences*, 22, 239-271.

- Jackson, J., and McKenzie, D. (1988). The relationship between plate motions and seismic moment tensors and the rates of active deformation in the Mediterranean and Middle East. *Geophysical Journal*, 93, 45-73.
- Jolivet, L., & Brun, J.-P. (2010). Cenozoic geodynamic evolution of the Aegean, *International Journal of Earth Sciences*, 99(1), 109-138. <https://doi.org/10.1007/s00531-008-0366-4>
- Jolivet, L., & Faccenna, C. (2000). Mediterranean extension and the Africa-Eurasia collision, *Tectonics*, 19(6), 1095–106. <https://doi.org/10.1029/2000TC900018>
- Jolivet, L., Faccenna, C., Goffé, B., Burov, E., & Agard, P. (2003). Subduction tectonics and exhumation of high-pressure metamorphic rocks in the Mediterranean orogens. *American Journal of Science*, 303, 353–409. <https://doi.org/10.2475/ajs.303.5.353>
- Jolivet, L., Faccenna, C., Huet, B., Labrousse, L., Le Pourhiet, L., et al. (2013). Aegean tectonics: Strain localisation, slab tearing and trench retreat. *Tectonophysics*, 597-598, 1-33.
- Jolivet, L., Menant, A., Clerc, C., Sternai, P., Bellahsen, N., Leroy, S., Pik, R., Stab, M., Faccenna, C., & Gorini, C. (2018). Extensional crustal tectonics and crust-mantle coupling, a view from the geological record. *Earth-Science Reviews*, 185, 1187-1209. <https://doi.org/10.1016/j.earscirev.2018.09.010>.
- Jolivet, L., Menant, A., Sternai, P., Rabillard, A., Arbaret, L., Augier, R., Laurent, V., Beaudoin, A., Grasemann, B., Huet, B., Labrousse, L., & Le Pourhiet, L. (2015). The geological signature of a slab tear below the Aegean, *Tectonophysics*, 659, 166-182. <https://doi.org/10.1016/j.tecto.2015.08.004>
- Jost, M.L., Knabenbauer, O., Cheng, J., & Harjes, H-P. (2002). Fault plane solutions of microearthquakes and small events in the Hellenic arc, *Tectonophysics*, 356 (1–3), 87-114. [https://doi.org/10.1016/S0040-1951\(02\)00378-5](https://doi.org/10.1016/S0040-1951(02)00378-5)
- Karabulut, H., Paul, A., Özbakır, A.D., Ergün, T., & Şentürk, S. (2019). A new crustal model of the Anatolia–Aegean domain: evidence for the dominant role of isostasy in the support of the Anatolian plateau. *Geophysical Journal International*, 218(1), 57–73. <https://doi.org/10.1093/gji/ggz147>
- Karagianni, E.E., Papazachos, C.B., Panagiotopoulos, D.G., Suhadolc, P., Vuan, A., & Panza, G.F. (2005). Shear velocity structure in the Aegean area obtained by inversion of Rayleigh waves. *Geophysical Journal International*, 160(1), 127–143. <https://doi.org/10.1111/j.1365-246X.2005.02354.x>
- Karaoğlu, Ö., & Helvacı, C. (2014). Isotopic evidence for a transition from subduction to slab-tear related volcanism in western Anatolia, Turkey. *Lithos*, 192–195, 226-239. <https://doi.org/10.1016/j.lithos.2014.02.006>
- Kárason, H., & van der Hilst R. (2000). Constraints on mantle convection from seismic tomography. *Geophysical Monograph*, 121, 277–288.
- Karato, S.-I. (1995). Effects of water on seismic wave velocities in the upper mantle. *Proceedings of the Japan Academy, Ser. B*, 71, 61–66.
- Kassarav, I., Makropoulos, K., Bourova, E., Pedersen, H., & Hatzfeld, D. (2005). 3 - Upper mantle structure of the Aegean derived from two-station phase velocities of fundamental mode Rayleigh waves. In: Editor(s): Fytikas, M., & Vogioukalakis, G.E. (Eds). *Developments in Volcanology*, 7, 19-45. [https://doi.org/10.1016/S1871-644X\(05\)80031-2](https://doi.org/10.1016/S1871-644X(05)80031-2)
- Kastens, K.A. (1991). Rate of outward growth of the Mediterranean Ridge accretionary complex, *Tectonophysics*, 199, 25–50.

- Kastens, K.A., Breen, N.A., & Cita, M.B. (1992). Progressive deformation of an evaporite-bearing accretionary complex: SeaMARC I, SeaBeam and piston-core observations from the Mediterranean Ridge. *Marine Geophysics Research*, 14, 249–298. <https://doi.org/10.1007/BF01203620>.
- Kaya, O. (1981). Miocene reference section for the coastal parts of West Anatolia, *Newsletters on Stratigraphy*, 10, 164–191.
- Kempler, D., & Ben-Avraham, Z. (1987). The tectonic evolution of the Cyprean Arc, *Annales Tectonicae*, 1, 58–71.
- Kenyon, N.H., Belderson, R.H., & Stride, A.H. (1982). Detailed tectonic trends on the central part of the Hellenic outer ridge and in the Hellenic trench system. *Geological Society of London*, 10, 335–343.
- Keranen, K., & Klemperer, S.L. (2008). Discontinuous and diachronous evolution of the Main Ethiopian Rift: Implications for development of continental rifts. *Earth and Planetary Science Letters*, 265(1–2), 96–111. <https://doi.org/10.1016/j.epsl.2007.09.038>
- Ketin, I. (1948). Über die tektonisch-mechanischen Folgerungen aus den großen anatolischen Erdbeben des letzten Dezenniums. *Geologische Rundschau = International Journal of Earth Sciences*, 36, 77–83. <https://doi.org/10.1007/BF01791916>
- Kind, R., Eken, T., Tilmann, F., Sodoudi, F., Taymaz, T., Bulut, F., Yuan, X., Can, B., & Schneider, F. (2015). Thickness of the lithosphere beneath Turkey and surroundings from S-receiver functions, *Solid Earth*, 6, 971–984. <https://doi.org/10.5194/se-6-971-2015>
- Kinnaird, T., & Robertson, A. (2012). Tectonic and sedimentary response to subduction and incipient continental collision in southern Cyprus, easternmost Mediterranean region. *Geological Society, London, Special Publications*, 372, 585–614. <https://doi.org/10.1144/SP372.10>
- Kokinou, E., Papadimitriou, E., Karakostas, V., Kamberis, E., & Vallianatos, F. (2006). The Kefalonia Transform Zone (offshore Western Greece) with special emphasis to its prolongation towards the Ionian Abyssal Plain. *Marine Geophysical Researches*, Dordrecht, 27(4), 241–252. <https://doi.org/10.1007/s11001-006-9005-2>
- Kokkalas, S., Paraskevas, X., Koukouvelas, I., & Doutsos, T. (2006). Postcollisional contractional and extensional deformation in the Aegean region. *Special Paper of the Geological Society of America*, 409, 97–123. <https://doi.org/10.1130/0-8137-2409-0.97>
- Kopf, A.J., (2002). Significance of mud volcanism. *Reviews in Geophysics*, 40(2), 1005, <https://doi.org/10.1029/2000RG000093>
- Kopf, A., Mascle, J., & Klaeschen, D. (2003). The Mediterranean Ridge: A mass balance across the fastest growing accretionary complex on Earth, *Journal of Geophysical Research*, 108(B8), 2372. <https://doi.org/10.1029/2001JB000473>
- Kreemer, C., Chamot-Rooke, N., & Le Pichon, X. (2004). Constraints on the evolution and vertical coherency of deformation in the Northern Aegean from a comparison of geodetic, geologic and seismologic data, *Earth and Planetary Science Letters*, 225(3–4), 329–346. <https://doi.org/10.1016/j.epsl.2004.06.018>
- Laigle, M., Hirn, A., Sachpazi, M., & Clément, C. (2002). Seismic coupling and structure of the Hellenic subduction zone in the Ionian Islands region. *Earth and Planetary Science Letters*, 200(3–4), 243–253. [https://doi.org/10.1016/S0012-821X\(02\)00654-4](https://doi.org/10.1016/S0012-821X(02)00654-4)
- Laigle, M., Sachpazi, M., and Him, A. (2004). Variation of seismic coupling with slab detachment and upper plate structure along the western Hellenic subduction zone, *Tectonophysics*, 391, 85–95.

- Lamont, T.N., Searle, M.P., Roberts, N.M.W., Waters, D.J., Palin, R.M., Smye, A.J., Dyck, B., Gopon, P., Weller, O.M., St-Onge, M. (2019). Compressional origin to the Naxos metamorphic core complex, Greece: structure, petrography and thermobarometry. *Geological Society of America Bulletin*, 132, 149-197. <https://doi.org/10.1130/B31978.1>
- Legendre, C.P., Meier, T., Lebedev, S., Friederich, W., & Viereck-Götte, L. (2012). A shear wave velocity model of the European upper mantle from automated inversion of seismic shear and surface waveforms, *Geophysical Journal International*, 191(1), 282–304. <https://doi.org/10.1111/j.1365-246X.2012.05613.x>
- Le Pichon, X., & Angelier, J. (1979). The Hellenic arc and trench system; A key to the neotectonic evolution of the eastern Mediterranean area. *Tectonophysics*, 60, 1-42.
- Le Pichon X., & Angelier J. (1981). The Aegean Sea. *Philosophical Transactions of the Royal Society of London. Series A., Mathematical and Physical Sciences*, 300, 357-372, <http://doi.org/10.1098/rsta.1981.0069>
- Le Pichon, X., Chamot-Rooke, N., & Lallemand, S. (1995). Geodetic determination of the kinematics of central Greece with respect to Europe: Implications for eastern Mediterranean tectonics. *Journal of Geophysical Research*, 100, 12,675-12,690.
- Le Pichon, X., Chamot-Rooke, N., Lallemand, S., Noomen, R., & Veis, G. (1995). Geodetic determination of the kinematics of central Greece with respect to Europe: Implications for eastern Mediterranean tectonics. *Journal of Geophysical Research*, 100(B7), 12675–12690. <https://doi.org/10.1029/95JB00317>
- Le Pichon X., Henry P., & Lallemand S. (1993). Accretion and erosion in subduction zones — the role of fluids. *Annual Reviews in Earth and Planetary Science*, 21, 307-331.
- Le Pichon, X., Lallemand, S.J., Chamot-Rooke, N., Lemeur, D., & Pascal, G. (2002). The Mediterranean Ridge backstop and the Hellenic nappes, *Marine Geology*, 186(1-2), 111-125. [https://doi.org/10.1016/S0025-3227\(02\)00175-5](https://doi.org/10.1016/S0025-3227(02)00175-5)
- Le Pichon X., Lyberis, N., Angelier, J., & Renard, V. (1982). Strain distribution over the East Mediterranean Ridge: a synthesis incorporating new Sea-Beam Data. *Tectonophysics*, 86, 243-274. [https://doi.org/10.1016/0040-1951\(82\)90069-5](https://doi.org/10.1016/0040-1951(82)90069-5)
- Le Pichon, X., Şengör, A.M.C., & İmren, C. (2019). A new approach to the opening of the eastern Mediterranean Sea and the origin of the Hellenic subduction zone. Part 2: The Hellenic subduction zone. *Canadian Journal of Earth Sciences*, 56(11), 1144-1162. <https://doi.org/10.1139/cjes-2018-0315>
- Limonov, A.F., Woodside, J.M., Cita, M.B., & Ivanov, M.K. (1996). The Mediterranean Ridge and related mud diapirism: a background. *Marine Geology*, 132(1-4), 7-19. [https://doi.org/10.1016/0025-3227\(96\)00150-8](https://doi.org/10.1016/0025-3227(96)00150-8)
- Lyberis, N. (1984). Tectonic evolution of the North Aegean trough, *Geological Society, London, Special Publications*, 17, 709-725.
- Lykousis, V., Alexandri, S., Woodside, J., de Lange, G., Dählmann, A., Perissoratis, C., Heeschen, K., Ioakim, C., Sakellariou, D., Nomikou, P., Rousakis, G., Casas, D., Ballas, D., & Ercilla, G. (2009). Mud volcanoes and gas hydrates in the Anaximander mountains (Eastern Mediterranean Sea). *Marine and Petroleum Geology*, 26(6), 854-872. <https://doi.org/10.1016/j.marpetgeo.2008.05.002>
- Maffione, M., & van Hinsbergen, D. J. J. (2018). Reconstructing plate boundaries in the Jurassic Neo-Tethys from the East and West Vardar Ophiolites (Greece and Serbia). *Tectonics*, 37, 858– 887. <https://doi.org/10.1002/2017TC004790>

- Maggini, M., & Caputo, R. (2020). Sensitivity analysis for crustal rheological profiles: examples from the Aegean Region. *Annals of Geophysics*, 63(3), GT334. <http://dx.doi.org/10.4401/ag-8244>
- Mahatsente, R., Alemdar, S., Çemen, I. (2017). Effect of Slab-Tear on Crustal Structure in Southwestern Anatolia. In: Çemen, I, Yilmaz, Y. (Eds.) *Active Global Seismology. Geophysical Monograph Series*, 103-119. <https://doi.org/10.1002/9781118944998.ch4>
- Makris, J. (1978). The crust and upper mantle of the Aegean region from deep seismic soundings, *Tectonophysics*, 46(3–4), 269-284. [https://doi.org/10.1016/0040-1951\(78\)90207-X](https://doi.org/10.1016/0040-1951(78)90207-X)
- Malandri, C., Soukis, C., Maffione, M., Özkaptan, M., Vassilakis, E., Lozios, S., & van Hinsbergen, D.J.J. (2017). Vertical-axis rotations accommodated along the Mid-Cycladic lineament on Paros Island in the extensional heart of the Aegean orocline (Greece). *Lithosphere*, 9(1), 78–99. <https://doi.org/10.1130/L575>
- Mantovani, E., Albarello, D., Tamburelli, C., Babbucci, D., & Viti, M. (1997). Plate convergence, crustal delamination, extrusion tectonics and minimization of shortening work as main controlling factors of the recent Mediterranean deformation pattern. In E. Mantovani (Ed.), *Geodynamics of the Mediterranean region and its implications for seismic and volcanic risk. Annali di Geofisica (Vol. 40, pp. 611-643).*
- Marone, F., van der Meijde, M., van der Lee, S., Giardini, D. (2003). Joint inversion of local, regional and teleseismic data for crustal thickness in the Eurasia–Africa plate boundary region. *Geophysical Journal International*, 154(2), 499–514. <https://doi.org/10.1046/j.1365-246X.2003.01973.x>
- Marone, F., Van Der Lee, S., & Giardini, D. (2004a). Three-dimensional upper-mantle S-velocity model for the Eurasia–Africa plate boundary region. *Geophysical Journal International*, 158(1), 109–130. <https://doi.org/10.1111/j.1365-246X.2004.02305.x>
- Marone F., Van Der Lee S., & Giardini D. (2004b). Shallow anisotropy in the Mediterranean mantle from surface waves. *Geophysical Research Letters*, 31(6). <https://doi.org/10.1029/2003GL018948>
- Marroni, M., Frassi, C., Göncüoğlu, M.C., Di Vincenzo, G., Pandolfi, L., Rebay, G., Ellero, A., & Ottria, G. (2014). Late Jurassic amphibolite-facies metamorphism in the Intra-Pontide Suture Zone (Turkey): an eastward extension of the Vardar Ocean from the Balkans into Anatolia? *Journal of the Geological Society*, 171(5), 605–608. <https://doi.org/10.1144/jgs2013-104>
- Marsellos, A.E., Kidd, W.S.F., & Garver, J.I. (2010). Extension and exhumation of the HP/LT rocks in the Hellenic forearc ridge. *American Journal of Science* 310(1), 1-36. <https://doi.org/10.2475/01.2010.01>
- Mascle, J., & Chaumillon, E. (1997). Pre-collisional geodynamics of the Mediterranean Sea: the Mediterranean Ridge and the Tyrrhenian Sea. *Annales Geophysicae*, 40(3), <https://www.annalsofgeophysics.eu/index.php/annals/article/view/3891>
- Mascle, J., Huguen, C., Benkhelil, J., ChamotRooke, N., Chaumillon, E., Foucher, J.P., Griboulard, R., Kopf, A., Lamarche, G., Volkonskaia, A., Woodside, J., & Zitter, T. (1999). Images may show start of European-African Plate collision. *Eos Transactions AGU*, 80(37), 421-436.
- Mascle, J., Le Cleac'h, A., & Jongsma, D. (1986). The eastern Hellenic margin from Crete to Rhodes: Example of progressive collision. *Marine Geology*, 73(1-2), 145-168. [https://doi.org/10.1016/0025-3227\(86\)90116-7](https://doi.org/10.1016/0025-3227(86)90116-7)

- McClusky S. et al. (2000). Global positioning system constraints on plate kinematics and dynamics in the eastern Mediterranean and Caucasus. *Journal of Geophysical Research*, 105(B3), 5695–5719. <https://doi.org/10.1029/1999JB900351>
- McKenzie, D. (1972). Active Tectonics of the Mediterranean Region. *Geophysical Journal of the Royal Astronomical Society*, 30, 109-185. <https://doi.org/10.1111/j.1365-246X.1972.tb02351.x>
- McKenzie, D. (1978). Active tectonics of the Alpine-Himalayan belt: the Aegean Sea and surrounding regions. *Geophysics Journal of the Royal Astronomical Society*, 55, 217-254.
- Meier, T., Becker, D., Endrun, B., Rische, M., Bohnhoff, M., Stöckhert, B., & Harjes, H.-P. (2007). A model for the Hellenic subduction zone in the area of Crete based on seismological investigations. *Geological Society, London, Special Publications*, 291, 183-199. <https://doi.org/10.1144/SP291.9>
- Meier, T., Dietrich, K., Stöckhert, B., & Harjes, H-P. (2004). One-dimensional models of shear wave velocity for the eastern Mediterranean obtained from the inversion of Rayleigh wave phase velocities and tectonic implications, *Geophysical Journal International*, 156(1), 45–58. <https://doi.org/10.1111/j.1365-246X.2004.02121.x>
- Meighan, H. E., ten Brink, U., & Pulliam, J. (2013), Slab tears and intermediate-depth seismicity, *Geophysical Research Letters*, 40, 4244– 4248. <https://doi.org/10.1002/grl.50830>
- Meijer, P. T., and Wortel, M. J. R. (1999), Cenozoic dynamics of the African plate with emphasis on the Africa-Eurasia collision, *J. Geophys. Res.*, 104(B4), 7405– 7418, doi:10.1029/1999JB900009.
- Mercier, J.L. (1981). Extensional-compressional tectonics associated with the Aegean Arc: comparison with the Andean Cordillera of south Peru-north Bolivia. *Philosophical Transactions of the Royal Society of London A (Mathematical and Physical Sciences)*, 300(1454), 337-355.
- Messini, A., Papadimitriou, E., Karakostas, V., & Baskoutas, I. (2007). Stress interaction between thrust faults along the SW Hellenic arc (Greece). *Bulletin of the Geological Society of Greece*, 40(1), 386-398. doi:<https://doi.org/10.12681/bgsg.16625>
- Meulenkamp, J.E., Wortel, M.J.R., van Wamel, W.A., Spakman, W., & Hoogerduyn, S.E. (1988). On the Hellenic subduction zone and the geodynamic evolution of Crete since the late middle Miocene. *Tectonophysics*, 146, 203-215.
- Moix, P., Beccaletto, L., Kozur, H.W., Hochard, C., Rosselet, F., & Stampfli, G.M. (2008). A new classification of the Turkish terranes and sutures and its implication for the paleotectonic history of the region. *Tectonophysics*, 451(1–4), 7-39. <https://doi.org/10.1016/j.tecto.2007.11.044>
- Morris, A., & Robertson, A.H.F. (1993). Miocene remagnetisation of carbonate platform and Antalya Complex units within the Isparta angle, SW Turkey, *Tectonophysics*, 220(1–4), 243-266. [https://doi.org/10.1016/0040-1951\(93\)90234-B](https://doi.org/10.1016/0040-1951(93)90234-B)
- Nikitas, A., Triantaphyllou, M.V., Rousakis, G., Panagiotopoulos, I., Pasadakis, N., Hatzianestis, I., & Gogou, A. (2021). Pre-Messinian Deposits of the Mediterranean Ridge: Biostratigraphic and Geochemical Evidence from the Olimpi Mud Volcano Field. *Water*, 13(10), 1367. <https://doi.org/10.3390/w13101367>
- Nyst, M., & Thatcher, W. (2004). New constraints on the active tectonic deformation of the Aegean. *Journal of Geophysical Research*, 109(11), 23. <https://doi.org/10.1029/2003JB002830>

- Okal, E.A., Synolakis, C.E., Uslu, B., Kalligeris, N., & Voukouvalas, E. (2009). The 1956 earthquake and tsunami in Amorgos, Greece, *Geophysical Journal International*, 178(3), 1533–1554. <https://doi.org/10.1111/j.1365-246X.2009.04237.x>
- Okay, A.I. (1980a). Mineralogy, petrology and phase relations of glaucophane–lawsonite zone blueschists from the Tavşanlı region, northwest Turkey. *Contributions to Mineralogy and Petrology*, 72, 243–255.
- Okay, A.I. (1980b). Lawsonite zone blueschists and a sodic amphibole producing reaction in the Tavşanlı region, northwest Turkey. *Contributions to Mineralogy and Petrology*, 75, 179–186.
- Okay, A.I. (1982). Incipient blueschist metamorphism and metasomatism in the Tavşanlı region, northwest Turkey. *Contributions to Mineralogy and Petrology*, 79, 361–367.
- Okay, A.I. (1984). Distribution and characteristics of the north-west Turkish blueschists. In J.E. Dixon, A.H.F. Robertson (Eds.), *The Geological Evolution of the Eastern Mediterranean*. Geological Society of London Special Publications, 17, 455–466.
- Okay, A.I. (1986). High-pressure/low-temperature metamorphic rocks of Turkey. In B.W. Evans, E.H. Brown (Eds.), *Blueschists and eclogites*. Memoir-Geological Society of America, 164, 333–347.
- Okay, A.I. (2008). Geology of Turkey: A synopsis. *Anschnitt*, 21, 19–42.
- Okay, A.I., & Kelley, S.P. (1994). Tectonic setting, petrology and geochronology of jadeite+glaucophane and chloritoid+glaucophane schists from north-west Turkey. *Journal of Metamorphic Geology*, 12, 455–466.
- Okay A.I., & Satir, M. (2000). Coeval plutonism and metamorphism in a latest Oligocene metamorphic core complex in northwest Turkey. *Geological Magazine*, 137, 495–516.
- Okay, A.I., & Satır, M. (2006). Geochronology of Eocene plutonism and metamorphism in northwest Turkey: evidence for a possible magmatic arc. *Geodinamica Acta* 19, 251–265.
- Okay A.I., & Tüysüz, O. (1999). Tethyan sutures of northern Turkey. In B. Durand, L. Jolivet, F. Horváth, M. Séranne (Eds.), *The Mediterranean Basins: Tertiary Extension with the Alpine Orogen*. Geological Society, London, Special Publications (Vol. 156, pp. 475–515). <https://doi.org/10.1144/GSL.SP.1999.156.01.22>
- Okay, A.I., & Whitney, D.L. (2010). Blueschists, eclogites, ophiolites and suture zones in northwest Turkey: A review and a field excursion guide. *Ophioliti*, 35(2), 131–172.
- Okay, A.I., Satır, M., Maluski, H., Siyako, M., Monie, P., Metzger, R., & Akyüz, S. (1996). Paleo- and Neo-Tethyan events in northwestern Turkey: geologic and geochronologic constraints. In T.M. Harrison (Ed.), *The Tectonic Evolution of Asia*. Cambridge University Press, Cambridge, pp. 420–441.
- Okay, A.I., Satir, M., Tuysuz, O., Akyuzm S., & Chen, F. (2001). The tectonics of Strandja Massif: late-Variscan and mid-Mesozoic deformation and metamorphism in the northern Aegean. *Geologische Rundschau = International Journal of Earth Sciences*, 90, 217–233.
- Okay, A.I., Siyako, M., & Burkan, K.A. (1991). Geology and tectonic evolution of the Biga Peninsula, northwest Turkey. *Bulletin of Technical University Istanbul*, 44, 191–256.
- Okay, A.I., Sunal, G., Sherlock, S., Altnr, D., Tuysuz, O., Kylander-Clark, A.R.C., & Aygul, M. (2013). Early Cretaceous sedimentation and orogeny on the active margin of Eurasia; southern-central Pontides, Turkey. *Tectonics*, 32, 1247–1271.
- Okay, A.I., Sunal, G., Sherlock, S., Kylander-Clark, A. R. C., & Özcan, E. (2020). İzmir-Ankara suture as a Triassic to Cretaceous plate boundary—Data from central Anatolia. *Tectonics*, 39, e2019TC005849. <https://doi.org/10.1029/2019TC005849>

- Oner, Z., Dilek, Y., & Kadioglu, Y.K. (2010). Geology and geochemistry of the synextensional Salihli granitoid in the Menderes core complex, western Anatolia, Turkey. *International Geology Review*, 52(2-3), 336-368. <https://doi.org/10.1080/00206810902815871>
- Oral, M.B., R. Reilinger, M. Toksöz, R.W. King, A. Barka, I. Kinik and O. Lenk (1995). Global positioning system offers evidence of plate motions in eastern Mediterranean. *Eos* 76: 9-11.
- Özacar, A.A., Zandt, G., Gilbert, H., & Beck, S.L. (2010). Seismic images of crustal variations beneath the East Anatolian Plateau (Turkey) from teleseismic receiver functions. *Geological Society, London, Special Publications*, 340, 485-496. <https://doi.org/10.1144/SP340.21>
- Özbakır, A.D., Govers, R., & Fichtner, A. (2020). The Kefalonia Transform Fault: A STEP fault in the making. *Tectonophysics*, 787, 228471. <https://doi.org/10.1016/j.tecto.2020.228471>.
- Özbakır, A.D., Şengör, A.M.C., Wortel, M.J.R., & Govers, R. (2013). The Pliny–Strabo trench region: A large shear zone resulting from slab tearing. *Earth and Planetary Science Letters*, 375, 188-195. <https://doi.org/10.1016/j.epsl.2013.05.025>
- Özdamar, S., Billor, M.Z., Sunal, G., Esenli, F., & Roden, N.F. (2013). First U–Pb SHRIMP zircon and $40\text{Ar}/39\text{Ar}$ ages of metarhyolites from the Afyon–Bolkardag Zone, SW Turkey: Implications for the rifting and closure of the Neo-Tethys, *Gondwana Research*, 24(1), 377-391. <https://doi.org/10.1016/j.gr.2012.10.006>
- Papadimitriou, E.E., and Karakostas, V.G. (2005). Faulting geometry and seismic coupling of the southwest part of the Hellenic subduction zone, Abstract in 33rd IASPEI General Assembly, Santiago, Chile, 2-8 October 2005.
- Papadimitriou, E.E., Karakostas, V.G. (2008). Rupture model of the great AD 365 Crete earthquake in the southwestern part of the Hellenic Arc. *Acta Geophys.* 56, 293–312. <https://doi.org/10.2478/s11600-008-0001-6>
- Papadopoulos, G.A. (1997). On the interpretation of large-scale seismic tomography images in the Aegean sea area. *Annals of Geophysics*, 40(1), 37-42. <https://doi.org/10.4401/ag-3933>
- Papadopoulos, G.A., Daskalaki, E., Fokaefs, A., & Giraleas, N. (2007). Tsunami hazards in the Eastern Mediterranean: Strong earthquakes and tsunamis in the East Hellenic Arc and Trench system. *Natural Hazards Earth System Science*, 7, 57–64. <https://doi.org/10.5194/nhess-7-57-2007>
- Papadopoulos, G.A., Daskalaki, E., Fokaefs, A., & Giraleas, N. (2010). Tsunami hazard in the Eastern Mediterranean Sea: Strong earthquakes and tsunamis in the West Hellenic arc and Trench System. *Journal of Earthquake and Tsunami*, 4(3), 145-179.
- Papanikolaou, D. (2010). Major paleogeographic, tectonic and geodynamic changes from the last stage of the Hellenides to the actual Hellenic arc and trench system. *Bulletin of the Geological Society of Greece*, 43(1), 72-85. <https://doi.org/10.12681/bgsg.11161>
- Papanikolaou, D.I. (2021). The Mediterranean. In: *The Geology of Greece. Regional Geology Reviews*. Springer, Cham. https://doi.org/10.1007/978-3-030-60731-9_3
- Papanikolaou, D.J., & Royden, L.H. (2007). Disruption of the Hellenic arc: Late Miocene extensional detachment faults and steep Pliocene-Quaternary normal faults—Or what happened at Corinth? *Tectonics*, 26, TC5003. <https://doi.org/10.1029/2006TC002007>
- Papazachos B.C., & Comninakis P.E. (1971). Geophysical and tectonic features of the Aegean Arc. *Journal of Geophysical Research*, 76(8517).

- Papazachos, B.C., & Dimitriu, P.P. (1991). Tsunamis in and near Greece and their relation to the earthquake focal mechanisms. *Natural Hazards*, 4, 161–170.
<https://doi.org/10.1007/BF00162785>
- Papazachos B.C., Karakostas, V.G., Papazachos, C.B., & Scordilis, E.M. (2000). The geometry of the Wadati–Benioff zone and lithospheric kinematics in the Hellenic arc. *Tectonophysics*, 319(4), 275–300. [https://doi.org/10.1016/S0040-1951\(99\)00299-1](https://doi.org/10.1016/S0040-1951(99)00299-1)
- Papazachos, C.B. (2019). Deep Structure and Active Tectonics of the South Aegean Volcanic Arc. *Elements*, 15(3), 153–158. <https://doi.org/10.2138/gselements.15.3.153>
- Papazachos, C.B., & Nolet, G. (1997). P and S deep velocity structure of the Hellenic area obtained by robust nonlinear inversion of travel times. *Journal of Geophysical Research*, 102(B4), 8349–8367.
- Pearce, F.D., Rondenay, S., Sachpazi, M., Charalampakis, M., & Royden, L. H. (2012). Seismic investigation of the transition from continental to oceanic subduction along the western Hellenic Subduction Zone. *Journal of Geophysical Research*, 117, B07306, <https://doi.org/10.1029/2011JB009023>
- Peterek, A., & Schwarze, J. (2004). Architecture and Late Pliocene to recent evolution of outer-arc basins of the Hellenic subduction zone (south-central Crete, Greece). *Journal of Geodynamics*, 38(1), 19–55. <https://doi.org/10.1016/j.jog.2004.03.002>
- Peters, K., Smit, M., van Hinsbergen, D.J.J., van Roermund, H., & Brouwer, F. (2017). The longevity of Neotethyan metamorphic soles from Lu–Hf garnet chronology. 19th EGU General Assembly, EGU2017, Vienna, Austria., p.15673.
<https://ui.adsabs.harvard.edu/abs/2017EGUGA..1915673P/abstract>
- Philippon, M., Brun, J.-P., Gueydan, F., & Sokoutis, D. (2014). The interaction between Aegean back-arc extension and Anatolia escape since Middle Miocene, *Tectonophysics*, 631, 176–188. <https://doi.org/10.1016/j.tecto.2014.04.039>
- Piromallo, C., & Morelli, A. (1997). Imaging the Mediterranean upper mantle by P-wave travel time tomography. *Annali Di Geofisica*, XL(4), 963–979. <https://doi.org/10.4401/ag-3890>
- Piromallo, C., & Morelli, A. (2003). P wave tomography of the mantle under the Alpine-Mediterranean area. *Journal of Geophysical Research*, 108, 2065.
<https://doi.org/10.1029/2002JB001757>.
- Plunder, A., Agard, P., Chopin, C., & Okay, A.I. (2013). Geodynamics of the Tavşanlı zone, western Turkey: Insights into subduction/obduction processes, *Tectonophysics*, 608, 884–903. <https://doi.org/10.1016/j.tecto.2013.07.028>
- Polat, G., & Ozel, M.N. (2007). Deep azimuthal seismic anisotropy in the western Anatolia and Aegean subduction zone. In Ubertini, L., Manciola, P., Casadei, S., Grimaldi, S. (Eds.) *International Union of Geodesy and Geophysics General Assembly*, 24, http://kocaeli2007.kocaeli.edu.tr/kocaeli2007/TAM_METIN_NUMARALI-SIRALI-PDF/141-145.pdf.
- Portner, D.E., Delph, J.R., Biryol, C.B., Beck, S.L., Zandt, G., Özacar, A.A., Sandvol, E., & Türkelli, N. (2018). Subduction termination through progressive slab deformation across Eastern Mediterranean subduction zones from updated P-wave tomography beneath Anatolia. *Geosphere*, 14(3), 907–925. <https://doi.org/10.1130/GES01617.1>
- Pourteau, A., Candan, O., & Oberhänsli, R. (2010). High-pressure metasediments in central Turkey: Constraints on the Neotethyan closure history. *Tectonics*, 29, 1–18.
- Pourteau, A., Oberhänsli, R., Candan, O., Barrier, E., & Vrielynck, B. (2016). Neotethyan closure history of western Anatolia: a geodynamic discussion. *Geologische Rundschau* =

- International Journal of Earth Sciences, 105, 203–224. <https://doi.org/10.1007/s00531-015-1226-7>
- Pourteau, A., Scherer, E. E., Schorn, S., Bast, R., Schmidt, A., & Ebert, L. (2019). Thermal evolution of an ancient subduction interface revealed by Lu–Hf garnet geochronology, Halilbağı Complex (Anatolia). *Geoscience Frontiers*, 10(1), 127–148.
- Pourteau, A., Sudo, M., Candan, O., Lanari, P., Vidal, O., & Oberhaensli, R. (2013). Neotethys closure history of Anatolia; insights from ⁴⁰Ar–³⁹Ar geochronology and P–T estimation in highpressure metasedimentary rocks. *Journal of Metamorphic Geology*, 31(6), 585–606.
- Rabayrol, F., & Hart, C.J.R. (2021). Petrogenetic and tectonic controls on magma fertility and the formation of post-subduction porphyry and epithermal mineralization along the late Cenozoic Anatolian Metallogenic Trend, Turkey. *Mineralium Deposita*, 56, 279–306. <https://doi.org/10.1007/s00126-020-00967-9>
- Rabayrol, F., Hart, C.J.R., & Creaser, R.A. (2019). Tectonic Triggers for Postsubduction Magmatic-Hydrothermal Gold Metallogeny in the Late Cenozoic Anatolian Metallogenic Trend, Turkey. *Economic Geology*, 114 (7), 1339–1363. <https://doi.org/10.5382/econgeo.4682>
- Reilinger, R., et al. (2006). GPS constraints on continental deformation in the Africa-Arabia-Eurasia continental collision zone and implications for the dynamics of plate interactions, *Journal of Geophysical Research*, 111, B05411. <https://doi.org/10.1029/2005JB004051>
- Reilinger, R., McClusky, S., Paradissis, D., Ergintav, S., & Vernant, P. (2010). Geodetic constraints on the tectonic evolution of the Aegean region and strain accumulation along the Hellenic subduction zone. *Tectonophysics*, 488(1–4), 22–30. <https://doi.org/10.1016/j.tecto.2009.05.027>
- Robertson, A.H.F., & Ustaömer, T. (2004). Tectonic evolution of the Intra-Pontide suture zone in the Armutlu Peninsula, NW Turkey. *Tectonophysics*, 381(1–4), 175–209. <https://doi.org/10.1016/j.tecto.2002.06.002>
- Romanowicz, B. (2003). 3D structure of the Earth's lower mantle. *Comptes Rendus Geoscience*, 335(1), 23–35. [https://doi.org/10.1016/S1631-0713\(03\)00012-9](https://doi.org/10.1016/S1631-0713(03)00012-9)
- Rotstein, &, & Ben-Avraham, Z. (1985). Accretionary processes at subduction zones in the eastern Mediterranean. *Tectonophysics*, 112, 551–561.
- Royden, L.H. (1993). The tectonic expression slab pull at continental convergent boundaries, *Tectonics*, 12(2), 303– 325. <https://doi.org/10.1029/92TC02248>
- Royden L.H., & Husson L. (2009). Subduction with Variations in Slab Buoyancy: Models and Application to the Banda and Apennine Systems. In S. Lallemand, F. Funiciello (Eds.), *Subduction Zone Geodynamics. Frontiers in Earth Sciences*. Springer, Berlin, Heidelberg. https://doi.org/10.1007/978-3-540-87974-9_2
- Royden, L.H., & Papanikolaou, D.J. (2011). Slab segmentation and late Cenozoic disruption of the Hellenic arc. *Geochemistry, Geophysics, Geosystems, AGU and the Geochemical Society*, 12(3), <https://doi.org/10.1029/2010GC003280>
- Ruff, L., Kanamori, H. (1983). Seismic coupling and uncoupling at subduction zones. *Tectonophysics*, 99(2–4), 99–117. [https://doi.org/10.1016/0040-1951\(83\)90097-5](https://doi.org/10.1016/0040-1951(83)90097-5)
- Sachpazi, M., Laigle, M., Charalampakis, M., Diaz, J., Kissling, E., Gesret, A., Becel, A., Flueh, E., Miles, P., & Hirn, A. (2016). Segmented Hellenic slab rollback driving Aegean deformation and seismicity, *Geophysical Research Letters*, 43, 65–658. <https://doi.org/10.1002/2015GL066818>

- Salaün, G., Pedersen, H.A., Paul, A., Farra, V., Karabulut, H., Hatzfeld, D., Papazachos, C., Childs, D.M., Pequegnat, C., & SIMBAAD Team (2012). High-resolution surface wave tomography beneath the Aegean-Anatolia region: constraints on upper-mantle structure. *Geophysical Journal International*, 190(1), 406–420. <https://doi.org/10.1111/j.1365-246X.2012.05483.x>
- Saunders, P., Priestley, K., & Taymaz, T. (1998). Variations in the crustal structure beneath western Turkey. *Geophysical Journal International*, 134(2), 373–389. <https://doi.org/10.1046/j.1365-246x.1998.00571.x>
- Savostin, L.A., Sibuet, J-C., Zonenshain, L.P., Le Pichon, X., & Roulet, M-J. (1986). Kinematic evolution of the Tethys belt from the Atlantic ocean to the pamirs since the Triassic, *Tectonophysics*, 123(1-4), 1-35. [https://doi.org/10.1016/0040-1951\(86\)90192-7](https://doi.org/10.1016/0040-1951(86)90192-7)
- Sayit, K., Marroni, M., Göncüoğlu, M.C., Pandolfi, L., Ellero, A., Ottria, G., & Frassi, C. (2016). Geological setting and geochemical signatures of the mafic rocks from the Intra-Pontide Suture Zone: implications for the geodynamic reconstruction of the Mesozoic Neotethys. *Geologische Rundschau = International Journal of Earth Sciences*, 105(1), 39-64.
- Schellart, W., Freeman, J., Stegman, D. et al. (2007). Evolution and diversity of subduction zones controlled by slab width. *Nature*, 446, 308–311. <https://doi.org/10.1038/nature05615>
- Schivardi R., & Morelli A. (2009). Surface wave tomography in the European and Mediterranean region. *Geophysical Journal International*, 177(3), 1050–1066.
- Scholl, D.W., & von Huene, R. (2009). Implications of estimated magmatic additions and recycling losses at the subduction zones of accretionary (non-collisional) and collisional (suturing) orogens. *Geological Society, London, Special Publications*, 318, 105-125, <https://doi.org/10.1144/SP318.4>
- Scholz, C.H. (1990). *The mechanics of earthquakes and faulting*, Cambridge, Cambridge University press, 439pp
- Scholz, C. H., and Campos, J. (2012), The seismic coupling of subduction zones revisited, *J. Geophys. Res.*, 117, B05310, <https://doi.org/10.1029/2011JB009003>
- Scoon R.N. (2021). Volcanoes of the Hellenic Volcanic Arc, Greece. In: *The Geotraveller*. Springer, Cham. https://doi.org/10.1007/978-3-030-54693-9_12
- Şengör, A.M.C., & Yılmaz, Y. (1981). Tethyan evolution of Turkey: A plate tectonic approach. *Tectonophysics*, 75, 181–241.
- Şengör, A.M.C., & Zabcı, C. (2019). The North Anatolian Fault and the North Anatolian Shear Zone. In: *Landscapes and Landforms of Turkey*, Kuzucuoğlu, C., Çiner, A., Kazancı, N. (Eds.), Springer International Publishing, *World Geomorphological Landscapes*. https://doi.org/10.1007/978-3-030-03515-0_27
- Şengün, F., Yiğitbaş, E., Tunç, I.O. (2011). Geology and tectonic emplacement of eclogite and blueschists, Biga Peninsula, northwest Turkey. *Turkish Journal of Earth Sciences*, 20, 273–285.
- Shaw, B., & Jackson, J. (2010). Earthquake mechanisms and active tectonics of the Hellenic subduction zone. *Geophysical Journal International*, 181, 966-984. <https://doi.org/10.1111/j.1365-246X.2010.04551.x>
- Shaw, B., Ambraseys, N., England, P. et al. (2008). Eastern Mediterranean tectonics and tsunami hazard inferred from the AD 365 earthquake. *Nature Geoscience*, 1, 268–276. <https://doi.org/10.1038/ngeo151>

- Shephard, G.E., Matthews, K.J., Hosseini, K., & Domeier, M. (2017). On the consistency of seismically imaged lower mantle slabs. *Scientific Reports*, 7, 10976. <https://doi.org/10.1038/s41598-017-11039-w>
- Sherlock, S., Kelley, S., Inger, S., Harris, N., & Okay, A. (1999). ^{40}Ar - ^{39}Ar and Rb-Sr geochronology of high-pressure metamorphism and exhumation history of the Tavşanlı Zone, NW Turkey. *Contributions to Mineralogy and Petrology*, 137, 46–58.
- Shin, T.A., Catlos, E.J., Jacob, L., & Black, K. (2013). Relationships between very high pressure subduction complex assemblages and intrusive granitoids in the Tavşanlı Zone, Sivrihisar Massif, central Anatolia. *Tectonophysics*, 595–596, 183–197.
- Sigl, M., Winstrup, M., McConnell, J. et al. (2015). Timing and climate forcing of volcanic eruptions for the past 2,500 years. *Nature* 523, 543–549. <https://doi.org/10.1038/nature14565>
- Silver, P.G., Russo, R.M., Lithgow-Bertelloni, C. (1998). Coupling of South American and African Plate Motion and Plate Deformation. *Science*, 279, 60-63.
- Snieder, R. (1988). Large-scale waveform inversions of surface waves for lateral heterogeneity: 2. Application to surface waves in Europe and the Mediterranean. *Journal of Geophysical Research-Solid Earth*, 93, 12067–12080.
- Snopek, K., Meier, T., Endrun, B., Bohnhoff, M., & Casten, U. (2007). Comparison of gravimetric and seismic constraints on the structure of the Aegean lithosphere in the forearc of the Hellenic subduction zone in the area of Crete. *Journal of Geodynamics*, 44(3–5), 173-185. <https://doi.org/10.1016/j.jog.2007.03.005>.
- Sodoudi, F., Kind, R., Hatzfeld, D., Priestley, K., Hanka, W., Wylegalla, K., Stavrakakis, G., Vafidis, A., Harjes, H.-P., & Bohnhoff, M. (2006). Lithospheric structure of the Aegean obtained from P and S receiver functions, *Journal of Geophysical Research*, 111, B12307. <https://doi.org/10.1029/2005JB003932>
- Solheim, L.P., & Peltier, W.R. (1994). Avalanche effects in phase transition modulated thermal convection: a model of Earth's mantle. *Journal of Geophysical Research*, 99, 6997-7018.
- Spakman, W. (1990). Tomographic images of the upper mantle below central Europe and the Mediterranean. *Terra Nova*, 2, 542-553. <https://doi.org/10.1111/j.1365-3121.1990.tb00119.x>
- Spakman, W. (1991). Delay-time tomography of the upper mantle below Europe, the Mediterranean, and Asia Minor, *Geophysical Journal International*, 107(2), 309–332. <https://doi.org/10.1111/j.1365-246X.1991.tb00828.x>
- Spakman, W., van der Lee, S., & van der Hilst, R. (1993). Travel-time tomography of the European-Mediterranean mantle down to 1400 km. *Physics of the Earth and Planetary Interiors*, 79(1–2), 3-74. [https://doi.org/10.1016/0031-9201\(93\)90142-V](https://doi.org/10.1016/0031-9201(93)90142-V)
- Spakman, W., Wortel, M.J.R., & Vlaar, N.J. (1988). The Hellenic subduction zone: A tomographic image and its geodynamic implications. *Geophysical Research Letters*, 15, 60–63.
- Speciale, P.A., Catlos, E.J., Yıldız, G.O., Shin, T.A., & Black, K.N. (2012). Zircon ages from the Beypazari granitoid pluton (north central Turkey): tectonic implications. *Geodinamica Acta*, 25(3-4), 162-182. <https://doi.org/10.1080/09853111.2013.858955>.
- Stampfli, G.M. (2000). Tethyan oceans. In E. Bozkurt, J.A. Winchester, J.D.A. Piper (Eds.), *Tectonics and magmatism in Turkey and surrounding area*. Geological Society of London, Special Publication, London (Vol. 173, pp. 1-23). <https://doi.org/10.1144/GSL.MEM.2006.032.01.04>

- Stampfli, G. M., & Borel, G.D. (2002). A plate tectonic model for the Paleozoic and Mesozoic constrained by dynamic plate boundaries and restored synthetic oceanic isochrons. *Earth and Planetary Science Letters*, 196, 17–33. [https://doi.org/10.1016/S0012-821X\(01\)00588-X](https://doi.org/10.1016/S0012-821X(01)00588-X)
- Stampfli, G.M., & Kozur, H.W. (2006). Europe from the Variscan to the Alpine cycles. *Geological Society, London, Memoirs*, 32, 57-82.
- Stern, R.J. (2004). Subduction initiation: spontaneous and induced. *Earth and Planetary Science Letters*, 226, 275-292.
- Stern, R.J., & Gerya, T. (2018). Subduction initiation in nature and models: A review. *Tectonophysics*, 746, 173-198. <https://doi.org/10.1016/j.tecto.2017.10.014>
- Stöckhert B. (1999). The Hellenic subduction zone, a world site to study the mechanics of roll back., CIESM Workshop Series 6, Workshop on Mediterranean Scientific Drilling Prospectives, Granada 10–12 June 1999, 103–108.
- Suckale, J., Rondenay, S., Sachpazi, M., Charalampakis, M., Hosa, A., Royden, L.H. (2009). High-resolution seismic imaging of the western Hellenic subduction zone using teleseismic scattered waves, *Geophysical Journal International*, 178(2), 775–791. <https://doi.org/10.1111/j.1365-246X.2009.04170.x>
- Symeou, V., Homberg, C., Nader, F. H., Darnault, R., Lecomte, J.-C., & Papadimitriou, N. (2018). Longitudinal and temporal evolution of the tectonic style along the Cyprus Arc system, assessed through 2-D reflection seismic interpretation. *Tectonics*, 37, 30– 47. <https://doi.org/10.1002/2017TC004667>
- Taylor, G., Thompson, D.A., Cornwell, D., & Rost, S. (2019). Interaction of the Cyprus/Tethys slab with the mantle transition zone beneath Anatolia, *Geophysical Journal International*, 216(3), 1665–1674. <https://doi.org/10.1093/gji/ggy514>
- Taymaz, T., Jackson, J., & Westaway, R. (1990). Earthquake mechanisms in the Hellenic Trench near Crete. *Geophysics Journal International*, 102, 695-731.
- Taymaz, T., Yilmaz, Y., & Dilek, Y. (2007). The geodynamics of the Aegean and Anatolia: introduction. *Geological Society, London, Special Publications*, 291(1-16). <https://doi.org/10.1144/SP291.1>
- Tekin, U.K., Göncüoğlu, M.C., & Turhan, N. (2002). First evidence of Late Carnian radiolarians from the Izmir–Ankara suture complex, central Sakarya, Turkey: implications for the opening age of the Izmir–Ankara branch of Neo-Tethys. *Geobios*, 35(1), 127-135. [https://doi.org/10.1016/S0016-6995\(02\)00015-3](https://doi.org/10.1016/S0016-6995(02)00015-3)
- ten Veen, J. H., & Kleinspehn, K. L. (2002). Geodynamics along an increasingly curved convergent plate margin: Late Miocene-Pleistocene Rhodes, Greece, *Tectonics*, 21(3), <https://doi.org/10.1029/2001TC001287>
- Tezel, T., Shibusani, T., & Kaypak, B. (2013). Crustal thickness of Turkey determined by receiver function. *Journal of Asian Earth Sciences*, 75, 36-45. <https://doi.org/10.1016/j.jseaes.2013.06.016>
- Thomson S.N., Stöckhert N., & Brix M.R. (1998). Thermochronology of the high-pressure metamorphic rocks of Crete, Greece: implications for the speed of tectonic processes. *Geology*, 26, 259–262.
- Tinti, S., Armigliato, A., Pagnoni, G., & Zaniboni, F. (2005). Scenarios of giant tsunamis of tectonic origin in the Mediterranean. *ISET Journal of Earthquake Technology*, Paper No. 464, 42(4), 171-188.

- Tirel, C., Gueydan, F., Tiberi, C., & Brun, J-P. (2004). Aegean crustal thickness inferred from gravity inversion. *Geodynamical implications. Earth and Planetary Science Letters*, 228(3–4), 267–280. <https://doi.org/10.1016/j.epsl.2004.10.023>.
- Triantafyllou, I., Novikova, T., Charalampakis, M. et al. (2019). Quantitative tsunami risk assessment in terms of building replacement cost based on tsunami modelling and GIS methods: The case of Crete Isl., Hellenic Arc. *Pure Applied Geophysics*, 176, 3207–3225. <https://doi.org/10.1007/s00024-018-1984-9>
- Ustaömer, T., Ustaömer, P.A., Robertson, A.H.F., & Gerdes, A. (2020). U-Pb-Hf isotopic data from detrital zircons in late Carboniferous and Mid-Late Triassic sandstones, and also Carboniferous granites from the Tauride and Anatolide continental units in S Turkey: implications for Tethyan palaeogeography. *International Geology Review*, 62(9), 1159–1186. <https://doi.org/10.1080/00206814.2019.1636415>
- van der Meer, D.G., van Hinsbergen, D.J.J., & Spakman, W. (2018). Atlas of the underworld: Slab remnants in the mantle, their sinking history, and a new outlook on lower mantle viscosity. *Tectonophysics*, 723, 309–448. <https://doi.org/10.1016/j.tecto.2017.10.004>.
- van der Meijde, M., Marone, F., Giardini, D., & van der Lee, S. (2003). Seismic evidence for water deep in Earth's upper mantle. *Science*, 300, 1556–1558.
- van Hinsbergen, D.J.J., Hafkenscheid, E., Spakman, W., Meulenkaamp, J.E., & Wortel, R. (2005). Nappe stacking resulting from subduction of oceanic and continental lithosphere below Greece. *Geology*, 33 (4), 325–328. <https://doi.org/10.1130/G20878.1>
- van Hinsbergen, D.J.J., Kaymakçi, N., Spakman, W., & Torsvik, T.H. (2010). Reconciling the geological history of western Turkey with plate circuits and mantle tomography, *Earth and Planetary Science Letters*, 297(3–4), 674–686.
- van Hinsbergen, D.J.J., & Schmid, S.M. (2012). Map view restoration of Aegean–West Anatolian accretion and extension since the Eocene, *Tectonics*, 31, TC5005. <https://doi.org/10.1029/2012TC003132>
- van Hinsbergen, D.J.J., Steinberger, B., Guilmette, C. et al. (2021). A record of plume-induced plate rotation triggering subduction initiation. *Nature Geoscience*, 14, 626–630. <https://doi.org/10.1038/s41561-021-00780-7>
- Vanderhaeghe, O., Hirsch, C., Siebenaller, L., Duchêne, S., de St Blanquat, M., Kruckenberg, S., Fotiadis, A., & Martin, L. (2007). Penrose Conference - Extending a Continent - Naxos Field Guide. In: (Eds.) Gordon Lister, Marnie Forster, and Uwe Ring, *Inside the Aegean Metamorphic Core Complexes*, *Journal of the Virtual Explorer*, 27(4). <https://doi.org/10.3809/jvirtex.2007.00175>
- Vaněk, J., Hanuš, V., Christova, C., & Simeonova, S. (1987). Morphology of the wadati-benioff zone in the hellenic arc. *Journal of Geodynamics*, 8(1), 79–93. [https://doi.org/10.1016/0264-3707\(87\)90059-7](https://doi.org/10.1016/0264-3707(87)90059-7)
- Vassilakis, E., Royden, L., & Papanikolaou, D. (2011). Kinematic links between subduction along the Hellenic trench and extension in the Gulf of Corinth, Greece: A multidisciplinary analysis. *Earth and Planetary Science Letters*, 303(1–2), 108–120.
- Ventouzi, C., Papazachos, C., Hatzidimitriou, P., Papaioannou, C., & EGELADOS Working Group, (2018). Anelastic P- and S- upper mantle attenuation tomography of the southern Aegean Sea subduction area (Hellenic Arc) using intermediate-depth earthquake data. *Geophysical Journal International*, 215(1), 635–658. <https://doi.org/10.1093/gji/ggy292>

- Vernant, P., Reilinger, R., & McClusky, S. (2014). Geodetic evidence for low coupling on the Hellenic subduction plate interface. *Earth and Planetary Science Letters*, 385, 122-129. <https://doi.org/10.1016/j.epsl.2013.10.018>.
- Von Huene R., & Scholl D.W. (1991). Observations at convergent margins concerning sediment subduction, subduction erosion, and the growth of continental-crust. *Reviews in Geophysics*, 29(3), 279-316.
- Von Huene R., & Scholl D.W. (1993). The return of sialic material to the mantle indicated by terrigenous material subducted at convergent margins. *Tectonophysics*, 219(1-3), 163-175/
- Wallace, L.M., Ellis, S., & Mann, P. (2008). Tectonic block rotation, arc curvature, and back-arc rifting: insights into these processes in the Mediterranean and the western Pacific. *IOP Conference Series: Earth and Environmental Sciences*, 2, 012010. <https://doi.org/10.1088/1755-1307/2/1/012010>
- Wallace, L.M., McCaffrey, R., Beavan, J., & Ellis S. (2005). Rapid microplate rotations and backarc rifting at the transition between collision and subduction. *Geology*, 33 (11), 857-860. <https://doi.org/10.1130/G21834.1>
- Wei, W., Zhao, D., Wei, F., Bai, X., & Xu, J. (2019). Mantle dynamics of the Eastern Mediterranean and Middle East: Constraints from P-wave anisotropic tomography. *Geochemistry, Geophysics, Geosystems*, 20, 4505-4530. <https://doi.org/10.1029/2019GC008512>
- Westaway, R. (1994). Present-day kinematics of the Middle East and eastern Mediterranean, *Journal of Geophysical Research*, 99(B6), 12071-12090. <https://doi.org/10.1029/94JB00335>
- Westbrook, G.K., & Reston, T.J. (2002). The accretionary complex of the Mediterranean Ridge: tectonics, fluid flow and the formation of brine lakes -an introduction to the special issue of *Marine Geology*. *Marine Geology*, 186, 1-8.
- White, D.A., Roeder, D.H., Nelson, T.H., & Crowell, J.C. (1970). Subduction. *Geological Society of America Bulletin*, 81, 3431-3432. [https://doi.org/10.1130/0016-7606\(1970\)81\[3431:S\]2.0.CO;2](https://doi.org/10.1130/0016-7606(1970)81[3431:S]2.0.CO;2)
- Widiyantoro, S., van der Hilst, R.D., & Wenzel F. (2004). Deformation of the Aegean Slab in the Mantle Transition Zone. *International Journal of Tomography & Statistics*, D04, 1-14.
- Wolfenden, E., Ebinger, C., Yirgu, G., Deino, A., & Ayalew, D. (2004). Evolution of the northern Main Ethiopian rift: birth of a triple junction. *Earth and Planetary Science Letters*, 224(1-2), 213-228. <https://doi.org/10.1016/j.epsl.2004.04.022>
- Wolfenden, E., Ebinger, C., Yirgu, G., Renne, P.R., & Kelley, S.P. (2005). Evolution of a volcanic rifted margin: Southern Red Sea, Ethiopia. *Geological Society of America Bulletin*, 117 (7-8), 846-864. <https://doi.org/10.1130/B25516.1>
- Woodside, J.M., Mascle, J., Huguen, C., & Volkonskaia, A. (2000). The Rhodes Basin, a post-Miocene tectonic trough. *Marine Geology*, 165, 1-12.
- Woodside, J.M., Mascle, J., Zitter, T.A.C., Limonov, A.F., Ergün, M., & Volkonskaia, A. (2002). The Florence Rise, the Western Bend of the Cyprus Arc. *Marine Geology*, 185(3-4), 177-194. [https://doi.org/10.1016/S0025-3227\(02\)00194-9](https://doi.org/10.1016/S0025-3227(02)00194-9)
- Wortel, M.J.R., & Spakman, W. (1992). Structure and dynamics of subducted lithosphere in the Mediterranean region. *Proceedings of the Koninklijke Nederlandse Akademie van Wetenschappen*, 95(3), 325-347.

- Wortel, M.J.R., & Spakman, W. (2000). Subduction and slab detachment in the Mediterranean-Carpathian region. *Science*, 290, 1910-1917. <https://doi.org/10.1126/science.290.5498.1910>
- Wortel, M.J.R., Goes, S.D.B., Spakman, W. (1990). Structure and seismicity of the Aegean subduction zone. *Terra Nova*, 2, 554-562.
- Yang, T., Gurnis, M., & Zahirovic, S. (2018). Slab avalanche-induced tectonics in self-consistent dynamic models. *Tectonophysics*, 746, 251-265. <https://doi.org/10.1016/j.tecto.2016.12.007>
- Yiğitbaş, E., Kerrich, R., Yılmaz, Y., Elmas, A., & Xie, Q. (2004). Characteristics and geochemistry of Precambrian ophiolites and related volcanics from the Istanbul–Zonguldak Unit, Northwestern Anatolia, Turkey: following the missing chain of the Precambrian South European suture zone to the east, *Precambrian Research*, 132(1-2), 179-206. <https://doi.org/10.1016/j.precamres.2004.03.003>
- Yılmaz, Y., Genc, C., Karacik, Z., & Altunkaynak, S. (2001). Two contrasting magmatic associations of NW Anatolia and their tectonic significance. *Journal of Geodynamics*, 31(3), 243-271.
- Yılmaz, Y., Tüysüz, O., Yiğitbaş, E., Genç, C., & Şengör, A.M.C. (1997). Geology and tectonic evolution of the Pontides. In A.G. Robinson (Ed.), *Regional and petroleum geology of the Black Sea and surrounding region*. American Association of Petroleum Geology Memiors (Vol. 68, pp. 183-226), Tulsa, Oklahoma.
- Yolsal, S., Taymaz, T., & Yalçiner, A.C. (2007). Understanding tsunamis, potential source regions and tsunami-prone mechanisms in the Eastern Mediterranean. *Geological Society, London, Special Publications*, 291, 201-230. <https://doi.org/10.1144/SP291.10>
- Yoshioka, T. (1996). Evolution of fault geometry and development of strike-slip basins: Comparative studies on the transform zones in Turkey and Japan. *Island Arc*, 5, 407-419. <https://doi.org/10.1111/j.1440-1738.1996.tb00162.x>
- Zanchetta G, Sulpizio R, Roberts N, et al. (2011). Tephrostratigraphy, chronology and climatic events of the Mediterranean basin during the Holocene: An overview. *The Holocene*, 21(1), 33-52. <https://doi.org/10.1177/0959683610377531>
- Zhu, L., Mitchell, B. J., Akyol, N., Çemen, I., & Kekovali, K. (2006). Crustal thickness variations in the Aegean region and implications for the extension of continental crust, *Journal of Geophysical Research*, 111, B01301. <https://doi.org/10.1029/2005JB003770>
- Zielhuis, A., & Nolet G. (1994). Shear-wave velocity variations in the upper-mantle beneath Central Europe. *Geophysics Journal International*, 117(3), 695–715.
- Zitter, T.A.C., Woodside, J.M., & Mascle, J. (2003). The Anaximander Mountains: a clue to the tectonics of southwest Anatolia. *Geological Journal*, 38, 375-394. <https://doi.org/10.1002/gj.961>




Figure 1. EMODnet Digital Bathymetry maps with some structures overlain. The circular inset shows higher magnification of the region as indicated by the arrow. The Aegean and Anatolian microplate boundaries are shown in grey after Nyst & Thatcher (2004). Other structures after Hall et al. (1984) and (2009), Woodside et al. (2002), Peterek & Schwarze (2004), Meier et al. (2007), Kinnaird & Robertson (2012), and Symeou et al. (2018). Abbreviations: AM= Anaximander Mountains; BT= Backthrust; ES = Eratosthenes Seamount; IAESZ = Izmir-Ankara-Erzincan Suture Zone; IPS= Intra-Pontide Suture; KFZ = Kephallonia Fault Zone; KM= Kırşehir Massif; MP = Methana Peninsula; MM= Menderes Massif; NHSZ = North Hellenic Shear Zone; PTF = Paphos Transform Fault; SHSZ = South Hellenic Shear Zone. See supplementary documents for this figure in color.




Figure 2. North-south generalized cross-section through the Hellenic arc system showing the key structural elements. Map of the Mediterranean Ridge after Westbrook & Reston (2002).




Figure 3. Map of the geometry of the Hellenic and part of the Cyprus subduction zones after Hayes (2018). The figure shows both the depth (up to 440 km) and strike of the subducted African plate (African, Nubian, Aegean, or Hellenic slab) based on the Slab2 model. Overlain in red are fault systems from NOAFaults, a digital database of active faults in Greece (Ganas et al., 2013). Countries and some features are indicated. See supplementary documents for this figure in color.




Figure 4. (A) Depth vs. latitude of earthquakes taken from a line of the section of 28°-43° and longitude of 24°-28°. Events were extracted from the Turkish Ministry of the Interior, Disaster and Emergency Management Presidency, Earthquake Department Earthquake Catalog ($M \geq 4.0$), 1900-20XX (<https://deprem.afad.gov.tr/depremkatalogu>) from 01/24/1900 to 6/17/2021. We indicate the largest event (6/26/1926, 19:46). The legend shows how the size of the earthquake correlates to the symbol. (B) Cross-section of the Aegean anomaly interpreted as the African slab using the UUP07 P-wave model (Amaru, 2007). The depths of the dashed lines are 410, 660, 1000 km from the surface. Interpretations of the geology below 1000 are debated and discussed in the text. Image created using Hosseini et al. (2018). See text for references to additional tomographic images.




Figure 5. Geological map of Western Anatolia focusing on the ophiolite and granite assemblages along the boundary between the Aegean and Anatolia microplates. Plate boundary after Nyst & Thatcher (2004). Terrane boundaries, major fault systems, and transfer zones after Okay (2008), Akbayram et al. (2016), Oner et al. (2010), and Karaoğlu & Helvacı (2014). Abbreviations: RPA= Rhodope -Pontide Arc; İBTZ = Izmir –Balıkesir Transfer Zone (also sometimes referred to as the Western Anatolia Transfer Zone, Gessner et al., 2013; 2017); SWASZ= South West Anatolian Shear Zone; IPS= Intra -Pontide suture zone; IAESZ = Izmir -Ankara -Erzincan suture zone; PS = Pamphylian suture zone; UMTZ= Uşak -Muğla Transfer Zone.




Figure 6. North-south cross section across the Aegean region from c. 95 Ma to <2 Ma. In this scenario, the Hellenic arc forms due to the transition from a divergent plate boundary to convergence sometime between 54 Ma and 25 Ma (Dilek & Altunkaynak, 2007). Scale is only a rough approximation. See Dilek & Altunkaynak (2007) for a color image of this figure. Abbreviations: ATC= Anatolide-Tauride Continent; IASZ= Izmir-Ankara Suture Zone; IPSZO= Intra-Pontide Suture Zone ophiolites; MG= Magmatic Granites; NAFZ= North Anatolian Fault Zone; SC = Sakarya Continent; SSZ = supra-subduction zone; SZG = Suture Zone Granitoids; RPC= Rhodope-Pontide Continent.




Figure 7. Lithospheric-scale cross sections of a model of the Aegean region through Western Anatolia in which the region records and experiences progressively younger subduction zones after Shin et al. (2013). The subduction zone cartoon is adapted from Cloos et al. (2005). Scale is only a rough approximation. Panel (A) shows the development of the IAESZ during the Early to Late Cretaceous. Panel (B) records the peak metamorphism of the zone and the development of the Afyon Zone subduction zone at ~85 Ma. Panel (C) shows granitoid arc magmas from the Afyon Zone intrude the IAESZ at ~45 Ma, emplacing granite bodies that intrude the ophiolite sequences. Panel (D) records the break off of the IAESZ slab and additional magma emplacement at 30 Ma. In this scenario, the Hellenic Arc experiences ongoing subduction.




Figure 8. Schematic overview of development of nappe stack and subduction during Alpine orogeny in Greece from van Hinsbergen et al. (2005). The cross-section displays the single subduction zone model for the development of the Hellenic arc. See van Hinsbergen et al. (2005) for a color version of this figure.