

Effects of snow and remineralization processes on nutrient distributions in multi-year Antarctic landfast sea ice

Reishi Sahashi¹, Daiki Nomura¹, Takenobu Toyota¹, Manami Tozawa¹, Masato Ito², Pat Wongpan³, Kazuya Ono¹, Daisuke Simizu², Kazuhiro Naoki⁴, Yuichi Nosaka⁴, Takeshi Tamura², Shigeru Aoki¹, and Shuki Ushio²

¹Hokkaido University

²National Institute of Polar Research

³Australian Antarctic Program Partnership, University of Otago

⁴Tokai University

November 22, 2022

Abstract

We elucidated the effects of snow and remineralization processes on nutrient distributions in multi-year landfast sea ice (fast ice) in Lützow-Holm Bay, East Antarctica. Based on sea-ice salinity, oxygen isotopic ratios, and thin section analyses, we found that the multi-year fast ice grew upward due to the year-by-year accumulation of snow. Compared to ice of seawater origin, nutrient concentrations in shallow fast ice were low due to replacement by clean and fresh snow. In deeper ice of seawater origin (the lower half of the multi-year fast ice column), remineralization was dominated by the degradation of organic matter. In addition, denitrification was detected in the low brine volume fraction and impermeable multi-year ice due to their disconnection from gas and water exchanges with the atmosphere and under-ice water. By comparison with first-year ice, we develop a conceptual image of the evolution of nutrient concentrations, with biological uptake dominating in relatively young ice and remineralization dominating in older, multi-year ice under the physical process of brine drainage.

Effects of snow and remineralization processes on nutrient distributions in multi-year Antarctic landfast sea ice

Reishi Sahashi¹, Daiki Nomura^{2,1,3*}, Takenobu Toyota⁴, Manami Tozawa¹, Masato Ito⁵, Pat Wongpan^{6,4,7}, Kazuya Ono⁴, Daisuke Simizu⁵, Kazuhiro Naoki⁸, Yuichi Nosaka⁹, Takeshi Tamura^{5,10}, Shigeru Aoki⁴, Shuki Ushio^{5,10}

¹*Faculty of Fisheries Sciences, Hokkaido University, Hakodate, Japan*

²*Field Science Center for Northern Biosphere, Hokkaido University, Hakodate, Japan*

³*Arctic Research Center, Hokkaido University, Sapporo, Japan*

⁴*Institute of Low Temperature Science, Hokkaido University, Sapporo, Japan*

⁵*National Institute of Polar Research, Tachikawa, Japan*

⁶*Institute for Marine and Antarctic Studies, University of Tasmania, Hobart, Australia*

⁷*JSPS International Research Fellow, Japan Society for the Promotion of Science, Tokyo, Japan*

⁸*Research and Information Center, Tokai University, Tokyo, Japan*

⁹*School of Biological Sciences, Tokai University, Sapporo, Hokkaido, Japan*

Abstract

We elucidated the effects of snow and remineralization processes on nutrient distributions in multi-year landfast sea ice (fast ice) in Lützow-Holm Bay, East Antarctica. Based on sea-ice salinity, oxygen isotopic ratios, and thin section analyses, we found that the multi-year fast ice grew upward due to the year-by-year accumulation of snow. Compared to ice of seawater origin, nutrient concentrations in shallow fast ice were low due to replacement by clean and fresh snow. In deeper ice of seawater origin (the lower half of the multi-year fast ice column), remineralization was dominated by the degradation of organic matter. In addition, denitrification was detected in the low brine volume fraction and impermeable multi-year ice due to their disconnection from gas and water exchanges with the atmosphere and under-ice water. By comparison with first-year ice, we develop a conceptual image of the evolution of nutrient concentrations, with biological uptake dominating in relatively young ice and remineralization dominating in older, multi-year ice under the physical process of brine drainage.

Plain language summary

Multi-year landfast sea ice (fast ice) is abundant around the coast of Antarctica. Fast ice is an important component of Antarctic coastal marine ecosystems, providing a prolific habitat for ice algal communities. Although nutrients are essential for biological productivity within sea ice, the status of nutrients and processes affecting nutrient concentrations were previously not known for multi-year fast ice. Here, we collected sea-ice cores from multi-year fast ice in Lützow-Holm Bay, East Antarctica, and we analyzed nutrient concentrations using physical and biogeochemical parameters. Nutrient concentrations in the upper parts of the sea ice decreased due to the accumulation of clean (nutrient-free) snow through melting and refreezing processes, contributing to upward ice growth. In deeper parts of the sea ice, nutrient concentrations were greatly affected by biological processes such as remineralization during the degradation of organic matter within sea ice.

Keywords:

Sea ice, snow, nutrients, remineralization, multi-year landfast ice, fast ice, East Antarctica

Key point:

*Antarctic multi-year landfast ice grew upward due to the year-by-year accumulation of snow

*Nutrient concentrations decreased in the upper sea ice due to the replacement by clean snow

*In deeper sea ice, remineralization by degradation of organic matter drove nutrient concentrations

1. Introduction

Multi-year landfast ice (fast ice) is abundant around Syowa Station, Lützow-Holm Bay, East Antarctica (Fraser et al., 2012; 2021; Ushio, 2006). This region receives relatively abundant snowfall and high snow accumulation rates over sea ice (Toyota et al., 2016), forming ice of snow origin such as ‘snow ice’ and ‘superimposed ice’ (Kawamura et al., 1997; Nomura et al., 2012, 2018). In general, snow ice forms under the weight of overlying snow on the sea ice, which depresses the ice–snow interface to below sea level; the subsequent flow of seawater into the snow forms a flooded slush layer that then freezes (Maksym & Jeffries, 2000; Sturm & Massom, 2017). During summer, the snow surface melts in the sunlight and warmer temperatures; the snowmelt then infiltrates the slush layer, decreasing its salinity to well below that of seawater (Nomura et al., 2012, 2018). However, in Lützow-Holm Bay, ice thickening due to the formation of snow ice does not necessarily depress the sea ice–snow interface to below sea level. In this case, when the shallowest snow thaws, the snowmelt permeates the sea ice–snow interface below the freezing point and refreezes, forming superimposed ice (Kawamura et al., 1997; Nomura et al., 2018). The difference between these two types of ice is that snow ice forms from a mixture of snow and seawater, whereas superimposed

ice forms only from snow. Due to the abundance of snow around Syowa Station, it is considered that superimposed ice forms only after sea ice stops sinking under the weight of overlying snow.

Because sea ice forms from seawater, the biogeochemical components in seawater are preserved in sea ice upon its formation. Various changes affect these biogeochemical components, particularly nutrients, after their entrapment in sea ice. For example, NO_3^- , NO_2^- , NH_4^- , and PO_4^{3-} are involved in biological activities such as photosynthesis and remineralization by microorganisms such as bacteria and phytoplankton, notably ice algae, phytoplankton inhabiting the bottommost layers of sea ice (Arrigo et al., 2010; Gleitz et al., 1995). Diatoms, which are considered to be the main primary producers in sea ice, also use $\text{Si}(\text{OH})_4$ to form their frustules (Arrigo et al., 2010; Treguer & De La Rocha, 2013). In addition to the consumption of nutrients by ice algae, high-salinity and high-nutrient brines are discharged beneath the sea ice over time (Fripiat et al., 2017). Nutrients are resupplied by remineralization during the degradation of organic matter within sea ice by heterotrophs and bacteria (Roukaerts et al., 2021; Thomas et al., 1995) as well as by convection through brine channels (Nomura et al., 2009; Vancoppenolle et al., 2010).

It has also been reported that denitrification reactions occur within sea ice (Kaartokallio, 2001; Nomura et al., 2010; Nomura et al., 2018; Rysgaard & Glud, 2004; Rysgaard et al., 2008; Staley & Gosink, 1999). Sea ice can locally become isolated from the atmosphere and under-ice water, resulting in the depletion of oxygen in the ice and promoting denitrification. Previous studies have reported denitrifying bacteria in Antarctic sea ice (Staley & Gosink, 1999) and have suggested or confirmed denitrification reactions in Arctic and sub-Arctic sea ice, e.g., in the Baltic Sea (Kaartokallio, 2001), along the coast of Greenland and Arctic Ocean (Rysgaard & Glud, 2004; Rysgaard et al., 2008), and in the southern Sea of Okhotsk (Nomura et al., 2010).

Primary production is also active in the nutrient-rich, seawater-flooded slush layer formed at the sea-ice surface during the formation of snow ice (Ackley & Sullivan, 1994; Fritsen et al., 1994; Kattner et al., 2004; Nomura et al., 2018; Schnack-Schiel et al., 2001). The snow cover on sea ice therefore plays an important role in increasing the productivity of the sea-ice surface (e.g., Saenz and Arrigo, 2014). When the slush freezes, it becomes snow ice, preserving the biogeochemical formations (Kawamura et al., 1997; Nomura et al., 2018). However, the thick sea ice produced in areas receiving abundant snowfall prevents seawater from flowing along the sea-ice surface, limiting productivity; therefore, snow cover can only increase productivity in the slush layer by so much. Furthermore, whereas snow-free sea ice grows downward due to the freezing of seawater, snow-covered sea ice grows upward due to the formation of snow ice and superimposed ice, such that melting progresses from the bottom of the sea ice (Kawamura et al., 1997). Thus, snow cover greatly affects biogeochemical distributions and circulations in sea ice.

Fast ice plays a crucial role in the biogeochemical cycle and marine ecosystem functions of the Antarctic coast because it hosts well-established habitats (in terms of nutrients, temperature, salinity, and light) for stable primary production (Arrigo, 2017; Wongpan et al., 2018). Because primary production in sea ice promotes the proliferation of zooplankton, i.e., secondary and higher producers, it is essential to know how the biogeochemical composition of fast ice changes. However, although many studies have explored nutrient changes in first-year fast ice (e.g., Lim et al., 2019), research remains limited on biogeochemical components in multi-year fast ice that has continued to grow year-by-year. Furthermore, of the few studies that have focused on nutrient dynamics in multi-year land-fast ice (Nomura et al., 2018), their sea-ice samples provide only single-year observations. Therefore, no study has rigorously investigated the processes of nutrient cycling in multi-year sea ice. Because multi-year fast ice has remained fixed to the coast for a long time (Fraser et al., 2012; 2021), it is ideal for examining the temporal evolution and interannual variation of nutrient concentrations and other biogeochemical parameters in a single location.

The Japanese Antarctic Research Expedition (JARE) has monitored the multi-year fast ice near Syowa Station in Lutzow-Holm Bay for many years. In recent years, it has been reported that the multi-year fast ice in Lutzow-Holm Bay periodically breaks up and flows out to sea (Aoki, 2017; Ushio, 2006). Because ice thickness reflects the strength of sea ice, sea-ice monitoring efforts have recently collected annual ice core samples from an easily accessible and stable area near Syowa Station, ideal samples for tracking year-to-year

changes. Additionally, because new ice formed after the partial ice breaks up and outflows, it is possible to track the cycling of biogeochemical components through the ice by comparing early stages of sea-ice formation with the initial multi-year ice.

In this study, we analyzed multi-year fast ice near Syowa Station to evaluate the effects of snow and biological activity on the vertical distribution of nutrient concentrations in sea ice. We investigated the factors affecting nutrient concentrations in the multi-year fast ice by collecting and comparing nearby first-year ice. Specifically, based on annual sampling during the 56–60th JAREs (2015–2019), we analyzed: (1) the structure and formation process of ice using thin section photographs, salinity, and stable oxygen isotopic ratios ($\delta^{18}\text{O}$); (2) the vertical distribution of nutrient concentrations in the fast ice; (3) the influence of snow (based on snow fraction) on nutrient distributions; and (4) the cycling of nutrient concentrations due to biological activity within fast ice using sea-ice salinity, nutrient concentrations, and nutrient ratios as proxies.

2. Materials and Methods

2.1. Sampling and sample preparation

Sea ice observations of multi-year fast ice were performed during the summers of 2015–2019 as part of the 56–60th JAREs. Sea-ice cores were collected near Syowa Station (Figure 1, Table 1). Ice cores were sampled using an ice corer (Geo Tecs Co., Ltd., Chiba, Japan) with an internal diameter of 0.09 m. Cores were immediately placed in tubular polyethylene bags and kept horizontal in a cooler box along with refrigerants to maintain below-freezing temperatures and minimize brine drainage from the core. Fresh snow samples were collected with an acid-washed polycarbonate shovel and placed into polyethylene zipper storage bags. For comparison, four cores of first-year fast ice were also collected (Table 1).

Note that in Table 1, cores JARE58_KU3 and JARE59_2 were sampled only the top 2.78 and 2.47 m of ice, respectively, because the ice was too thick to sample the entire column. As part of the ice monitoring at Syowa Station, an electromagnetic (EM) ice thickness survey was conducted near our coring site in January 2018; the ice thickness was approximately 6 m.

Under-ice water was collected through the coring holes with a Teflon water sampler (GL Science Inc., Japan) from 1 m below the bottom of the sea ice during JARE60 (2019). Water samples were collected approximately 30 minutes after the ice core at JARE60_KU2 was drilled to avoid potential artifacts associated with the disturbance of drilling. All collected water samples were subsampled into (1) a 120-mL glass vial (Maruemu Co., Ltd., Osaka, Japan) for salinity measurement, (2) a 15-mL glass screw-cap vial (Nichiden-Rika Glass Co. Ltd, Kobe, Japan) for $\delta^{18}\text{O}$ analysis, (3) a 10-mL polyethylene screw-cap vial (Eiken Chemical Co. Ltd, Tokyo, Japan) for measurement of major inorganic nutrient concentrations ($\text{NO}_3^- + \text{NO}_2^-$, NH_4^+ , PO_4^{3-} , and $\text{Si}(\text{OH})_4$), and (4) a 500-mL Nalgene polycarbonate bottle (Thermo Fisher Scientific Inc., Waltham, MA, USA) for measurement of chlorophyll a (chl.a) concentrations. All samples were returned to the laboratory of the Japan Maritime Self-Defense Force icebreaker *Shirase* immediately after sampling. Samples for salinity and $\delta^{18}\text{O}$ analyses were stored in a refrigerator at $+4^\circ\text{C}$. Samples for nutrient analyses were stored in a freezer at -30°C . Samples for chl.a measurements were immediately filtered through 25-mm-diameter Whatman GF/F filters. Pigments on the filters were then extracted with *N,N*-dimethylformamide (Suzuki & Ishimaru, 1990) for more than 24 h and stored at -30°C until analysis.

Sea ice cores were stored in a freezer at -30°C during their transport to the Institute of Low Temperature Science (ILTS), Hokkaido University, Sapporo, Japan. In the ILTS cold room (-15°C), cores were split lengthwise into two halves with an electric band saw: the first half for ice textural analysis and the second half for measurements of bulk ice salinity, $\delta^{18}\text{O}$ composition, nutrient and chl.a concentrations, and cell counting for ice algae community assemblage.

2.2. Textural analysis and the age of an ice core

We examined ice thin sections to analyze ice growth processes. Ice sections 0.7 cm thick were attached to glass plates and planed down to 0.1-cm thickness with a microtome (Model SM2400, Leica Microsystems, Wetzlar,

Germany). We photographed ice crystallographic structures by illuminating the thin sections placing on a light table in between the cross-polarized sheets.

Ice growth processes were examined based on our classification of the ice cores as granular, columnar, or mixed ice (Eicken & Lange, 1989). Granular ice was subdivided into frazil ice and snow ice (or superimposed ice) based on its $\delta^{18}\text{O}$ value. We defined snow ice or superimposed ice as granular ice with a $\delta^{18}\text{O}$ value below that of its parent seawater (see Granskog et al., 2004; Jeffries et al., 1994), taken as the $\delta^{18}\text{O}$ value of the under-ice seawater sampled during JARE60 (-0.7‰ granular crystal textures) and fine-grained opaque ice originate from superimposed and snow ice, respectively (Kawamura et al., 1997), we hereafter refer to both as “snow-origin ice” unless obviously clear ice was observed in thin section, which was described as superimposed ice. We note that the analyzed snow was thawed and moistened, and refrozen ice was observed. The snow crystal structure was an aggregate of fine particles in thin section but did not contain salt (salinity = 0); the structure sometimes became sandwiched between superimposed ice, which we refer to as “fine-grained superimposed-ice” (see details in Section 3.1).

The ice type was determined according to WMO (2014) after checking satellite imagery of the sampling point, visually observing the sea ice condition at Syowa Station, and confirming the vertical profile of textural parameters (including thin section analysis) in the core. First-year ice was collected in locations where fast ice had broken up and flowed out during the previous year (Figure 1d–f), and sea ice had reformed within the year prior to collection. According to Syowa Station records, multi-year ice has not been broken up and flowed out since March 1980 (Higashi et al., 1982; Ushio, 2006).

2.3. Sample analysis

To measure ice salinity, water $\delta^{18}\text{O}$ composition, and nutrient and chl.a concentrations, the second half of the core was trimmed into a rectangle with a cross-section of 4.5 cm \times 2.5 cm and then sliced into 12–18-cm-thick sections. To avoid contamination during sampling and handling, the outer 0.3 cm of each ice section was removed with a stainless steel plane. The ice sample was then put in a polyethylene bag (GL Science Inc., Japan) and melted in a cold room at +4. Snow samples were melted in the same bags and in the same cold room. Snow and sea ice melt samples were analyzed with the same protocol as for under-ice water samples, except for nutrient analyses; in snow and sea ice melt samples, nutrient samples were filtered with a 0.22- μm Durapore PVDF Membrane filter (MILLEX GV Filter unit, Merck Millipore Ltd., Germany). For ice algae community assemblage, sea ice melt water for the bottom ice section for JARE60 KU1 and 2 was subsampled into 100-mL polyethylene screw-cap vial (AS ONE Corporation, Tokyo, Japan), and an aliquot of 5-mL of Lugol’s solution was added to preserve the sample.

The salinities of under-ice water and melted snow and sea-ice samples were measured with a conductivity sensor (Cond 315i, WTW GmbH, Germany). The oxygen isotopic ratio of under-ice water and melted snow and sea-ice samples were determined by mass spectrometry (DELTA plus; Finnigan MAT, San Jose, CA) via the equilibration method. The $\delta^{18}\text{O}$ value (the $^{18}\text{O}/^{16}\text{O}$ ratio of standard mean ocean water. The standard deviation (SD) on $\delta^{18}\text{O}$ values was 0.026‰ water with $\delta^{18}\text{O} = 0.241(\text{NO}_3^- + \text{NO}_2^-, \text{NH}_4^+, \text{PO}_4^{3-}, \text{and Si(OH)}_4)$ were measured with an auto-analyzer system (Quattro; Bran+Luebbe, Norderstedt, Germany) according to the spectrophotometric method reported by the Joint Global Ocean Flux Study (JGOFS, 1994). SDs on nutrient concentrations were 0.53, 0.18, 0.07, and 1.1 $\mu\text{mol L}^{-1}$ for $\text{NO}_3^- + \text{NO}_2^-$, NH_4^+ , PO_4^{3-} , and Si(OH)_4 , respectively, calculated from 10 subsamples of a reference water sample with respective known concentrations of 52.53, 9.99, 5.00, and 117.5 $\mu\text{mol L}^{-1}$. Chl.a concentrations were determined by a fluorometer (Model 10AU, Turner Designs, Inc., Sunnyvale, CA) using the method of Parsons et al. (1984). Standards (0.28–282.30 $\mu\text{g L}^{-1}$ chl.a) prepared from a liquid chl.a standard (Wako Pure Chemical Industries Ltd., Osaka, Japan) by stepwise dilution with *N,N*-dimethylformamide were used to calibrate the fluorometer before analyses. The cell counting for ice algae community assemblage was examined with a microscope (Olympus, BH-T, Tokyo, Japan) 10 \times oculars and 40 \times objective.

2.4. Mass fraction of snow in snow-origin ice

To determine the mass fraction of snow in snow-origin ice (hereafter, ‘snow fraction’), we used a mass balance

equation (Jeffries et al., 1994, 2001):

$$f_{\text{snow}} + f_{\text{sea}} = 1, (1)$$

$$f_{\text{snow}} \delta_{\text{snow}} + f_{\text{sea}} \delta_{\text{sea}} = \delta_{\text{obs}}, (2)$$

where f represents the mass fraction of snow or seawater, δ represents their $\delta^{18}\text{O}$ values, and the subscripts ‘snow’, ‘sea’, and ‘obs’ indicate snow, under-ice seawater, and observed bulk ice, respectively. For δ_{sea} , we used the parent water $\delta^{18}\text{O}$ values collected during JARE60 (-0.7‰ rather than the value that includes the fractionation factor during seawater freezing (see Jeffries et al., 1994, 2001) because we assumed that all granular ice with a $\delta^{18}\text{O}$ value below that of the under-ice water was snow ice (Granskog et al., 2004; Jeffries et al., 1994). In addition, we assumed that granular ice with a $\delta^{18}\text{O}$ value below that of under-ice water was snow-origin ice. Based on the result obtained from the JARE60 samples, we used $\delta_{\text{sea}} = -0.7$ ‰ the mean $\delta^{18}\text{O}$ value of snow samples (-20.3‰ during JARE56–60. We estimated the errors on the snow fractions to be less than 12.2% by calculating the effect of changing the snow $\delta^{18}\text{O}$ values by their standard deviation (± 2.1).

3. Results and Discussion

3.1. *ξρτισαλ προφιλες οφ σεα-ιςε σαλινιτις, $\delta^{18}\text{O}$ αλινες, ανδ σεα-ιςε στρυςτυρε*

Over the study period, first-year ice and snow on ice were ranged between 1.32 and 1.55 m thick and 0.05 and 0.20 m deep, respectively (Table 1). Multi-year ice and snow on ice were ranged between >2.47 and ca. 6 m thick and 0.34 and 1.68 m deep, respectively (Table 1). Figure 2 shows vertical salinity and $\delta^{18}\text{O}$ profiles for both first-year and multi-year ice cores. The salinity of the first-year ice was close to 0 near the ice–snow interface and increased with increasing depth until reaching a maximum, after which it decreased with increasing depth (Figure 2a). Salinity again increased at the bottom of some cores, giving the salinity profiles of first-year ice a characteristic “C” shape (excluding the topmost low-salinity ice). $\delta^{18}\text{O}$ values in first-year ice were as low as -20‰ (Figure 2b) and increased with increasing depth, reaching values above that of the under-ice water (-0.7‰) profiles record low salinity and $\delta^{18}\text{O}$ values near the ice–snow interface because the cores were collected in late summer (Table 1), by which time the accumulated snow had become snow-origin ice due to melting of the sea-ice surface in above-freezing air temperatures and strong sunlight (Nomura et al., 2011a). Furthermore, in late summer, the nighttime air temperature becomes negative in this area (Nomura et al., 2011a), freezing any water that had melted during the day. Because the snow depths were quite small relative to the ice thickness (Table 1) and the salinity was extremely low, especially near the ice surface, we considered that the ice formed was superimposed ice rather than snow ice because snow ice should only form when the snow depth is large with respect to the ice thickness (Maksym & Jeffries, 2000; Sturm & Massom, 2017). Indeed, in thin section, we observed large, polygonal, granular crystal textures (photograph in Figure 2a) characteristic of superimposed ice (Kawamura et al., 2004) near the ice surface.

The salinity of multi-year ice was 0 from the ice–snow interface to depths of around -100 cm (Figure 2c), then generally increased with increasing depth. The maximum observed salinity was 3.8 in core JARE58-KU3. At the bottom of the ice column, salinities were about 1–3. The $\delta^{18}\text{O}$ values of sea ice were similar to those of the overlying snow from the ice–snow interface to depths of -200 cm (Figure 2d), then increased with increasing depth. The ice $\delta^{18}\text{O}$ values reached that of the under-ice water at depths of -300 cm in core JARE56-NMS and -380 cm in core JARE57-P31.5. These results indicate that snow cover affects to a maximum depth of -380 cm.

Figure 3 compares the salinity and $\delta^{18}\text{O}$ profiles of multi-year ice core JARE57-P31.5 to the interpreted ice structure and photographs of representative thin sections. Up to a depth of -117 cm, salinity and $\delta^{18}\text{O}$ were maintained at 0 and -20‰ respectively (Figure 3a). Because most of this upper section of the core comprised coarse-grained (large polygonal) granular ice, we concluded that it was superimposed ice (Figure 3b, c; Kawamura et al., 2004). In addition, fine-grained granular ice was present at depths from -100 to -117 cm (Figure 3b, d). Based on the salinity and textures down to a depth of -117 cm, it is unlikely that snow ice was formed by the infiltration of seawater. In addition, the small crystal size at depths of -100 to -117 cm

(Figure 3d) indicated that this ice corresponded to superimposed ice formed from wet, unmelted snow that had re-frozen at negative temperatures, which we refer to as fine-grained superimposed ice.

At depths of -117 to -170 cm, sea ice salinity and $\delta^{18}\text{O}$ values increased (Figure 3a) and the ice structure became columnar (Figure 3b, e), with larger grains than in the overlying fine-grained granular ice. However, the columns were neither as wide nor as vertically extended as those of columnar ice formed from seawater (Figure 3f, see next paragraph). Based on the salinity and $\delta^{18}\text{O}$ values of the ice, this structural transformation occurred because the ice formed snowmelt with a slight seawater influence. Therefore, we suspect that this ice formed from a refrozen mixture of seawater and snowmelt. Indeed, in this area, it has been reported that a layer of superimposed ice forms under the snow, and that a slush layer (puddles in the sea ice that are likely snowmelt ponds) forms under the superimposed ice (Nomura et al., 2018). In that study, the salinity and $\delta^{18}\text{O}$ values of the slush layer were $[-1, -20]$ (as snow or superimposed ice), and a small amount of seawater was transported upwards through a crack and then horizontally transported into the slush layer. The salinity and $\delta^{18}\text{O}$ values of their slush layer were approximately the same as those of the columnar ice measured in our study (Figure 3a). The slush layer is about 10–20 cm thick (Nomura et al., 2012) and, in winter, is thought to freeze from the top faster than columnar ice develops at greater depths in the sea ice. Therefore, we consider that the columns in this zone are thinner and shorter than those in columnar ice formed from seawater (Figure 3e, f) because the freezing speed affects the crystal structure with slower growth on the c-axis direction than the basal plane (e.g., Hilling, 1959). Both fine- and coarse-grained granular ice have random crystal texture (Figure 3c-d). As the freezing develops, geometric selection sweep away crystals with c-axis perpendicular to the ice–ocean interface (Kolmogorov, 1949). The crystals in the slush layer that have the c-axes pointing radially (perpendicular to the ice–ocean interface) grow faster than those with c-axes pointing parallel to the ice–ocean interface. Therefore, it seems likely that the columns in slush layer was thinner due to the initial stage of geometric selection and shorter than those in columnar ice formed from seawater constrained by the slush layer thickness. In this study, we refer to this particular columnar ice texture as ‘slush-origin columnar ice’, in contrast to columnar ice formed from seawater (‘seawater-origin columnar ice’).

A distinct unit comprising a fine-grained granular ice layer overlying a coarse-grained granular ice layer was sandwiched between slush-origin columnar ices at depths of -170 to -190 cm. This unit formed because superimposed ice formed under the snow (Nomura et al., 2018) and a slush layer below it. This is thought to reflect greater amounts of snowfall in following years, with the slush layer being repeatedly formed and refrozen. The underlying slush-origin columnar ice layer extended to a depth of -300 cm, where the ice structure became columnar, but of seawater origin.

In contrast, the first-year ice was predominantly columnar (Table 2). From the above discussion, snow accumulates on the ice year-by-year, causing flooding and slush formation, then refreezing in winter. Furthermore, when the ice thickness increases, flooding ceases, and the snowmelt is refrozen. This upward growth is characteristic of Antarctic multi-year ice (Kawamura et al., 1997; Nomura et al., 2018), although the collection of a nearly 5-m sea-ice core has never been reported. This is thus the first time that the characteristics of the upward growth of multi-year fast ice have been shown based on multiple years of repeated ice coring.

3.2. Vertical profiles of sea-ice nutrient concentrations

Figure 4 presents vertical profiles of NO_3^- , PO_4^{3-} , and Si(OH)_4 concentrations in first-year and multi-year ice. NO_3^- concentrations in first-year ice varied widely with depth in core JARE60_KU1 (0 – $2.6 \mu\text{mol L}^{-1}$), but were in the range 0 – $1 \mu\text{mol L}^{-1}$ in other cores (Figure 4a). NO_3^- concentrations in multi-year ice were 0 – $0.5 \mu\text{mol L}^{-1}$ in the top half of the ice column, and the first maximum at those depths, peaking at $\sim 1.0 \mu\text{mol L}^{-1}$, was observed at a depth of -200 cm in some ice cores (Figure 4e). At greater depths in the cores, NO_3^- concentrations were as low as 0 – $0.5 \mu\text{mol L}^{-1}$, and the maximum of $3.5 \mu\text{mol L}^{-1}$ was observed at the bottom of the sea ice. In first-year ice, PO_4^{3-} concentrations varied between cores but were fairly constant at shallow to medial depths, and were greatest at the bottom of the ice column (Figure 4b). In multi-year ice, however, PO_4^{3-} concentrations were extremely low from the ice–snow interface to a depth of -150 cm,

and the maximum concentration of $1.6 \mu\text{mol L}^{-1}$ was observed at a depth of -360 cm in core JARE57_P31.5 (Figure 4f). Si(OH)_4 concentrations varied little with depth in the first-year ice (Figure 5c), but mimicked the PO_4^{3-} profile in multi-year ice.

Our results show clear differences between the vertical nutrient profiles in first-year and multi-year ice (Figure 4). First-year ice generally has the lowest nutrient concentrations in snow, and concentrations increase slightly in sea ice, but they remain relatively constant until peaking at the bottom of the ice column (Figure 4a–c). In contrast, in multi-year ice, the upper half of the ice column was nutrient poor, similar to the overlying snow, and nutrient concentrations increased from the middle of the column to maximum concentrations in the bottom half of the column (Figure 4e–g). Because nutrient concentrations in snow were consistently low (Figure 4e–g), the upper half of the multi-year ice columns, comprising snow-origin ice, also contained low nutrient concentrations.

Such vertical nutrient distributions are considered to be closely related to physical characteristics such as salinity, $\delta^{18}\text{O}$ value, and ice structure (discussed in Section 3.3). Furthermore, the high concentrations of various nutrients at the bottom of both first-year and multi-year ice columns are most likely related to the high concentrations of chl.a in the bottommost ice column (Figure 4d, h), which undergoes both photosynthesis by the ice algae and remineralization by the degradation of organic matter (Roukaerts et al., 2021; Rysgaard & Glud, 2004; Rysgaard et al., 2008; Thomas et al., 1995) and receives nutrients supplied from under-ice water (Nomura et al., 2009; Vancoppenolle et al., 2010).

The mean value and standard deviation of each nutrient component and physical parameter are reported in Table 3 for snow and each sea-ice type. Nutrient and chl.a concentrations were low in snow and superimposed ice and particularly high in columnar ice. In contrast, snow-ice, which is affected by both snow and seawater, had intermediate nutrient and chl.a concentrations. Our results clearly indicate that a greater snow influence results in lower nutrient and chl.a concentrations in sea ice.

3.3. Variations of nutrient concentrations in snow-origin ice

In Section 3.2, we showed that in multi-year ice columns, nutrient concentrations were lower in snow-origin ice than in columnar ice (Table 3). To quantify the effect of snow on sea-ice nutrient concentrations, we calculated the snow fraction in snow-origin ice (Eqs. 1 and 2) to examine the relationship between nutrient concentrations and snow fraction (Figure 5). Although the regressions differed between nutrients, nutrient concentrations tended to decrease with increasing snow fraction. This is due to the purity of the air near Antarctica, being far from areas of human activity, compared to that in the Northern Hemisphere. Therefore, the atmospheric transfer and deposition of nutrients onto sea ice is limited in Antarctica, and the top surface of the sea ice is replaced by clean, snow-origin ice. In contrast, in the Northern Hemisphere, atmospheric transfer and deposition of polluted snow is known to be a potential source of nutrients for sea ice (Granskog & Kaartokallio, 2004; Granskog et al., 2003; Kaartokallio, 2001; Krell et al., 2003; Nomura et al., 2010; 2011b; Rahm et al., 1995).

Although nutrient concentrations were generally negatively correlated with snow fraction (Figure 5a, b, d), NO_3^- concentrations notably differed, with the negative correlation between snow fraction and NO_3^- concentration being markedly low compared to the other nutrients (Figure 5c). Because most of the sea-ice samples in this study were collected near Syowa Station, it is possible that this difference reflects slight pollution from local human activities, such as exhaust from the research facility and/or snowmobiles. Therefore, greater NO_3^- concentrations might be detected (Table 3) in areas of greater snow fraction (Figure 5). However, many NO_3^- data points plot near zero at snow fractions of 30–80% (Figure 5c), potentially indicating that NO_3^- was selectively removed by sea ice, which can be explored by considering denitrification.

Denitrification is a reaction in which NO_3^- is used by bacteria in anoxic conditions to decompose organic substances within sea ice, converting NO_3^- to NO_2^- , NO, and eventually N_2O and N_2 (Kaartokallio, 2001; Nomura et al., 2010; 2011b; 2018; Rysgaard & Glud, 2004; Rysgaard et al., 2008; Staley & Gosink, 1999). Snow-origin ice with small snow fractions, i.e., containing some, albeit small, amounts of seawater, must be older than snow-origin ice not containing any seawater (e.g., superimposed ice) because it is formed at an

earlier stage of ice growth. Therefore, we consider that denitrification occurred in the older parts of the ice column where the snow fraction was low (Figure 5c; see Section 3.4 for details).

Due to the formation of snow-origin ice, the snow fraction in multi-year ice increased year-by-year; the mean snow fractions in the top 200 cm of multi-year ice gradually increased from about 88% in 2015 to 100% by 2018 (Figure 6). Therefore, our sea-ice core samples record a steady increase in the contribution of snow to sea ice. Although we have not collected multi-year ice cores at distant locations within the same year (Figure 1c) and thus have not been able to unequivocally determine that there is no spatial bias in our sampling, the nutrient concentrations do not differ greatly between sampling locations, and the trends and profiles are similar from year to year (Figures 2, 4).

3.4. Variations of nutrient concentrations in seawater-origin ice

Although nutrient concentrations decreased as the snow fraction increased in the topmost sea ice, the distributions of nutrient concentrations differed among the different nutrient components (Figure 5). Therefore, we here evaluate the effects of primary production, organic matter remineralization, and denitrification on nutrient concentrations in both first-year and multi-year ice. We first compared salinity to nutrient concentrations (Figure 7). Sea-ice salinity is a conservative component that only changes through physical processes such as brine drainage or exchange with under-ice water. Therefore, we investigated deviations from a dilution line based on the salinity and nutrient concentrations of the under-ice water collected during JARE60 to understand the biological processes affecting the nutrient distributions in the sea ice. Data plotting above the dilution line indicate nutrient supply due to the remineralization of organic matter by heterotrophic organisms and bacteria in the sea ice or from external sources (e.g., the atmosphere); conversely, data points below the dilution line indicate nutrient uptake by ice algae during primary production (e.g., Thomas et al., 1995).

Si(OH)_4 concentrations in first-year ice locate significantly below the dilution line (Figure 7a), whereas those in multi-year ice are closer to or above the dilution line (Figure 7b). The distributions of chl.a concentrations in the sea ice (Figure 4d, h) indicate the presence of ice algae, which would mainly be diatoms (*Fragilariopsis* sp. and *Pleurosigma* sp. (photo in 4c, d) based on the cell counting for ice algae community assemblage in this study. Indeed, in first-year ice, primary production by diatoms consumes a large amount of Si(OH)_4 (Tréguer & De La Rocha, 2013), and the slow dissolution of diatom frustules should limit the resupply of Si(OH)_4 . In contrast, in multi-year ice, the more protracted dissolution of diatom frustules should resupply Si(OH)_4 , eventually balancing the consumption of Si(OH)_4 by primary production. However, in core JARE57_P31.5 (the deepest core studied), Si(OH)_4 concentrations plot well above the dilution line (Figure 7b), indicating that the resupply of Si(OH)_4 by dissolution of diatom frustules exceeded consumption.

Consistent with a previous sampling at the same location (Nomura et al., 2018), PO_4^{3-} concentrations plot above the dilution line in both first-year and multi-year ice (Figure 7c, d), suggesting resupply by the remineralization of organic matter. The difference in the resupply rates of PO_4^{3-} and Si(OH)_4 is due to the fact that the remineralization of PO_4^{3-} is faster than the dissolution of Si(OH)_4 (Fripiat et al., 2017).

NO_3^- concentrations in most first-year and multi-year ice cores plot below the dilution line (Figure 7e, f). This is expected for first-year ice, in which primary production consumes NO_3^- , as with Si(OH)_4 . In contrast, remineralization in multi-year ice should resupply NO_3^- , although our data do not reflect this. Therefore, we also measured NO_2^- concentrations (Figure 7g, h), which plot close to the dilution line for first-year ice and are shifted upward for multi-year ice. From this result, we expect that denitrification resupplied NO_2^- over longer periods in the multi-year ice. Indeed, denitrifying bacterial species has been confirmed in Antarctic ice (Staley & Gosink, 1999), which consume NO_3^- and supply NO_2^- (Rysgaard et al., 2008). To better understand the effect of denitrification on the nutrient balance, we calculated DIN:P, the ratio of dissolved inorganic nitrogen (DIN, i.e., $\text{NO}_3^- + \text{NO}_2^-$) to PO_4^{3-} (Figure 8). In most cores, DIN:P values were lower than the Redfield ratio (16:1) and the diatom nutrient uptake ratio (12:1; Takeda, 1998), which is thought to reflect the transformation of NO_3^- and NO_2^- to NO , N_2O , and N_2 by denitrification. The value of N^* , which has been suggested as a proxy for denitrification, is calculated as follows (Gruber & Sarmiento, 1997;

Nishioka et al., 2021; Yoshikawa et al., 2006):

$$N^* = ([NO_3^-] - 16 [PO_4^{3-}] + 2.9) \times 0.87. \quad (3)$$

Figure 9 shows N^* profiles of multi-year ice cores JARE56_NMS and JARE57_P31.5. In the lower parts of these cores, N^* values decreased to negative values, reaching a minimum of $-12.3 \mu\text{mol L}^{-1}$ in core JARE57_P31.5. Chl.a concentrations are also negatively correlated with N^* . Previous studies have reported that denitrification occurred in oxygen-depleted and organic matter-rich parts of the ice column (Rysgaard et al., 2008). It is therefore reasonable that denitrification occurred at depths where high chl.a concentrations indicate abundant organic matter. To our knowledge, because multi-year ice formed a long time ago in this area (about 40 years ago) (Higashi et al., 1982; Ushio, 2006), salinity is low (Figure 2c), and it is unlikely that brine channels physically drain the ice. Therefore, the air and brine pockets in multi-year ice has remained isolated and disconnected from exchanges with the atmosphere and under-ice water.

To examine the possibility of oxygen exchange through the brine channel network in the sea ice, we focused on the brine volume fraction, a proxy of sea-ice permeability. Calculating brine volume fraction requires the ice temperature and salinity; however, we could not obtain realistic ice temperature data due to the difficulty of inserting a temperature sensor in the sea ice, especially in low-salinity, snow-origin ice. The ice temperature increased during coring due to friction, and the air temperature was often positive (Nomura et al., 2011a), precluding accurate measurement of the sea ice temperature. Therefore, for the brine volume calculation, we assumed the temperature of the sea ice to be -2 to 0°C , a reasonable estimate given the positive air temperature. We obtained brine volume fractions, estimated following Leppäranta and Manninen (1988), of $2.4 \pm 2.6\%$ (mean and standard deviation) at all depths in the multi-year ice. Typically, sea ice is considered permeable when the brine volume fraction exceeds 5 – 7.5% (Golden et al., 1998; Pringle et al., 2009; Zhou et al., 2013). Therefore, the multi-year ice collected in this study was impermeable, precluding any exchanges through brine channels. Therefore, gas components such as oxygen must have been consumed in a closed system within the ice by heterotrophic organisms and bacteria. Furthermore, the high density of the hard, low-salinity superimposed ice suggests that gases could not be exchanged with the atmosphere, consistent with previous studies that were unable to detect gas fluxes over superimposed ice at the same location (Nomura et al., 2012, 2013). Therefore, oxygen in the isolated interior sea ice could only have been depleted by heterotrophic organisms and bacteria, promoting denitrification in multi-year ice (Figure 9).

Based on our results, we developed a conceptual illustration for the evolving nutrient concentrations in multi-year fast ice (Figure 10). Immediately after ice formation, when nutrient concentrations have not yet been affected by biological processes, the ratio of salinity to nutrients in the brine discharged from the growing sea ice should plot along the dilution line. Then, during the first year, nutrient uptake by primary production of ice algae decreases nutrient concentrations in the ice while the salinity is reduced by brine convection. Over multiple years, remineralization by the degradation of organic matter resupplies nutrients to the ice while salinity continues to decrease. Once ice algae have exhausted the nutrient supply and decompose, and brine exchange with seawater is established at the bottom of the ice column, the nutrient concentrations surpass the dilution line (Figure 10). In addition, denitrification decreases NO_3^- concentrations and modifies the speciation of nitrogen within the ice.

4. Conclusions

We investigated the effects of snow cover on the vertical distribution of nutrient concentrations and the long-term remineralization process in multi-year, Antarctic fast sea ice. We analyzed thin sections to determine the ice structure, which confirmed the upward growth of fast ice by the year-by-year formation of snow-origin ice. Vertical profiles of biogeochemical components in the ice showed that the high nutrient concentrations of first-year ice are transported downward by the subsequent formation of new snow-origin ice, eventually forming multi-year ice. Because the atmospheric nutrient supply is limited by the purity of Antarctic snow, nutrient concentrations in snow-origin ice decreased as the snow fraction increased.

Nutrients incorporated into columnar ice of seawater origin were dominantly processed by biological uptake via primary production during the first year, but remineralization of degrading organic matter by

heterotrophic organisms and bacteria resupplied nutrients to multi-year ice. By analyzing data on first-year and multi-year ice spanning several years in the same area, it was possible for the first time to determine the long-term cycling of nutrients in fast ice.

Acknowledgements

We express our heartfelt thanks to the sea ice team, members of the Japanese Antarctic Research Expedition, the crew of the Japan Maritime Self-Defense Force icebreaker *Shirase*, and cruise members for their support in conducting field work. We thank Prof. K. Kawashima, K. Murakami, H. Nakajima, M. Kiuchi, T. Ishino, and T. P. Tamura for their support in the low temperature room, M. Hirabayashi for oxygen isotopic analyses, and A. Ooki for nutrient analyses. We benefitted greatly from advice and useful comments on the nutrient data offered by A. Ooki and T. Ishino. This work was supported by grants from the Japan Society for the Promotion of Science (#17H04715, #17H04710, #17H06317, #17H06322, #18F18794, #20H04345, #21H04931),

the Science Program of the Japanese Antarctic Research Expedition (JARE) as Prioritized Research AJ0902 (ROBOTICA), the National Institute of Polar Research (NIPR) through Project Research KP-303 and General Collaboration Project #28-14, #2-13 and the grant for young scientist, the Center for the Promotion of Integrated Sciences of SOKENDAI, and the Joint Research Program of the Institute of Low Temperature Science, Hokkaido University. The data used in this study are available in the supplementary information (Table S1) and are archived in the data repository of the National Institute of Polar Research, Tokyo, Japan (<https://ads.nipr.ac.jp>), is underway.

References

- Ackley, S. F., & Sullivan, C. W. (1994). Physical controls on the development and characteristics of Antarctic sea ice biological communities— A review and synthesis. *Deep Sea Research Part I : Oceanographic Research Papers*, 41, 1583–1604. [https://doi.org/10.1016/0967-0637\(94\)90062-0](https://doi.org/10.1016/0967-0637(94)90062-0)
- Aoki, S. (2017). Breakup of land-fast sea ice in Lutzow-Holm Bay, East Antarctica, and its teleconnection to tropical Pacific sea surface temperatures. *Geophysical Research. Letters*, 44, 3219–3227. doi:10.1002/2017GL072835
- Arrigo, K. R., Mock, T., & Lizotte, M. P. (2010). Primary producers and sea ice. In D.N. Thomas., & G.S. Dieckmann. (Eds.), *Sea Ice (Second edition)*, Wiley-Blackwell, Oxford, 283-325. <https://doi.org/10.1002/9781444317145.ch8>
- Arrigo, K. R. (2017). Sea ice as a habitat for primary producers. In D. N. Thomas (Ed.), *Sea ice (Third edition)*, 352– 369. <https://doi.org/10.1002/9781118778371.ch14>
- Eicken, H., & Lange, M. (1989). Development and properties of sea ice in the coastal regime of the southern Weddell Sea. *Journal of Geophysical Research: Oceans*, 94, 8193–8206. <https://doi.org/10.1029/JC094iC06p08193>
- Fraser, A. D., Massom, R. A., Michael, K. J., Galton-Fenzi, B. K., & Lieser, J. L. (2012). East Antarctic landfast sea ice distribution and variability, 2000–08. *Journal of Climate*, 25, 1137–1156. DOI:10.1175/JCLI-D-10-05032.1
- Fraser, A. D., Massom, R. A., Handcock, M., Reid, P., Ohshima, K. I., Raphael, M., et al. (2021). Eighteen-year record of circum-Antarctic landfast-sea-ice distribution allows detailed baseline characterisation and reveals trends and variability. *The Cryosphere*, 15, 11, 5061–5077. <http://dx.doi.org/10.5194/tc-15-5061-2021> Retrieved from <https://escholarship.org/uc/item/4gb0869t>
- Fripiat, F., Meiners, K. M., Vancoppenolle, M., Papadimitrio, S., Thomas, D.N., Ackley, S. F., et al. (2017). Macro-nutrient concentrations in Antarctic pack ice: Overall patterns and overlooked processes. *Elementa Science of Anthropocene*, 5-13. <https://doi.org/10.1525/elementa.217>

- Fritsen, C. H., Lytle, V. I., Ackley, S. F., & Sullivan, C. W. (1994). Autumn bloom of Antarctic pack-ice algae. *Science* , 266, 782–784. DOI: 10.1126/science.266.5186.782
- Gleitz, M., Vonderlo, M. R., Tomas, D. N., Dieckmann, G. S., & Millero, F. J. (1995). Comparison of summer and winter inorganic carbon, oxygen and nutrient concentrations in Antarctic sea ice brine. *Marine Chemistry* , 51, 81–89. [https://doi.org/10.1016/0304-4203\(95\)00053-T](https://doi.org/10.1016/0304-4203(95)00053-T)
- Golden, K. M., Ackley, S. F., & Lytle, V. I. (1998). The percolation phase transition in sea ice. *Science* , 282, 2238–2241. DOI: 10.1126/science.282.5397.2238
- Granskog, M. A., Kaartokallio, H., & Shirasawa, K. (2003). Nutrient status of Baltic Sea ice: evidence for control by snow–ice formation, ice permeability, and ice algae. *Journal of Geophysical Research: Oceans* , 108 (C8), 3253. doi:10.1029/2002JC001386
- Granskog, M. A., & Kaartokallio, H. (2004). An estimation of the potential fluxes of nitrogen, phosphorus, cadmium and lead from sea ice and snow in the northern Baltic Sea. *Water Air Soil Pollution* , 154, 331–347.
- Granskog, M. A., Lepparanta, M., Kawamura, T., Ehn, J., & Shirasawa, K. (2004). Seasonal development of the properties and composition of landfast sea ice in the Gulf of Finland, the Baltic Sea. *Journal of Geophysical Research: Oceans* , 109 (C2), C02020. DOI:10.1029/2003JC001874
- Gruber, N., & Sarmiento, J. L. (1997). Global patterns of marine nitrogen fixation and denitrification. *Global Biogeochemical Cycles* , 11, pp. 235–266. <https://doi.org/10.1029/97GB00077>
- Higashi, A., Goodman, D. J., Kawaguchi, S., & Mae, S. (1982). The cause of the breakup of fast ice on March 18, 1980 near Syowa station, East Antarctica. *Memoirs of National Institute of Polar Research* , 24, 222–231.
- Hillig, W. (1959). Kinetics of solidification from nonmetallic liquids. In Kingery W, ed. *Kinetics of high-temperature processes* . MIT Press, Cambridge, MA, 127–135.
- Jeffries, M., Shaw, R., Morris, K., Veazey, A., & Krouse, H. (1994). Crystal structure, stable isotopes ($\delta^{18}\text{O}$), and development of sea ice in the Ross, Amundsen, and Bellingshausen seas, Antarctica. *Journal of Geophysical Research: Oceans* , 99 (C1), 985–995. <https://doi.org/10.1029/93JC02057>
- Jeffries, M. O., Krouse, H. R., Hurst-Cushing, B., & Maksym, T. (2001). Snow-ice accretion and snow-cover depletion on Antarctic first-year sea-ice floes. *Annals of Glaciology* , 33, 51–60.
- Kaartokallio, H. (2001). Evidence for active microbial nitrogen transformations in sea ice (Gulf of Bothnia, Baltic Sea) in midwinter. *Polar Biology* , 24:21–28. <https://doi.org/10.1007/s0030000000169>
- Kattner, G., Thomas, D. N., Haas, C., Kennedy, H., & Dieckmann, G. S. (2004). Surface ice and gap layers in Antarctic sea ice: Highly productive habitats. *Marine Ecology Progress Series* , 277, 1–12.
- Kawamura, T., Ohshima, K. I., Takizawa, T., & Ushio, S. (1997). Physical, structural, and isotopic characteristics and growth processes of fast sea ice in Lützow-Holm Bay-Holm Bay, Antarctica. *Journal of Geophysical Research: Oceans* , 102, 3345–3355.
- Kawamura, T., Jeffries, M., Tison, J., & Krouse, H. (2004). Superimposed-ice formation in summer on Ross Sea pack-ice floes. *Annals of Glaciology* , 39 , 563–568. doi:10.3189/172756404781814168
- Kolmogorov, A. N. (1976). Geometric selection of crystals. *Doklady Akad Nauk Minerologia USSR*.
- Krell, A., Ummenhofer, C., Kattner, G., Naumov, A., Evans, D., Dieckmann, G. S., et al. (2003). The biology and chemistry of land fast ice in the White Sea, Russia—A comparison of winter and spring conditions. *Polar Biology* , 26, 707–719.

- Leppäranta, M., & Manninen, T. (1988). The brine and gas content of sea ice with attention to low salinities and high temperatures (Internal Rep. 88-2). Helsinki, Finland: Finnish Institute of Marine Research.
- Lim, S. M., Moreau, S., Vancoppenolle, M., Deman, F., Roukaerts, A., Meiners, K. M., et al. (2019). Field observations and physical-biogeochemical modeling suggest low silicon affinity for Antarctic fast-ice diatoms. *Journal of Geophysical Research: Oceans* , 124, 11, 7837–7853. <https://doi.org/10.1029/2018jc014458>
- Maksym, T., & Jeffries, M. O. (2000). A one-dimensional percolation model of flooding and snow-ice formation on Antarctic sea ice. *Journal of Geophysical Research: Oceans* , 105(C11), 26313–26331. <https://doi.org/10.1029/2000JC900130>
- Nishioka, J., Hirawake, T., Nomura, D., Yamashita, Y., Ono, K., Murayama, A., et al. (2021). Iron and nutrient dynamics along the East Kamchatka Current, western Bering Sea Basin and Gulf of Anadyr. *Progress in Oceanography* , 198, 10.1016/j.pocean.2021.102662, 2021.
- Nomura, D., Takatsuka, T., Ishikawa, M., Kawamura, T., Shirasawa, K., Yoshikawa-Inoue, H. (2009). Transport of chemical components in sea ice and under-ice water during melting in the seasonally ice-covered Saroma-ko Lagoon, Hokkaido, Japan. *Estuarine, Coastal and Shelf Science* , 81, 201–209.
- Nomura, D., Nishioka, J., Granskog, M. A., Krell, A., Matoba, S., Toyota, T., et al. (2010). Nutrient distributions associated with snow and sediment-laden layers in sea ice of the southern Sea of Okhotsk. *Marine Chemistry* , 119, 1–4. <https://doi.org/10.1016/j.marchem.2009.11.005>
- Nomura, D., Simizu, D., Shinagawa, H., Oouchida, C., & Fukuchi, M. (2011a). Biogeochemical properties of water in surface ponds on Antarctic fast ice and relationship to underlying sea ice properties. *Journal of Glaciology* , 57, 205, 1–9. DOI: <https://doi.org/10.3189/002214311798043825>
- Nomura, D., McMinn, A., Hattori, H., Aoki, S., & Fukuchi, M. (2011b). Incorporation of nitrogen compounds into sea ice from atmospheric deposition. *Marine Chemistry* , 127, 90–99. <https://doi.org/10.1016/j.marchem.2011.08.002>
- Nomura, D., Koga, S., Kasamatsu, N., Shinagawa, H., Simizu, D., Wada, M., et al. (2012). Direct measurements of DMS flux from Antarctic fast sea ice to the atmosphere by a chamber technique. *Journal of Geophysical Research-Oceans* , 117, C04011, doi: 10.1029/2010JC006755
- Nomura, D., Aoki, S., Simizu, D., & Iida, T. (2018). Influence of sea ice crack formation on the spatial distribution of nutrients and microalgae in flooded Antarctic multiyear ice. *Journal of Geophysical Research: Oceans* , 123, 939–951. <https://doi.org/10.1002/2017JC012941>
- Parsons, T. R., Takahashi, M., & Hargrave, B. (1984). Biological oceanographic processes (3rd ed., 330 pp.). Oxford, UK: Pergamon.
- Pringle, D. J., Miner, J. E., Eicken, H., & Golden, K. M. (2009). Pore space percolation in sea ice single crystals. *Journal of Geophysical Research: Oceans* , 114, C12017. doi:10.1029/2008JC005145
- Rahm, L., Hakansson, B., Larsson, P., Fogelqvist, E., Bremle, G., & Valderrama, J. (1995). Nutrient and persistent pollutant deposition on the Bothnian Bay ice and snow fields. *Water Air and Soil Pollution* , 84, 187–201.
- Roukaerts, A., Deman, F., Linden, F. V., Carnat, G., Bratkic, A., Moreau, S., et al. (2021). The biogeochemical role of a microbial biofilm in sea ice: Antarctic landfast sea ice as a case study. *Elementa: Science of the Anthropocene* , 9: 1. DOI: <https://doi.org/10.1525/elementa.2020.00134>
- Rysgaard, S., & Glud, R. N. (2004). Anaerobic N₂ production in Arctic sea ice. *Limnology and Oceanography* , 49:86–94. <https://doi.org/10.4319/lo.2004.49.1.0086>
- Rysgaard, S., Glud, R. N., Sej, M. K., Blicher, M. E., & Stahl, H. J. (2008). Denitrification activity and oxygen dynamics in Arctic sea ice. *Polar Biology* , 31, 527–537. doi: 10.1007/s00300-007-0384-x

- Saenz, B., T., & Arrigo, K. R. (2014). Annual primary production in Antarctic sea ice during 2005–2006 from a sea ice state estimate. *Journal of Geophysical Research: Oceans* , 119, 3645– 3678, <https://doi.org/10.1002/2013JC009677>
- Schnack-Schiel, B., Thomas, D. N., Haas, C., Dieckmann, G. S., & Alheit, R. (2001). The occurrence of the copepods *Stephos longipes* (Calanoida) and *Drescheriella glacialis* (Harpacticoida) in summer sea ice in the Weddell Sea, Antarctica. *Antarctic Science* , 13, 2, 150–157.
- Staley, J. T., & Gosink, J. J. (1999). Poles apart: biodiversity and biogeography of sea ice bacteria. *Annual Review of Microbiology* , 53, 189-215. doi: 10.1146/annurev.micro.53.1.189
- Sturm, M., & Massom, R. A. (2017). Snow and sea ice. D.N. Thomas, G.S. Dieckmann (Ed.), *Sea Ice (Second edition)* , Wiley-Blackwell, Oxford, 153-204. <https://doi.org/10.1002/9781444317145.ch5>
- Suzuki, R., & Ishimaru, T. (1990). An improved method for the determination of phytoplankton chlorophyll using N, N-dimethylformamide. *Journal of the Oceanographic Society of Japan* , 46, 190–194. <https://doi.org/10.1007/BF02125580>
- Takeda, S. (1998). Influence of iron availability on nutrient consumption ratio of diatoms in oceanic waters. *Nature* , 393, 774–777.
- Thomas, D. N., Lara, R. J., Eicken, H., Kattner, G., & Skoog, A. (1995). Dissolved organic matter in Arctic multi-year sea ice during winter: Major components and relationship to ice characteristics. *Polar Biology* , 15(7), 477–483.
- Toyota, T., Massom, R., Lecimte, O., Nomura, D., Heil, P., Tamura, T., Fraser, A. D. (2016). On the extraordinary snow on the sea ice off East Antarctica in late winter, 2012. *Deep-Sea Research II* , 131, (2016), 53-67. <https://doi.org/10.1016/j.dsr2.2016.02.003>
- Treguer, P. J. & De La Rocha, C. L. (2013). The World Ocean silica cycle. *Annual Review of Marine Science*, 5: 477–501. DOI: <https://doi.org/10.1146/annurev-marine-121211-172346>
- Yoshikawa, C., Nakatsuka, T., & Wakatsuchi, M. (2006). Distribution of N* in the Sea of Okhotsk and its use as a biogeochemical tracer of the Okhotsk Sea Intermediate Water formation process. *Journal of Marine Systems* . 63 (1–2), 49–62. <https://doi.org/10.1016/j.jmarsys.2006.05.008>
- Ushio, S. (2006). Factors affecting fast-ice break-up frequency in Lutzow-Holm Bay, Antarctica. *Annals of Glaciology* , 44, 177–182. DOI: <https://doi.org/10.3189/172756406781811835>
- Vancoppenolle, M., Goosse, H., de Montety, A., Fichet, T., Tremblay, B., & Tison, J.-L. (2010). Modeling brine and nutrient dynamics in Antarctic sea ice: The case of dissolved silica, *Journal of Geophysical Research: Oceans* , 115, C02005, doi:10.1029/2009JC005369
- WMO (2014). Sea Ice Nomenclature. World Meteorological Organisation, Geneva.
- Wongpan, P., Meiners, K. M., Langhorne, P. J., Heil, P., Smith, I. J., Leonard, G. H., et al. (2018). Estimation of Antarctic land-fast sea ice algal biomass and snow thickness from under-ice radiance spectra in two contrasting areas. *Journal of Geophysical Research: Oceans* , 123(3), 1907–1923. doi: 10.1002/2017jc013711
- Zhou, J., Delille, B., Eicken, H., Vancoppenolle, M., Brabant, F., Carnat, G., et al. (2013). Physical and biogeochemical properties in Landfast sea ice (Barrow, Alaska): Insights on brine and gas dynamics across seasons. *Journal of Geophysical Research: Oceans* , 118, 6, 3172–3189. <https://doi.org/10.1002/jgrc.20232>

Figure captions

Figure 1. (a) The study area and (b–f) sampling locations in Lutzow-Holm Bay, Eastern Antarctica. MODIS-Terra satellite images were obtained on (b) 19 January 2019, (d) 24 April 2015 at site JARE57_EM, (e) 25 April 2017 at site JARE59_POPs, and (f) 18 March 2018 at sites JARE60_KU1 and JARE60_KU2. The images in (d–f) show the sea-ice condition before winter at each coring site; dark areas indicate where the sea

ice was broken, and the cores obtained from these sites were first-year ice. The background image in (c) is an Advanced Land Observing Satellite (ALOS)/Panchromatic Remote-sensing Instrument for Stereo Mapping (PRISM) image obtained on 26 January 2010 by the Japan Aerospace Exploration Agency (JAXA).

Figure 2. Vertical profiles of (a, c) salinity and (b, d) $\delta^{18}\text{O}$ values in snow, first-year ice cores (left), and multi-year ice cores (right). Horizontal solid lines at 0 cm depth indicate the snow–ice interface. Dashed lines in (b, d) show the $\delta^{18}\text{O}$ value of under-ice water (-0.7‰ JARE60). The photo in (a) is a vertical thin section of the top 5 cm of core JARE60_KU2.

Figure 3. (a) Vertical profiles of salinity and $\delta^{18}\text{O}$ values in core JARE57_P31.5 compared to (b) its ice structure. (c–f) Representative vertical thin section photographs are provided for each ice structure.

Figure 4. Vertical profiles of (a, e) NO_3^- , (b, f) PO_4^{3-} , (c, g) Si(OH)_4 , and (d, h) chl.a concentrations in first-year ice (top) and multi-year ice cores (bottom). Horizontal solid lines at a depth of 0 cm indicate the snow–ice interface. The photo in (c) and (d) is the ice algae community assemblage for the bottom of core JARE60_KU1.

Figure 5. Plots of (a) PO_4^{3-} , (b) Si(OH)_4 , (c) NO_3^- , and (d) NO_2^- concentrations versus snow fraction in multi-year ice cores.

Figure 6. Time series of the mean snow fraction in the top 200 cm of each multi-year ice core from 2015 (core JARE56_NMS) to 2018 (core JARE59_2).

Figure 7. Plots of (a, b) Si(OH)_4 , (c, d) PO_4^{3-} , (e, f) NO_3^- , and (g, h) NO_2^- concentrations versus salinity in first-year (left) and multi-year ice cores (right). Dilution lines are based on the nutrient concentrations and salinity of under-ice water.

Figure 8. Dissolved inorganic nitrogen ($\text{DIN} = \text{NO}_3^- + \text{NO}_2^-$) versus PO_4^{3-} concentrations in multi-year ice cores. The red solid line indicates the Redfield ratio ($\text{DIN:P} = 16:1$) and the black dashed line indicates the ratio for uptake by diatoms ($\text{DIN:P} = 12:1$) (Takeda, 1998).

Figure 9. Vertical profiles of N^* (Equation 3) and chl.a concentrations in multi-year ice cores (a) JARE56_NMS and (b) JARE57_P31.5.

Figure 10. Conceptual illustration of the evolution of nutrient concentrations produced by photosynthesis and remineralization during the desalination process from new ice to first-year ice to multi-year ice.

Effects of snow and remineralization processes on nutrient distributions in multi-year Antarctic landfast sea ice

Reishi Sahashi¹, Daiki Nomura^{2,1,3*}, Takenobu Toyota⁴, Manami Tozawa¹, Masato Ito⁵, Pat Wongpan^{6,4,7}, Kazuya Ono⁴, Daisuke Simizu⁵, Kazuhiro Naoki⁸, Yuichi Nosaka⁹, Takeshi Tamura^{5,10}, Shigeru Aoki⁴, Shuki Ushio^{5,10}

¹*Faculty of Fisheries Sciences, Hokkaido University, Hakodate, Japan*

²*Field Science Center for Northern Biosphere, Hokkaido University, Hakodate, Japan*

³*Arctic Research Center, Hokkaido University, Sapporo, Japan*

⁴*Institute of Low Temperature Science, Hokkaido University, Sapporo, Japan*

⁵*National Institute of Polar Research, Tachikawa, Japan*

⁶*Institute for Marine and Antarctic Studies, University of Tasmania, Hobart, Australia*

⁷*JSPS International Research Fellow, Japan Society for the Promotion of Science, Tokyo, Japan*

⁸*Research and Information Center, Tokai University, Tokyo, Japan*

⁹*School of Biological Sciences, Tokai University, Sapporo, Hokkaido, Japan*

¹⁰*Graduate University for Advanced Studies (SOKENDAI), Tachikawa, Japan*

*Corresponding author: (email: daiki.nomura@fish.hokudai.ac.jp)

Abstract

We elucidated the effects of snow and remineralization processes on nutrient distributions in multi-year landfast sea ice (fast ice) in Lützow-Holm Bay, East Antarctica. Based on sea-ice salinity, oxygen isotopic ratios, and thin section analyses, we found that the multi-year fast ice grew upward due to the year-by-year accumulation of snow. Compared to ice of seawater origin, nutrient concentrations in shallow fast ice were low due to replacement by clean and fresh snow. In deeper ice of seawater origin (the lower half of the multi-year fast ice column), remineralization was dominated by the degradation of organic matter. In addition, denitrification was detected in the low brine volume fraction and impermeable multi-year ice due to their disconnection from gas and water exchanges with the atmosphere and under-ice water. By comparison with first-year ice, we develop a conceptual image of the evolution of nutrient concentrations, with biological uptake dominating in relatively young ice and remineralization dominating in older, multi-year ice under the physical process of brine drainage.

Plain language summary

Multi-year landfast sea ice (fast ice) is abundant around the coast of Antarctica. Fast ice is an important component of Antarctic coastal marine ecosys-

tems, providing a prolific habitat for ice algal communities. Although nutrients are essential for biological productivity within sea ice, the status of nutrients and processes affecting nutrient concentrations were previously not known for multi-year fast ice. Here, we collected sea-ice cores from multi-year fast ice in Lützow-Holm Bay, East Antarctica, and we analyzed nutrient concentrations using physical and biogeochemical parameters. Nutrient concentrations in the upper parts of the sea ice decreased due to the accumulation of clean (nutrient-free) snow through melting and refreezing processes, contributing to upward ice growth. In deeper parts of the sea ice, nutrient concentrations were greatly affected by biological processes such as remineralization during the degradation of organic matter within sea ice.

Keywords:

Sea ice, snow, nutrients, remineralization, multi-year landfast ice, fast ice, East Antarctica

Key point:

Antarctic multi-year landfast ice grew upward due to the year-by-year accumulation of snow

Nutrient concentrations decreased in the upper sea ice due to the replacement by clean snow

In deeper sea ice, remineralization by degradation of organic matter drove nutrient concentrations

1. Introduction

Multi-year landfast ice (fast ice) is abundant around Syowa Station, Lützow-Holm Bay, East Antarctica (Fraser et al., 2012; 2021; Ushio, 2006). This region receives relatively abundant snowfall and high snow accumulation rates over sea ice (Toyota et al., 2016), forming ice of snow origin such as ‘snow ice’ and ‘superimposed ice’ (Kawamura et al., 1997; Nomura et al., 2012, 2018). In general, snow ice forms under the weight of overlying snow on the sea ice, which depresses the ice–snow interface to below sea level; the subsequent flow of seawater into the snow forms a flooded slush layer that then freezes (Maksym & Jeffries, 2000; Sturm & Massom, 2017). During summer, the snow surface melts in the sunlight and warmer temperatures; the snowmelt then infiltrates the slush layer, decreasing its salinity to well below that of seawater (Nomura et al., 2012, 2018). However, in Lützow-Holm Bay, ice thickening due to the formation of snow ice does not necessarily depress the sea ice–snow interface to below sea level. In this case, when the shallowest snow thaws, the snowmelt permeates the sea ice–snow interface below the freezing point and refreezes, forming superimposed ice (Kawamura et al., 1997; Nomura et al., 2018). The difference between these two types of ice is that snow ice forms from a mixture of snow and seawater, whereas superimposed ice forms only from snow. Due to the abundance of

snow around Syowa Station, it is considered that superimposed ice forms only after sea ice stops sinking under the weight of overlying snow.

Because sea ice forms from seawater, the biogeochemical components in seawater are preserved in sea ice upon its formation. Various changes affect these biogeochemical components, particularly nutrients, after their entrapment in sea ice. For example, NO_3^- , NO_2^- , NH_4^- , and PO_4^{3-} are involved in biological activities such as photosynthesis and remineralization by microorganisms such as bacteria and phytoplankton, notably ice algae, phytoplankton inhabiting the bottommost layers of sea ice (Arrigo et al., 2010; Gleitz et al., 1995). Diatoms, which are considered to be the main primary producers in sea ice, also use Si(OH)_4 to form their frustules (Arrigo et al., 2010; Tréguer & De La Rocha, 2013). In addition to the consumption of nutrients by ice algae, high-salinity and high-nutrient brines are discharged beneath the sea ice over time (Fripiat et al., 2017). Nutrients are resupplied by remineralization during the degradation of organic matter within sea ice by heterotrophs and bacteria (Roukaerts et al., 2021; Thomas et al., 1995) as well as by convection through brine channels (Nomura et al., 2009; Vancoppenolle et al., 2010).

It has also been reported that denitrification reactions occur within sea ice (Kaartokallio, 2001; Nomura et al., 2010; Nomura et al., 2018; Rysgaard & Glud, 2004; Rysgaard et al., 2008; Staley & Gosink, 1999). Sea ice can locally become isolated from the atmosphere and under-ice water, resulting in the depletion of oxygen in the ice and promoting denitrification. Previous studies have reported denitrifying bacteria in Antarctic sea ice (Staley & Gosink, 1999) and have suggested or confirmed denitrification reactions in Arctic and sub-Arctic sea ice, e.g., in the Baltic Sea (Kaartokallio, 2001), along the coast of Greenland and Arctic Ocean (Rysgaard & Glud, 2004; Rysgaard et al., 2008), and in the southern Sea of Okhotsk (Nomura et al., 2010).

Primary production is also active in the nutrient-rich, seawater-flooded slush layer formed at the sea-ice surface during the formation of snow ice (Ackley & Sullivan, 1994; Fritsen et al., 1994; Kattner et al., 2004; Nomura et al., 2018; Schnack-Schiel et al., 2001). The snow cover on sea ice therefore plays an important role in increasing the productivity of the sea-ice surface (e.g., Saenz and Arrigo, 2014). When the slush freezes, it becomes snow ice, preserving the biogeochemical formations (Kawamura et al., 1997; Nomura et al., 2018). However, the thick sea ice produced in areas receiving abundant snowfall prevents seawater from flowing along the sea-ice surface, limiting productivity; therefore, snow cover can only increase productivity in the slush layer by so much. Furthermore, whereas snow-free sea ice grows downward due to the freezing of seawater, snow-covered sea ice grows upward due to the formation of snow ice and superimposed ice, such that melting progresses from the bottom of the sea ice (Kawamura et al., 1997). Thus, snow cover greatly affects biogeochemical distributions and circulations in sea ice.

Fast ice plays a crucial role in the biogeochemical cycle and marine ecosystem functions of the Antarctic coast because it hosts well-established habitats (in

terms of nutrients, temperature, salinity, and light) for stable primary production (Arrigo, 2017; Wongpan et al., 2018). Because primary production in sea ice promotes the proliferation of zooplankton, i.e., secondary and higher producers, it is essential to know how the biogeochemical composition of fast ice changes. However, although many studies have explored nutrient changes in first-year fast ice (e.g., Lim et al., 2019), research remains limited on biogeochemical components in multi-year fast ice that has continued to grow year-by-year. Furthermore, of the few studies that have focused on nutrient dynamics in multi-year land-fast ice (Nomura et al., 2018), their sea-ice samples provide only single-year observations. Therefore, no study has rigorously investigated the processes of nutrient cycling in multi-year sea ice. Because multi-year fast ice has remained fixed to the coast for a long time (Fraser et al., 2012; 2021), it is ideal for examining the temporal evolution and interannual variation of nutrient concentrations and other biogeochemical parameters in a single location.

The Japanese Antarctic Research Expedition (JARE) has monitored the multi-year fast ice near Syowa Station in Lützow-Holm Bay for many years. In recent years, it has been reported that the multi-year fast ice in Lützow-Holm Bay periodically breaks up and flows out to sea (Aoki, 2017; Ushio, 2006). Because ice thickness reflects the strength of sea ice, sea-ice monitoring efforts have recently collected annual ice core samples from an easily accessible and stable area near Syowa Station, ideal samples for tracking year-to-year changes. Additionally, because new ice formed after the partial ice breaks up and outflows, it is possible to track the cycling of biogeochemical components through the ice by comparing early stages of sea-ice formation with the initial multi-year ice.

In this study, we analyzed multi-year fast ice near Syowa Station to evaluate the effects of snow and biological activity on the vertical distribution of nutrient concentrations in sea ice. We investigated the factors affecting nutrient concentrations in the multi-year fast ice by collecting and comparing nearby first-year ice. Specifically, based on annual sampling during the 56–60th JAREs (2015–2019), we analyzed: (1) the structure and formation process of ice using thin section photographs, salinity, and stable oxygen isotopic ratios (^{18}O); (2) the vertical distribution of nutrient concentrations in the fast ice; (3) the influence of snow (based on snow fraction) on nutrient distributions; and (4) the cycling of nutrient concentrations due to biological activity within fast ice using sea-ice salinity, nutrient concentrations, and nutrient ratios as proxies.

2. Materials and Methods

2.1. Sampling and sample preparation

Sea ice observations of multi-year fast ice were performed during the summers of 2015–2019 as part of the 56–60th JAREs. Sea-ice cores were collected near Syowa Station (Figure 1, Table 1). Ice cores were sampled using an ice corer (Geo Tecs Co., Ltd., Chiba, Japan) with an internal diameter of 0.09 m. Cores were immediately placed in tubular polyethylene bags and kept horizontal in a cooler box along with refrigerants to maintain below-freezing temperatures

and minimize brine drainage from the core. Fresh snow samples were collected with an acid-washed polycarbonate shovel and placed into polyethylene zipper storage bags. For comparison, four cores of first-year fast ice were also collected (Table 1).

Note that in Table 1, cores JARE58_KU3 and JARE59_2 were sampled only the top 2.78 and 2.47 m of ice, respectively, because the ice was too thick to sample the entire column. As part of the ice monitoring at Syowa Station, an electromagnetic (EM) ice thickness survey was conducted near our coring site in January 2018; the ice thickness was approximately 6 m.

Under-ice water was collected through the coring holes with a Teflon water sampler (GL Science Inc., Japan) from 1 m below the bottom of the sea ice during JARE60 (2019). Water samples were collected approximately 30 minutes after the ice core at JARE60_KU2 was drilled to avoid potential artifacts associated with the disturbance of drilling. All collected water samples were subsampled into (1) a 120-mL glass vial (Maruemu Co., Ltd., Osaka, Japan) for salinity measurement, (2) a 15-mL glass screw-cap vial (Nichiden-Rika Glass Co. Ltd, Kobe, Japan) for ^{18}O analysis, (3) a 10-mL polyethylene screw-cap vial (Eiken Chemical Co. Ltd, Tokyo, Japan) for measurement of major inorganic nutrient concentrations ($\text{NO}_3^- + \text{NO}_2^-$, NH_4^+ , PO_4^{3-} , and Si(OH)_4), and (4) a 500-mL Nalgene polycarbonate bottle (Thermo Fisher Scientific Inc., Waltham, MA, USA) for measurement of chlorophyll a (chl.a) concentrations. All samples were returned to the laboratory of the Japan Maritime Self-Defense Force icebreaker *Shirase* immediately after sampling. Samples for salinity and ^{18}O analyses were stored in a refrigerator at $+4^\circ\text{C}$. Samples for nutrient analyses were stored in a freezer at -30°C . Samples for chl.a measurements were immediately filtered through 25-mm-diameter Whatman GF/F filters. Pigments on the filters were then extracted with *N,N*-dimethylformamide (Suzuki & Ishimaru, 1990) for more than 24 h and stored at -30°C until analysis.

Sea ice cores were stored in a freezer at -30°C during their transport to the Institute of Low Temperature Science (ILTS), Hokkaido University, Sapporo, Japan. In the ILTS cold room (-15°C), cores were split lengthwise into two halves with an electric band saw: the first half for ice textural analysis and the second half for measurements of bulk ice salinity, ^{18}O composition, nutrient and chl.a concentrations, and cell counting for ice algae community assemblage.

2.2. Textural analysis and the age of an ice core

We examined ice thin sections to analyze ice growth processes. Ice sections 0.7 cm thick were attached to glass plates and planed down to 0.1-cm thickness with a microtome (Model SM2400, Leica Microsystems, Wetzlar, Germany). We photographed ice crystallographic structures by illuminating the thin sections placing on a light table in between the cross-polarized sheets.

Ice growth processes were examined based on our classification of the ice cores as granular, columnar, or mixed ice (Eicken & Lange, 1989). Granular ice was subdivided into frazil ice and snow ice (or superimposed ice) based on its ^{18}O

value. We defined snow ice or superimposed ice as granular ice with a ^{18}O value below that of its parent seawater (see Granskog et al., 2004; Jeffries et al., 1994), taken as the ^{18}O value of the under-ice seawater sampled during JARE60 (-0.7‰). Although coarse-grained clear ice (large, polygonal, granular crystal textures) and fine-grained opaque ice originate from superimposed and snow ice, respectively (Kawamura et al., 1997), we hereafter refer to both as “snow-origin ice” unless obviously clear ice was observed in thin section, which was described as superimposed ice. We note that the analyzed snow was thawed and moistened, and refrozen ice was observed. The snow crystal structure was an aggregate of fine particles in thin section but did not contain salt (salinity = 0); the structure sometimes became sandwiched between superimposed ice, which we refer to as “fine-grained superimposed-ice” (see details in Section 3.1).

The ice type was determined according to WMO (2014) after checking satellite imagery of the sampling point, visually observing the sea ice condition at Syowa Station, and confirming the vertical profile of textural parameters (including thin section analysis) in the core. First-year ice was collected in locations where fast ice had broken up and flowed out during the previous year (Figure 1d–f), and sea ice had reformed within the year prior to collection. According to Syowa Station records, multi-year ice has not been broken up and flowed out since March 1980 (Higashi et al., 1982; Ushio, 2006).

2.3. Sample analysis

To measure ice salinity, water ^{18}O composition, and nutrient and chl.a concentrations, the second half of the core was trimmed into a rectangle with a cross-section of $4.5\text{ cm} \times 2.5\text{ cm}$ and then sliced into 12–18-cm-thick sections. To avoid contamination during sampling and handling, the outer 0.3 cm of each ice section was removed with a stainless steel plane. The ice sample was then put in a polyethylene bag (GL Science Inc., Japan) and melted in a cold room at $+4^\circ\text{C}$. Snow samples were melted in the same bags and in the same cold room. Snow and sea ice melt samples were analyzed with the same protocol as for under-ice water samples, except for nutrient analyses; in snow and sea ice melt samples, nutrient samples were filtered with a 0.22- μm Durapore PVDF Membrane filter (MILLEX GV Filter unit, Merck Millipore Ltd., Germany). For ice algae community assemblage, sea ice melt water for the bottom ice section for JARE60 KU1 and 2 was subsampled into 100-mL polyethylene screw-cap vial (AS ONE Corporation, Tokyo, Japan), and an aliquot of 5-mL of Lugol’s solution was added to preserve the sample.

The salinities of under-ice water and melted snow and sea-ice samples were measured with a conductivity sensor (Cond 315i, WTW GmbH, Germany). The oxygen isotopic ratio of under-ice water and melted snow and sea-ice samples were determined by mass spectrometry (DELTA plus; Finnigan MAT, San Jose, CA) via the equilibration method. The ^{18}O value (‰) of a sample was calculated against the $^{18}\text{O}/^{16}\text{O}$ ratio of standard mean ocean water. The standard deviation (SD) on ^{18}O values was 0.026‰ based on analyses of 10 subsamples of a reference water with $^{18}\text{O} = 0.241\text{‰}$. Nutrient concentrations (NO_3^-

+ NO_2^- , NH_4^+ , PO_4^{3-} , and $\text{Si}(\text{OH})_4$) were measured with an auto-analyzer system (Quattro; Bran+Luebbe, Norderstedt, Germany) according to the spectrophotometric method reported by the Joint Global Ocean Flux Study (JGOFS, 1994). SDs on nutrient concentrations were 0.53, 0.18, 0.07, and 1.1 mol L^{-1} for $\text{NO}_3^- + \text{NO}_2^-$, NH_4^+ , PO_4^{3-} , and $\text{Si}(\text{OH})_4$, respectively, calculated from 10 subsamples of a reference water sample with respective known concentrations of 52.53, 9.99, 5.00, and 117.5 mol L^{-1} . Chl.a concentrations were determined by a fluorometer (Model 10AU, Turner Designs, Inc., Sunnyvale, CA) using the method of Parsons et al. (1984). Standards (0.28–282.30 g L^{-1} chl.a) prepared from a liquid chl.a standard (Wako Pure Chemical Industries Ltd., Osaka, Japan) by stepwise dilution with *N,N*-dimethylformamide were used to calibrate the fluorometer before analyses. The cell counting for ice algae community assemblage was examined with a microscope (Olympus, BH-T, Tokyo, Japan) 10 \times oculars and 40 \times objective.

2.4. Mass fraction of snow in snow-origin ice

To determine the mass fraction of snow in snow-origin ice (hereafter, ‘snow fraction’), we used a mass balance equation (Jeffries et al., 1994, 2001):

$$f_{\text{snow}} + f_{\text{sea}} = 1, (1)$$

$$f_{\text{snow}} \delta_{\text{snow}} + f_{\text{sea}} \delta_{\text{sea}} = \delta_{\text{obs}}, (2)$$

where f represents the mass fraction of snow or seawater, δ represents their ^{18}O values, and the subscripts ‘snow’, ‘sea’, and ‘obs’ indicate snow, under-ice seawater, and observed bulk ice, respectively. For δ_{sea} , we used the parent water ^{18}O values collected during JARE60 (-0.7‰) rather than the value that includes the fractionation factor during seawater freezing (see Jeffries et al., 1994, 2001) because we assumed that all granular ice with a ^{18}O value below that of the under-ice water was snow ice (Granskog et al., 2004; Jeffries et al., 1994). In addition, we assumed that granular ice with a ^{18}O value below that of under-ice water was snow-origin ice. Based on the result obtained from the JARE60 samples, we used $\delta_{\text{sea}} = -0.7\text{‰}$. δ_{snow} was taken as the mean ^{18}O value of snow samples (-20.3‰) collected during JARE56–60. We estimated the errors on the snow fractions to be less than 12.2% by calculating the effect of changing the snow ^{18}O values by their standard deviation ($\pm 2.1\text{‰}$).

3. Results and Discussion

3.1. Vertical profiles of sea-ice salinity, ^{18}O values, and sea-ice structure

Over the study period, first-year ice and snow on ice were ranged between 1.32 and 1.55 m thick and 0.05 and 0.20 m deep, respectively (Table 1). Multi-year ice and snow on ice were ranged between >2.47 and ca. 6 m thick and 0.34 and 1.68 m deep, respectively (Table 1). Figure 2 shows vertical salinity and ^{18}O profiles for both first-year and multi-year ice cores. The salinity of the first-year ice was close to 0 near the ice–snow interface and increased with increasing depth until reaching a maximum, after which it decreased with increasing depth (Figure 2a). Salinity again increased at the bottom of some cores, giving the salinity

profiles of first-year ice a characteristic “C” shape (excluding the topmost low-salinity ice). ^{18}O values in first-year ice were as low as -20‰ near the ice–snow interface (Figure 2b) and increased with increasing depth, reaching values above that of the under-ice water (-0.7‰) by the middle of the core. The profiles record low salinity and ^{18}O values near the ice–snow interface because the cores were collected in late summer (Table 1), by which time the accumulated snow had become snow-origin ice due to melting of the sea-ice surface in above-freezing air temperatures and strong sunlight (Nomura et al., 2011a). Furthermore, in late summer, the nighttime air temperature becomes negative in this area (Nomura et al., 2011a), freezing any water that had melted during the day. Because the snow depths were quite small relative to the ice thickness (Table 1) and the salinity was extremely low, especially near the ice surface, we considered that the ice formed was superimposed ice rather than snow ice because snow ice should only form when the snow depth is large with respect to the ice thickness (Maksym & Jeffries, 2000; Sturm & Massom, 2017). Indeed, in thin section, we observed large, polygonal, granular crystal textures (photograph in Figure 2a) characteristic of superimposed ice (Kawamura et al., 2004) near the ice surface.

The salinity of multi-year ice was 0 from the ice–snow interface to depths of around -100 cm (Figure 2c), then generally increased with increasing depth. The maximum observed salinity was 3.8 in core JARE58_KU3. At the bottom of the ice column, salinities were about 1–3. The ^{18}O values of sea ice were similar to those of the overlying snow from the ice–snow interface to depths of -200 cm (Figure 2d), then increased with increasing depth. The ice ^{18}O values reached that of the under-ice water at depths of -300 cm in core JARE56_NMS and -380 cm in core JARE57_P31.5. These results indicate that snow cover affects to a maximum depth of -380 cm.

Figure 3 compares the salinity and ^{18}O profiles of multi-year ice core JARE57_P31.5 to the interpreted ice structure and photographs of representative thin sections. Up to a depth of -117 cm, salinity and ^{18}O were maintained at 0 and -20‰ , respectively (Figure 3a). Because most of this upper section of the core comprised coarse-grained (large polygonal) granular ice, we concluded that it was superimposed ice (Figure 3b, c; Kawamura et al., 2004). In addition, fine-grained granular ice was present at depths from -100 to -117 cm (Figure 3b, d). Based on the salinity and textures down to a depth of -117 cm, it is unlikely that snow ice was formed by the infiltration of seawater. In addition, the small crystal size at depths of -100 to -117 cm (Figure 3d) indicated that this ice corresponded to superimposed ice formed from wet, unmelted snow that had re-frozen at negative temperatures, which we refer to as fine-grained superimposed ice.

At depths of -117 to -170 cm, sea ice salinity and ^{18}O values increased (Figure 3a) and the ice structure became columnar (Figure 3b, e), with larger grains than in the overlying fine-grained granular ice. However, the columns were neither as wide nor as vertically extended as those of columnar ice formed from seawater (Figure 3f, see next paragraph). Based on the salinity and ^{18}O values of the ice,

this structural transformation occurred because the ice formed snowmelt with a slight seawater influence. Therefore, we suspect that this ice formed from a refrozen mixture of seawater and snowmelt. Indeed, in this area, it has been reported that a layer of superimposed ice forms under the snow, and that a slush layer (puddles in the sea ice that are likely snowmelt ponds) forms under the superimposed ice (Nomura et al., 2018). In that study, the salinity and ^{18}O values of the slush layer were 1 and -20‰ , respectively, because snowmelt was supplied mainly from above (as snow or superimposed ice), and a small amount of seawater was transported upwards through a crack and then horizontally transported into the slush layer. The salinity and ^{18}O values of their slush layer were approximately the same as those of the columnar ice measured in our study (Figure 3a). The slush layer is about 10–20 cm thick (Nomura et al., 2012) and, in winter, is thought to freeze from the top faster than columnar ice develops at greater depths in the sea ice. Therefore, we consider that the columns in this zone are thinner and shorter than those in columnar ice formed from seawater (Figure 3e, f) because the freezing speed affects the crystal structure with slower growth on the c-axis direction than the basal plane (e.g., Hilling, 1959). Both fine- and coarse-grained granular ice have random crystal texture (Figure 3c-d). As the freezing develops, geometric selection sweep away crystals with c-axis perpendicular to the ice–ocean interface (Kolmogorov, 1949). The crystals in the slush layer that have the c-axes pointing radially (perpendicular to the ice–ocean interface) grow faster than those with c-axes pointing parallel to the ice–ocean interface. Therefore, it seems likely that the columns in slush layer was thinner due to the initial stage of geometric selection and shorter than those in columnar ice formed from seawater constrained by the slush layer thickness. In this study, we refer to this particular columnar ice texture as ‘slush-origin columnar ice’, in contrast to columnar ice formed from seawater (‘seawater-origin columnar ice’).

A distinct unit comprising a fine-grained granular ice layer overlying a coarse-grained granular ice layer was sandwiched between slush-origin columnar ices at depths of -170 to -190 cm. This unit formed because superimposed ice formed under the snow (Nomura et al., 2018) and a slush layer below it. This is thought to reflect greater amounts of snowfall in following years, with the slush layer being repeatedly formed and refrozen. The underlying slush-origin columnar ice layer extended to a depth of -300 cm, where the ice structure became columnar, but of seawater origin.

In contrast, the first-year ice was predominantly columnar (Table 2). From the above discussion, snow accumulates on the ice year-by-year, causing flooding and slush formation, then refreezing in winter. Furthermore, when the ice thickness increases, flooding ceases, and the snowmelt is refrozen. This upward growth is characteristic of Antarctic multi-year ice (Kawamura et al., 1997; Nomura et al., 2018), although the collection of a nearly 5-m sea-ice core has never been reported. This is thus the first time that the characteristics of the upward growth of multi-year fast ice have been shown based on multiple years of repeated ice coring.

3.2. Vertical profiles of sea-ice nutrient concentrations

Figure 4 presents vertical profiles of NO_3^- , PO_4^{3-} , and Si(OH)_4 concentrations in first-year and multi-year ice. NO_3^- concentrations in first-year ice varied widely with depth in core JARE60_KU1 ($0\text{--}2.6\text{ mol L}^{-1}$), but were in the range $0\text{--}1\text{ mol L}^{-1}$ in other cores (Figure 4a). NO_3^- concentrations in multi-year ice were $0\text{--}0.5\text{ mol L}^{-1}$ in the top half of the ice column, and the first maximum at those depths, peaking at $\sim 1.0\text{ mol L}^{-1}$, was observed at a depth of $\sim 200\text{ cm}$ in some ice cores (Figure 4e). At greater depths in the cores, NO_3^- concentrations were as low as $0\text{--}0.5\text{ mol L}^{-1}$, and the maximum of 3.5 mol L^{-1} was observed at the bottom of the sea ice. In first-year ice, PO_4^{3-} concentrations varied between cores but were fairly constant at shallow to medial depths, and were greatest at the bottom of the ice column (Figure 4b). In multi-year ice, however, PO_4^{3-} concentrations were extremely low from the ice-snow interface to a depth of $\sim 150\text{ cm}$, and the maximum concentration of 1.6 mol L^{-1} was observed at a depth of $\sim 360\text{ cm}$ in core JARE57_P31.5 (Figure 4f). Si(OH)_4 concentrations varied little with depth in the first-year ice (Figure 5c), but mimicked the PO_4^{3-} profile in multi-year ice.

Our results show clear differences between the vertical nutrient profiles in first-year and multi-year ice (Figure 4). First-year ice generally has the lowest nutrient concentrations in snow, and concentrations increase slightly in sea ice, but they remain relatively constant until peaking at the bottom of the ice column (Figure 4a–c). In contrast, in multi-year ice, the upper half of the ice column was nutrient poor, similar to the overlying snow, and nutrient concentrations increased from the middle of the column to maximum concentrations in the bottom half of the column (Figure 4e–g). Because nutrient concentrations in snow were consistently low (Figure 4e–g), the upper half of the multi-year ice columns, comprising snow-origin ice, also contained low nutrient concentrations.

Such vertical nutrient distributions are considered to be closely related to physical characteristics such as salinity, ^{18}O value, and ice structure (discussed in Section 3.3). Furthermore, the high concentrations of various nutrients at the bottom of both first-year and multi-year ice columns are most likely related to the high concentrations of chl.a in the bottommost ice column (Figure 4d, h), which undergoes both photosynthesis by the ice algae and remineralization by the degradation of organic matter (Roukaerts et al., 2021; Rysgaard & Glud, 2004; Rysgaard et al., 2008; Thomas et al., 1995) and receives nutrients supplied from under-ice water (Nomura et al., 2009; Vancoppenolle et al., 2010).

The mean value and standard deviation of each nutrient component and physical parameter are reported in Table 3 for snow and each sea-ice type. Nutrient and chl.a concentrations were low in snow and superimposed ice and particularly high in columnar ice. In contrast, snow-ice, which is affected by both snow and seawater, had intermediate nutrient and chl.a concentrations. Our results clearly indicate that a greater snow influence results in lower nutrient and chl.a concentrations in sea ice.

3.3. Variations of nutrient concentrations in snow-origin ice

In Section 3.2, we showed that in multi-year ice columns, nutrient concentrations were lower in snow-origin ice than in columnar ice (Table 3). To quantify the effect of snow on sea-ice nutrient concentrations, we calculated the snow fraction in snow-origin ice (Eqs. 1 and 2) to examine the relationship between nutrient concentrations and snow fraction (Figure 5). Although the regressions differed between nutrients, nutrient concentrations tended to decrease with increasing snow fraction. This is due to the purity of the air near Antarctica, being far from areas of human activity, compared to that in the Northern Hemisphere. Therefore, the atmospheric transfer and deposition of nutrients onto sea ice is limited in Antarctica, and the top surface of the sea ice is replaced by clean, snow-origin ice. In contrast, in the Northern Hemisphere, atmospheric transfer and deposition of polluted snow is known to be a potential source of nutrients for sea ice (Granskog & Kaartokallio, 2004; Granskog et al., 2003; Kaartokallio, 2001; Krell et al., 2003; Nomura et al., 2010; 2011b; Rahm et al., 1995).

Although nutrient concentrations were generally negatively correlated with snow fraction (Figure 5a, b, d), NO_3^- concentrations notably differed, with the negative correlation between snow fraction and NO_3^- concentration being markedly low compared to the other nutrients (Figure 5c). Because most of the sea-ice samples in this study were collected near Syowa Station, it is possible that this difference reflects slight pollution from local human activities, such as exhaust from the research facility and/or snowmobiles. Therefore, greater NO_3^- concentrations might be detected (Table 3) in areas of greater snow fraction (Figure 5). However, many NO_3^- data points plot near zero at snow fractions of 30–80% (Figure 5c), potentially indicating that NO_3^- was selectively removed by sea ice, which can be explored by considering denitrification.

Denitrification is a reaction in which NO_3^- is used by bacteria in anoxic conditions to decompose organic substances within sea ice, converting NO_3^- to NO_2^- , NO, and eventually N_2O and N_2 (Kaartokallio, 2001; Nomura et al., 2010; 2011b; 2018; Rysgaard & Glud, 2004; Rysgaard et al., 2008; Staley & Gosink, 1999). Snow-origin ice with small snow fractions, i.e., containing some, albeit small, amounts of seawater, must be older than snow-origin ice not containing any seawater (e.g., superimposed ice) because it is formed at an earlier stage of ice growth. Therefore, we consider that denitrification occurred in the older parts of the ice column where the snow fraction was low (Figure 5c; see Section 3.4 for details).

Due to the formation of snow-origin ice, the snow fraction in multi-year ice increased year-by-year; the mean snow fractions in the top 200 cm of multi-year ice gradually increased from about 88% in 2015 to 100% by 2018 (Figure 6). Therefore, our sea-ice core samples record a steady increase in the contribution of snow to sea ice. Although we have not collected multi-year ice cores at distant locations within the same year (Figure 1c) and thus have not been able to unequivocally determine that there is no spatial bias in our sampling, the nutrient concentrations do not differ greatly between sampling locations, and

the trends and profiles are similar from year to year (Figures 2, 4).

3.4. Variations of nutrient concentrations in seawater-origin ice

Although nutrient concentrations decreased as the snow fraction increased in the topmost sea ice, the distributions of nutrient concentrations differed among the different nutrient components (Figure 5). Therefore, we here evaluate the effects of primary production, organic matter remineralization, and denitrification on nutrient concentrations in both first-year and multi-year ice. We first compared salinity to nutrient concentrations (Figure 7). Sea-ice salinity is a conservative component that only changes through physical processes such as brine drainage or exchange with under-ice water. Therefore, we investigated deviations from a dilution line based on the salinity and nutrient concentrations of the under-ice water collected during JARE60 to understand the biological processes affecting the nutrient distributions in the sea ice. Data plotting above the dilution line indicate nutrient supply due to the remineralization of organic matter by heterotrophic organisms and bacteria in the sea ice or from external sources (e.g., the atmosphere); conversely, data points below the dilution line indicate nutrient uptake by ice algae during primary production (e.g., Thomas et al., 1995).

Si(OH)_4 concentrations in first-year ice locate significantly below the dilution line (Figure 7a), whereas those in multi-year ice are closer to or above the dilution line (Figure 7b). The distributions of chl.a concentrations in the sea ice (Figure 4d, h) indicate the presence of ice algae, which would mainly be diatoms (*Fragilariopsis* sp. and *Pleurosigma* sp. (photo in 4c, d) based on the cell counting for ice algae community assemblage in this study. Indeed, in first-year ice, primary production by diatoms consumes a large amount of Si(OH)_4 (Tréguer & De La Rocha, 2013), and the slow dissolution of diatom frustules should limit the resupply of Si(OH)_4 . In contrast, in multi-year ice, the more protracted dissolution of diatom frustules should resupply Si(OH)_4 , eventually balancing the consumption of Si(OH)_4 by primary production. However, in core JARE57_P31.5 (the deepest core studied), Si(OH)_4 concentrations plot well above the dilution line (Figure 7b), indicating that the resupply of Si(OH)_4 by dissolution of diatom frustules exceeded consumption.

Consistent with a previous sampling at the same location (Nomura et al., 2018), PO_4^{3-} concentrations plot above the dilution line in both first-year and multi-year ice (Figure 7c, d), suggesting resupply by the remineralization of organic matter. The difference in the resupply rates of PO_4^{3-} and Si(OH)_4 is due to the fact that the remineralization of PO_4^{3-} is faster than the dissolution of Si(OH)_4 (Fripiat et al., 2017).

NO_3^- concentrations in most first-year and multi-year ice cores plot below the dilution line (Figure 7e, f). This is expected for first-year ice, in which primary production consumes NO_3^- , as with Si(OH)_4 . In contrast, remineralization in multi-year ice should resupply NO_3^- , although our data do not reflect this. Therefore, we also measured NO_2^- concentrations (Figure 7g, h), which plot

close to the dilution line for first-year ice and are shifted upward for multi-year ice. From this result, we expect that denitrification resupplied NO_2^- over longer periods in the multi-year ice. Indeed, denitrifying bacterial species has been confirmed in Antarctic ice (Staley & Gosink, 1999), which consume NO_3^- and supply NO_2^- (Rysgaard et al., 2008). To better understand the effect of denitrification on the nutrient balance, we calculated DIN:P, the ratio of dissolved inorganic nitrogen (DIN, i.e., $\text{NO}_3^- + \text{NO}_2^-$) to PO_4^{3-} (Figure 8). In most cores, DIN:P values were lower than the Redfield ratio (16:1) and the diatom nutrient uptake ratio (12:1; Takeda, 1998), which is thought to reflect the transformation of NO_3^- and NO_2^- to NO , N_2O , and N_2 by denitrification. The value of N^* , which has been suggested as a proxy for denitrification, is calculated as follows (Gruber & Sarmiento, 1997; Nishioka et al., 2021; Yoshikawa et al., 2006):

$$\text{N}^* = ([\text{NO}_3^-] - 16 [\text{PO}_4^{3-}] + 2.9) \times 0.87. \quad (3)$$

Figure 9 shows N^* profiles of multi-year ice cores JARE56_NMS and JARE57_P31.5. In the lower parts of these cores, N^* values decreased to negative values, reaching a minimum of -12.3 mol L^{-1} in core JARE57_P31.5. Chl.a concentrations are also negatively correlated with N^* . Previous studies have reported that denitrification occurred in oxygen-depleted and organic matter-rich parts of the ice column (Rysgaard et al., 2008). It is therefore reasonable that denitrification occurred at depths where high chl.a concentrations indicate abundant organic matter. To our knowledge, because multi-year ice formed a long time ago in this area (about 40 years ago) (Higashi et al., 1982; Ushio, 2006), salinity is low (Figure 2c), and it is unlikely that brine channels physically drain the ice. Therefore, the air and brine pockets in multi-year ice has remained isolated and disconnected from exchanges with the atmosphere and under-ice water.

To examine the possibility of oxygen exchange through the brine channel network in the sea ice, we focused on the brine volume fraction, a proxy of sea-ice permeability. Calculating brine volume fraction requires the ice temperature and salinity; however, we could not obtain realistic ice temperature data due to the difficulty of inserting a temperature sensor in the sea ice, especially in low-salinity, snow-origin ice. The ice temperature increased during coring due to friction, and the air temperature was often positive (Nomura et al., 2011a), precluding accurate measurement of the sea ice temperature. Therefore, for the brine volume calculation, we assumed the temperature of the sea ice to be -2 to $0 \text{ }^\circ\text{C}$, a reasonable estimate given the positive air temperature. We obtained brine volume fractions, estimated following Leppäranta and Manninen (1988), of $2.4 \pm 2.6\%$ (mean and standard deviation) at all depths in the multi-year ice. Typically, sea ice is considered permeable when the brine volume fraction exceeds 5–7.5% (Golden et al., 1998; Pringle et al., 2009; Zhou et al., 2013). Therefore, the multi-year ice collected in this study was impermeable, precluding any exchanges through brine channels. Therefore, gas components such as oxygen must have been consumed in a closed system within the ice by het-

erotrophic organisms and bacteria. Furthermore, the high density of the hard, low-salinity superimposed ice suggests that gases could not be exchanged with the atmosphere, consistent with previous studies that were unable to detect gas fluxes over superimposed ice at the same location (Nomura et al., 2012, 2013). Therefore, oxygen in the isolated interior sea ice could only have been depleted by heterotrophic organisms and bacteria, promoting denitrification in multi-year ice (Figure 9).

Based on our results, we developed a conceptual illustration for the evolving nutrient concentrations in multi-year fast ice (Figure 10). Immediately after ice formation, when nutrient concentrations have not yet been affected by biological processes, the ratio of salinity to nutrients in the brine discharged from the growing sea ice should plot along the dilution line. Then, during the first year, nutrient uptake by primary production of ice algae decreases nutrient concentrations in the ice while the salinity is reduced by brine convection. Over multiple years, remineralization by the degradation of organic matter resupplies nutrients to the ice while salinity continues to decrease. Once ice algae have exhausted the nutrient supply and decompose, and brine exchange with seawater is established at the bottom of the ice column, the nutrient concentrations surpass the dilution line (Figure 10). In addition, denitrification decreases NO_3^- concentrations and modifies the speciation of nitrogen within the ice.

4. Conclusions

We investigated the effects of snow cover on the vertical distribution of nutrient concentrations and the long-term remineralization process in multi-year, Antarctic fast sea ice. We analyzed thin sections to determine the ice structure, which confirmed the upward growth of fast ice by the year-by-year formation of snow-origin ice. Vertical profiles of biogeochemical components in the ice showed that the high nutrient concentrations of first-year ice are transported downward by the subsequent formation of new snow-origin ice, eventually forming multi-year ice. Because the atmospheric nutrient supply is limited by the purity of Antarctic snow, nutrient concentrations in snow-origin ice decreased as the snow fraction increased.

Nutrients incorporated into columnar ice of seawater origin were dominantly processed by biological uptake via primary production during the first year, but remineralization of degrading organic matter by heterotrophic organisms and bacteria resupplied nutrients to multi-year ice. By analyzing data on first-year and multi-year ice spanning several years in the same area, it was possible for the first time to determine the long-term cycling of nutrients in fast ice.

Acknowledgements

We express our heartfelt thanks to the sea ice team, members of the Japanese Antarctic Research Expedition, the crew of the Japan Maritime Self-Defense Force icebreaker *Shirase*, and cruise members for their support in conducting field work. We thank Prof. K. Kawashima, K. Murakami, H. Nakajima, M. Kiuchi, T. Ishino, and T. P. Tamura for their support in the low temperature

room, M. Hirabayashi for oxygen isotopic analyses, and A. Ooki for nutrient analyses. We benefitted greatly from advice and useful comments on the nutrient data offered by A. Ooki and T. Ishino. This work was supported by grants from the Japan Society for the Promotion of Science (#17H04715, #17H04710, #17H06317, #17H06322, #18F18794, #20H04345, #21H04931),

the Science Program of the Japanese Antarctic Research Expedition (JARE) as Prioritized Research AJ0902 (ROBOTICA), the National Institute of Polar Research (NIPR) through Project Research KP-303 and General Collaboration Project #28-14, #2-13 and the grant for young scientist, the Center for the Promotion of Integrated Sciences of SOKENDAI, and the Joint Research Program of the Institute of Low Temperature Science, Hokkaido University. The data used in this study are available in the supplementary information (Table S1) and are archived in the data repository of the National Institute of Polar Research, Tokyo, Japan (<https://ads.nipr.ac.jp>), is underway.

References

- Ackley, S. F., & Sullivan, C. W. (1994). Physical controls on the development and characteristics of Antarctic sea ice biological communities— A review and synthesis. *Deep Sea Research Part I: Oceanographic Research Papers*, 41, 1583–1604. [https://doi.org/10.1016/0967-0637\(94\)90062-0](https://doi.org/10.1016/0967-0637(94)90062-0)
- Aoki, S. (2017). Breakup of land-fast sea ice in Lützow-Holm Bay, East Antarctica, and its teleconnection to tropical Pacific sea surface temperatures. *Geophysical Research Letters*, 44, 3219–3227. doi:10.1002/2017GL072835
- Arrigo, K. R., Mock, T., & Lizotte, M. P. (2010). Primary producers and sea ice. In D.N. Thomas., & G.S. Dieckmann. (Eds.), *Sea Ice (Second edition)*, Wiley-Blackwell, Oxford, 283-325. <https://doi.org/10.1002/9781444317145.ch8>
- Arrigo, K. R. (2017). Sea ice as a habitat for primary producers. In D. N. Thomas (Ed.), *Sea ice (Third edition)*, 352– 369. <https://doi.org/10.1002/9781118778371.ch14>
- Eicken, H., & Lange, M. (1989). Development and properties of sea ice in the coastal regime of the southern Weddell Sea. *Journal of Geophysical Research: Oceans*. 94, 8193–8206. <https://doi.org/10.1029/JC094iC06p08193>
- Fraser, A. D., Massom, R. A., Michael, K. J., Galton-Fenzi, B. K., & Lieser, J. L. (2012). East Antarctic landfast sea ice distribution and variability, 2000–08. *Journal of Climate*, 25, 1137–1156. DOI:10.1175/JCLI-D-10-05032.1
- Fraser, A. D., Massom, R. A., Handcock, M., Reid, P., Ohshima, K. I., Raphael, M., et al. (2021). Eighteen-year record of circum-Antarctic landfast-sea-ice distribution allows detailed baseline characterisation and reveals trends and variability. *The Cryosphere*, 15, 11, 5061–5077. <http://dx.doi.org/10.5194/tc-15-5061-2021> Retrieved from <https://escholarship.org/uc/item/4gb0869t>
- Fripiat, F., Meiners, K. M., Vancoppenolle, M., Papadimitrio, S., Thomas, D.N., Ackley, S. F., et al. (2017). Macro-nutrient concentrations in Antarctic pack ice:

- Overall patterns and overlooked processes. *Elementa Science of Anthropocene*, 5-13. <https://doi.org/10.1525/elementa.217>
- Fritsen, C. H., Lytle, V. I., Ackley, S. F., & Sullivan, C. W. (1994). Autumn bloom of Antarctic pack-ice algae. *Science*, 266, 782–784. DOI: 10.1126/science.266.5186.782
- Gleitz, M., Vonderlo, M. R., Tomas, D. N., Dieckmann, G. S., & Millero, F. J. (1995). Comparison of summer and winter inorganic carbon, oxygen and nutrient concentrations in Antarctic sea ice brine. *Marine Chemistry*, 51, 81–89. [https://doi.org/10.1016/0304-4203\(95\)00053-T](https://doi.org/10.1016/0304-4203(95)00053-T)
- Golden, K. M., Ackley, S. F., & Lytle, V. I. (1998). The percolation phase transition in sea ice. *Science*, 282, 2238–2241. DOI: 10.1126/science.282.5397.2238
- Granskog, M. A., Kaartokallio, H., & Shirasawa, K. (2003). Nutrient status of Baltic Sea ice: evidence for control by snow–ice formation, ice permeability, and ice algae. *Journal of Geophysical Research: Oceans*, 108 (C8), 3253. doi:10.1029/2002JC001386
- Granskog, M. A., & Kaartokallio, H. (2004). An estimation of the potential fluxes of nitrogen, phosphorus, cadmium and lead from sea ice and snow in the northern Baltic Sea. *Water Air Soil Pollution*, 154, 331–347.
- Granskog, M. A., Leppäranta, M., Kawamura, T., Ehn, J., & Shirasawa, K. (2004). Seasonal development of the properties and composition of landfast sea ice in the Gulf of Finland, the Baltic Sea. *Journal of Geophysical Research: Oceans*, 109 (C2), C02020. DOI:10.1029/2003JC001874
- Gruber, N., & Sarmiento, J. L. (1997). Global patterns of marine nitrogen fixation and denitrification. *Global Biogeochemical Cycles*, 11, pp. 235–266. <https://doi.org/10.1029/97GB00077>
- Higashi, A., Goodman, D. J., Kawaguchi, S., & Mae, S. (1982). The cause of the breakup of fast ice on March 18, 1980 near Syowa station, East Antarctica. *Memoirs of National Institute of Polar Research*, 24, 222–231.
- Hillig, W. (1959). Kinetics of solidification from nonmetallic liquids. In Kingery W, ed. *Kinetics of high-temperature processes*. MIT Press, Cambridge, MA, 127–135.
- Jeffries, M., Shaw, R., Morris, K., Veazey, A., & Krouse, H. (1994). Crystal structure, stable isotopes (^{18}O), and development of sea ice in the Ross, Amundsen, and Bellingshausen seas, Antarctica. *Journal of Geophysical Research: Oceans*, 99 (C1), 985–995. <https://doi.org/10.1029/93JC02057>
- Jeffries, M. O., Krouse, H. R., Hurst-Cushing, B., & Maksym, T. (2001). Snow-ice accretion and snow-cover depletion on Antarctic first-year sea-ice floes. *Annals of Glaciology*, 33, 51–60.

- Kaartokallio, H. (2001). Evidence for active microbial nitrogen transformations in sea ice (Gulf of Bothnia, Baltic Sea) in midwinter. *Polar Biology*, 24:21–28. <https://doi.org/10.1007/s003000000169>
- Kattner, G., Thomas, D. N., Haas, C., Kennedy, H., & Dieckmann, G. S. (2004). Surface ice and gap layers in Antarctic sea ice: Highly productive habitats. *Marine Ecology Progress Series*, 277, 1–12.
- Kawamura, T., Ohshima, K. I., Takizawa, T., & Ushio, S. (1997). Physical, structural, and isotopic characteristics and growth processes of fast sea ice in Lützow-Holm Bay-Holm Bay, Antarctica. *Journal of Geophysical Research: Oceans*, 102, 3345–3355.
- Kawamura, T., Jeffries, M., Tison, J., & Krouse, H. (2004). Superimposed-ice formation in summer on Ross Sea pack-ice floes. *Annals of Glaciology*, 39, 563–568. doi:10.3189/172756404781814168
- Kolmogorov, A. N. (1976). Geometric selection of crystals. Doklady Akad Nauk Minerologiya USSR.
- Krell, A., Ummenhofer, C., Kattner, G., Naumov, A., Evans, D., Dieckmann, G. S., et al. (2003). The biology and chemistry of land fast ice in the White Sea, Russia—A comparison of winter and spring conditions. *Polar Biology*, 26, 707–719.
- Leppäranta, M., & Manninen, T. (1988). The brine and gas content of sea ice with attention to low salinities and high temperatures (Internal Rep. 88-2). Helsinki, Finland: Finnish Institute of Marine Research.
- Lim, S. M., Moreau, S., Vancoppenolle, M., Deman, F., Roukaerts, A., Meiners, K. M., et al. (2019). Field observations and physical-biogeochemical modeling suggest low silicon affinity for Antarctic fast-ice diatoms. *Journal of Geophysical Research: Oceans*. 124, 11, 7837–7853. <https://doi.org/10.1029/2018jc014458>
- Maksym, T., & Jeffries, M. O. (2000). A one-dimensional percolation model of flooding and snow-ice formation on Antarctic sea ice. *Journal of Geophysical Research: Oceans*, 105(C11), 26313–26331. <https://doi.org/10.1029/2000JC900130>
- Nishioka, J., Hirawake, T., Nomura, D., Yamashita, Y., Ono, K., Murayama, A., et al. (2021). Iron and nutrient dynamics along the East Kamchatka Current, western Bering Sea Basin and Gulf of Anadyr. *Progress in Oceanography*, 198, 10.1016/j.pocean.2021.102662, 2021.
- Nomura, D., Takatsuka, T., Ishikawa, M., Kawamura, T., Shirasawa, K., Yoshikawa-Inoue, H. (2009). Transport of chemical components in sea ice and under-ice water during melting in the seasonally ice-covered Saroma-ko Lagoon, Hokkaido, Japan. *Estuarine, Coastal and Shelf Science*, 81, 201–209.
- Nomura, D., Nishioka, J., Granskog, M. A., Krell, A., Matoba, S., Toyota, T., et al. (2010). Nutrient distributions associated with snow and sediment-laden

layers in sea ice of the southern Sea of Okhotsk. *Marine Chemistry*, 119, 1–4. <https://doi.org/10.1016/j.marchem.2009.11.005>

Nomura, D., Simizu, D., Shinagawa, H., Oouchida, C., & Fukuchi, M. (2011a). Biogeochemical properties of water in surface ponds on Antarctic fast ice and relationship to underlying sea ice properties. *Journal of Glaciology*, 57, 205, 1–9. DOI: <https://doi.org/10.3189/002214311798043825>

Nomura, D., McMinn, A., Hattori, H., Aoki, S., & Fukuchi, M. (2011b). Incorporation of nitrogen compounds into sea ice from atmospheric deposition. *Marine Chemistry*, 127, 90–99. <https://doi.org/10.1016/j.marchem.2011.08.002>

Nomura, D., Koga, S., Kasamatsu, N., Shinagawa, H., Simizu, D., Wada, M., et al. (2012). Direct measurements of DMS flux from Antarctic fast sea ice to the atmosphere by a chamber technique. *Journal of Geophysical Research-Oceans*, 117, C04011, doi: 10.1029/2010JC006755

Nomura, D., Aoki, S., Simizu, D., & Iida, T. (2018). Influence of sea ice crack formation on the spatial distribution of nutrients and microalgae in flooded Antarctic multiyear ice. *Journal of Geophysical Research: Oceans*, 123, 939–951. <https://doi.org/10.1002/2017JC012941>

Parsons, T. R., Takahashi, M., & Hargrave, B. (1984). Biological oceanographic processes (3rd ed., 330 pp.). Oxford, UK: Pergamon.

Pringle, D. J., Miner, J. E., Eicken, H., & Golden, K. M. (2009). Pore space percolation in sea ice single crystals. *Journal of Geophysical Research: Oceans*, 114, C12017. doi:10.1029/2008JC005145

Rahm, L., Håkansson, B., Larsson, P., Fogelqvist, E., Bremle, G., & Valderrama, J. (1995). Nutrient and persistent pollutant deposition on the Bothnian Bay ice and snow

fields. *Water Air and Soil Pollution*, 84, 187–201.

Roukaerts, A., Deman, F., Linden, F. V., Carnat, G., Bratkic, A., Moreau, S., et al. (2021). The biogeochemical role of a microbial biofilm in sea ice: Antarctic landfast sea ice as a case study. *Elementa: Science of the Anthropocene*, 9: 1. DOI: <https://doi.org/10.1525/elementa.2020.00134>

Rysgaard, S., & Glud, R. N. (2004). Anaerobic N₂ production in Arctic sea ice. *Limnology and Oceanography*, 49:86–94. <https://doi.org/10.4319/lo.2004.49.1.0086>

Rysgaard, S., Glud, R. N., Sejr, M. K., Blicher, M. E., & Stahl, H. J. (2008). Denitrification activity and oxygen dynamics in Arctic sea ice. *Polar Biology*, 31, 527–537. doi: 10.1007/s00300-007-0384-x

Saenz, B., T., & Arrigo, K. R. (2014). Annual primary production in Antarctic sea ice during 2005–2006 from a sea ice state estimate. *Journal of Geophysical Research: Oceans*, 119, 3645– 3678, <https://doi.org/10.1002/2013JC009677>

- Schnack-Schiel, B., Thomas, D. N., Haas, C., Dieckmann, G. S., & Alheit, R. (2001). The occurrence of the copepods *Stephos longipes* (Calanoida) and *Drescheriella glacialis* (Harpacticoida) in summer sea ice in the Weddell Sea, Antarctica. *Antarctic Science*, 13, 2, 150–157.
- Staley, J. T., & Gosink, J. J. (1999). Poles apart: biodiversity and biogeography of sea ice bacteria. *Annual Review of Microbiology*, 53, 189–215. doi: 10.1146/annurev.micro.53.1.189
- Sturm, M., & Massom, R. A. (2017). Snow and sea ice. D.N. Thomas, G.S. Dieckmann (Ed.), *Sea Ice (Second edition)*, Wiley-Blackwell, Oxford, 153–204. <https://doi.org/10.1002/9781444317145.ch5>
- Suzuki, R., & Ishimaru, T. (1990). An improved method for the determination of phytoplankton chlorophyll using N, N-dimethylformamide. *Journal of the Oceanographic Society of Japan*, 46, 190–194. <https://doi.org/10.1007/BF02125580>
- Takeda, S. (1998). Influence of iron availability on nutrient consumption ratio of diatoms in oceanic waters. *Nature*, 393, 774–777.
- Thomas, D. N., Lara, R. J., Eicken, H., Kattner, G., & Skoog, A. (1995). Dissolved organic matter in Arctic multi-year sea ice during winter: Major components and relationship to ice characteristics. *Polar Biology*, 15(7), 477–483.
- Toyota, T., Massom, R., Lecimte, O., Nomura, D., Heil, P., Tamura, T., Fraser, A. D. (2016). On the extraordinary snow on the sea ice off East Antarctica in late winter, 2012. *Deep-Sea Research II*, 131, (2016), 53–67. <https://doi.org/10.1016/j.dsr2.2016.02.003>
- Tréguer, P. J. & De La Rocha, C. L. (2013). The World Ocean silica cycle. *Annual Review of Marine Science*, 5: 477–501. DOI: <https://doi.org/10.1146/annurev-marine-121211-172346>
- Yoshikawa, C., Nakatsuka, T., & Wakatsuchi, M. (2006). Distribution of N* in the Sea of Okhotsk and its use as a biogeochemical tracer of the Okhotsk Sea Intermediate Water formation process. *Journal of Marine Systems*. 63 (1–2), 49–62. <https://doi.org/10.1016/j.jmarsys.2006.05.008>
- Ushio, S. (2006). Factors affecting fast-ice break-up frequency in Lützow-Holm Bay, Antarctica. *Annals of Glaciology*, 44, 177–182. DOI: <https://doi.org/10.3189/172756406781811835>
- Vancoppenolle, M., Goosse, H., de Montety, A., Fichefet, T., Tremblay, B., & Tison, J.-L. (2010). Modeling brine and nutrient dynamics in Antarctic sea ice: The case of dissolved silica, *Journal of Geophysical Research: Oceans*, 115, C02005, doi:10.1029/2009JC005369
- WMO (2014). Sea Ice Nomenclature. World Meteorological Organisation, Geneva.
- Wongpan, P., Meiners, K. M., Langhorne, P. J., Heil, P., Smith, I. J., Leonard, G. H., et al. (2018). Estimation of Antarctic land-fast sea ice algal biomass and

snow thickness from under-ice radiance spectra in two contrasting areas. *Journal of Geophysical Research: Oceans*, 123(3), 1907–1923. doi: 10.1002/2017jc013711

Zhou, J., Delille, B., Eicken, H., Vancoppenolle, M., Brabant, F., Carnat, G., et al. (2013). Physical and biogeochemical properties in Landfast sea ice (Barrow, Alaska): Insights on brine and gas dynamics across seasons. *Journal of Geophysical Research: Oceans*, 118, 6, 3172–3189. <https://doi.org/10.1002/jgrc.20232>

Figure captions

Figure 1. (a) The study area and (b–f) sampling locations in Lützow-Holm Bay, Eastern Antarctica. MODIS-Terra satellite images were obtained on (b) 19 January 2019, (d) 24 April 2015 at site JARE57_EM, (e) 25 April 2017 at site JARE59_POps, and (f) 18 March 2018 at sites JARE60_KU1 and JARE60_KU2. The images in (d–f) show the sea-ice condition before winter at each coring site; dark areas indicate where the sea ice was broken, and the cores obtained from these sites were first-year ice. The background image in (c) is an Advanced Land Observing Satellite (ALOS)/Panchromatic Remote-sensing Instrument for Stereo Mapping (PRISM) image obtained on 26 January 2010 by the Japan Aerospace Exploration Agency (JAXA).

Figure 2. Vertical profiles of (a, c) salinity and (b, d) ^{18}O values in snow, first-year ice cores (left), and multi-year ice cores (right). Horizontal solid lines at 0 cm depth indicate the snow–ice interface. Dashed lines in (b, d) show the ^{18}O value of under-ice water (-0.7‰) sampled during JARE60. The photo in (a) is a vertical thin section of the top 5 cm of core JARE60_KU2.

Figure 3. (a) Vertical profiles of salinity and ^{18}O values in core JARE57_P31.5 compared to (b) its ice structure. (c–f) Representative vertical thin section photographs are provided for each ice structure.

Figure 4. Vertical profiles of (a, e) NO_3^- , (b, f) PO_4^{3-} , (c, g) $\text{Si}(\text{OH})_4$, and (d, h) chl.a concentrations in first-year ice (top) and multi-year ice cores (bottom). Horizontal solid lines at a depth of 0 cm indicate the snow–ice interface. The photo in (c) and (d) is the ice algae community assemblage for the bottom of core JARE60_KU1.

Figure 5. Plots of (a) PO_4^{3-} , (b) $\text{Si}(\text{OH})_4$, (c) NO_3^- , and (d) NO_2^- concentrations versus snow fraction in multi-year ice cores.

Figure 6. Time series of the mean snow fraction in the top 200 cm of each multi-year ice core from 2015 (core JARE56_NMS) to 2018 (core JARE59_2).

Figure 7. Plots of (a, b) $\text{Si}(\text{OH})_4$, (c, d) PO_4^{3-} , (e, f) NO_3^- , and (g, h) NO_2^- concentrations versus salinity in first-year (left) and multi-year ice cores (right). Dilution lines are based on the nutrient concentrations and salinity of under-ice water.

Figure 8. Dissolved inorganic nitrogen ($\text{DIN} = \text{NO}_3^- + \text{NO}_2^-$) versus PO_4^{3-} concentrations in multi-year ice cores. The red solid line indicates the Redfield

ratio ($\text{DIN:P} = 16:1$) and the black dashed line indicates the ratio for uptake by diatoms ($\text{DIN:P} = 12:1$) (Takeda, 1998).

Figure 9. Vertical profiles of N^* (Equation 3) and chl.a concentrations in multi-year ice cores (a) JARE56_NMS and (b) JARE57_P31.5.

Figure 10. Conceptual illustration of the evolution of nutrient concentrations produced by photosynthesis and remineralization during the desalination process from new ice to first-year ice to multi-year ice.

Supporting Information for

Effects of snow and remineralization processes on nutrient distributions in multi-year Antarctic landfast sea ice

Reishi Sahashi¹, Daiki Nomura^{2,1,3*}, Takenobu Toyota⁴, Manami Tozawa¹, Masato Ito⁵, Pat Wongpan^{6,4,7}, Kazuya Ono⁴, Daisuke Simizu⁵, Kazuhiro Naoki⁸, Yuichi Nosaka⁹, Takeshi Tamura^{5,10}, Shigeru Aoki⁴, Shuki Ushio^{5,10}

¹*Faculty of Fisheries Sciences, Hokkaido University, Hakodate, Japan*

²*Field Science Center for Northern Biosphere, Hokkaido University, Hakodate, Japan*

³*Arctic Research Center, Hokkaido University, Sapporo, Japan*

⁴*Institute of Low Temperature Science, Hokkaido University, Sapporo, Japan*

⁵*National Institute of Polar Research, Tachikawa, Japan*

⁶*Institute for Marine and Antarctic Studies, University of Tasmania, Hobart, Australia*

⁷*JSPS International Research Fellow, Japan Society for the Promotion of Science, Tokyo, Japan*

⁸*Research and Information Center, Tokai University, Tokyo, Japan*

⁹*School of Biological Sciences, Tokai University, Sapporo, Hokkaido, Japan*

¹⁰*Graduate University for Advanced Studies (SOKENDAI), Tachikawa, Japan*

Contents of this file

Table S1

Introduction

The supporting information file describes biogeochemical data in the landfast sea ice, snow and under-ice water in Lützow-Holm Bay, East Antarctica collected during the 56–60th JAREs (2015–2019).

Sea ice observations of multi-year fast ice were performed during the summers of 2015–2019 as part of the 56–60th JAREs. Sea-ice cores were collected near Syowa Station. Ice cores were sampled using an ice corer (Geo Tecs Co., Ltd., Chiba, Japan) with an internal diameter of 0.09 m. Cores were immediately placed in tubular polyethylene bags and kept horizontal in a cooler box along with refrigerants to maintain below-freezing temperatures and minimize brine drainage from the core. Fresh snow samples were collected with an acid-washed polycarbonate shovel and placed into polyethylene zipper storage bags.

Sea ice cores were stored in a freezer at -30°C during their transport to the Institute of Low Temperature Science (ILTS), Hokkaido University, Sapporo, Japan. In the ILTS cold room (-15°C), cores were split lengthwise into two halves with an electric band saw: the first half for ice textural analysis and the second half for measurements of bulk ice salinity, $\delta^{18}\text{O}$ composition, nutrient and chl.a concentrations, and cell counting for ice algae community assemblage.

Under-ice water was collected through the coring holes with a Teflon water sampler (GL Science Inc., Japan) from 1 m below the bottom of the sea ice during JARE60 (2019). Water samples were collected approximately 30 minutes after the ice core was drilled at JARE60_KU_2 to avoid potential artifacts associated with the disturbance of drilling. All collected water samples were subsampled into (1) a 120-mL glass vial (Maruemu Co., Ltd., Osaka, Japan) for salinity measurement, (2) a 15-mL glass screw-cap vial (Nichiden-Rika Glass Co. Ltd, Kobe, Japan) for $\delta^{18}\text{O}$ analysis, (3) a 10-mL polyethylene screw-cap vial (Eiken Chemical Co. Ltd, Tokyo, Japan) for measurement of major inorganic nutrient concentrations ($\text{NO}_3^- + \text{NO}_2^-$, NH_4^+ , PO_4^{3-} , and $\text{Si}(\text{OH})_4$), and (4) a 500-mL Nalgene polycarbonate bottle (Thermo Fisher Scientific Inc., Waltham, MA, USA) for measurement of chlorophyll a (chl.a) concentrations. All samples were returned to the laboratory of the Japan Maritime Self-Defense Force icebreaker *Shirase* immediately after sampling.

Table S1. Dataset. “–” indicates no data.

Sample type	Station	Upper Limit of Depth range (cm)	Lower Limit of Depth range (cm)	Salinity	$\delta^{18}\text{O}$ (‰)	NO_3^- ($\mu\text{mol L}^{-1}$)	NO_2^- ($\mu\text{mol L}^{-1}$)	PO_4^{3-} ($\mu\text{mol L}^{-1}$)	$\text{Si}(\text{OH})_4$ ($\mu\text{mol L}^{-1}$)	Chl.a ($\mu\text{g L}^{-1}$)
Snow	JARE58_KU3	168	112	0.0	–21.8	0.0	0.0	0.0	0.0	0.0
Snow	JARE58_KU3	112	56	0.0	–22.6	0.1	0.0	0.0	0.0	0.0
Snow	JARE58_KU3	56	0	0.0	–22.2	0.1	0.0	0.0	0.0	0.0
Snow	JARE59_2	78	52	0.0	–22.3	0.1	0.0	0.0	0.1	0.0
Snow	JARE59_2	52	26	0.0	–19.8	0.1	0.0	0.0	0.1	0.0
Snow	JARE59_2	26	0	0.0	–21.0	0.0	0.0	0.0	0.1	0.0
Snow	JARE59_POPs	8	0	0.0	–16.1	0.4	0.0	0.0	0.0	0.0
Snow	JARE59_POPs	20	8	0.0	–17.4	0.3	0.0	0.0	0.0	0.3
Snow	JARE60_KU1	20	0.0	0.0	–21.2	0.3	0.0	0.0	0.0	0.0
Snow	JARE60_KU2	8	0	0.0	–18.9	0.3	0.0	0.0	0.0	0.0
Sea ice	JARE56_NMS	0	–13	0.0	–18.9	0.1	0.0	0.0	0.1	0.0
Sea ice	JARE56_NMS	–13	–25	0.0	–18.9	0.4	0.0	0.0	0.1	0.0
Sea ice	JARE56_NMS	–25	–38	0.0	–18.9	0.1	0.0	0.0	0.1	0.0
Sea ice	JARE56_NMS	–38	–50	0.0	–18.7	0.2	0.0	0.0	0.1	0.0
Sea ice	JARE56_NMS	–50	–63	0.0	–18.9	0.1	0.0	0.0	0.2	0.0
Sea ice	JARE56_NMS	–63	–76	0.0	–18.8	0.1	0.0	0.1	0.2	0.0
Sea ice	JARE56_NMS	–76	–88	0.0	–22.8	0.0	0.0	0.0	0.2	0.0
Sea ice	JARE56_NMS	–88	–101	0.2	–20.2	0.1	0.0	0.0	0.4	0.0
Sea ice	JARE56_NMS	–101	–113	0.0	–16.5	0.1	0.0	0.0	0.1	0.0
Sea ice	JARE56_NMS	–113	–126	0.0	–16.2	0.1	0.0	0.0	0.2	0.0
Sea ice	JARE56_NMS	–126	–139	0.2	–15.6	0.1	0.0	0.0	0.3	0.0
Sea ice	JARE56_NMS	–139	–151	0.5	–15.5	0.2	0.0	0.1	0.6	0.1
Sea ice	JARE56_NMS	–151	–164	0.6	–15.6	0.1	0.0	0.1	0.8	0.2
Sea ice	JARE56_NMS	–164	–176	0.2	–16.6	0.0	0.0	0.2	0.5	0.3
Sea ice	JARE56_NMS	–176	–189	0.4	–21.2	0.0	0.0	0.1	0.6	0.5

Sea ice	JARE56_NMS	-189	-201	1.8	-14.7	0.0	0.0	0.5	2.1	1.2
Sea ice	JARE56_NMS	-201	-214	1.5	-12.6	0.0	0.1	0.4	1.5	1.2
Sea ice	JARE56_NMS	-214	-227	0.6	-9.9	0.0	0.0	0.2	0.7	0.6
Sea ice	JARE56_NMS	-227	-239	0.5	-7.8	0.0	0.0	0.2	0.7	0.4
Sea ice	JARE56_NMS	-239	-252	1.0	-6.7	0.0	0.0	0.4	1.4	1.5
Sea ice	JARE56_NMS	-252	-264	0.9	-6.1	0.0	0.0	0.3	1.3	1.9
Sea ice	JARE56_NMS	-264	-277	1.0	-5.2	0.1	0.0	0.4	0.9	5.0
Sea ice	JARE56_NMS	-277	-290	1.9	-3.1	0.3	0.1	0.9	1.4	15.3
Sea ice	JARE56_NMS	-290	-302	2.3	-2.3	1.1	0.1	0.9	1.7	18.0
Sea ice	JARE56_NMS	-302	-315	1.4	-0.6	1.1	0.0	0.2	0.9	4.1
Sea ice	JARE56_NMS	-315	-327	1.2	0.8	1.0	0.0	0.2	1.0	4.8
Sea ice	JARE56_NMS	-327	-340	2.7	1.5	2.4	0.0	0.2	2.8	2.2
Sea ice	JARE57_P.31.5	0	-12	0.0	-19.8	0.1	0.0	0.0	0.0	0.0
Sea ice	JARE57_P.31.5	-12	-25	0.0	-19.9	0.0	0.0	0.0	0.0	0.0
Sea ice	JARE57_P.31.5	-25	-37	0.0	-20.6	—	0.0	0.0	0.1	0.0
Sea ice	JARE57_P.31.5	-37	-49	0.0	-20.7	0.2	0.0	0.0	0.0	0.0
Sea ice	JARE57_P.31.5	-49	-62	0.0	-20.7	0.2	0.0	0.0	0.0	0.0
Sea ice	JARE57_P.31.5	-62	-74	0.0	-22.4	0.0	0.0	0.0	0.1	0.0
Sea ice	JARE57_P.31.5	-74	-86	0.0	-22.2	0.0	0.0	0.0	0.0	0.0
Sea ice	JARE57_P.31.5	-86	-98	0.0	-21.1	0.0	0.0	0.0	0.1	0.0
Sea ice	JARE57_P.31.5	-98	-111	0.0	-22.6	0.0	0.0	0.0	0.1	0.0
Sea ice	JARE57_P.31.5	-111	-123	0.0	-23.8	0.0	0.0	0.0	0.2	0.0
Sea ice	JARE57_P.31.5	-123	-135	1.9	-20.1	0.0	0.0	0.1	1.7	0.1
Sea ice	JARE57_P.31.5	-135	-148	0.4	-17.5	0.0	0.0	0.1	0.5	0.1
Sea ice	JARE57_P.31.5	-148	-160	0.3	-13.5	0.0	0.0	0.2	0.8	0.3
Sea ice	JARE57_P.31.5	-160	-172	1.2	-14.9	0.0	0.0	0.4	1.6	0.6
Sea ice	JARE57_P.31.5	-172	-185	1.9	-15.5	0.0	0.0	0.4	3.0	0.9
Sea ice	JARE57_P.31.5	-185	-197	1.2	-19.8	0.0	0.0	0.3	1.4	0.6
Sea ice	JARE57_P.31.5	-197	-209	0.8	-18.2	0.0	0.0	0.1	1.1	0.5
Sea ice	JARE57_P.31.5	-209	-221	1.8	-16.8	0.0	0.0	0.3	2.9	1.7
Sea ice	JARE57_P.31.5	-221	-234	2.6	-15.3	0.0	0.0	0.4	4.8	1.9
Sea ice	JARE57_P.31.5	-234	-246	2.3	-15.8	0.0	0.0	0.3	4.5	1.5
Sea ice	JARE57_P.31.5	-246	-258	2.2	-15.7	0.0	0.0	0.4	4.9	1.8
Sea ice	JARE57_P.31.5	-258	-271	1.9	-15.3	0.0	0.0	0.5	4.1	3.7

Sea ice	JARE57_P.31.5	-271	-283	2.3	-15.0	0.0	0.1	0.7	6.6	7.0
Sea ice	JARE57_P.31.5	-283	-295	1.5	-14.3	0.0	0.1	0.8	11.5	8.2
Sea ice	JARE57_P.31.5	-295	-308	0.6	-13.8	0.0	0.1	1.1	8.3	13.9
Sea ice	JARE57_P.31.5	-308	-320	1.0	-6.3	0.0	0.0	0.5	9.3	4.8
Sea ice	JARE57_P.31.5	-320	-332	1.4	-3.9	0.4	0.0	0.7	12.3	6.0
Sea ice	JARE57_P.31.5	-332	-344	1.8	-2.0	1.6	0.0	0.6	7.6	5.8
Sea ice	JARE57_P.31.5	-344	-357	2.1	-1.0	2.4	0.0	0.8	9.2	6.0
Sea ice	JARE57_P.31.5	-357	-369	2.3	0.4	1.3	0.1	1.6	16.0	14.8
Sea ice	JARE57_P.31.5	-369	-381	1.6	1.6	0.4	0.1	1.2	8.5	11.5
Sea ice	JARE57_P.31.5	-381	-394	1.2	1.8	0.6	0.0	0.5	1.5	5.2
Sea ice	JARE57_P.31.5	-394	-406	1.4	1.0	1.9	0.0	0.3	1.9	3.3
Sea ice	JARE57_P.31.5	-406	-418	1.3	-0.4	2.1	0.0	0.4	1.6	3.6
Sea ice	JARE57_P.31.5	-418	-431	1.4	1.1	2.2	0.0	0.3	1.8	1.9
Sea ice	JARE57_P.31.5	-431	-443	2.1	1.7	3.5	0.0	0.3	2.9	0.6
Sea ice	JARE57_P.31.5	-443	-455	1.5	2.0	2.4	0.0	0.2	1.9	0.6
Sea ice	JARE57_P.31.5	-455	-467	1.6	2.0	2.5	0.0	0.2	1.9	0.6
Sea ice	JARE57_P.31.5	-467	-480	2.6	1.7	3.0	0.0	0.3	3.6	0.3
Sea ice	JARE57_P.31.5	-480	-492	1.2	-7.1	1.3	0.0	0.7	1.3	0.6
Sea ice	JARE57_EM	0	-15	0.6	-16.6	0.6	0.0	0.0	1.1	0.1
Sea ice	JARE57_EM	-15	-29	2.6	-1.9	0.4	0.0	0.1	1.7	0.2
Sea ice	JARE57_EM	-29	-44	2.3	-0.4	0.5	0.1	0.1	1.7	0.2
Sea ice	JARE57_EM	-44	-59	2.2	0.6	0.2	0.0	0.1	1.4	0.2
Sea ice	JARE57_EM	-59	-73	1.7	1.2	0.1	0.0	0.1	1.4	0.1
Sea ice	JARE57_EM	-73	-88	1.8	1.4	0.2	0.0	0.1	1.2	0.7
Sea ice	JARE57_EM	-88	-103	1.9	1.4	0.5	0.0	0.2	1.6	3.3
Sea ice	JARE57_EM	-103	-117	2.4	1.4	0.5	0.0	1.0	2.2	4.4
Sea ice	JARE57_EM	-117	-132	2.8	1.2	1.3	0.1	2.0	5.0	99.2
Sea ice	JARE58_KU3	0	-15	0.0	-20.6	—	0.0	0.0	0.0	0.1
Sea ice	JARE58_KU3	-15	-29	0.0	-20.6	—	0.0	0.0	0.0	0.1
Sea ice	JARE58_KU3	-29	-44	0.0	-21.7	—	0.0	0.0	0.0	0.0
Sea ice	JARE58_KU3	-44	-59	0.0	-20.6	0.1	0.0	0.0	0.0	0.0
Sea ice	JARE58_KU3	-59	-73	0.0	-19.8	0.1	0.0	0.0	0.0	0.0
Sea ice	JARE58_KU3	-73	-88	0.0	-20.0	0.1	0.0	0.0	0.0	0.1
Sea ice	JARE58_KU3	-88	-102	0.0	-21.6	0.2	0.0	0.0	0.0	0.1

Sea ice	JARE58_KU3	-102	-117	0.1	-20.6	0.2	0.0	0.0	0.0	0.1
Sea ice	JARE58_KU3	-117	-132	0.1	-20.2	0.2	0.0	0.0	0.0	0.0
Sea ice	JARE58_KU3	-132	-146	0.1	-20.2	0.1	0.0	0.0	0.0	0.1
Sea ice	JARE58_KU3	-146	-161	1.0	-17.9	0.2	0.0	0.1	1.0	1.3
Sea ice	JARE58_KU3	-161	-176	1.4	-16.7	0.2	0.0	0.0	1.3	0.8
Sea ice	JARE58_KU3	-176	-190	2.5	-18.2	0.2	0.0	0.1	2.6	0.5
Sea ice	JARE58_KU3	-190	-205	3.1	-20.9	0.4	0.0	0.1	2.8	1.4
Sea ice	JARE58_KU3	-205	-219	3.8	-21.4	0.7	0.0	0.1	3.9	1.1
Sea ice	JARE58_KU3	-219	-234	3.6	-21.4	0.2	0.0	0.2	2.8	1.4
Sea ice	JARE58_KU3	-234	-249	3.2	-22.6	0.3	0.0	0.1	2.1	1.3
Sea ice	JARE58_KU3	-249	-263	0.9	-23.3	0.3	0.0	0.0	0.3	0.8
Sea ice	JARE58_KU3	-263	-278	0.5	-22.6	0.1	0.0	0.0	0.3	1.0
Sea ice	JARE59_2	0	-18	0.0	-21.6	—	0.0	0.0	0.1	0.0
Sea ice	JARE59_2	-18	-35	0.0	-21.2	—	0.0	0.0	0.1	0.0
Sea ice	JARE59_2	-35	-53	0.0	-21.0	—	—	0.0	0.0	0.0
Sea ice	JARE59_2	-53	-71	0.0	-20.0	—	0.0	0.0	0.0	0.0
Sea ice	JARE59_2	-71	-88	0.0	-20.8	0.2	0.0	0.0	0.0	0.0
Sea ice	JARE59_2	-88	-106	0.0	-20.7	0.2	0.0	0.0	0.1	0.0
Sea ice	JARE59_2	-106	-123	0.0	-21.1	0.2	0.0	0.0	0.2	0.0
Sea ice	JARE59_2	-123	-141	0.0	-21.3	0.2	0.0	0.0	0.1	0.0
Sea ice	JARE59_2	-141	-159	0.0	-19.9	0.3	0.0	0.0	0.0	0.0
Sea ice	JARE59_2	-159	-176	0.0	-19.2	0.3	0.0	0.0	0.1	0.0
Sea ice	JARE59_2	-176	-194	0.3	-20.1	0.3	0.0	0.1	0.2	0.0
Sea ice	JARE59_2	-194	-212	0.6	-18.6	0.3	0.0	0.3	0.7	0.2
Sea ice	JARE59_2	-212	-229	0.4	-14.4	0.6	0.0	0.3	0.6	0.2
Sea ice	JARE59_2	-229	-247	0.5	-17.5	0.7	0.0	0.3	0.7	0.2
Sea ice	JARE59_POPs	0	-16	0.4	-16.4	0.2	0.0	0.1	0.1	0.0
Sea ice	JARE59_POPs	-16	-32	1.8	-4.5	0.3	0.0	0.3	1.2	0.3
Sea ice	JARE59_POPs	-32	-48	1.9	-1.5	0.3	0.0	0.3	0.8	1.2
Sea ice	JARE59_POPs	-48	-64	1.7	-1.1	0.5	0.0	0.3	0.5	1.1
Sea ice	JARE59_POPs	-64	-80	1.9	-0.1	0.2	0.0	0.4	0.5	1.0
Sea ice	JARE59_POPs	-80	-96	2.3	0.3	0.5	0.0	0.2	0.5	1.3
Sea ice	JARE59_POPs	-96	-112	2.9	0.2	0.4	0.0	0.5	1.1	1.8
Sea ice	JARE59_POPs	-112	-128	3.3	0.7	0.5	0.0	0.6	1.5	4.0

Sea ice	JARE59_POPs	−128	−144	3.2	1.3	0.5	0.0	1.1	1.7	11.2
Sea ice	JARE60_KU1	0	−16	4.9	−1.5	0.1	0.0	0.6	0.9	7.3
Sea ice	JARE60_KU1	−16	−33	5.8	−0.9	0.2	0.0	0.7	1.1	13.9
Sea ice	JARE60_KU1	−33	−49	5.6	0.0	1.6	0.0	0.5	0.4	13.6
Sea ice	JARE60_KU1	−49	−65	4.8	0.6	1.8	0.0	0.8	0.6	17.7
Sea ice	JARE60_KU1	−65	−82	5.1	0.6	0.7	0.0	0.5	0.4	10.7
Sea ice	JARE60_KU1	−82	−98	3.8	1.4	0.8	0.0	0.6	0.6	10.4
Sea ice	JARE60_KU1	−98	−114	4.2	1.6	2.6	0.0	1.1	1.1	56.4
Sea ice	JARE60_KU2	0	−16	3.3	−11.4	0.3	0.1	0.2	0.3	2.9
Sea ice	JARE60_KU2	−16	−31	3.0	−7.5	0.5	0.1	0.3	0.5	1.6
Sea ice	JARE60_KU2	−31	−47	3.8	−6.2	0.3	0.0	0.2	0.5	1.6
Sea ice	JARE60_KU2	−47	−62	4.6	−2.7	0.1	0.0	0.4	1.1	3.5
Sea ice	JARE60_KU2	−62	−78	4.2	−1.3	0.2	0.0	0.6	1.4	6.0
Sea ice	JARE60_KU2	−78	−94	4.1	−0.1	0.3	0.0	0.5	1.2	10.1
Sea ice	JARE60_KU2	−94	−109	3.2	1.1	0.2	0.0	0.2	0.6	7.0
Sea ice	JARE60_KU2	−109	−125	2.9	1.4	0.2	0.0	0.1	0.4	3.0
Sea ice	JARE60_KU2	−125	−140	3.6	1.4	0.5	0.0	0.6	1.3	13.7
Under-ice water (1 m from ice bottom)	JARE60_KU2	—	—	33.9	−0.7	28.0	0.2	1.8	60.6	—