

# Long-term (2000-2020) variability of in situ time series of Carbonyl Sulfide

Carmine Serio<sup>1</sup>, Stephen A. Montzka<sup>2</sup>, Guido Masiello<sup>1</sup>, and Vincenzo Carbone<sup>3</sup>

<sup>1</sup>Universita della Basilicata

<sup>2</sup>NOAA-ESRL-GMD

<sup>3</sup>Universita della Calabria

November 24, 2022

## Abstract

The monthly time series of carbonyl sulfide (OCS) atmospheric mole fractions measured at NOAA network stations (2000 to 2020) have been analyzed, and the long-term behaviour has been assessed based on the Empirical Mode Decomposition (EMD). EMD is a fully non-parametric analysis of frequency modes and trends in a given series and is based on the data alone. We have found that the OCS atmospheric mole fraction, after an increasing phase up to ~2015, with a temporary decline around 2009, is now decreasing at all stations, reflecting a recent imbalance in its total sources and losses. Our analysis has revealed a characteristic time scale for variation of 8-10 years. The variance associated with this long-term behaviour ranges from 15 to 40% of the total strength of the signal, depending on location. To our knowledge, this low-frequency mode is a novel result not assessed in previous studies. Apart from this complex long-term behaviour, the OCS time series show a strong annual cycle, which primarily results from summertime OCS uptake by vegetation. In addition, we have also found one more frequency of minor variance intensity in the measured mole fraction time-history, which corresponds to periods in the range of 2 to 3 years. This inter-annual variability of OCS may be linked to the Quasi-Biennial Oscillation or QBO.

1           **Long-term (2000–2020) variability of in situ time series of Carbonyl Sulfide**

2  
3                   **C. Serio<sup>1\*</sup>, S. Montzka<sup>2</sup>, G. Masiello<sup>1</sup>, V. Carbone<sup>3</sup>**

4  
5           <sup>1</sup>Scuola di Ingegneria, Università degli studi della Basilicata, Potenza, Italy.

6           <sup>2</sup>NOAA Earth System Research Laboratory, Boulder, Colorado, USA

7           <sup>3</sup>Dipartimento di Fisica, Università della Calabria, Rende, Italy.

8  
9           \*Corresponding author address: Carmine Serio (carmine.serio@unibas.it)

10           **Key Points:**

- 11           • Atmospheric Carbonyl Sulfide is decreasing at NOAA network stations
- 12           • Time Series Analysis and Characteristic Scales encompassing one year to 8-10 years
- 13           • Empirical Mode Decomposition shows a reach wealth of frequencies, some compatible
- 14           with Quasi Biennial Oscillation
- 15

## 16 **Abstract**

17 The monthly time series of carbonyl sulfide (OCS) atmospheric mole fractions measured at NOAA  
18 network stations (2000 to 2020) have been analyzed, and the long-term behaviour has been  
19 assessed based on the Empirical Mode Decomposition (EMD). EMD is a fully non-parametric  
20 analysis of frequency modes and trends in a given series and is based on the data alone. We have  
21 found that the OCS atmospheric mole fraction, after an increasing phase up to ~2015, with a  
22 temporary decline around 2009, is now decreasing at all stations, reflecting a recent imbalance in  
23 its total sources and losses. Our analysis has revealed a characteristic time scale for variation of 8-  
24 10 years. The variance associated with this long-term behaviour ranges from ~15 to 40% of the  
25 total strength of the signal, depending on location. To our knowledge, this low-frequency mode is  
26 a novel result not assessed in previous studies. Apart from this complex long-term behaviour, the  
27 OCS time series show a strong annual cycle, which primarily results from summertime OCS  
28 uptake by vegetation. In addition, we have also found one more frequency of minor variance  
29 intensity in the measured mole fraction time-history, which corresponds to periods in the range of  
30 2 to 3 years. This inter-annual variability of OCS may be linked to the Quasi-Biennial Oscillation  
31 or QBO.

32 .

## 33 **Plain Language Summary**

34 Carbonyl sulfide (OCS) is the most abundant sulfur-containing trace gas in the atmosphere and  
35 accounts for a significant part of sulfur in the stratospheric aerosol. OCS has recently emerged as  
36 a putative proxy for the terrestrial photosynthetic uptake of CO<sub>2</sub> because OCS and CO<sub>2</sub> have the  
37 same diffusion pathway into leaves. The OCS hydration reaction in this process is irreversible. For  
38 this reason, a better understanding of its time scales of variability can improve our knowledge of  
39 the carbon cycle. The study has analyzed OCS at 14 cooperative stations, which are distributed all  
40 around the world. We have found a characteristic time scale for 8-10 years variation. To our  
41 knowledge, this low-frequency mode is a novel result not assessed in previous studies. Apart from  
42 this complex long-term behaviour, the OCS time series show a robust yearly cycle, primarily from  
43 summertime OCS uptake by vegetation. Finally, we have also found one more frequency, which  
44 corresponds to periods in the range of 2 to 3 years. This inter-annual variability of OCS may be  
45 linked to the Quasi-Biennial Oscillation, which is an almost periodic oscillation of the winds of  
46 the equatorial stratosphere.

47

48

## 49 **1. Introduction**

50 The importance of carbonyl sulfide in the study of terrestrial vegetative ecosystems has clearly  
51 emerged in recent studies (Campbell et al., 2008, 2017; Maseyk et al., 2014; Montzka et al., 2007).  
52 OCS is the most abundant sulfur-containing trace gas in the atmosphere and accounts for a  
53 significant part of sulfur in the stratospheric aerosol (Brühl et al., 2012). Essential sources of OCS  
54 are natural, and among them, oceans, soils, and volcanic eruptions play a dominant role.  
55 Otherwise, anthropogenic sources have been recognized as secondary contributors: biomass  
56 burning and industrial activities (Campbell et al., 2008). The main sink of OCS has been identified  
57 as vegetation uptake, whose magnitude is also influenced by seasonal trends in terrestrial  
58 vegetative photosynthesis. Conversely, in the stratosphere, the photochemical loss is the prominent

59 removal process, but at a substantially slower rate than vegetative uptake (Aydin et al., 2020; Berry  
60 et al., 2013; Glatthor et al., 2015; Kettle, 2002; Whelan et al., 2018).

61 Moreover, OCS has recently emerged as a putative proxy for the terrestrial photosynthetic  
62 uptake of CO<sub>2</sub> because OCS and CO<sub>2</sub> have the same diffusion pathway into leaves (Campbell et  
63 al., 2008; Montzka et al., 2007), and OCS hydration reaction in this process is irreversible. In  
64 addition to these earlier studies, more recent works (Berry et al., 2013; Campbell et al., 2015) have  
65 shown that carbonyl sulfide holds great promise for studies of carbon cycle processes because it is  
66 an atmospheric tracer of photosynthetic Gross Primary Production (GPP). According to (Berry et  
67 al., 2013; Campbell et al., 2015; Montzka et al., 2007), the uptake of OCS from the atmosphere is  
68 dominated by carbonic anhydrase (CA), an enzyme abundant in leaves that also catalyzes CO<sub>2</sub>  
69 hydration during photosynthesis. However, as a continuation of previous studies, it has been shown  
70 in (Ogée et al., 2016) that soils can also effectively exchange OCS with the atmosphere, which can  
71 complicate the retrieval of GPP from atmospheric budgets for some regions and scales. Some  
72 agricultural fields can take up large amounts of OCS from the atmosphere as soil microorganisms  
73 contain CA. OCS emissions from soils have been reported in agricultural fields or anoxic soils  
74 (Ogée et al., 2016). On a global scale, uptake by vegetation and soils account for more than 90%  
75 of the removal of OCS from the atmosphere, the remaining 10% being assigned to OH oxidation  
76 and transport to the stratosphere (Aydin et al., 2020; Berry et al., 2013; Glatthor et al., 2015; Kettle,  
77 2002; Whelan et al., 2018).

78 Apart from seasonal variations, the OCS atmospheric mole fraction had remained relatively  
79 stable, e.g., within 7% (Montzka et al., 2007) for the period 2000-2005, when OCS routinely began  
80 measured at the 18 NOAA stations and aircraft profiling sites. Ice core and firm air measurements,  
81 e.g., (Aydin et al., 2020) and references therein, have been used to reconstruct atmospheric  
82 carbonyl sulfide's preindustrial and industrial history. The more recent atmospheric OCS  
83 abundance surveys use a panoply of complementary ground-based, airborne, and satellite  
84 observations, e.g., (Camy-Peyret et al., 2017; Krysztofiak et al., 2015; Lejeune et al., 2017;  
85 Montzka et al., 2007).

86 Almost all analyses of historical and contemporary data sets (Campbell et al., 2017) have  
87 been interpreted with models that simulate changes in OCS concentration according to changes in  
88 its global budget of natural and anthropogenic sources (from oceans and soils, from industry and  
89 biomass burning, respectively), and biogenic sinks (from plant photosynthesis and soils) as  
90 reviewed by (Whelan et al., 2018). Although anthropogenic emissions have exerted a dominant  
91 influence in driving secular atmospheric abundance changes since the 19<sup>th</sup> century (Aydin et al.,  
92 2020; Campbell et al., 2015; Montzka et al., 2007) found that long-term changes in the  
93 observation-based OCS record were most consistent with simulations of climate and the carbon  
94 cycle that assume large growth in plant photosynthesis during the twentieth century. However,  
95 these analyses did not encompass the most recent trends in atmospheric OCS, e.g., since 2014-  
96 2015.

97 This study analyzes OCS measurements from the NOAA's global flask network, whose  
98 observing stations are spread around the globe but are more numerous in the North Hemisphere  
99 (NH), where anthropogenic sources are localized. A qualitative inspection of these data shows that  
100 the atmospheric OCS has entered a decline phase at all stations in recent years. Overall, we will  
101 show that a long-term behaviour with a characteristic time scale of ~8-10 years characterizes OCS  
102 time series from all sites analyzed in this study. However, over-imposed to this trend, there are  
103 cyclic behaviors with annual and inter-annual scales of variability.

104 The paper is organized as follows. Section 2 describes data and methods. Then, section 3  
 105 is devoted to presenting and discussing results. Finally, conclusions are taken in section 4.  
 106

## 107 2. Data and Methods


### 108 2.1. Data

109 For many years, OCS measurements from flasks have been obtained at approximately weekly  
 110 intervals at 14 NOAA and cooperative stations (Montzka et al., 2007). The sampling process  
 111 involves simultaneously pressurizing air into a pair of stainless steel or glass flasks that are  
 112 subsequently shipped to the Boulder laboratory for analysis. Here we consider monthly mean mole  
 113 fractions, and the data span different periods according to the station. The longest OCS time series  
 114 at these sites extends from March 2000 to December 2020. Table 1 summarizes the basic  
 115 information about the 14 stations, whereas Fig. 1 shows the position of the measurement stations  
 116 around the globe.


117  
 118 **Table 1.** NOAA stations whose OCS measurements from flasks have been analyzed in this study. The table  
 119 also gives the percentage of missing data, as monthly means, for each time series. These gaps result from  
 120 a lack of availability of flasks at a site and larger-than-acceptable differences in simultaneously filled  
 121 flasks.

Station	Code	Lat [°N]	Lon [°W]	Elevation [masl]	Time Interval endpoints	% Missing data
Alert, Nunavut, Canada	ALT	82.4508	62.5072	185	May 2000-October 2020	12.60
Point Barrow, USA	BRW	71.3230	156.6114	11	March 2000-December 2020	2.40
Cape Grim, Tasmania	CGO	-40.683	144.6900	94	February 2000-December 2020	3.60
Harvard Forest, USA	HFM	42.5378	72.1714	340	March 2000-December 2020	2.40
Cape Kumukahi, USA	KUM	19.7371	155.0116	0.30	March 2000-December 2020	0.80
Park Falls, USA	LEF	45.9451	90.2732	472	May 2000-December 2020	2.01
Mace Head, Ireland	MHD	53.3260	9.899	5.00	May. 2001-December 2020	7.60
Mauna Loa, USA	MLO	19.5362	155.5763	3397	March 2000-December2020	0.40
Niwot Ridge, USA	NWR	40.0531	105.5864	3523	March 2000-December 2020	3.20
Palmer Station, Antarctica	PSA	-64.7742	64.0527	10	May 2000-December 2020	14.50
Tutuila, American Samoa	SMO	-14.2474	170.5644	42	March 2000-December 2020	2.80
South Pole, Antarctica	SPO	-89.98	24.8	2810	May 2000-December 2020	11.29
Summit, Greenland	SUM	72.5962	38.422	3209	June 2004-December2020	5.03
Trinidad Head, USA	THD	41.0541	124.151	107	April 2002-December 2020	0.44

122  
 123 The time series can have occasional missing data (see also the last column in Tab. 1); when needed,  
 124 gaps in the OCS sequences have been filled by cubic spline interpolation. An example is shown in  
 125 Fig. 2. Because the sampling of OCS “events” is not the same at all stations and can vary at the  
 126 same station, month-to-month, the event measurements are averaged to form monthly means. The  
 127 analysis is then performed on these monthly time series. Also, we clarify that the gaps shown in  
 128 Tab. 1 are assessed on the basis of the monthly time series. We also note that the uneven sample  
 129 frequency at the same station adds a sampling noise, which the EMD methodology is capable of  
 130 filtering out, as it will be shown further in the paper.



131  
 132 **Figure 1.** Location of the 14 NOAA stations considered in this work  
 133  
 134



135  
 136 **Figure 2.** Example of a monthly OCS time series showing the gap-filling with cubic spline  
 137 interpolation. The case shown in the figure refers to the HFM station.  
 138

139 **2.2. Methods: trends identification**

140 The long-term behaviours or trends in data are identified through the empirical mode  
 141 decomposition (EMD) technique, developed to process nonlinear and nonstationary data (Huang  
 142 et al., 1998) and successfully applied in many different fields, e.g., (Capparelli et al., 2013;  
 143 Coughlin & Tung, 2004; Echeverría et al., 2001; Laurenza et al., 2012; Lee & Ouarda, 2011, 2012;  
 144 Loh, 2004; Y. Wu & Shen, 2016). EMD decomposes a time series into a finite number of intrinsic  
 145 mode functions (IMFs) and a residual by using an adaptive basis derived from the time series  
 146 through a so-called “sifting” process, namely,  
 147

$$X(t) = \sum_{j=1}^m c_j(t) + r(t) \quad (1)$$

where  $t$  is the time,  $m$  the number of modes, and  $X(t)$  denotes a generic time series;  $c_j$  is  $j$ -th IMF, and finally,  $r$  is the residual, which can be either the *mean* trend or a constant. Because the series  $X$  is sampled at discrete time  $t = j\Delta t, j = 1, \dots, N$ , with  $N$  the total number of discrete measurements, we have that the whole time span of the series is  $N\Delta t$ . In our case,  $\Delta t = 1$  month. Furthermore, to simplify notation, hereafter, we will write  $j$  for  $j\Delta t$  and  $N$  for  $N\Delta t$ .

In conventional trend analysis, it is often assumed, e.g., that the trend is linear, and therefore, it can be extracted with formal regression analysis (e.g. Gardiner et al., 2008; Lejeune et al., 2017). Furthermore, in non-parametric methods, the trend is analyzed through digital filtering techniques, e.g., the Fourier transform and low-pass filters to smooth the selected data and separate the low-frequency components from the seasonal cycle (e.g. Thoning et al., 1989).

In the present analysis, the trend is defined by considering all the components of the signal which show frequency modes lower than a given threshold frequency  $f_{th}$ ; in this study, the default value is  $f_{th} = 3/N$ , that is the frequency corresponding to a period equal to  $N/3$ . Because in our analysis, the OCS time series is 17 to 20 years long,  $N/3$  yields approximately 5–7 years. The threshold has been selected by trial and error and has been checked to provide a consistent analysis for the various stations. Also, OCS has a tropospheric lifetime of  $\sim 2$ –7 years (Blake, 2004), therefore frequencies lower than  $f_{th}$  characterizes long-term behaviour with timescales longer than the finite lifetime of OCS.

With this in mind, the trend,  $\tau$  is defined according to,

$$\tau(t) = \sum_{j=1}^m c_j(t) + r(t) \quad (2)$$

with  $f_m < \dots < f_l \leq f_{th}$ . Again, this definition is consistent with the idea that the trend has to capture the low-frequency variability of the signal.

The characteristic frequency of a given mode,  $c_j(t)$  can be identified with the usual computation of the classical Fourier variance spectrum analysis or Power Spectral Density (PSD). Later in this study (see section 3.2), we will show examples of how the frequency components within each IMF can be analyzed through the Hilbert transform (Huang et al., 1998). However, in case we are interested in determining the dominant frequency of each mode, we can resort to the classical PSD.

This work uses the EMD algorithm included in Matlab distribution 2020b, which implements all prescriptions and stopping criteria, as suggested by (Wang et al., 2010), to avoid the decomposition to run endlessly toward the limit with many infinite iterations of sifting, e.g., (Z. Wu & Huang, 2010). However, the black-box usage of the tool is not recommended. Even with the stopping criteria, there is no way to prevent the code from decomposing part of the trend in the lower frequency modes. Therefore there are at least three aspects that need to be carefully

190 addressed when using the Matlab software package: a) how to fix the number of modes,  $m$ ; b) how  
 191 to prevent mode splitting and mode mixing; c) how to handle problems with the boundaries or end  
 192 effects because of the finiteness of the series.

193

194 For issue a), we limit the number of modes to  $m=4$ , which is based on physical insights.  
 195 We know that the observations are affected by noise; therefore, the first mode will fit the high  
 196 oscillatory component of the noise. The second IMF or mode is expected to fit the annual cycle.  
 197 The third is devoted to representing inter-annual variability, which is likely to be found in the  
 198 series. Finally, the fourth last mode is to model possible lower frequency oscillations and,  
 199 therefore, long-term trend structures. For this reason, by default, we have the threshold criterion  
 200  $f_{th} = 3/N$  in defining the trend: everything with frequency lower than  $f_{th} = 3/N$  is moved to the  
 201 trend. The threshold  $f_{th}$  can be changed in case we are interested in looking at EMD reconstruction  
 202 of the signal, which includes specific frequencies.

203

204 For issue b), we use the EEMD (Ensemble Empirical Mode Decomposition, e.g., (Z. Wu  
 205 & Huang, 2009)) strategy of adding noise to the observations. For a given sample of observations,  
 206  $X(j), j = 1, \dots, N$  we build up the noise sample  $\tilde{X}(j) = X(j) + w(j)$ , with  $w$  a Gaussian noise term  
 207 with zero mean and standard deviation,  $\sigma_w$ .  $\tilde{X}(j)$  is EMD decomposed, and the operation is  
 208 repeated  $nsamples$  time. Finally, the four IMF and the residual are taken by considering the  
 209 average over the corresponding  $nsamples$ . However, before performing EMD on  $\tilde{X}(j)$ , we first  
 210 extend the signal to account for possible boundary effects.

211

212 To this end, - issue c) -, we use the strategy proposed by (Stallone et al., 2020). The series  
 213  $\tilde{X}(j)$  is symmetrically extended outside the boundaries, producing, on both sides, an extended  
 214 signal  $\tilde{X}_{ext}(j)$  which is, on each side,  $N$  times longer than the original one. Then,  $\tilde{X}_{ext}(j)$  is  
 215 multiplied by a function  $\chi(j)$ , which is one for the original signal  $\tilde{X}(j)$  and tends smoothly to zero  
 216 as we approach the two left and right ends of the extended signal. In this way, the signal  $\tilde{X}_{ext}(j)$   
 217 is periodic at the boundaries.

218

219 For completeness, the last word has to be said for  $\sigma_w$ . We know that the observation noise  
 220 of the OCS measurement is below 2 ppt or less than 0.5% on average. Therefore,  $\sigma_w$  is taken equal  
 221 to 1.5 ppt to preserve the original structure of the series.

222

223 For the benefit of the reader, we summarize the algorithm we have devised to apply EMD to  
 224 the OCS time series.


225

- 226 1. Set  $f_{th} = 3/N$  (default value,  $3/N$ ) and  $\sigma_w$  (default value, =1.5 ppt)
- 227 2. Set the maximum number of modes (default,  $m = 4$ )
- 228 3. Set the number of random samples, (default,  $nsamples = 1000$ )
- 229 4. Generate the noisy series  $\tilde{X}(j), j = 1, \dots, N$
- 230 5. Generate the extended series  $\tilde{X}_{ext}(j), j = 1, \dots, 3N$
- 231 6. EMD the series  $\tilde{X}_{ext}(j)$
- 232 7. Store the IMFs and the residual over the original range of the signal,  $j = N + 1, \dots, 2N$
- 233 8. Repeat steps 4 to 7  $nsamples$  times
- 234 9. Compute the final IMFs and residual by considering the average over the  $nsamples$  of  
 235 the corresponding functions calculated at step 7.

- 236 10. Compute the pdf or power density function of the four IMF (we use the tool *pcov* in the  
 237 Matlab distribution 2020b).  
 238 11. Compute the frequency peak of each IMF and related uncertainty  
 239 12. Compute the trend according to Eq. (2).  
 240


241 It should be stressed that the above procedure has been finalized, and the sensitivity of the  
 242 procedure to the various parameters checked by trial and error, simulations, and applications to the  
 243 time series at hand.

244 To explain how the EMD decomposition is applied and used in this study, we show its  
 245 application to the MLO series (monthly averages from March 2000 to December 2020 ( $N =$   
 246 250 months)). The decomposition consists of 4 modes, and a residual and is shown in Fig. 3, and  
 247 it is possible to see that the higher mode numbers are associated with lower frequency variability.



248 **Figure 3.** Exemplifying the EMD analysis applied to MLO monthly mean mole fractions measured  
 249 for OCS (in ppt). Top to bottom, signal, IMFs and residual.  
 250  
 251

252 As expected, the first IMF extracts the high oscillatory component of the noise. The second  
 253 component is an almost perfect harmonic of the constant period, although the amplitude can  
 254 change with time. To better understand the relevant frequencies in the third and fourth modes, the  
 255 PDFs of the four IMFs in Fig. 3 are shown in Fig. 4.  
 256



257  
 258 **Figure 4.** MLO station. Power density functions of the four IMF corresponding to the EMD  
 259 decomposition of the MLO monthly time series; a) IMF1; b) IMF2; 3) IMF3; d) IMF4. The figure  
 260 also shows the peak frequency of IMF2-4.  
 261

262 From Fig. 4, we see that the first IMF has a flat spectrum as expected for white noise, and  
 263 its spectral density is two orders of magnitude lower than the sharp power of the annual cycle  
 264 (there is a ratio 100:1 in the y-axis scale of IMF2 vs IMF1). Compared with Fig. 3, it is possible  
 265 to see that the EMD methodology can filter out the random component in the data.  
 266

267 The second IMF extracted from the MLO record yields a frequency peak almost exactly at  
 268  $1/12 \cong 0.0833$  in units of 1/month. IMF2 has the most prominent spectral density, and in fact,  
 269 from Fig. 3, we see that the mode is close to a pure harmonic with a period equal to 12 months.  
 270 We also see that the amplitude is not exactly the same from year-to-year, suggesting the presence  
 271 of interannual variability. To perform an assessment of how close IMF2 is to a pure harmonic, we  
 272 have fitted it with the model  
 273

$$A \sin\left(\frac{2\pi(t-\tau)}{T}\right) \quad (3)$$


274  
 275 with the time  $t$  in units of months and  $T = 12$  months; the amplitude and delay  $A, \tau$  are fit  
 276 parameters. A Least Squares fitting procedure of the model of Eq. (3) to the IMF2 data shown in  
 277 Fig. 3 (to phase the harmonic with the calendar year, the fit considers the data from January 2001,  
 278 ( $t = 1$ ) up to December 2020 ( $t = 240$ )) yields,  $A = 17.27$  ppt, with a 95% confidence interval  
 279 of [16.51, 18.03] ppt and,  $\tau = 1.93$  months, with a 95% confidence interval of [1.85, 2.02] months.  
 280

281 The goodness of the fit has been assessed through the correlation coefficient, and we found  $R^2 =$   
282 0.90. The delay  $\tau \sim 2$  months says that the peak value is attained in May, whereas the trough is in  
283 November. Finally, on average, the annual cycle's peak-to-peak amplitude is equal to  $\sim 34$  ppt in  
284 the MLO measurement record.

285  
286 The third IMF is close to 2 years, although its uncertainty is as large as  $\sim 6$  months, and its  
287 spectral density is 1-2 order of magnitude lower than that of the annual cycle. However, although  
288 of less intensity, the IMF3 power maximizes at a value which is in good agreement with the QBO  
289 (Quasi Biennial Oscillation) mean cycle, which has a periodicity of 28-29 months, or  $\sim 0.4$  per year,  
290 e.g., see (Ray et al., 2020). Finally, the fourth mode is more peaked than the third. It has a larger  
291 density but corresponds to a period close to 10 years. Therefore this mode is moved to the trend or  
292 long-term behaviour, which is shown in Fig. 5. In passing, we note that the frequency uncertainty  
293 shown in Fig. 4 is computed as the Half-Width at Half-maximum of the corresponding spectral  
294 line.


295  
296 According to the definition of Eq. (2), the EMD trend,  $\tau$  is prescribed to show time scales  
297 larger than those corresponding to the threshold frequency,  $f_{th} = 3/N$ , which for the MLO station  
298 corresponds to  $\sim$  seven years. From Fig. 5 we see that on time scales larger than 7 years, the decline  
299 of the OCS in recent years is clearly seen. Again in Fig. 5, the EMD trend is compared with the  
300 other two smoothing, non-parametric and non-linear, algorithms. These are the *lowess*,  $\tau_l$  (an  
301 acronym of locally weighted scatter plot smoothing, e.g., (Cleveland and Devlin, 1988)) and the  
302 moving average,  $\tau_{ma}$ . They are both prescribed with a span of  $N/3$  to properly compare with the  
303 time scales designed for the EMD trend. The *lowess* smoothing is based on a local least-squares  
304 fitting and generalizes the smoothing average method, which is also shown in Fig. 5. It is seen that  
305 the moving average filter shows a high-frequency ringing close to the boundaries of the signal,  
306 where it tends to collapse on the data points. In contrast, the *lowess*,  $\tau_l$  is much more consistent at  
307 the boundaries, although it provides a smoother version than the EMD,  $\tau$ . Nevertheless, the  
308 comparison exemplifies how EMD yields a methodology to determine and control the  
309 characteristic scales we want to include in the reconstruction of the signal.

310




311  
 312 **Figure 5.** OCS monthly averages (2000 to 2020) for the MLO station and trend analysis according  
 313 to EMD, lowess and moving average filters (e.g., see Eq. (2) and the text in the paper).  
 314

315 To exemplify this aspect, Fig. 6 provides a reconstruction of the signal, which also includes  
 316 the IMF3. Therefore, this is equivalent to using a threshold  $f_{th} = 1/12$ , in order to remove the  
 317 annual cycle from the original data. Based on the pdf analysis in Fig. 4, the EMD reconstruction  
 318 in Fig. 6 includes all characteristics scales larger than  $\sim 2$  years. For comparison, Fig. 6 also shows  
 319 the representation of the data after their smoothing with a moving average filter with a span of 12,  
 320 24, and 36 months, respectively. From Fig. 6, we see that the moving average still retains a high  
 321 oscillatory component, likely due to the observation noise. Conversely, EMD reconstruction  
 322 appears smoother because the noise has been filtered through the IMF1, which is not included in  
 323 the reconstruction. EMD clearly identifies the very large peak-to-peak variation across 2004-2005  
 324 and the relative trough in 2009-2010. These features are attenuated in the moving average filters.  
 325 Finally, the distance among peaks of the EMD reconstruction suggests variability scales of 2 – 4  
 326 years, which, as discussed above, could be linked to QBO.  
 327



328  
 329 **Figure 6.** OCS monthly averages (2000 to 2020) for the MLO station and trend analysis according  
 330 to the EMD reconstruction with the removal of the annual cycle. For comparison, the figure also  
 331 shows the results with a moving average filter with three different time spans, 12, 24 and 36  
 332 months, respectively.

333  
 334 Before closing this section, we also note that a conventional Fourier analysis of the signal  
 335 does not detect all of the modes evidenced for OCS by the EMD analysis. Figure 7 shows the PDF  
 336 of the MLO time series, whose EMD composition has been exemplified through Fig. 3 to Fig. 4.  
 337 It is seen that the Fourier analysis is capable of extracting the annual cycle. In contrast, the  
 338 remaining modes, which EMD identifies in Fig. 3, are lost in a broad low pass spectrum with a  
 339 zero-frequency peak. Figure 7 also shows, for comparison, the PDF of the second IMF, which  
 340 extracts the annual cycle from the original signal. It can be seen that the PDF of the second IMF  
 341 exactly matches the peak of the annual cycle in the PDF signal, which allows us to stress the  
 342 property of EMD to extract the relevant modes from the signal. An analysis based solely on the  
 343 PDF of the signal would conclude the presence of a single dominant mode and a low-pass  
 344 component with a peak at zero frequency, which parallels the EMD residue and IMF4. In contrast,  
 345 EMD can correctly identify the annual cycle but can also reveal cyclic mode in the lower frequency  
 346 range with a characteristic time of  $\sim 10$  years (IMF4). In addition, EMD reveals an intermediate  
 347 mode that can be linked to inter-annual variability of characteristic time scales of 2-3 years, which,  
 348 in turn, may be associated with influences from the QBO, e.g. (Ray et al., 2020).



349  
 350 **Figure 7.** Power Density Function of the whole signal derived from a Fourier analysis of the MLO  
 351 OCS monthly mean mole fraction time series over the past 20 years (“Signal PDF”), and the second  
 352 IMF extracted through the EMD analysis.


353

### 354 **3. Results for the NOAA network: OCS measurements for the year range 2000-2020**

#### 355 3.1. The long-term EMD component, $\tau$


356 The OCS monthly mole fraction time series from the NOAA and cooperative sampling  
 357 stations listed in Tab. 1 has been processed to identify EMD analysis trends computed according  
 358 to Eq. (2). The EMD trend,  $\tau$ , results for the North-Hemisphere stations north of  $30^{\circ}\text{N}$  are shown  
 359 in Fig. 8. The EMD  $\tau$  yields the long-range behaviour with frequency lower than the threshold  
 360  $f_{th} = 3/N$ . The decomposition is shown in Fig. 8. All Northern stations consistently show a  
 361 decreasing atmospheric OCS mole fraction from 2015-2020.

362




363  
 364 **Figure 8.** EMD-derived trend determination,  $\tau$  component (Eq. (2)), for the NOAA stations in the  
 365 Northern Hemisphere at latitudes greater than 30N.  
 366

367 The long-term component is always relevant in terms of explained variance, as shown in  
 368 Tab. 2. In terms of standard deviation, the trend  $\tau$  explains more than  $\sim 15\%$  of the variability of  
 369 the whole signal,  $X(j), j = 1, \dots, N$ . We stress that the long-term components' variability in Fig. 8  
 370 reflects a good general agreement with original data. The overall mean is not distorted and long-  
 371 term local features at the scale of the threshold frequency are well reproduced. This is exemplified  
 372 in Fig. 9 for the case of the NWR station. In Fig. 9 we also show a comparison with the *lowess*  
 373 trend,  $\tau_l$ , which as for the case of the MLO station smooths the features at the scale of the threshold  
 374 frequency,  $f_{th} = 3/N$ . For the sake of brevity, the comparison between  $\tau$  and  $\tau_l$  is not shown in  
 375 the paper for all stations. However, the supplemental material has provided this comparison for  
 376 the interested reader. Here we stress that the *lowess* smoothing agrees with EMD in detecting a  
 377 decline in OCS atmospheric column amount since 2015-2016.




378  
 379 **Figure 9.** OCS monthly averages (2000 to 2020) for the NWR station and trend analysis according  
 380 to EMD (Eq. (2)) and the non-parametric *lowess* approach (see text in the paper).  
 381

382 The results for the stations between 30N and 30S are shown in Fig. 10. Consistent with  
 383 what has been shown for the Northern Hemisphere, we see a decreasing trend for the three stations  
 384 since 2015-2016.  
 385



386  
 387 **Figure 10.** EMD-derived trend determination,  $\tau$  component (Eq. (2)) for the NOAA stations  
 388 between 30°N and 30°S.  
 389

390 Finally, Fig. 11 shows the results for the three stations in the Southern Hemisphere. Also  
 391 in this case, we have that the three stations show a negative trend since 2015-2016, which is  
 392 strongly consistent with the findings we have shown for the other NOAA stations after these years.



393

394 **Figure 11.** Trend analysis for the NOAA stations in the Southern Hemisphere. We note that for  
 395 PSA, the trend seems to have reversed from a decreasing one since about 2010. However, it is  
 396 likely that the trend at PSA may be influenced by contamination in sampling equipment used at  
 397 that site in the first half of the record (2000-2010). The record is certainly quite a bit noisier prior  
 398 to 2010 than after it.

399

400

401 An essential aspect of the analysis we have shown with the 20-year long time series is  
 402 the presence of a relatively large variance of the OCS signal at frequencies below the threshold  
 403  $f_{th} = \frac{3}{N}$  which may reflect scales of the general atmospheric circulation, the climate forcing or  
 404 even the long-term changes in the magnitude of overall or total OCS emissions, e.g, (Zumkehr et  
 405 al., 2018).

406

407 The low-frequency variability is shown in Tab. 2 in terms of the standard deviation, i.e.,  
 408 the variability strength, of the EMD trend  $\tau$  (computed according to Eq. (2) ) and the original  
 409 monthly observations,  $X(j), j = 1, \dots, N$ .

410

411 **Table 2.** Variability (in terms of standard deviation) of the EMD trend  $\tau$  (Eq. (2)) and the original signal,  
 412  $X(j), j = 1, \dots, N$ , for the 20 year-long time series analyzed in this paper.

Station	Code	Lat [°N]	Lon [°W]	Elevation [masl]	Variability [ppt]		
					Trend, $\tau$	Signal, $X(j)$	% Ratio Trend/Signal
Alert, Nunavut, Canada	ALT	82.4508	62.5072	185	5.38	39.64	13.6
Point Barrow, USA	BRW	71.3230	156.6114	11	6.34	40.87	15.5
Cape Grim, Tasmania	CGO	-40.683	144.6900	94	4.26	14.76	28.8
Harvard Forest, USA	HFM	42.5378	72.1714	340	8.34	49.53	16.8
Cape Kumukahi, USA	KUM	19.7371	155.0116	0.30	6.10	22.48	27.1
Park Falls, USA	LEF	45.9451	90.2732	472	9.62	44.77	21.4
Mace Head, Ireland	MHD	53.3260	9.899	5.00	7.00	33.35	20.9

Mauna Loa, USA	MLO	19.5362	155.5763	3397	6.98	17.90	38.9
Niwot Ridge, USA	NWR	40.0531	105.5864	3523	7.65	19.91	38.4
Palmer Station, Antarctica	PSA	-64.7742	64.0527	10	8.52	20.03	42.5
Tutuila, American Samoa	SMO	-14.2474	170.5644	42	5.80	12.85	45.1
South Pole, Antarctica	SPO	-89.98	24.8	2810	2.65	14.69	6.40
Summit, Greenland	SUM	72.5962	38.422	3209	8.24	34.19	24.1
Trinidad Head, USA	THD	41.0541	124.151	107	9.95	41.40	24.0

413

414

415

416

417

418

419

From Tab. 2, we see that the trend or long-term variability is in between ~15-40% of the total power of the signal. Therefore, this component is not negligible with respect to the yearly cycle. In effect, from Fig.s 8-11, we see that the variability has consistently increased in the last few years, which leads us to conclude that the OCS mole fraction has entered a worldwide phase of decline. These findings suggest a recent broad-scale atmospheric decline that is captured by measurements at all of the NOAA sites.

420

421

422

423

In conclusion, we can say that the twenty-year OCS record at all sites shows a consistent low-frequency component, which yields a complex behaviour with a generally increasing trend up to 2015, a temporary decrease during 2009, and finally a decline in the last 6-7 years.

424

### 3.2. Oscillatory modes

425

426

427

428

429

430

431

432

433

434


435

436

437

438


Although this study focuses mainly on assessing low-frequency components in NOAA's OCS records over the past twenty years, EMD also extracts other relevant modes in the time series. Throughout the paper, we have already noticed the strong presence of the annual cycle, which is due to the summer OCS drawdown by vegetation. However, EMD analysis has also revealed modes of frequency  $> \frac{3}{N}$ . In principle, this rich variability could be associated with climate characteristic scales such as the QBO (~2 years), (El-Nino (~2 – 7 years), or simply interannual variability linked to biogenic activities. The in-depth analysis of these modes is not the present study's focus. However, we highlight them here for the benefit of the reader and to incite further studies. The peak frequencies of the IMF 2 to 4 are shown in Fig. 12 as a function of the station. From Fig. 11, we see a great consistency among the various stations. The IMF2 represents the annual cycle, with frequency  $f_o = \frac{1}{12} = 0.0833 \text{ month}^{-1}$ , and we see that IMF2 at all stations is peaked at this frequency. In Fig. 11, we have also drawn the sub-tone frequency,  $f_o/2$ ,  $f_o/4$  and  $f_o/10$  to help to identify where the observed peak frequencies accumulate.



439  
 440 **Figure 12.** Peak frequencies of the IMF from 2 to 4 as a function of the station. The figure also  
 441 shows the subtone frequencies of the annual cycle  $f_0$ , that is  $f_0/2$ ,  $f_0/4$  and  $f_0/10$  to identify where  
 442 the observed peak frequencies accumulate. The grey area gives the range of the QBO mean cycle,  
 443 which has a periodicity of 28-29 months.  
 444


445 It is seen that the IMF4 tends to accumulate at the frequency  $f_0/10$ , which is lower than  
 446 the threshold frequency,  $f_{th}$ . In effect, the IMF4 has been moved to the trend  $\tau$ , according to its  
 447 definition of Eq. (2). Much more interesting is the behavior of the IMF3. This is the faintest among  
 448 the three shown in Fig. 11 and shows good consistency with the QBO mean cycle, potentially  
 449 related to its influence on atmospheric mixing processes (Ray et al., 2020), or on the natural the  
 450 balance of OCS sources and sinks. The presence of this frequency within the range  $f_0/4$ ,  $f_0/2$  is  
 451 consistent with the IMF3 being linked to QBO.

452 Before ending this section, we also show examples of the Huang-Hilbert transforms or *hht*  
 453 used to check for time-dependence of the frequency. The transform is exemplified in Fig. 13 for  
 454 the second mode or IMF2 related to the EMD decomposition of the MLO OCS time series. The  
 455 transform gives the frequency as a function of the time (expressed in months in Fig. 12), and each  
 456  $(t, f)$  pair has assigned an instantaneous strength or amplitude (in ppt) according to the color bar.  
 457 In the case of a pure sine wave of frequency,  $A \sin(2\pi ft)$ , the *hht* would give a flat line equal to  
 458  $f$  and a constant amplitude equal to  $A$ . To clarify the meaning of the transform, in Fig. 13 we also  
 459 show the flat line corresponding to the annual cycle, that is  $1/12 \text{ month}^{-1}$



460  
 461 **Figure 13.** The plot shows the *hht* transform which represents the instantaneous frequency  
 462 spectrum of the IMF2 component decomposed from the original mixed signal for the MLO station.  
 463 For comparison, the plot also shows the line corresponding to the annual cycle.  
 464

465 We see that the frequency oscillates around the annual cycle, showing that just one  
 466 dominant harmonic governs the time dependence of IMF2. For IMF3, see Fig. 14, we have that  
 467 the intensity of the amplitude is fainter and again is close to the peak frequency. For IMF3, we see  
 468 an amplitude increase around 2005 (~58 months in Fig. 14). The frequency tends to increase, and  
 469 in fact, if we go back to Fig. 3, it appears that around 2005, the frequency of the oscillations of the  
 470 3<sup>rd</sup> mode tends to increase. However, for IMF3 the *hht* transform is close to the frequency 0.042  
 471 1/month, computed with the PDF analysis shown in Fig. 4.



472  
 473 **Figure 14.** As Fig. 12 but for IMF3. For comparison, the plot also shows the flat line corresponding  
 474 to frequency  $0.042 \text{ month}^{-1}$ , which the PDF analysis has extracted from the mode IMF3.  
 475

476 Note that the slight bump we see at around 2005 (month 58) is a transient phenomenon,  
 477 which seems to relax back to a stationary behaviour in a time span of the basic period of 1 year.

478

#### 479 **4. Discussion and Conclusions**

480 In this study, monthly average time series of OCS have been analyzed using data from the  
 481 NOAA/GML network covering 2000 to 2020. The analysis has been performed by using the  
 482 Empirical Mode Decomposition, which decomposes a given time series in its primary cycles plus  
 483 a trend. The method is non-parametric, and there is no need to specify a trend model as generally  
 484 done with other approaches.  
 485

486 EMD is more suitable than traditional methods for the analysis of nonlinear and  
 487 nonstationary signals. However, the straightforward applications of the technique could lead to  
 488 misuse if its known limitations and basic assumptions are not carefully considered. EMD still has  
 489 some open issues about its formal characterization when operating on a broadband signal, such as  
 490 white noise, e.g., (Z. Wu & Huang, 2010). In our analysis, this issue has been minimized by  
 491 resorting to decomposition, which, while non-exact, still provides an approximation of the given  
 492 signal (Wang et al., 2010). The EMD method we use to calculate the decomposition has been  
 493 implemented with the two basic stopping criteria recommended by (Wang et al., 2010) *to obtain*  
 494 *physically meaningful results*. The stopping rules include a Cauchy criterion, e.g., (Wang et al.,  
 495 2010) to stop the iteration from getting a given IMF and an Energy ratio criterion, e.g., (Wang et  
 496 al., 2010) to stop the EMD decomposition. In this way, as stressed by (Wang et al., 2010), the  
 497 EMD implementation yields an approximation with respect to the cubic spline basis but avoids  
 498 resulting in IMFs that have no physical significance.

499

500 In addition, we remark that other problems could affect EMD performance in practice  
501 (Huang et al., 1998, 2003), especially in measurement noise. One limitation is the difficulty of  
502 carrying out a clean separation in IMFs when their local frequencies are too close, e.g., (Stallone  
503 et al., 2020). In some cases, this separation could be improved by applying the so-called Ensemble  
504 Empirical Mode Decomposition (EEMD) (Z. Wu & Huang, 2009), an approach taken in this paper,  
505 which adds random noise the observations.

506 Finally, we constrain EMD by specifying the maximum number of modes and a frequency  
507 threshold to separate lower frequencies from the annual cycle. In effect, the stopping criteria  
508 (Wang et al., 2010) embedded in the most updated EMD software tool by Matlab (we used the  
509 release 2020b in this study) do not provide a reliable strategy to separate the trend from pure  
510 modes. Therefore, we have shown that frequency thresholding and a suitable limitation of modes  
511 are *best practices* for the successful use of Empirical Mode Decomposition.

512

513 With this in mind, the decomposition in cyclic modes of the OCS series has shown the  
514 presence of low-frequency time scales of  $\sim 10$  years. Furthermore, the low-frequency component  
515 yields a long-range time evolution, indicating a decline in OCS concentration in the atmosphere in  
516 the last 6-7 years. The reduction is seen in data obtained from all stations examined in the present  
517 work, consistent with a recent imbalance in total global OCS sources and losses. Moreover, we  
518 have shown that the OCS records exhibit a cyclic mode between 2-4 years, which may be linked  
519 to the Quasi Biennial Oscillation (QBO)).

520

521 In conclusion, a decreasing trend of OCS mole fraction has been observed in the last 6-7  
522 years at all NOAA/GML measurement sites. No matter the origin of the present OCS decay, the  
523 carbonyl sulfide atmospheric budget is currently unbalanced. We think that further analysis with  
524 global transport models could yield new insights in light of these most recent changes that we have  
525 identified and assessed in this study.

## 526 Acknowledgments

527 Measurements from flasks from the fourteen stations listed in Tab. 1 have been downloaded from the  
528 website <http://www.esrl.noaa.gov/gmd/hats/gases/OCS.html>. We give credit to the National  
529 Oceanic and Atmospheric Administration (NOAA)/Global Monitoring Laboratory (GML) in  
530 Boulder as a source of the data and appreciate the technical assistance of B. Hall, C. Siso,  
531 and I. Vimont, for assisting with the creation and maintenance of a reliable long-term  
532 atmospheric history of OCS mole fraction. This research was carried out in the framework of  
533 the project 'OT4CLIMA' which was funded by the Italian Ministry of Education, University and  
534 Research (D.D. 2261 del 6.9.2018, PON R&I 2014-2020 and FSC).

535

## 536 Open Research

537 The OCS data used in the paper are freely available from the website  
538 <http://www.esrl.noaa.gov/gmd/hats/gases/OCS.html>.

539

540

541 **References**

542 Aydin, M., Britten, G. L., Montzka, S. A., Buizert, C., Primeau, F., Petrenko, V., et al. (2020).

543 Anthropogenic Impacts on Atmospheric Carbonyl Sulfide Since the 19th Century

544 Inferred From Polar Firn Air and Ice Core Measurements. *Journal of Geophysical*545 *Research: Atmospheres*, 125(16). <https://doi.org/10.1029/2020JD033074>

546 Berry, J., Wolf, A., Campbell, J. E., Baker, I., Blake, N., Blake, D., et al. (2013). A coupled

547 model of the global cycles of carbonyl sulfide and CO<sub>2</sub>: A possible new window on the548 carbon cycle. *Journal of Geophysical Research: Biogeosciences*, 118(2), 842–852.549 <https://doi.org/10.1002/jgrg.20068>

550 Blake, N. J. (2004). Carbonyl sulfide and carbon disulfide: Large-scale distributions over the

551 western Pacific and emissions from Asia during TRACE-P. *Journal of Geophysical*552 *Research*, 109(D15), D15S05. <https://doi.org/10.1029/2003JD004259>

553 Brühl, C., Lelieveld, J., Crutzen, P. J., &amp; Tost, H. (2012). The role of carbonyl sulphide as a

554 source of stratospheric sulphate aerosol and its impact on climate. *Atmospheric Chemistry*555 *and Physics*, 12(3), 1239–1253. <https://doi.org/10.5194/acp-12-1239-2012>

556 Campbell, J. E., Carmichael, G. R., Chai, T., Mena-Carrasco, M., Tang, Y., Blake, D. R., et al.

557 (2008). Photosynthetic Control of Atmospheric Carbonyl Sulfide During the Growing

558 Season. *Science*, 322(5904), 1085–1088. <https://doi.org/10.1126/science.1164015>

559 Campbell, J. E., Whelan, M. E., Seibt, U., Smith, S. J., Berry, J. A., &amp; Hilton, T. W. (2015).

560 Atmospheric carbonyl sulfide sources from anthropogenic activity: Implications for

561 carbon cycle constraints. *Geophysical Research Letters*, 42(8), 3004–3010.562 <https://doi.org/10.1002/2015GL063445>

- 563 Campbell, J. E., Berry, J. A., Seibt, U., Smith, S. J., Montzka, S. A., Launois, T., et al. (2017).  
564 Large historical growth in global terrestrial gross primary production. *Nature*, *544*(7648),  
565 84–87. <https://doi.org/10.1038/nature22030>
- 566 Camy-Peyret, C., Liuzzi, G., Masiello, G., Serio, C., Venafrà, S., & Montzka, S. A. (2017).  
567 Assessment of IASI capability for retrieving carbonyl sulphide (OCS). *Journal of*  
568 *Quantitative Spectroscopy and Radiative Transfer*, *201*, 197–208.  
569 <https://doi.org/10.1016/j.jqsrt.2017.07.006>
- 570 Capparelli, V., Franzke, C., Vecchio, A., Freeman, M. P., Watkins, N. W., & Carbone, V.  
571 (2013). A spatiotemporal analysis of U.S. station temperature trends over the last century:  
572 On the Temperature Trends over the United States. *Journal of Geophysical Research:*  
573 *Atmospheres*, *118*(14), 7427–7434. <https://doi.org/10.1002/jgrd.50551>
- 574 Coughlin, K. T., & Tung, K. K. (2004). 11-Year solar cycle in the stratosphere extracted by the  
575 empirical mode decomposition method. *Advances in Space Research*, *34*(2), 323–329.  
576 <https://doi.org/10.1016/j.asr.2003.02.045>
- 577 Echeverría, J. C., Crowe, J. A., Woolfson, M. S., & Hayes-Gill, B. R. (2001). Application of  
578 empirical mode decomposition to heart rate variability analysis. *Medical & Biological*  
579 *Engineering & Computing*, *39*(4), 471–479. <https://doi.org/10.1007/BF02345370>
- 580 Gardiner, T., Forbes, A., de Mazière, M., Vigouroux, C., Mahieu, E., Demoulin, P., et al. (2008).  
581 Trend analysis of greenhouse gases over Europe measured by a network of ground-based  
582 remote FTIR instruments. *Atmospheric Chemistry and Physics*, *8*(22), 6719–6727.  
583 <https://doi.org/10.5194/acp-8-6719-2008>
- 584 Glatthor, N., Höpfner, M., Baker, I. T., Berry, J., Campbell, J. E., Kawa, S. R., et al. (2015).  
585 Tropical sources and sinks of carbonyl sulfide observed from space: TROPICAL

586 SOURCES AND SINKS OF COS. *Geophysical Research Letters*, 42(22), 10,082-  
587 10,090. <https://doi.org/10.1002/2015GL066293>

588 Huang, N. E., Shen, Z., Long, S. R., Wu, M. C., Shih, H. H., Zheng, Q., et al. (1998). The  
589 empirical mode decomposition and the Hilbert spectrum for nonlinear and non-stationary  
590 time series analysis. *Proceedings of the Royal Society of London. Series A:  
591 Mathematical, Physical and Engineering Sciences*, 454(1971), 903–995.  
592 <https://doi.org/10.1098/rspa.1998.0193>

593 Huang, N. E., Wu, M.-L. C., Long, S. R., Shen, S. S. P., Qu, W., Gloersen, P., & Fan, K. L.  
594 (2003). A confidence limit for the empirical mode decomposition and Hilbert spectral  
595 analysis. *Proceedings of the Royal Society of London. Series A: Mathematical, Physical  
596 and Engineering Sciences*, 459(2037), 2317–2345.  
597 <https://doi.org/10.1098/rspa.2003.1123>

598 Kettle, A. J. (2002). Global budget of atmospheric carbonyl sulfide: Temporal and spatial  
599 variations of the dominant sources and sinks. *Journal of Geophysical Research*,  
600 107(D22), 4658. <https://doi.org/10.1029/2002JD002187>

601 Krysztofíak, G., Té, Y. V., Catoire, V., Berthet, G., Toon, G. C., Jégou, F., et al. (2015).  
602 Carbonyl Sulphide (OCS) Variability with Latitude in the Atmosphere. *Atmosphere-  
603 Ocean*, 53(1), 89–101. <https://doi.org/10.1080/07055900.2013.876609>

604 Laurenza, M., Vecchio, A., Storini, M., & Carbone, V. (2012). Quasi-Biennial Modulation of  
605 Galactic Cosmic Rays. *The Astrophysical Journal*, 749(2), 167.  
606 <https://doi.org/10.1088/0004-637X/749/2/167>

- 607 Lee, T., & Ouarda, T. B. M. J. (2011). Prediction of climate nonstationary oscillation processes  
608 with empirical mode decomposition. *Journal of Geophysical Research*, *116*(D6),  
609 D06107. <https://doi.org/10.1029/2010JD015142>
- 610 Lee, T., & Ouarda, T. B. M. J. (2012). Stochastic simulation of nonstationary oscillation  
611 hydroclimatic processes using empirical mode decomposition: Simulation of NSO Using  
612 EMD. *Water Resources Research*, *48*(2). <https://doi.org/10.1029/2011WR010660>
- 613 Lejeune, B., Mahieu, E., Vollmer, M. K., Reimann, S., Bernath, P. F., Boone, C. D., et al.  
614 (2017). Optimized approach to retrieve information on atmospheric carbonyl sulfide  
615 (OCS) above the Jungfraujoch station and change in its abundance since 1995. *Journal of*  
616 *Quantitative Spectroscopy and Radiative Transfer*, *186*, 81–95.  
617 <https://doi.org/10.1016/j.jqsrt.2016.06.001>
- 618 Loh, C.-H. (2004). Application of the Empirical Mode Decomposition-Hilbert Spectrum Method  
619 to Identify Near-Fault Ground-Motion Characteristics and Structural Responses. *Bulletin*  
620 *of the Seismological Society of America*, *91*(5), 1339–1357.  
621 <https://doi.org/10.1785/0120000715>
- 622 Maseyk, K., Berry, J. A., Billesbach, D., Campbell, J. E., Torn, M. S., Zahniser, M., & Seibt, U.  
623 (2014). Sources and sinks of carbonyl sulfide in an agricultural field in the Southern  
624 Great Plains. *Proceedings of the National Academy of Sciences*, *111*(25), 9064–9069.  
625 <https://doi.org/10.1073/pnas.1319132111>
- 626 Montzka, S. A., Calvert, P., Hall, B. D., Elkins, J. W., Conway, T. J., Tans, P. P., & Sweeney, C.  
627 (2007). On the global distribution, seasonality, and budget of atmospheric carbonyl  
628 sulfide (COS) and some similarities to CO<sub>2</sub>. *Journal of Geophysical Research*, *112*(D9),  
629 D09302. <https://doi.org/10.1029/2006JD007665>

- 630 Ogée, J., Sauze, J., Kesselmeier, J., Genty, B., Van Diest, H., Launois, T., & Wingate, L. (2016).  
631 A new mechanistic framework to predict OCS fluxes from soils. *Biogeosciences*, *13*(8),  
632 2221–2240. <https://doi.org/10.5194/bg-13-2221-2016>
- 633 Ray, E. A., Portmann, R. W., Yu, P., Daniel, J., Montzka, S. A., Dutton, G. S., et al. (2020). The  
634 influence of the stratospheric Quasi-Biennial Oscillation on trace gas levels at the Earth’s  
635 surface. *Nature Geoscience*, *13*(1), 22–27. <https://doi.org/10.1038/s41561-019-0507-3>
- 636 Stallone, A., Cicone, A., & Materassi, M. (2020). New insights and best practices for the  
637 successful use of Empirical Mode Decomposition, Iterative Filtering and derived  
638 algorithms. *Scientific Reports*, *10*(1), 15161. <https://doi.org/10.1038/s41598-020-72193-2>
- 639 Thoning, K. W., Tans, P. P., & Komhyr, W. D. (1989). Atmospheric carbon dioxide at Mauna  
640 Loa Observatory: 2. Analysis of the NOAA GMCC data, 1974-1985. *Journal of*  
641 *Geophysical Research: Atmospheres*, *94*(D6), 8549–8565.  
642 <https://doi.org/10.1029/JD094iD06p08549>
- 643 Wang, G., Chen, X.-Y., Qiao, F.-L., Wu, Z., & Huang, N. E. (2010). On Intrinsic Mode  
644 Function. *Advances in Adaptive Data Analysis*, *02*(03), 277–293.  
645 <https://doi.org/10.1142/S1793536910000549>
- 646 Whelan, M. E., Lennartz, S. T., Gimeno, T. E., Wehr, R., Wohlfahrt, G., Wang, Y., et al. (2018).  
647 Reviews and syntheses: Carbonyl sulfide as a multi-scale tracer for carbon and water  
648 cycles. *Biogeosciences*, *15*(12), 3625–3657. <https://doi.org/10.5194/bg-15-3625-2018>
- 649 Wu, Y., & Shen, B.-W. (2016). An Evaluation of the Parallel Ensemble Empirical Mode  
650 Decomposition Method in Revealing the Role of Downscaling Processes Associated with  
651 African Easterly Waves in Tropical Cyclone Genesis. *Journal of Atmospheric and*  
652 *Oceanic Technology*, *33*(8), 1611–1628. <https://doi.org/10.1175/JTECH-D-15-0257.1>

653 Wu, Z., & Huang, N. E. (2009). Ensemble Empirical Mode Decomposition: A Noise-Assisted

654 Data Analysis Method. *Advances in Adaptive Data Analysis*, 01(01), 1–41.

655 <https://doi.org/10.1142/S1793536909000047>

656 Wu, Z., & Huang, N. E. (2010). On the Filtering Properties of the Empirical Mode

657 Decomposition. *Advances in Adaptive Data Analysis*, 02(04), 397–414.

658 <https://doi.org/10.1142/S1793536910000604>

659

**Long-term (2000–2020) variability of in situ time series of Carbonyl Sulfide**

*C. Serio<sup>1\*</sup>, S. Montzka<sup>2</sup>, G. Masiello<sup>1</sup>, V. Carbone<sup>3</sup>*

<sup>1</sup>Scuola di Ingegneria, Università degli studi della Basilicata, Potenza, Italy.

<sup>2</sup>NOAA Earth System Research Laboratory, Boulder, Colorado, USA

<sup>3</sup>Dipartimento di Fisica, Università della Calabria, Rende, Italy.


]

**Contents of this file**


Figures S1 to S14

**Introduction**


The supplemental material compares the OCS monthly times series (flask measurements), the *lowess* trend, and the EMD trend for the 14 stations examined in the present work. The list of the 14 stations and related acronyms can be found in Tab. 1 of the paper.




**Figure S1.:** ALT station: Original OCS time series, *lowess* trend,  $\tau_l$  and EMD trend  $\tau$ .




**Figure S2:** BRW station: Original OCS time series, *lowess* trend,  $\tau_l$  and EMD trend  $\tau$ .




**Figure S3:** CGO station: Original OCS time series, *lowess* trend,  $\tau_l$  and EMD trend  $\tau$ .




**Figure S4:** HFM station: Original OCS time series, *lowess* trend,  $\tau_l$  and EMD trend  $\tau$ .




**Figure S5:** KUM station: Original OCS time series, *lowess* trend,  $\tau_l$  and EMD trend  $\tau$ .




**Figure S6:** LEF station: Original OCS time series, *lowess* trend,  $\tau_l$  and EMD trend  $\tau$ .




**Figure S7:** MHD station: Original OCS time series, *lowess* trend,  $\tau_l$  and EMD trend  $\tau$ .




**Figure S8:** MLO station: Original OCS time series, *lowess* trend,  $\tau_l$  and EMD trend  $\tau$ .




**Figure S9:** NWR station: Original OCS time series, *lowess* trend,  $\tau_l$  and EMD trend  $\tau$ .




**Figure S10:** PSA station: Original OCS time series, *lowess* trend,  $\tau_l$  and EMD trend  $\tau$ .




**Figure S11:** SMO station: Original OCS time series, *lowess* trend,  $\tau_l$  and EMD trend  $\tau$ .



**Figure S12:** SPO station: Original OCS time series, *lowess* trend,  $\tau_l$  and EMD trend  $\tau$ .



**Figure 13:** SUM station: Original OCS time series, *lowess* trend,  $\tau_l$  and EMD trend  $\tau$ .



**Figure S14:** THD station: Original OCS time series, *lowess* trend,  $\tau_l$  and EMD trend  $\tau$ .