

Global-scale shifts in Anthropocene rooting depths pose unexamined consequences for critical zone functioning

Hauser Emma¹, Sullivan Pamela L², Flores Alejandro N.³, Hirmas Daniel⁴, and Billings Sharon A¹

¹University of Kansas

²Oregon State University

³Boise State University

⁴University of California, Riverside

November 16, 2022

Abstract

Rooting depth is an ecosystem trait that determines the extent of soil development and carbon (C) and water cycling. Recent hypotheses propose that human-induced changes to Earth's biogeochemical cycles propagate deeply due to rooting depth changes from agricultural and climate-induced land cover changes. Yet, the lack of a global-scale quantification of rooting depth responses to human activity limits knowledge of hydrosphere-atmosphere-lithosphere feedbacks in the Anthropocene. Here we use land cover datasets to demonstrate that root depth distributions are changing globally as a consequence of agricultural expansion truncating depths above which 99% of root biomass occurs (D99) by ~60 cm, and woody encroachment linked to anthropogenic climate change extending D99 in other regions by ~38 cm. The net result of these two opposing drivers is a global reduction of D99 by 5%, or ~8 cm, representing a loss of ~11,600 km³ of rooted volume. Projected land cover scenarios in 2100 suggest additional future D99 shallowing of up to 30 cm, generating further losses of rooted volume of ~43,500 km³, values exceeding root losses experienced to date and suggesting that the pace of root shallowing will quicken in the coming century. Losses of Earth's deepest roots — soil-forming agents — suggest unanticipated changes in fluxes of water, solutes, and C. Two important messages emerge from our analyses: dynamic, human-modified root distributions should be incorporated into earth systems models, and a significant gap in deep root research inhibits accurate projections of future root distributions and their biogeochemical consequences.

Global-scale shifts in Anthropocene rooting depths pose unexamined consequences for critical zone functioning

Emma Hauser¹, Pamela L. Sullivan², Alejandro N. Flores³, Daniel Hirmas⁴, Sharon A. Billings¹

¹Department of Ecology and Evolutionary Biology and Kansas Biological Survey and Center for Ecological Research, The University of Kansas, Lawrence, KS, USA.

²College of Earth, Ocean, and Atmospheric Science, Oregon State University, Corvallis, OR, USA.

³Department of Geosciences, Boise State University, Boise, ID, USA

⁴ Department of Environmental Sciences, University of California, Riverside, CA, USA.

Corresponding author: Emma Hauser, Sharon Billings (emma.hauser@umt.edu, sharon.billings@ku.edu)

Key Points:

- Rooting depths are changing globally; the depth to which 99% of crop roots extend is shallower by ~ 60 cm compared to natural systems.
- In other regions, such as those experiencing woody encroachment, roots are deepening by ~38 cm compared to previous dominant vegetation.
- These opposing phenomena result in average rooting depths that are ~8 cm shallower today and projected to become ~30 cm shallower by 2100.

Abstract

Rooting depth is an ecosystem trait that determines the extent of soil development and carbon (C) and water cycling. Recent hypotheses propose that human-induced changes to Earth's biogeochemical cycles propagate deeply due to rooting depth changes from agricultural and climate-induced land cover changes. Yet, the lack of a global-scale quantification of rooting depth responses to human activity limits knowledge of hydrosphere-atmosphere-lithosphere feedbacks in the Anthropocene. Here we use land cover datasets to demonstrate that root depth distributions are changing globally as a consequence of agricultural expansion truncating depths above which 99% of root biomass occurs (D99) by ~60 cm, and woody encroachment linked to anthropogenic climate change extending D99 in other regions by ~38 cm. The net result of these two opposing drivers is a global reduction of D99 by 5%, or ~8 cm, representing a loss of ~11,600 km³ of rooted volume. Projected land cover scenarios in 2100 suggest additional future D99 shallowing of up to 30 cm, generating further losses of rooted volume of ~43,500 km³, values exceeding root losses experienced to date and suggesting that the pace of root shallowing will quicken in the coming century. Losses of Earth's deepest roots — soil-forming agents — suggest unanticipated changes in fluxes of water, solutes, and C. Two important messages emerge from our analyses: dynamic, human-modified root distributions should be incorporated into earth systems models, and a significant gap in deep root research inhibits accurate projections of future root distributions and their biogeochemical consequences.

Plain Language Summary

The distribution of plant roots helps determine the extent of nutrient, C and water cycling beneath Earth's surface. Human activities, including land use and climate change, can change the distribution of plant roots and their activities across the globe. Here, we used global land cover datasets in combination with field-generated rooting depth equations to estimate global scale changes to roots both now and into the future. Globally, roots are shallower than they would be in the absence of human activity due to extensive land conversion to agriculture. In some regions, human-promoted woody encroachment induces root elongation, but this effect is overwhelmed by the spatial extent of agricultural conversion. In the future, roots likely will become shallower at an even faster pace. In future projections, deep roots appear especially vulnerable to loss, prompting numerous questions for additional field- and modeling-based

studies about the ways nutrients, C, and water will cycle in a future with fewer deep roots. We provide a foundation for those questions by demonstrating human influence on the roots that shape the character of Earth's skin.

1 Introduction

Roots are subsurface engineers, and their distributions drive ecosystem-scale processes (Maeght et al., 2013; Pierret et al., 2016; Sullivan et al., 2022) such as soil development (Brantley et al., 2017; Hasenmueller et al., 2017; Austin et al., 2018), release of mineral-bound nutrients (Jobbagy & Jackson, 2001; Hasenmueller et al., 2017; Austin et al., 2018), subsoil water flow paths and residence time (Zhang et al., 2015; Fan et al., 2017), and deep C fluxes (Richter and Markewitz, 1995; Schenk, 2007; Pierret et al., 2016; Fan et al., 2017; Billings et al., 2018). The dominant drivers of rooting distributions are plant functional type (PFT, Jackson et al., 1996) and variation in water availability (Schenk, 2007; Nippert et al., 2007; Fan et al., 2017), both of which are changing in response to anthropogenic land cover conversion, as well as altered atmospheric composition and concomitant changes in climate (Edgeworth et al., 2001; Cramer et al., 2010; Ellis et al., 2010). This observation suggests that rooting depth distributions are likely undergoing changes due to human activities in the critical zone (CZ, Earth's living skin, Jordan et al., 2001).

Quantifying large-scale, human-induced changes to rooting distributions and how they may differ regionally is a critical step towards a greater understanding of how roots govern large-scale, sub-surface and surface processes. In spite of widespread recognition of the importance of root depth (Maeght et al., 2013; Pierret et al., 2016) and a growing recognition of the great depths to which roots can penetrate (Stone & Kalisz, 1991; Nepstad et al., 1994; Canadell et al., 1996; Schenk & Jackson, 2002a; Schenk & Jackson et al., 2002b; Fan et al., 2017), large-scale responses of rooting depths to anthropogenic perturbations of the biosphere have been poorly characterized. This knowledge gap is due in part to the challenges of accessing relatively deep soil horizons (Maeght et al., 2013), as well as the challenge of unraveling the vast complexity of Earth's subsurface systems. One consequence of poorly defined rooting distributions at large spatial scales is generalized representations of rooting parameters in land models (McCormack et al., 2015; Iversen et al., 2017; McCormack et al., 2017). Although many land models, such as the Community Land Model (CLM), represent changes to roots with land use change (Lawrence et

al., 2019), some land cover types are not well represented in these models. For example, crops in CLMs are assigned the same rooting depth as C3 grasses (Lawrence et al., 2019), though row crops, in particular, typically have far shallower roots than perennial plants (Canadell et al., 1996; DuPont et al., 2014; Billings et al., 2018). Given the plethora of CZ functions influenced by roots (Maeght et al., 2013; Pierret et al., 2016), poor characterization of rooting depths likely limits the accuracy of projected responses of the coupled terrestrial water, energy, and C cycles to climate in the Anthropocene.

Two Anthropocene phenomena occur at sufficient magnitude to potentially alter rooting distributions at the global scale. First, many regions have experienced conversion to annual row crops (Ramankutty & Foley, 1999; Ellis et al., 2010), a process that induces mortality of deep perennial root systems and replaces them with relatively shallow roots (Billings et al., 2018). In contrast, climate change and increasing atmospheric CO₂ concentrations are linked to root extension of extant woody plants (Iversen, 2010), and shifting ecoregion ranges may increase rooting depths where more deeply rooted woody vegetation becomes increasingly abundant in grasslands and tundra (Jackson et al., 1996; Harsch et al., 2009; Stevens et al., 2017; Wang et al., 2019). Studies exploring rooting depth typically focus on absolute rooting depths and their responses to climate or atmospheric CO₂ (Kleidon & Heimann, 1998; Kleidon, 2003) or, separately, land cover changes in specific regions of interest (Jaramillo et al., 2003; Hertel et al., 2009; DuPont et al., 2010). Despite known changes in global land cover (Ellis et al., 2010) that are associated with distinct rooting depths (Jackson et al., 1996; Zeng, 2001), as well as global analyses of the maximum extent of contemporary root depths (Schenk & Jackson, 2002a; Schenk & Jackson, 2002b; Schenk & Jackson, 2005), to date, no one has directly quantified the net change in rooting distributions at the global scale as a consequence of these opposing human activities.

Here we provide a first estimate of the extent to which rooting depths increase or decrease in response to land use and climate change and the volume of soil affected by this change. We also project how rooting depths and rooted volumes may change throughout the 21st century as more land is converted to agricultural and urban use, and as biome ranges continue to shift with changing climate. We emphasize that our focus is not on maximum rooting depths. Indeed, there

is a growing appreciation of the great depths to which vegetation can root (Stone & Kalisz 1991; Schenk & Jackson, 2002a; Schenk & Jackson, 2005; Maeght et al., 2013; Pierret et al., 2016; Fan et al., 2017) though the true maximum rooting depth may never be known in some systems (Kleidon, 2003; Pierret et al., 2016; Fan et al., 2017). Instead, we focus on the depths to which most or half (i.e., 99%, 95%, and 50%) of the root biomass of an ecosystem extends (Zeng, 2001), as well as changes to rooted soil volume. These metrics highlight the depths within which most roots reside as well as the soil volume through which most root distribution changes occur, both functionally consequential measures. Additionally, these metrics represent those for which much data exist, enabling the cross-system comparisons necessary to estimate the spatial extent of rooting depth changes in the Anthropocene. Our work thus reveals how anthropogenic, global-scale changes in rooting depth metrics are changing, thereby illuminating critical next steps to help us understand future CZ functioning.

2 Materials and Methods

We estimated the volume of soil influenced by human-promoted modification of root distributions. To do this, we estimated potential (i.e., no human influence), contemporary, and projected root distributions at the global scale by combining biome-specific rooting depth functions derived from empirical studies (described below) with spatially explicit land cover datasets. As a part of this process, we examined multiple datasets that, in theory, could help us estimate how humans modify rooting distributions. First, we offer a description of selected datasets followed by an explanation of our selection from those available.

We used satellite-derived, potential vegetation representing 15 land cover classes (Haxeltine & Prentice, 1996) and their potential global distribution in the absence of human activity at a 5-minute spatial resolution (Ramankutty & Foley, 1999). We compared potential vegetation classes to contemporary land cover as defined by the Global Land Cover 2000 (GLC2000) dataset (Bartolome & Belward, 2005). GLC2000 represents 22 land cover types, which are designated according to plant functional types ascribed to satellite images and ground-truthed by regional analysts. We aligned contemporary vegetation classifications with potential vegetation classes according to previously published frameworks for ecoregion designation (Bartolome & Belward, 2005), and augmented these classes to include a class for permafrost regions where

rooting depth is likely limited (Billings et al., 1997; Boike et al., 2018). These efforts resulted in 25 distinct land cover types for which rooting depths were assigned. Projected vegetation classes were similarly developed for four Shared Socioeconomic Pathway (SSP) and Representative Concentrations Pathway (RCP) scenarios using spatial projections of gridded, $0.5^\circ \times 0.5^\circ$ resolution land covers for the year 2100 (Hurtt et al., 2011). All maps were adjusted to the same resolution for analyses using the Raster package in R (Hijmans et al., 2019).

For all vegetation datasets except those above 60°N (described below), we estimated biome-specific rooting depths by assigning rooting depth functions derived from empirical data compiled in the Fine Root Ecology Database (FRED) and the National Ecological Observatory Network (NEON) database (Iversen et al., 2021; NEON 2021). These datasets have recently expanded rooting depth knowledge beyond earlier works (e.g., Jackson et al., 1996; Zeng, 2001; Schenk and Jackson, 2005) by accumulating new datapoints detailing root trait and distribution patterns in diverse biomes (Krasowski et al., 2018; Montagnoli et al., 2018; Lozanova et al., 2019; Andrade et al., 2020). However, to date no one has harmonized and analyzed these datasets to produce equations describing global rooting depth distributions. Their use here thus represents an advance in the ways we represent rooting depths and their distributions across the globe. Specifically, we used these datasets to estimate the depths by which rooting systems exhibit 50% (D50), 95% (D95), and 99% (D99) of their total biomass in each land cover type. To generate rooting depth functions, we assigned FRED and NEON rooting depth data to biomes according to the position of each datapoint on our modified GLC2000 land cover map. Each set of points was checked using Google Earth to ensure that datapoints were correctly assigned. Due to the resolution of the GLC2000 map, some shrubland and woodland categories were incorrectly identified as cropland; for these points, we reassigned shrub-covered areas to the open-closed deciduous shrubland class and woodlands to the open broadleaved deciduous forest class. We then fit depth-decay curves to each set of points for each biome using the model presented by Zeng (2001). Parameter values and their confidence intervals were obtained for depth-decay curves using a bootstrap procedure where curves were fit to randomly-selected samples (with replacement) of each set of points 1200 times as recommended by Lander (2013). By using the Zeng (2001) model, we assumed that rooting depth distributions remain similar for each vegetation functional type in the potential, contemporary, and future scenarios. The merit of

174 this assumption may vary with time but keeping the rooting depth of each biome's vegetation
175 type consistent across the Holocene and into the future allows us to parse the influence of land
176 cover change on rooting depths from that of less well-characterized phenomena.

177
178 To match the land cover classifications used in potential and contemporary vegetation maps to
179 biome classifications for which we have rooting depth equations, we modified estimated rooting
180 depth distributions for several land covers based on findings from region-specific literature. For
181 example, potential land cover datasets combine both polar and mid-latitude deserts into a single
182 desert category based on hydrologic regimes, yet rooting depths in polar deserts are often
183 constrained by permafrost. We thus separated these two desert regions, reassigning deserts in
184 polar regions to the 'tundra' classification above 60°N (Zhang et al., 2008). Further, in potential
185 and contemporary vegetation datasets, we reassigned evergreen forest and mixed vegetation
186 classes above 50°N to the 'boreal' vegetation classification given previously generated
187 vegetation maps of northern region forests (Brandt et al., 2013; Price et al., 2013), and also
188 assigned herbaceous and shrubland classes above 60°N to the class 'tundra' because these
189 regions exhibit low stature vegetation and lie in previously described tundra areas (Zhang et al.,
190 2008). To generate maps of rooting depth, we gave potential vegetation above 60°N that was
191 previously assigned to the polar desert class a rooting depth specific to permafrost-underlain
192 regions, where roots typically do not penetrate deeper than 30 cm and 50% of root biomass is
193 typically found within 10 cm (Billings et al., 1997; Zhang et al., 2008; Boike et al., 2018; Keuper
194 et al. 2020). For contemporary rooting depth maps, regions above 60°N were all assigned to
195 either a permafrost underlain tundra class or boreal class, which reflect recent measurements in
196 FRED and NEON datasets. Finally, because many remote sensing-based studies of regional
197 ecosystem fluxes omit large, lower latitude desert regions from their analyses due to the lack of
198 quantifiable ecosystem productivity in these systems (Zhao et al., 2005b), we omitted mid-
199 latitude deserts from rooting depth averages reported in the main text. Instead, we present rooting
200 depth metrics that incorporate the potential contribution of these mid-latitude deserts to global
201 root averages in Table 1 of the Supporting Information. Comparison of these results with those
202 reported in the text reveal an inflated influence of mid-latitude desert rooting depth estimates on
203 global averages that likely does not represent reality due to the low density of plants in true
204 deserts (Whitford & Duval, 2019). Ice-covered regions were also omitted from the analyses.

To assess potential effects of global-scale perturbations projected by the year 2100 on rooting depth distributions, we examined multiple SSP and RCP land cover projections from the Intergovernmental Panel on Climate Change (IPCC). Projected vegetation classes were developed for 4 SSP RCP scenarios (SSP2 RCP4.5, SSP1 RCP2.6, SSP4 RCP6.0, SSP5 RCP8.5). Landuse Harmonization datasets designate land cover classes more coarsely than either GLC2000 or potential vegetation datasets, delineating primary and secondary forest regions, primary and secondary non-forest regions, five agricultural classes, pastureland, rangeland, and urban regions (Hurtt et al., 2011). We assigned a rooting depth equation derived from agricultural croplands in the FRED and NEON datasets to all five agricultural classes in the Landuse Harmonization dataset. For secondary non-forests, pastures, and rangelands we assigned rooting depth equations representing herbaceous and grassland systems in the FRED and NEON datasets. Because most secondary forests in these scenarios were in the boreal region, we assigned secondary forests the average root depth value (107.5 cm) of mixed forests (130 cm) and boreal forests (85 cm). Primary forests were assigned depth values generated from the average of all forest classes in the contemporary dataset, and primary non-forests were assigned depths generated by averaging contemporary grassland and shrubland classes. Reflecting anticipated warming and large projected losses of permafrost in the northern hemisphere (Lawrence & Slater, 2005), rooting depths assigned in all future scenarios removed permafrost constraints.

We examined multiple datasets describing contemporary global root distributions (Schenk and Jackson, 2009) and landcover scenarios across time (Hurtt et al., 2011) as potential candidates for addressing the degree to which humans modify the rooted volume of Earth's subsurface. Such datasets have been pivotal in developing our understanding of and appreciation for the depths of deep roots (Schenk & Jackson, 2005; Schenk, 2005; Pierret et al., 2016), and the Landuse Harmonization (LUH) scenarios represent the best available data for future land cover classifications to date (Hurtt et al., 2020). However, the Schenk and Jackson dataset does not describe roots in agricultural lands, ploughed and fertilized lands, or wetlands (Schenk and Jackson, 2005), and is not divided into land cover classes that can be integrated with datasets describing potential and future land cover scenarios. The LUH scenarios combine land cover

classes in ways that result in the loss of important nuances in root distribution estimates in past and contemporary scenarios. For example, all forest types in LUH scenarios are grouped into ‘secondary’ and ‘primary’ forest rather than more region-specific forest classifications (Hurt et al., 2020). In contrast, employing the GLC2000 vegetation classes with rooting depths derived from FRED and NEON data, which include data from Jackson et al. (1996), permitted us to examine two key features of interest. First, this approach permitted incorporation of agricultural land cover classes — a feature that is absent in datasets featuring root distributions alone. Second, the Ramankutty and Foley (1999) dataset serves as the only spatially quantified representation of the potential land cover in the absence of human activity at a 5-minute resolution, allowing for detailed backcasting of estimates of human-induced changes to roots.

Using the R raster package (RStudio Team, 2017; Hijmans et al., 2019), we assigned rooting depth values to each land cover classification of the potential, contemporary, and projected vegetation maps, and calculated global means of each depth metric. After determining the differences in rooting depths across scenarios, we examined the spatial extent of depth changes to determine differences in rooted volume across scenarios. We then compared metrics across time using 95% confidence intervals of the mean estimates of global rooting depth metrics. Estimates of rooting depth, reflect measurement uncertainty, particularly at deeper depths (Schenk and Jackson, 2002b). However, because we applied root measurements in a consistent manner across potential, contemporary, and projected vegetation maps, we can assess relative differences of root distributions across these different scenarios. We performed correlated t-tests on pairs of rasterized parameter estimate maps (i.e., potential vs. contemporary, and contemporary vs. projected) to determine whether differences between these estimated rooting depth metrics are significantly different from zero. Data were assessed to ensure they met the assumptions of correlated t-tests, including independence of observations, normal distribution of the dependent variable, and no dependent variable outliers. Where data did not meet the assumptions, we ran Wilcoxon tests on the dataset pairs to assess differences in root depth metrics and reported the V-statistics and associated *P*-values generated from those tests.

3 Results

Comparisons of potential and contemporary land cover (Figures 1a and b) and their estimated rooting depths (Figures 1c and d) suggest that spatially averaged, global values of D99 are the net result of two competing phenomena: shallowing of roots in agricultural regions and deepening of roots in regions experiencing woody encroachment. Specifically, the global average D99 is 5% shallower (8 cm) under contemporary land cover distributions than if potential vegetation cover types covered Earth's terrestrial surface ($V = 7.11 \times 10^{11}$, Wilcoxon $P < 0.0001$; Figures 1c and d, Table S1). This represents a loss of rooted volume of $\sim 11,600 \text{ km}^3$. Values of D95 for contemporary land cover also express similar trends of root shallowing (6% or 5 cm, loss of $\sim 7250 \text{ km}^3$; $V = 7.06 \times 10^{11}$, Wilcoxon $P < 0.0001$; Figures S1a and b). Depth to 50% root biomass (D50), by comparison, displays relatively greater variation between contemporary and potential land cover, becoming 21% shallower (1.5 cm, 1300 km^3 , $V = 5.32 \times 10^{11}$, Wilcoxon $P < 0.0001$) on average (Figure S2).




Figure 1. Land cover and associated rooting depths under potential vegetation in the absence of human influence (left column) and current vegetation distribution (right column). (a) Potential vegetation cover in the absence of

human activity (b) Contemporary land cover distribution from Global Land Cover 2000 (GLC2000), modified to correspond to potential vegetation land cover classifications. (c) and (d) depict depths by which 99% of rooting biomass occurs (D99) under potential (c) and contemporary (d) land cover types. Inset histogram displays rooting depth distributions. Blue histogram reflects potential vegetation data, and red histogram contemporary land cover. Dashed vertical lines represent means. Appearance of a distinct color change from dark blue to light grey in Asia and Canada at 60°N in (c) is an artifact of restricting maximum rooting depth assignments to reflect well-characterized limitations imposed by frozen soils; this distinction is less evident in contemporary D99 maps (d) because of the higher spatial resolution of the GLC2000 dataset. Appearance of a distinct line at 50°N, especially evident in (d), reflects reassignment of mixed forests to the boreal forest class above this latitude (Brandt et al., 2013; Price et al., 2013). See text for reassignment details. While these lines are unrealistic, it reflects our current knowledge about root depths in northern regions and demonstrates the remaining need for additional work combining cryospheric studies and soil science to characterize root systems at relatively high latitudes.

Agricultural land conversion serves as a dominant influence on these global trends (Figures 2 and 3). Regions where roots experienced shallowing during the shift from potential to contemporary land cover are on average 43 cm shallower (23%) than potential vegetation distributions and represent ~48% of Earth's land surface (7.01×10^7 ha; Fig. 3). Thirty three percent of shallowing regions (2.28×10^7 ha) experience agricultural expansion. In these areas, perennial vegetation has been converted to agricultural land (defined here as annual crops and managed pasture), such that D99 has decreased by as much as 33% (60 cm). The remaining shallowing occurs primarily in some northern and arid regions, possibly due to increased disturbance (Harsch et al., 2009; Wang et al., 2020, Hurtt et al., 2020), urbanization (Lindsey & Bassuk, 1992; Day et al. 2010) and desertification (Lal, 2001, Zhao et al., 2005b). Where woody encroachment is evident in contemporary land cover data, D99 increased relative to potential vegetation by up to 39% (38 cm; note that here we use the phrase 'woody encroachment' to refer to both shrubland encroachment into grasslands, and forest encroachment into Arctic and alpine tundra). This result may overestimate current rooting depths if the rooting depths we assigned were derived from well-established, mature systems, given that woody plants in recently encroached systems likely have not yet achieved such depths (Stevens et al., 2017; Billings et al., 2018). Despite this possible overestimation, root deepening via woody encroachment does not overcome the effect of root shallowing, in part because of the smaller total fraction of Earth's terrestrial surface experiencing woody encroachment (35% or 5.06×10^7 ha).




Figure 2. Representation of rooting depth elongation due to woody encroachment (a and b) and rooting depth truncation due to agricultural expansion (c and d). Blue region in B demonstrates the belowground increase in roots shown in blue in Figure 3. Red region in D exemplifies loss of rooting system depth for red regions in Figure 3.




Figure 3. Mapped differences between potential and contemporary rooting depths. Red cells indicate a decrease in the depth to 99% of rooting biomass (D99) while blue cells indicate an increase in D99 resulting from contemporary vegetation distributions. Appearance of a distinct color change from dark blue to light grey and red in Asia and

Canada at 50°N reflects reassignment of mixed forests to the boreal forest class above this latitude (Brandt et al., 2013; Price et al., 2013). See Figure 1 caption for additional explanation.

Changes to rooting distributions by the year 2100 vary under different potential scenarios of climate and land use change as well as different societal responses to those changes. The SSP scenarios examined here represent global narratives including a scenario with few roadblocks to both mitigation of and adaptation to climate change (SSP1), moderate challenges to mitigation and adaptation (SSP2), a scenario of social inequality with many challenges to adaptation but few for mitigation (SSP4), and a strategy of fossil fuel dependence with many challenges to mitigation but few to social adaptation (SSP 5, Riahi et al., 2017). These narratives are used in conjunction with projected land use and climate (RCP) scenarios to model future societal and ecological conditions, on which we rely for our rooting distribution estimates.

Projections for the year 2100 suggest that the scenario with the largest cropland increase and relatively low radiative forcing enhancement from current levels (SSP1 RCP2.6, Figure 4a) generates the most extreme reduction of deep roots, truncating values of D99 by 30 cm ($V = 2.16 \times 10^{10}$, Wilcoxon $P < 0.0001$). The smallest shallowing of D99, 22.3 cm ($V = 1.77 \times 10^{10}$, Wilcoxon $P < 0.0001$), occurs under the highest emissions scenario (SSP5 RCP8.5, Figure 4b). As a result, the future rooted volume will be reduced by $\sim 32,400 \text{ km}^3$ to $\sim 43,500 \text{ km}^3$.




Figure 4. Projected changes of depth to 99% rooting biomass (D99) by the year 2100 relative to contemporary rooting depth distributions. Projections are based on land use and emissions changes under two combinations of Shared Socioeconomic Pathways (SSP) and Representative Concentration Pathways (RCP), SSP1 RCP2.6 (a) and SSP5 RCP8.5 (b). These two maps represent the scenario of greatest and least projected change, respectively. Red colors indicate root depth truncation or shallowing, and blue indicates elongation or deepening. Appearance of a

distinct color change from dark red to light grey in Asia at 50°N reflects reassignment of mixed forests to the boreal forest class above this latitude (Brandt et al., 2013; Price et al., 2013; see text for reassignment details).

Values of D50 for the year 2100 experience a shallowing of 3 cm across all assessed scenarios ($V = 2.47 \times 10^{10}$, Wilcoxon $P < 0.0001$; Figure S5), representing a loss of rooted soil volume of $\sim 4400 \text{ km}^3$. Though small relative to changes in deep root systems, this D50 shallowing is double that occurring during the previous $\sim 10,000 \text{ y}$ (Gupta, 2004) of anthropogenic land conversion (Figure S6).

4 Discussion

Our estimates of rooting depth and rooted soil volume suggest that root biomass throughout Earth's soils, even deep in the subsurface, has been and will continue to be vulnerable to human influence (Figures 2, 3, 4). Although maximum rooting depths are poorly characterized and are likely deeper than is typically appreciated (Maeght et al., 2013; Pierret et al., 2016; Fan et al., 2017), we demonstrate that the depths to which most or half of all rooting biomass reach (i.e., D99, D95, and D50) currently reflect human-induced, global-scale changes in land cover (Figure 1). We further demonstrate that root shallowing in agricultural regions ($\sim 60 \text{ cm}$ across $2.28 \times 10^7 \text{ ha}$ for D99) and root deepening in regions experiencing woody encroachment ($\sim 38 \text{ cm}$ across $5.06 \times 10^7 \text{ ha}$ for D99) result in a globally-averaged estimate of net 8 cm shallowing of D99 values. This represents a net loss of $\sim 11,600 \text{ km}^3$ of rooted volume to date in the Anthropocene.

In the future, rooting depth scenarios might be expected to reflect the elongating effects of woody encroachment on D99, D95, D50 and rooted soil volume to a yet greater extent, given the apparent role of rising atmospheric CO_2 concentrations in promoting woody encroachment (Devine et al. 2017). However, the four IPCC scenarios explored here suggest that by 2100, globally-averaged rooting distributions may become yet shallower relative to contemporary rooting depths (Figures 4, S4 and S5). Reduced rooting depths by 2100 are driven by substantial root shallowing across regions of Africa, the Middle East, Asia and Australia (Fig. 4), where deeply rooted shrublands are projected to transition to herbaceous grasslands and where there is continued agricultural and pasture expansion (Hurtt et al. 2020). In both cases, a more shallowly rooted, herbaceous vegetation cover replaces the current, more deeply rooted vegetation, either

as a consequence of shifting climate or land cover change. These transitions result in a nearly three-fold decrease in our two relatively deep rooting depth metrics (D95 and D99) and a two-fold decrease in D50 by the year 2100, suggesting that roots across Earth's subsurface will be subject to extensive additional anthropogenic changes in the future and that the deepest roots appear especially vulnerable to loss.

The global patterns we observed are strongly driven by trends in boreal and tundra regions, where mapped scenarios suggest patterns of both root shallowing and deepening (Figs. 2, 4, S5, and S6), and thus uncertainty about temporal dynamics of roots. While some studies hint that roots may deepen as soils currently designated as permafrost thaw (Harsch et al., 2009; Sistla et al., 2013; Malhotra et al., 2020; Wang et al., 2020), others suggest that long term changes in snowpack will produce extremes in soil freeze/thaw cycles that will reduce vegetation survival and rooting depth (Groffman et al., 2001; Blume-Werry et al., 2016). Most of our scenarios suggest deepening of D99 and D95 in northern regions over time, lending support to findings of deepening roots as permafrost thaws (Figs. 3 and 4) . However, contemporary D50 maps demonstrate shallowing relative to potential vegetation in these same regions (Fig. S6), implying that roots in boreal and tundra regions may be experiencing a more general change in the curvature of rooting depth distributions instead of consistently deepening over time. These observations support findings of altered root distributions where permafrost experiences altered seasonal cycles, such as longer growing seasons (Blume-Werry et al. 2019). Data describing rooting depths in these regions are more limited than in many other ecoregions (Iversen et al., 2021; NEON 2021), resulting in less certainty about future rooting depths in areas currently underlain by permafrost, and likely leading to the varied findings in our maps.

In maps of D50, additional regions also suggest that rooting depth distributions are undergoing a general change in curvature as a response to anthropogenic change. Shallowing D50 values are evident across potential, contemporary, and future scenarios (Figs. 3, 4, S5 and S6), and these D50 metrics appear to become shallower to a greater extent between contemporary and future (i.e., 2100) scenarios compared to the D50 changes that appear to have taken place already. This finding suggests that anthropogenically-induced changes in the root abundances of surficial soil horizons within the coming decades will likely exceed those of the past several millennia.

Shallowing D50 values occur alongside both shallowing and deepening of D99 and D95 values in different regions of the globe, hinting of a trend of reshaped root distributions. Recently collected data from the FRED and NEON databases make this change in curvature more apparent than some of the individual datasets on which they build (Canadell et al., 1996; Zeng, 2001; Schenk and Jackson, 2005), highlighting the importance of continuing to characterize the distribution of roots across the globe for understanding both the depths to which roots proliferate, and the shape of their depth distributions. These most recent advances in FRED and NEON D50 data emphasize that even relatively shallow soil horizons (*i.e.*, those expressed by D50), where both natural and agricultural species root, will undergo redistribution in the coming decades, with roots shifting the curvature of their distributions in response to regional changes in land use and climate.

There are myriad feasible consequences of altered rooting depth distributions for biogeochemical and hydrological fluxes that prompt intriguing hypotheses. For example, roots beneath the zone of maximum rooting density are attributed to developing the soils that mantle Earth's surface, so much so that they are referred to as the planet's biotic weathering front, where life — roots and microbes — promotes the dissolution of bedrock (Richter & Markewitz, 1995; Berner et al., 2003; Brantley et al., 2012; Pawlik, 2013; Dontsova et al., 2020). Results from the current study suggest that these biotic weathering forces in many temperate and tropical regions do not reach as deeply into the regolith as they did prior to human influence (Figure 3), prompting the hypothesis that the intensity of biotic processes responsible for soil formation at the bottom of the soil profile have declined in the Anthropocene. Further, a smaller volume of soil explored by rooting systems of some regions prompts the hypothesis that soil water storage capacity, nutrient replenishment, and solute losses from freshly weathered material have similarly declined (Swank, 1986; Nepstad et al., 1994; Berner, 1998). In contrast, in regions where root deepening is occurring, we might expect increases in the influences of biotic weathering deep in the soil profile.

Our findings serve as a useful starting point for formulating and probing these hypotheses. Although this study makes a first attempt at measuring the extent of anthropogenically-induced changes in rooting systems at a global scale, it also points to key knowledge gaps. The

uncertainty embedded in the projections reported here highlights the substantial need for better quantification of rooting distributions in diverse biomes, particularly for deep roots, and how we quantify their future dynamics. One challenge to global root quantification is the lack of correspondence between potential, contemporary and future land cover classifications. These incongruencies sometimes result in estimated changes in regionally-specific rooting depths that contrast with current knowledge about anticipated vegetation transitions. In the current study, place-based literature provided invaluable constraints on rooting depths for many ecosystems, but rooting depths in many regions of Asia, Australia, and Africa remain understudied. A lack of data describing contemporary rooting depth distributions in northern regions and estimates of vegetative cover and associated rooting depths in the future also emerged as important knowledge gaps (see especially Figure 1c). Additionally, there is a great deal of uncertainty in estimates of the deepest roots worldwide (Shenk & Jackson, 2002). Indeed, many of the deepest roots have been observed incidentally, suggesting that we have not yet sampled roots to their fullest extent (Fan et al., 2017).

We suggest that CZ research combining empirical and modeling approaches could help focus future research efforts on these critical gaps. First, empirical studies clarifying the ways in which global rooting distributions are changing could help with the development of decadal- to centennial-scale responses of extant ecosystems to climate change. Specifically, the leveraging of on-going climate experiments (e.g., Caplan et al., 2019), naturally existing climatic gradients (e.g., Ziegler et al., 2017), and chronosequences (e.g., Billings et al., 2018) could demonstrate how rooting depths respond to global changes to temperature and precipitation, as well as reveal quantitative relationships between rooting depth distributions and their impacts on soil formation processes, especially at depth. Focusing these studies in regions with relatively less research will improve our understanding of root-induced processes at the global scale.

Additionally, empirical and modeling studies examining the biogeochemical consequences of rooting depth change are critical. More extensive work either directly measuring subsurface biogeochemical fluxes as they respond to changes in rooting depth distributions, or modeling of biogeochemical processes that project such fluxes, will be invaluable for generating input parameters representing subsurface biogeochemical fluxes in ESMs. Because terrestrial

vegetation exerts a fundamental global control on land-atmosphere exchanges of water, energy, C, and other elements, improved representation of rooting distributions in global land models such as the Community Land Model (Lawrence et al., 2019) is of critical importance. This is particularly true as more sophisticated aboveground and belowground vegetation and biogeochemical processes are incorporated into these models (e.g., Tang et al., 2013; Fisher et al., 2017; Kennedy et al., 2019). With improved fidelity to biophysical and biogeochemical processes comes the corresponding opportunity to explore the potential consequences of changes in global rooting depths on land-atmosphere exchanges of water, energy, and C, and the large-scale ramifications that changes in rooting depths have for climate. Well-designed numerical experiments could elucidate the relative impacts of exogenous (e.g., agricultural conversion, woody encroachment) versus endogenous (e.g., water and nutrient limitation) drivers of changes in rooting depths on terrestrial cycling of water, energy, and C. These modeling efforts can feedback into empirical studies by illuminating regions where rooting depth knowledge is not sufficient and by pointing toward parameters requiring more explicit definition to improve future predictions. Such integrative studies would strengthen the nascent interactions between ESM and CZ communities to address pressing questions about global change that cannot be solved without substantial input from both disciplines (National Academy of Sciences, Engineering and Medicine, 2020). The improved representation of changing rooting depth distributions can link these research communities, representing a critical collaboration for understanding current and future functioning of Earth's CZ and climate.

5 Conclusion

Losses of relatively deep roots suggest an overlooked and subtle mechanism by which humans alter soil and ecosystem development. It is well established that humans accelerate losses of surface soil via erosion, which can result in a thinning of Earth's skin of soil (Wilkinson and McElroy, 2007). In contrast, altered rooting depths deep in soil profiles and associated shifts in rooted volume due to anthropogenic land use and climate change suggest a means by which human actions may govern soil thickness near the bottom of soil profiles. These shifts in rooting distributions support the idea that signals of the Anthropocene penetrate deeply into the subsurface even in naturally-occurring elemental cycles (Billings et al., 2018). Indications of widespread human transformation of land cover across millennia (Edgeworth et al., 2015) imply

that reductions in deep root abundances have been underway in multiple regions for a similar length of time. Though improving process representation in land models continues apace (Fisher and Koven, 2020), the representation of rooting depth distributions remains largely a static function of only PFT (cf. Drewniak, 2019). We present an opportunity to advance a dynamic representation of roots in land models by better constraining how rooting depth distributions vary with global change, as well as by identifying specific ecological processes particularly suited to better quantifying the dynamics of rooting, both past and future (e.g., regions of woody encroachment). Co-designed modeling, field and lab studies are needed to help clarify the consequences of rooting depth changes for contemporary and future CZ development. Such studies can elucidate the ways in which surficial anthropogenic activities radiate deep within Earth's subsurface, altering the developmental pace and character of Earth's CZ.

Acknowledgments

We thank Drs. Jorge Soberón, A. Townsend Peterson, Daniel Richter and Pauly Cartwright for offering helpful insights. We extend appreciation to Dr. Dan Markewitz for sharing his thoughts about root extension into the subsoil, and to multiple anonymous reviewers for their comments. Funding for this work was provided by NSF 1331846 and NSF 1841614. This material is also based upon work supported by the National Science Foundation under Award No. OIA-1656006 and matching support from the State of Kansas through the Kansas Board of Regents.

Data Availability and Code Availability

The original GLC2000 dataset modified for this analysis can be accessed at <https://forobs.jrc.ec.europa.eu/products/glc2000/products.php>. The unmodified potential vegetation data can be found at https://daac.ornl.gov/cgi-bin/dsviewer.pl?ds_id=961. All future land use projections can be accessed through the Landuse Harmonization data portal at <http://luh.umd.edu/data.shtml>. Rasters modified as described in Methods for contemporary and potential land cover, along with root depth assignment .csv files and code are available on

Zenodo (<https://doi.org/10.5281/zenodo.6522673>).

Author Contributions

SAB and EMH conceived of the idea with input from PLS. Analyses were developed and implemented by EMH and SAB. The manuscript was written by EMH and SAB with input from PLS, ANF and DH.

References

- Andrade, E., Rosa, G. Q., Almeida, A. M., Silva, A. G. R. D., & Sena, M. G. T. (2020). Rainfall regime on fine root growth in a seasonally dry tropical forest. *Revista Caatinga*, 33, 458-469.
- Austin, J.J.C., Perry, A., Richter, D. D. & Schroeder, P. A. (2018). Modifications of 2:1 clay minerals in a kaolinite-dominated Ultisol under changing landuse regimes. *Clays and Clay Minerals*, 66, 61-73. doi: 10.1346/CCMN.2017.064085
- Bartolome, E. & Belward, A.S. (2005). GLC2000: A new approach to global land cover mapping from Earth observation data. *International Journal of Remote Sensing*, 26, 1959-1977. doi: 10.1080/01431160412331291297
- Berner, R. A. (1998). The carbon cycle and carbon dioxide over Phanerozoic time: the role of land plants. *Philosophical Transactions of the Royal Society of London. Series B: Biological Sciences*, 353, 75-82.
- Berner, E. K., Berner, R. A., & Moulton, K. L. (2003). Plants and mineral weathering: present and past. *TrGeo*, 5, 605. doi: 10.1098/rstb.1998.0192
- Billings, S.A., Hirmas, D., Sullivan, P. L., Lehmeier, C. A., Bagchi, S., Min, K., Brecheisen, Z., Hauser, E., Stair, R., Flournoy, R. & Richter, D.D. (2018). Loss of deep roots limits agents of soil development that are only partially restored by decades of forest regeneration. *Elementa Science of the Anthropocene*, 6, 34. doi:10.1525/elementa.287
- Billings, W. D., Peterson, K. M., Shaver, G. R., & Trent, A. W. (1977). Root growth, respiration, and carbon dioxide evolution in an arctic tundra soil. *Arctic and Alpine Research*, 9, 129-137.
- Boike, J., Juszak, I., Lange, S., Chadburn, S., Burke, E., Paul Overduin, P., ... & Gouttevin, I. (2018). A 20-year record (1998–2017) of permafrost, active layer and meteorological conditions at a high Arctic permafrost research site (Bayelva, Spitsbergen). *Earth System Science Data*, 10, 355-390. doi: 10.5194/essd-10-355-2018

- Blume-Werry, G., Kreyling, J., Laudon, H., & Milbau, A. (2016). Short-term climate change manipulation effects do not scale up to long-term legacies: Effects of an absent snow cover on boreal forest plants. *Journal of Ecology*, 104, 1638-1648.
- Blume-Werry, G., Milbau, A., Teuber, L. M., Johansson, M., & Dorrepaal, E. (2019). Dwelling in the deep—strongly increased root growth and rooting depth enhance plant interactions with thawing permafrost soil. *New Phytologist*, 223, 1328-1339.
- Brandt, J.P., Flannigan, M.D., Maynard, D.G., Thompson, I.D. & Volney, W.J.A. (2013). An introduction to Canada's boreal zone: ecosystem processes, health, sustainability and environmental issues. *Environmental Reviews*, 21. doi:10. 1139/er-2013-0040..
- Brantley, S. L., Eissenstat, D. M., Marshall, J. A., Godsey, S. E., Balogh-Brunstad, Z., Karwan, D. L., Papuga, S. A., Roering, J., Dawson, T. E., Evaristo, J., Chadwick, O., McDonnell, J. J., & Weathers, K. C. (2017). Reviews and syntheses: On the roles trees play in building and plumbing the Critical Zone. *Biogeosciences*, 14, 5115-5142. doi: [10.5194/bg-14-5115-2017](https://doi.org/10.5194/bg-14-5115-2017)
- Brantley, S. L., Lebedeva, M., & Hausrath, E. M. (2012). A geobiological view of weathering and erosion. *Fundamentals of Geobiology*, 205-227. doi: 10.1002/9781118280874.ch12
- Canadell, J., Jackson, R. B., Ehleringer, J. B., Mooney, H. A., Sala, O. E. & Schulze E-D. (1996). Maximum rooting depth of vegetation types at the global scale. *Oecologia*, 108, 583-595. doi: 10.1007/BF00329030
- Caplan, J.S., Gimenez, D., Hirmas, D.R., Brunsell, N.A., Blair, J.M. & Knapp, A.K. (2019). Decadal-scale shifts in soil hydraulic properties as induced by altered precipitation. *Sci Adv.* 5 :eaau6635. doi:10.1126/sciadv.aau6635.
- Cramer, W., Bondeau, A., Woodward, I., Prentice, I., Betts, R., Brovkin, V., Cox, P., Fisher, V., Foley, J., Friend, A., Kucharik, C., Lomas, M., Ramankutty, N., Sitch, S., Smith, B., White, A. & Young-Molling, C.(2001). Global response of terrestrial ecosystem structure and function to CO2 and climate change: results from six dynamic global vegetation models. *Global Change Biology*, 7, 357-373. doi:10.1046/j.13652486.2001.00383.x
- Day, S. D., Wiseman, P. E., Dickinson, S. B., & Harris, J. R. (2010). Contemporary concepts of root system architecture of urban trees. *Arboriculture & Urban Forestry*, 36, 149-159.
- Devine, A.P., McDonald, R.A., Quaife, T. & Maclean, I.M.D. (2017). Determinants of woody Encroachment and cover in African savannas. *Oecologia*, 183, 939-951.
- Dontsova, K., Balogh-Brunstad, Z. & Chorover, J. (2020). Plants as Drivers of Rock Weathering. In *Biogeochemical Cycles* (eds K. Dontsova, Z. Balogh-Brunstad and G. Le Roux). doi:10.1002/9781119413332.ch2

- Drewniak, B. A. (2019). Simulating Dynamic Roots in the Energy Exascale Earth System Land Model. *Journal of Advances in Modeling Earth Systems*, 11, 338–359. doi: 10.1029/2018MS001334
- DuPont, S. T., Culman, S. W., Ferris, H., Buckley, D. H., & Glover, J. D. (2010). No-tillage conversion of harvested perennial grassland to annual cropland reduces root biomass, decreases active carbon stocks, and impacts soil biota. *Agriculture, ecosystems & environment*, 137, 25–32. doi: 10.1016/j.agee.2009.12.021
- DuPont, S. T., Beniston, J., Glover, J. D., Hodson, A., Culman, S. W., Lal, R., & Ferris, H. (2014). Root traits and soil properties in harvested perennial grassland, annual wheat, and never-tilled annual wheat. *Plant and Soil*, 38, 405–420.
- Edgeworth, M., Richter, D. D., Waters, C., Haff, P., Neal, C. & Price, S. J. (2015). Diachronous beginnings of the Anthropocene: The lower bounding surface of anthropogenic deposits. *The Anthropocene Review*, 2, 33–58. doi: 10.1177/2053019614565394
- Ellis, E.C., Goldewijk, K. K. , Siebert, S., Lightman, D. & Ramankutty, N. (2010). Anthropogenic transformation of the biomes, 1700 to 2000. *Global Ecology and Biogeography*, 19, 589–606. doi: 10.1111/j.1466-8238.2010.00540.x
- Fan, Y., Miguez-Macho G., Jobbágy, E. G., Jackson, R. B. & Otero-Casal, C. (2017). Hydrologic regulation of plant rooting depth. *Proceedings of the National Academy of Science USA*, 114, 10572–10577. doi: 10.1073/pnas.1712381114
- Fisher, R. A., & Koven, C. D. (2020). Perspectives on the Future of Land Surface Models and the Challenges of Representing Complex Terrestrial Systems. *Journal of Advances in Modeling Earth Systems*, 12, 1–24. doi: [10.1029/2018MS001453](https://doi.org/10.1029/2018MS001453)
- Fisher, R. A., Koven, C. D., Anderegg, W. R. L., Christoffersen, B. O., Dietze, M. C., Farrior, C. E., Holm, J. A., Hurtt, G. C., Knox, R. G., Lawrence, P. J., Lichstein, J. W., Longo, M., Matheny, A. M., Medvigy, D., Muller-Landau, H. C., Powell, T. L., Serbin, S. P., Sato, H., Shuman, J. K., ... Moorcroft, P. R. (2017). Vegetation demographics in Earth System Models: A review of progress and priorities. *Global Change Biology*, 24, 35–54. doi: 10.1111/gcb.13910
- Groffman, P. M., Driscoll, C. T., Fahey, T. J., Hardy, J. P., Fitzhugh, R. D., & Tierney, G. L. (2001). Colder soils in a warmer world: a snow manipulation study in a northern hardwood forest ecosystem. *Biogeochemistry*, 56, 135–150.
- Gupta, A.K. (2004). Origin of agriculture and domestication of plants and animals linked to early Holocene climate amelioration. *Current Science*, 87, 54–59.
- Harsch, M. A., Hulme, P. E., McGlone, M. S., & Duncan, R. P. (2009). Are treelines advancing? A global meta-analysis of treeline response to climate warming. *Ecology Letters*, 12, 1040–1049. doi: 10.1111/j.1461-0248.2009.01355.x
- Hasenmueller, E.A., Gu, X., Weitzman, J. N., Adams, T. S., Stinchcomb, G. E., Eissenstat, D. M., Drohan, P. J., Brantley, S. L., Kaye, J. P. (2017). Weathering of rock to regolith: The

- activity of deep roots in bedrock fractures. *Geoderma*, 300, 11-31.
doi: 10.1016/j.geoderma.2017.03.020
- Haxeltine, A., & Prentice, I. C. (1996). BIOME3: An equilibrium terrestrial biosphere model based on ecophysiological constraints, resource availability, and competition among plant functional types. *Global Biogeochemical Cycles*, 10, 693-709. doi: 10.1029/96GB02344
- Hertel, D., Harteveld, M. A., & Leuschner, C. (2009). Conversion of a tropical forest into agroforest alters the fine root-related carbon flux to the soil. *Soil Biology and Biochemistry*, 41, 481-490. doi: 10.1016/j.soilbio.2008.11.020
- Hijmans, R. J., van Etten, J., Sumner, M., Cheng, J., Baston, D., Bevan, A., et al. (2019). "Geographic Data Analysis and Modeling," (CRAN, v. 3.0-7, 2019; <https://cran.r-project.org/web/packages/raster/raster.pdf>).
- Hurt, G. C., Chini, L. P., Frolking, S., Betts, R. A., Feddema, J., Fischer, G., ... & Jones, C. D. (2011). Harmonization of land-use scenarios for the period 1500–2100: 600 years of global gridded annual land-use transitions, wood harvest, and resulting secondary lands. *Climatic Change*, 109, 117. doi: 10.1007/s10584-011-0153-2
- Hurt, G. C., Chini, L., Sahajpal, R., Frolking, S., Bodirsky, B. L., Calvin, K., Doelman, J. C., Fisk, J., Fujimori, S., Goldewijk, K. K., Hasegawa, T., Havlik, P., Heinemann, A., Humpenöder, F., Jungclaus, J., Kaplan, J., Kennedy, J., Kristzin, T., Lawrence, D., Lawrence, P., Ma, L., Mertz, O., Pongratz, J., Popp, A., Poulter, B., Riahi, K., Shevliakova, E., Stehfest, E., Thornton, P., Tubiello, F. N., van Vuuren, D. P., and Zhang, X. (2020). Harmonization of Global Land-Use Change and Management for the Period 850–2100 (LUH2) for CMIP6. *Geosci. Model Dev. Discuss.*, 13, 5425-5464. <https://doi.org/10.5194/gmd-2019-360>.
- Iversen, C.M. (2010). Digging deeper: Fine-root responses to rising atmospheric CO₂ concentration in forested ecosystems. *New Phytologist*, 186, 346-357.
doi: 10.1111/j.1469-8137.2009.03122.x
- Iversen, C. M., McCormack, M. L., Powell, A. S., Blackwood, C. B., Freschet, G. T., Kattge, J., ... & van Bodegom, P. M. (2017). A global fine-root ecology database to address below-ground challenges in plant ecology. *New Phytologist*, 215, 15-26.
- Iversen, C.M., McCormack, M. L., Baer, J.K., Powell, A.S., Chen, W., Collins, C., ... & Zadworny, M. (2021). Fine-Root Ecology Database (FRED): A global collection of root trait data with coincident site, vegetation, edaphic, and climatic data, version 3. Oak Ridge National Laboratory, TES SFA, U.S. Department of Energy, Oak Ridge, Tennessee, U.S.A. Access on-line at: <https://doi.org/10.25581/ornlsfa.014/1459186>.
- Jackson, R.B., Canadell, J., Ehleringer, J. R., Mooney, H. A., Sala, O.E. & Schulze, E. D. (1996). A global analysis of root distributions for terrestrial biomes. *Oecologia*, 108, 389-411.
doi: 10.1007/BF00333714
- Jaramillo, V. J., Ahedo-Hernández, R., & Kauffman, J. B. (2003). Root biomass and carbon in a tropical evergreen forest of Mexico: changes with secondary succession and forest conversion to pasture. *Journal of Tropical Ecology*, 19, 457-464. doi: 10.1017/S0266467403003493

- Jobbagy, E. G. & Jackson, R. B. (2001) The distribution of soil nutrients with depth: Global patterns and the imprint of plants. *Biogeochemistry*, 53, 51-77. doi: 10.1023/A:1010760720215
- Jordan, T. Ashley G. M., Barton, M. D., Burges, S. J., Farley, K. A., Freeman, K. H., Jeanloz, R., Marshall, C. R., Orcutt, J.A., Richter, F.M. et al. (Eds.). (2001). *Basic research opportunities in Earth Science*. Washington D.C.: National Academy Press.
- Kennedy, D., Swenson, S., Oleson, K. W., Lawrence, D. M., Fisher, R., Lola da Costa, A. C., & Gentine, P. (2019). Implementing Plant Hydraulics in the Community Land Model, Version 5. *Journal of Advances in Modeling Earth Systems*, 11, 485–513. doi: [10.1029/2018MS001500](https://doi.org/10.1029/2018MS001500)
- Keuper, F., Wild, B., Kumm, M., Beer, C., Blume-Werry, G., Fontaine, S., ... & Dorrepaal, E. (2020). Carbon loss from northern circumpolar permafrost soils amplified by rhizosphere priming. *Nature Geoscience*, 13, 560-565.
- Kleidon, A. (2003). Global datasets of rooting zone depth inferred from inverse methods. *Journal of Climate*, 17, 2741-2722. doi: 10.1175/1520-0442(2004)017<2714:GDORZD>2.0.CO;2
- Kleidon, A. & Heimann M. (1998). A method of determining rooting depth from a terrestrial biosphere model and its impacts on the global water and carbon cycle. *Global Change Biology*, 4, 275-286. doi: 10.1046/j.1365-2486.1998.00152.x
- Krasowski, M. J., Lavigne, M. B., Szuter, M. A., Olesinski, J., Kershaw, J. A., & McGarrigle, E. (2018). Age-related changes in survival and turnover rates of balsam fir (*Abies balsamea* (L.) Mill.) fine roots. *Tree physiology*, 38, 865-876.
- Lal, R. (2001). Potential of desertification control to sequester carbon and mitigate the greenhouse effect. *Climatic change*, 51, 35-72.
- Lander, J.P. (2013). *R for everyone: Advanced analytics and graphics*. Addison-Wesley, Boston, MA.
- Lawrence, D. M., Fisher, R. A., Koven, C. D., Oleson, K. W., Swenson, S. C., Bonan, G., Collier, N., Ghimire, B., Kampenhout, L., Kennedy, D., Kluzek, E., Lawrence, P. J., Li, F., Li, H., Lombardozzi, D., Riley, W. J., Sacks, W. J., Shi, M., Vertenstein, M., ... Zeng, X. (2019). The Community Land Model Version 5: Description of New Features, Benchmarking, and Impact of Forcing Uncertainty. *Journal of Advances in Modeling Earth Systems*, 11, 4245–4287. doi: 10.1029/2018MS001583

- Lawrence, D. M., & Slater, A. G. (2005). A projection of severe near-surface permafrost degradation during the 21st century. *Geophysical Research Letters*, 32, L24401. doi:10.1029/2005GL025080.
- Lindsey, P., & Bassuk, N. (1992). Redesigning the urban forest from the ground below: A new approach to specifying adequate soil volumes for street trees. *Arboricultural Journal*, 16, 25-39.
- Lozanova, L., Zhiyanski, M., Vanguelova, E., Doncheva, S., Marinov, M. P., & Lazarova, S. (2019). Dynamics and vertical distribution of roots in European Beech forests and Douglas Fir plantations in Bulgaria. *Forests*, 10, 1123.
- Maeght, J. L., Rewald, B. & Pierret, A. (2013). How to study deep roots and why it matters. *Frontiers in Plant Science*, 4, 299. doi: 10.3389/fpls.2013.00299
- Malhotra, A., Brice, D. J., Childs, J., Graham, J. D., Hobbie, E. A., Vander Stel, H., ...&Iversen, C. M. (2020). Peatland warming strongly increases fine-root growth. *Proceedings of the National Academy of Sciences*, 117, 17627-17634.
- McCormack, M. L., Crisfield, E., Raczka, B., Schnekenburger, F., Eissenstat, D. M., & Smithwick, E. A. (2015). Sensitivity of four ecological models to adjustments in fine root turnover rate. *Ecological modelling*, 297, 107-117.
- McCormack, M. L., Guo, D., Iversen, C. M., Chen, W., Eissenstat, D. M., Fernandez, C. W., ... & Reich, P. B. (2017). Building a better foundation: Improving root-trait measurements to understand and model plant and ecosystem processes. *New Phytologist*, 215, 27-37.
- Montagnoli, A., Terzaghi, M., Giussani, B., Scippa, G. S., & Chiatante, D. (2018). An integrated method for high-resolution definition of new diameter-based fine root sub-classes of *Fagus sylvatica* L. *Annals of forest science*, 75, 1-13.
- National Academy of Sciences, Engineering and Medicine. (2020). *A Vision for NSF Earth Sciences 2020-2030: Earth in Time*. Washington, DC: The National Academies Press. <https://doi.org/10.17226/25761>.
- NEON (National Ecological Observatory Network). Root biomass and chemistry, Megapit, RELEASE-2021 (DP1.10066.001). <https://doi.org/10.48443/s6k6-8h54>. Dataset accessed from <https://data.neonscience.org> on November 22, 2021
- Nepstad, D.C., Carvalho, C. R. d., Davidson, E. A., Jipp, P. H., Lefebvre, P. A., Negreiros, G. H., da Silva, E. D., Stone, T. A., Trumbore, S. E. & Vieira, S. (1994). The role of deep roots in the hydrological and carbon cycles of Amazonian forests. *Nature*, 372, 666-669. doi: 10.1038/372666a0
- Nippert, J.B. & Knapp, A.K. (2007). Linking water uptake with rooting patterns in grassland species. *Oecologia*, 153, 261-272. doi: 10.1007/s00442-007-0745-8
- Pawlik, L. (2013). The role of trees in the geomorphic system of forested hillslopes—A review.

Earth-Science Reviews, 126, 250-265. doi: 10.1016/j.earscirev.2013.08.007

- Pierret, A., Maeght, J. L., Clement, C., Montoroi, J. P., Hartman, C., & Gonkhamdee, S. (2016). Understanding deep roots and their functions in ecosystems: An advocacy for more unconventional research. *Annals of Botany*, 118, 621-625.
<https://doi.org/10.1093/aob/mcw130>
- Price, D.T., Alfaro, R.I., Brown, K.J., Flannigan, M.D., Fleming, R.A., Hogg, E.H., ... & Venier, L.A. (2013). Anticipating the consequences of climate change for Canada's boreal forest ecosystems. *Environmental Reviews*, 21, 322-326.
- Ramankutty, N. & Foley, J.A. (1999). Estimating historical changes in global land cover: Croplands from 1700 to 1992. *Global Biogeochemical Cycles*, 13, 997-1027.
[doi: 10.1029/1999GB900046](https://doi.org/10.1029/1999GB900046)
- Richter, D. D. & Markewitz, D. (1995). How deep is soil? *BioScience*, 45, 600-609. doi: 10.2307/1312764
- Riahi, K., Van Vuuren, D. P., Kriegler, E., Edmonds, J., O'Neill, B. C., Fujimori, S., ... & Lutz, W. (2017). The shared socioeconomic pathways and their energy, land use, and greenhouse gas emissions implications: an overview. *Global Environmental Change*, 42, 153-168.
- RStudio Team. (2017). "RStudio: Integrated development for R," (RStudio Inc, Boston, <http://www.rstudio.com/>).
- Schenk, J. H. (2007). The shallowest possible water extraction profile: A null model for global root distributions. *Vadose Zone Journal*, 7, 1119-1124.
doi: 10.2136/vzj2007.0119
- Schenk, H. J. & Jackson, R.B. (2002a). The global biogeography of roots. *Ecological Monographs*, 72, 311-328.
- Schenk, H. J. & Jackson R.B. (2002b). Rooting depths, lateral root spreads, and belowground/aboveground allometries of plants in water limited ecosystems. *Journal of Ecology*, 90, 480-494.
- Schenk, H. J. & Jackson, R.B. (2003). Global Distribution of Root Profiles in Terrestrial Ecosystems. Data set with documentation. Available from the Oak Ridge National

Laboratory Distributed Active Archive Center, Oak Ridge, Tennessee, U.S.A. On-line at
<http://www.daac.ornl.gov>.

Schenk, H. J., & Jackson, R.B.. (2005). Mapping the global distribution of deep roots in
 relation to climate and soil characteristics. *Geoderma*, 126, 129-140.

Schenk, H.J., & Jackson, R.B. 2009. ISLSCP II Ecosystem Rooting Depths. In Hall, Forrest G.,
 G. Collatz, B. Meeson, S. Los, E. Brown de Colstoun, and D. Landis (eds.). ISLSCP
 Initiative II Collection. Data set. Available on-line [<http://daac.ornl.gov/>] from Oak Ridge
 National Laboratory Distributed Active Archive Center, Oak Ridge, Tennessee, U.S.A.
 doi:10.3334/ORNLDAAAC/929

Sistla, S. A., Moore, J. C., Simpson, R. T., Gough, L., Shaver, G. R., & Schimel, J. P.
 (2013). Long-term warming restructures Arctic tundra without changing net soil
 carbon storage. *Nature*, 497, 615-618.

Stevens, N., Lehmann, C. E. R., Murphy, B. P. & Durigan, G. (2017). Savanna woody
 encroachment is widespread across three continents. *Global Change Biology*, 23, 235-
 244. doi: 10.1111/gcb.13409

Stone, E. L., & Kalisz, P. J. (1991). On the maximum extent of tree roots. *Forest ecology and
 management*, 46, 59-102.

Sullivan, P. L., Billings, S. A., Hirmas, D., Li, L., Zhang, X., Ziegler, S., ... & Wen, H. (2022).
 Embracing the dynamic nature of soil structure: A paradigm illuminating the role of life
 in critical zones of the Anthropocene. *Earth-Science Reviews*, 225, 103873.

Swank, W. T. (1986). Biological control of solute losses from forest ecosystems (Vol. 85). John
 Wiley & Sons, New York.

Tang, J. Y., Riley, W. J., Koven, C. D., & Subin, Z. M. (2013). CLM4-BeTR, a generic
 biogeochemical transport and reaction module for CLM4: Model development,
 evaluation, and application. *Geosci. Model Dev.*, 6, 127–140. doi: 10.5194/gmd-6-127-
 2013

Wang, J. A., Sulla-Menashe, D., Woodcock, C. E., Sonnentag, O., Keeling, R. F., & Friedl, M.
 A. (2020). Extensive land cover change across Arctic–Boreal Northwestern North
 America from disturbance and climate forcing. *Global Change Biology*, 26, 807-822. doi:
[10.1111/gcb.14804](https://doi.org/10.1111/gcb.14804)

Whitford, W. G., & Duval, B. D. (2019). *Ecology of desert systems*. Academic Press.

Wilkinson, B. H., & McElroy, B. J. (2007). The impact of humans on continental erosion and
 sedimentation. *Geological Society of America Bulletin*, 119, 140-156. doi:
 10.1130/B25899.1

- Zhang, T., Barry, R. G., Knowles, K., Heginbottom, J. A., & Brown, J. (2008). Statistics and characteristics of permafrost and ground-ice distribution in the Northern Hemisphere. *Polar Geography*, 31, 47-68. doi: 10.1080/10889370802175895
- Zhang, Y., Niu, J., Yu, X., & Zhu, W. (2015). Effects of fine root length density and root biomass on soil preferential flow in forest ecosystems. *Forest Systems*, 24, e012. doi: [10.5424/fs/2015241-06048](https://doi.org/10.5424/fs/2015241-06048)
- Zhao, H. L., Zhao, X. Y., Zhou, R. L., Zhang, T. H., & Drake, S. (2005a). Desertification processes due to heavy grazing in sandy rangeland, Inner Mongolia. *Journal of arid environments*, 62, 309-319.
- Zhao, M., Heinsch, F. A., Nemani, R. R., & Running, S. W. (2005b). Improvements of the MODIS terrestrial gross and net primary production global data set. *Remote sensing of Environment*, 95, 164-176. doi: 10.1016/j.rse.2004.12.011
- Zeng, X. (2001). Global vegetation root distribution for land modeling. *American Meteorological Society*, 2, 525-530. doi: 10.1175/1525-7541(2001)002<0525:GVRDFL>2.0.CO;2
- Ziegler, S.E., Benner, R., Billings, S.A., Edwards, K.A., Philben, M., Zhu, X. & Laganière, J. (2017). Climate warming can accelerate carbon fluxes without changing soil carbon stocks. *Frontiers in Earth Science*, 5, 2. doi:10.3389/feart.2017.00002

Global-scale shifts in rooting depths in the Anthropocene present unexamined consequences in critical zone functioning

Emma Hauser^{1,2*}, Pamela L. Sullivan³, Alejandro Flores⁴, Daniel Hirmas⁵, Sharon A. Billings^{1,2*}

¹Department of Ecology and Evolutionary Biology, The University of Kansas, Lawrence, KS, USA.

²The Kansas Biological Survey, Lawrence, KS, USA.

³College of Earth, Ocean, and Atmospheric Science, Oregon State University, Corvallis, OR, USA.

⁴Department of Geosciences, Boise State University, Boise, ID, USA

⁵Department of Environmental Science, University of California, Riverside, CA

Contents of this file

Figures S1 to S6
Table S1

Introduction

Here we include additional figures and a data table, generated via the same methods detailed in the main text. The figures represent an extension of the figures presented in text and the table represents alternate ways of calculating rooting depth metrics using different representations of rooting depths for desert biomes.




Figure S1. Comparison of potential (A) and contemporary (B) distributions of depth to 95% rooting biomass (D95) across the globe. Inset histograms display the distribution of rooting depths in each map, with dashed lines marking the means of the two datasets (blue histogram represents potential vegetation, red contemporary). Appearance of a distinct color change from dark blue to light grey in Asia and Canada at 60°N is an artifact of restricted maximum rooting depth assignments at northern latitudes used in our calculations to account for growth limitations imposed by frozen soils. Appearance of a distinct color change from dark blue to light blue and grey in Asia and Canada at 50°N reflects reassignment of mixed forests to the boreal forest class above this latitude (Brandt et al., 2013; Price et al., 2013).




Figure S2. Comparison of potential (A) and contemporary (B) distributions of depth to 50% rooting biomass (D50) across the globe. Inset histograms display the distribution of rooting depths in each map, with dashed lines marking the means of the two datasets (blue histogram represents potential vegetation, red contemporary). The appearance of a distinct color change from blue to light grey in Asia and Canada at 60°N is an artifact of restricted maximum rooting depth assignments at northern latitudes used in our calculations to account for growth limitations imposed by frozen soils. Appearance of a distinct color change from dark blue to light blue and grey in Asia and Canada at 50°N

reflects reassignment of mixed forests to the boreal forest class above this latitude (Brandt et al., 2013; Price et al., 2013).




Figure S3. Change in depth to 95% rooting biomass (D95) due to differences in potential vegetation distributions compared to contemporary vegetation distributions. Red regions denote shallower roots in contemporary systems, while blue regions denote deeper roots in contemporary systems when compared to their potential vegetation distributions. Appearance of a distinct color change from blue to light grey in Asia and Canada at 50°N reflects reassignment of mixed forests to the boreal forest class above this latitude (Brandt et al., 2013; Price et al., 2013).




Figure S4. Change in depth to 99% rooting biomass (D99) due to differences between contemporary rooting depth distributions and anticipated rooting distributions under two projected SSP RCP scenarios for the year 2100; SSP2 RCP4.5 (a) and SSP4 RCP6.0 (b). Grey and red colors indicate root depth truncation and blue indicates elongation.




Figure S5. Change in depth to 50% rooting biomass (D50) due to differences between contemporary rooting depth distributions and anticipated rooting distributions under two projected SSP RCP scenarios for the year 2100 (results are similar for both scenarios); SSP1 RCP2.6 (a) and SSP5 RCP8.5 (b). These two maps represent scenarios of greatest projected change and least projected change. Grey and red colors indicate root depth truncation and blue indicates elongation.




Figure S6. Change in depth to 50% rooting biomass (D50) due to differences in potential vegetation distributions compared to contemporary vegetation distributions. Red regions denote shallower roots in contemporary systems, while blue regions denote deeper roots in contemporary systems when compared to their potential vegetation distributions. Appearance of a distinct color change from dark blue to light grey in Asia and Canada at 50°N reflects reassignment of mixed forests to the boreal forest class above this latitude (Brandt et al., 2013; Price et al., 2013)

Table S1. Mean global rooting depth metrics with 95% confidence intervals for potential and contemporary land cover distributions under two scenarios of user assumptions. The third column displays the difference in cm between potential and contemporary root distributions and the percent change in parentheses. The first three rows indicate global means excluding true desert regions. The second three rows include true deserts in calculations of global mean rooting depth metrics but set roots in those systems to a depth of zero.

Metric	Potential Mean Rooting Depth (m, 95% CI)	Contemporary Mean Rooting Depth (m, 95% CI)	Change From potential (cm)
D99 (Desert excluded)	1.50 (+/- 0.001)	1.41 (+/- 0.0001)	-8.15 (5.4%)
D95(Desert excluded)	0.88 (+/- 0.0006)	0.82(+/- 0.00006)	-4.93 (5.6%)
D50 (Desert excluded)	0.07 (+/- 0.00007)	0.057 (+/- 0.000008)	-1.45 (20.7%)
D99 (Desert roots set to 0m)	1.35 (+/- 0.0011)	1.22 (+/- 0.0001)	-14.7 (10.9%)
D95(Desert roots set to 0m)	0.79 (+/- 0.0006)	0.72 (+/- 0.00007)	-11.1 (14.1%)
D50 (Desert roots set to 0m)	0.066 (+/- 0.00007)	0.05 (+/- 0.000007)	-1.7 (25.8%)