

# Diel to interannual variation in carbon dioxide emissions from lakes and reservoirs

Malgorzata Golub<sup>1</sup>, Nikaan Koupaie-Abyazani<sup>2</sup>, Timo Vesala<sup>3</sup>, Ivan Mammarella<sup>4</sup>, Anne Ojala<sup>5</sup>, Gil Bohrer<sup>6</sup>, Gesa A Weyhenmeyer<sup>7</sup>, Peter D. Blanken<sup>8</sup>, Werner Eugster<sup>9</sup>, Franziska Koebsch<sup>10</sup>, Jiquan Chen<sup>11</sup>, Kevin P. Czajkowski<sup>12</sup>, Chandrashekhar Deshmukh<sup>13</sup>, Frédéric Guérin<sup>14</sup>, Jouni Heiskanen<sup>4</sup>, Elyn Humphreys<sup>15</sup>, Anders Jonsson<sup>16</sup>, Jan Karlsson<sup>17</sup>, George W. Kling<sup>18</sup>, Xuhui Lee<sup>19</sup>, Heping Liu<sup>20</sup>, Annalea Lohila<sup>21</sup>, Erik Johannes Lundin<sup>22</sup>, Timothy Hector Morin<sup>23</sup>, Eva Podgrajsek<sup>24</sup>, Maria Provenzale<sup>4</sup>, Anna Rutgersen<sup>25</sup>, Torsten Sachs<sup>26</sup>, Erik Sahlée<sup>27</sup>, Dominique Serça<sup>28</sup>, Changliang Shao<sup>29</sup>, Christopher Spence<sup>30</sup>, Ian B. Strachan<sup>31</sup>, Wei Xiao<sup>32</sup>, and Ankur Rashmikan Desai<sup>2</sup>

<sup>1</sup>Dundalk Institute of Technology

<sup>2</sup>University of Wisconsin-Madison

<sup>3</sup>University of Helsinki, Institute for Atmospheric and Earth System Research

<sup>4</sup>University of Helsinki

<sup>5</sup>Natural Resources Institute

<sup>6</sup>Ohio State University

<sup>7</sup>Ecology and Genetics/Limnology

<sup>8</sup>University of Colorado Boulder

<sup>9</sup>ETH Zurich

<sup>10</sup>GFZ German Research Centre for Geosciences

<sup>11</sup>Michigan State University

<sup>12</sup>University of Toledo

<sup>13</sup>APRIL Asia

<sup>14</sup>IRD - Marseille, France.

<sup>15</sup>Carleton University

<sup>16</sup>Department of Ecology and Environmental Science

<sup>17</sup>Umea University

<sup>18</sup>University of Michigan-Ann Arbor

<sup>19</sup>Yale University, School of Forestry and Environmental Studies

<sup>20</sup>Washington State University

<sup>21</sup>Finnish Meteorological Institute

<sup>22</sup>Swedish Polar Research Secretariat

<sup>23</sup>State University of New York College of Environmental Science and Forestry

<sup>24</sup>OX2

<sup>25</sup>Uppsala University

<sup>26</sup>Helmholtz Centre Potsdam - German Research Centre for Geosciences (GFZ)

<sup>27</sup>Earth Sciences

<sup>28</sup>Laboratoire d'Aérodologie, Université de Toulouse, CNRS, UPS, France

<sup>29</sup>Institute of Agricultural Resources and Regional Planning, Chinese Academy of

Agricultural Sciences

<sup>30</sup>Environment and Climate Change Canada

<sup>31</sup>McGill University

<sup>32</sup>Nanjing University of Information Science and Technology


November 23, 2022



## Abstract

Accounting for temporal changes in carbon dioxide (CO<sub>2</sub>) emissions from freshwaters remains a challenge for global and regional carbon budgets. Here, we synthesize 171 site-months of eddy covariance flux measurements of CO<sub>2</sub> from 13 lakes and reservoirs in the Northern Hemisphere (NH) and quantify dynamics at multiple temporal scales. We found pronounced sub-annual variability in CO<sub>2</sub> flux at all sites. Accounting for diel variation, only 11% of site-months were net daily sinks of CO<sub>2</sub>. Annual CO<sub>2</sub> emissions had an average of 25% (range 3-58%) interannual variation. Nighttime emissions regularly exceeded daytime emissions. Sources of CO<sub>2</sub> flux variability were delineated through mutual information analysis. Sample analysis of CO<sub>2</sub> fluxes indicate importance of continuous sampling. Constraining short- and long-term variability is necessary to improve detection of temporal changes of CO<sub>2</sub> fluxes in response to natural and anthropogenic drivers.

## Hosted file

golub\_20220626\_supplement.docx available at <https://authorea.com/users/544248/articles/601614-diel-to-interannual-variation-in-carbon-dioxide-emissions-from-lakes-and-reservoirs>





# 1 Diel to interannual variation in carbon

## 2 dioxide emissions from lakes and reservoirs

Name	Email	Affiliation-1
Malgorzata Golub	malgorzata.golub@dkit.ie	Dundalk Institute of Technology, Centre for Freshwater and Environmental Studies, Dundalk, Ireland
Nikaan Koupaei-Abyazani	koupaeiabyaz@wisc.edu	University of Wisconsin-Madison, Dept. of Atmospheric and Oceanic Sciences, Madison, WI, USA
Timo Vesala	timo.vesala@helsinki.fi	University of Helsinki, Institute for Atmospheric and Earth System Research (INAR)/Physics, Faculty of Science, Finland
Ivan Mammarella	ivan.mammarella@helsinki.fi	University of Helsinki, Institute for Atmospheric and Earth System Research (INAR)/Physics, Faculty of Science, Finland
Anne Ojala	anne.ojala@luke.fi	Natural Resources Institute Finland, Helsinki
Gil Bohrer	bohrer.17@osu.edu	The Ohio State University, Dept. of Civil, Environmental and Geodetic Engineering, Columbus, OH, USA
Gesa A. Weyhenmeyer	Gesa.Weyhenmeyer@ebc.uu.se	Uppsala University, Dept. of Ecology and Genetics/Limnology, Uppsala, Sweden
Peter D. Blanken	blanken@colorado.edu	University of Colorado, Dept. of Geography, Boulder, CO, USA
Werner Eugster	werner.eugster@usys.ethz.ch	ETH Zürich, Zürich, Switzerland
Franziska Koebsch	Franziska.Koebsch@uni-goettingen.de	GFZ German Research Centre for Geosciences, Potsdam, Germany
Jiquan Chen	jqchen@msu.edu	Michigan State University, Department of Geography, Environment and Spatial Science, East Lansing, MI; USA
Kevin Czajkowski	kevin.czajkowski@utoledo.edu	University of Toledo, Dept. of Geography and Planning, Toledo, OH, USA
Chandrashekhhar Deshmukh	csd.cae@gmail.com	APRIL Asia, Laboratoire d'Aérodynamique, Observatoire Midi-Pyrénées, Toulouse, France
Frederic Guérin	frederic.guerin@ird.fr	IRD - Marseille, Toulouse, France.
Jouni Heiskanen	jouni.heiskanen@helsinki.fi	University of Helsinki, Faculty of Biological and Environmental Sciences, Helsinki, Finland
Elyn Humphreys	elyn.humphreys@carleton.ca	Carleton University, Geography and Environmental Studies, Ottawa, Canada
Anders Jonsson	anders.jonsson@emg.umu.se	Umeå University, Dept. of Ecology and Environmental Science, Umeå, Sweden

Jan Karlsson	jan.p.karlsson@emg.umu.se	Umeå University, Climate Impacts Research Centre (CIRC), Dept. of Ecology and Environmental Science, Umeå, Sweden
George Kling	gwk@umich.edu	University of Michigan, Dept. of Ecology and Evolutionary Biology, Ann Arbor, MI, USA
Xuhui Lee	xuhui.lee@yale.edu	Yale University, School of the Environment, New Haven, CT, USA
Heping Liu	heping.liu@wsu.edu	Washington State University, Dept. of Civil and Environmental Engineering, Pullman, WA, USA
Annalea Lohila	annalea.lohila@fmi.fi	Finnish Meteorological Institute, Climate System Research, Helsinki, Finland
Erik Lundin	erik.lundin@polar.se	Swedish Polar Research Secretariat, Abisko Scientific Research Station, Abisko, Sweden
Tim Morin	thmorin@esf.edu	State University of New York, College of Environmental Science and Forestry, Syracuse, New York, USA
Eva Podgrajsek	eva.podgrajsek@ox2.com	OX2, Stockholm
Maria Provenzale	mariaprovenzale@icloud.com	University of Helsinki, Institute for Atmospheric and Earth System Research (INAR)/Dept. Physics, Helsinki, Finland
Anna Rutgersson	Anna.Rutgersson@met.uu.se	Uppsala University, Dept. of Earth Sciences, Uppsala, Sweden
Torsten Sachs	torsten.sachs@gfz-potsdam.de	GFZ German Research Centre for Geosciences, Potsdam, Germany
Erik Sahleé	Erik.Sahlee@met.uu.se	Uppsala University, Dept. of Earth Sciences, Uppsala, Sweden
Dominique Serça	Dominique.Serca@aero.obs-mip.fr	Laboratoire d'Aérodynamique, Université de Toulouse, CNRS, UPS, France
Changliang Shao	shaochangliang@caas.cn	Institute of Agricultural Resources and Regional Planning, Chinese Academy of Agricultural Sciences, Beijing, China
Christopher Spence	chris.spence@canada.ca	Environment and Climate Change Canada, Saskatoon, SK, Canada
Ian B. Strachan	ian.strachan@mcgill.ca	McGill University, Dept. of Natural Resource Sciences, Ste. Anne de Bellevue, QC, Canada
Wei Xiao	wei.xiao@nuist.edu.cn	Nanjing University of Information Science and Technology, Yale-NUIST Center on Atmospheric Environment, Nanjing, Jiangsu, China
Ankur R. Desai	desai@aos.wisc.edu	University of Wisconsin-Madison, Dept. of Atmospheric and Oceanic Sciences, Madison, WI, USA

3

4

\*Corresponding author: Ankur Desai [desai@aos.wisc.edu](mailto:desai@aos.wisc.edu), +1-608-520-0305

## Abstract

Accounting for temporal changes in carbon dioxide (CO<sub>2</sub>) emissions from freshwaters remains a challenge for global and regional carbon budgets. Here, we synthesize 171 site-months of eddy covariance flux measurements of CO<sub>2</sub> from 13 lakes and reservoirs in the Northern Hemisphere (NH) and quantify dynamics at multiple temporal scales. We found pronounced sub-annual variability in CO<sub>2</sub> flux at all sites. Accounting for diel variation, only 11% of site-months were net daily sinks of CO<sub>2</sub>. Annual CO<sub>2</sub> emissions had an average of 25% (range 3-58%) interannual variation. Nighttime emissions regularly exceeded daytime emissions. Sources of CO<sub>2</sub> flux variability were delineated through mutual information analysis. Sample analysis of CO<sub>2</sub> fluxes indicate importance of continuous sampling. Constraining short- and long-term variability is necessary to improve detection of temporal changes of CO<sub>2</sub> fluxes in response to natural and anthropogenic drivers.

## Plain Language Summary

Lakes and reservoirs around the world are a major component of the global carbon cycle. Recent syntheses of measurements find their contributions to be on the order of 2-6% of total global fossil fuel emissions. However, these estimates are primarily derived from compilations with low frequency of sampling, from a few times a year up to weekly, often restricted to a single season, and with limited regard to year-to-year variations. Here, we conduct the first analysis of a globally distributed network of sub-hourly, multi-year lake and reservoir carbon dioxide emissions. These measurements were made using eddy-covariance flux towers, which continuously sample these emissions year-round. Across our 13 study sites, we found nighttime emissions regularly exceeding daytime emissions and persistent sub-monthly variability regardless of lake size or nutrient status. For sites with multiple years of data, we found an average 25% variation in estimated annual emissions depending on the year chosen. Together, these results point to a need for improved, systematic sub-weekly sampling of freshwater systems to better understand dynamics, reduce uncertainty in landscape to global carbon budgets, and project changes to atmospheric greenhouse gas burdens in a warming climate.

**Index terms (5):** 0428 Carbon cycling, 0426 Biosphere/atmosphere interactions, 0438 Diel, seasonal, and annual cycles, 0434 Data sets, 0458 Limnology

**Keywords (6):** eddy covariance; freshwater systems; lakes; reservoirs; carbon flux; synthesis

### Key Points:

- Synthesis of high-frequency aquatic freshwater carbon dioxide flux observations reveals large diel, sub-annual, and interannual variation
- At all sites, nighttime emissions are larger than daytime, sub-monthly variability is present, and year-to-year variation averaged 25%
- Under-sampling of these dynamics leads to potential bias in estimates of contribution of freshwater systems to the global carbon cycle

# 1. Introduction

The global carbon budget is rapidly changing in response to human emissions (Friedlingstein *et al.*, 2020; Hanson *et al.*, 2006). Prior studies have estimated that 0.14-0.64 Pg C-CO<sub>2</sub> is annually released to the atmosphere through lakes and reservoirs (Aufdenkampe *et al.*, 2011; Ciais *et al.*, 2013; Cole *et al.*, 1994, 2007; DelSontro *et al.*, 2018; Drake *et al.*, 2018; Holgerson *et al.*, 2016; Raymond *et al.*, 2013). However, most of these estimates are made with relatively limited sampling, generally constrained to the open-water or summer season during the daytime, and with limited consideration of interannual and shorter-scale variation (Butman *et al.*, 2018; Ran *et al.*, 2021).

Underrepresentation of temporal CO<sub>2</sub> flux variability in existing CO<sub>2</sub> flux inventories may bias estimates of lake CO<sub>2</sub> emissions (Deemer *et al.*, 2016; Klaus *et al.*, 2019). Recent studies have found nighttime emissions exceeding daytime emissions or uptake in reservoirs (Liu *et al.*, 2016) and rivers (Gómez-Gener *et al.*, 2021). A lack of frequent and long-term CO<sub>2</sub> observations also limits our ability to differentiate natural CO<sub>2</sub> flux variations from the consequences of anthropogenic perturbations (Hasler *et al.*, 2016). Multiyear-scale time series that capture sub-annual variability of the aquatic CO<sub>2</sub> flux remain rare (Finlay *et al.*, 2019; Huotari *et al.*, 2011). Traditional in-situ aquatic sampling methods for CO<sub>2</sub> concentrations and fluxes in natural and artificial freshwaters also come with high uncertainty (Baldocchi *et al.*, 2020; Golub *et al.*, 2017), with one source being the heterogeneity of littoral and pelagic lake CO<sub>2</sub> fluxes (Spafford and Risk, 2018).

Advances in the past several decades, however, have enabled more long-term, continuous high-frequency (hourly) measurements in freshwater ecosystems, which are capable of capturing the dynamics of air-water fluxes at time scales of hours to years (Eugster *et al.*, 2003; Huotari *et al.*, 2011; Morales-Pineda *et al.*, 2014). At these time scales, CO<sub>2</sub> fluxes have been shown to respond to variations in photosynthesis and respiration rates (Cole *et al.*, 2007), wind speed and direction (Podgrasjek *et al.*, 2015), carbonate equilibria (Atilla *et al.*, 2011), ecosystem metabolism (Provenzale *et al.*, 2018), convective mixing (Eugster *et al.*, 2003; Mammarella *et al.*, 2015), internal waves (Heiskanen *et al.*, 2014), ice phenology (Reed *et al.*, 2018), and hydrological and carbon inflows/outflows (Rantakari *et al.*, 2005; Weyhenmeyer *et al.*, 2015). These sources of

variation may be overlooked by low-frequency and season-restricted sampling that dominate freshwater science (Desai *et al.*, 2015).

Many previous studies were conducted using eddy covariance (EC) flux towers, which measure ecosystem-scale air-water CO<sub>2</sub> fluxes (Vesala *et al.*, 2006). This method has also gained prominence for use in freshwaters (Vesala *et al.*, 2012). While its application over lakes has mostly covered short periods of time (e.g., Eugster *et al.*, 2003; Podgrajsek *et al.*, 2015; Vesala *et al.*, 2006), an increasing number of sites are now measuring lake-atmosphere fluxes continuously over multiple years (Franz *et al.*, 2016; Huotari *et al.*, 2011; Mammarella *et al.*, 2015; Reed *et al.*, 2018; Eugster *et al.*, 2020). Other methods for high frequency sampling have also included the use of forced diffusion autochambers (Spafford and Risk, 2018). Here, to identify modes of CO<sub>2</sub> flux variability missed by infrequent sampling that may lead to biases in estimates of annual CO<sub>2</sub> flux from lakes and reservoirs quantify diel to inter-annual dynamics of CO<sub>2</sub> fluxes, directly measured by EC from 13 lakes and reservoirs representing a broad nutrient-humic spectrum of sites in the Northern Hemisphere.

## 2. Materials and Methods

### 2.1 Study sites

Data on air-water CO<sub>2</sub> exchange and meteorological drivers were acquired from nineteen study sites across the Northern Hemisphere with at least one season of observations between 2005-2015, of which 13 were retained here for analysis (Table 1 and S1). The six remaining submitted sites were withheld for challenges in meeting uncertainty and gap filling criteria (see Supplemental Methods). These sites were collected based on organization of a workshop (Desai *et al.*, 2015) and an open call through listservs. Selected sites included 9 lakes and 4 reservoirs, mostly located between 40-68°N latitude, coinciding with the largest area of Earth covered with lakes. Eight sites had data available over multiple seasons, but only a few also had measurements during winter ice cover. Lake area ranged from 0.036 km<sup>2</sup> to 623 km<sup>2</sup> (median: 15.2 km<sup>2</sup>), with median mean depth of 6 m (range: 0.6 to 11 m); most developed a seasonal thermocline and were



dimictic or monomictic (Table S1). Two water bodies had a significant fraction of submerged and emergent macrophytes (SE-Tam and DE-Zrk) within the footprint of the flux tower.

## 2.2 Measurements

The EC technique directly measures the exchange of momentum, heat and matter (water vapor, CO<sub>2</sub>, or other trace gasses) at the air-water interface and is a reliable method for measuring surface exchanges with the atmosphere (Vesala *et al.*, 2006). The flux towers were located on floating platforms, lake shoals or islands, or on shore depending on the site (Table S1). The towers were additionally equipped with instruments providing half-hourly to hourly measurements of biophysical variables (e.g. net radiation (R<sub>net</sub>), air temperature (TA) and humidity, photosynthetically-active radiation (PAR), 2-D wind direction and speed, water surface temperature (TW), aquatic CO<sub>2</sub> or O<sub>2</sub> concentration, and water level), although data availability and frequency varied among the sites. Data were harmonized to uniform formats and units, screened for fetch, de-spiked, and gap-filled using a common flux post-processing standard prior to calculation of diel and monthly averages (Pastorello *et al.*, 2020 and supporting material text). Note that a negative CO<sub>2</sub> flux indicates uptake by the ecosystem from the atmosphere and a positive flux means the reverse. All data are published in the Environmental Data Initiative repository (Golub *et al.*, 2022).

## 2.3 Data analysis

We analyzed the half-hourly CO<sub>2</sub> fluxes and three major groups of biophysical covariates. The first group included variables related to wind forcing acting on the water surface (i.e. friction velocity, wind speed, momentum flux). The second group encompassed variables related to temperature cycles and proxies of energy in the system (i.e. TA, TW,  $\Delta T$  (TW - TA), sensible (H) and latent heat (LE) fluxes). The last group included the variables associated with solar radiation -- proxies for primary productivity (i.e.  $\Delta p\text{CO}_2$  ( $p\text{CO}_{2\text{water}} - p\text{CO}_{2\text{air}}$ ), PAR). To determine the standardized difference between two means with repeated unpaired measurements and imbalanced population sizes, we used the Cohen's *d* test where the mean difference between the mean daily CO<sub>2</sub> fluxes is divided by the pooled variance. A coefficient *d* of 0.20, 0.50, 0.80

indicates small, medium, and large standardized differences between the two means, respectively.

To determine the degree of NEE predictability by biophysical drivers (i.e. TA, TW, H and LE, friction velocity ( $U_{star}$ ), and  $R_{net}$ ), we also performed mutual information analysis (MI). Ultimately, this method can reveal dependencies between two variables with co-varying factors, making it a useful approach for ascertaining NEE dependencies on ecosystem variables (Knox et al., 2021). To take into account driver impacts on different temporal scales, we utilized a wavelet-based time scale decomposition approach to decompose half-hourly data into four temporal scales, hourly, diel, multiday, and seasonal, with further details in the supplement and Sturtevant *et al.* (2016).

Finally, a sample analysis was conducted on the oligotrophic (US-Too; 2012), mixotrophic (FI-Van; 2016), and eutrophic (DE-Zrk; 2014) lakes with the smallest data gaps. One thousand random samples without replacement were taken for each of the following times: daytime-only (DT), daytime/nighttime-only (DT/NT), summer mid-day (SMD), growing season (GS), and annual. DT and NT were defined as 10am-3:30pm and 10pm-3:30am (local times) respectively. Hours between 11am and 1:30pm were considered mid-day while the GS counted fluxes between March 1<sup>st</sup> and September 30<sup>th</sup>. Each sample contained either 1, 5, or 10 counts of fluxes. To obtain a single flux value, the samples containing 5 and 10 fluxes were averaged. This sampling algorithm was created using Python version 3.8.3.

## 3. Results

### 3.1 Magnitude of CO<sub>2</sub> fluxes from lakes and reservoirs

Study sites represented a wide range of nutrient-color status and physical characteristics of water bodies, and as a result spanned a range of daily CO<sub>2</sub> fluxes, though with some common elements (Fig. 1). The mean daily CO<sub>2</sub> flux across all sites was  $0.43 \pm 0.34 \mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$  (range:  $-0.075$  to  $1.25 \mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$ ) with only 6% of observations indicating neutral fluxes or net CO<sub>2</sub> uptake. The spread of time-resolved fluxes varied between 102 and 798% of the site-specific

daily mean (Fig. 1). Reservoirs had smaller but more variable fluxes relative to the lakes ( $0.32 \pm 0.71$  vs.  $0.41 \pm 0.31 \mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$ ), though the reservoir sample size is smaller and more geographically restricted. Two thirds of sites had at least 66% of daily fluxes within the cross-site flux mean  $\pm 1$  SD (Cohen's  $d$ :  $0.02 < d < 0.76$ ).

Annually, all sites were  $\text{CO}_2$  sources to the atmosphere, except for DE-Zrk and LA-NT2, with large variability across sites (Fig. 2). This was also the case when comparing the same lake or reservoir type. On a single day (on average), the mixotrophic and eutrophic lakes and reservoirs were the largest and smallest C sources respectively. While most sites were a greater carbon source during the nighttime relative to the daytime, the difference in hourly fluxes was small (range  $\sim 0.5 \text{ C } \mu\text{mol m}^{-2} \text{ s}^{-1}$ ), with the exception of DE-Zrk.

## 3.2 Temporal variability of $\text{CO}_2$ fluxes from lakes and reservoirs

Averaged diel  $\text{CO}_2$  curves had regular patterns of daytime minima and nighttime maxima across all sites in most months (Fig. 2a). Daytime hourly fluxes were on average 35% (range 7-60%) lower than nighttime fluxes, though in 94% of site days, those were still net positive emissions. Despite the commonly observed daytime  $\text{CO}_2$  flux dip, the flux decrease was large enough to convert our sites to daily net sinks of  $\text{CO}_2$  in only 11% of site-months (Fig. 2a). The mean uncertainty of diel  $\text{CO}_2$  was strongly influenced by extreme observations, with 192% mean uncertainty, but only 79% median uncertainty (Fig. 2b).

Maximum diel flux amplitudes typically occurred in July and August and ranged  $0.24$ - $1.09 \mu\text{mol CO}_2 \text{ m}^{-2} \text{ s}^{-1}$ . Relative to the summer amplitudes, shoulder season  $\text{CO}_2$  flux amplitudes were on average 44-49% smaller in May and September and 26-37% in April and October. Diel variation was negligible at both ends of the ice-free season.

Monthly to sub-annual  $\text{CO}_2$  flux variability was nearly twofold compared to diel flux variation. Surprisingly, we found frequent sub-monthly (20-30-day) variability across all water bodies, regardless of the system's physical or biogeochemical conditions. While most site-level variability fluctuated around the  $\text{CO}_2$  flux averages, for some, amplitudes scaled with flux minima and maxima (Fig. S1).

Sites with multi-year data had relatively consistent sub-annual patterns across years, although the timing and amplitudes of sub-monthly variability varied among lake-years. When integrated over time-resolved daily CO<sub>2</sub> fluxes, both sub-monthly and sub-annual modes of variability accounted for two thirds of the site-level daily CO<sub>2</sub> flux variability (range 10-190%). Mean and median uncertainty were 167% and 67% of mean daily CO<sub>2</sub> flux, respectively .

Once scaled to ice-free season annual emissions, and assuming zero fluxes during ice cover, we found all water bodies were net sources of CO<sub>2</sub>, despite missing any ice off/on related fluxes (Table 1). The cross-site mean and standard deviation of 23 site-years was 95±49 gC m<sup>-2</sup> yr<sup>-1</sup> (range: 14 to 224 gC m<sup>-2</sup> yr<sup>-1</sup>). Inter-annual variability (IAV) was calculated as a standard deviation of annual CO<sub>2</sub> flux for each site with multi-year data (Supplemental Fig. S2). The mean cross-site IAV was 22 gC m<sup>-2</sup> yr<sup>-1</sup> (25%) and ranged between 4 and 44 gC m<sup>-2</sup> yr<sup>-1</sup> (3-58%).

### 3.3 Drivers of CO<sub>2</sub> fluxes from lakes and reservoirs

While the continuous data allowed capturing CO<sub>2</sub> flux variability at different temporal scales, we still had a limited capacity to attribute which factors and processes governed the observed patterns of CO<sub>2</sub> flux. We found small standardized differences between CO<sub>2</sub> fluxes among site groups belonging to the three humic states ( $d < 0.01$ ), medium differences between oligotrophic and eutrophic states ( $d = 0.24$ ), and large CO<sub>2</sub> differences between mesotrophic and oligotrophic states ( $d = 0.66$ ), and between mesotrophic and eutrophic states ( $d = 0.72$ ). Commonly observed biophysical covariates explained an average of 32% of variance in half-hourly CO<sub>2</sub> fluxes (Fig. 4g). Wind-related variables were identified as key to explaining CO<sub>2</sub> flux variability in eight out of 13 sites. Biophysical variables related to exchanges of heat at the air-water interface, particularly  $\Delta T$  and turbulent energy exchange (H and LE), correlated with CO<sub>2</sub> flux. The fitted regressions were non-linear and highly variable across sites, owing to ecosystem differences and presence of confounding factors (e.g. differential responses to co-dependent covariates).

Mutual information analysis revealed different drivers to be responsible for CO<sub>2</sub> fluxes on different temporal scales (Supplemental Fig. S3). On hourly scales, NEE at all sites was predicted mostly by TA and TW. The strongest links were found to occur at LA-NT2 and DE-Zrk (both eutrophic). Analysis on diel scales yielded a similar result. On multi-day scales,

however, more linkage between NEE and drivers was found at CA-Dar, SE-Mer, and FI-Pal (all oligotrophic). While the seasonal scale MI analysis was subject to many gaps, it did show a more uniform NEE prediction magnitude across all sites and drivers relative to other timescales.

### 3.4 Sample Analysis

Random sampling among different temporal resolutions resulted in large differences between mean sampled NEE and mean continuous annual NEE (Fig 3). For DE-Zrk and FI-Van, the greatest percent error (PE) was for samples taken during SMD, calculated to be  $868 \pm 26\%$  and  $38 \pm 2\%$  (mean  $\pm$  range), respectively. US-Too experienced the largest error during NT sampling, with a PE of  $87 \pm 31\%$ . Increasing the number of NEE values per sample (i.e. going from 1 to 5 to 10 samples with the latter two NEE values calculated as the average) gave sporadic results, in that, agreement sometimes improved (FI-Van during growing season) and sometimes worsened (US-Too during nighttime). DT/NT and annual sampling were the most representative of continuous annual NEE among all sites regardless of lake/reservoir type. GS sampling showed PE that was well within the typical uncertainty for EC flux measurements ( $\sim 20\%$ ) for FI-Van and US-Too. Sampling on an annual scale further constrained PE, including even DE-Zrk in addition to FI-Van and US-Too.

## 4. Discussion

### 4.1 Unresolved temporal variation in CO<sub>2</sub> fluxes

CO<sub>2</sub> fluxes from lakes and reservoirs exhibited large variability at diel to (inter)-annual scales, which could comprise unresolved sources of uncertainty or bias in current estimates of annual CO<sub>2</sub> fluxes from infrequent and season-restricted sampling. Though our study lakes were not randomly selected and cannot be directly used to upscale (Stanley *et al.*, 2019), they were broadly reflective of common mid-latitude freshwater systems spanning a broad range of humic-status and mixing regimes. Additional considerations for sampling across lake size and

catchment area (Hanson *et al.*, 2007; Holgerson *et al.*, 2016) and hydrological setting (Jones *et al.*, 2018) would be required to design a representative estimate for global upscaling.

We were able to investigate, however, the role of temporal variation on a range of systems that broadly reflect many lakes and reservoirs. Our reported continuous daily fluxes corresponded to the upper end (88<sup>th</sup> percentile) of previously published flux magnitudes (Table S2). The observed temporal variation suggests that temporal restrictions in sampling may add a significant source of underestimation bias in existing inventories of CO<sub>2</sub> fluxes from lakes and reservoirs of similar type and size (Klaus *et al.*, 2019).

In particular, we noted significant diel variation found in all study sites, with routinely higher emissions at night, consistent with a recent study over rivers (Gómez-Gener *et al.*, 2021). The diel reduction of dissolved CO<sub>2</sub> concentrations and fluxes are often associated with ecosystem metabolism (Hanson *et al.*, 2003) and was supported by negative correlations with PAR (Fig. 4g). Water temperature (Provenzale *et al.*, 2018), carbonate equilibria fluctuations (Atilla *et al.*, 2011), water-side convection (Eugster *et al.*, 2003; Mammarella *et al.*, 2015; Podgrajsek *et al.*, 2015), and internal waves (Heiskanen *et al.*, 2014) can additionally govern diel CO<sub>2</sub> dynamics. Our observed diel amplitudes were within 21-43% of sub-hourly flux amplitudes derived from dissolved CO<sub>2</sub> concentrations (Hanson *et al.*, 2003; Morales-Pineda *et al.*, 2014) or previously published EC-measured fluxes (Liu *et al.*, 2016; Vesala *et al.*, 2006). Our results support the notion that existing global lake carbon budgets are underestimates of net emissions.

We also found common sub-monthly modes of CO<sub>2</sub> flux variability across all of our sites. Similar variability in the continuous observations have been reported for dissolved CO<sub>2</sub> (Atilla *et al.*, 2011; Huotari *et al.*, 2009; Morales-Pineda *et al.*, 2014; Vachon and del Giorgio, 2014) and CO<sub>2</sub> fluxes (Franz *et al.*, 2016; Eugster *et al.*, 2020), indicating the prevalence of oscillatory patterns in CO<sub>2</sub> time series at both sides of the air-water interface. Variability has been previously attributed to the interplay of wind forcing (Liu *et al.*, 2016), upwellings of CO<sub>2</sub>-rich waters (Morales-Pineda *et al.*, 2014), biologically-driven (metabolic and trophic) changes in carbonate equilibria (Atilla *et al.*, 2011), convective mixing (Huotari *et al.*, 2009) and TW (Atilla *et al.*, 2011). However, this is the first study to find a consistent pattern in a wide range of systems, regardless of size. We also observed changes to the prevalence of underlying sub-

monthly CO<sub>2</sub> flux variability through the year at several sites, likely reflecting seasonal ecosystem changes, such as spring/fall turnover (Baehr *et al.*, 2004), radiative and heat exchanges (Heiskanen *et al.*, 2014), and hydrological inflows (Vachon *et al.*, 2017).

## 4.2 Implications for the global carbon budget

After our daily fluxes were scaled to annual totals, our estimates of annual CO<sub>2</sub> emissions were in the upper end reported for lakes and reservoirs (Table S2). All systems were sources of CO<sub>2</sub> in most years, though there have been sites that reported significant carbon sinks (e.g., Shao *et al.* 2015; Reed *et al.*, 2018) and additional propagation of uncertainty from data gap filling and filtering (e.g., of nighttime uptake) can push some of our study sites toward sinks, though weakly. While our lakes are not fully representative for all lakes on Earth, we postulate that improved temporal resolution of site-level CO<sub>2</sub> fluxes is one of the sources of differences between this study and published annual fluxes (Table S2). The results also imply that a proposed recommended number of samples per year (4-8) (Klaus *et al.*, 2019; Natchimuthu *et al.*, 2017) is likely insufficient to constrain annual CO<sub>2</sub> fluxes from lakes and reservoirs. Rather, approaches to increase nighttime, open-water season, weekly, and generally higher-frequency sampling would increase the accuracy of annual estimates, given our observed diel and sub-monthly variations.

Additionally, sites with multiple years of data all showed non-trivial interannual variation. The estimate of average IAV of CO<sub>2</sub> fluxes (25%) is modest compared to that (88%) observed in terrestrial ecosystems (Baldocchi *et al.*, 2018), probably reflecting the lower number and diversity of ecosystems with multi-year measurements or more buffering against climate extremes by large water bodies. However, given that CO<sub>2</sub> flux from freshwaters positively scales with the productivity of terrestrial ecosystems at shorter timescales (Butman *et al.*, 2016; Hastie *et al.*, 2018; Walter *et al.*, 2021), it is possible that the interannual variation of carbon displaced from land will propagate onto CO<sub>2</sub> outgassed through freshwaters (Drake *et al.*, 2018; McDonald *et al.*, 2013), providing a possible pathway to constrain freshwater IAV. Neglecting this variation is an additional source of bias in our current view on global CO<sub>2</sub> emissions from lakes and reservoirs.



Given that EC CO<sub>2</sub> fluxes are affected at both sides of the air-water interface (Wanninkhof *et al.*, 2009), a better constraint of the contribution of lakes to the global carbon cycle will also require reporting and synthesis of additional continuous waterside data (e.g. temperature, dissolved CO<sub>2</sub> and O<sub>2</sub>), site-level ecosystem characteristics (e.g. nutrient-color legacies, ecosystem metabolism, and aquatic vegetation such as algae) and sampling an increased site diversity within climatic zones (Lehner and Döll, 2004). With more frequent air and aquatic observations, we will better constrain CO<sub>2</sub> fluxes at different time scales, assess the prevalence of temporal patterns in CO<sub>2</sub> fluxes, and reduce uncertainty in eddy flux measurements over freshwaters (e.g., Ejarque *et al.*, 2021) and therefore improve model estimates of responses of these ecosystems to climate change. Such work will be needed to quantify and evaluate landscape (Buffam *et al.*, 2011; Zwart *et al.*, 2018) to global (DelSontro *et al.*, 2018) carbon budget components from lakes and reservoirs.

### 4.3 Lake and reservoir carbon flux drivers among types

In this study, water temperature (TW) has been shown to be a large predictor of lake and reservoir NEE, agreeing with past work (Zwart *et al.*, 2019; Eugster *et al.*, 2020). There is a high degree of spatiotemporal variability between these two variables. For example, NEE at LA-NT2 and DE-Zrk (eutrophic reservoir and eutrophic shallow lake respectively) was most highly predicted by TW on short timescales (hourly and diel), indicating these ecosystems to be most susceptible to releasing carbon in the future due to a warming climate. This large link may also be explicable through lake type. Eutrophic lakes are defined as being nutrient rich, meaning that they contain larger phosphorus, nitrogen, or dissolved organic carbon concentrations than their oligotrophic counterparts (Reed *et al.*, 2018). On multiday timescales, however, the distinguishability of the NEE/TW linkage is absent relative to all other sites. At least for these two sites, this points to the greatest relative NEE impact of TW to be on short timescales, suggesting a rapid influence on the carbon cycle at these two eutrophic reservoirs. Predictability was seemingly weaker at the other lake sites. Another variable with high NEE predictability was air temperature (TA). This was particularly true for the same sites and timescales. However, it is certainly possible that these fluxes have an indirect relationship with TA in the form of DOC concentration magnitudes (Sobek *et al.*, 2005).



## 5. Conclusions

Across 13 study sites with EC flux observations, on average all lakes and reservoirs were net annual sources of CO<sub>2</sub> to the atmosphere. However, the time series revealed large diel to (sub)-monthly CO<sub>2</sub> flux variability across sites, among a broad range of biogeochemical and physical site characteristics. These modes of variability accounted for two thirds of daily and a quarter of annual CO<sub>2</sub> flux variation, with sub-annual variability dominating over diel and inter-annual flux variabilities. After integrating these modes of variability into time-resolved fluxes, the CO<sub>2</sub> flux estimates were at the upper end of published CO<sub>2</sub> emissions for lakes and reservoirs. Our results support the idea that long-term, frequent measurements at both day and night of carbon dynamics in freshwater aquatic systems are critical to resolve lake C flux magnitudes and detect long-term trends of lake carbon fluxes. Omitting these temporal scales will not only limit our knowledge of lake C fluxes, but also restrict our understanding of biophysical driver impacts.

We advocate for establishing and maintaining a long-term observation network that combines EC flux measurements with highly detailed site-specific carbon budget studies over key lake and reservoir ecosystems representing broader geographical gradients.

## Acknowledgements

We thank all project participants for kindly sharing data and time. All flux tower data and code for processing data will be publicly available in the Environmental Data Initiative depository (DOI pending). MG and ARD acknowledge support from the U.S. National Science Foundation North Temperate Lakes LTER (NSF DEB-1440297, NTL LTER). Funding for US-UM3 was provided by the U.S. Department of Energy's Office of Science. IM and TV thank the support by the EU-Horizon Europe project 101056921 — GreenFeedBack, Academy Professor projects (312571 and 282842), ACCC Flagship funded by the Academy of Finland (337549) and the grant of the Tyumen region, Russia, Government in accordance with the Program of the World-Class West Siberian Interregional Scientific and Educational Center (National Project "Nauka"). GAW was financially supported by the Swedish Research Council (VR: Grant No. 2016-04153 and 2020-03222). The deployment of the EC at the Nam Theun 2 Reservoir (Lao PDR) was

funded by Electricité de France (EDF) and Nam Theun Power Company (NTPC). TS and FK were supported by the Helmholtz Association of German Research Centres through grants to TS (grant VH-NG-821) and FK (grant PD-129), the Helmholtz Climate Initiative REKLIM (Regional Climate Change), and infrastructure funding through the Terrestrial Environmental Observatories Network (TERENO). TS and FK thank the (staff of the) Department Chemical Analytics and Biogeochemistry at the Leibniz-Institute of Freshwater Ecology and Inland Fisheries (Berlin) for providing water chemistry data for DE-Zrk. WE and GK acknowledge support from NSF-DEB 1637459. The deployment of the flux tower at CA-Eastmain was supported by Hydro Quebec. Flux observations at US-OWC were funded by the Ohio Department of Natural Resources, by NOAA's National Estuarine Research Reserves' Davidson Fellowship, and by US Department of Energy awards DE-SC0021067 and DE-SC0022191. The co-authors express gratitude for the kindness and contributions of posthumous co-author Werner Eugster.

## Open Research

We have deposited all EC lake observations and gap-filled values in the Environmental Data Initiative repository Golub *et al.* (2022). Several sites are also accessible from Fluxnet affiliated archives as noted in Table S2.

## Author Contribution Statement

M.G. designed experimental protocol and conducted the data syntheses. N.K.-A. conducted additional analyses and revisions. A.R.D, N.K.-A., and M.G. wrote the manuscript. T.V., I.M., G.B., and G.W. supervised research, contributed observations, and edited the manuscript. All other authors contributed with flux observations and commented on the manuscript.

## Competing Financial Interests

The authors declare no competing financial interests.

## 367    References

- 368    Atilla, N. *et al.* (2011). Observed variability of Lake Superior  $p\text{CO}_2$ . *Limnology and*  
369        *Oceanography* 56, 775–786.
- 370    Aufdenkampe, A. K. *et al.* (2011). Riverine coupling of biogeochemical cycles between land,  
371        oceans, and atmosphere. *Frontiers in Ecology and the Environment* 9, 53–60..
- 372    Baehr, M. M. & DeGrandpre, M. D. (2004). In Situ  $p\text{CO}_2$  and  $\text{O}_2$  Measurements in a Lake during  
373        Turnover and Stratification: Observations and Modeling. *Limnology and Oceanography* 49,  
374        330–340.
- 375    Baldocchi, A., Reed, D.E., Loken, L., Stanley, E., Huerd, H., Desai, A.R. (2020). Resolving  
376        space and time variation of lake-atmosphere carbon dioxide fluxes using multiple methods. *J.*  
377        *Geophys. Res.-Biogeosciences* 125, e2019JG005623, doi:10.1029/2019JG005623.
- 378    Baldocchi, D., Chu, H. & Reichstein, M. (2018). Inter-annual variability of net and gross  
379        ecosystem carbon fluxes: A review. *Agricultural and Forest Meteorology* 249, 520–533.
- 380    Buffam, I. *et al.* (2011). Integrating aquatic and terrestrial components to construct a complete  
381        carbon budget for a north temperate lake district. *Global Change Biology* 17, 1193–1211.
- 382    Butman, D. *et al.* (2016). Aquatic carbon cycling in the conterminous United States and  
383        implications for terrestrial carbon accounting. *Proceedings of the National Academy of*  
384        *Sciences* 113, 58–63.
- 385    Butman, D. *et al.* (2018). Chapter 14: Inland waters. in *Second State of the Carbon Cycle Report*  
386        *(SOCCR2): A Sustained Assessment Report* (eds. Cavallaro, N. *et al.*) (U.S. Global Change  
387        Research Program, Washington, DC, USA).
- 388    Ciais, P. *et al.* (2013). Carbon and Other Biogeochemical Cycles. in *Climate Change 2013: The*  
389        *Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of*  
390        *the Intergovernmental Panel on Climate Change* (eds. Stocker, T. F. *et al.*) (Cambridge  
391        University Press, Cambridge, United Kingdom and New York, NY, USA.).
- 392    Cole, J. J., Caraco, N. F., Kling, G. W. & Kratz, T. K. (1994). Carbon Dioxide Supersaturation in  
393        the Surface Waters of Lakes. *Science*, 265, 1568–1570.
- 394    Cole, J. J. *et al.* (2007). Plumbing the Global Carbon Cycle: Integrating Inland Waters into the  
395        Terrestrial Carbon Budget. *Ecosystems* 10, 172–185.
- 396    Deemer, B. R. *et al.* (2016). Greenhouse Gas Emissions from Reservoir Water Surfaces: A New  
397        Global Synthesis. *BioScience* 66, 949–964.
- 398    DelSontro, T., Beaulieu, J. J. & Downing, J. A. (2018). Greenhouse gas emissions from lakes  
399        and impoundments: Upscaling in the face of global change. *Limnology and Oceanography*  
400        *Letters* 3, 64–75.
- 401    Desai, A.R., Vesala, T., and Rantakari, M. (2015). Measurements, modeling, and scaling of  
402        inland water gas exchange. *EOS* 96, doi:10.1029/2015EO022151.
- 403    Drake, T. W., Raymond, P. A. & Spencer, R. G. M. (2018). Terrestrial carbon inputs to inland  
404        waters: A current synthesis of estimates and uncertainty. *Limnology and Oceanography*  
405        *Letters* 3, 132–142.
- 406    Ejarque, E., Scholz, K., Wohlfahrt, G., Battin, T.J., Kainz, M.J., and Schelker, J. (2021).  
407        Hydrology controls the carbon mass balance of a mountain lake in the eastern European  
408        Alps. *Limnology and Oceanography*, doi:10.1002/lno.11712.
- 409    Eugster, W. *et al.* (2003)  $\text{CO}_2$  exchange between air and water in an Arctic Alaskan and  
410        midlatitude Swiss lake: Importance of convective mixing. *Journal of Geophysical Research*  
411        108, 4362.

- Eugster, W., T. DelSontro, G. R. Shaver, and G. W. Kling. 2020. Interannual, summer, and diel variability of CH<sub>4</sub> and CO<sub>2</sub> effluxes from Toolik Lake, Alaska, during the ice-free periods 2010–2015. *Environmental Science: Processes & Impacts* 22, 2181–2198. DOI: 10.1039/D0EM00125B
- Finlay, K., Vogt, R. J., Simpson, G. L. & Leavitt, P. R. (2019). Seasonality of *p*CO<sub>2</sub> in a hard-water lake of the northern Great Plains: The legacy effects of climate and limnological conditions over 36 years. *Limnology and Oceanography* 64, S118–S129.
- Franz, D., Koebsch, F., Larmanou, E., Augustin, J. & Sachs, T. (2016). High net CO<sub>2</sub> and CH<sub>4</sub> release at a eutrophic shallow lake on a formerly drained fen. *Biogeosciences* 13, 3051–3070.
- Friedlingstein, P. *et al.* (2020). Global Carbon Budget 2020. *Earth System Science Data* 12, 3269–3340, doi:10.5194/essd-12-3269-2020.
- Golub, M., Desai, A. R., McKinley, G. A., Remucal, C. K. & Stanley, E. H. (2017). Large Uncertainty in Estimating *p*CO<sub>2</sub> From Carbonate Equilibria in Lakes. *Journal of Geophysical Research: Biogeosciences* 122, 2909–2924.
- Golub, M., A.R. Desai, T. Vesala, I. *et al.* (2022). Half-hourly gap-filled Northern Hemisphere lake and reservoir carbon flux and micrometeorology, 2006 - 2015 ver 1. Environmental Data Initiative. <https://doi.org/10.6073/pasta/87a35ca843d8739d75882520c724e99e>
- Gómez-Gener, L., Rocher-Ros, G., Battin, T. *et al.* (2021). Global carbon dioxide efflux from rivers enhanced by high nocturnal emissions. *Nat. Geosci.* doi:10.1038/s41561-021-00722-3.
- Hanson, P. C., Bade, D. L., Carpenter, S. R. & Kratz, T. K. (2003). Lake Metabolism: Relationships with Dissolved Organic Carbon and Phosphorus. *Limnology and Oceanography* 48, 1112–1119.
- Hanson, P. C., Carpenter, S. R., Armstrong, D. E., Stanley, E. H. & Kratz, T. K. (2006). Lake Dissolved Inorganic Carbon and Dissolved Oxygen: Changing Drivers from Days to Decades. *Ecological Monographs* 76, 343–363.
- Hanson, P. C., Carpenter, S. R., Cardille, J. A., Coe, M. T. and Winslow, L. A. (2007). Small lakes dominate a random sample of regional lake characteristics. *Freshwater Biology* 52 814–822. doi:10.1111/j.1365-2427.2007.01730.x.
- Hasler, C. T., Butman, D., Jeffrey, J. D. & Suski, C. D. (2016). Freshwater biota and rising *p*CO<sub>2</sub>? *Ecology Letters* 19, 98–108.
- Hastie, A. *et al.* (2018). CO<sub>2</sub> evasion from boreal lakes: Revised estimate, drivers of spatial variability, and future projections. *Global Change Biology* 24, 711–728.
- Heiskanen, J. J. *et al.* (2014). Effects of cooling and internal wave motions on gas transfer coefficients in a boreal lake. *Tellus B: Chemical and Physical Meteorology* 66, 22827.
- Holgerson, M. A. & Raymond, P. A. (2016). Large contribution to inland water CO<sub>2</sub> and CH<sub>4</sub> emissions from very small ponds. *Nature Geoscience* 9, 222–226.
- Huotari, J. *et al.* (2009). Temporal variations in surface water CO<sub>2</sub> concentration in a boreal humic lake based on high-frequency measurements. *Boreal Environment Research* 14, 48–60.
- Huotari, J. *et al.* (2011) Long-term direct CO<sub>2</sub> flux measurements over a boreal lake: Five years of eddy covariance data. *Geophysical Research Letters* 38, L18401.
- Jones, S.E., Zwart, J.A., Kelly, P.T. and Solomon, C.T. (2018). Hydrologic setting constrains lake heterotrophy and terrestrial carbon fate. *Limnol. Oceanogr.* 3 256–264. doi:10.1002/lol2.10054.
- Klaus, M., Seekell, D. A., Lidberg, W. & Karlsson, J. (2019). Evaluations of Climate and Land Management Effects on Lake Carbon Cycling Need to Account for Temporal Variability in CO<sub>2</sub> Concentrations. *Global Biogeochemical Cycles* 33, 243–265.

- Knox *et al.* (2021). Identifying dominant environmental predictors of freshwater wetland methane fluxes across diurnal to seasonal time scales. *Global Change Biology*, 27(15), pp.3582-3604. doi: 10.1111/gcb.15661
- Lehner, B. & Döll, P. (2004). Development and validation of a global database of lakes, reservoirs and wetlands. *Journal of Hydrology* 296, 1–22.
- Liu, H. *et al.* (2016). Large CO<sub>2</sub> effluxes at night and during synoptic weather events significantly contribute to CO<sub>2</sub> emissions from a reservoir. *Environmental Research Letters* 11, 064001.
- Mammarella, I. *et al.* (2015). Carbon dioxide and energy fluxes over a small boreal lake in Southern Finland: CO<sub>2</sub> and Energy Fluxes Over Lake. *Journal of Geophysical Research: Biogeosciences* 120, 1296–1314.
- McDonald, C. P., Stets, E. G., Striegl, R. G. & Butman, D. (2013). Inorganic carbon loading as a primary driver of dissolved carbon dioxide concentrations in the lakes and reservoirs of the contiguous United State. *Global Biogeochemical Cycles* 27, 285–295.
- Morales-Pineda, M., Cózar, A., Laiz, I., Úbeda, B. & Gálvez, J. Á. (2014). Daily, biweekly, and seasonal temporal scales of pCO<sub>2</sub> variability in two stratified Mediterranean reservoirs. *Journal of Geophysical Research: Biogeosciences* 119, 509–520.
- Natchimuthu, S., Sundgren, I., Gålfalk, M., Klemedtsson, L. & Bastviken, D. (2017). Spatiotemporal variability of lake pCO<sub>2</sub> and CO<sub>2</sub> fluxes in a hemiboreal catchment. *Journal of Geophysical Research: Biogeosciences* 122, 30–49.
- Podgrajsek, E., Sahlée, E. & Rutgersson, A. (2015). Diel cycle of lake-air CO<sub>2</sub> flux from a shallow lake and the impact of waterside convection on the transfer velocity. *Journal of Geophysical Research: Biogeosciences* 120, 29–38.
- Provenzale, M. *et al.* (2018). High-frequency productivity estimates for a lake from free-water CO<sub>2</sub> concentration measurements. *Biogeosciences* 15, 2021–2032.
- Ran, L., Butman, D.E., Battin, T.J., *et al.* (2021). Substantial decrease in CO<sub>2</sub> emissions from Chinese inland waters due to global change. *Nat Commun* 12, 1730, doi:10.1038/s41467-021-21926-6.
- Rantakari, M. & Kortelainen, P. (2005). Interannual variation and climatic regulation of the CO<sub>2</sub> emission from large boreal lakes. *Global Change Biology* 11, 1368–1380.
- Raymond, P. A. *et al.* (2013). Global carbon dioxide emissions from inland waters. *Nature* 503, 355–359.
- Reed, D.R., Dugan, H., Flannery, A., and Desai, A.R. (2018). Carbon sink and source dynamics of a eutrophic deep lake using multiple flux observations over multiple years. *Limnology and Oceanography Letters* 3, 285-292, doi:10.1002/lol2.10075.
- Shao, C., Chen, J., Stepien, C. A., Chu, H., Ouyang, Z., Bridgeman, T. B., Czajkowski, K. P., Becker, R. H., and John, R. (2015), Diurnal to annual changes in latent, sensible heat, and CO<sub>2</sub> fluxes over a Laurentian Great Lake: A case study in Western Lake Erie. *J. Geophys. Res. Biogeosci.* 120, 1587– 1604, doi:10.1002/2015JG003025.
- Sobek, S *et al.* (2005), Temperature Independence Of Carbon Dioxide Supersaturation In Global Lakes. *Global Biogeochemical Cycles*, 19, <https://doi.org/10.1029/2004gb002264>.
- Spafford, L., & Risk, D. (2018). Spatiotemporal variability in lake-atmosphere net CO<sub>2</sub> exchange in the littoral zone of an oligotrophic lake. *Journal of Geophysical Research: Biogeosciences*, 123, 1260–1276. <https://doi.org/10.1002/2017JG004115>
- Stanley, E.H., Collins, S.M., Lottig, N.R., Oliver, S.K., Webster, K.E., Cheruvilil, K.S. and



Soranno, P.A. (2019). Biases in lake water quality sampling and implications for macroscale research. *Limnol. Oceanogr.* 64, 1572-1585. doi:10.1002/lno.11136.

Sturtevant, C., Ruddell, B., Knox, S., Verfaillie, J., Matthes, J., Oikawa, P. and Baldocchi, D., 2016. Identifying scale-emergent, nonlinear, asynchronous processes of wetland methane exchange. *Journal of Geophysical Research: Biogeosciences*, 121(1), pp.188-204.

Vachon, D. & del Giorgio, P. A. (2014). Whole-Lake CO<sub>2</sub> Dynamics in Response to Storm Events in Two Morphologically Different Lakes. *Ecosystems* 17, 1338–1353.

Vachon, D., Solomon, C. T. & del Giorgio, P. A. (2017). Reconstructing the seasonal dynamics and relative contribution of the major processes sustaining CO<sub>2</sub> emissions in northern lakes: Lake CO<sub>2</sub> seasonal dynamics. *Limnology and Oceanography* 62, 706–722.

Vesala, T. *et al.* (2006). Eddy covariance measurements of carbon exchange and latent and sensible heat fluxes over a boreal lake for a full open-water period. *Journal of Geophysical Research* 111, D11101.

Vesala, T., Eugster, W. & Ojala, A. (2012). Eddy covariance measurements over lakes. In *Eddy Covariance: A Practical Guide to Measurement and Data Analysis* (eds. Aubinet, M., Vesala, T. & Papale, D.) 365–376 (Springer, Dordrecht).

Walter, J. A., Fleck, R., Kastens, J. H. *et al.* (2012). Temporal Coherence Between Lake and Landscape Primary Productivity. *Ecosystems* 24, 502–515, doi:10.1007/s10021-020-00531-6

Wanninkhof, R., Asher, W. E., Ho, D. T., Sweeney, C. & McGillis, W. R. (2009). Advances in Quantifying Air-Sea Gas Exchange and Environmental Forcing. *Annual Review of Marine Science* 1, 213–244.


Weyhenmeyer, G. A. *et al.* (2015). Significant fraction of CO<sub>2</sub> emissions from boreal lakes derived from hydrologic inorganic carbon inputs. *Nature Geoscience* 8, 933–936.

Zwart, J. A. *et al.* (2018). Spatially Explicit, Regional-Scale Simulation of Lake Carbon Fluxes. *Global Biogeochemical Cycles* 32, 1276– 1293.


Zwart, J. A., Hanson, Z. J., Read, J. S., Fienen, M. N., Hamlet, A. F., Bolster, D., & Jones, S. E. (2019). Cross-scale interactions dictate regional lake carbon flux and productivity response to future climate. *Geophysical Research Letters*, 46, 8840–8851. <https://doi.org/10.1029/2019GL083478>

# Figures

**Fig. 1.** Normalized histograms of daily CO<sub>2</sub> fluxes over ice-free season in nine lakes and four reservoirs, showing that all studied ecosystems emitted CO<sub>2</sub> to atmosphere in the majority of site-days. Vertical solid lines and their numerical representation indicate mean daily CO<sub>2</sub> flux. Shaded areas show observations with negative CO<sub>2</sub> flux, which by convention, indicate net CO<sub>2</sub> uptake.




541 **Fig. 2.** Daily (A), cumulative (B), and summer diel cycle (C) of NEE for all 13 sites.  
 542 Oligotrophic, mixotrophic, and eutrophic lakes and reservoirs are represented by red, blue, and  
 543 green lines respectively. Averaged NEE is reported for sites with multiple years of data.





**Fig. 3.** Sample analysis for the mixotrophic (blue), eutrophic (green), and oligotrophic (red) lakes and reservoirs with the least data-gaps. Each bar shows percent error between randomly sampled mean NEE (without replacement) and mean continuous annual NEE. A zoomed in version of the plot is shown to better distinguish differences between FI-Van and US-To.



## 551 Tables

552 **Table 1.** Comparison of ice-free CO<sub>2</sub> flux at temporal (i.e. annual, seasonal, diurnal and  
553 nocturnal) scales derived from high-frequency eddy covariance measurements over lakes and  
554 reservoirs. One standard deviation of the mean represents uncertainty of sub-annual CO<sub>2</sub> fluxes.  
555 The numbers in brackets represent the number of observations integrated at a given time scale.

Lake ID	Name	Year	Air-water CO <sub>2</sub> fluxes			
			Annual Totals [gC m <sup>-2</sup> yr <sup>-1</sup> ]	Seasonal daily mean [mgC m <sup>-2</sup> d <sup>-1</sup> ]	Daytime flux [mgC m <sup>-2</sup> hr <sup>-1</sup> ]	Nighttime flux [mgC m <sup>-2</sup> hr <sup>-1</sup> ]
CA-Dar	Daring Lake	2006	na	89±157 (n=95)	0.8±10.7 (n=1685)	12.2±7.5 (n=497)
CA-Est	Eastmain Reservoir	2008	119.1 (n=214)	581±398 (n=214)	22.4±27.5 (n=2790)	26.2±23.4 (n=2117)
		2009	137.2 (n=214)	610±433 (n=214)	21.9±24.2 (n=2786)	30.1±25.1 (n=2127)
		2010	na	431±335 (n=214)	18±20.8 (n=2804)	17.9±19.5 (n=2108)
		2011	75.9 (n=214)	367±272 (n=173)	15.2±18.8 (n=2399)	15.7±15.7 (n=1568)
		2012	na (n=214)	na	na	na
DE-Zrk	Zarnekow Polder Reservoir	2013	-126.1 (n=214)	81± 880 (n=170)	-78.6±111.6 (n=2240)	103.1±47.5 (n=1678)
		2014	-190.7 (n=214)	-250± 835 (n=214)	-86±104.1 (n=2817)	81.4±42 (n=2098)
		2015	-29.5 (n=214)	396±1148 (n=214)	-41.2±101.2 (n=2791)	84.2±54.6 (n=2139)
FI-Kui	Kuivajarvi Lake	2010	31.4 (n=214)	643±140 (n= 58)	22.8±13.9 (n= 670)	30.2±13.8 (n= 656)
		2011	107.9 (n=214)	1047±304 (n=153)	39.7±17.1 (n=2075)	48.3±21.2 (n=1455)
		2012	91.5 (n=241)	684±274 (n=169)	24.4±16.5 (n=1981)	32.4±18.4 (n=1893)
FI-Pal	Pallasjärvi Lake	2013	21.9 (n=173)	304±154 (n=93)	8.8±9.8 (n=1201)	17.2±9.9 (n=939)
FI-VKa	Valkea-Kotinen Lake	2003	59.7 (n=209)	544±155 (n=208)	22±7 (n=2385)	23.4± 8.8 (n=1848)
		2004	46.4 (n=239)	450±261 (n=238)	16.5±16.4 (n=2986)	21±13.6 (n=2464)
		2005	31.1 (n=227)	384±215 (n=226)	11.4±15.3 (n=2940)	22.6±9.1 (n=2103)
		2006	40.6 (n=254)	472±263 (n=253)	15.8± 13 (n=2983)	23.4±13.4 (n=2824)
		2007	43.6 (n=222)	539±232 (n=221)	20.8±11.3 (n=3033)	24.5±13.5 (n=2038)
		2008	-10.9 (n=101)	na	na	na
		2009	na	na	na	na
FI-Van	Vänajavesi Lake	2016	105 (n=237)	457±334 (n=237)	17.6±18.7 (n=2943)	20.8±17.8 (n=2505)
		2017	na	na	na	na
LA-NT2	NamTheun 2 Reservoir	2008	na	1762±186 (n=10)	61±17.8 (n=125)	87±39.2 (n=106)
		2009	na	1623±345 (n=15)	73.5±28.2 (n=146)	63.2±29.7 (n=200)
		2010	na	861±183 (n= 4)	36±16.3 (n= 47)	35.3±13.5 (n= 46)
		2011	na	na	na	na
SE-Mer	Merasjärvi Lake	2005	9 (n=165)	145±149 (n=117)	4.7±9.4 (n=1877)	8.6±9.3 (n=835)
SE-Tam	Tamnaren Lake	2010	8.5 (n=216)	189±125 (n= 49)	6.9± 9.9 (n= 493)	8.6±11.4 (n= 628)
		2011	28.5 (n=291)	124±161 (n=290)	4.9±11.2 (n=3619)	5.3±10 (n=3027)
		2012	na	386±176 (n=105)	10.8±12.3 (n=1663)	27.4±16.2 (n= 743)
US-UM3	Douglas Lake	2013	46.8 (n=275)	432±318 (n=102)	10.5±25.9 (n=1374)	28.5± 39 (n= 965)
		2014	60.1 (n=275)	412±313 (n=142)	9.6±24.9 (n=1889)	27.7±38.7 (n=1380)
US-RBa	Ross Barnett Reservoir	2007	20.3 (n=365)	162±308 (n=129)	5±23.5 (n=1324)	8.4±27.4 (n=1659)
US-Too	Toolik Lake	2012	na	304±130 (n=62)	8±13 (n=1120)	28±27.5 (n=308)

Figure 1.






Figure 2.

**A** **Daily NEE**



**B** **Cumulative NEE**



- Oligotrophic
- Mixotrophic
- Eutrophic

**C** **Summer Diel Cycle**





Figure 3.



