

Exotic tree plantations in the Chilean Coastal Range: Balancing effects of discrete disturbances, connectivity and a persistent drought on catchment erosion

Violeta Tolorza¹, Christian Mohr², Mauricio Zambrano-Bigiarini³, Benjamin Sotomayor⁴, Dagoberto Poblete-Caballero⁵, Sebastien Carretier⁶, Mauricio Galleguillos⁷, and Oscar Seguel⁵

¹Universidad de La Frontera

²University of Potsdam, Germany

³Universidad de La Frontera, Center for Climate and Resilience Research (CR2)

⁴Dron Aerogeomática SpA, Spatial Data and Analysis in Aysén, Chile

⁵Universidad de Chile

⁶Geosciences Environnement Toulouse

⁷Universidad Adolfo Ibáñez

August 10, 2023

Abstract

The Coastal Range in the Mediterranean segment of Chile is a soil mantled landscape with potential to store valuable supplies of fresh water and support a biodiverse native forest. Nevertheless, human intervention has been increasing soil erosion for ~ 200 yr, with intensive management of exotic tree plantations during the last ~ 45 yr. At the same time, this landscape has been affected by a prolonged megadrought, and is not yet well understood how the combined effect of anthropogenic disturbances and hydrometeorologic trends affect sediment transport at the catchment scale.

In this study we calculate a decadal-scale catchment erosion rate from suspended sediment loads and compare it with a 10⁴-year-scale catchment denudation rate estimated from detritic ¹⁰Be. We then contrast these rates against the effects of discrete anthropogenic disturbances and hydroclimatic trends. Erosion/denudation rates are similar on both time scales, i.e. 0.018 ± 0.005 mm/yr and 0.024 ± 0.004 mm/yr, respectively. Recent human-made disturbances include logging operations during each season and a dense network of forestry roads, which increase structural sediment connectivity. Other disturbances include two widespread wildfires (2015 and 2017) and one Mw 8.8 earthquake (2010).

We observe a decrease in suspended sediment load during the wet seasons for the period 1986-2018 coinciding with declines in streamflow, baseflow and rainfall. The low 10⁴-year denudation rate agrees with a landscape dominated by slow diffusive soil creep. However, the low 10-year-scale erosion rate and the decrease in suspended sediments are not in agreement with the expected effect of intensive anthropogenic disturbances and increased structural (sediment) connectivity. These paradox suggest that, either suspended sediment loads and, thus, catchment erosion, are underestimated, and/or that decennial sediment detachment and transport were smeared by decreasing rainfall and streamflow. Our findings indicate that human-made disturbances and hydrometeorologic trends may result in opposite, partially offsetting effects on recent erosion, but both contribute to the landscape degradation.

Exotic tree plantations in the Chilean Coastal Range: Balancing effects of discrete disturbances, connectivity and a persistent drought on catchment erosion

Violeta Tolorza¹, Christian H Mohr², Mauricio Zambrano-Bigiarini^{1,3}, Benjamín Sotomayor⁴, Dagoberto Poblete-Caballero⁵, Sebastien Carretier⁶, Mauricio Galleguillos^{3,7}, and Oscar Seguel⁵

¹Universidad de La Frontera. 4811230 Temuco, Chile

²Institute of Environmental Science and Geography, University of Potsdam, Germany

³Center for Climate and Resilience Research (CR2). Blanco Encalada 2002, 4th floor, Santiago, Chile

⁴Dron Aerogeomática SpA, Spatial Data and Analysis in Aysén, Chile

⁵School of Agronomic Sciences, Universidad de Chile. 11315 Santiago, Chile

⁶Geosciences Environnement Toulouse, IRD, OMP, UPS, CNRS, Université de Toulouse, France

⁷Facultad de Ingeniería y Ciencias, Universidad Adolfo Ibáñez, Peñalolen, Chile

Correspondence: Violeta Tolorza (violeta.tolorza@ufrontera.cl)

Abstract. The Coastal Range in the Mediterranean segment of Chile is a soil mantled landscape with potential to store valuable supplies of fresh water and support a biodiverse native forest. Nevertheless, human intervention has been increasing soil erosion for ~200 yr, with intensive management of exotic tree plantations during the last ~45 yr. At the same time, this landscape has been affected by a prolonged megadrought, and is not yet well understood how the combined effect of anthropogenic disturbances and hydrometeorologic trends affect sediment transport at the catchment scale.

In this study we calculate a decadal-scale catchment erosion rate from suspended sediment loads and compare it with a 10⁴-year-scale catchment denudation rate estimated from detritic ¹⁰Be. We then contrast these rates against the effects of discrete anthropogenic disturbances and hydroclimatic trends. Erosion/denudation rates are similar on both time scales, i.e. 0.018 ±0.005 mm/yr and 0.024 ±0.004 mm/yr, respectively. Recent human-made disturbances include logging operations during each season and a dense network of forestry roads, which increase structural sediment connectivity. Other disturbances include two widespread wildfires (2015 and 2017) and one Mw 8.8 earthquake (2010).

We observe a decrease in suspended sediment load during the wet seasons for the period 1986-2018 coinciding with declines in streamflow, baseflow and rainfall. The low 10⁴-year denudation rate agrees with a landscape dominated by slow diffusive soil creep. However, the low 10-year-scale erosion rate and the decrease in suspended sediments are not in agreement with the expected effect of intensive anthropogenic disturbances and increased structural (sediment) connectivity. These paradox suggest that, either suspended sediment loads and, thus, catchment erosion, are underestimated, and/or that decennial sediment detachment and transport were smeared by decreasing rainfall and streamflow. Our findings indicate that human-made disturbances and hydrometeorologic trends may result in opposite, partially offsetting effects on recent erosion, but both contribute to the landscape degradation.

Over 75% of Earth's ice-free land has been altered by humans (Ellis and Ramankutty, 2008), with severe consequences for sediment transport during the Anthropocene (Syvitski et al., 2022). Land Use and Land Cover Changes (LULCC) are important in increasing soil erosion (Borrelli et al., 2020). Human-made forests – or better, tree plantations (DellaSala, 2020) – are frequently disturbed by logging and the implementation of forestry roads. Such disturbances may intensify soil erosion (e.g., Schuller et al., 2013; Sidle and Ziegler, 2012), as may heavy machinery traffic (e.g., Malmer and Grip, 1990), wildfires and terracing (e.g., Martins et al., 2013). Short rotational cycles, i.e. the period between planting, harvesting, and replanting of tree plantations, also change hillslope stability by cycles of root strength decay and recovery, which in turn promote landsliding and debris flows (Imaizumi et al., 2008; Montgomery et al., 2000). Ultimately, all such processes may modify sediment trajectories and storage on hillslopes and along rivers (Wainwright et al., 2011) with long-lasting impacts on sediment yields for periods of 10-100 years (Moody and Martin, 2009; Bladon et al., 2014).

The Chilean Coastal Range (CCR) in its Mediterranean section (35-37.5° S) is a landscape of gentle and largely convex hillslopes (Fig. 1). Here, forests, soils and water are closely coupled (Galleguillos et al., 2021). This morphology results from relatively slow denudation rates by soil creep on regolith-mantled landscapes (Roering et al., 2007), yet modified by the underlying bedrock (Gabet et al., 2021). Currently, the remnants of secondary native forest stand on soils as thick as 2 m (Soto et al., 2019), suggesting such minimum soil depths under undisturbed conditions. In the absence of snow storage, these soils form a major fresh water supply along the Mediterranean CCR, which many rural communities rely on. Thus, decision-making regarding land management is strategic for the resilience of these communities (e.g., Gimeno et al., 2022), especially under recent (Garreaud et al., 2020) and projected (IPCC, 2021) conditions of water scarcity.

The CCR has experienced deforestation for more than 200 years (Armesto et al., 2010) intensifying soil erosion, as has been recognized by Bianchi-Gundian (1947) and Chilean governments in the middle of 20th century (IREN, 1965). From the beginning of 20th century, governments blamed environmental issues due to deforestation to promote the expansion of tree plantations (e.g., CONAF and MINAGRI, 2016; Pizarro et al., 2020). The most relevant transformation of land cover began with the law DL 701 (1974) to subsidize the forestry sector (Manuschevich, 2020). This law and following political action accelerated LULCC, which in practice transformed degraded lands, shrublands and native forest into industrially managed tree plantations (Heilmayr et al., 2016). From ~450,000 ha of tree plantations in 1974 (Barros, 2018), their spatial extent increased to at least some 2.8 ± 0.2 million ha in 2011 (Heilmayr et al., 2016), mostly within the Mediterranean CCR (Fig. 1).

In Chile, tree plantations are managed mostly as monocultures of fast-growing *Eucalyptus* spp or *Pinus Radiata*. The rotation cycles are as short as 9-12 and 18-25 years, respectively (INFOR, 2004; Gerding, 1991). Harvesting commonly occurs by clear-cutting, and their extent usually expand over entire hillslopes (Fig. 2, supplementary video S1). Such practice is permitted by current Chilean law, as clear-cutting requires environmental impact assessments only for harvest areas ≥ 500 ha/yr or $\geq 1,000$ ha/yr in Mediterranean and Temperate regions, respectively (*Artículo Primero, Título I, Artículo 3, m.1* at Chilean Law 19.300, 2013). As a consequence, the CCR ranks among the highest worldwide in terms of combined forest loss and gain (Hansen et al., 2013).




Figure 1. Study region. A. Land cover in the Coastal Range (Zhao et al., 2016) and catchments with published detrital ^{10}Be denudation rates outlined in black (Carretier et al., 2018). The Purapel catchment, which denudation rate is presented in this work, is represented in red. B. Purapel catchment. All the detected forestry roads intersecting tree plantations and the position of photos in C are shown. Elevation data comes from a 5-m resolution LiDAR DTM obtained in 2009. C. Photos captured on hillslopes of Purapel catchment.

Tree plantations frequently are intersected by dense networks of logging roads. These roads are intended to facilitate access and use of heavy forest machinery, storage and transport of timber, as well as the subsequent (re-)plantation. At the storm to yearly scale (10^{-4} - 10^0 yr), logged hillslopes, like logging roads, are important sediment sources and routes during storms and after wet-season clear-cutting (Schuller et al., 2013, 2021; Aburto et al., 2020). For example, Aburto et al. (2020) reported highest post-harvest soil loss in a catchment sustaining a one-year-old plantation. Post-harvest erosion is mainly rainfall triggered (Aburto et al., 2020; Schuller et al., 2013) and after exceeding specific rainfall intensity thresholds (Mohr et al., 2013). This is not surprising, since they remain bare and prone to compaction by heavy machinery transit. These roads often intersect streams, which form bypasses to preferentially route sediment (Fig. 2), increasing the efficacy of mass transfer within a geomorphic system, or sediment connectivity (Wohl et al., 2019). In this case, road networks modify the pathways of runoff and sediments, and may also modify thresholds of rainfall to trigger sediment detachment and transport (for example, due to soil compaction), potentially affecting the structural and functional components of sediment connectivity, as defined by Wainwright et al. (2011). This shift is also relevant to constraining off-site impacts of soil erosion (Boardman et al., 2019). The erosional work efficacy depends on the logging season, which is higher for wet season logging (Mohr et al., 2014). At the storm to yearly




Figure 2. Details of forest roads in the Purapel catchment under different stages of the tree plantation rotational cycle and their connection to streams. A-D Google images (©Google Maps 2016, ©Google Maps 2021, ©Google Maps 2022, ©Google Earth 2016, respectively), E-F pictures of a gravel road and its connection to a stream during a storm in July 2017.

scale (10^{-4} - 10^0 yr), roads are prime sources and routers of sediments in catchments covered by tree plantations (Schuller et al., 2013).

Despite the increase in structural connectivity, sediment mobilization depends mostly on specific thresholds of rainfall. For example, hydrologic connectivity to initiate runoff in recently logged areas required a threshold of 20 mm/hr in rainfall simulations on tree plantations near Nacimiento (Mohr et al., 2013). In the absence of long term records of rainfall intensities, hydro climatic trends on rainfall and streamflow are relevant to interpret catchment erosion. In Central Chile (30-39°S), rainfall decreased at ca. 4% per decade between 1960 and 2016 (Boisier et al., 2018), culminating in an unprecedented megadrought starting 2010 (Garreaud et al., 2020).

While the erosional response of logging is largely indisputable, hydrologic responses to tree harvest are ambiguous. On the one hand, logging may increase streamflow discharge in general and peak flow in particular (Iroumé et al., 2006). On the other hand, logging may also decrease streamflow discharge due to enhanced groundwater recharge immediately after logging (Mohr,




Figure 3. Annual and seasonal rainfall and annual aridity index (AI=P/PET) at Purapel catchment. Main monotonic trends are tested with Mann-Kendall and LOWESS smoothing for 1962-2015 (blue), 1979-2019 (red), 1986-2018 (purple) and 2000-2018 (green). Unfilled circles are discarded data. A. Annual rainfall and AI time series. B. Seasonal time series for Nirivilo station.

2013). The distinct responses may most likely vary with tree species and age, harvest size, forestry treatment (thinning, clear cutting, replanting), riparian buffer width, and especially, with the moisture storage decrease under recent drought conditions, which exacerbated declines in runoff (Iroumé et al., 2021).

In addition to the megadrought, recent increase in both magnitude and frequency in wildfire affects relatively more tree plantations compared to alternative land cover (Bowman et al., 2019). This is likely because fuel is more abundant under dense plantation cover that connect large continuous tracts of the landscape. Instead, native species are more patchy (Gómez-González et al., 2017, 2018).

While the observed disturbances affecting the vegetation cover commonly increase sediment yields in rivers (e.g., Reneau et al., 2007; Brown and Krygier, 1971), the long and persistent decline in rainfall (Méndez-Freire et al., 2022; Tolorza et al., 2019) together with the high water demands of tree plantations is expected to reduce sediment detachment and mobilization assuming fluvial transport-limited conditions. To evaluate the impacts of these opposite responses and their potential effects in land degradation, we explore the catchment scale erosion of the Purapel river (406 km² of drainage area). To this end, we combine two distinct temporal scales (10⁴ and 10¹ yr), explore discrete disturbance events (2017 and 2015 wildfire, wet-season logging), and calculate sediment connectivity associated with forestry roads. In a recent analysis of the same suspended sediment data, Pizarro et al. (2023) concluded that the concomitant afforestation and the decrease in sediment discharge of




Figure 4. Mean seasonal streamflow, baseflow and suspended sediment concentrations at “Purapel en Sauzal” station. Main monotonic trends are tested with Mann-Kendall and LOWESS smoothing for 1986-2018 (purple) and 2000-2018 (green). Unfilled circles are discarded data. A. Streamflow at Purapel en Sauzal station. B. Baseflow at Purapel en Sauzal station. C. Suspended sediment concentrations at Purapel en Sauzal station.




Figure 5. Denudation and specific SSD at Purapel en Sauzal gauge. Distributions of SSD for individual hydrologic years (March to Feb). Purple circles show percentiles (0.05, 0.25, 0.5, 0.75 and 0.95) and purple triangles show the mean. Filled symbols represent years with more than 185 daily data. Catchment erosion/denudation rates are indicated in red. Solid line is the sediment yield equivalent to the ¹⁰Be denudation rate, dashed line is the average of all suspended sediment records.




Figure 6. Daily hydrometric data for pre- and post- fire hydrologic years for the two large wildfire events. Rainfall, streamflows, SSC and SSD of pre- and post-fire hydrologic years were contrasted in homoscedasticity with the Fligner-Killeen test using an $\alpha = 0.05$. Heterocedastic distributions were compared with Welch t-test. Homocedastic distributions were compared with Wilcoxon-Mann-Whitney test.

the catchment could be interpreted as a causal relationship. Here, we analyze a wide range of meteorological and vegetational indexes over the whole record period and we show that factors different from afforestation can explain the observed decline in sediment discharge, in particular the drought that this catchment has experienced since 2010.

2 Materials and methods

2.1 The Purapel catchment

The Purapel river drains the eastern flank of the CCR. The climate is Mediterranean type. Mean annual rainfall is 845 mm, mean minimum and maximum air temperatures are 7.2 and 20.3°C, respectively, and the flood regime is exclusively pluvial bound (Álvarez-Garretón et al., 2018). The catchment is 406 km² and dominated by metamorphic (47.5%) and granitic (44.3%) lithologies. Elevation ranges between 164 and 747 m a.s.l. Most hillslopes are gentle (hillslope gradients around 16%), largely convex, and incised by gullies that converted this landscape into badlands (Fig. 1). CIREN (2021) classified most of those hillslopes as severely affected by soil erosion. The dominant soil types are Inceptisols and Alfisols (Bonilla and Johnson,

Table 1. Published and new detrital ^{10}Be denudation rates in the Mediterranean CCR. Denudation rates and their uncertainties were calculated with procedures described in Carretier et al. (2018). Characteristic time refers to the quartz residence time within a particle mean free path in rocks of 60 cm, and represent a timescale for steadily erosion (Lal, 1991). ^{10}Be concentrations and their uncertainties were analyzed in the CEREGE laboratory with the indicated standard.

Name	Denudat. rate (mm/yr)	Denudat. rate unc. (mm/yr)	Char. time (kyr)	Lat	Lon	Catch. area (km ²)	Analyzed grain size (mm)	[¹⁰ Be] (at/g)	[¹⁰ Be] unc. (at/g)	Standard material	Source
SAUR	0.024	0.004	25	-35.6197	-72.0171	406	[0.5,1]	143751	5469	STD-11	This work
Mi2	0.037	0.006	16	-37.0488	-72.8614	235	[0.5,1]	93772	4280	4325	Carretier et al. (2018)
C3-2	0.039	0.007	15	-37.4052	-72.7976	357	[0.5,1]	97896	8272	4325	Carretier et al. (2018)
C1-1b	0.041	0.010	14	-37.4652	-72.7495	739	[0.5,1]	113680	20735	4325	Carretier et al. (2018)

2012). Soil properties are highly variable in space. Yet, soils under tree plantations are generally thinner, more depleted in soil organic matter and with lower invertebrate diversity than soils under native forest (Cifuentes-Croquevielle et al., 2020; Soto et al., 2019). Throughout the entire soil profile, the soil bulk densities of Eucalyptus (1.38 ± 0.08 to 1.58 ± 0.12 g/cm³) and Pine (1.28 ± 0.18 to 1.53 ± 0.13 g/cm³) stands are higher than under native forests (0.89 ± 0.27 to 1.25 ± 0.24 g/cm³) for depths between 0 to 60 cm (Soto et al., 2019).

2.2 Hydrometeorologic data and analysis

Daily rainfall and potential evapotranspiration data (1979-2020) for the Purapel catchment were downloaded from CAMELS-CL dataset (<https://camels.cr2.cl/>). Daily rainfall data are derived from the CR2METv2 precipitation product, which merges the ERA5 reanalysis, local topographic data, and is calibrated with an updated national rain-gauge network (DGA, 2017). In addition, longer rainfall daily time series (1956-2019) at the Nirivilo rain gauge were downloaded from Mawüm (<https://mawun.cr2.cl/>). We analyzed trends on annual and seasonal rainfall for several periods using both in situ (Nirivilo rain gauge) and gridded (CR2MET) data. We also calculated annual aridity trends using the Aridity Index (AI), computed as the annual rainfall divided by annual potential evapotranspiration ($AI=P/PET$) using CAMELS-CL data. For annual analysis we included years with less than 330 data points and all months with more than 27 data points. For seasonal analysis (~90 days) we included data for seasons with more than 60 daily data (more than ~66.6% of data). Seasons considered here were Autumn (MAM), Winter (JJA), Spring (SON), and Summer (DJF). For the pre- and post-fire hydrologic years (Mar-Feb) we analyzed daily rainfall too. To this end, we used the gap-free gridded CR2METv2 product.

Streamflow and suspended sediment data are available in the Chilean General Directorate of Water (DGA) site (<https://snia.mop.gob.cl/BNAConsultas/reportes>). The DGA estimated daily streamflows from single gauge stage readings using calibrated rating curves. Roughly once a month, the rating curves are updated by manual current meter measurements. The suspended sediment concentrations (SSC) were sampled on a daily-basis, too. All samples were obtained close to the water surface in the vicinity of the water stage. The samples were filtered using a cotton linter cellulose paper with 80% of collection efficiency for

particles larger than $0.3\ \mu\text{m}$ (Advantec Qualitative Filter Papers 2, written communication from DGA operator). Then, they were dried, and burnt for 2 hours at $550\text{--}600^\circ\text{C}$ in DGA laboratories (Solar, 1999).

We calculated daily suspended sediment discharge (SSD, t/day) as the product of streamflow discharge (m^3/s) and SSC (mg/l), assuming those measurements as representative of the entire day (Pepin et al., 2010). This assumption is based in the catchment size and measurements of streamflow velocities we had performed in this landscape (supplemental material S1). In addition, we calculated the number of data, the percentiles and the mean value of SSD for single hydrologic years. We calculated the daily baseflow at Purapel en Sauzal station with the Lyne and Hollick filter (Ladson et al., 2013), which is a standard approach used in several studies (e.g. Li et al., 2022; Huang et al., 2021; Teutschbein et al., 2015; Zhang et al., 2017). For the baseflow separation we used several α values between 0.5 and 0.95 and $n.\text{reflected}=30$ days as parameters.

The streamflow and SSC time series contain gaps, which are not seasonally clustered. In particular, gaps during the dry season are mostly related to ceased streamflow (personal communication from DGA operator). We calculated the mean value and the number of daily streamflow, baseflow and SSC data in each month and season. Based on the number of data (supplemental material S2), we discarded the monthly analysis of trends. Then, we calculated seasonal trends on streamflows and suspended sediments for seasons with more than 60 data points. All trends are computed with the Mann-Kendall test (Helsel et al., 2020) and plotted with an additional LOWESS smoothing (Cleveland, 1981). In addition, we computed quantiles and averages of SSD for hydrologic years with more than 185 data points.

We also examined the existence of differences between the daily data for the hydrologic years before and after the fire, i.e., 2014 and 2016 versus 2015 and 2017, respectively ($n>262$ for each group of data). Given the non-normal distribution of all hydrological datasets, we used the Wilcoxon-Mann-Whitney test instead of the traditional t-test. That test requires the two compared populations have the same shape and variance (homoscedasticity). We used the non-parametric Fligner-Killeen test to judge whether our samples were homoscedastic (p-value larger than 0.05) or not (p-value lower than 0.05). If the compared populations had unequal shape or variance (heteroscedasticity), we used the Welch t-test (Skovlund and Fenstad, 2001).

2.3 Catchment-wide erosion and denudation rates

We obtained catchment-wide erosion rates for Purapel river at the gauge “Río Purapel en Sauzal” using two approaches for different time scales, short-term (decadal) from suspended sediments and long-term (10^3 to 10^4 yrs) from detrital ^{10}Be . We calculated the long-term denudation rate as a benchmark to compare the recent sediment yields against. In most fluvial catchments the long-term rates exceed the short-term rates (Covault et al., 2013). This picture, however, may flip vice versa if soil erosion is high (Hewawasam et al., 2003; Vanacker et al., 2007). A limitation of our approach is the fact that detrital ^{10}Be rates include physical erosion and chemical weathering rates (von Blanckenburg and Willenbring, 2014), while suspended sediment yields account only for physical erosion of very fine sediment (Summerfield and Hulton, 1994), which excludes bedload and dissolved load. Thus, we regard our short-term erosion rates as minimum rates for landscape lowering.

For the short-term, we calculated the mean specific SSD ($\text{t}/\text{km}^2/\text{yr}$) as the average of all records (06/1985 to 11/2018) on a yearly scale and normalized by catchment area (Pepin et al., 2010). We estimated resulting erosion rate (mm/year) assuming a mean soil bulk density of $2.6\ \text{g}/\text{cm}^3$ (Carretier et al., 2018).

160 For the long-term, we assume the ^{10}Be concentrations within fluvial sands are proportional for catchment-wide averaged denudation rate (von Blanckenburg, 2005; Granger and Schaller, 2014). This rate integrates over a characteristic timescale that is inversely proportional to the denudation rate. These timescales are commonly longer than 10^3 years (Covault et al., 2013). We therefore regard the ^{10}Be derived rates as a reference that largely excludes recent human disturbances but includes low frequency and high magnitude erosion events (Kirchner et al., 2001). We obtained a bulk sample of fluvial sands from the
 165 active river bed along a cross section close to the water stage “Río Purapel en Sauzal”, collecting sands from the surface at three locations within ~ 10 m distance. We mixed all samples and sieved to a grain size fraction 0.5-1 mm.

The mixed sand sample was processed at the French AMS ASTER facility in CEREGE (Standard STD-11). In order to convert the ^{10}Be concentration C into catchment mean denudation rate, we neglect radioactive decay and assume steady state of ^{10}Be concentration, leading to the classical following equation

$$170 \quad \epsilon = \frac{1}{\rho C} P_{SLHL} (f_{sp} S_{sp} \Lambda_{sp} + f_{sm} S_{sm} \Lambda_{sm} + f_{fm} S_{fm} \Lambda_{fm}) \quad (1)$$

where $P_{SLHL} = 4$ at/g/yr is the sea-level-high-latitude total production rate of the considered nuclide. f_{sp} , f_{sm} and f_{fm} are the fractions of this production rate due to spallation, slow muons capture and fast muons averaged over the catchment area, respectively (Braucher et al., 2011). S_{sp} , S_{sm} , S_{fm} are scaling factors depending on latitude and elevation averaged over the catchment area (Stone, 2000), and $\rho = 2.6$ g/cm³.

175 2.4 Land cover changes

The Purapel catchment has experienced high rates of LULCC since the 19th century. This was largely due to the extensive increase in wheat production caused by the gold rushes in California and Australia (Cortés et al., 2022). Later on, between 1955 and 2014 tree plantations increased from (a minimum of) 10.27 (Hermosilla-Palma et al., 2021) to 203.5 km² (Zhao et al., 2016). Recently, two large wildfires burned the catchment: In 2015 14% of the catchment area burned. In 2017 almost
 180 the entire catchment burned (95%) (Tolorza et al., 2022).

To describe recent LULCC in this catchment, we use land cover maps both from compiled sources (1955, 1975 and 2017) and from our own (1986, 2000, 2005, 2010 and 2015):

- The 1955 and 1975 land cover maps of Hermosilla-Palma et al. (2021) cover the headwaters of the Purapel catchment (157 km²). These maps were made interpreting the land cover from the 1:70,000 aerial photograph (Hycon flight) for
 185 1955, and from the 60 m resolution Landsat-2 MMS and the 1:30,000 aerial photographs of 1978 (CH-30 flight) for 1975.
- We used Landsat Surface Reflectance products to identify land cover classes during dry seasons of 1986, 2000, 2005, 2010 and 2015. We classified unburned land cover using the Maximum Likelihood Classifier (Chuvieco, 2008) which we trained and validated with 20 and 10 polygons for each class, respectively. We validated the results with field observations
 190 during 2014-2015. We sub-classified burned surfaces into low, moderate and severe fire according to the differences in

NBR index of pre- and post- fire images (thresholds <0.1 - 0.269>, <0.27 - 0.659>, <0.66 - 1.3> Key and Benson, 2006).

- The Land cover map of 2017 was made by Tolorza et al. (2022) with pre-fire Sentinel and LiDAR data. Here, this classification was resampled to 30 m resolution, to be compatible with LANDSAT classifications.

195 2.5 Logging roads and sediment connectivity

To identify changes in the structural connectivity we applied the Connectivity Index (IC , dimensionless) using the weighting factor (W , dimensionless) of (Cavalli et al., 2013). IC is a semi-quantitative approach to describe the degree of coupling between hillslopes and a target (for example, the stream network):

$$IC = \log_{10} \left(\frac{\overline{W} \overline{S} \sqrt{A}}{\sum_i \frac{d_i}{W_i S_i}} \right) \quad (2)$$

200 , where \overline{W} and \overline{S} (m/m) are the average weighting factor and slope gradients on the upslope contributing area (A , m²), respectively. d_i (m), W_i (dimensionless) and S_i (m/m) are the path length, the weighting factor and the slope gradient on the i th cell in downslope towards a target.

W is calculated from a DTM to account for the effect of topographic roughness. The Roughness Index (RI , m) is the standard deviation of the residual topography. The residual topography refers to the difference between the original DTM and
205 a smoothed version obtained by averaging DTM values on a 5×5 (=25) cell moving window:

$$RI = \sqrt{\frac{\sum_i^{25} (x_i - x_m)^2}{25}} \quad (3)$$

, where x_i (m) is the value of one specific cell of the residual topography within the moving window, and x_m (m) is the mean of all 25 window cells. The weighting factor is calculated as:

$$W = 1 - \frac{RI}{RI_{max}} \quad (4)$$

210 , where RI_{max} is the maximum value of RI in the study area.

We quantified changes in sediment connectivity due to the forestry roads, RC , as

$$RC = IC_{rs} - IC_s \quad (5)$$

, where the subscripts s and rs refer to the stream network and to the stream network including roads. We fed the model with a mapped forestry road network obtained from images available in the OpenLayers plugin of QGIS and post-2017-fire
215 Sentinel compositions.

2.6 Disturbances in vegetation

We used the Breaks For Additive Season and Trend algorithm (BFAST, Verbesselt et al., 2010) on a LANDSAT collection to detect disturbances in vegetation at the pixel scale, i.e. $\geq 30\text{m}$. In the Purapel catchment, disturbances $> 30\text{ m}$ are mostly due to wildfires and/or clear-cuts. Such disturbances lean on the seasonal behavior of the NDVI index on a time series of LANDSAT surface reflectance (Level 2, Collection 2, Tier 1) for the period from 09/1999 to 10/2021. Clouds were filtered using the QA band which uses the CFMask algorithm (Foga et al., 2017). We used the same parameter set as Cabezas and Fassnacht (2018), namely the threshold value for disturbances set to 93 manually labeled reference polygons with fire events, clear-cuts and constant tree-cover. It is worth mentioning, that we applied a sieve filter to the results. Hence, only disturbances greater than 1 ha were considered. We trained the algorithm with the Landsat time series of 1999 to 2001. Given the disturbance regime of Purapel catchment (two large wildfires and possible loggings each 9 to 25 years) we run BFAST anticipating three possible breaks for the period 2002-2021. The accuracy assessment was performed on 35 manually drawn polygons that were randomly distributed across the catchment.

3 Results

3.1 Hydro climatic trends

At the annual scale at Nirivilo rainfall station, most data of the period 1962-2015 passed our completeness assessment criteria (53 of 54 years). In the case of CR2MET, the longest period analyzed here is 1979-2019. Judging Mann-Kendall tests and LOWESS smoothing, we did not find a single trend for the longest interval of records (1962-2015). Nevertheless, for the period after 1979, we obtained decreasing non-monotonic trends for rainfall, both at Nirivilo station and for the CR2MET product. That decrease is steeper for 2000-2019, but less pronounced for intermediate intervals such as 1986-2018. During 1986-2018, however, a decrease in seasonal rainfall is observed for Autumn, at the beginning of the hydrologic year (Fig. 3). Generally, the AI follows similar decreasing trends as is the case for rainfall, thus indicating persistently dry conditions across this catchment. For only 2 years (1982 and 2002) the AI was higher than 1. During all other years, potential evapotranspiration was greater than rainfall.

Streamflow data is available at Purapel en Nirivilo between 1979 and 2019 and at Purapel en Sauzal between 1981-2019. For Purapel en Sauzal streamflow data, results of baseflow separation are in the supplement material 3. We selected the results obtained with $\alpha=0.7$ for further trend analysis, given the observed magnitudes and shape of the baseflow time series. For the Suspended Sediment Concentration data, the longest period is 1985-2018. The seasonal analysis for “Purapel en Sauzal” station is in Fig. 4. Although none of those time-series is monotonic, the sharp decrease in suspended sediment concentrations is clear for the three wetter seasons (Autumn, Winter and Spring).

245 3.2 Catchment-wide erosion and denudation rates

^{10}Be denudation rate resulted in 0.024 ± 0.004 mm/yr (Table 1), assuming a soil particle density of 2.6 t/m^3 . This rate translates into a specific sediment yield of $62.4 \pm 10.4 \text{ t km}^{-2}\text{yr}^{-1}$. This rate integrates over a characteristic timescale of ~ 25 kyrs. Together with published data, detrital ^{10}Be denudation rates in the CCR point to between 0.02 and 0.05 mm/yr (table 1 Carretier et al., 2018).

250 Given the data completeness test, we assume the decadal catchment-wide erosion rate from suspended sediments to be a conservative estimate. Following Pepin et al. (2010) we calculated the mean specific SSD for all the records between 1985 and 2018 and a 30% of error (Pepin et al., 2010). For the Purapel catchment we estimate $47 \pm 14.1 \text{ t km}^{-2}\text{yr}^{-1}$, equal to 0.018 ± 0.005 mm/yr, assuming the same soil bulk density. Both rates do not statistically differ (Fig. 5).

3.3 Decennial trends in SSD and comparison of pre- and post-fire hydrometric data

255 On a decennial time-scale, we observe decreasing trends in mean and high (p95 and p75) annual values of SSD, judging from Mann-Kendall p-values for $\alpha = 0.05$ (Fig. 5). Since 2010, high and medium percentiles of SSD are lower than most of the previous hydrologic years. Only the lower percentiles of SSD revert the decreasing trend at the end of the time series. Such behavior corresponds to baseflow conditions.

On a shorter time-scale, in Fig. 6 we compare pre- and post-fire hydrometric data for the 2015 and 2017 wildfires. Compared
260 series resulted to be homoscedastic for rainfall in the two comparisons, and for SSC in the comparison before and after the 2015 wildfire. No changes in median rainfall values can be interpreted from the large p-value of their Wilcoxon-Mann-Whitney tests. Before and after the 2015 wildfire, judging from p-values for $\alpha = 0.05$ on Welch t-test or Wilcoxon-Mann-Whitney tests, mean streamflow diminished, but median SSC and mean SSD increased. Before and after the 2017 wildfire, mean streamflow increased, as well as mean SSC and mean SSD.

265 3.4 Recent land cover changes

We developed five land cover maps for the period 1986-2015. The overall classification accuracy ranged between 83% and 92%. We distinguished between tree plantations, native forests, shrublands and seasonal grasslands. Seasonal grasslands included bare surfaces, seasonal pasture and sparse vegetation. We also classified seasonal grasslands to separate recently logged areas (clear-cuts) from other poorly vegetated areas.

270 Fig. 7 shows that the upper catchment was covered by a minimum of 1,000 ha of tree plantations and 5,500 ha of shrublands in 1955 (Hermosilla-Palma et al., 2021). Between the 1980s and the beginning of 21st century, the most prominent change comprised the transition from seasonal grasslands and shrublands into tree plantations. The first two classes covered a minimum of 23,550 ha in 1986 and 13,050 ha on 2005. During the same period, tree plantations expanded from 8,090 to 20,980 ha. Between the wildfires of 2015 and 2017, seasonal grasslands and shrublands together expanded to $\sim 20,300$ ha (Fig. 7).

The result of our mapped road network is illustrated in Fig. 1. Using this road network on a 5 m resolution LiDAR, we estimate some 18,000 ha of increased sediment connectivity, resulting in $RC > 0$ (Fig. 8). RC values exceeding the 95-percentile (> 3.12) are 1,986 ha. That area of high RC is mostly located on hilltops: 1,966 ha (i.e. 99%) and resulted in upstream contributing area < 1 ha. Particularly these topographic settings exceed an empirical threshold between high and low connectivity
 280 for a mountain catchment, i.e. -2.32 (Martini et al., 2022). In the Purapel catchment the area above this threshold increased from 1,120 to as much as 6,570 ha simply due to the dense road network. This quantification, however, is done with a digital terrain model of coarser resolution compared to the original study of Martini et al. (2022) (5 m vs 0.5 to 2.5 m).

Based on our BFAST modeling, we obtained monthly time series of disturbances for 2002-2019 that we aggregate to the seasonal scale. We achieved a confusion matrix with a balanced accuracy of 0.86 and a F1 score of 0.69. For the complete
 285 period (Fig. 9A) 13,640 ha of the Purapel catchment (33.7%) experienced one break in the NDVI time series, 16,810 ha (41.5%) showed two breaks and 5,010 ha (12%) presented three breaks. The undisturbed 12.8% included tree plantation stands that remained unlogged, and seasonal grasslands that remained poorly vegetated. Considering the seasonality (Fig. 9B), the periods with largest disturbed areas were the summers (dry season) of 2015 and 2017 due to the wildfires. Both wildfires were detected in $\sim 5,000$ and $24,000$ ha, respectively. The periods that follow in disturbed area were the Autumn (wet season) of
 290 2002 ($\sim 2,000$ ha) and 2007 (1,910 ha). The largest surface disturbed during a Winter and a Spring were 770 ha each in 2006 and 2009, respectively.

Compared to the dNBR classification for 2017 (Tolorza et al., 2022), the BFAST results detected less burned areas for the 2017 wildfire (33,618 vs 24,299 ha). This difference could be explained by the better capabilities of the dNBR index to detect burned areas, since it is a dedicated method to classify burned surfaces based in the NBR index of a pre- and a post- fire image
 295 (Key and Benson, 2006), while the BFAST algorithm was applied here on the NDVI, which is a index more suitable to detect the density of vegetation, and thus more sensitive to clear cuts.

4 Discussion

Both ^{10}Be denudation rate and suspended sediment erosion rate are surprisingly similar (Fig. 5). Hence, we argue that the suspended sediment samples capture at least the effects of erosion events recorded on the long-term. Both rates are low for
 300 fluvial catchments between $100\text{--}1,000\text{ km}^2$ (Covault et al., 2013). Yet, those rates are similar to 3 tributaries of the Biobío river draining the eastern CCR, which are between 0.037 ± 0.006 and $0.042 \pm 0.008\text{ mm/yr}$ (Carretier et al., 2018). The low ^{10}Be denudation rate agrees with a landscape dominated by slow soil creep with occasional mass wasting triggered by intense rainfall events and earthquakes, yet there were only two reported landslides after the 2010 within the catchment area (Serey et al., 2019) with no effects in suspended sediments (Tolorza et al., 2019). Short term erosion does not exceed the long term
 305 denudation, as in other highly human-disturbed catchments (Hewawasam et al., 2003; Vanacker et al., 2007). Considering the numerous gaps on streamflow and suspended sediments, and the absence of sub-daily or depth-integrated measurements of sediment concentrations, we regard the decadal sediment data as a conservative estimate for recent catchment erosion. This

was also reported for suspended sediments from other rivers of the western Andes (Vanacker et al., 2020; Carretier et al., 2018). In addition, suspended sediments do not record the effects of chemical weathering on denudation rates. This process
310 seems to be relevant in the CCR: in the absence of spatially resolved data of regolith thickness, single observations suggest thick saprolite layers (at least) locally (Vázquez et al., 2016; Mohr et al., 2012; Krone et al., 2021). Thus, depending on the magnitude of mass loss due to chemical weathering, which ranges between 0 and $\sim 240 \text{ t/km}^2/\text{a}$ in other latitudes of the CCR (Schaller and Ehlers, 2022), total denudation in the short term can be equal or even higher than the long term denudation. Yet, we do not have quantitative estimates of local chemical weathering and soil production rates to test that hypothesis.

315 The Purapel river catchment has been a staging ground for rapid expansion of tree plantations and a number of disturbances during the period of suspended sediment monitoring. This landscape has been affected by clear-cuts, two widespread wildfires and one Mw 8.8 earthquake. The expansion of tree plantations was mostly at the expense of poorly vegetated surfaces (Fig. 7). Yet, their management includes extensive logging operations (supplementary videos) – mostly during wet seasons (Fig. 9) – and the construction and maintaining of forestry roads used by heavy machinery. The distribution and density of a road
320 network by itself means an increase in structural sediment connectivity (Fig. 8). Higher connectivity facilitates the routing of detached soils, even from hilltops, where soil production rate is slower compared to the mid-slope or toe positions across the CCR (Schaller and Ehlers, 2022). Thus, hilltop soils may be more difficult to recover during human time scales. The increase in sediment connectivity is distributed along all the hillslopes and more than the half of the catchment experienced at least 2 disturbance events between 2002 and 2019. Here, we emphasize that the connectivity index may be a minimum estimate, as
325 we used a coarser digital terrain model compared to the original study of Martini et al. (2022).

Despite the disturbances, the decennial trends in mean and high annual SSD (Fig. 5) and the mean SSC during wet seasons (Fig. 4) are decreasing. After the 2015 wildfire, SSC increased respect to the previous year, while the mean streamflow decreased (Fig. 6). There we can unambiguously interpret a response of the erosion to the wildfire. Nevertheless, the described changes in pre- and post-fire suspended sediments are limited, since high and medium percentiles of SSD after the fire are
330 lower than most of the hydrologic years previous to 2010, the beginning of the megadrought (Fig. 5).

If the suspended sediment record is representative of the sediment yields on Purapel river, the disturbance regime contrasts with expected responses in sediment mobilization, given observations reported in other landscapes (e.g., Reneau et al., 2007; Brown and Krygier, 1971). Nevertheless, the arid conditions, i.e. the AI ratio between annual precipitation and evapotranspiration, indicate increasingly scarce water. A decrease can be also interpreted for the streamflow and the baseflow of the wet
335 seasons, mostly in the Autumn. The sediment detachment and transport likely coincide with these negative trends. Sediment mobilization both on hillslopes and streams depends mostly on specific thresholds of rainfall intensity and water discharge, while the unprecedented drought starting in 2010, together with high root water uptake by fast-growing tree plantations resulted in a reduction in water availability. In this scenario, a lack of minimum rainfall intensity required to trigger runoff and soil erosion on hillslopes (Mohr et al., 2013) and/or an increase in the residence time of sediments stored within the valleys is
340 plausible. As rainfall and direct runoff control sediment fluxes at the catchment scale (Andermann et al., 2012; Tolorza et al., 2014), sediment mobilization under the current hydrological regime may stay low despite landscape disturbances. A recent model in post-fire sediment cascades indicates that, even when post-fire erosion may be severe in source areas, a substantial

fraction of the detached sediment load may (intermittently) remain stored within valleys with only moderate delivery to the river network (Murphy et al., 2019). Assuming transport limitation under the current drought conditions, prolonged residence times of sediments are also expected. Indeed, both tree plantations and drought conditions reduced the recharge to deep soil water reservoirs (Iroumé et al., 2021; Huber et al., 2010). Also the loss of soils due to erosion may further reduce the water-storage capacity (Ratta and Lal, 1998). The long deficit of water due to the drought and the tree plantations may reduce groundwater storage, which is consistent with the observed negative trend in baseflow. Such sharp reduction in water availability may limit the sediment transport in channels. The possible increase of the sediment transport only for the lower percentiles supports the notion that sediment transport is largely restricted to baseflow conditions during the study period. Hence, we cannot unambiguously quantify the overall effect of landscape disturbances on sediment fluxes. Sediment fluxes are more efficient during periods of high flows which correspond to wetter conditions (e.g., Mohr et al., 2013). Consequently, the sediment stored in the valleys, highly rich in nutrients and carbon, can be re-suspended during higher discharge events, causing temporarily delayed off-site problems for several decades to come.

The expansion of tree plantations has been proposed as a tool to mitigate soil erosion (CONAF and MINAGRI, 2016). Recently, plantations have been favored as a better solution to mitigate soil erosion compared to native forests for the same Purapel catchment (Pizarro et al., 2020, 2023). At Purapel catchment, a direct comparison between native forest and plantations cannot be done for the period 1986-2018, because the major land cover transition was from poorly vegetated surfaces to tree plantations (Fig. 7). Nevertheless, we can discuss whether the observed land management is a suitable solution for soil erosion mitigation in the CCR. There is abundant evidence of increased soil erosion in Chilean tree plantations, such as truncated soil profiles in an eucalyptus stand at 36°37'S (Banfield et al., 2018), a fourfold increase in net soil loss under pine stands relative to native forest at Talcamavida (37°7'S) and Nacimiento (37°30'S) (Aburto et al., 2020) or changes in nutrient cycles and increased sedimentation rates in coastal lakes, such as Matanza (33°45'S, Fuentealba et al., 2020), Vichuquén (34°S, Fuentealba et al., 2021), San Pedro (36°51', Cisternas et al., 2001), and Lanalhue (37°S, Alaniz et al., 2021). Based on such strong empirical evidence along CCR and our own results (Fig. 2, 7, 8 and 9), we argue that the observed ongoing forest management of tree plantations promotes soil erosion and landscape degradation. In addition, soils in tree plantations are depleted in carbon and nutrients (Soto et al., 2019; Banfield et al., 2018), and inhibit lower invertebrate diversity (Cifuentes-Croquevielle et al., 2020) compared to soils under native forest. As a result, C and N stocks are relatively lower in tree plantations up to deep soil compartments (>120 cm) (Crovo et al., 2021). Soil organic matter is a key component for soil formation (Bernhard et al., 2018). For that reason alone, native forests rather than exotic tree plantations are a more appropriate land cover to regenerate soils and reverse or, at least, decelerate 200 years of intense soil erosion. Indeed, the protection and conservation of natural vegetation has the strongest effect on improving soil quality after water erosion (Vanacker et al., 2022). Also, empirical restoration examples available show that the transition from former Eucalyptus plantations to native forest is promising in terms of improving water availability (Lara et al., 2021).

The Purapel catchment, as other similar catchments along the CCR, denudates slowly on scales of 10^3 to 10^4 years. The averaged suspended sediment discharge is similar in magnitude, although likely underestimating total denudation. The Purapel catchment, as other similar catchments along the CCR, denudates slowly on scales of 10^3 to 10^4 years. The averaged suspended sediment discharge is similar in magnitude, although likely underestimating total denudation. Then, depending on the magnitude of the unmeasured portion of the denudation, decadal lowering of Earth Surface may be equal or even higher than the long-term average.

Suspended sediment transport decreased during the wet seasons between 1986 and 2018, which, at first glance, conflicts with the disturbances observed in vegetation, especially the intense and widespread wildfires. The decrease in several hydroclimatic variables, including baseflow and aridity, coincides with lower suspended sediment loads. We argue that the low range of recent suspended sediment discharge resulted from limitations in the detachment and transport of sediments due to the observed water decline. Or in other words: the drought offsets the effects of disturbances and higher connectivity. Without sufficient water, the residence times of sediments are long, despite the increased sediment connectivity on hillslopes. The contribution of tree plantations to reduce erosion, if any, is more related to their impact in water availability than directly in soil protection.

Because the surface lowering in the last three decades is similar to or higher than the long-term benchmark, and those measures are spread along a specific dry period, we argue it may be considered high for this specific system. That conclusion and the documented effects of tree plantations on SOC and soil biodiversity are clear indicators of a degrading landscape.

Code availability. R scripts used in this study for data analysis are accessible upon request by contacting Violeta Tolorza (violeta.tolorza@ufrontera.cl)

Data availability. Supplemental data-sets related to this submission are available at:

395 <https://doi.org/10.5281/zenodo.6958544>

<https://doi.org/10.5281/zenodo.6974312>

<https://doi.org/10.5281/zenodo.7328071>

Video supplement. Time-lapses showing disturbances in vegetation are available at:

<https://doi.org/10.5446/62703>

400 <https://doi.org/10.5446/62704>

Author contributions. VT: Conceptualization, funding acquisition, project administration, methodology, field work, validation, investigation, writing - original draft; CHM: conceptualization, methodology, investigation, writing - deep review and editing; MZB, SC, MG and OS: methodology, investigation, writing - review and editing, BS and DP: field work, validation, writing - review and editing

Competing interests. The contact author has declared that none of the authors has any competing interests.

405 *Acknowledgements.* This paper arises mostly from research funded by ANID-FONDECYT project 11190864 and the UFRO Postdoc grant VRIP20P001, and received contributions from ANID-FONDAP project 15110009 and DFG project 493703771. We thanks to Claudio Ramirez Bravo (DGA) for describing details of hydrometric monitoring in Purapel river, and for provide the actual location of the gauge *Río Purapel en Sauzal*.

References

- 410 Aburto, F., Cartes, E., Mardones, O., and Rubilar, R.: Hillslope soil erosion and mobility in exotic pine plantations and native deciduous forest in the coastal range of south-central Chile, *Land Degradation & Development*, p. ldr.3700, <https://doi.org/10.1002/ldr.3700>, 2020.
- Alaniz, A. J., Abarzúa, A. M., Martel-Cea, A., Jarpa, L., Hernández, M., Aquino-López, M. A., and Smith-Ramírez, C.: Linking sedimentological and spatial analysis to assess the impact of the forestry industry on soil loss: The case of Lanalhue Basin, Chile, *Catena*, 207, <https://doi.org/10.1016/j.catena.2021.105660>, 2021.
- 415 Álvarez-Garretón, C., Mendoza, P. A., Boisier, J. P., Addor, N., Galleguillos, M., Zambrano-Bigiarini, M., Lara, A., Puelma, C., Cortes, G., Garreaud, R., McPhee, J., and Ayala, A.: The CAMELS-CL dataset: catchment attributes and meteorology for large sample studies – Chile dataset, *Hydrology and Earth System Sciences Discussions*, pp. 1–40, <https://doi.org/10.5194/hess-2018-23>, 2018.
- Andermann, C., Crave, A., Gloaguen, R., Davy, P., and Bonnet, S.: Connecting source and transport: Suspended sediments in the Nepal Himalayas, *Earth and Planetary Science Letters*, 351–352, 158–170, <https://doi.org/10.1016/j.epsl.2012.06.059>, 2012.
- 420 Armesto, J. J., Manuschevich, D., Mora, A., Smith-Ramírez, C., Rozzi, R., Abarzúa, A. M., and Marquet, P. a.: From the Holocene to the Anthropocene: A historical framework for land cover change in southwestern South America in the past 15,000 years, *Land Use Policy*, 27, 148–160, <https://doi.org/10.1016/j.landusepol.2009.07.006>, 2010.
- Banfield, C. C., Braun, A. C., Barra, R., Castillo, A., and Vogt, J.: Erosion proxies in an exotic tree plantation question the appropriate land use in Central Chile, *CATENA*, 161, 77–84, <https://doi.org/10.1016/j.catena.2017.10.017>, 2018.
- 425 Barros, S.: Evolución de las plantaciones forestales en Chile. Forestación y reforestación, *Ciencia e Investigación Forestal*, 24, 89–115, <https://bibliotecadigital.infor.cl/handle/20.500.12220/28235>, 2018.
- Bernhard, N., Moskwa, L.-M., Schmidt, K., Oeser, R. A., Aburto, F., Bader, M. Y., Baumann, K., von Blanckenburg, F., Boy, J., van den Brink, L., Brucker, E., Büdel, B., Canessa, R., Dippold, M. A., Ehlers, T. A., Fuentes, J. P., Godoy, R., Jung, P., Karsten, U., Köster, M., Kuzyakov, Y., Leinweber, P., Neidhardt, H., Matus, F., Mueller, C. W., Oelmann, Y., Oses, R., Osses, P., Paulino, L., Samolov, E., Schaller, M., Schmid, M., Spielvogel, S., Spohn, M., Stock, S., Stroncik, N., Tielbörger, K., Übernickel, K., Scholten, T., Seguel, O., Wagner, D., and Kühn, P.: Pedogenic and microbial interrelations to regional climate and local topography: New insights from a climate gradient (arid to humid) along the Coastal Cordillera of Chile, *CATENA*, 170, 335–355, <https://doi.org/10.1016/j.catena.2018.06.018>, 2018.
- 430 Bianchi-Gundian, V.: Erosión. Cáncer del suelo., Imprenta Universitaria, Santiago de Chile, 1947.
- Bladon, K. D., Emelko, M. B., Silins, U., and Stone, M.: Wildfire and the Future of Water Supply, *Environmental Science & Technology*, 48, 8936–8943, <https://doi.org/10.1021/es500130g>, 2014.
- 435 Boardman, J., Vandaele, K., Evans, R., and Foster, I. D.: Off-site impacts of soil erosion and runoff: Why connectivity is more important than erosion rates, *Soil Use and Management*, 35, 245–256, <https://doi.org/10.1111/sum.12496>, 2019.
- Boisier, J. P., Alvarez-Garretón, C., Cordero, R. R., Damiani, A., Gallardo, L., Garreaud, R. D., Lambert, F., Ramallo, C., Rojas, M., and Rondanelli, R.: Anthropogenic drying in central-southern Chile evidenced by long-term observations and climate model simulations, *Elementa*, 6, <https://doi.org/10.1525/elementa.328>, 2018.
- 440 Bonilla, C. A. and Johnson, O. I.: Soil erodibility mapping and its correlation with soil properties in Central Chile, *Geoderma*, 189–190, 116–123, <https://doi.org/10.1016/j.geoderma.2012.05.005>, 2012.
- Borrelli, P., Robinson, D. A., Panagos, P., Lugato, E., Yang, J. E., Alewell, C., Wuepper, D., Montanarella, L., and Ballabio, C.: Land use and climate change impacts on global soil erosion by water (2015–2070), *Proceedings of the National Academy of Sciences of the United States of America*, 117, 21 994–22 001, <https://doi.org/10.1073/pnas.2001403117>, 2020.
- 445

- Bowman, D. M. J. S., Moreira-Muñoz, A., Kolden, C. A., Chávez, R. O., Muñoz, A. A., Salinas, F., González-Reyes, A., Rocco, R., de la Barrera, F., Williamson, G. J., Borchers, N., Cifuentes, L. A., Abatzoglou, J. T., and Johnston, F. H.: Human–environmental drivers and impacts of the globally extreme 2017 Chilean fires, *Ambio*, 48, 350–362, <https://doi.org/10.1007/s13280-018-1084-1>, 2019.
- Braucher, R., Merchel, S., Borgomano, J., and Bourlès, D.: Production of cosmogenic radionuclides at great depth: A multi element approach, *Earth and Planetary Science Letters*, 309, 1–9, <https://doi.org/10.1016/j.epsl.2011.06.036>, 2011.
- Brown, G. W. and Krygier, J. T.: Clear-Cut Logging and Sediment Production in the Oregon Coast Range, *Water Resources Research*, 7, 1189–1198, <https://doi.org/10.1029/WR007i005p01189>, 1971.
- Cabezas, J. and Fassnacht, F. E.: Reconstructing the Vegetation Disturbance History of a Biodiversity Hotspot in Central Chile Using Landsat, Bfast and Landtrends, in: *IGARSS 2018 - 2018 IEEE International Geoscience and Remote Sensing Symposium*, pp. 7636–7639, IEEE, <https://doi.org/10.1109/IGARSS.2018.8518863>, 2018.
- Carretier, S., Tolorza, V., Regard, V., Aguilar, G., Bermúdez, M., Martinod, J., Guyot, J.-L., Hérail, G., and Riquelme, R.: Review of erosion dynamics along the major N-S climatic gradient in Chile and perspectives, *Geomorphology*, 300, 45–68, <https://doi.org/10.1016/j.geomorph.2017.10.016>, 2018.
- Cavalli, M., Trevisani, S., Comiti, F., and Marchi, L.: Geomorphometric assessment of spatial sediment connectivity in small Alpine catchments, *Geomorphology*, 188, 31–41, <https://doi.org/10.1016/j.geomorph.2012.05.007>, 2013.
- Chilean Law 19.300: Decreto 40. Reglamento del Sistema de Evaluación de Impacto Ambiental, <http://bcn.cl/2f8a8>, 2013.
- Chuvieco, E.: *Teledetección Ambiental: la observación de la tierra desde el espacio*. 3a edición., Ariel Ciencia, Barcelona, 2008.
- Cifuentes-Croquevielle, C., Stanton, D. E., and Armesto, J. J.: Soil invertebrate diversity loss and functional changes in temperate forest soils replaced by exotic pine plantations, *Scientific Reports*, 10, 7762, <https://doi.org/10.1038/s41598-020-64453-y>, 2020.
- CIREN: Cuenca Río Purapel (Estados erosivos actuales), <http://erosionmaule.ciren.cl/documents/348>, 2021.
- Cisternas, M., Araneda, A., Martínez, P., and Pérez, S.: Effects of historical land use on sediment yield from a lacustrine watershed in central Chile, *Earth Surface Processes and Landforms*, 26, 63–76, [https://doi.org/10.1002/1096-9837\(200101\)26:1<63::AID-ESP157>3.0.CO;2-J](https://doi.org/10.1002/1096-9837(200101)26:1<63::AID-ESP157>3.0.CO;2-J), 2001.
- Cleveland, W. S.: LOWESS: A Program for Smoothing Scatterplots by Robust Locally Weighted Regression, *The American Statistician*, 35, 54, <https://doi.org/10.2307/2683591>, 1981.
- CONAF and MINAGRI: Decreto Ley 701 de 1974, Cuarenta años de incentivos a la forestación (1975-2015), <https://biblioteca.digital.gob.cl/handle/123456789/2334>, 2016.
- Cortés, L., Hernández, H. J., and Silva, P.: Historic Land Cover Change assesment of Chilean Mediterranean Coast: Did forest plantations really caused fragmentation?, *ISPRS Annals of the Photogrammetry, Remote Sensing and Spatial Information Sciences*, V-3-2022, 383–388, <https://doi.org/10.5194/isprs-annals-V-3-2022-383-2022>, 2022.
- Covault, J. A., Craddock, W. H., Romans, B. W., Fildani, A., and Gosai, M.: Spatial and Temporal Variations in Landscape Evolution: Historic and Longer-Term Sediment Flux through Global Catchments, *The Journal of Geology*, 121, 35–56, <https://doi.org/10.1086/668680>, 2013.
- Crovo, O., Aburto, F., Alborno, M. F., and Southard, R.: Soil type modulates the response of C, N, P stocks and stoichiometry after native forest substitution by exotic plantations, *CATENA*, 197, 104 997, <https://doi.org/10.1016/j.catena.2020.104997>, 2021.
- DellaSala, D. A.: “Real” vs. “Fake” Forests: Why Tree Plantations Are Not Forests, in: *Encyclopedia of the World’s Biomes*, vol. 3-5, pp. 47–55, Elsevier, <https://doi.org/10.1016/B978-0-12-409548-9.11684-7>, 2020.

- DGA: Actualización del balance hídrico nacional, SIT N°417 Ministerio de Obras Públicas, Tech. rep., Universidad de Chile, Pontificia Universidad Católica de Chile, Ministerio de Obras Públicas, Dirección General de Aguas, División de Estudios y Planificación, Santiago, Chile, <https://snia.mop.gob.cl/sad/REH5796v1.pdf>, 2017.
- 485 Ellis, E. C. and Ramankutty, N.: Putting people in the map: anthropogenic biomes of the world, *Frontiers in Ecology and the Environment*, 6, 439–447, <https://doi.org/10.1890/070062>, 2008.
- Foga, S., Scaramuzza, P. L., Guo, S., Zhu, Z., Dilley, R. D., Beckmann, T., Schmidt, G. L., Dwyer, J. L., Joseph Hughes, M., and Laue, B.: Cloud detection algorithm comparison and validation for operational Landsat data products, *Remote Sensing of Environment*, 194, 379–390, <https://doi.org/10.1016/j.rse.2017.03.026>, 2017.
- 490 Fuentealba, M., Latorre, C., Frugone-Álvarez, M., Sarricolea, P., Giralt, S., Contreras-Lopez, M., Prego, R., Bernárdez, P., and Valero-Garcés, B.: A combined approach to establishing the timing and magnitude of anthropogenic nutrient alteration in a mediterranean coastal lake-watershed system, *Scientific Reports*, 10, 5864, <https://doi.org/10.1038/s41598-020-62627-2>, 2020.
- Fuentealba, M., Latorre, C., Frugone-Álvarez, M., Sarricolea, P., Godoy-Aguirre, C., Armesto, J., Villacís, L. A., Laura Carrevedo, M., Meseguer-Ruiz, O., and Valero-Garcés, B.: Crossing a critical threshold: Accelerated and widespread land use changes
495 drive recent carbon and nitrogen dynamics in Vichuquén Lake (35°S) in central Chile, *Science of the Total Environment*, 791, <https://doi.org/10.1016/j.scitotenv.2021.148209>, 2021.
- Gabet, E. J., Mudd, S. M., Wood, R. W., Grieve, S. W. D., Binnie, S. A., and Dunai, T. J.: Hilltop Curvature Increases With the Square Root of Erosion Rate, *Journal of Geophysical Research: Earth Surface*, 126, 1–16, <https://doi.org/10.1029/2020jf005858>, 2021.
- Galleguillos, M., Gimeno, F., Puelma, C., Zambrano-Bigiarini, M., Lara, A., and Rojas, M.: Disentangling the effect of future land use
500 strategies and climate change on streamflow in a Mediterranean catchment dominated by tree plantations, *Journal of Hydrology*, 595, <https://doi.org/10.1016/j.jhydrol.2021.126047>, 2021.
- Garreaud, R. D., Boisier, J. P., Rondanelli, R., Montecinos, A., Sepúlveda, H. H., and Veloso-Aguila, D.: The Central Chile Mega Drought (2010–2018): A climate dynamics perspective, *International Journal of Climatology*, 40, 421–439, <https://doi.org/10.1002/joc.6219>, 2020.
- Gerding, V.: Manejo de las plantaciones de *Pinus radiata* D. Don en Chile, *Bosque*, 12, 3–10, 1991.
- 505 Gimeno, F., Galleguillos, M., Manuschevich, D., and Zambrano-Bigiarini, M.: A coupled modeling approach to assess the effect of forest policies in water provision: A biophysical evaluation of a drought-prone rural catchment in south-central Chile, *Science of The Total Environment*, 830, 154 608, <https://doi.org/10.1016/j.scitotenv.2022.154608>, 2022.
- Gómez-González, S., Paula, S., Cavieres, L. A., and Pausas, J. G.: Postfire responses of the woody flora of Central Chile: Insights from a germination experiment, *PLoS ONE*, 12, 1–12, <https://doi.org/10.1371/journal.pone.0180661>, 2017.
- 510 Gómez-González, S., Ojeda, F., and Fernandes, P. M.: Portugal and Chile: Longing for sustainable forestry while rising from the ashes, *Environmental Science & Policy*, 81, 104–107, <https://doi.org/10.1016/j.envsci.2017.11.006>, 2018.
- Granger, D. E. and Schaller, M.: Cosmogenic Nuclides and Erosion at the Watershed Scale, *Elements*, 10, 369–373, <https://doi.org/10.2113/gselements.10.5.369>, 2014.
- Hansen, M. C., Potapov, P. V., Moore, R., Hancher, M., Turubanova, S. A., Tyukavina, A., Thau, D., Stehman, S. V., Goetz, S. J., Loveland,
515 T. R., Kommareddy, A., Egorov, A., Chini, L., Justice, C. O., and Townshend, J. R. G.: High-Resolution Global Maps of 21st-Century Forest Cover Change, *Science*, 342, 850–853, <https://doi.org/10.1126/science.1244693>, 2013.
- Heilmayr, R., Echeverría, C., Fuentes, R., and Lambin, E. F.: A plantation-dominated forest transition in Chile, *Applied Geography*, 75, 71–82, <https://doi.org/10.1016/j.apgeog.2016.07.014>, 2016.

- Helsel, D. R., Hirsch, R. M., Ryberg, K. R., Archfield, S. A., and Gilroy, E. J.: Statistical methods in water resources, Tech. rep., USGS, Reston, VA, <https://doi.org/10.3133/tm4A3>, 2020.
- Hermosilla-Palma, K., Plischoff, P., and Folchi, M.: Sixty years of land-use and land-cover change dynamics in a global biodiversity hotspot under threat from global change, *Journal of Land Use Science*, 16, 467–478, <https://doi.org/10.1080/1747423X.2021.2011970>, 2021.
- Hewawasam, T., von Blanckenburg, F., Schaller, M., and Kubik, P.: Increase of human over natural erosion rates in tropical highlands constrained by cosmogenic nuclides, *Geology*, 31, 597–600, [https://doi.org/10.1130/0091-7613\(2003\)031<0597:IOHONE>2.0.CO;2](https://doi.org/10.1130/0091-7613(2003)031<0597:IOHONE>2.0.CO;2), 2003.
- Huang, S., Dong, Q., Zhang, X., and Deng, W.: Catchment natural driving factors and prediction of baseflow index for Continental United States based on Random Forest technique, *Stochastic Environmental Research and Risk Assessment*, 35, 2567–2581, <https://doi.org/10.1007/s00477-021-02057-2>, 2021.
- Huber, A., Iroumé, A., Mohr, C., and Frêne, C.: Effect of *Pinus radiata* and *Eucalyptus globulus* plantations on water resource in the Coastal Range of Biobio region, Chile, *Bosque (Valdivia)*, 31, 219–230, <https://doi.org/10.4067/S0717-92002010000300006>, 2010.
- Imaizumi, F., Sidle, R. C., and Kamei, R.: Effects of forest harvesting on the occurrence of landslides and debris flows in steep terrain of central Japan, *Earth Surface Processes and Landforms*, 33, 827–840, <https://doi.org/10.1002/esp.1574>, 2008.
- INFOR: Informe Técnico 165: *Eucalyptus nitens* en Chile: primera monografía., Tech. rep., INFOR, Valdivia, <https://bibliotecadigital.infor.cl/handle/20.500.12220/7741?show=full>, 2004.
- IPCC: Regional Fact Sheet – Central and South America in: Sixth Assessment Report. Working Group I – The Physical Science Basis, Ipcc, p. 2, <https://www.ipcc.ch/assessment-report/ar6/>, 2021.
- IREN: Evaluación de la erosión Cordillera de la Costa entre Valparaíso y Cautín, Tech. rep., Instituto de Investigación de Recursos Naturales, Instituto de Investigación de Recursos Naturales, Santiago, Chile, <http://bibliotecadigital.ciren.cl/handle/123456789/14283>, 1965.
- Iroumé, A., Mayen, O., and Huber, A.: Runoff and peak flow responses to timber harvest and forest age in southern Chile, *Hydrological Processes*, 20, 37–50, <https://doi.org/10.1002/hyp.5897>, 2006.
- Iroumé, A., Jones, J., and Bathurst, J. C.: Forest operations, tree species composition and decline in rainfall explain runoff changes in the Nacimiento experimental catchments, south central Chile, *Hydrological Processes*, 35, 1–21, <https://doi.org/10.1002/hyp.14257>, 2021.
- Key, C. H. and Benson, N. C.: Landscape Assessment (LA) sampling and analysis methods, in: Lutes, Duncan C.; Keane, Robert E.; Caratti, John F.; Key, Carl H.; Benson, Nathan C.; Sutherland, Steve; Gangi, Larry J. FIREMON: Fire effects monitoring and inventory system. Gen. Tech. Rep. RMRS-GTR-164-CD, pp. LA1–LA51, US Department of Agriculture, Forest Service, Rocky Mountain Research Station, <https://www.fs.usda.gov/treearch/pubs/24066>, 2006.
- Kirchner, J. W., Finkel, R. C., Riebe, C. S., Granger, D. E., Clayton, J. L., King, J. G., and Megahan, W. F.: Mountain erosion over 10 yr, 10 k.y., and 10 m.y. time scales, *Geology*, 29, 591, [https://doi.org/10.1130/0091-7613\(2001\)029<0591:MEOYKY>2.0.CO;2](https://doi.org/10.1130/0091-7613(2001)029<0591:MEOYKY>2.0.CO;2), 2001.
- Krone, L. V., Hampl, F. J., Schwerdhelm, C., Bryce, C., Ganzert, L., Kitte, A., Übernickel, K., Dielforder, A., Aldaz, S., Osés-Pedraza, R., Perez, J. P. H., Sanchez-Alfaro, P., Wagner, D., Weckmann, U., and von Blanckenburg, F.: Deep weathering in the semi-arid Coastal Cordillera, Chile, *Scientific Reports*, 11, <https://doi.org/10.1038/s41598-021-90267-7>, 2021.
- Ladson, A., Brown, R., Neal, B., and Nathan, R.: A standard approach to baseflow separation using the Lyne and Hollick filter, *Australian Journal of Water Resources*, 17, <https://doi.org/10.7158/W12-028.2013.17.1>, 2013.
- Lal, D.: Cosmic ray labeling of erosion surfaces: in situ nuclide production rates and erosion models, *Earth and Planetary Science Letters*, 104, 424–439, [https://doi.org/10.1016/0012-821X\(91\)90220-C](https://doi.org/10.1016/0012-821X(91)90220-C), 1991.
- Lara, A., Jones, J., Little, C., and Vergara, N.: Streamflow response to native forest restoration in former *Eucalyptus* plantations in south central Chile, *Hydrological Processes*, 35, <https://doi.org/10.1002/hyp.14270>, 2021.

- Li, M., Li, S., Liu, Q., Kang, Y., Liang, L., Yuan, X., Zhang, J., Wang, X., and Li, C.: Assessment of hydrological response to multiyear drought: Insights from lag characteristics and shift magnitude, *Hydrological Processes*, 36, <https://doi.org/10.1002/hyp.14636>, 2022.
- Malmer, A. and Grip, H.: Soil disturbance and loss of infiltrability caused by mechanized and manual extraction of tropical rainforest in Sabah, Malaysia, *Forest Ecology and Management*, 38, 1–12, [https://doi.org/10.1016/0378-1127\(90\)90081-L](https://doi.org/10.1016/0378-1127(90)90081-L), 1990.
- Manuschevich, D.: Land use as a socio-ecological system: Developing a transdisciplinary approach to studies of land use change in South-Central Chile, in: *Ecological Economic and Socio Ecological Strategies for Forest Conservation: A Transdisciplinary Approach Focused on Chile and Brazil*, pp. 79–97, Springer International Publishing, https://doi.org/10.1007/978-3-030-35379-7_5, 2020.
- Martini, L., Cavalli, M., and Picco, L.: Predicting sediment connectivity in a mountain basin: A quantitative analysis of the index of connectivity, *Earth Surface Processes and Landforms*, <https://doi.org/10.1002/esp.5331>, 2022.
- Martins, M. A., Machado, A. I., Serpa, D., Prats, S. A., Faria, S. R., Varela, M. E., González-Pelayo, O., and Keizer, J. J.: Runoff and inter-rill erosion in a Maritime Pine and a Eucalypt plantation following wildfire and terracing in north-central Portugal, *Journal of Hydrology and Hydromechanics*, 61, 261–268, <https://doi.org/10.2478/johh-2013-0033>, 2013.
- Méndez-Freire, V., Villaseñor, T., and Mellado, C.: Spatial and temporal changes in suspended sediment fluxes in central Chile induced by the mega drought: The case of the Itata River Basin (36°–37°S), *Journal of South American Earth Sciences*, 118, <https://doi.org/10.1016/j.jsames.2022.103930>, 2022.
- Mohr, C.: Hydrological and erosion responses to man-made and natural disturbances – Insights from forested catchments in South-central Chile, Ph.D. thesis, University of Potsdam, https://publishup.uni-potsdam.de/opus4-ubp/frontdoor/deliver/index/docId/6782/file/mohr_diss.pdf, 2013.
- Mohr, C. H., Montgomery, D. R., Huber, A., Bronstert, A., and Iroumé, A.: Streamflow response in small upland catchments in the Chilean coastal range to the Mw 8.8 Maule earthquake on 27 February 2010, *Journal of Geophysical Research*, 117, F02032, <https://doi.org/10.1029/2011JF002138>, 2012.
- Mohr, C. H., Coppus, R., Iroumé, A., Huber, A., and Bronstert, A.: Runoff generation and soil erosion processes after clear cutting, *Journal of Geophysical Research: Earth Surface*, 118, 814–831, <https://doi.org/10.1002/jgrf.20047>, 2013.
- Mohr, C. H., Zimmermann, A., Korup, O., Iroumé, A., Francke, T., and Bronstert, A.: Seasonal logging, process response, and geomorphic work, *Earth Surface Dynamics*, 2, 117–125, <https://doi.org/10.5194/esurf-2-117-2014>, 2014.
- Montgomery, D. R., Schmidt, K. M., Greenberg, H. M., and Dietrich, W. E.: Forest clearing and regional landsliding, *Geology*, 28, 311–314, [https://doi.org/10.1130/0091-7613\(2000\)28<311:FCARL>2.0.CO;2](https://doi.org/10.1130/0091-7613(2000)28<311:FCARL>2.0.CO;2), 2000.
- Moody, J. A. and Martin, D. A.: Synthesis of sediment yields after wildland fire in different rainfall regimes in the western United States, *International Journal of Wildland Fire*, 18, 96, <https://doi.org/10.1071/WF07162>, 2009.
- Murphy, B. P., Czuba, J. A., and Belmont, P.: Post-wildfire sediment cascades: A modeling framework linking debris flow generation and network-scale sediment routing, *Earth Surface Processes and Landforms*, 44, 2126–2140, <https://doi.org/10.1002/esp.4635>, 2019.
- Pepin, E., Carretier, S., Guyot, J. L., and Escobar, F.: Specific suspended sediment yields of the Andean rivers of Chile and their relationship to climate, slope and vegetation, *Hydrological Sciences Journal*, 55, 1190–1205, <https://doi.org/10.1080/02626667.2010.512868>, 2010.
- Pizarro, R., García-Chevesich, P., Pino, J., Ibáñez, A., Pérez, F., Flores, J. P., Sharp, J. O., Ingram, B., Mendoza, R., Neary, D. G., Sangüesa, C., and Vallejos, C.: Stabilization of stage–discharge curves following the establishment of forest plantations: Implications for sediment production, *River Research and Applications*, 36, 1828–1837, <https://doi.org/10.1002/rra.3718>, 2020.

Pizarro, R., García-Chevesich, P., Ingram, B., Sangüesa, C., Pino, J., Ibáñez, A., Mendoza, R., Vallejos, C., Pérez, F., Flores, J. P., Vera, M., Balocchi, F., Bustamante-Ortega, R., and Martínez, G.: Establishment of Monterrey Pine (*Pinus radiata*) Plantations and Their Effects on Seasonal Sediment Yield in Central Chile, *Sustainability*, 15, 6052, <https://doi.org/10.3390/su15076052>, 2023.

Ratta, R. and Lal, R.: Soil quality and soil erosion, CRC press, 1998.

Reneau, S. L., Katzman, D., Kuyumjian, G. A., Lavine, A., and Malmon, D. V.: Sediment delivery after a wildfire, *Geology*, 35, 151–154, <https://doi.org/10.1130/G23288A.1>, 2007.

Roering, J. J., Perron, J. T., and Kirchner, J. W.: Functional relationships between denudation and hillslope form and relief, *Earth and Planetary Science Letters*, 264, 245–258, <https://doi.org/10.1016/j.epsl.2007.09.035>, 2007.

Schaller, M. and Ehlers, T. A.: Comparison of soil production, chemical weathering, and physical erosion rates along a climate and ecological gradient (Chile) to global observations, *Earth Surface Dynamics*, 10, 131–150, <https://doi.org/10.5194/esurf-10-131-2022>, 2022.

Schuller, P., Walling, D. E., Iroumé, A., Quilodrán, C., Castillo, A., and Navas, A.: Using ¹³⁷Cs and ²¹⁰Pbex and other sediment source fingerprints to document suspended sediment sources in small forested catchments in south-central Chile, *Journal of Environmental Radioactivity*, 124, 147–159, <https://doi.org/10.1016/j.jenvrad.2013.05.002>, 2013.

Schuller, P., Walling, D. E., Iroumé, A., Quilodrán, C., and Castillo, A.: Quantifying the temporal variation of the contribution of fine sediment sources to sediment yields from Chilean forested catchments during harvesting operations, *Bosque (Valdivia)*, 42, 231–244, <https://doi.org/10.4067/S0717-92002021000200231>, 2021.

Serey, A., Piñero-Feliciangeli, L., Sepúlveda, S. A., Poblete, F., Petley, D. N., and Murphy, W.: Landslides induced by the 2010 Chile megathrust earthquake: a comprehensive inventory and correlations with geological and seismic factors, *Landslides*, <https://doi.org/10.1007/s10346-019-01150-6>, 2019.

Sidle, R. C. and Ziegler, A. D.: The dilemma of mountain roads, <https://doi.org/10.1038/ngeo1512>, 2012.

Skovlund, E. and Fenstad, G. U.: Should we always choose a nonparametric test when comparing two apparently nonnormal distributions?, *Journal of Clinical Epidemiology*, 54, 86–92, [https://doi.org/10.1016/S0895-4356\(00\)00264-X](https://doi.org/10.1016/S0895-4356(00)00264-X), 2001.

Solar, W.: Manual de terreno y centros de filtrado. Dirección General de Aguas, <https://snia.mop.gob.cl/sad/SED4939.pdf>, 1999.

Soto, L., Galleguillos, M., Seguel, O., Sotomayor, B., and Lara, A.: Assessment of soil physical properties' statuses under different land covers within a landscape dominated by exotic industrial tree plantations in south-central Chile, *Journal of Soil and Water Conservation*, 74, 12–23, <https://doi.org/10.2489/jswc.74.1.12>, 2019.

Stone, J. O.: Air pressure and cosmogenic isotope production, *Journal of Geophysical Research*, 105, 23 753, <https://doi.org/10.1029/2000JB900181>, 2000.

Summerfield, M. a. and Hulton, N. J.: Natural controls of fluvial denudation rates in major world drainage basins, *Journal of Geophysical Research*, 99, 13 871–13 883, <https://doi.org/10.1029/94JB00715>, 1994.

Syvitski, J., Ángel, J. R., Saito, Y., Overeem, I., Vörösmarty, C. J., Wang, H., and Olago, D.: Earth's sediment cycle during the Anthropocene, *Nature Reviews Earth & Environment*, <https://doi.org/10.1038/s43017-021-00253-w>, 2022.

Teutschbein, C., Grabs, T., Karlsen, R. H., Laudon, H., and Bishop, K.: Hydrological response to changing climate conditions: Spatial streamflow variability in the boreal region, *Water Resources Research*, 51, 9425–9446, <https://doi.org/10.1002/2015WR017337>, 2015.

Tolorza, V., Carretier, S., Andermann, C., Ortega-Culaciati, F., Pinto, L., and Mardones, M.: Contrasting mountain and piedmont dynamics of sediment discharge associated with groundwater storage variation in the Biobío river, *Journal of Geophysical Research: Earth Surface*, 119, 2730–2753, <https://doi.org/10.1002/2014JF003105>, 2014.

- 630 Tolorza, V., Mohr, C. H., Carretier, S., Serey, A., Sepúlveda, S. A., Tapia, J., and Pinto, L.: Suspended Sediments in Chilean Rivers Reveal Low Postseismic Erosion After the Maule Earthquake (Mw 8.8) During a Severe Drought, *Journal of Geophysical Research: Earth Surface*, m, 2018JF004 766, <https://doi.org/10.1029/2018JF004766>, 2019.
- Tolorza, V., Poblete-Caballero, D., Banda, D., Little, C., Leal, C., and Galleguillos, M.: An operational method for mapping the composition of post-fire litter, *Remote Sensing Letters*, 13, 511–521, <https://doi.org/10.1080/2150704X.2022.2040752>, 2022.
- 635 Vanacker, V., von Blanckenburg, F., Govers, G., Molina, A., Poesen, J., Deckers, J., and Kubik, P.: Restoring dense vegetation can slow mountain erosion to near natural benchmark levels, *Geology*, 35, 303, <https://doi.org/10.1130/G23109A.1>, 2007.
- Vanacker, V., Guns, M., Clapuyt, F., Balthazar, V., Tenorio, G., and Molina, A.: Spatio-temporal patterns of landslides and Erosion in tropical Andean catchments, *Pirineos*, 175, <https://doi.org/10.3989/PIRINEOS.2020.175001>, 2020.
- Vanacker, V., Molina, A., Rosas-Barturen, M., Bonnesoeur, V., Román-Dañobeytia, F., Ochoa-Tocachi, B. F., and Buytaert, W.: The effect of
640 natural infrastructure on water erosion mitigation in the Andes, *SOIL*, 8, 133–147, <https://doi.org/10.5194/soil-8-133-2022>, 2022.
- Vázquez, M., Ramírez, S., Morata, D., Reich, M., Braun, J. J., and Carretier, S.: Regolith production and chemical weathering of granitic rocks in central Chile, *Chemical Geology*, 446, 87–98, <https://doi.org/10.1016/j.chemgeo.2016.09.023>, 2016.
- Verbesselt, J., Hyndman, R., Newnham, G., and Culvenor, D.: Detecting trend and seasonal changes in satellite image time series, *Remote Sensing of Environment*, 114, 106–115, <https://doi.org/10.1016/j.rse.2009.08.014>, 2010.
- 645 von Blanckenburg, F.: The control mechanisms of erosion and weathering at basin scale from cosmogenic nuclides in river sediment, *Earth and Planetary Science Letters*, 237, 462–479, <https://doi.org/10.1016/j.epsl.2005.06.030>, 2005.
- von Blanckenburg, F. and Willenbring, J. K.: Cosmogenic Nuclides: Dates and Rates of Earth-Surface Change, *Elements*, 10, 341–346, <https://doi.org/10.2113/gselements.10.5.341>, 2014.
- Wainwright, J., Turnbull, L., Ibrahim, T. G., Lexartza-Artza, I., Thornton, S. F., and Brazier, R. E.: Linking environmental régimes, space and time: Interpretations of structural and functional connectivity, *Geomorphology*, 126, 387–404, <https://doi.org/10.1016/j.geomorph.2010.07.027>, 2011.
- 650 Wohl, E., Brierley, G., Cadol, D., Coulthard, T. J., Covino, T., Fryirs, K. A., Grant, G., Hilton, R. G., Lane, S. N., Magilligan, F. J., Meitzen, K. M., Passalacqua, P., Poeppl, R. E., Rathburn, S. L., and Sklar, L. S.: Connectivity as an emergent property of geomorphic systems, *Earth Surface Processes and Landforms*, 44, 4–26, <https://doi.org/10.1002/esp.4434>, 2019.
- Zhang, J., Zhang, Y., Song, J., and Cheng, L.: Evaluating relative merits of four baseflow separation methods in Eastern Australia, *Journal of Hydrology*, 549, 252–263, <https://doi.org/10.1016/j.jhydrol.2017.04.004>, 2017.
- Zhao, Y., Feng, D., Yu, L., Wang, X., Chen, Y., Bai, Y., Hernández, H. J., Galleguillos, M., Estades, C., Biging, G. S., Radke, J. D., and Gong, P.: Detailed dynamic land cover mapping of Chile: Accuracy improvement by integrating multi-temporal data, *Remote Sensing of Environment*, 183, 170–185, <https://doi.org/10.1016/j.rse.2016.05.016>, 2016.




Figure 7. Land cover classification and transitions. Maps of 1955 and 1975 from Hermosilla-Palma et al. (2021), 1986-2015 from this work, and 2017 from Tolorza et al. (2022).




Figure 8. Sediment connectivity index (Cavalli et al., 2013) calculated using (A) the streams and (B) the streams and forest roads as targets. (C) is the difference between both models. (D-F) Details of hilltops with highest values of RC .




Figure 9. Detected disturbances from BFAST (A) map of the number of disturbances in vegetation detected for the period 2002-2019 (B) Seasonality of disturbance area detected within the Purapel catchment.

S1 Timing for sediment mobilization at the Purapel catchment

The available data on suspended sediments in the studied river have a daily resolution. Catchment size and flow velocity on first-order rivers are important in estimating whether or not a daily sampling may reproduce the main events of erosion. To estimate ranges for the travel time of particles in the studied landscape, we measured streamflows in a small river of the region (~ 750 ha drained area) on two dates (high and low flow). We obtained a maximum flow velocity of 0.58 m/s during a flood in the wet season (August 2021), and a minimum of 0.03 m/s during the dry season (March 2022). Using that range of flow velocities, we estimated a travel time at constant velocity for all the distribution of distances to the gauge *Purapel en Sauzal* (Figure S1). For the flood, the maximum travel time is close to 26 hrs. During low flows it is of hundreds of hours.

We also estimated the Time of Concentration using several approaches, recognizing the very large uncertainties in its estimation (Grimaldi et al., 2012, Table S1). The highest T_c (Viparelli, 1961-1963) was very close to the maximum travel time estimated from the distances to the outlet.

Based on those results, we argue that a daily sampling can capture sediment mobilization for a wide range of hydrologic events affecting the entire catchment. To calculate suspended sediment discharge, we should assume a distribution which is unknown, since sub-daily data is not available. In the absence of that sub-daily data, we assumed a constant value for the entire day, which is obtained from the instant measurements published by the DGA.




Figure S1: Estimations for travel time of particles between the hillslopes and the stream gauge at constant flow velocity. Distances to the outlet were calculated using TauDEM version 5 (<https://hydrology.usu.edu/taudem/taudem5/>)

Table S1: Several estimations for Time of Concentration. See Grimaldi et al. (2012) for details

Source of equation	Tc(Hours)	Tc(minutes)					
'Department of Public Works' (1995)	8,42892	505,7	L=	34,529	H=	1930	
Giandotti (1934)	14,6534	879,2	A=	406	L=	43,28	H= 154,1
Kirpich (1940)	2,02502	121,5	L=	141995	S=	0,26	
Viparelli (1961, 1963)	26,6133	1597	L=	55569	V=	0,58	

S2 Analysis of missing data of streamflow and suspended sediment concentration in the *Río Purapel en Sauzal* Station

The gauge *Río Purapel en Sauzal* was monitored by the Chilean Directorate of Water since 1981 on a daily basis. Streamflow (Q , m^3/s) records includes the period 1981-06-02 to 2019-06-30, while Suspended Sediment Concentration (SSC, mg/l) was recorded between 1985-06-06 and 2018-11-04. The Suspended Sediment Discharge, calculated as the product $Q \times \text{SSC}$, can be obtained for a total of 10,105 daily data. That means 33.06% of missing data.

Previous studies on Central-South Chile had reported a decline in sediment transport together with water in rivers after 2010 (Méndez-Freire et al., 2022; Tolorza et al., 2019). Under presumable non-stationary conditions, we discarded the usage of gap-filling techniques for trend analysis. Instead, we decided to carry out a monthly analysis using only months with less than 15 missing data (i.e., more than 50% of available data). The monthly amount of data (%) is in figure S2, and we noticed that our threshold of 50% discard a large amount of months, which difficult the analysis of trends.

To obtain more reliable results, we decided to aggregate the data by season, but increasing the threshold percent of available data from 50% to 66.6% (i.e., accepting a maximum of 30 missing daily data in each 90-day season period). Here we define seasons as Autumn (MAM), Winter (JJA), Spring (SON) and Summer (DJF). The percent of data by season is in figure S3.




Figure S2: Percent of monthly data on a daily basis




Figure S3: Percent of seasonal data on a daily basis

S3 Baseflow separation on daily streamflow data at Purapel en Sauzal station

We calculated the daily baseflow at Purapel en Sauzal station with the Lyne and Hollick filter (Ladson et al., 2013) by using several α values between 0.5 and 0.95 and $n.reflected=30$ days as parameters. The results for each α for 1994 is in Fig. S4 as an example. Daily CR2met precipitation is plotted for reference. All resulted baseflows are available in the supporting information Baseflow.RData file. We selected the results obtained with $\alpha=0.7$ for further trend analysis, given the observed magnitudes and shape of the baseflow time series.




Figure S4: Streamflow (red), baseflow (blue) and rainfall (black) for the hydrologic year (M-F) 1994

References

- 'Department of Public Works'. *California culvert practice, second edition*. DPW, Division of Highways, Sacramento, CA, 1995.
- M. Giandotti. Previsione delle piene e delle magre dei corsi d'acqua. *Istituto Poligrafico dello Stato*, 8: 107–117, 1934.
- S. Grimaldi, A. Petroselli, F. Tauro, and M. Porfiri. Time of concentration: a paradox in modern hydrology. *Hydrological Sciences Journal*, 57(2):217–228, 2 2012. ISSN 0262-6667. doi: 10.1080/02626667.2011.644244. URL <https://www.tandfonline.com/doi/full/10.1080/02626667.2011.644244>.
- Z. P. Kirpich. Time of concentration of small agricultural watersheds. *Civil engineering*, 10(6):362, 1940.
- A. Ladson, R. Brown, B. Neal, and R. Nathan. A standard approach to baseflow separation using the Lyne and Hollick filter. *Australian Journal of Water Resources*, 17(1), 2013. ISSN 13241583. doi: 10.7158/W12-028.2013.17.1. URL http://www.engineersmedia.com.au/journals/ajwr/2013/17_1/W12_028.html.
- V. Méndez-Freire, T. Villaseñor, and C. Mellado. Spatial and temporal changes in suspended sediment fluxes in central Chile induced by the mega drought: The case of the Itata River Basin (36°-37°S). *Journal of South American Earth Sciences*, 118, 10 2022. ISSN 08959811. doi: 10.1016/j.jsames.2022.103930.
- V. Tolorza, C. H. Mohr, S. Carretier, A. Serey, S. A. Sepúlveda, J. Tapia, and L. Pinto. Suspended Sediments in Chilean Rivers Reveal Low Postseismic Erosion After the Maule Earthquake (Mw 8.8) During a Severe Drought. *Journal of Geophysical Research: Earth Surface*, m:2018JF004766, 6 2019. ISSN 2169-9003. doi: 10.1029/2018JF004766. URL <https://onlinelibrary.wiley.com/doi/abs/10.1029/2018JF004766>.
- C. Viparelli. *Ricostruzione dell'idrogramma di piena*. Napoli:Istituto di Idraulica dell'Università di Palermo, Stab. Tip.Genovese, 1961.
- C. Viparelli. Ricostruzione dell'idrogramma di piena. *L'Energia Elettrica*, 6:421–428, 1963.