

# Ultra-Slow Discharges That Precede Lightning Initiation

Sterpka Christopher Francis<sup>1</sup>, Dwyer Joseph R<sup>1</sup>, Liu Ningyu<sup>2</sup>, Demers Nicholas R<sup>1</sup>, Hare Brian<sup>3</sup>, Scholten Olaf<sup>3</sup>, and ter Veen Sander<sup>4</sup>

<sup>1</sup>University of New Hampshire

<sup>2</sup>The University of New Hampshire

<sup>3</sup>University of Groningen

<sup>4</sup>ASTRON

November 16, 2022

## Abstract

We report on ultra-slowly propagating discharge events with speeds in the range 1-13 km/s, much lower than any known lightning process. The propagation speeds of these discharges are orders of magnitude slower than leader or streamer speeds, but faster than the ion drift speed. For one particular event, a lightning leader forms about 40 ms later within 50 m of the discharge, likely within the same high field region. A second slow event forms 9 ms prior to the initiation, and leads into the negative leader. Most slow events appear to not be directly involved with lightning initiation. This suggests that the classic streamer cascade model of initiation is not always a definitive process. In this work we describe these discharge events displaying unique behavior, their relation to common lightning discharges, and their implications for lightning initiation.

# Ultra-Slow Discharges That Precede Lightning Initiation

C. Sterpka<sup>1\*</sup>, J. Dwyer<sup>1</sup>, N. Liu<sup>1</sup>, N. Demers<sup>1</sup>, B.M. Hare<sup>2</sup>, O. Scholten<sup>2,3</sup>,  
S. ter Veen<sup>4</sup>

<sup>1</sup>Space Science Center (EOS), Department of Physics and Astronomy, University of New Hampshire,  
Durham NH, USA

<sup>2</sup>University Groningen, Kapteyn Astronomical Institute, Landleven 12, 9747 AD Groningen, The  
Netherlands

<sup>3</sup>Interuniversity Institute for High-Energy, Vrije Universiteit Brussel, Pleinlaan 2, 1050 Brussels, Belgium

<sup>4</sup>Netherlands Institute for Radio Astronomy (ASTRON), Dwingeloo, The Netherlands

## Key Points:

- The ultra-slowly propagating events travel at speeds at least an order of magnitude slower than the slowest positive leaders.
- In one observed case, the slow propagation led directly into the formation of a lightning leader.
- In most cases, these discharges are not connected with lightning initiation.

## Abstract

We report on ultra-slowly propagating discharge events with speeds in the range 1-13 km/s, much lower than any known lightning process. The propagation speeds of these discharges are orders of magnitude slower than leader or streamer speeds, but faster than the ion drift speed. For one particular event, a lightning leader forms about 40 ms later within 50 m of the discharge, likely within the same high field region. A second slow event forms 9 ms prior to the initiation, and leads into the negative leader. Most slow events appear to not be directly involved with lightning initiation. This suggests that the classic streamer cascade model of initiation is not always a definitive process. In this work we describe these discharge events displaying unique behavior, their relation to common lightning discharges, and their implications for lightning initiation.

## Plain Language Summary

While lightning is generally a very fast process, here we report on ultra-slow discharges which may be a new and unexpected method of lightning initiation. These discharges travel at uncharacteristically low speeds and are observed in conjunction with lightning initiation in two cases, while in three different cases they are not. This indicates that these events are also evidence of failed lightning leader formation, which complicates the current understanding of how lightning initiates. Additionally, the velocity of these events is slow enough that in principle the propagation can be observed by the unaided eye - challenging the colloquial notion of “fast as lightning.”

## Introduction


Lightning is generally a very fast process, with each discharge having a range of associated speeds. The slowest reported speeds are positive leaders, which are commonly reported in the range of  $1.6\text{--}3\times 10^4$  m/s, with an average velocity of about  $2\times 10^4$  m/s (van der Velde & Montanyà, 2013) (with an exception for one esoteric reference to rocket lightning, which travels “about as fast as a rocket” (Everett, 1903)). In 2D video observations, the speeds reported are possibly as low as 10 km/s (Kong et al., 2008). Negative leaders, which are a branched lightning process that expands outward as it approaches ground or another positively charged region, propagate at speeds between  $1\text{--}6\times 10^5$  m/s (Hill et al., 2011). Streamers, which are a cold-plasma phenomena underpinning many

discharge processes in lightning including both initiation and propagation, have been shown to possibly have speeds as low as  $10^5$  m/s just above the critical field for streamer formation (Liu & Dwyer, 2013; Koile et al., 2020; Dwyer & Uman, 2014), but have been observed as fast as  $1 \times 10^7$  m/s in sprites (McHarg et al., 2007; Phelps & Griffiths, 1976). In typical lightning initiation processes, however, it has been shown that positive streamers grow in VHF at  $4.8 \times 10^6$  m/s (Sterpka et al., 2021). Anvil crawlers, also known as spider lightning, are mistaken for slowly propagating leaders where the propagation can be observed by eye. However, this is only due to the spatial extent in which they cover, their travel speeds are between  $2-4 \times 10^5$  m/s (Mazur et al., 1998; Peterson et al., 2021).

In this work, imaging for all figures is performed with the Time Resolved Interferometric 3D (TRI-D) imager, which provides location, intensity, and polarization of sources (Scholten et al., 2021b). This is possible in part due to LOFAR's thousands of VHF (30-80 MHz) antennas, of which hundreds are selected to allow for extraordinary sensitivity through interferometric beamforming, and also enable detection of lightning features with meter scale precision and intensities that are below the level of galactic background (Sterpka et al., 2021; Hare et al., 2018; Scholten et al., 2021a). The intensity units used within this work are orders of the galactic background (gb), and they represent the normalized noise level for individual antennas (Sterpka et al., 2021; Scholten et al., 2021b). In this paper we will present four ultra-slowly propagating discharge events imaged by the LOFAR radio telescope.

## Results

The first slow propagation event was discovered in close proximity to a flash that took place on June 27, 2020 at about 14:51 UTC, denoted as flash 20B-10. Sources develop about 65 ms before the initiation of a lightning leader (Figure 1) and about 50 m south east from the initiation location. The event took place 19 km west, 12 km north, and at an altitude of about 6 km from the LOFAR core. On the top left of the figure, the altitude versus time for the slow propagation is displayed in a wagon wheel style TRI-D plot. The top right of the same plot shows the initial development of the lightning leader, about 65 ms after the start of the slow propagation. The size of each wagon wheel indicates the relative VHF intensity. Note that the intensity of the slow propagation is similar to the initial intensity of the first few sources of the lightning leader.




**Figure 1.** This figure displays the ultra-slow propagation and the lightning leader that follows 40 ms after the cessation via a wagon-wheel TRI-D plot. The slow propagation and lightning leader sources are labeled with arrows in both the altitude versus time (top) and ground projection (bottom). The color indicates the relative timing, and the size qualitatively indicates relative intensity. The initial development of the slow propagation is followed by the onset of the lightning leader in green. Note that overlap of the sources in the ground projection indicates the close proximity of the separate discharges.

The second figure displays quadratic fits for the ultra-slow propagation via a least squares regression. The fits excludes sources that are more than 2.0 standard deviations from the central curve along each axis (omitted sources are indicated by a red "x"), and the intensity is cut to 2.0 gb; these cuts ensure sources which are artifacts of sidebeams and/or different distributions are excluded from the fit. The points indicate the source locations along the Easting (top panel), Northing (second panel from top), and altitude axes (third panel from top). The bottom panel provides a histogram of the spread density and a normal distribution with a 1.1 m standard deviation. The fit reveals that the discharge begins with a speed of about 1.9 km/s and decelerates to a speed of about 0.5 km/s, with an overall acceleration of  $-91 \text{ km/s}^2$ . Initially, there are several clusters of sources which form less than 1 ms apart (indicated by the purple, orange, and red source groupings on the top panel of Figure 1), then as the discharge progresses there are several sources which develop either individually or with only one or two adjacent sources to form a cluster. This continues until the cessation of the discharge 25 ms later. The propagation moves downward about 21 m, with a slight lateral velocity on both the North and East axes with displacements of 1.8 m and 1.1 m respectively.


Figure 3 shows a zoom in of the lightning leader, which is of significant interest as the initiation also begins with an ultra-slowly propagating discharge. Note that from 66-68.5 ms is linear with a speed of about  $1.5 \times 10^3 \text{ m/s}$ ; this abruptly changes to  $1.2 \times 10^6 \text{ m/s}$  slightly after 68.5 ms with the onset of the lightning leader. What is additionally of interest is that subsequent bursts of the first 6 ms (see supplemental figure S1) of this discharge are separated from the previous by about only 0.25-0.5 m for an average speed of only about  $300 \pm 200 \text{ m/s}$ . Alternatively, one could interpret this to mean that the sources are stationary within the margin of error of LOFAR.

A third slow propagation event (supplemental figure S3) was also found within the same data set and appears to be unconnected with local lightning activity. Sources developed 18 km west, 8 km north, and at an altitude of about 10 km from the LOFAR core. The discharge has a linear speed of about 1.0 km/s. Initially, there are only a few sources that develop, with the largest burst of activity forming 15 ms after the discharge starts. The closest lightning activity to this event is 2.5 km lower in altitude and south of the discharge about 700 ms before the slow propagation starts. What is particularly surprising about this discharge is that the propagation is not along the vertical axis, which is the usual electric field direction. While both negative and positive leaders are observed



**Figure 2.** First slow propagation, approximately 65 ms before the initiation of flash 20B-10.

Top panel shows the Easting versus time, middle top shows Northing vs time, middle bottom shows altitude versus time, and the bottom panel provides the spread density and corresponding normal probability distribution. The overall acceleration is  $91 \text{ km/s}^2$  with  $v_0 = 1.9 \text{ km/s}$ ,  $v_f = 0.5 \text{ km/s}$ . Sources outside two standard deviations along each axis are excluded from the fit and are indicated by a red 'x'. The black arrow denotes a burst that propagates away from the main trajectory.



**Figure 3.** Zoom of initialization of leader initiation displayed in figure 2. Note that discharge begins with ultra-slow propagation on left prior to the formation of the lightning leader on right. Since we are fitting the overall motion of the ultra-slow propagation, the weak sources that form the vertical lines are considered part of a different distribution and are ignored. Additionally, note that only the final 2.5 ms of the slow propagation are shown.

to grow horizontally in thunderstorms, the trajectory of the slow propagation is mainly along the north-south axis from the inception point, which is not typical of lightning discharges (Yuan et al., 2019; Wu et al., 2015).

The fourth observed slow propagation event (Supplemental Figure S4) was found in close proximity to a flash that took place on June 18th (colloquially known as The Netherlands Apocalypse Storm) at 17:46 UTC in 2021, denoted flash 21C-1. The event occurred 20 km west, 16 km South, and at an altitude of about 11 km from the LOFAR core. The discharge took place 800 ms before the nearest lightning event. This lightning discharge formed an intensely radiating negative leader (IRNL)(Scholten et al., 2021), about 150 m to the East. The slow discharge began with a slightly higher speed of about 12.5 km/s and quickly decelerated to a speed of about 1.7 km/s with an overall rate of change in velocity of  $-1158 \text{ km/s}^2$ .

## Discussion

The initial speeds of these discharges are typically of the order of  $1 \times 10^4 \text{ m/s}$ , but in some cases deceleration brings the speeds possibly as low as 100 m/s. For some of the ultra-slowly propagating discharges, the standard deviation of the trajectories is of the order of 1 m, indicating that the diameter of the channels is of the order of our resolution or less. These events have intermittent bursts where the average location of each



burst collectively forms an overall motion typically characterized by a decelerating quadratic trajectory; although for the discharge which initiates lightning, this was not the case.

While the ultra-slow discharges typically decelerate, the trajectory that preceded lightning had three distinct stages. Initially, the sources effectively remained in a fixed location over the first 6 ms (see supplemental figure S1). This was followed by an abrupt transition to a constant velocity of about  $1.5 \times 10^3$  before another abrupt change in velocity to  $1.2 \times 10^6$  m/s as the leader forms. The final two stages are analogous to results of previous observations of lightning initiation events (Sterpka et al., 2021). However, the major differences are the ultra-slow speed, the intensity profile of the initiating event remaining relatively constant throughout the trajectory, and that the constituent bursts are initially sparse, but then the density of sources increases.

As mentioned previously, since these events are likely within the same high field region of the thunderstorm and lead into an initiation event, this adds potential complications to the classic Griffiths and Phelps model (Griffiths & Phelps, 1976); if streamers form on hydrometeors within the same high field, why is it that in one location 50 m from the initiation a slow propagation forms without lightning initiation, however at the exact location it leads into leader formation? One would expect that the hydrometeor density and fields within this region should be of similar magnitude, otherwise the initiation would not take place. Additionally, previous studies (Tilles et al., 2019; Rison et al., 2016) have reported that lightning initiation begins with fast breakdown, but if lightning can initiate with an ultra-slow discharge or possibly even stationary discharge, how would this modify the understanding of virgin air breakdown? Typically observed streamer cascade initiation events have been shown to initiate with velocities between 2-4 orders of magnitude higher than the ultra-slow propagations (Sterpka et al., 2021; Rison et al., 2016; Tilles et al., 2019). This implies that if the discharges are related to the classic Griffiths and Phelps streamer cascade processes and they more often fail to trigger lightning than successfully initiate lightning, then this model cannot be a straightforward process in all cases (Griffiths & Phelps, 1976; Attanasio et al., 2019). Or, to be more explicit, simply having a field above the level required for breakdown and a high enough hydrometeor density to enable the formation of streamers may be necessary, but not sufficient conditions for the formation of a lightning leader (Dwyer & Uman, 2014). Lastly, as the slow propagation that forms in conjunction with the lightning leader leads into the initiation, what is the cause of the spontaneous transition?

Since these ultra-slowly propagating discharges are not always found in conjunction with an initiation event, they could also be connected with failed leader initiation (Shao et al., 1995; Kolmašová et al., 2020). Every other time that lightning initiation has been observed it's been a fast process, but these events do not necessitate the formation of lightning. The temptation is to think that the E-fields are below the breakdown threshold, however it is not clear that that would resolve the issue as leaders have been observed in low electric fields. Additionally, no leaders have been observed this slow, and certainly nothing that travels this slow for up to a hundred meters and for up to 70 ms (van der Velde & Montanyà, 2013; Hill et al., 2011; Kong et al., 2008). One final note is that the number of clusters are decreasing for the ultra-slow events that do not initiate lightning, but for the event that does initiate lightning the number of clusters increases with time. The natural inclination is to think that this would be indicative of an increase in hydrometeor density within the initiation region, but the issue with this conjecture is that this would mean that the density of hydrometeors would be changing on millisecond timescales, which seems highly unlikely.

What is surprising about these events in addition to their ultra-slow speeds is that there are gaps in VHF activity during the event that can last from fractions of a millisecond to tens of milliseconds. Additionally, sometimes bursts form propagating features that are nearly perpendicular to the trajectory (see for example the sources indicated by the black arrow in figure 2), similar to previously discovered positive leader needles (Hare et al., 2019). Optical measurements of positive leader velocities have shown that they may travel as slow as  $1.0 \times 10^4$  m/s. These connections are interpreted as only analogous features, as the overall propagation follows the expected upward trajectory of a negative leader for this altitude. This does however lead to the question of whether the structure and/or the frequency of the bursts are indicative of successful versus unsuccessful lightning initiation events.

One of the explanations that has been proposed and rejected is that this trajectory is somehow related to the ion drift velocity. This hypothesis was implausible, due to the fact that the ion drift speed at 6 km altitude is expected to only be about 600 m/s, so even the slowest event reported here would already exceed this by a factor of 2 (Dwyer & Uman, 2014).

## Conclusions

Within this work we highlight the features of these ultra-slowly propagating discharges through true 3D VHF beamforming that is only possible with the sensitivity of the LOFAR array. Future work will need to address the following questions:

1. Are the ultra-slowly propagating discharge events a common or uncommon method of lightning initiation and/or failed initiation?
2. Do the bursts form disorganized clusters or do they share features of streamer or leader discharges? Consequently, do the burst structures and/or frequency suggest whether the propagation leads to initiation versus failed initiation?
3. Are there associated environmental differences between the events that fail to initiate lightning versus those that succeed?
4. Most importantly, what are the physical processes that produces their ultra-slow speeds and corresponding implications for the Griffiths and Phelps model, given their role in initiation?

We have identified discharges that are remarkable both in their slow speeds and frequency in occurrence within LOFAR data. While only three events are described within this work, seven have been observed within three different data sets. The events presented here suggest a new form of lightning initiation and/or failed initiation characterized by velocities orders of magnitude slower than any known discharge process. This is supported by the fact that in at least one case the slow discharge leads directly into the formation of a lightning leader, although most of the observed propagations do not lead to lightning initiation. Given these facts, it is essential that further study address the outstanding questions to find their proper role in both initiation and failed initiation as well as the underlying physics behind their ultra-slow propagation speeds.

## Acknowledgments

Computations were performed on Marvin, a Cray CS500 supercomputer at UNH supported by the NSF MRI program under grant AGS-1919310.

## Author Contributions

C.S. drafted the manuscript and completed data analysis. J.D., N.L., O.S., and B.M.H. contributed to critical review of main text and interpretation of results. N.D. improved image calibration and provided edits to manuscript text and images. O.S and B.M.H. developed the interferometry software for this study. S.t.V. performed data calibration and acquisition.

## Open Research

Figures in this work were created with the Matplotlib Python package (Caswell et al., 2019). Data are located on the LOFAR Long Term Archive and can be downloaded after setting up a LOFAR LTA account and through following the instructions for "Staging Transient Buffer Board Data" (ASTRON, n.d.) using the wget software package as follows: `wget --no-check-certificate https://lofar-download.grid.surfsara.nl/lofigrid/SRMFifoGet.py?surl=srm://srm.grid.sara.nl/pnfs/grid.sara.nl/data/lofar/ops/TBB/lightning/L786655\_D20200627T145100.178Z\_\"station\"\_R000\_tbb.h5` and "station" is replaced with one of the names of the LOFAR stations: CS001, CS002, CS003, CS004, CS005, CS006, CS007, CS011, CS013, CS017, CS021, CS024, CS026, CS028, CS030, CS031, CS032, CS101, CS103, RS106, CS201, RS205, RS208, RS210, CS301, CS302, RS305, RS306, RS307, RS310, CS401, RS406, RS407, RS409, CS501, RS503, RS508, or RS509.

## Funding

This research was supported in part through the University of New Hampshire AFOSR Grants No. FA9550-16-1-0396 and No. FA9550-18-1-0358 and by NWO (VI.VENI.192.071).

## References

- ASTRON, L. (n.d.). *LOFAR Long Term Archive Access*.  
[https://www.astron.nl/lofarwiki/doku.php?id=public:lta\\_howto](https://www.astron.nl/lofarwiki/doku.php?id=public:lta_howto).
- Attanasio, A., Krehbiel, P. R., & da Silva, C. L. (2019). Griffiths and Phelps Lightning Initiation Model, Revisited. *Journal of Geophysical Research: Atmo-*

- spheres, 124(14), 8076–8094. doi: 10.1029/2019JD030399
- Caswell, T. A., Droettboom, M., Lee, A., Hunter, J., Firing, E., Stansby, D., ...  
Ivanov, P. (2019, December). *Matplotlib/matplotlib v3.1.2*. Zenodo. doi:  
10.5281/zenodo.3563226
- Dwyer, J. R., & Uman, M. A. (2014, January). The physics of lightning. *Physics Re-  
ports*, 534(4), 147–241. doi: 10.1016/j.physrep.2013.09.004
- Everett, W. H. (1903, October). Rocket Lightning. *Nature*, 68(1773), 599–599. doi:  
10.1038/068599c0
- Griffiths, R. F., & Phelps, C. T. (1976). A model for lightning initiation arising from  
positive corona streamer development. *Journal of Geophysical Research (1896-  
1977)*, 81(21), 3671–3676. doi: 10.1029/JC081i021p03671
- Hare, B. M., Scholten, O., Bonardi, A., Buitink, S., Corstanje, A., Ebert, U., ...  
Winchen, T. (2018). LOFAR Lightning Imaging: Mapping Lightning With  
Nanosecond Precision. *Journal of Geophysical Research: Atmospheres*, 123(5),  
2861–2876. doi: 10.1002/2017JD028132
- Hare, B. M., Scholten, O., Dwyer, J., Trinh, T. N. G., Buitink, S., ter Veen, S.,  
... Zucca, P. (2019, April). Needle-like structures discovered on pos-  
itively charged lightning branches. *Nature*, 568(7752), 360–363. doi:  
10.1038/s41586-019-1086-6
- Hill, J. D., Uman, M. A., & Jordan, D. M. (2011). High-speed video observations  
of a lightning stepped leader. *Journal of Geophysical Research: Atmospheres*,  
116(D16). doi: 10.1029/2011JD015818
- Koile, J., Shi, F., Liu, N., Dwyer, J., & Tilles, J. (2020). Negative Streamer  
Initiation From an Isolated Hydrometeor in a Subbreakdown Electric  
Field. *Geophysical Research Letters*, 47(15), e2020GL088244. doi:  
10.1029/2020GL088244
- Kolmašová, I., Santolík, O., Defer, E., Kašpar, P., Kolínská, A., Pedebay, S., & Co-  
quillat, S. (2020). Two Propagation Scenarios of Isolated Breakdown Lightning  
Processes in Failed Negative Cloud-to-Ground Flashes. *Geophysical Research  
Letters*, 47(23), e2020GL090593. doi: 10.1029/2020GL090593
- Kong, X., Qie, X., & Zhao, Y. (2008). Characteristics of downward leader in a  
positive cloud-to-ground lightning flash observed by high-speed video cam-  
era and electric field changes. *Geophysical Research Letters*, 35(5). doi:

10.1029/2007GL032764

- Liu, N., & Dwyer, J. R. (2013). Modeling terrestrial gamma ray flashes produced by relativistic feedback discharges. *Journal of Geophysical Research: Space Physics*, 118(5), 2359–2376. doi: 10.1002/jgra.50232
- Mazur, V., Shao, X.-M., & Krehbiel, P. R. (1998). “Spider” lightning in intracloud and positive cloud-to-ground flashes. *Journal of Geophysical Research: Atmospheres*, 103(D16), 19811–19822. doi: 10.1029/98JD02003
- McHarg, M. G., Stenbaek-Nielsen, H. C., & Kammer, T. (2007). Observations of streamer formation in sprites. *Geophysical Research Letters*, 34(6). doi: 10.1029/2006GL027854
- Peterson, M., Light, T. E. L., & Shao, X.-M. (2021). Combined Optical and Radio-Frequency Measurements of a Lightning Megaflash by the FORTE Satellite. *Journal of Geophysical Research: Atmospheres*, 126(15), e2020JD034411. doi: 10.1029/2020JD034411
- Phelps, C. T., & Griffiths, R. F. (1976, July). Dependence of positive corona streamer propagation on air pressure and water vapor content. *Journal of Applied Physics*, 47(7), 2929–2934. doi: 10.1063/1.323084
- Rison, W., Krehbiel, P. R., Stock, M. G., Edens, H. E., Shao, X.-M., Thomas, R. J., ... Zhang, Y. (2016, February). Observations of narrow bipolar events reveal how lightning is initiated in thunderstorms. *Nature Communications*, 7(1), 10721. doi: 10.1038/ncomms10721
- Scholten, O., Hare, B., Dwyer, J., Liu, N., Sterpka, C., Buitink, S., ... ter Veen, S. (2021a, April). Time resolved 3D interferometric imaging of a section of a negative leader with LOFAR. *arXiv:2104.11908 [astro-ph, physics:physics]*.
- Scholten, O., Hare, B. M., Dwyer, J., Liu, N., Sterpka, C., Buitink, S., ... ter Veen, S. (2021b, September). Time resolved 3D interferometric imaging of a section of a negative leader with LOFAR. *Physical Review D*, 104(6), 063022. doi: 10.1103/PhysRevD.104.063022
- Scholten, O., Hare, B. M., Dwyer, J., Liu, N., Sterpka, C., Kolmašová, I., ... Winchen, T. (2021, August). A distinct negative leader propagation mode. *Scientific Reports*, 11(1), 16256. doi: 10.1038/s41598-021-95433-5
- Shao, X. M., Krehbiel, P. R., Thomas, R. J., & Rison, W. (1995). Radio interferometric observations of cloud-to-ground lightning phenomena in Florida. *Jour-*

- 315 *nal of Geophysical Research: Atmospheres*, 100(D2), 2749–2783. doi: 10.1029/  
316 94JD01943
- 317 Sterpka, C., Dwyer, J., Liu, N., Hare, B. M., Scholten, O., Buitink, S., ... Nelles, A.  
318 (2021). The Spontaneous Nature of Lightning Initiation Revealed. *Geophysical*  
319 *Research Letters*, 48(23), e2021GL095511. doi: 10.1029/2021GL095511
- 320 Tilles, J. N., Liu, N., Stanley, M. A., Krehbiel, P. R., Rison, W., Stock, M. G., ...  
321 Wilson, J. (2019, April). Fast negative breakdown in thunderstorms. *Nature*  
322 *Communications*, 10(1), 1–12. doi: 10.1038/s41467-019-09621-z
- 323 van der Velde, O. A., & Montanyà, J. (2013). Asymmetries in bidirectional leader  
324 development of lightning flashes. *Journal of Geophysical Research: Atmo-*  
325 *spheres*, 118(24), 13,504–13,519. doi: 10.1002/2013JD020257
- 326 Wu, T., Yoshida, S., Akiyama, Y., Stock, M., Ushio, T., & Kawasaki, Z. (2015).  
327 Preliminary breakdown of intracloud lightning: Initiation altitude, propa-  
328 gation speed, pulse train characteristics, and step length estimation. *Jour-*  
329 *nal of Geophysical Research: Atmospheres*, 120(18), 9071–9086. doi:  
330 10.1002/2015JD023546
- 331 Yuan, S., Jiang, R., Qie, X., Sun, Z., Wang, D., & Srivastava, A. (2019). Develop-  
332 ment of Side Bidirectional Leader and Its Effect on Channel Branching of the  
333 Progressing Positive Leader of Lightning. *Geophysical Research Letters*, 46(3),  
334 1746–1753. doi: 10.1029/2018GL080718

# Supporting Information for Ultra-Slow Discharges That Precede Lightning Initiation

C. Sterpka<sup>1\*</sup>, J. Dwyer<sup>1</sup>, N. Liu<sup>1</sup>, N. Demers<sup>1</sup>, B.M. Hare<sup>2</sup>, O. Scholten<sup>2,3</sup>,

S. ter Veen<sup>4</sup>

<sup>1</sup>Space Science Center (EOS), Department of Physics and Astronomy, University of New Hampshire, Durham NH, USA

<sup>2</sup>University Groningen, Kapteyn Astronomical Institute, Landleven 12, 9747 AD Groningen, The Netherlands

<sup>3</sup>Interuniversity Institute for High-Energy, Vrije Universiteit Brussel, Pleinlaan 2, 1050 Brussels, Belgium

<sup>4</sup>Netherlands Institute for Radio Astronomy (ASTRON), Dwingeloo, The Netherlands

## Contents of this file


1. Figures S1 to S5

---




## Introduction


The following TRI-D images display a few additional ultra-slowly propagating electrical discharge events. While each of these discharges have unique and interesting characteristics, they are still seen as supplemental to the main text and therefore reside here.




**Figure S1.** Zoom in of the first 6 ms of the slow propagation event displayed in figure 3 in the main text. Shown are the height versus time on the top panel and the ground projection on the bottom panel. As shown, the first source (yellow) and the second source (light orange) are separated by about 1 ms, and a distance of about 250 cm. The second source is separated by about 500 cm from the fourth source, which is in turn separated by the final source (green) by about 300 cm and 5 ms. This is consistent with a stationary source within the margin of error of LOFAR.




**Figure S2.** Displayed here is a TRI-D image of a slow discharge also contained within the 20B-10 flash dataset. The trajectory has a standard deviation of 2.4 m. This event Follows a nearby lightning discharge by 400 ms after activity has ceased. The entire event is 55 ms long and is roughly 200 m from the associated flash. Two interesting features of this event are that between -50 ms and -20 ms there is a gap in activity and that overall motion of the discharge during this period is only about 3 m.



**Figure S3.** Isolated slow discharge event, with no observed flash within  $\pm 700$  ms from the propagation. The nearest lightning activity is about 2.5 km from the discharge. Velocity is about 1.0 km/s and the standard deviation is about 2.0 m. The most surprising feature of this particular event is that the trajectory is along the North-South axis, which is not the typical propagation direction of a negative leader within a thunderstorm.



**Figure S4.** This slow discharge precedes a lightning initiation event by about 800 ms and is about 100 m from the location of the initiation event. It starts with a velocity of 12.5 km/s and decelerates to 1.7 km/s about 20 ms later. Lower quality fit with a standard deviation of 2.8 m is attributed to simultaneous remote flash about 100 km from this discharge.



**Figure S5.** Shown here is an impulsive image of the slow propagation shown in figure S4 (yellow) and corresponding IRNL that forms 800 ms later (blue-green). The top plot indicates the altitude versus time in ms, the middle shows height versus easting, and the bottom details the northing versus easting or ground projection of the event. Note in the ground projection the discharges are within roughly separated by 100 m despite being 800 ms apart in time.