Is there a semi-molten layer at the base of the lunar mantle?

Michaela Walterová^{1,1}, Marie Behounkova^{2,2}, and Michael Efroimsky^{3,3}

¹DLR Institute of Planetary Research ²Charles University ³United States Naval Observatory

December 7, 2022

Abstract

Parameterised by the Love number k_2 and the tidal quality factor Q, and inferred from lunar laser ranging (LLR), tidal dissipation in the Moon follows an unexpected frequency dependence often interpreted as evidence for a highly dissipative, melt-bearing layer encompassing the core-mantle boundary. Within this, more or less standard interpretation, the basal layer's viscosity is required to be of order 10^{15} to 10^{16} Pa.s and its outer radius is predicted to extend to the zone of deep moonquakes. While the reconciliation of those predictions with the mechanical properties of rocks might be challenging, alternative lunar interior models without the basal layer are said to be unable to fit the frequency dependence of tidal Q.

The purpose of our paper is to illustrate under what conditions the frequency-dependence of lunar tidal Q can be interpreted without the need for deep-seated partial melt. Devising a simplified lunar model, in which the mantle is described by the Sundberg-Cooper rheology, we predict the relaxation strength and characteristic timescale of elastically-accommodated grain boundary sliding in the mantle that would give rise to the desired frequency dependence. Along with developing this alternative model, we test the traditional model with basal partial melt; and we show that the two models cannot be distinguished from each other by the available selenodetic measurements. Additional insight into the nature of lunar tidal dissipation can be gained either by measurements of higher-degree Love numbers and quality factors or by farside lunar seismology.

Is there a semi-molten layer at the base of the lunar mantle?

Michaela Walterová¹, Marie Běhounková², Michael Efroimsky³

 $^1 \rm Institute$ of Planetary Research, German Aerospace Center (DLR), Berlin, Germany $^2 \rm Department$ of Geophysics, Faculty of Mathematics and Physics, Charles University, Prague, Czech Republic ³US Naval Observatory, Washington DC, USA

Key	Points:
-----	----------------

1

2

3

4 5

> 6 7

8

9	• A lunar mantle governed by the Andrade model fits selenodetic constraints only
10	with a very weak frequency dependence of tidal dissipation
11	• We seek the parameters of the Sundberg-Cooper model that would explain the anoma
12	lous frequency dependence of tidal Q measured by LLR
13	• Both a dissipative basal layer and elastically-accommodated grain-boundary slid-
14	ing in the deep mantle result in the same tidal response

Corresponding author: Michaela Walterová, kanovami@gmail.com

15 Abstract

Parameterised by the Love number k_2 and the tidal quality factor Q, and inferred from 16 lunar laser ranging (LLR), tidal dissipation in the Moon follows an unexpected frequency 17 dependence often interpreted as evidence for a highly dissipative, melt-bearing layer en-18 compassing the core-mantle boundary. Within this, more or less standard interpretation, 19 the basal layer's viscosity is required to be of order 10^{15} to 10^{16} Pa s and its outer ra-20 dius is predicted to extend to the zone of deep moonquakes. While the reconciliation of 21 those predictions with the mechanical properties of rocks might be challenging, alterna-22 tive lunar interior models without the basal layer are said to be unable to fit the frequency 23 dependence of tidal Q. 24

The purpose of our paper is to illustrate under what conditions the frequency-dependence 25 of lunar tidal Q can be interpreted without the need for deep-seated partial melt. De-26 vising a simplified lunar model, in which the mantle is described by the Sundberg-Cooper 27 rheology, we predict the relaxation strength and characteristic timescale of elastically-28 accommodated grain boundary sliding in the mantle that would give rise to the desired 20 frequency dependence. Along with developing this alternative model, we test the tra-30 ditional model with basal partial melt; and we show that the two models cannot be dis-31 tinguished from each other by the available selenodetic measurements. Additional in-32 sight into the nature of lunar tidal dissipation can be gained either by measurements of 33 higher-degree Love numbers and quality factors or by farside lunar seismology. 34

³⁵ Plain Language Summary

As the Moon raises ocean tides on the Earth, the Earth itself gives rise to periodic 36 deformation of the Moon. Precise measurements of lunar shape and motion can reveal 37 those deformations and even relate them to our natural satellite's interior structure. In 38 this work, we discuss two interpretations of those measurements. According to the first 39 one, the lunar interior is hot and there is a thick layer of partial melt or other weak ma-40 terial buried more than 1000 km deep under the lunar surface. According to the second 41 one, there is no such layer, and the measured deformation can be explained by the be-42 haviour of solid rocks at relatively low temperatures. We show that the two possibili-43 ties cannot be distinguished from each other by the existing data. 44

45 **1** Motivation

Fitting of the lunar laser ranging (LLR) data to the quality-factor power scaling law $Q \sim \chi^p$ rendered a small *negative* value of the exponential: p = -0.19 (Williams et al., 2001). Further attempts by the JPL team to reprocess the data led to p = -0.07. According to Williams and Boggs (2009),

50 51

⁵¹ " Q for rock is expected to have a weak dependence on tidal period, but it is ex-⁵² pected to decrease with period rather than increase."

The most recent estimates of the tidal contribution to the lunar physical librations (Williams & Boggs, 2015) still predict a mild increase of Q with period: from $Q = 38\pm$ 4 at one month to $Q = 41 \pm 9$ at one year, yielding $p = -0.03 \pm 0.09$.

Efroimsky (2012a, 2012b) suggested that since the frequency-dependence of k_2/Q always has a kink shape, like in Figure 1, the negative slope found by the LLR measurements could be consistent with the peak of the kink residing between the monthly and annual frequencies. This interpretation entails, for a Maxwell or Andrade moon, very low values of the mean viscosity, indicating the presence of partial melt. Our goal now is to devise an interpretation based on the Sundberg-Cooper model. Within that model, the kink contains not one but two peaks, and we are considering the possibility that the negative slope of our interest is due to the monthly and annual frequencies bracketing either this peak or the local inter-peak minimum. (It is unlikely that both of these frequencies are located on the negative-slope side of the peak, because the slope of that peak is too steep.)

67 2 Introduction

68

2.1 Overview of Previous Works

The knowledge of the interior structure of the Moon is essential for understand-69 ing its thermal, geochemical, and orbital evolution as well as the coupled evolution of 70 the Earth-Moon system. The proximity of our natural satellite to the Earth has also made 71 it a frequent target of geophysical exploration. A large amount of data was collected by 72 lunar seismic stations, deployed by the Apollo missions, that were functional for several 73 years between 1972 and 1977 (for a review, see, e.g., Garcia et al., 2019; Nunn et al., 2020). 74 Other constraints are being placed by selenodetic measurements or by geochemical and 75 petrological considerations. However, the deepest interior of the Moon still remains some-76 what mysterious. Although different models based on the inversion of seismic travel times 77 generally agree on the lunar mantle structure down to ~ 1200 km, below these depths 78 they start to diverge greatly (Garcia et al., 2019). 79

After the acquisition of the first data by the lunar seismic network, it was pointed 80 out by Nakamura et al. (1973, 1974) that direct shear-waves from the farside of the Moon 81 are not being detected by some of the near-side seismometers. Moreover, deep moonquakes, 82 a class of tidally-triggered seismic events originating at around 1000 km depth, were al-83 most absent on the farside. This puzzling phenomenon was interpreted by Nakamura et 84 al. (1973) as an indication for a shear-wave shadow zone caused by a highly attenuat-85 ing region around the core. Later, Nakamura (2005) reported his further efforts to find 86 farside moonquakes among the discovered nests of deep moonquakes. Having had iden-87 tified about 30 nests likely to be on the farside, his updated analysis still demonstrated 88 that either the region of the Moon's deep interior within about 40 degrees from the an-89 tipodes (the centre of the farside) is nearly aseismic or a portion of lunar lower mantle 90 severely attenuates or deflects seismic waves. Lunar seismic data were also reprocessed 91 by Weber et al. (2011) and Garcia et al. (2011). However, while Weber et al. (2011) also 92 found evidence for deep mantle layering and a strongly attenuating zone at the mantle 93 base, Garcia et al. (2011) did not include such a feature in their lunar interior model. 94 The discussion about the seismic evidence for a strongly attenuating zone is thus still 95 ongoing (Garcia et al., 2019). 96

Several authors argued for the existence of a low-velocity zone (LVZ) at the base 97 of the mantle also on other than seismological grounds. They linked it to partial melt-98 ing in deep lunar interior, which might be triggered either by tidal dissipation (Harada 99 et al., 2014), or by the presence of incompatible, radiogenic elements buried after an an-100 cient mantle overturn. The idea of an overturn has been suggested by numerical mod-101 elling of magma ocean solidification with the emplacement of ilmenite-bearing cumulates 102 above core-mantle boundary. Moreover, it is potentially supported by observations of 103 near-surface gravity anomalies (Zhang et al., 2013). 104

Evidence for a low-rigidity/low-viscosity zone has also been sought in the lunar libration signal obtained by LLR (e.g., Williams et al., 2001; Williams & Boggs, 2015), and in selenodetic measurements (including orbiter tracking) that are sensitive to the lunar gravity field and tidal deformation (e.g., Konopliv et al., 2013; Lemoine et al., 2013). One of the most surprising findings resulting from fitting the LLR data was the low value and unexpected frequency dependence of the tidal quality factor Q, as mentioned in Section 1 above. The inferred frequency dependence can be explained by a low effective vis-

cosity of the Moon (Efroimsky, 2012a, 2012b), or by the presence of a secondary peak 112 in the dissipation spectrum (e.g., Williams & Boggs, 2015), possibly caused by the pu-113 tative basal layer (Harada et al., 2014; Matsumoto et al., 2015). Earlier results from LLR 114 indicated that the lunar core-mantle boundary (CMB) might still be out of equilibrium, 115 which would imply long relaxation times and high lower-mantle viscosities, in contra-116 diction to the presence of a partial melt. However, this hypothesis is not supported by 117 more recent evaluations of LLR data (Viswanathan et al., 2019), showing a CMB at hy-118 drostatic equilibrium. 119

Despite relative consistency of the evidence for and the theoretical expectation of 120 a highly dissipative basal layer, alternative models of a "melt-free" Moon have been pro-121 posed (Nimmo et al., 2012; Karato, 2013; Matsuyama et al., 2016). One argument for 122 high values of lower-mantle viscosities comes from the observations of deep moonquakes. 123 Kawamura et al. (2017) reevaluated an ensemble of moonquakes occurring at depths be-124 tween 750 and 1200 km and found a brittle-ductile transition temperature of approxi-125 mately 1240-1275 K, implying a cold lunar interior with temperatures below solidus of 126 dry peridotite. Moreover, the employment of a realistic, microphysically substantiated 127 models of the tidal response (Nimmo et al., 2012) can explain the low tidal Q and the 128 observed k_2 of the Moon without requiring the existence of a weak basal layer, which is 129 necessitated in some of the other studies by the model settings and the simplified rhe-130 ological assumptions. 131

A feature of the selenodetic measurements that is difficult to explain without the 132 existence of a highly dissipative basal layer is the aforementioned frequency dependence 133 of the lunar Q, repeatedly derived from LLR measurements in the series of works by Williams 134 et al. (2001); Williams and Boggs (2009); Williams et al. (2014), and Williams and Boggs 135 (2015). Even an independent implementation of the LLR software by Pavlov et al. (2016) 136 predicts the same value of Q for the monthly period as for the annual period, which is 137 still not consistent with the expected frequency dependence of tidal dissipation in melt-138 free silicates. 139

In the absence of other than LLR-based data on the lunar Q, the most plausible explanation for the unexpected frequency dependence might still be an observational uncertainty, rather than an effect contained in a tidal model. Nevertheless, in this work, we shall explore two possible implications of the frequency dependence under the explicit assumption that the fitted values are a result of a natural phenomenon and not of a model's limitations or an observation error.

146

2.2 A Putative Weak Basal Layer: Pros and Contras

The following paragraphs review the last ten years of discussion about the presence or absence of a low-viscosity basal layer, with the argumentation derived mainly from the lunar tidal response.

We begin by noting that a negative value of the exponent in $Q \sim \chi^p$ is impossi-150 ble for the *seismic* quality factor of rocks obeying simple rheologies like the Maxwell or 151 And rade models. This can be easily understood if we express the seismic Q via the real 152 and imaginary parts of the complex compliance (Efroimsky, 2015, eqn 46). By insert-153 ing into this expression either the Maxwell model or any other simple model lacking peaks, 154 we obtain a monotonic function $Q(\chi)$. On the other hand, even for simple rheologies the 155 exponential p can assume negative values if we are fitting to the $Q \sim \chi^p$ law not a seis-156 mic but a *tidal* quality factor (Efroimsky, 2015, eqn 45). The tidal Q tends to zero at 157 both very low and very high loading frequencies χ , and has a maximum in between. The 158 maximum is called into being by interplay of rheology and self-gravity. 159

This theoretical frequency dependence of the tidal quality factor motivated Efroimsky (2012a, Section 5.2) to hypothesise that the small negative exponent p reported by Williams et al. (2001) and Williams and Boggs (2009) may result from a proximity of the major tidal frequencies in the Moon to the frequency delimiting the peak dissipation. Efroimsky (2012a, Section 5.7) also noted that this interpretation would imply a low effective viscosity of the Moon (modeled with a homogeneous body governed by the Maxwell or the combined Maxwell-Andrade rheology), with an estimated value of $\eta = 3 \times 10^{15}$ Pa s. Such a low viscosity would support seismic models containing a layer of partial melt (Nakamura et al., 1974; Weber et al., 2011).

Nimmo et al. (2012) aimed at answering the question whether basal partial melt 169 is indeed required for reproducing the tidal data, and studied the effect of lunar ther-170 171 mal structure on the seismic and tidal Q. They described the rheology of the lunar interior with the extended Burgers model of Jackson et al. (2010), which contains an ab-172 sorption band corresponding to high-temperature background, as well as an additional 173 low-temperature peak. The peak represents the elastically-accommodated grain bound-174 ary sliding, a phenomenon that will be considered also in our work, although within an-175 other rheology. Nimmo et al. (2012) further considered a radially heterogeneous elastic 176 structure of the mantle and accounted for the temperature, pressure, and grain-size-177 dependence of the characteristic relaxation times. Using this model, they were able to 178 match the tidal Love numbers k_2 and h_2 and the monthly quality factor, and they also 179 deduced that the lower-mantle viscosity should be as high as 10^{23} Pa s and must be in-180 creasing towards the surface. However, the model used did not succeed in fitting the un-181 expected slope of Q as a function of frequency. Although the authors showed that a model 182 case with grain size of $1 \,\mathrm{mm}$ (instead of their baseline value of $1 \,\mathrm{cm}$) would imply a neg-183 ative value of the exponential, p = -0.02, they dismissed this model as a poor fit to 184 both k_2 and Q. Moreover, they argued that the smaller grain size would not match the 185 tentative observation of unrelaxed CMB (Williams et al., 2012). 186

An original explanation of the high tidal dissipation in the Moon was provided by 187 Karato (2013), who linked the measurements of electrical conductivity and Q to the wa-188 ter content in the lunar mantle. That the water content might not be as low as had been 189 presumed in earlier models was illustrated by geochemical studies of lunar samples, and 190 Karato (2013) combined this observation with his own results to propose a new theory 191 of lunar formation. Using the observational constraints on Q and electrical conductiv-192 ity, he further concluded that the temperature at a 800 km depth of the lunar mantle is 193 ~ 1200–1500 K for a water content between 10^{-3} and 10^{-2} wt.%. Karato (2013) was scep-194 tical to the idea of partial melting at the base of the lunar mantle. He argued that the 195 melt-bearing seismic model of Weber et al. (2011) would require more than $\sim 1\%$ of melt 196 and that retaining such an amount of melt would be difficult due to efficient compaction. 197 Regarding the frequency-dependence of Q, Karato (2013) rejected the models of Efroimsky 198 (2012a) and Nimmo et al. (2012) and suggested that the negative exponent p might be 199 caused by non-linear anelasticity of the monthly tide and linear anelasticity of the an-200 nual tide. However, this idea was partly based on the incorrect assumption that the tide 201 at the annual frequency is due to Sun-raised tidal deformation of the Moon. As explained 202 by Williams et al. (2001), the annual modulation is produced by solar perturbations to 203 lunar orbit only. The annual tide is thus raised by the Earth, just as the monthly tide. 204 Still, the remark on a possible non-linearity of the lunar tide remains valid. 205

Adopting the density and rigidity profiles from a 10-layer structural model by Weber 206 et al. (2011), Harada et al. (2014) explored the possible effects of a low-viscosity layer 207 at the base of the mantle. To keep the number of unknowns reasonable, the authors set 208 constant viscosity values for the lithosphere, mantle, low-viscosity layer, outer core, and 209 inner core, and applied the Maxwell rheological model. They then calculated the tidal 210 parameters for various thicknesses (outer radii 450-500 km) and viscosities ($10^9-10^{21} \text{ Pa s}$) 211 of the basal layer, at both the monthly and annual tidal frequencies, assuming that the 212 rest of the mantle has a constant viscosity of $\eta = 10^{21}$ Pa s. With the highest consid-213 ered basal layer thickness ($D_{\rm LVZ} = 170 \,\rm km$) and a viscosity of about $2 \times 10^{16} \,\rm Pa$ s, Harada 214 et al. (2014) were able to reproduce the quality factors given by Williams et al. (2001) 215 as well as their frequency dependence. Their value for the Love number at the monthly 216 period falls into the interval $k_2 = 0.0242 \pm 0.0004$ suggested by Yan et al. (2012), while 217

their value of the Love number at the annual period fits into the interval $k_2 = 0.0255 \pm$ 0.0016 observed by Goossens et al. (2011). Viscoelastic, the model of Harada et al. (2014) rendered different values of k_2 at the monthly and annual frequencies. This said, neither Yan et al. (2012) nor Goossens et al. (2011) considered frequency-dependence of their empirical values of k_2 .

An updated version of the forward-modelling approach by Harada et al. (2014) was 223 presented in Harada et al. (2016). Using the improved set of tidal parameters (limits on 224 Q at four tidal frequencies and the values of k_2 , k_3 , and h_2 at the monthly frequency), 225 the estimate of the basal layer's outer radius was expanded from $500\,\mathrm{km}$ to $540-560\,\mathrm{km}$ 226 (i.e., layer thickness $D_{\rm LVZ} = 210 - 230 \,\rm km$ for a core radius of $330 \,\rm km$) and the corre-227 sponding basal viscosity slightly changed to 3×10^{16} Pa s. In a recent follow-up study, 228 Tan and Harada (2021) considered full radial profile of the lunar interior (Weber et al.. 229 2011; Garcia et al., 2011) and assumed a temperature-dependent viscosity structure of 230 the basal layer. The viscosity structure either followed a convective temperature profile 231 (viscosity almost constant with depth) or a conductive profile (linear decrease of viscos-232 ity with depth). Since the former model was shown to match the selenodetic data bet-233 ter, the authors argued that the low-viscosity layer should be locally convecting. More-234 over, they concluded that the layer's outer radius reaches 560 or 580 km (that is, to the 235 depths of ~ 1160 km) and that the viscosity is the same as found by Harada et al. (2016). 236

The question whether a basal partial melt is required by the selenodetic data was 237 also raised by Khan et al. (2014), though with an answer different from Nimmo et al. 238 (2012). Khan et al. (2014) concentrated on detailed modelling of the lunar mantle petrol-239 ogy, and performed a Bayesian inversion of the mean density, the moment of inertia, the 240 apparent resistivity, and the tidal data $(k_2 \text{ and } Q)$ at the monthly period. To model the 241 tidal response of the lunar mantle within a purely elastic model, they calculated an anelas-242 tic correction to k_2 based on a homogeneous spherical model and the power-law depen-243 dence of tidal dissipation, which is valid for large *seismic* quality factors (or weak seis-244 mic wave attenuation; Zharkov & Gudkova, 2005). For cases with the Andrade param-245 eter $\alpha > 0.1$, the resulting elastic k_2 clearly implied the existence of a partial melt in 246 a basal layer with the thickness of 150-200 km (i.e., a depth range $\sim 1250-1400$ km 247 or the outer radii between \sim 340-490 km). Khan et al. (2014) also found that, in order 248 to be neutrally buoyant, the partially molten material should be enriched in FeO and 249 TiO_2 with respect to the bulk mantle. In addition to the models with a partially molten 250 layer, the authors tested a model with a fully solid mantle: this model still fitted all ob-251 servations, except for the anelastically-corrected k_2 . 252

Similarly, Matsumoto et al. (2015) performed a Bayesian inversion of seismic travel 253 times and a set of available selenodetic data (mean density, moment of inertia, k_2 , and 254 Q at the monthly and annual frequencies), to infer the interior structure of an eight-layered 255 lunar model. As in Harada et al. (2014), the authors considered the Maxwell rheolog-256 ical model, in which the existence of a low-viscosity layer is required not only by the slope 257 of Q's frequency dependence but also by the magnitude of k_2 . The viscosity of the solid 258 mantle was always set to 10^{21} Pa s; otherwise, Matsumoto et al. (2015) varied a wide range 259 of parameters. While their inverted structure of the shallow mantle agrees with the re-260 sults of Weber et al. (2011) and Garcia et al. (2011), the lower mantle, mainly constrained 261 by selenodetic data, slightly differs from the melt-containing model of Weber et al. (2011). 262 The outer radius of the highly dissipative layer is around 570 km and the predicted viscosity in that region reaches $2.5^{+1.5}_{-0.9} \times 10^{16}$ Pa s. The authors noted that with the model 263 264 used, k_2 and the annual Q are slightly biased from the observed values, although not be-265 yond 1σ . Matsumoto et al. (2015) also reported a trade-off between the outer core ra-266 dius and the LVZ thickness. The thickness of the LVZ corresponding to the calculated 267 outer radius is at least 170 km and, for the core size estimate of Weber et al. (2011), it 268 may reach 240 km. 269

In a paper presenting their interpretation of LLR data, Williams and Boggs (2015) compared several rheological models and endeavoured to fit the lunar k_2/Q at the monthly

and annual tidal periods, considering physical libration at five periods (1 month, 206 days, 272 1 year, 3 years, and 6 years). Aware of the complex properties of the lunar interior and 273 the possible unmodelled effects of its lateral heterogeneity, the authors proposed a model 274 consisting of an absorption band and a narrow Debye peak: the former characterising 275 the dissipation in the solid mantle, the latter describing the contribution of the partially 276 molten layer suggested by Harada et al. (2014). For the thickness of the partially molten 277 layer, Williams and Boggs (2015) obtained $D_{LVZ} \ge 205 \text{ km}$, placing its outer radius at 278 $\geq 535 \,\mathrm{km}.$ 279

280 The results of Williams and Boggs (2015) are relatively consistent with the predictions by Harada et al. (2014); Matsumoto et al. (2015), and Harada et al. (2016). As 281 in the other studies containing a LVZ, they indicate that if partial melt is present, it might 282 extend to the zone of deep moonquakes. On the one hand, the coexistence of partially 283 molten material with seismic sources is hard to imagine: while the former requires that 284 the lower-mantle temperatures exceed solidus, the latter should be concentrated in re-285 gions where the mantle rocks undergo brittle deformation, limited to lower temperatures. 286 On the other hand, the movement of small amounts of melt to the zone of moonquake 287 nests might be considered one of the mechanisms triggering seismic events. Frohlich and 288 Nakamura (2009) proposed an explanation for the periodic occurrence of deep moonquakes, 289 which combines dehydration embrittlement due to partial melting and crack opening by 290 moving fluids. The authors pointed out the correlation between tidal loading and seis-291 mic events associated with magma movements in terrestrial volcanoes and remarked that 292 a similar process may be active in the lunar interior. Tentative evidence for a link be-293 tween deep moonquakes and magma movements might also be seen in the correlation 294 between the locations of deep moonquake nests and lunar maria (Qin et al., 2012). How-295 ever, a definitive answer to the question of whether a rheologically weak layer and seis-296 mic sources can exist at comparable depths awaits further modelling efforts. 297

The specific effect of a partially-molten basal layer on the *elastic* Love number $k_{2,e}$ 298 was discussed in the study of Raevskiy et al. (2015), which combined seismic and geode-299 tic data with models of lunar mantle composition. Depending on the model used, the 300 rigidity of the basal layer was required to be 20-50% lower than the rigidity of the over-301 lying solid mantle and the outer radius of that zone was determined to reach 530-550 km. 302 From the petrological perspective, the authors argued that partial melting of a peridotite/harzburgite 303 mantle above the core-mantle boundary (CMB) would require temperatures in the depth 304 of 1000 km to be in the range of 1350–1400 °C, unless the temperature gradients in the 305 lower mantle become steeper. Furthermore, they concluded that the seismic velocities 306 of Weber et al. (2011) are inconsistent with temperature profiles approaching solidus at 307 the CMB. Although the models of Raevskiy et al. (2015) assume elastic response of the 308 Moon, the authors also mentioned that anelasticity might explain the observed Love num-309 ber without the need for a basal semi-molten layer. 310

Matsuyama et al. (2016) constrained their lunar interior models by the elastic Love 311 numbers k_2 and h_2 (calculated using the same anelastic correction for Q at the monthly 312 period as in Khan et al., 2014), the mean density of the Moon, and the moment of in-313 ertia. After carrying out MCMC-type inversion, the authors concluded that although 314 the chosen observables do not rule out the existence of a semi-molten layer, there is a 315 strong preference for higher, solid-mantle-like values of the lower-mantle rigidity. If the 316 semi-molten layer exists, its thickness calculated by Matsuyama et al. (2016) is $D_{\rm LVZ} = 194^{+66}_{-186}$ km, its rigidity is $\mu_{\rm LVZ} = 43^{+26}_{-9}$ GPa, and its density may reach exceptionally high values, $\rho_{\rm LVZ} = 4676^{+410}_{-1179}$ kg m⁻³. 317 318 319

Recently, the combined geochemical, seismic, and selenogetic ensemble of Raevskiy et al. (2015) was further studied by Kronrod et al. (2022), who extended the former work by considering explicitly a viscoelastic lunar interior. Regarding the division into interior layers and the adopted rheological model, the authors followed Matsumoto et al. (2015); i.e., they assumed the Maxwell model for the mantle and included a semi-molten basal layer. Besides the main results of their Bayesian analysis, indicating a major difference in the chemical composition of the bulk silicate Earth and the Moon, Kronrod et al. (2022) presented probability distributions for the seismic wave velocities, mean density, and the thickness of the basal layer. The resulting distributions are wide, constraining the basal layer's density to $3400-3800 \text{ kg m}^{-3}$ and the thickness to 100-350 km, depending on the mantle composition. As in Khan et al. (2014), the authors conclude that the layer should be enriched in TiO₂ and FeO, if it is present.

In summary, the literature discussing the unexpected frequency dependence of lu-332 nar tidal Q as well as the properties of a hypothetical semi-molten layer atop the lunar 333 core is rich, and the proposed values of the layer's thickness range from 0 to 350 km. Mod-334 els considering linear viscoelastic Maxwell rheology (both for the basal layer and for the 335 bulk mantle; Harada et al., 2014, 2016; Matsumoto et al., 2015; Tan & Harada, 2021) 336 typically arrive at viscosities of order 10^{16} Pa s. If the semi-molten layer exists, its up-337 per radius extends to the depths of ~ 1150 km, i.e., just below the regions that are rel-338 atively well mapped by seismological studies and contain the nests of tidally-triggered 339 deep moonquakes. Nevertheless, the existence of a low-viscosity layer is not necessarily 340 required by selenodetic measurements at the best accessible, monthly period (Nimmo 341 et al., 2012; Matsuvama et al., 2016). The main advantage of melt-bearing models lies 342 in their ability to explain the possible increase in tidal Q from the monthly to the an-343 nual period. 344

2.3 Lunar k_2 and Q

345

Here, we shall use the potential tidal Love number derived from the GRAIL mis-346 sion tracking data. Two independent analyses performed by the JPL group (Konopliv 347 et al., 2013, the GL0660B solution) and the GSFC group (Lemoine et al., 2013, the GRGM660PRIM 348 solution) yielded two possible values of the parameter: $k_2 = 0.02405 \pm 0.000176$ and 349 $k_2 = 0.02427 \pm 0.00026$, respectively. The unweighted mean of the two alternative val-350 ues is $k_2 = 0.02416 \pm 0.000222$ for a reference radius of 1738 km, and $k_2 = 0.02422 \pm 0.02422$ 351 0.000222 for the actual mean radius of 1737.151 km (Williams et al., 2014). For compar-352 ison, the recent analysis of the data from the Chang'e 5T1 mission gives $k_2 = 0.02430 \pm$ 353 0.0001 (Yan et al., 2020). We note that the value obtained from satellite tracking data 354 corresponds, in particular, to the real part of the complex Love number introduced later 355 in Subsection 4.1. The GRAIL data are dominated by data arcs collected throughout 356 a one-month time interval, and the resulting k_2 is thus interpreted as indicative of the 357 deformation at monthly frequency (A. Konopliv, private communication). 358

The tidal quality factor Q was obtained by fitting tidal contribution to lunar phys-359 ical libration measured by LLR (Williams et al., 2001, 2014; Williams & Boggs, 2015). 360 Interpreting the measurements of physical libration presents a highly complex problem, 361 depending on cross interactions of tides raised by the Earth and the Sun, precise mod-362 eling of the lunar orbit and of the instantaneous positions of the Earth-based stations 363 and the Moon-based retroreflectors, and on an adequate incorporation of the lunar core-364 mantle friction (Williams et al., 2001). In practice, the tidal time delay at a monthly pe-365 riod and the dissipation-related corrections to the periodic latitudinal and longitudinal 366 variations in the Moon's orientation are outputted and related analytically to linear com-367 binations of k_2/Q at a number of loading frequencies. Since many of the loading frequen-368 cies are close to each other, the periodic corrections enable approximate estimation of 369 the leading dissipation terms. Specifically, the strongest correction (compared to its un-370 certainty) is related to the annual longitudinal libration. Assuming a fixed k_2 at the monthly 371 frequency, equal to the above-mentioned unweighted average, and using a complex rhe-372 373 ological model best fitting the dissipation-related corrections to libration angles, Williams and Boggs (2015) derived the following frequency-dependent values of tidal quality fac-374 tor: $Q = 38 \pm 4$ at the period of 1 month, $Q = 41 \pm 9$ at 1 year, and lower bounds of 375 $Q \ge 74$ at 3 years and $Q \ge 58$ at 6 years. The tidal quality factors at other than the 376 monthly frequency are model-dependent because the actual quantities extracted from 377

the dissipation-related corrections to libration angles are the ratios $(k_2/Q)_{\chi}/(k_2/Q)_{\text{monthly}}$, where χ denotes frequency.

Williams and Boggs (2015) also attempted to find the frequency-dependence of k_2 ; however, the effect could not be detected by existing measurements. We note that in contrast to the unexpected frequency dependence of Q found with the JPL-based software (Williams et al., 2001, 2014; Williams & Boggs, 2015), an independent implementation of the fitting tool with different preset solutions for part of the geophysical phenomena (Pavlov et al., 2016) predicted Q = 45 at both the monthly and the annual frequencies.

As an additional, though a relatively weak constraint on the lunar interior struc-387 ture, we consider the degree-3 potential tidal Love number k_3 and the degree-2 defor-388 mational Love number h_2 corresponding to radial deformation. The former has been de-389 rived from GRAIL mission tracking data and, as with k_2 above, we adopt the unweighted 390 average of the two existing independent solutions (Lemoine et al., 2013; Konopliv et al., 391 2013): $k_3 = 0.0081 \pm 0.0018$. The latter has been measured by LLR and by laser al-392 timetry (Mazarico et al., 2014; Pavlov et al., 2016; Viswanathan et al., 2018; Thor et al., 393 2021), the most recent value, presented by Thor et al. (2021), being $h_2 = 0.0387 \pm 0.0025$. 394

We would finally mention the reason why the constraints on the lunar interior from the measurements of k_3 are weak. A degree-*l* component of the internal tidal potential is proportional to r^l , where *r* is the distance between the centres of mass of the tidally perturbed body and the perturber. For this reason, with increasing degree *l*, the shallower depths contribute more and more to the Love numbers k_l . The sensitivity of the higher-degree Love numbers to the deep interior is, therefore, limited as compared to degree 2.

402

2.4 Outline of This Work

After an overview of the models and interpretations proposed in recent literature 403 (with the focus on the last ten years of the discussion), we are ready to continue with 404 the central part of this project. Our plan is to provide an interpretation of the unexpected 405 frequency dependence of tidal Q which does not require partial melting (in a way sim-406 ilar to Nimmo et al., 2012) and compare it with a model containing a highly dissipative 407 basal layer (Harada et al., 2014; Matsumoto et al., 2015). Section 3 introduces and gives a justification for the rheological model employed. Namely, it discusses the Sundberg-409 Cooper extension of the Andrade model and the dissipation related to elastically accom-410 modated grain-boundary sliding (GBS). The following Section 4 links the non-elastic rhe-411 ology to Love numbers and tidal quality factors. In Section 5, we first illustrate the ex-412 pected position of a secondary peak in the dissipation spectrum of a homogeneous Moon, 413 and then attempt to find the parameters of two- or three-layered lunar models that would 414 produce the values of the monthly tidal Q and annual k_2/Q reported by Williams and 415 Boggs (2015). At the same time, we fit the empirical values of lunar k_2 , k_3 , and h_2 given 416 in Subsection 2.3. Section 6 discusses implication of both our models, and the results 417 are briefly summarised in Section 7. 418

3 General Facts on Rheologies

420 **3.1 Constitutive Equation**

⁴²¹ Rheological properties of a material are encoded in a constitutive equation inter-⁴²² connecting the present-time deviatoric strain tensor $u_{\gamma\nu}(t)$ with the values that have ⁴²³ been assumed by the deviatoric stress $\sigma_{\gamma\nu}(t')$ over the time period $t' \leq t$. Under linear deformation, the equation has the form of convolution, in the time domain:

$$2 u_{\gamma\nu}(t) = \hat{J}(t) \ \sigma_{\gamma\nu} = \int_{-\infty}^{t} \dot{J} \ (t - t') \ \sigma_{\gamma\nu}(t') \ dt' \ , \tag{1}$$

⁴²⁶ and the form of product, in the frequency domain:

$$2 \bar{u}_{\gamma\nu}(\chi) = \bar{J}(\chi) \bar{\sigma}_{\gamma\nu}(\chi) \quad . \tag{2}$$

Here $\bar{u}_{\gamma\nu}(\chi)$ and $\bar{\sigma}_{\gamma\nu}(\chi)$ are the Fourier images of strain and stress, while the complex compliance $\bar{J}(\chi)$ is a Fourier image of the kernel $\dot{J}(t-t')$ of the integral operator (1), see, e.g., Efroimsky (2012a, 2012b) for details.

3.2 The Maxwell and Andrade Models

At low frequencies, deformation of most minerals is viscoelastic and obeys the Maxwell model:

$$\dot{\mathbb{U}} = \frac{1}{2\,\mu} \,\,\dot{\mathbb{S}} + \frac{1}{2\,\eta}\,\mathbb{S} \tag{3a}$$

435 or, equivalently:

425

427

431

434

436

440

443

446

449

455

$$\dot{\mathbb{S}} + \frac{1}{\tau_{_{\mathrm{M}}}} \mathbb{S} = 2\,\mu \,\,\dot{\mathbb{U}} \quad, \tag{3b}$$

⁴³⁷ U and S being the deviatoric strain and stress; η and μ denoting the viscosity and ⁴³⁸ rigidity. (Below, we shall address the question as to whether μ is the unrelaxed or re-⁴³⁹ laxed rigidity.) The *Maxwell time* is introduced as

$$\tau_{\rm M} \equiv \frac{\eta}{\mu} \ . \tag{4}$$

For this rheological model, the kernel of the convolution operator (1) is a time derivative of the compliance function

$$^{(M)}J(t-t') = \left[J_{e} + (t-t') \frac{1}{\eta}\right] \Theta(t-t') \quad , \tag{5}$$

where $\Theta(t - t')$ is the Heaviside step function, while the elastic compliance J_e is the inverse of the shear rigidity μ :

$$J_{\rm e} \equiv \frac{1}{\mu} \ . \tag{6}$$

In the frequency domain, equation (3) can be cast into form (2), with the complex compliance given by

$$^{(M)}\bar{J}(\chi) = J_e - \frac{i}{\eta\chi} = J_e \left(1 - \frac{i}{\chi\tau_{\rm M}}\right) , \qquad (7)$$

and the terms $J_{\rm e}$ and $-i/(\eta \chi)$ being the elastic and viscous parts of deformation, correspondingly. So a Maxwell material is elastic at high frequencies, viscous at low.

452 More general is the combined Maxwell-Andrade rheology, often referred to simply 453 as the Andrade rheology. It comprises inputs from elasticity, viscosity, and anelastic pro-454 cesses: (A) = t

$$^{A)}J(t) = J_{e} + \beta t^{\alpha} + \frac{t}{\eta} , \qquad (8)$$

456 the corresponding complex compliance being

⁴⁵⁹ =
$$J_{\rm e} + \beta (i\chi)^{-\alpha} \Gamma (1+\alpha) - i J (\chi \tau_{\rm M})^{-1}$$
, (9b)

where Γ is the Gamma function, while α and β denote the dimensionless and dimensional Andrade parameters.

Expressions (9a - 9b) suffer an inconvenient feature, the fractional dimensions of the parameter β . It was therefore suggested in Efroimsky (2012a, 2012b) to shape the compliance into a more suitable form

$$^{(A)}J(t) = J_{\rm e} \left[1 + \left(\frac{t}{\tau_{\rm A}}\right)^{\alpha} + \frac{t}{\tau_{\rm M}}\right] \Theta(t - t') , \qquad (10)$$

466 467

465

$${}^{(A)}\bar{J}(\chi) = J_{\rm e} \left[1 + (i\,\chi\,\tau_{\rm A})^{-\alpha}\,\Gamma(1+\alpha) - i\,(\chi\,\tau_{\rm M})^{-1} \right] , \qquad (11)$$

with the parameter τ_A christened as the Andrade time and linked to β through

$$\beta = J_e \tau_{\scriptscriptstyle A}^{-\alpha} \quad . \tag{12}$$

⁴⁷⁰ Compliance (11) is identical to (9a) and (9b), but is spared of the parameter β of fractional dimensions.

472

476

481

483

484

485

486

469

3.3 Why the Maxwell and Andrade Models Require Refinement

In the literature, it is common to postulate that both the rigidity and compliance assume their *unrelaxed* values denoted with $\mu_{\rm U}$ and $J_{\rm U}$.

475 This convention is reasonable for sufficiently high frequencies:

$$\chi$$
 is high $\implies \mu = \mu_{\rm U}$ and $J_{\rm e} = J_{\rm U}$. (13)

The convention, however, becomes unjustified for low frequencies. In that situation, the material has, at each loading cycle, enough time to relax, wherefore both the rigidity modulus and its inverse assume values different from the unrelaxed ones. In the zero-frequency limit, they must acquire the relaxed values:

$$\chi \to 0 \implies \mu \to \mu_{\rm R} \quad \text{and} \quad J_{\rm e} \to J_{\rm R} \quad .$$
 (14)

⁴⁸² This fact must be taken care of, both within the Maxwell and Andrade models.

3.4 Generalisation of the Maxwell and Andrade Models, According to Sundberg and Cooper (2010)

The simplest expression for the time relaxation of the elastic part of the compliance is

$$J_{\rm e}(t) = J_{\rm U} + (J_{\rm R} - J_{\rm U}) \left[1 - e^{-t/\tau}\right]$$
 (15a)

491

49

 $= J_{\rm U} \left[1 + \Delta \left(1 - e^{-t/\tau} \right) \right] , \qquad (15b)$

⁴⁹⁰ where the so-called relaxation strength is introduced as

$$\Delta \equiv \frac{J_{\rm R}}{J_{\rm U}} - 1 \quad , \tag{16}$$

⁴⁹² while τ is the characteristic relaxation time. When relaxation of $J_{\rm e}$ is due to elastically ⁴⁹³ accommodated grain-boundary sliding, this time can be calculated as

$$\tau = \tau_{\rm gbs} = \frac{\eta_{\rm gb} d}{\mu_{\rm U} \delta} , \qquad (17)$$

where $\eta_{\rm gb}$ is the grain-boundary viscosity, d is the grain size, while δ is the structural width of the grain boundary. In the frequency domain, this compliance writes as

$$\bar{J}_{\rm e}(\chi) = J_{\rm U} \left[1 + \frac{\Delta}{1 + \chi^2 \tau^2} + i \frac{\chi \tau \Delta}{1 + \chi^2 \tau^2} \right] , \qquad (18)$$

its imaginary part demonstrating a Debye peak. Our goal is to trace how this Debye peak 499 translates into the frequency-dependence of the inverse tidal quality factor 1/Q and of 500 k_2/Q of a near-spherical celestial body. 501

Substitution of formula (18) into the overall expression (11) for the Andrade com-502 plex compliance will produce the Sundberg and Cooper (2010) rheology: 503

$$\bar{J}(\chi) = J_{\rm U} \left[1 + \frac{\Delta}{1 + \chi^2 \tau^2} - i \frac{\chi \tau \Delta}{1 + \chi^2 \tau^2} + (i \chi \tau_{\rm A})^{-\alpha} \Gamma(1 + \alpha) - i (\chi \tau_{\rm M})^{-1} \right]$$
(19a)

50

497

$$= J_{\rm U} \left[1 + \frac{\Delta}{1 + \chi^2 \tau^2} + \Gamma(1 + \alpha) \zeta^{-\alpha} (\chi \tau_{\rm M})^{-\alpha} \cos\left(\frac{\alpha \pi}{2}\right) \right]$$
(19b)

508

511

516

507

$$- i J_{\rm U} \left[\frac{\chi \tau \Delta}{1 + \chi^2 \tau^2} + \Gamma(1 + \alpha) \zeta^{-\alpha} (\chi \tau_{\rm M})^{-\alpha} \sin\left(\frac{\alpha \pi}{2}\right) + (\chi \tau_{\rm M})^{-1} \right] ,$$

where we introduced the dimensionless Andrade time 510

$$\zeta = \frac{\tau_{\rm A}}{\tau_{\rm M}} \quad . \tag{20}$$

Be mindful that in expression (10) it is only the first term, $J_{\rm e}$, that is changed to func-512 tion (15b). Accordingly, in equation (11), it is only the first term, $J_{\rm e}$, that is substituted 513 with function (18). In the other terms, both the Maxwell and Andrade times are still 514 introduced through the unrelaxed value $J_{\rm e} = J_{\rm U}$: 515

$$\tau_{\rm M} \equiv \eta J_{\rm U} , \quad \tau_{\rm A} \equiv \left(\frac{J_{\rm U}}{\beta}\right)^{1/\alpha} .$$
(21)

Had we combined the elastic relaxation rule (18) with the Maxwell model (7) in-517 stead of Andrade, we would have arrived at the Burgers model — which would be equa-518 tion (19) with the Andrade terms omitted, i.e. with $\tau_{\rm A} \longrightarrow \infty$. Simply speaking, in the 519 absence of transient processes, Andrade becomes Maxwell, while Sundberg-Cooper be-520 comes Burgers. 521

The presently standard term "Sundberg-Cooper rheology" was coined by Renaud 522 and Henning (2018) who studied tidal heating in mantles obeying this rheological law. 523

Along with the dimensionless Andrade time ζ , we shall employ below the relative 524 relaxation time 525 526

$$t_{\rm rel} = \frac{\tau}{\tau_{\rm M}} \tag{22}$$

relating the relaxation timescale for the compliance $J_{\rm e}$ to the Maxwell time. 527

3.5 Further Options

The characteristic relaxation time τ can be replaced with a distribution $D(\tau)$ of 529 times spanning an interval from a lower bound $\tau_{\rm L}$ to an upper bound $\tau_{\rm H}$. So the relax-530 ation of the elastic part of the compliance will be not 531

$$J_{\rm e}(t) = J_{\rm U} \left[1 + \Delta \left(1 - e^{-t/\tau} \right) \right]$$
(23)

but 533

528

532

534

$$J_e(t) = J_U \left[1 + \Delta \int_{\tau_L}^{\tau_H} D(\tau) \left[1 - \exp\left(-\frac{t}{\tau}\right) \right] d\tau \right]$$
(24)

If the relaxation is due to elastically-accommodated GBS, this distribution would be a consequence of variable grain-boundary viscosity, grain sizes and shapes, and non-uniform orientation of grain boundaries with respect to the applied stress (see also Lee & Morris, 2010).

Insertion of expression (24) in the Maxwell model (5) or in the Andrade model (10) produces the *extended Burgers model* or the *extended Sundberg-Cooper model*, correspondingly. For details, see Bagheri et al. (2022) and references therein.

4 Complex Love Numbers and Quality Functions

The perturbing potential wherewith the Earth is acting on the Moon can be decomposed in series over Fourier modes ω_{lmpq} parameterised with four integers lmpq. If the tidal response of the Moon is linear, both the produced deformation and the resulting additional tidal potential of the Moon are expandable over the same Fourier modes, as proved in Efroimsky and Makarov (2014, Appendix C). The proof is based on the fact that a linear integral operator (convolution) in the time domain corresponds to a product of Fourier images in the frequency domain.

While the Fourier modes can be of either sign, the physical forcing frequencies in the body are

$$\chi_{lmpq} = |\omega_{lmpq}| . \tag{25}$$

An extended discussion of this fact can be found in Section 4.3 of Efroimsky and Makarov (2013).

⁵⁵⁵ Wherever this causes no confusion, we omit the subscript to simplify the notation:

$$\omega \equiv \omega_{lmpq}$$
 , $\chi \equiv \chi_{lmpq}$. (26)

557 4.1 The Complex Love Number

552

55

559

565

568

569

571

558 Writing the degree-l complex Love number as

$$\bar{k}_{l}(\omega) = \Re \left[\bar{k}_{l}(\omega) \right] + i \Im \left[\bar{k}_{l}(\omega) \right] = \left| \bar{k}_{l}(\omega) \right| e^{-i\epsilon_{l}(\omega)} , \qquad (27)$$

we conventionally denote the phase as $-\epsilon_l$, with a "minus" sign. This convention imparts ϵ_l with the meaning of phase lag. We also introduce the so-called *dynamical Love number*

$$k_l(\omega) = |\bar{k}_l(\omega)| \quad . \tag{28}$$

A key role in the tidal theory is played by the *quality functions*

$$K_{l}(\omega) \equiv -\Im\left[\bar{k}_{l}(\omega)\right] = \bar{k}_{l}(\omega) \sin \epsilon_{l}(\omega)$$
(29a)

entering the series expansions for tidal forces, torques, dissipation rate (Efroimsky & Makarov,
 2014), and orbital evolution (Boué & Efroimsky, 2019)

Since $\operatorname{Sign} \epsilon_l(\omega) = \operatorname{Sign} \omega$ (Efroimsky & Makarov, 2013), they can be written as

$$K_l(\omega) \equiv -\Im\left[\bar{k}_l(\omega)\right] = \frac{k_l(\omega)}{Q_l(\omega)} \operatorname{Sign}\omega$$
, (29b)

⁵⁷⁰ where the tidal quality factor is introduced via

$$Q_l^{-1}(\omega) = |\sin \epsilon_l(\omega)| .$$
(30)

The dependency $\sin \epsilon_l(\omega)$ being odd, the function $Q_l(\omega)$ is even. Also, even is the function $k_l(\omega)$. Therefore, for any sign of ω and ϵ_l , it is always possible to treat both $Q_l(\omega)$ and $k_l(\omega)$ as functions of the forcing frequency $\chi \equiv |\omega|$:

$$Q_l(\omega) = Q_l(\chi) , \quad k_l(\omega) = k_l(\chi) .$$
(31)

Often attributed to Biot (1954), though known yet to Sir George Darwin (1879), 576 the so-called *correspondence principle*, or the *elastic-viscoelastic analogy*, is a valuable 577 key to numerous problems of viscoelasticity. It enables one to derive solutions to these 578 problems from the known solutions to analogous static problems. In application to bod-579 ily tides, this principle says that the complex Love number of a uniform spherical vis-580 coelastic body, $\bar{k}_l(\chi)$, is linked to the complex compliance $\bar{J}(\chi)$ by the same algebraic 581 expression through which the static Love number k_l of that body is linked to the relaxed 582 compliance $J_{\rm R}$: 583

$$\bar{k}_l(\chi) = \frac{3}{2(l-1)} \frac{1}{1 + \mathcal{B}_l/\bar{J}(\chi)}$$
, (32)

585 where

575

584

586

594

5

$$\mathcal{B}_{l} \equiv \frac{(2l^{2} + 4l + 3)}{l \, g \, \rho \, R} = \frac{3 \, (2l^{2} + 4l + 3)}{4 \, l \, \pi \, G \, \rho^{2} \, R^{2}} \quad , \tag{33}$$

 ρ , R, and g being the density, radius, and surface gravity of the body, and G being Newton's gravitational constant.

As an aside, we would mention that while $-\Im[k_l(\omega)]$ emerges in the tidal torque, the real part of the complex Love number, $\Re[k_l(\omega)] = k_l(\omega) \cos \epsilon_l(\omega)$, shows up in the expansion for the tidal potential. Not considered further in the present study, the general expression for this product and its version for the Maxwell and other rheologies can be found in Efroimsky (2015, Appendix A6).

4.2 $k_l(\chi)/Q_l(\chi)$ and $1/Q_l(\chi)$ for an Arbitrary Rheology

595 Expression (32) entails:

$$K_{l}(\chi) = k_{l}(\chi) \sin \epsilon_{l}(\chi) = - \frac{3}{2(l-1)} \frac{\mathcal{B}_{l} \Im \left[\bar{J}(\chi) \right]}{\left(\Re \left[\bar{J}(\chi) \right] + \mathcal{B}_{l} \right)^{2} + \left(\Im \left[\bar{J}(\chi) \right] \right)^{2}} , \qquad (34)$$

the coefficients \mathcal{B}_l rendered by equation (33). We see that for a homogeneous incompressible sphere, the information needed to calculate the quality function comprises the radius, the density, and the rheological law $\bar{J}(\chi)$.

The inverse tidal quality factor of degree l is given by (Efroimsky, 2015)

$$Q_l(\chi)^{-1} \equiv |\sin \epsilon_l(\chi)| \quad , \tag{35}$$

601 602

600

$$\sin \epsilon_l(\chi) = -\frac{\mathcal{B}_l \Im \left[\bar{J}(\chi)\right]}{\sqrt{\left(\Re \left[\bar{J}(\chi)\right]\right)^2 + \left(\Im \left[\bar{J}(\chi)\right]\right)^2}} \sqrt{\left(\Re \left[\bar{J}(\chi)\right] + \mathcal{B}_l\right)^2 + \left(\Im \left[\bar{J}(\chi)\right]\right)^2} \quad . \tag{36}$$

All new is well-forgotten old. As we were writing this paper, it became known to us that for the Maxwell rheology the frequency-dependence of $\sin \epsilon_2$ was studied yet by Gerstenkorn (1967, Fig. 2) in a work that went virtually unnoticed. Because of different notation and Gerstenkorn's terse style, it is not apparent if his values for the peak's magnitude and location are the same as ours. However, the overall shape of the frequency-dependence of $\sin \epsilon_2$ obtained by Gerstenkorn (1967) seems right.

4.3 Notational Point: Q and Q_2

In publications where both seismic and tidal dissipation are considered, it is necessary to distinguish between the seismic and tidal quality factors. In that situation, the letter Q without a subscript is preserved for the seismic factor.

In the literature on tides, it is common to employ Q as a shorter notation for the quadrupole tidal factor Q_2 . We shall follow the latter convention:

$$Q \equiv Q_2 , \qquad (37)$$

and shall use the two notations intermittently.

4.4 The frequency-dependencies of k_l/Q_l and $1/Q_l$ for the Maxwell and Andrade models

For a homogeneous sphere composed of a Maxwell or Andrade material, the quality function $K_l(\omega)$ has a kink form, as in Figure 1. The function $\sin \epsilon_l(\omega)$ is shaped similarly.



Figure 1. A typical shape of the quality function $K_l(\omega) = k_l(\omega) \sin \epsilon_l(\omega)$, where ω is a shortened notation for the tidal Fourier mode ω_{lmpq} . (From Noyelles et al., 2014).

622

625

627

630

616

Insertion of expression (7) into equation (34) shows that for a spherical Maxwell body the extrema of the kink $K_l(\omega)$ are located at

$$\omega_{\text{peak}_l} = \pm \frac{\tau_{\text{M}}^{-1}}{1 + \mathcal{B}_l \,\mu} \tag{38}$$

⁶²⁶ the corresponding extrema assuming the values

$$K_l^{(\text{peak})} = \pm \frac{3}{4(l-1)} \frac{\mathcal{B}_l \mu}{1 + \mathcal{B}_l \mu} , \qquad (39)$$

where from $|K_l| < \frac{3}{4(l-1)}$.

Inside the interval between peaks, the quality functions are near-linear in ω :

$$|\omega| < |\omega_{\text{peak}_l}| \implies K_l(\omega) \simeq \frac{3}{2(l-1)} \frac{\mathcal{B}_l \mu}{1 + \mathcal{B}_l \mu} \frac{\omega}{|\omega_{\text{peak}_l}|} .$$
(40)

Outside the inter-peak interval, they fall off as about ω^{-1} :

638

643

 $|\omega| > |\omega_{\text{peak}_l}| \implies K_l(\omega) \simeq \frac{3}{2(l-1)} \frac{\mathcal{B}_l \mu}{1 + \mathcal{B}_l \mu} \frac{|\omega_{\text{peak}_l}|}{\omega} .$ (41)

⁶³³ While the peak magnitudes (39) are ignorant of the viscosity η , the spread between ⁶³⁴ the peaks scales as the inverse η , as evident from expression (38). The lower the mean ⁶³⁵ viscosity, the higher the peak frequency $|\omega_{\text{peak}_l}|$.

It can be demonstrated using equation (36) that for a homogeneous Maxwell body the extrema of $\sin \epsilon_l(\omega)$ are located at

$$\omega_{\text{peak of } \sin \epsilon_1} = \pm \frac{\tau_{\text{M}}^{-1}}{\sqrt{1 + \mathcal{B}_l \mu}} . \tag{42}$$

For the Moon, this peak is located within a decade from its counterpart for K_l given by formula (38).

In many practical situations, only the quadrupole (l = 2) terms matter. The corresponding peaks are located at

$$\omega_{\text{peak}_2} = \pm \frac{\tau_{\text{M}}^{-1}}{1 + \mathcal{B}_2 \mu} \approx \pm \frac{1}{\mathcal{B}_2 \eta} = \pm \frac{8 \pi G \rho^2 R^2}{57 \eta} \quad . \tag{43}$$

The approximation in this expression relies on the inequality $\mathcal{B}_l \mu \gg 1$, fulfilment whereof depends on the size of the body. For a Maxwell Moon with $\mu = 6.4 \times 10^{10}$ Pa and $G(\rho R)^2 \approx$ 2.24 × 10⁹ Pa, we have $\mathcal{B}_2 \mu \approx 64.5$, so the approximation works.

⁶⁴⁷ While for the Maxwell and Andrade models each of the functions $K_l(\omega)$ and $\sin \epsilon_l(\omega)$ ⁶⁴⁸ possesses only one peak for a positive argument, the situation changes for bodies of a ⁶⁴⁹ more complex rheology. For example, the existence of an additional peak is ensured by ⁶⁵⁰ the insertion of the Sundberg-Cooper compliance (19) into expressions (34) or (36).

⁶⁵¹ 5 Application to the Moon

652

661

66

5.1 The "Wrong" Slope Interpreted with the Maxwell Model

As we explained in Section 1, fitting of the LLR-obtained quadrupole tidal quality factor $Q = Q_2$ to the power law $Q \sim \chi^p$ resulted in small negative value of the exponential p (Williams & Boggs, 2015). An earlier attempt to explain this phenomenon implied an identification of this slightly negative slope with the incline located to the left of the maximum of the quality function $(k_2/Q_2)(\chi)$, see Figure 1. Within this interpretation, $\chi_{\text{peak}} \equiv |\omega_{\text{peak}}|$ should be residing somewhere between the monthly and annual frequencies explored in Williams and Boggs (2015). As was explained in Efroimsky (2012a) , this sets the mean viscosity of the Moon as low as

$$\eta \approx 3 \times 10^{15} \text{ Pa s} \quad , \tag{44}$$

The extrema of $(1/Q_2)(\chi)$ are close to those of $(k_2/Q_2)(\chi)$, as can be observed from equations (19) and (45) Efroimsky (2015). Therefore, had we used instead of the maximum of k_2/Q_2 given by (43) the maximum of $1/Q_2$ given by (42), the ensuing value would have been only an order higher:

$$\eta \approx 4 \times 10^{16} \text{ Pa s}$$
 (45)

⁶⁶⁷ Such values imply a high concentration of the partial melt in the mantle – quite in ac-⁶⁶⁸ cordance with the seismological models by Nakamura et al. (1974) and Weber et al. (2011).

However, employment of a rheology more realistic than Maxwell may entail not so
 low a viscosity — in which case the existence of a semi-molten layer may be questioned.

5.2 Frequency Dependence of Tidal Dissipation in the Sundberg-Cooper Model

The Debye peak emerging in the imaginary part of \bar{J}_{e} (equation (18)) will, obviously, show itself also in the shape of the imaginary part of the overall \bar{J} , the bottom line of equation (19b). Consequently, substitution of expression (19) in equations (34) and (36) will entail the emergence of a Debye warp on the kinks for k_l/Q_l and $1/Q_l$. Where will the additional peak be located for realistic values of the relaxation timescale τ ? What values for the mean viscosity will it entail?

In the end of Section 3.4, we introduced the relative relaxation time as $t_{\rm rel} \equiv \tau/\tau_{\rm M}$. Figure 2 illustrates specifically the effect of $t_{\rm rel}$ in the Sundberg-Cooper model on the position of the additional Debye peak for a homogeneous lunar interior with an arbitrarily chosen high mean viscosity $\eta_{\rm Moon} = 10^{22}$ Pa s. The emergence of another local maximum in the k_2/Q_2 and $1/Q_2$ functions may naturally explain the decrease in dissipation (or increase in the quality factor Q) with frequency, even within a homogeneous and highly viscous model.



Figure 2. The negative imaginary part of the Love number (left) and the inverse quality factor (right) for different ratios between the timescale τ and the Maxwell time $\tau_{\rm M}$ (indicated by the shades of blue). The yellow and red vertical lines show the Q_2 values given by Williams and Boggs (2015) for the annual and the monthly component, respectively. In this case, we consider a homogeneous lunar interior model governed by the Sundberg-Cooper rheology. The mantle viscosity was set to 10^{22} Pa s and the mantle rigidity to 80 GPa.

686

671

672

5.3 Constructing a Multi-layered Model

Section 4 introduced the complex Love number $k_l(\chi)$ for an arbitrary linear anelas-687 tic or viscoelastic rheology assuming a homogeneous incompressible sphere. While such 688 a model can reasonably approximate the response of the Moon with a homogeneous man-689 tle and a small core (see also Figure 4), its application to a body with a highly dissipa-690 tive basal layer would not be accurate (Bolmont et al., 2020). Planetary interior with 691 a highly dissipative layer can still be approximated by a homogeneous model with an ad-692 ditional absorption peak or band in the underlying rheological law. However, we would 693 need to know the mapping between the parameters of the dissipative layer and the pa-694 rameters of the additional peak (Gevorgyan, 2021). 695

Therefore, in the following sections, we will complement the homogeneous model with three models consisting of two or three layers and we will calculate the corresponding complex Love numbers numerically, using a matrix method based on the normal mode

theory (e.g., Takeuchi & Saito, 1972; Wu & Peltier, 1982; Sabadini & Vermeersen, 2004). 699 For the sake of simplicity, we consider all layers in the numerical model (linearly) vis-700 coelastic and we model the response of liquid layers by the Maxwell model with $J_{\rm e}$ in 701 equation (7) approaching 0. This method has also been tested against another imple-702 mentation of the same model, in which the liquid layers were inputted through differ-703 ent boundary conditions; the results obtained within the two approaches are virtually 704 the same. Using the outputted complex Love numbers for various rheological parame-705 ters, we then proceed by fitting the empirical values. If not stated differently for illus-706 trative purposes, the three alternative models will always comprise a liquid core with a 707 low viscosity ($\eta_c = 1 \text{ Pa s}$), a constant density ($\rho_c = 5000 \text{ kg m}^{-3}$), and an outer ra-708 dius identical to the mean value reported by Weber et al. (2011), $R_c = 330$ km. 709

Although the existence of an inner core is possible and even indicated by the stacked seismograms presented by Weber et al. (2011), its response to tidal loading would be decoupled from the rest of the mantle, and it would contribute to the resulting tidal deformation only negligibly. Therefore, we do not include the inner core in our modelling.

Subsection 5.4 makes use of a two-layered model consisting of the liquid core and a homogeneous mantle, the response of which is described by the Andrade rheology. For the mantle density, we prescribe a constant value of $\rho_{\rm m} = 3300 \,\mathrm{kg} \,\mathrm{m}^{-3}$, and Andrade parameter ζ is set to 1, implying comparable timescales for viscous and anelastic relaxation. Other values of ζ were also tested and their effect on the results is discussed in the aforementioned Subsection. The viscosity $\eta_{\rm m}$, rigidity $\mu_{\rm m}$, and Andrade parameter α of the mantle are treated as free parameters and fitted to the data.

The second model, considered in Subsection 5.5, comprises a liquid core and a Sundberg-Cooper homogeneous mantle. The mantle density is always set to the average value $\rho_{\rm m} =$ 3300 kg m⁻³. Rheological parameters $\eta_{\rm m}$, $\mu_{\rm m}$, τ , and Δ are fitted, while the Andrade empirical parameters α and ζ are held constant during each run of the inversion. We have also tested the effect of varying α in the range [0.1, 0.4] and of magnifying or reducing ζ by one order of magnitude.

The model with a basal dissipative layer, which is discussed in Subsection 5.6, con-727 tains a core and a two-layered mantle. Each layer of the mantle is assumed to be homo-728 geneous. The basal layer is described by the Maxwell model with fitted parameters $\mu_{\rm LVZ}$ 729 and η_{LVZ} ; additionally, we fit its outer radius R_{LVZ} . For the overlying bulk mantle, we 730 consider the Andrade model with free (fitted) parameters $\eta_{\rm m}$, $\mu_{\rm m}$ and with α , ζ kept con-731 stant during each run of the inversion. Both mantle layers have a prescribed density of 732 $\rho_{\rm LVZ} = \rho_{\rm m} = 3300 \, \rm kg \, m^{-3}$. The reason for using the simple Maxwell model instead 733 of the Andrade model in the basal layer is the following: in order to fit the measured tidal 734 quality factor Q at the monthly and the annual frequency, the peak dissipation from the 735 basal layer should be located either between these frequencies, or above the monthly fre-736 quency. At the same time, in the vicinity of the peak dissipation, the Andrade and Maxwell 737 rheologies are almost indistinguishable from each other. (Comparing the last two terms 738 on the final line of equation (19), we observe that the viscous term exceeds the Andrade term when $\tau_{\rm M} \chi \ll (\tau_{\rm A}/\tau_{\rm M})^{\alpha/(1-\alpha)}$. In realistic situations, $\tau_{\rm M} \chi_{\rm peak}$ satisfies this con-739 740 dition safely. So, near the peak the Andrade term is virtually irrelevant, and the regime 741 is almost Maxwell.) Hence, we chose the simpler of the two rheological models. This de-742 cision will also facilitate the comparison of our results for the basal layer's characteris-743 tics with the predictions by Harada et al. (2014, 2016), and Matsumoto et al. (2015), who 744 likewise modeled the basal layer with the Maxwell model. In contrast to our study, they 745 applied the same model to the mantle as well. 746

In this work, we are not predicting the mineralogy of the mantle — and the composition of the basal layer, if present, is only briefly discussed in Subsection 6.2. Our use
of a homogeneous mantle layer (or two homogeneous mantle layers) reflects our lack of
information on the exact chemical and mineralogical composition, the grain size, the thermal structure, and the presence of water. Instead, we characterise the mantle with a sin-

gle, "effective", rigidity and viscosity, which can be later mapped to a detailed interior 752 structure (see also Dumoulin et al., 2017; Bolmont et al., 2020, who discussed the effect 753 of approximating a radially stratified mantle with a homogeneous one for Venus and ter-754 restrial exoplanets). Furthermore, we neglect any lateral heterogeneities in the lunar in-755 terior. We also assume that the lunar mantle is incompressible and can be reasonably 756 described by a linear viscoelastic model — which is valid at low stresses. Given the mag-757 nitude of tidal stresses in the Moon, this assumption might have to be lifted in future 758 works, though (Karato, 2013). 759

760 Since the radial structure of our models is deliberately simplified, we do not attempt to fit either the mean density or the moment of inertia given for the Moon. (The mean 761 density of our lunar toy-models is less than 1% lower than the actual value.) The inver-762 sions presented below are only performed for the tidal parameters, namely k_2 and tidal 763 Q at the monthly frequency, k_2/Q at the annual frequency, and k_3 , h_2 at the monthly 764 frequency. A list of the model parameters in the reference cases discussed in the follow-765 ing sections is presented in Table 1. The empirical values considered are then given in 766 Table 2. 767

Parameter	Type	Value	Unit
Common par	ameters		
$\overline{\text{Core size } R_{c}}$	const.	330	km
Core viscosity η_c	const.	1	Pa s
Core density $\rho_{\rm c}$	const.	5,000	${ m kg}~{ m m}^{-3}$
Mantle viscosity $\eta_{\rm m}$	fitted	$10^{15} - 10^{30}$	Pa s
Mantle rigidity $\mu_{\rm m}$	fitted	$10^9 - 10^{12}$	Pa
Mantle density $\rho_{\rm m}$	const.	3,300	${ m kg}~{ m m}^{-3}$
And rade parameter ζ	const.	1	_
Two-layered model I	(Andrad	le mantle)	
And rade parameter α	fitted	0 - 0.5	
Two-layered model II (Su	undberg-0	Cooper mantle	2)
Andrade parameter α	const.	0.2	
Relaxation strength Δ	fitted	$10^{-5} - 10^{0}$	
Relative relaxation time $t_{\rm rel}$	fitted	$10^{-7} - 10^{0}$	
Three-layered model	(Andrao	le mantle)	
Andrade parameter α	const.	0.2	
Thickness of the basal layer $D_{\rm LVZ}$	fitted	0 - 370	$\rm km$
Viscosity of the basal layer $\eta_{\rm LVZ}$	fitted	$10^0 - 10^{30}$	Pa s
Rigidity of the basal layer $\mu_{\rm LVZ}$	fitted	$0 - \mu_{\rm m}$	Pa

Table 1. Parameters of the three models considered in this work.

⁷⁶⁸ 5.4 Applicability of the Andrade Model

Before discussing the two interior models able to fit the anomalous frequency dependence of lunar tidal dissipation, we first attempt to use the full set of tidal parameters given in Table 2 to constrain a simpler model, which only contains a liquid core and a viscoelastic mantle governed by the Andrade rheology (equation (11)). Such a model, accounting neither for a basal dissipative layer nor for elastically-accommodated GBS, might still be able to fit the data. Thanks to the large uncertainty on the lunar qual-

Parameter	Value	Reference
$ \frac{k_2, \text{ monthly}}{Q, \text{ monthly}^{a}} \\ \frac{k_2/Q, \text{ annual}^{a}}{k_3, \text{ monthly}^{b}} \\ \frac{k_2}{h_2, \text{ monthly}} $	$\begin{array}{c} 0.02422\pm 0.00022\\ 38\pm 4\\ (6.2\pm 1.4)\times 10^{-4}\\ 0.0081\pm 0.0018\\ 0.0387\pm 0.0025 \end{array}$	Williams et al. (2014) Williams and Boggs (2015) Williams and Boggs (2015) Konopliv et al. (2013); Lemoine et al. (2013) Thor et al. (2021)

^a The standard deviations from this table are only used in Subsection 5.4. In the rest of the paper, we arbitrarily set the uncertainties to 1% of the mean value. ^b Listed is the unweighted mean of the values given in references.

 Table 2.
 Observational constraints used in this work.

⁷⁷⁵ ity factor (more than 10% at the monthly frequency and 20% at the annual frequency, ⁷⁷⁶ Williams & Boggs, 2015), we may not need to introduce any additional complexities to ⁷⁷⁷ interpret the tidal response of the Moon. The error bars of the tidal quality factors are ⁷⁷⁸ so wide that they allow, at least in principle, for a situation where $Q_{2, \text{ annual}}$ is smaller ⁷⁷⁹ than $Q_{2, \text{ monthly}}$.

To find the parameters of this preliminary model, we performed a Bayesian inver-780 sion using the MCMC approach and assuming Gaussian distribution of observational un-781 certainties (e.g., Mosegaard & Tarantola, 1995). In particular, we employed the emcee 782 library for Python (Foreman-Mackey et al., 2013), which is based on the sampling meth-783 ods proposed by Goodman and Weare (2010). The algorithm was instructed to look for 784 the mantle viscosity $\eta_{\rm m}$, the mantle rigidity $\mu_{\rm m}$, and the Andrade parameter α fitting 785 the empirical values of $k_{2,\text{monthly}}$, $k_{3,\text{monthly}}$, $h_{2,\text{monthly}}$, $Q_{2,\text{monthly}}$, and $(k_2/Q_2)_{\text{annual}}$, 786 while the other Andrade parameter was set to $\zeta = 1$. We generated ~ 30,000 random 787 samples until the model converged. Specifically, the convergence was tested against the 788 autocorrelation time of each variable in the ensemble, the total length of all chains be-789 ing required to exceed 100 times the longest autocorrelation time. Moreover, in order 790 to filter out the influence of initial conditions, we neglected the first $\sim 3,000$ samples 791 (our burn-in period was, therefore, 10 times the autocorrelation time). 792

The posterior probabilities of the fitted parameters are depicted in Figure 3, using the *Python* library *corner* (Foreman-Mackey, 2016). In line with a similar model by Nimmo et al. (2012), we find a relatively high lunar mantle viscosity of $\log \eta$ [Pa s] = 22.99^{+0.89}_{-1.35} and rigidity of $\log \mu$ [Pa] = 10.92±0.06, the Andrade parameter α being as low as $0.06^{+0.04}_{-0.02}$.

Treating the Andrade parameter ζ as a free parameter in the Bayesian inversion 797 has a negligible effect on the predicted values of α and $\mu_{\rm m}$. However, it essentially de-798 termines the fitted mantle viscosity. If the transient deformation prevails over the vis-799 cous creep ($\zeta \ll 1$), the response of the lunar mantle to tidal loading is almost elastic 800 (with viscosity up to $\eta \approx 10^{27}$ Pa s). On the other hand, if the dissipation is preferen-801 tially due to viscous creep ($\zeta \gg 1$), the mantle viscosity allowed by the observational 802 data has to be much lower, $\eta \approx 10^{21}$ Pa s. This latter case is equivalent to the assump-803 tion that the mantle is governed by the Maxwell rheology, followed by Harada et al. (2014, 804 2016); Matsumoto et al. (2015); Tan and Harada (2021), and Kronrod et al. (2022). 805

If we compare the resulting Andrade parameter $\alpha = 0.06^{+0.04}_{-0.02}$ with the typical values reported in the literature (0.1 < α < 0.4; see, e.g., the overview by Castillo-Rogez et al., 2011; Efroimsky, 2012a, 2012b), we may notice that it is unusually small. This discrepancy between our prediction and the laboratory data already indicates that although it is, in principle, possible to fit the lunar tidal response with a simple model assuming Andrade rheology in the mantle, the required parameters of this model might not be realistic. A similar point has been made by Khan et al. (2014) and used as an ar-

gument in favour of their interior model containing basal partial melt. Following the same

line of argumentation, we will now focus our study on the Sundberg-Cooper model.



Figure 3. Posterior probabilities of the effective mantle rigidity $\mu_{\rm m}$, the mantle viscosity $\eta_{\rm m}$, and the Andrade parameter α satisfying the full set of observational constraints $(k_2, k_3, h_2, \text{ and} Q$ at the monthly period; k_2/Q at the annual period). A model with a liquid core and a viscoelastic mantle governed by the Andrade rheology, assuming $\zeta = 1$.

815

5.5 Lunar Mantle Governed by the Sundberg-Cooper Model

In the present Subsection, as well as in Subsection 5.6, we will specifically search 816 for lunar interior models that exhibit a second dissipation peak in the spectra of k_2/Q_2 817 and Q_2^{-1} . Since the current error bars of the empirical Qs allow for both a decrease and 818 increase of dissipation with frequency, and since our study focuses on the latter case, we 819 consider a hypothetical situation in which the uncertainty in Q_2 is comparable with the 820 present-day uncertainty in k_2 . The standard deviations of Q_2 at the monthly frequency 821 and k_2/Q_2 at the annual frequency are thus arbitrarily set to 1% of the mean value. As 822 in the previous inversion with Andrade mantle, we again employ the MCMC approach 823 and seek the parameters of the Sundberg-Cooper model $(\eta_m, \mu_m, \Delta, \text{ and } t_{rel})$ fitting the 824 empirical tidal parameters. Values of α and ζ are kept constant. For illustration pur-825 poses, we consider both 1) a two-layered interior structure consisting of a liquid core and 826 a viscoelastic (Sudberg-Cooper) mantle and 2) a homogeneous lunar interior. As we shall 827 see, the effect of the small lunar core $(R_c = 330 \text{ km})$ on the results is negligible. 828

In contrast with the previous inversion, and mainly due to the greater dimension of the explored parameter space, the model only succeeded to converge after generating ~ 700,000 random samples. The posterior distributions of the tidal quality factors typically presented two peaks: a higher one with $Q_{2,\text{monthly}} > Q_{2,\text{annual}}$ and a lower one with $Q_{2,\text{monthly}} < Q_{2,\text{annual}}$. Here, we only discuss the model parameters corresponding to the latter case.

Figure 4 illustrates the results of the inversion with Andrade parameters specifi-835 cally set to $\alpha = 0.2$ and $\zeta = 1$. Similarly as before, to filter-out the influence of ini-836 tial conditions, we neglected the first 70,000 samples. Then, 16% of the remaining, anal-837 ysed samples fulfilled the condition of quality factor decreasing with frequency. The mean 838 value of the predicted mantle viscosity lies close to 3.5×10^{22} Pa s and the predicted un-839 relaxed rigidity is around 60 - 120 GPa. In particular, for the nominal case with $\alpha =$ 840 0.2 and $\zeta = 1$ and for the arbitrarily chosen small standard deviation of empirical Q 841 and k_2/Q , the decadic logarithms of the predicted mantle viscosity and rigidity are log $\eta_{\rm m}[{\rm Pa~s}] =$ 842 $22.55^{+0.15}_{-0.54}$ and $\log \mu_{\rm m}$ [Pa] = 10.84^{+0.14}_{-0.02}. Increasing α by 0.1 or ζ by the factor of 10 re-843 sults in decreasing the mantle viscosity approximately by an order of magnitude (and 844 the same trend pertains to the other direction, when decreasing α or ζ). On the other 845 hand, the mantle rigidity, being dictated by the magnitude of k_2 , seems relatively robust 846 and its inverted value does not depend on α . 847

The parameters of the Debye peak are, in this story, the key to fitting the unex-848 pected slope of the frequency-dependent tidal dissipation. Independently of the consid-849 ered Andrade parameters, the relaxation timescale τ lies between 10⁴ and 10⁶ s (log τ [s] = 850 $4.89_{-0.72}^{+0.62}$), while the relaxation strength falls into the interval between 0.03 and 1 (log $\Delta = -1.17_{-0.35}^{+0.84}$). The exact values depend on the predicted viscosity and rigidity, which de-851 852 fine the position of the first peak, corresponding to the attenuation in the overlying man-853 tle. Such short relaxation timescales would indicate that the elastically accommodated 854 GBS is much faster than diffusion creep. For comparison, Sundberg and Cooper (2010) 855 mention a GBS relaxation timescale of 0.1 s as a reasonable value in their experiments, 856 using a material with $\tau_{\rm M} \sim 10-100 \, \text{s.}$ Our $\tau_{\rm M}$ in this specific case is in the order of $10^{10}-$ 857 10^{13} s; hence, the ratio of the two time scales for $\alpha = 0.2$ and $\zeta = 1$ reaches $t_{\rm rel} = 10^{-7} -$ 858 10^{-6} . A more detailed discussion of this result will be provided in Subsection 6.1. 859

860 861

5.6 Comparison of a Sundberg-Cooper Moon with an Andrade Moon Having a Weak Basal Layer

As was recently shown by Gevorgyan (2021), the tidal response of a homogeneous 862 Sundberg-Cooper planet mimics the response of a body consisting of two Andrade lay-863 ers with different relaxation times. This kind of aliasing may, in principle, be demonstrated 864 by the Moon. Figure 5 depicts the imaginary part of the tidal Love number (equal to 865 k_2/Q_2) and the inverse quality factor $1/Q_2$ as functions of frequency, for a homogeneous 866 Sundberg-Cooper moon and for a differentiated lunar interior with a rheologically weak 867 layer at the base of the mantle. In the second case, the basal layer is described by the 868 Maxwell model and the overlying mantle by the Andrade model. Both cases follow the same frequency dependence, implying that the existence of a weak basal layer cannot 870 be confirmed unequivocally by the tidal data. In a layered model containing a core, a 871 Sundberg-Cooper mantle, and a Maxwell basal semi-molten layer, the tidal response would 872 be characterised by three peaks (Figure 6). 873

For comparison with other models presented in the literature, we also seeked for the parameters of a three-layered lunar model comprising a liquid core, an Andrade mantle, and a Maxwell basal low-viscosity layer that would fit the empirical constraints. As in the previous Subsection, in order to reduce the number of unknowns, the parameters α and ζ of the Andrade model were kept constant. We also prescribed the same constant core radius of 330 km. The remaining quantities were treated as free parameters: we thus varied the rigidity and viscosity of the mantle and of the basal layer, and the outer ra-



Figure 4. Best-fitting models and the corresponding model parameters for a melt-free Moon with a liquid core and a Sundberg-Cooper mantle. Upper row: the real (left) and negative imaginary (right) parts of the complex Love number \bar{k}_2 , as functions of frequency. The red and yellow lines indicate the values provided by Williams and Boggs (2015). Lower row: model samples plotted in the parameter space, with the mantle rigidity μ_m depicted against viscosity η_m (left), the relaxation strength Δ depicted against the characteristic time τ of the elastically-accommodated GBS (centre), and the Maxwell time τ_M versus the characteristic time τ (right). The Andrade parameters are kept constant at $\alpha = 0.2$ and $\zeta = 1$. Gray dots in the lower left panel show the results obtained with a homogeneous model consisting only of a Sundberg-Cooper mantle, while black dots represent the default two-layered model.



Figure 5. The negative imaginary part of the Love number (left) and inverse quality factor (right) for three model cases: a homogeneous Andrade model (dashed red line), a homogeneous Sundberg-Cooper model (blue line), and a three-layered model (solid red line) comprising a core, an Andrade mantle and a Maxwell semi-molten layer at the base of the mantle.



Figure 6. The negative imaginary part of the Love number (left) and inverse quality factor (right) of a three-layered lunar model comprising a core, a Sundberg-Cooper mantle, and a Maxwell semi-molten basal layer. Different shades of blue correspond to different ratios between the timescale τ and the Maxwell time $\tau_{\rm M}$. For illustrative purposes, the semi-molten basal layer is made unrealistically thick (500 km).

dius of the basal layer. Due to the higher dimensionality of the parameter space, the inverse problem took longer to converge; therefore, we generated 10,000,000 random samples satisfying all constraints from Table 2. Since the longest autocorrelation time in this case was 500,000 steps, we discarded the first 5,000,000 samples and then applied the condition $Q_{2,\text{monthly}} < Q_{2,\text{annual}}$, being left with 11% of the generated samples.

As illustrated in Figure 7, and in line with the discussion above, the frequency de-886 pendencies of $\Re[k_2]$ and $-\Im[k_2]$ in the model with a low-viscosity basal layer closely re-887 semble those of the previous one, in which we considered the Sundberg-Cooper model. 888 Similarly to the earlier predictions of the basal layer's viscosity and thickness (e.g., Harada 889 et al., 2014, 2016; Matsumoto et al., 2015), we find that the observed frequency depen-890 dence of lunar Q_2^{-1} can be explained by the viscosity $\eta_{\rm LVZ}$ in the range from ~ 10¹⁵ to 891 $\sim 3 \times 10^{16}$ Pa s and the thickness $D_{\rm LVZ}$ in the range from 70 km to the maximum value 892 considered in our model (370 km). The parameter dependencies of all model samples are 893 plotted on Figure 8. For the nominal case with $\alpha = 0.2$ and $\zeta = 1$, and considering 894 the condition on Q mentioned in the above paragraph, we obtain the following rigidity 895 and viscosity of the overlying mantle and of the LVZ: $\log \eta_{\rm m}[{\rm Pa~s}] = 22.79^{+0.19}_{-0.06}, \ \mu_{\rm m}[{\rm Pa}] = 10.89 \pm 0.03, \ \eta_{\rm LVZ}[{\rm Pa~s}] = 15.20^{+0.53}_{-0.21}, \ \mu_{\rm LVZ}[{\rm Pa}] = 10.23^{+0.37}_{-0.34}.$ The corresponding outer radius of the LVZ is $R_{\rm LVZ}[{\rm km}] = 599.39^{+65.83}_{-84.46}.$ 897 898

Similarly to the "melt-free" case with the Sundberg-Cooper model, increasing α to 0.3 results in an order-of-magnitude decrease in the fitted mantle viscosity. Decreasing α to 0.1 leads to a mantle viscosity two orders of magnitude greater. On the other hand, the predicted properties of the semi-molten layer remain almost the same.

903 6 Discussion

In the previous section, we have compared the frequency dependence of lunar Qwithin the widely accepted lunar interior model containing a highly dissipative layer at the base of the mantle (e.g., Nakamura et al., 1973; Williams et al., 2001; Harada et al., 2014) and within an alternative model taking into account the time relaxation of the elastic compliance J_{e} . On the following lines, we discuss the implications of each of the considered models for the lunar interior properties. Keep in mind that the inversions per-



Figure 7. Overview of best-fitting models for the case with a basal low-viscosity zone. The red and yellow lines indicate the values provided by Williams and Boggs (2015). As in the previous inversion, the Andrade parameters are kept constant at $\alpha = 0.2$ and $\zeta = 1$, and the core size is fixed to 330 km.



Figure 8. Model samples corresponding to Figure 7, plotted in the parameter space. The intensity indicates the sample count. Upper row: the rigidity vs. viscosity of the LVZ (left), the rigidity vs. viscosity of the mantle (centre), and the outer radius vs. viscosity of the LVZ (right). Lower row: the rigidity of the LVZ vs. rigidity of the mantle (left), the viscosity of the LVZ vs. viscosity of the mantle (centre), and the outer radius vs. rigidity of the LVZ (right).

formed in our study explicitly assumed that the value of Q at the monthly frequency and k_2/Q at the annual frequency are known with a high precision. This is not the case in reality. However, as we have seen in Subsection 5.4, a lunar mantle governed by the Andrade model without a basal dissipative layer can fit the data with the actual uncertainties only for unrealistically low values of parameter α .

6.1 Melt-free Lunar Interior

In the model cases considering a two-layered, "melt-free" lunar interior, where the 916 negative slope of the frequency dependence of k_2/Q is explained by a secondary dissi-917 pation peak induced by elastically accommodated GBS, we found that the logarithm of 918 the relaxation timescale, $\log \tau$, falls into the range of [4,6], corresponding to τ between 919 3 and 300 hours. In the reference case depicted in Figure 4, this would imply a ratio of 920 the characteristic timescales for the elastic and diffusional accommodation $t_{\rm rel} = \tau / \tau_M$ 921 to be of order from 10^{-7} to 10^{-6} . Are such ratios of the characteristic times observed 922 in any natural materials? 923

According to Jackson et al. (2014), grain boundary sliding comprises three processes. 924 The relative contribution of each of them to the energy dissipation in a sample depends 925 on the temperature and loading frequency. The processes are: (i) elastically accommo-926 dated GBS with a characteristic time τ , at high frequencies/low temperatures; (ii) dif-927 fusionally assisted GBS described by the power-law frequency-dependence of the seis-928 mic quality factor, $Q \propto \chi^p$; and (iii) diffusionally accommodated GBS at timescales 929 greater than the Maxwell time $\tau_{\rm M}$, where the seismic Q is a linear function of frequency, 930 $Q \propto \chi$. The value of $t_{\rm rel}$ thus determines the range of frequencies over which the dif-931 fusionally assisted sliding on spacial scales smaller than grain size occurs. Experimen-932 tal data for fine-grained polycrystals indicate that $t_{\rm rel} \ll 1$ (Morris & Jackson, 2009). 933

Jackson et al. (2014) presented results of laboratory experiments on fine-grained 934 olivine subjected to torsional oscillations at high pressures (P = 200 MPa) and rela-935 tively low temperatures $(T < 900 \,^{\circ}\text{C})$, i.e., around the threshold between elastic response 936 and elastically accommodated GBS. They found a GBS relaxation timescale of $\log \tau_{\rm B} =$ 937 $1.15\pm0.07\,\mathrm{s}$, where the subscript "R" now stands for "reference". Considering the ref-938 erence temperature $T_{\rm R} = 900$ °C, reference pressure $P_{\rm R} = 200$ MPa, reference grain 939 size $d_{\rm R} = 10 \,\mu{\rm m}$, activation volume $V^* = 10 \,{\rm cm}^3 \,{\rm mol}^{-1}$, and activation energy $E^* =$ 940 $259 \,\mathrm{kJ} \,\mathrm{mol}^{-1}$, as given by Jackson et al. (2014), we can extrapolate τ to the conditions 941 of the lunar mantle with the Arrhenius law (Jackson et al., 2010): 942

943

915

$$\tau = \tau_{\rm R} \left(\frac{d}{d_{\rm R}}\right)^m \exp\left\{\frac{E^*}{R} \left(\frac{1}{T} - \frac{1}{T_{\rm R}}\right)\right\} \exp\left\{\frac{V^*}{R} \left(\frac{P}{T} - \frac{P_{\rm R}}{T_{\rm R}}\right)\right\} \,. \tag{46}$$

In addition to the parameters introduced earlier, d is the grain size and m char-944 acterises the grain-size dependence of the process in question. We adopt the value m =945 1.31, found by Jackson et al. (2010) for anelastic processes. Figure 9 illustrates the ex-946 trapolation of $\tau_{\rm R}$ of Jackson et al. (2014) to lunar interior conditions, considering our 947 melt-free model and two depth-independent grain sizes. Over the colour-coded maps, we 948 also plot the steady-state heat conduction profiles of Nimmo et al. (2012). We note that 949 the conduction profiles were only chosen for illustration purposes: the discussion of the 950 thermal regime (conductive vs. convective) in the lunar mantle is beyond the scope of 951 this paper. 952

The laboratory measurements of Jackson et al. (2014) were performed on a single sample of fine-grained polycrystalline olivine under constant pressure $P_{\rm R}$ and the Arrhenian extrapolation of τ was only tested for temperature dependence. Nevertheless, if we accept the assumption that these results are applicable to the Moon, Figure 9 and the fitted relaxation time from Figure 4 (log $\tau \in [4, 6]$) can help us to identify the minimum depth in which elastically accomodated GBS contributes to the tidal dissipation. For the



Figure 9. Relaxation time τ (colour-coded) of elastically accommodated GBS, as given by Jackson et al. (2014) and extrapolated to lunar interior conditions using the Arrhenian equation (46). White lines demarcate the relaxation times resulting from our inversion. Blue lines indicate analytically-calculated conduction profiles proposed by Nimmo et al. (2012) for three different mantle heat productions (8, 9.5, and 11 nW m⁻³), crustal heat production of 160 nW m⁻³ crustal thickness of 45 km, and no heat exchange between core and mantle. Other parameters, such as the core size, core density, and mantle density, are adjusted to our melt-free model. Grain sizes are given in the upper right corner of each plot.

smaller grain size (d = 0.1 mm) and the reference profile of Nimmo et al. (2012) (solid 959 line, mantle heat production of $9.5 \,\mathrm{nW} \,\mathrm{m}^{-3}$), we predict the minimum depth of $400-500 \,\mathrm{km}$. 960 For the larger grain size (d = 1 cm), the minimum depth is 600–800 km. A conductive 961 profile corresponding to lower heat production than illustrated here would push the min-962 imum depth to even greater values. The occurrence of elastically accommodated GBS 963 in shallower depths would give rise to a relaxation peak (or to an onset of a relaxation band) at lower loading frequencies, which would not fit the observed annual and monthly 965 tidal Q. Although the MCMC inversion from the previous section was performed for a 966 model with a homogeneous mantle, i.e., assuming the occurrence of elastically-accommodated 967 GBS at all depths from the surface down to the core, we also checked that a model de-968 scribed by the Andrade rheology above the derived depths and by the Sundberg-Cooper 969 model below the derived depths would fit the considered observables under the condi-970 tion that $\log \tau \gtrsim$. For shorter τ , the estimated minimum depth of applicability of the 971 Sundberg-Cooper model would not match the Love numbers at monthly frequency. 972

Besides the timescale τ , we have derived the relaxation strength of the hypothetical secondary peak: $\log \Delta \in [-1.5, 0]$, or $\Delta \in [0.03, 1]$. Parameter Δ controls the height of the secondary dissipation peak in the Sundberg-Cooper model. Figure 10 shows the dependence of this Q^{-1} on the relaxation strength for all our models from Figure 4. Are these values consistent with theoretical prediction and laboratory data?

Sundberg and Cooper (2010) reported relaxation strengths of polycrystalline olivine 978 between 0.23 and 1.91, as found in different sources and under different assumptions on 979 the grain shapes (Kê, 1947; Raj & Ashby, 1971; Ghahremani, 1980). Their own mechan-980 ical tests on peridotite (olivine-orthopyroxene) at temperatures between 1200 and 1300 $^{\circ}$ C 981 were best fitted with $\Delta = 0.43$ and the corresponding dissipation associated with elastically-982 accommodated GBS in their sample was $Q^{-1} = 0.25 - 0.3$. On the other hand, Jackson 983 et al. (2014), who performed torsion oscillation experiments on olivine, found a relatively 984 low dissipation peak with $Q^{-1} \leq 0.02$. Low secondary dissipation peaks with $Q^{-1} \sim$ 985 10^{-2} were also predicted theoretically by Lee and Morris (2010) for a grain boundary 986 slope of 30°, while smaller slopes seem to allow Q^{-1} exceeding 1, especially when the in-987 dividual grains are of comparable sizes and the grain boundary viscosity does not vary 988

too much. Accordingly, Lee et al. (2011) note that Q^{-1} in the secondary peak depends strongly on the slope of the grain boundaries.

Following this brief discussion of dissipation arising due to elastically accommodated GBS, we can conclude that the relaxation strength Δ (or Q^{-1} in the secondary dissipation peak) is not well constrained and the values found in literature permit any of the Δ s predicted in our Subsection 5.5.



Figure 10. Seismic Q^{-1} of the mantle at the frequency of the secondary peak, plotted as a function of the relaxation strength Δ for models from Figure 4.

6.2 Highly Dissipative Basal Layer

995



Figure 11. Shear modulus prediction compared to seismic measurements. Shear modulus μ_{LVZ} for $R_{\text{LVZ}} = 400,500$ and 700 km (gray, yellow and orange areas). Shear modulus derived from seismic velocities and densities: green (Weber et al., 2011), red (Khan et al., 2014) and blue (Matsumoto et al., 2015), dashed lines: errors.

A highly dissipative layer located at any depth could also produce the desired secondary peak needed to explain the anomalous Q dependence. (Note, however, that a presence of a highly dissipative layer at a shallow depth may lead to changes in the body's response to tides and might be incompatible with the measured values of the Love numbers.) Petrological considerations combined with an indication of a basal low-velocity zone point to the presence of this anomalous layer in the deep interior. Therefore, as an
 alternative to the "melt-free" model, we tested the popular hypothesis of a putative highly
 dissipative layer at the base of the lunar mantle.

The derived rheological properties of the mantle and of the basal layer as well as 1004 the layer's thickness are poorly constrained and can be strongly biased. Firstly, the outer 1005 radius $R_{\rm LVZ}$ of the basal layer is correlated with the value of the mantle rigidity $\mu_{\rm m}$; the 1006 thicker the basal layer, the larger mantle rigidity can be expected to satisfy the model 1007 constraints. The mantle viscosity $\eta_{\rm m}$ depends on the empirical Andrade parameters, and 1008 an increase of α by 0.1 leads to a reduction of the fitted mantle viscosity approximately 1009 by one order of magnitude. On the other hand, the viscosity of the basal layer remains 1010 independent of the empirical Andrade parameters. The predicted contrast in viscosity 1011 between the two layers thus decreases with increasing α and/or ζ . Secondly, the range 1012 of acceptable basal rigidities μ_{INZ} widens with the basal layer thickness (Figure 11). We 1013 do not find an acceptable solution for $R_{\rm LVZ} \lesssim 400 \,\rm km$ due to our a priori requirement 1014 on the relationship between the mantle and basal layer's rigidities ($\mu_{LVZ} \leq \mu_m$). The 1015 range of acceptable $\mu_{\rm LVZ}$ values increases with the basal layer radius up to one and a half 1016 order of magnitude for the maximum $R_{\rm LVZ} = 700 \,\rm km$ considered here. Interestingly, 1017 the predicted rigidities of a basal layer with thickness $\sim 170 \,\mathrm{km} \,(R_{\mathrm{LVZ}} \approx 500 \,\mathrm{km})$ cor-1018 responds well with the seismic observations. Lastly, the basal viscosity is correlated with 1019 the basal layer thickness: the viscosity $\eta_{\rm LVZ}$ decreases from $3 \cdot 10^{16} \, {\rm Pas}$ for a thin weak 1020 layer $(R_{\rm LVZ} = 400 \,\rm km)$ to $< 10^{15} \,\rm Pas$ for the greatest considered thickness $(R_{\rm LVZ} =$ 1021 700 km). The basal layer viscosity is, therefore, always considerably lower than the man-1022 tle viscosity. However, this is not surprising, as the low viscosity of this layer is essen-1023 tial to predict the anomalous frequency dependence of the tidal quality factor, when the 1024 rest of the high-viscosity mantle is set to obey the Andrade law. 1025



Figure 12. Basal viscosity prediction compared to rheological properties. Predicted ranges of viscosities η_{LVZ} for $R_{\text{LVZ}} = 400, 500$ and 700 km are indicated by gray, yellow, and orange areas, respectively. Over the predicted ranges is plotted the temperature dependence of viscosity of ilmenite (blue, Dygert et al., 2016), dry olivine (red, Hirth & Kohlstedt, 1996), and ilmeniteolivine aggregate (2 - 16%), the latter corresponding either to isostress (blue area, harmonic mean, suggested for high strain) or Tullis (red area, geometric mean, suggested for low strain) models. Errors of experimentally determined viscosities not included; ilmenite error factor is ~ 5. Vertical lines delimit solidus temperatures for peridotite (Katz et al., 2003) and ilmenite-bearing material (Wyatt, 1977) at radii 330 km and 700 km. Left panel: temperature dependence for $\sigma_D = 1$ MPa, dry olivine. Right panel: temperature dependence for $\sigma_D = 1$ MPa, wet olivine.

Rigidity and viscosity magnitudes, and their contrast between the mantle and the basal layer values, can be indicative of the variations in the composition, in the presence of melt, and in temperature. A stable partially molten zone in the lunar interior would pose strong constraints on the composition (Khan et al., 2014). Given the absence of geologically recent volcanic activity, any melt residing in the deep lunar interior would have
to be neutrally or negatively buoyant. Using an experimental approach on the synthetic
equivalent of Moon samples, van Kan Parker et al. (2012) concluded that the condition
on the buoyancy below 1000 km is satisfied if high content of titanium dioxide is present
in the melt. We can expect the presence of a partially molten layer at any depth below
this neutral buoyancy level.

Moreover, evolutionary models suggest that high-density ilmenite bearing cumu-1036 lates enriched with TiO_2 and FeO are created towards the end of the shallow lunar magma 1037 ocean crystallisation, resulting in near-surface gravitational anomalies. This instability, 1038 combined with the low viscosity of those cumulates, might have eventually facilitated 1039 the mantle overturn, creating an ilmenite-rich layer at the base of the mantle (e.g., Zhang 1040 et al., 2013; Zhao et al., 2019; Li et al., 2019). Recently, Kraettli et al. (2022) suggested 1041 an alternative compositional model: $a \sim 70 \,\mathrm{km}$ thick layer of garnetite could have been 1042 created at the base of the mantle if two independently evolving melt reservoirs were present. 1043 The resulting high-density garnet, olivine, and FeTi-oxide assemblage is gravitationally 1044 stable and can contain a neutrally or negatively buoyant Fe-rich melt. The scenario of Kraettli et al. (2022) can also be accompanied by the mantle overturn, as suggested for 1046 the ilmenite-rich layer created at shallow depths. 1047

Rheologically weak ilmenite combined with appropriate lower-mantle temperature 1048 can help to explain the low basal viscosity (Figure 12). If the lower mantle were only made 1049 of dry olivine, the predicted viscosity would require temperature $\gtrsim 1800 \,\mathrm{K}$, whereas for 1050 wet olivine, the temperature range between ~ 1500 and ~ 1800 K would be sufficient. 1051 Creep experiments (Dygert et al., 2016) conclude that the viscosity of ilmenite is more 1052 than three orders of magnitude lower than dry olivine. Consequently, a lower-mantle tem-1053 perature $(1400 - 1700 \,\mathrm{K})$ might be acceptable to explain the predicted viscosities for 1054 pure ilmenite. The properties of ilmenite-olivine aggregates introduce yet another com-1055 plexity. The viscosity of aggregates is suggested to depend on the value of the strain: it 1056 follows the Tullis model for low strain, whereas it tends to follow the lower bound on Fig-1057 ure 12 (isostress model) for large strain (see, e.g., Dygert et al., 2016, for a deeper dis-1058 cussion). The acceptable temperature range for olivine-ilmenite aggregate is close to the 1059 values for the pure olivine in the case of the Tullis model. The prediction for the isostress 1060 model (minimum bound, Reuss model) is consistent with temperature values between 1061 1500–1800 K. Another obstacle in interpretation originates in the stress-sensitivity of 1062 the relevant creep. The viscosity can decrease by ~ 2.5 orders of magnitude while de-1063 creasing the differential stress by one order of magnitude. In terms of acceptable ther-1064 mal state, the temperature consistent with our prediction would decrease roughly by \sim 1065 100 K considering two-fold higher differential stress and increase by the same value for 1066 two-fold lower stress, respectively. 1067

Consequently, we find acceptable solutions both below and above the solidus. Our three-layered model thus cannot exclude or confirm a possible partial melt presence. An alternative explanation for the viscosity reduction can be the presence of water (see also Karato, 2013, for a deeper dicussion), which would also reduce the solidus temperature and facilitate partial melting. Both the enrichment in ilmenite and elevated water content can lead to the desired value of viscosity at lower temperatures compared to the dry and/or ilmenite-free models (Figure 12).

Focusing now on the elastic properties, we note that the rigidities of olivine (e.g. 1075 Mao et al., 2015), ilmenite (Jacobs et al., 2022), and garnetite (Kraettli et al., 2022) are 1076 comparable. The temperature has only a limited impact on their value (-0.01 GPa/K)1077 for olivine and ilmenite). Also, dependence on the water content (olivine-brucite) is only 1078 moderate (-1.3 GPa/wt%; Jacobsen et al., 2008). The magnitude of rigidity is, there-1079 fore, rather insensitive to possible constituents, temperature and water content. The up-1080 per bound of basal layer's rigidity predicted here ($\sim 60 \,\mathrm{GPa}$ for $R_{\mathrm{LVZ}} = 400 \,\mathrm{km}$, $\sim 70 \,\mathrm{GPa}$ 1081 for $R_{\rm LVZ} = 500$ km and ~ 85 GPa for $R_{\rm LVZ} = 700$ km) fits the elastic properties of all 1082

considered minerals—ilmenite, olivine, and garnet. However, the lower bound values (for 1083 $R_{\rm LVZ} > 500 \,\rm km$) are difficult to explain by the changes in composition, high temper-1084 ature, and/or water content. 1085



Figure 13. Impact of melt on the viscosity and rigidity contrast. The viscosity and rigidity contrast expressed as a function of the ϕ/ϕ_c (ϕ denotes the porosity and ϕ_c the critical porosity and parameterised using Kervazo et al. (2021); η_{solid} and μ_{solid} represents values with no melt present at the solidus temperature; no change in composition is considered. The shaded areas depict the predicted contrasts.

1086 1087

The magnitude of rigidity (Figure 13) is, nevertheless, sensitive to the presence of melt around or above the disintegration point (characterised by the critical porosity ϕ_c), which describes the transition from the solid to liquid behaviour and its typical values 1088 lie between 25-40%. Similarly, the viscosity value is very sensitive to the presence of 1089 melt for porosity higher than ϕ_c . For low porosities, it follows an exponential (Arrhe-1090 nian) dependence. Figure 13 suggest that the predicted rheological contrasts in the nom-1091 inal case are consistent with $\phi \lesssim 1.1 \phi_c$ for shear modulus contrast and with $\phi > 1.1 \phi_c$ 1092 for the viscosity contrast. This apparent inconsistency may be accounted for by the pres-1093 ence of melt accompanied by the changes in composition of the basal layer and by the 1094 susceptibility of viscosity to these changes. Consequently, the knowledge of the contrasts 1095 in both rheological parameters (rigidity and viscosity) could help tackle the trade-offs 1096 between porosity content and composition/temperature. Nevertheless, we must empha-1097 sise that the viscosity contrast predicted by our models is sensitive to the Andrade pa-1098 rameters of the mantle, leading to another uncertainty. 1099

The presence of a partially molten material would pose a strong constraint on the 1100 temperature and possible mode of the heat transfer in the lower mantle of the Moon, al-1101 lowing only models that reach the temperature between the solidus and liquidus (Fig-1102 ure 14). The traditional advective models predict stagnant-lid convection with a rela-1103 tively thick lid at present (e.g. Zhang et al., 2013). Below the stagnant lid, the temper-1104 ature follows the adiabatic or, for large internal heating, sub-adiabatic gradient. We es-1105 timate the temperature increase across the entire mantle due to the adiabatic gradient 1106 to be bounded by 100 K. Within those traditional models, it is plausible to reach solidus 1107 only in the lowermost thermal-compositional boundary layer. In the case of conductive 1108 models (e.g. Nimmo et al., 2012), the temperature gradient is steeper than the solidus 1109 gradient and the solidus temperature can be reached in the entire basal layer, given ap-1110 propriate internal heating (as demonstrated in Figure 14). Interestingly, the lunar se-1111 lenotherm determined by the inversions of lunar geophysical data combined with phase-1112 equilibrium computations (Khan et al., 2014) lies between the conductive and adiabatic 1113 gradients. 1114



Figure 14. Comparison of temperature profiles. Colour scale: conductive profile, calculated with the matrix propagator method; parameters as in Figure 9. Individual branches correspond to average heating 8, 9.5 and $11 \,\mathrm{nW/m^2}$ in the mantle. The coefficient f denotes the enrichment in the radiogenic elements of the basal layer ($R_{\rm LVZ} = 500 \,\mathrm{km}$) compared to the rest of the mantle. Gray area is the temperature profile adapted from Khan et al. (2014); darker blue lines: peridotite solidus (solid), water-saturated solidus (dotted), and liquidus (dashed) according to Katz et al. (2003); light blue lines: clinopyroxene+ilmenite solidus (solid) and liquidus (dashed) according to Wyatt (1977).

In the future, distinct sensitivity of rigidity, viscosity, and other transport properties to temperature, melt fraction, and composition may provide a way to separate the interior thermal and composition structure. At present, inversion errors and the uncertainties on material properties cannot confirm or rule out the existence of a partially molten basal layer. It therefore remains a valid hypothesis.

1120

6.3 Other Sources of Information

The two models discussed here — one with a highly dissipative basal layer and the 1121 other with elastically-accommodated GBS in the mantle — cannot be distinguished from 1122 each other by the available selenodetic measurements. To answer the question stated in 1123 the title of our paper, one would need to resort to other types of empirical data. Among 1124 all geophysical methods devised for the exploration of planetary interiors, seismology is 1125 of foremost importance. Therefore, a question that cannot be solved by the interpreta-1126 tion of lunar tidal response might be answered by comparing the arrival times and the 1127 phases detected at individual seismic stations. 1128

As we mentioned in Introduction, the Moon demonstrates a nearside-farside seis-1129 mic asymmetry. Judging by the currently available seismic data collected on the near 1130 side, the deep interior of the far side is virtually aseismic or, alternatively, the seismic 1131 waves emanating from it are strongly attenuated or deflected. The existence of an aseis-1132 mic area on the farside might not be entirely inconceivable. First, as pointed out by Nakamura 1133 (2005), there are large zones with no located nests of deep moonquakes even on the near-1134 side; and, in fact, most of the known deep seismic nests are part of an extended belt reach-1135 ing from the south-west to the north-east of the lunar face. Second, there exists a pro-1136 nounced dichotomy between the near side and far side of the Moon in terms of the crustal 1137 thickness, gravity field, and surface composition, which might point to a deeper, inter-1138 nal dichotomy as predicted by some evolutionary models (e.g., Laneuville et al., 2013; 1139 Zhu et al., 2019; Jones et al., 2022). 1140

An obvious way to illuminate the lack of deep farside moonquakes detected by the 1141 Apollo seismic stations would be to place seismometers on the far side of the Moon. They 1142 would observe the far side activity, and record the known repeating nearside moonquakes 1143 or events determined from impact flash observations. The Farside Seismic Suite (FSS) 1144 mission, recently selected for flight as part of the NASA PRISM program and planned 1145 for launch in 2024 or 2025, might provide such a measurement by delivering two seismome-1146 ters to Schrödinger Crater (Panning et al., 2021). While this crater is far from the an-1147 tipodes (in fact, close to the South pole), a seismometer residing in it should still be able 1148 to detect events from the far side, thereby addressing the hemispheric asymmetry in the 1149 Apollo observations. However, resolving polarisation of arrivals may be challenging for 1150 many moonquakes, meaning that many events will only have distance estimated, but not 1151 azimuth. (We are grateful to Mark P. Panning for an enlightening consultation on this 1152 topic.) 1153

A better site for this science objective would be the far side Korolev crater residing by the equator, about 23 degrees from the antipodes. It is now considered as one of the possible landing sites for the Lunar Geophysical Network (LGN) mission proposed to arrive on the Moon in 2030 and to deploy packages at four locations to enable geophysical measurements for 6 - 10 years (Fuqua Haviland et al., 2022).

Still, having a station or even an array of seismic stations at or near the antipodes would be ideal. Observed by such a station or stations, all events at distances less than 90 degrees from the antipodes could be confidently assigned to the far side. So we would recommend the near-antipodes zone as a high-priority landing site for some future mission, a perfect area to monitor the seismic activity on the far side and, especially, to observe if and how seismic waves proliferate through the base of the mantle.

In addition to seismic measurements, and similarly to what is predicted for Jupiter's 1165 volcanic moon Io or for icy moons with subsurface oceans, the presence of a highly dis-1166 sipative or a partially molten layer might be reflected in the tidal heating pattern on lu-1167 nar surface (e.g., Segatz et al., 1988; Tobie et al., 2005). However, as illustrated in the 1168 upper row of Figure 15, the positioning of the layer at the base of the mantle results in 1169 a very small difference between the surface heating patterns corresponding to the two 1170 alternative models. Both models show maxima of the average surface tidal heat flux Φ_{tide} 1171 on the lunar poles and minima on the "subterranean" ($\varphi = 0$) and antipodal ($\varphi = \pi$) 1172 points. Moreover, the magnitude of Φ_{tide} is generally very small, about three orders of 1173 magnitude lower than the flux produced by radiogenic heating of lunar interior (e.g., Siegler 1174 & Smrekar, 2014). The detection of any differences between the surface heat flux of the 1175 two models would be extremely challenging, if not impossible. 1176



Figure 15. Average surface tidal heat flux (top) and volumetric tidal heating (bottom) for a specific realisation of each of the two models discussed in this work: the model considering elastically-accommodated GBS through the Sundberg-Cooper rheological model (left) and the model with a basal low-viscosity zone (right). In particular, the volumetric tidal heating is plotted as a function of relative radius r/R and colatitude ϑ with longitude φ equal to 0.

The lower row of Figure 15 illustrates volumetric heat production due to tidal dis-1177 sipation. As pointed out by Harada et al. (2014), the presence of a low-viscosity zone 1178 at the base of the mantle results in considerable local increase of tidal heating with re-1179 spect to the rest of the mantle or to the model without the basal layer. While the tidal 1180 contribution to heat production in the high-viscosity parts of the mantle is around 10^{-11} W m⁻³, 1181 the tidal heat production in the basal layer reaches $\sim 10^{-8}$ W m⁻³. For comparison, the 1182 global average of mantle heat production by all sources (radiogenic and tidal) is estimated 1183 to be 6.3×10^{-9} W m⁻³ (Siegler & Smrekar, 2014). The predicted tidal dissipation in 1184 the basal layer can help to locally increase the temperature and exceed the solidus, es-1185 pecially if conductive heat transfer prevails in the lunar mantle. Combined with a high 1186 enrichment of the basal layer in heat producing elements, it may then contribute to main-1187 taining the presence of melt. 1188

Although virtually discarded in the beginning of this Subsection, let us nevertheless discuss possible insights provided by future high-precision tidal measurements. At present, the quality factor Q at tidal frequencies is obtained exclusively from fitting the ¹¹⁹² lunar physical libration, empirically determined by LLR. However, increased precision ¹¹⁹³ of satellite tracking (Dirkx et al., 2019; Hu et al., 2022; Stark et al., 2022) might even-¹¹⁹⁴ tually enable the determination of lunar tidal phase lag from the gravity field. Having ¹¹⁹⁵ an independent determination of tidal Q, which is related to the phase lag, would serve ¹¹⁹⁶ as a verification of the method used for fitting the LLR time series.

Among the quantities that we used in the inversion was degree-3 potential Love 1197 number k_3 . This parameter is currently only known with a large error bar but its refine-1198 ment would only help to discern between the two alternative models considered here if 1199 the elastically-accommodated GBS was contributing to the dissipation throughout the entire mantle (and not only in greater depths, as tentatively derived in Subsection 6.1). 1201 This is a consequence of a degree-dependent sensitivity of Love numbers to the interior 1202 structure. While degree-2 Love numbers and quality factors probe the lunar interior down 1203 to the core, higher-order quantities are only sensitive to shallower depths. The Love num-1204 ber k_3 —or the quality factor Q_3 —would thus not "see" the basal low-viscosity layer, but 1205 it might sense complex tidal response in the upper mantle. As a result, the detection of 1206 the unexpected frequency dependence of tidal dissipation even in Q_3 (accompanied by a relatively high $k_3 \sim 0.01$) would clearly point at a mechanism acting in shallow depths. 1208

Interestingly, the two alternative models can be better distinguished from each other 1209 in case the secondary peak of tidal dissipation, resulting either from the existence of a 1210 weak basal layer or from the Sundberg-Cooper model, lies at frequencies close to 10^{-4} rad s⁻¹. 1211 Then, provided that the elastically-accommodated GBS is only active below distinct depths (400-600 km), one could see a difference in predicted h_2 of the two models. Indepen-1213 dently on that depth, the models with secondary dissipation peak close to 10^{-4} rad s⁻¹ 1214 also differ in elastic Love number $k_{2,e}$, which can be calculated for interior structures ob-1215 tained from the inversion of seismic waves (as was done by Weber et al., 2011). Specif-1216 ically, $k_{2,e}$ in the melt-free model is then much lower that that of the model with a weak 1217 basal layer. The value reported by Weber et al. (2011), which is $k_{2,e} = 0.0232$, is at-1218 tained by both the alternative models for a secondary tidal dissipation peak lying at \sim 1219 $10^{-5.5}$ rad s⁻¹. In that case, the models are already indistinguishable. Seismic Q in the 1220 melt-free part of the mantle (at 1 Hz) for the models mentioned in the previous sentence 1221 is around 800 - 1000. 1222

Finally, we would like to note that any increase in the precision of Q determina-1223 tion will greatly help in answering the question whether any specific source of additional 1224 dissipation, be it a weak basal layer or elastic accommodation of strain at grain bound-1225 aries, is necessary in the first place. Recall that in order to fit the two alternative mod-1226 els to the tidal data, we assumed that the uncertainty on Q is of the order of 1% the mean value. In reality, the empirical Q at the monthly and the annual frequencies present an 1228 uncertainty between 10 and 20%. Keeping the original uncertainties, we were still able 1229 to fit the tidal data with the standard Andrade model, although with an unrealistically 1230 small exponential factor. 1231

1232 7 Conclusions

Tidal effects strongly depend not only on the interior density, viscosity, and rigidity profiles of celestial bodies, but also on the implied deformation mechanisms, which are reflected in the rheological models adopted. In this work, we attempted to illustrate that the unexpected frequency dependence of the tidal Q measured by LLR (Williams & Boggs, 2015) can be explained by lunar interior models both with and without a partially molten basal layer, and that each of the considered models leads to a different set of constraints on the interior properties.

As a first guess, we fitted the lunar tidal parameters (k_2, k_3, h_2, Q) at the monthly frequency and k_2/Q at the annual frequency) with a model consisting of a fluid core and a viscoelastic mantle governed by the Andrade rheology. Within that model, and setting $\zeta = 1$ (i.e., the time scales of viscoelastic and anelastic deformation were consid-

ered comparable) we found a mantle viscosity of $\log \eta_{\rm m}[{\rm Pa~s}] = 22.99^{+0.89}_{-1.35}$, mantle rigid-1244 ity of $\log \mu_{\rm m}[{\rm Pa}] = 10.92 \pm 0.06$, and the Andrade parameter α as low as $0.06^{+0.04}_{-0.02}$. The 1245 predicted value of α is generally lower than reported in the literature (0.1-0.4; e.g., Jack-1246 son et al., 2010; Castillo-Rogez et al., 2011; Efroimsky, 2012a, 2012b). This observation 1247 leads us to the conclusion that the tidal response of the Moon probably cannot be ex-1248 plained by the Andrade model alone and requires either a basal low-viscosity zone (in 1249 line with the conclusion of Khan et al., 2014) or an additional dissipation mechanism in 1250 the mantle (similar to Nimmo et al., 2012). 1251

Throughout Section 5, we have seen that the two alternative models expected to 1252 explain the anomalous frequency dependence of lunar Q (assumed to be known with an 1253 arbitrarily chosen high precision) cannot be distinguished from each other by the exist-1254 ing measurements of tidal deformation and dissipation alone. In the two-layered model 1255 consisting of a liquid core and a Sundberg-Cooper mantle, the fitting of tidal parame-1256 ters requires the relaxation time τ associated with elastically-accommodated GBS to be 1257 in the range from 3 to 300 hours. The corresponding relaxation strength Δ is predicted 1258 to lie in the interval [0.03, 1]. For a nominal case with Andrade parameters $\alpha = 0.2$ and $\zeta = 1$, we further obtain a mantle viscosity of log $\eta_m[Pa s] = 22.55^{+0.15}_{-0.54}$ and a mantle 1260 rigidity $\log \mu_{\rm m}[{\rm Pa}] = 10.84^{0.14}_{-0.02}$. 1261

In the three-layered model containing a liquid core, a low-rigidity basal layer, and 1262 an Andrade mantle, the tidal parameters are consistent with a wide range of basal layer 1263 thicknesses $D_{\rm LVZ}$ and rigidities $\mu_{\rm LVZ}$. As a general rule, a thicker layer implies weaker 1264 constraints on its rigidity, allowing both melt-like and solid-like behaviour. The predicted 1265 values of $\mu_{\rm LVZ}$ are consistent with elastic properties of all considered minerals (olivine, 1266 ilmenite, granite) and with a wide range of lower-mantle temperatures. In contrast to 1267 the rigidity, the viscosity η_{INZ} of the basal layer is constrained relatively well and falls 1268 into the range from about 10^{15} to 3×10^{16} Pa s, with a preference for the lower values 1269 $(\log \eta_{\text{LVZ}}[\text{Pa s}] = 15.20^{+0.53}_{-0.21})$. This is also in accordance with the results of Efroimsky 1270 (2012a, 2012b); Harada et al. (2014, 2016); Matsumoto et al. (2015); Tan and Harada 1271 (2021), and Kronrod et al. (2022). Nevertheless, even the viscosity is not able to pose 1272 strong constraints on the lower-mantle temperature, owing to the large uncertainties both 1273 on tidal Q and on the rheological properties of lunar minerals. For the viscosity and rigid-1274 ity of the overlying mantle in the nominal case, we get $\log \eta_{\rm m}[{\rm Pa~s}] = 22.79^{+0.19}_{-0.06}$ and 1275 $\log \mu_{\rm m}[{\rm Pa}] = 10.88 \pm 0.03.$ 1276

The existence of a basal weak or possibly semi-molten layer in the mantles of ter-1277 restrial bodies has been recently also suggested for Mercury (Steinbrügge et al., 2021) 1278 and for Mars (Samuel et al., 2021). In the case of Mercury, a lower mantle viscosity as low as 10^{13} Pa s was proposed to match the latest measurements of the moment of in-1280 ertia and of k_2 ; although this result was later critically reassessed by Goossens et al. (2022), 1281 who showed that more realistic values around 10^{18} Pa s might still explain the observa-1282 tions. In the case of Mars, the putative basal semi-molten layer was introduced by Samuel 1283 et al. (2021) to provide an alternative fit to seismic data which would not require the ex-1284 istence of a large core with unexpectedly high concentration of light elements (reported 1285 in Stähler et al., 2021). Lastly, large provinces of decreased shear seismic velocities also exist at the base of the Earth's mantle. These zones form a heterogeneous pattern in the 1287 deep terrestrial interior; however, according to numerical models, the formation of a con-1288 tinuous layer right above the core-mantle boundary is also possible for some values of 1289 model parameters (e.g., Dannberg et al., 2021). A new question thus arises: is a weak 1290 basal layer something common among terrestrial planet's mantles? Is it a natural and 1291 widely present outcome of magma ocean solidification and subsequent dynamical pro-1292 cesses? Or is it merely a popular explanation of the data available? 1293

Since the available tidal parameters were deemed insufficient to distinguish a weak basal layer above the lunar core from the manifestation of elastically accommodated GBS in the mantle, we conclude that an answer to the question stated in the title of our pa-

per awaits future lunar seismic experiments (ideally with a uniform distribution of seis-1297 mometers across the lunar surface) as well as a better understanding of elastic param-1298 eters of olivine-ilmenite assemblages near their melting point. Additionally, a tighter bound 1299 on the hypothetical basal layer parameters or on the strength and position of the secondary Debye peak in the alternative, Sundberg-Cooper model might be given by up-1301 dated values of tidal Q at multiple frequencies or by an independent inference of inte-1302 rior dissipation from the tidal phase lag and frequency-dependent k_2 , theoretically mea-1303 surable by laser altimetry or orbital tracking data (Dirkx et al., 2019; Hu et al., 2022; 1304 Stark et al., 2022). A combination of all those sources of information will probably still 1305 not provide a bright picture of deep lunar interior; however, it will help us to refute at 1306 least some of the many possible interior models. 1307

1308 Open Research

The software developed for the calculation of tidal Love numbers and quality factors of multi-layered bodies, the Python interface for running the MCMC inversion, and the plotting tools used for the figures presented in this study will be made available at the GitHub repository of the corresponding author (https://github.com/kanovami/ Lunar_Q) and preserved at [DOI to be added later during the peer review process] under the licence [to be added later during the peer review process].

1315 Acknowledgments

The authors would like to thank James G. Williams, Mark P. Panning, and Alexander 1316 S. Konopliv for extremely helpful conversations on various aspects of the lunar science. 1317 M.W. is grateful to Ana-Catalina Plesa and Martin Knapmeyer for discussions about 1318 the lunar interior and to Philipp A. Baumeister for introducing her to the Python libraries 1319 used in this work. She also gratefully acknowledges the financial support and endorse-1320 ment from the DLR Management Board Young Research Group Leader Program and 1321 the Executive Board Member for Space Research and Technology. M.B. received fund-1322 ing from Czech Science Foundation grant no. GA22-20388S. 1323

1324 References

1328

1329

- Bagheri, A., Efroimsky, M., Castillo-Rogez, J., Goossens, S., Plesa, A.-C., Rambaux,
 N., ... Giardini, D. (2022, August). Tidal insights into rocky and icy bodies:
 An introduction and overview. Advances in Geophysics, 63, 231-320.
 - Biot, M. A. (1954). Theory of stress-strain relations in anisotropic viscoelasticity and relaxation phenomena. *Journal of Applied Physics*, 25(11), 1385–1391.
- Bolmont, E., Breton, S. N., Tobie, G., Dumoulin, C., Mathis, S., & Grasset, O.
 (2020, December). Solid tidal friction in multi-layer planets: Application to
 Earth, Venus, a Super Earth and the TRAPPIST-1 planets. Potential approximation of a multi-layer planet as a homogeneous body. Astronomy &
 Astrophysics, 644, A165. doi: 10.1051/0004-6361/202038204
- Boué, G., & Efroimsky, M. (2019). Tidal evolution of the keplerian elements. Ce *lestial Mechanics and Dynamical Astronomy*, 131, 30. doi: 10.1007/s10569-019
 -9908-2
- 1338Castillo-Rogez, J. C., Efroimsky, M., & Lainey, V.(2011, September).The1339tidal history of Iapetus: Spin dynamics in the light of a refined dissipationmodel.Journal of Geophysical Research (Planets), 116(E9), E09008.doi:134110.1029/2010JE003664
- Dannberg, J., Myhill, R., Gassmöller, R., & Cottaar, S. (2021, November). The
 morphology, evolution and seismic visibility of partial melt at the core-mantle
 boundary: implications for ULVZs. *Geophysical Journal International*, 227(2),
 1028-1059. doi: 10.1093/gji/ggab242

1346 1347	Darwin, G. H. (1879, January). On the Analytical Expressions Which Give the History of a Fluid Planet of Small Viscosity, Attended by a Single Satellite. Pro-
1348	ceedings of the Royal Society of London Series I, 30, 255-278.
1349	Dirkx, D., Prochazka, I., Bauer, S., Visser, P., Noomen, R., Gurvits, L. I., & Ver-
1350	meersen, B. (2019, November). Laser and radio tracking for planetary sci-
1351	ence missions—a comparison. Journal of Geodesy, $93(11)$, 2405-2420. doi:
1352	10.1007/s00190-018-1171-x
1353	Dumoulin, C., Tobie, G., Verhoeven, O., Rosenblatt, P., & Rambaux, N. (2017,
1354	June). Tidal constraints on the interior of Venus. Journal of Geophysical
1355	Research (Planets), 122(6), 1338-1352. doi: 10.1002/2016JE005249
1356	Dygert, N., Hirth, G., & Liang, Y. (2016). A flow law for ilmenite in dis-
1357	location creep: Implications for lunar cumulate mantle overturn. Geo-
1358	physical Research Letters, 43(2), 532-540. Retrieved from https://
1359	agupubs.onlinelibrary.wiley.com/doi/abs/10.1002/2015GL066546 doi:
1360	https://doi.org/10.1002/2015GL066546
1361	Efroimsky, M. (2012a, March). Bodily tides near spin-orbit resonances. <i>Celestial</i>
1362	Mechanics and Dynamical Astronomy, $112(3)$, 283-330. doi: 10.1007/s10569
1363	-011-9397-4
1364	Efroimsky, M. (2012b, February). Tidal Dissipation Compared to Seismic Dissipa-
1365	tion: In Small Bodies, Earths, and Super-Earths. The Astrophysical Journal,
1366	74b(2), 150. doi: 10.1088/0004-037X/740/2/150
1367	Efroimsky, M. (2015, October). Tidal Evolution of Asteroidal Binaries. Ruled by
1368	Viscosity. Ignorant of Rigidity. The Astronomical Journal, 150(4), 98. doi: 10
1369	.1088/0004-0250/150/4/98
1370	Efformsky, M., & Makarov, V. V. (2013). Tidal friction and tidal lagging, applica-
1371	bility limitations of a popular formula for the tidal torque. The Astrophysical $L_{introphysical}$
1372	Journal, 104, 20. doi: 10.1088/0004-037A/704/1/20
1373	Enrollinsky, M., & Makarov, V. V. (2014, November). I Idal Dissipation in a Homogeneous Spherical Dady. I. Mathada, The Astrophysical Journal $205(1)$ 6, doi:
1374	neous Spherical Dody. I. Methods. The Astrophysical Journal, $795(1)$, 0. doi: 10.1088/0004.627X/705/1/6
1375	
1376	The MCMC Hammer Publications of the Astronomical Society of the Pacific
1377	125(925) 306 doi: 10.1086/670067
1370	Foreman-Mackey D (2016 iun) corner by: Scatterplot matrices in bython The
1379	Journal of Open Source Software 1(2) 24 Retrieved from https://doi org/
1381	10.21105/joss.00024 doi: 10.21105/joss.00024
1382	Frohlich, C., & Nakamura, Y. (2009, April). The physical mechanisms of deep
1383	moonquakes and intermediate-depth earthquakes: How similar and how dif-
1384	ferent? Physics of the Earth and Planetary Interiors, 173(3-4), 365-374. doi:
1385	10.1016/j.pepi.2009.02.004
1386	Fuqua Haviland, H., Weber, R. C., Neal, C. R., Lognonné, P., Garcia, R. F.,
1387	Schmerr, N., Bremner, P. M. (2022, February). The Lunar Geophysi-
1388	cal Network Landing Sites Science Rationale. The Planetary Science Journal,
1389	3(2), 40. doi: 10.3847/PSJ/ac0f82
1390	Garcia, R. F., Gagnepain-Beyneix, J., Chevrot, S., & Lognonné, P. (2011, Septem-
1391	ber). Very preliminary reference Moon model. Physics of the Earth and Plan-
1392	etary Interiors, 188(1), 96-113. doi: 10.1016/j.pepi.2011.06.015
1393	Garcia, R. F., Khan, A., Drilleau, M., Margerin, L., Kawamura, T., Sun, D.,
1394	Zhu, P. (2019, November). Lunar Seismology: An Update on Interior Structure
1395	Models. Space Science Reviews, 215(8), 50. doi: 10.1007/s11214-019-0613-y
1396	Gerstenkorn, H. (1967, January). The Earth as a Maxwell body. Icarus, $6(1-3)$, 92-
1397	99. doi: 10.1016/0019-1035(67)90006-1
1398	Gevorgyan, Y. (2021, June). Homogeneous model for the TRAPPIST-1e planet with
1399	an icy layer. Astronomy & Astrophysics, 650, A141. doi: $10.1051/0004-6361/$
1400	202140736

1401	Ghahremani, F. (1980). Effect of grain boundary sliding on anelasticity of poly-
1402	crystals. International Journal of Solids and Structures, 16(9), 825-845.
1403	Retrieved from https://www.sciencedirect.com/science/article/pii/
1404	0020768380900529 doi: https://doi.org/10.1016/0020-7683(80)90052-9
1405	Goodman, J., & Weare, J. (2010, January). Ensemble samplers with affine invari-
1406	ance. Communications in Applied Mathematics and Computational Science,
1407	5(1), 65-80. doi: $10.2140/camcos.2010.5.65$
1408	Goossens, S., Matsumoto, K., Liu, Q., Kikuchi, F., Sato, K., Hanada, H., Chen,
1409	M. (2011, April). Lunar gravity field determination using SELENE same-beam
1410	differential VLBI tracking data. Journal of Geodesy, 85(4), 205-228. doi:
1411	10.1007/s00190-010-0430-2
1412	Goossens, S., Renaud, J. P., Henning, W. G., Mazarico, E., Bertone, S., & Genova,
1413	A. (2022, February). Evaluation of Recent Measurements of Mercury's Mo-
1414	ments of Inertia and Tides Using a Comprehensive Markov Chain Monte Carlo
1415	Method. The Planetary Science Journal, 3(2), 37. doi: 10.3847/PSJ/ac4bb8
1416	Harada, Y., Goossens, S., Matsumoto, K., Yan, J., Ping, J., Noda, H., & Haruyama,
1417	J. (2014, August). Strong tidal heating in an ultralow-viscosity zone at the
1418	core-mantle boundary of the Moon. Nature Geoscience, 7(8), 569-572. doi:
1419	10.1038/ngeo2211
1420	Harada, Y., Goossens, S., Matsumoto, K., Yan, J., Ping, J., Noda, H., & Haruyama,
1421	J. (2016, September). The deep lunar interior with a low-viscosity zone: Re-
1422	vised constraints from recent geodetic parameters on the tidal response of the
1423	Moon. Icarus, 276, 96-101. doi: 10.1016/j.icarus.2016.04.021
1424	Hirth, G., & Kohlstedt, D. L. (1996, October). Water in the oceanic upper
1425	mantle: implications for rheology, melt extraction and the evolution of the
1426	lithosphere. Earth and Planetary Science Letters, 144(1-2), 93-108. doi:
1427	10.1016/0012-821X(96)00154-9
1428	Hu, X., Stark, A., Dirkx, D., Hussmann, H., Fienga, A., Fayolle-Chambe, M.,
1429	Oberst, J. (2022, May). Sensitivity analysis of frequency-dependent visco-
1430	elastic effects on lunar orbiters. In Egu general assembly conference abstracts
1431	(p. EGU22-9722). doi: 10.5194/egusphere-egu22-9722
1432	Jackson, I., Faul, U. H., & Skelton, R. (2014, March). Elastically accommodated
1433	grain-boundary sliding: New insights from experiment and modeling. <i>Physics</i>
1434	of the Earth and Planetary Interiors, 228, 203-210. doi: 10.1016/j.pepi.2013.11
1435	.014
1436	Jackson, I., Faul, U. H., Suetsugu, D., Bina, C., Inoue, T., & Jellinek, M. (2010,
1437	November). Grainsize-sensitive viscoelastic relaxation in olivine: Towards a ro-
1438	bust laboratory-based model for seismological application. Physics of the Earth
1439	and Planetary Interiors, 183(1-2), 151-163. doi: 10.1016/j.pepi.2010.09.005
1440	Jacobs, M. H. G., van den Berg, A. P., Schmid-Fetzer, R., de Vries, J., van Westre-
1441	nen, W., & Zhao, Y. (2022, July). Thermodynamic properties of geikielite
1442	$(MgTiO_3)$ and ilmenite $(FeTiO_3)$ derived from vibrational methods combined
1443	with Raman and infrared spectroscopic data. Physics and Chemistry of Miner-
1444	als, $49(7)$, 23. doi: 10.1007/s00269-022-01195-5
1445	Jacobsen, S. D., Jiang, F., Mao, Z., Duffy, T. S., Smyth, J. R., Holl, C. M., & Frost,
1446	D. J. (2008, July). Effects of hydration on the elastic properties of olivine.
1447	Geophysical Research Letters, $35(14)$, L14303. doi: $10.1029/2008$ GL034398
1448	Jones, M. J., Evans, A. J., Johnson, B. C., Weller, M. B., Andrews-Hanna, J. C.,
1449	Tikoo, S. M., & Keane, J. T. (2022, April). A South Pole–Aitken impact origin
1450	of the lunar compositional asymmetry. $Science Advances, 8(14), eabm8475.$
1451	doi: 10.1126 /sciadv.abm8475
1452	Karato, Si. (2013, December). Geophysical constraints on the water content of the
1453	lunar mantle and its implications for the origin of the Moon. Earth and Plane-
1454	tary Science Letters, 384, 144-153. doi: 10.1016/j.epsl.2013.10.001
1455	Katz, R. F., Spiegelman, M., & Langmuir, C. H. (2003). A new parameterization

1456	of hydrous mantle melting. $Geochemistry, Geophysics, Geosystems, 4(9).$
1457	Retrieved from https://agupubs.onlinelibrary.wiley.com/doi/abs/
1458	10.1029/2002GC000433 doi: https://doi.org/10.1029/2002GC000433
1459	Kawamura, T., Lognonné, P., Nishikawa, Y., & Tanaka, S. (2017, July). Evaluation
1460	of deep moonquake source parameters: Implication for fault characteristics and
1461	thermal state. Journal of Geophysical Research (Planets), 122(7), 1487-1504.
1462	doi: 10.1002/2016JE005147
1463	Kê, TS. (1947, April). Experimental Evidence of the Viscous Behavior of Grain
1464	Boundaries in Metals. <i>Physical Review</i> , 71(8), 533-546. doi: 10.1103/PhysRev
1465	.71.533
1466	Kervazo, M., Tobie, G., Choblet, G., Dumoulin, C., & Běhounková, M. (2021, June).
1467	Solid tides in lo's partially molten interior. Contribution of bulk dissipation.
1468	Astronomy & Astrophysics, 650 , A72. doi: $10.1051/0004-6361/202039433$
1469	Khan, A., Connolly, J. A. D., Pommier, A., & Noir, J. (2014, October). Geophysical
1470	evidence for melt in the deep lunar interior and implications for lunar evolu- tion L_{constant} (R_{constant}) ($10(10)$) (107) (2001) (100)
1471	tion. Journal of Geophysical Research (Planets), $119(10)$, $2197-2221$. doi: 10.1002/2014. IE004661
1472	Venerlin A C Dark D C Ven D N Arman C W Wething M M Williams
1473	L C Zuber M T (2013 July) The IPI Jupar gravity field to spherical
1475	harmonic degree 660 from the GRAIL Primary Mission — Journal of Ceonhusi-
1475	cal Research (Planets) 118(7) 1415-1434 doi: 10.1002/jgre 20097
1477	Kraettli G Schmidt M W & Liebske C (2022) Fractional crystallization of
1478	a basal lunar magma ocean: A dense melt-bearing garnetite laver above the
1479	core? Icarus, 371, 114699. Retrieved from https://www.sciencedirect.com/
1480	science/article/pii/S0019103521003547 doi: https://doi.org/10.1016/
1481	j.icarus.2021.114699
1482	Kronrod, E., Matsumoto, K., Kuskov, O. L., Kronrod, V., Yamada, R., & Kamata,
1483	S. (2022, April). Towards geochemical alternatives to geophysical models of the
1484	internal structure of the lunar mantle and core. Advances in Space Research,
1485	69(7), 2798-2824.doi: 10.1016/j.asr.2022.01.012
1486	Laneuville, M., Wieczorek, M. A., Breuer, D., & Tosi, N. (2013, July). Asymmet-
1487	ric thermal evolution of the Moon. Journal of Geophysical Research (Planets),
1488	118(7), 1435-1452. doi: 10.1002/jgre.20103
1489	Lee, L. C., & Morris, S. J. S. (2010, March). Anelasticity and grain boundary
1490	sliding. Proceedings of the Royal Society of London Series A, 466(2121), 2651-
1491	2671. doi: 10.1098/rspa.2009.0624
1492	Lee, L. C., Morris, S. J. S., & Wilkening, J. (2011, June). Stress concentrations,
1493	diffusionally accommodated grain boundary sliding and the viscoelasticity of
1494	polycrystals. Proceedings of the Royal Society of London Series A, 467 (2130),
1495	1024-1044. doi: $10.1098/rspa.2010.0447$
1496	Lemome, r. G., GOOSSENS, S., SADAKA, I. J., NICHOIAS, J. B., MAZATICO, E., KOW- lands D. D. Zuhor, M. T. (2013 August). High damage gravity models
1497	from CRAIL primary mission data Lowroal of Coonhusical Research (Planeta)
1498	118(8) 1676-1698 doi: 10.1002/jgrg 20118
1499	Li H Zhang N Liang V Wu B Dygert N I Huang I & Parmentier E M
1500	(2010) Lunar cumulate mantle overturn: A model constrained by ilmenite
1502	rheology Journal of Geonhusical Research: Planets 12/(5) 1357-1378
1503	Retrieved from https://agupubs.onlinelibrarv.wilev.com/doi/abs/
1504	10.1029/2018JE005905 doi: https://doi.org/10.1029/2018JE005905
1505	Mao, Z., Fan, D., Lin, JF., Yang, J., Tkachev, S. N., Zhuravlev, K., & Prakapenka
1506	V. B. (2015, September). Elasticity of single-crystal olivine at high pressures
1507	and temperatures. Earth and Planetary Science Letters, 426, 204-215. doi:
1508	10.1016/j.epsl.2015.06.045
1509	Matsumoto, K., Yamada, R., Kikuchi, F., Kamata, S., Ishihara, Y., Iwata, T.,
1510	Sasaki, S. (2015, September). Internal structure of the Moon inferred from

1511	Apollo seismic data and selenodetic data from GRAIL and LLR. Geophysical
1512	Research Letters, 42(18), 7351-7358. doi: 10.1002/2015GL065335
1513	Matsuyama, I., Nimmo, F., Keane, J. T., Chan, N. H., Taylor, G. J., Wieczorek,
1514	M. A., Williams, J. G. (2016, August). GRAIL, LLR, and LOLA con-
1515	straints on the interior structure of the Moon. Geophysical Research Letters,
1516	43(16), 8365-8375. doi: 10.1002/2016GL069952
1517	Mazarico, E., Barker, M. K., Neumann, G. A., Zuber, M. T., & Smith, D. E. (2014,
1518	April). Detection of the lunar body tide by the Lunar Orbiter Laser Altimeter.
1519	Geophysical Research Letters, 41(7), 2282-2288. doi: 10.1002/2013GL059085
1520	Morris, S. J. S., & Jackson, I. (2009, April). Diffusionally assisted grain-boundary
1521	sliding and viscoelasticity of polycrystals. Journal of Mechanics and Physics of
1522	Solids, 57(4), 744-761. doi: 10.1016/j.jmps.2008.12.006
1523	Mosegaard, K., & Tarantola, A. (1995, July). Monte Carlo sampling of solutions
1524	to inverse problems. Journal of Geophysical Research, 100(B7), 12,431-12,447.
1525	doi: 10.1029/94JB03097
1526	Nakamura, Y. (2005, January). Farside deep moonquakes and deep interior of the
1527	Moon. Journal of Geophysical Research (Planets), 110(E1), E01001. doi: 10
1528	.1029/2004JE002332
1529	Nakamura, Y., Lammlein, D., Latham, G., Ewing, M., Dorman, J., Press, F., &
1530	Toksoz, N. (1973, July). New Seismic Data on the State of the Deep Lunar
1531	Interior. Science, 181 (4094), 49-51. doi: 10.1126/science.181.4094.49
1532	Nakamura, Y., Latham, G., Lammlein, D., Ewing, M., Duennebier, F., & Dorman,
1533	J. (1974, January). Deep lunar interior inferred from recent seismic data.
1534	Geophysical Research Letters, $1(3)$, 137-140, doi: 10.1029/GL001j003p00137
1535	Nimmo, F., Faul, U. H., & Garnero, F. J. (2012, September). Dissipation at tidal
1536	and seismic frequencies in a melt-free Moon. Journal of Geophysical Research
1537	(<i>Planets</i>), 117(E9), E09005, doi: 10.1029/2012JE004160
1538	Novelles B Frouard J Makarov V V & Efroimsky M (2014 October) Spin-
1530	orbit evolution of Mercury revisited <i>Learns</i> 2/1 26-44 doi: 10.1016/j.jcarus
1540	.2014.05.045
1541	Nunn C. Garcia R. F. Nakamura Y. Marusiak A. G. Kawamura T. Sun D.
1542	
1543	Review. Space Science Reviews. 216(5), 89. doi: 10.1007/s11214-020-00709-3
1544	Panning, M., Kedar, S., Bowles, N., Calcutt, S., Cutler, J., Elliott, J.,, Yana, C.
1545	(2021, December). Farside Seismic Suite (FSS): First seismic data from the
1546	farside of the Moon delivered by a commercial lander. In Agu fall meeting ab-
1547	stracts (Vol. 2021, p. P54C-01). Retrieved from https://www.hou.usra.edu/
1548	meetings/lpsc2022/pdf/1576.pdf
1549	Pavlov, D. A., Williams, J. G., & Suvorkin, V. V. (2016, November). Determining
1550	parameters of Moon's orbital and rotational motion from LLR observations
1551	using GRAIL and IERS-recommended models. Celestial Mechanics and Dy-
1552	namical Astronomy, 126(1-3), 61-88, doi: 10.1007/s10569-016-9712-1
1553	Qin, C., Muirhead, A. C., & Zhong, S. (2012, July). Correlation of deep moon-
1554	quakes and mare basalts: Implications for lunar mantle structure and evolu-
1555	tion. <i>Icarus</i> , 220(1), 100-105, doi: 10.1016/j.jcarus.2012.04.023
1556	Baevskiv S N Gudkova T V Kuskov O L & Kronrod V A (2015 Jan-
1557	uary) On reconciling the models of the interior structure of the moon with
1558	gravity data. <i>Izvesting. Physics of the Solid Earth.</i> 51(1), 134-142. doi:
1559	10.1134/S1069351315010127
1560	Rai, R., & Ashby, M. F. (1971, January). On grain boundary sliding and diffusional
1561	creep. Metalluraical Transactions, 2, 1113-1127. doi: 10.1007/BF02664244
1562	Renaud, J. P., & Henning, W. G. (2018 April) Increased Tidal Dissipation Using
1563	Advanced Rheological Models: Implications for Io and Tidally Active Exoplan-
1564	ets. The Astrophysical Journal, 857(2), 98. doi: 10.3847/1538-4357/aab784
1565	Sabadini, R., & Vermeersen, B. (2004). Global Dynamics of the Earth Annlica-

1566	tions of Normal Mode Relaxation Theory to Solid-Earth Geophysics. Dodrech,
1567	the Netherlands: Knuwer Academic Publishers.
1568	Samuel, H., Balliner, M. D., Padovan, S., 10si, N., Rivoldini, A., & Plesa, AC.
1569	(2021, April). The Thermo Chemical Evolution of Mars with a Strongly Strat-
1570	ined Mantie. Journal of Geophysical Research (Flanets), 120(4), 600015. doi: 10.1020/2020 IE006612
1571	10.1029/2020JE000013
1572	Segatz, M., Spohn, T., Ross, M. N., & Schubert, G. (1988). Tidal Dissipation,
1573	Surface Heat Flow, and Figure of Viscoelastic Models of Io. $Icarus, 75(2),$
1574	18(-206, doi: 10.1016/0019-1035(88)90001-2
1575	Siegler, M. A., & Smrekar, S. E. (2014, January). Lunar heat flow: Regional
1576	(<i>Planets</i>) 119(1) 47-63 doi: 10.1002/2013 IE004453
1577	Stähler S C Khan A Banerdt W B Lognonné P Ciardini D Covlan S
1578	Stanler, S. C., Khan, A., Daherdt, W. D., Loghonne, T., Glardini, D., Oeylan, S., Smrokar, S. F. (2021, July) Soismic detection of the martian core. Science
1579	272(6553) 443 448 doi: 10.1126/scionce.pbi7730
1580	Stark A Viao H Hu X Fiange A Huggmann H Obergt I Saliby C
1581 1582	(2022, May). Measurement of tidal deformation through self-registration of
1583	laser profiles: Application to Earth's Moon. In Equ general assembly confer-
1584	ence abstracts (p. EGU22-10626). doi: 10.5194/egusphere-egu22-10626
1585	Steinbrügge, G., Dumberry, M., Rivoldini, A., Schubert, G., Cao, H., Schroeder,
1586	D. M., & Soderlund, K. M. (2021, February). Challenges on Mercury's In-
1587	terior Structure Posed by the New Measurements of its Obliquity and Tides.
1588	Geophys. Res. Lett., 48(3), e89895. doi: 10.1029/2020GL089895
1589	Sundberg, M., & Cooper, R. F. (2010). A composite viscoelastic model for incorpo-
1590	rating grain boundary sliding and transient diffusion creep; correlating creep
1591	and attenuation responses for materials with a fine grain size. <i>Philosophical</i>
1592	Magazine, 90(20), 2817-2840. Retrieved from https://doi.org/10.1080/
1593	14786431003746656 doi: 10.1080/14786431003746656
1594	Takeuchi, H., & Saito, M. (1972). Seismic Surface Waves. Methods in Compu-
1594 1595	Takeuchi, H., & Saito, M. (1972). Seismic Surface Waves. Methods in Compu- tational Physics: Advances in Research and Applications, 11, 217-295. doi: 10
1594 1595 1596	Takeuchi, H., & Saito, M. (1972). Seismic Surface Waves. Methods in Compu- tational Physics: Advances in Research and Applications, 11, 217-295. doi: 10 .1016/B978-0-12-460811-5.50010-6
1594 1595 1596 1597	 Takeuchi, H., & Saito, M. (1972). Seismic Surface Waves. Methods in Computational Physics: Advances in Research and Applications, 11, 217-295. doi: 10.1016/B978-0-12-460811-5.50010-6 Tan, Y., & Harada, Y. (2021, September). Tidal constraints on the low-viscosity
1594 1595 1596 1597 1598	 Takeuchi, H., & Saito, M. (1972). Seismic Surface Waves. Methods in Computational Physics: Advances in Research and Applications, 11, 217-295. doi: 10.1016/B978-0-12-460811-5.50010-6 Tan, Y., & Harada, Y. (2021, September). Tidal constraints on the low-viscosity zone of the Moon. Icarus, 365, 114361. doi: 10.1016/j.icarus.2021.114361
1594 1595 1596 1597 1598 1599	 Takeuchi, H., & Saito, M. (1972). Seismic Surface Waves. Methods in Computational Physics: Advances in Research and Applications, 11, 217-295. doi: 10.1016/B978-0-12-460811-5.50010-6 Tan, Y., & Harada, Y. (2021, September). Tidal constraints on the low-viscosity zone of the Moon. Icarus, 365, 114361. doi: 10.1016/j.icarus.2021.114361 Thor, R. N., Kallenbach, R., Christensen, U. R., Gläser, P., Stark, A., Steinbrügge,
1594 1595 1596 1597 1598 1599 1600	 Takeuchi, H., & Saito, M. (1972). Seismic Surface Waves. Methods in Computational Physics: Advances in Research and Applications, 11, 217-295. doi: 10.1016/B978-0-12-460811-5.50010-6 Tan, Y., & Harada, Y. (2021, September). Tidal constraints on the low-viscosity zone of the Moon. Icarus, 365, 114361. doi: 10.1016/j.icarus.2021.114361 Thor, R. N., Kallenbach, R., Christensen, U. R., Gläser, P., Stark, A., Steinbrügge, G., & Oberst, J. (2021, January). Determination of the lunar body tide
1594 1595 1596 1597 1598 1599 1600 1601	 Takeuchi, H., & Saito, M. (1972). Seismic Surface Waves. Methods in Computational Physics: Advances in Research and Applications, 11, 217-295. doi: 10.1016/B978-0-12-460811-5.50010-6 Tan, Y., & Harada, Y. (2021, September). Tidal constraints on the low-viscosity zone of the Moon. Icarus, 365, 114361. doi: 10.1016/j.icarus.2021.114361 Thor, R. N., Kallenbach, R., Christensen, U. R., Gläser, P., Stark, A., Steinbrügge, G., & Oberst, J. (2021, January). Determination of the lunar body tide from global laser altimetry data. Journal of Geodesy, 95(1), 4. Retrieved
1594 1595 1596 1597 1598 1599 1600 1601 1602	 Takeuchi, H., & Saito, M. (1972). Seismic Surface Waves. Methods in Computational Physics: Advances in Research and Applications, 11, 217-295. doi: 10.1016/B978-0-12-460811-5.50010-6 Tan, Y., & Harada, Y. (2021, September). Tidal constraints on the low-viscosity zone of the Moon. Icarus, 365, 114361. doi: 10.1016/j.icarus.2021.114361 Thor, R. N., Kallenbach, R., Christensen, U. R., Gläser, P., Stark, A., Steinbrügge, G., & Oberst, J. (2021, January). Determination of the lunar body tide from global laser altimetry data. Journal of Geodesy, 95(1), 4. Retrieved from https://www.hou.usra.edu/meetings/lpsc2022/pdf/1576.pdf doi:
1594 1595 1596 1597 1598 1599 1600 1601 1602 1603	 Takeuchi, H., & Saito, M. (1972). Seismic Surface Waves. Methods in Computational Physics: Advances in Research and Applications, 11, 217-295. doi: 10.1016/B978-0-12-460811-5.50010-6 Tan, Y., & Harada, Y. (2021, September). Tidal constraints on the low-viscosity zone of the Moon. Icarus, 365, 114361. doi: 10.1016/j.icarus.2021.114361 Thor, R. N., Kallenbach, R., Christensen, U. R., Gläser, P., Stark, A., Steinbrügge, G., & Oberst, J. (2021, January). Determination of the lunar body tide from global laser altimetry data. Journal of Geodesy, 95(1), 4. Retrieved from https://www.hou.usra.edu/meetings/lpsc2022/pdf/1576.pdf doi: 10.1007/s00190-020-01455-8
1594 1595 1596 1597 1598 1600 1600 1602 1603 1604	 Takeuchi, H., & Saito, M. (1972). Seismic Surface Waves. Methods in Computational Physics: Advances in Research and Applications, 11, 217-295. doi: 10.1016/B978-0-12-460811-5.50010-6 Tan, Y., & Harada, Y. (2021, September). Tidal constraints on the low-viscosity zone of the Moon. Icarus, 365, 114361. doi: 10.1016/j.icarus.2021.114361 Thor, R. N., Kallenbach, R., Christensen, U. R., Gläser, P., Stark, A., Steinbrügge, G., & Oberst, J. (2021, January). Determination of the lunar body tide from global laser altimetry data. Journal of Geodesy, 95(1), 4. Retrieved from https://www.hou.usra.edu/meetings/lpsc2022/pdf/1576.pdf doi: 10.1007/s00190-020-01455-8 Tobie, G., Grasset, O., Lunine, J. I., Mocquet, A., & Sotin, S. (2005). Tidal dis-
1594 1595 1596 1597 1598 1600 1601 1602 1603 1604 1605	 Takeuchi, H., & Saito, M. (1972). Seismic Surface Waves. Methods in Computational Physics: Advances in Research and Applications, 11, 217-295. doi: 10.1016/B978-0-12-460811-5.50010-6 Tan, Y., & Harada, Y. (2021, September). Tidal constraints on the low-viscosity zone of the Moon. Icarus, 365, 114361. doi: 10.1016/j.icarus.2021.114361 Thor, R. N., Kallenbach, R., Christensen, U. R., Gläser, P., Stark, A., Steinbrügge, G., & Oberst, J. (2021, January). Determination of the lunar body tide from global laser altimetry data. Journal of Geodesy, 95(1), 4. Retrieved from https://www.hou.usra.edu/meetings/lpsc2022/pdf/1576.pdf doi: 10.1007/s00190-020-01455-8 Tobie, G., Grasset, O., Lunine, J. I., Mocquet, A., & Sotin, S. (2005). Tidal dissipation within large icy satellites: Applications to Europa and Titan. Icarus,
1594 1595 1596 1597 1598 1600 1601 1602 1603 1604 1605 1606	 Takeuchi, H., & Saito, M. (1972). Seismic Surface Waves. Methods in Computational Physics: Advances in Research and Applications, 11, 217-295. doi: 10.1016/B978-0-12-460811-5.50010-6 Tan, Y., & Harada, Y. (2021, September). Tidal constraints on the low-viscosity zone of the Moon. Icarus, 365, 114361. doi: 10.1016/j.icarus.2021.114361 Thor, R. N., Kallenbach, R., Christensen, U. R., Gläser, P., Stark, A., Steinbrügge, G., & Oberst, J. (2021, January). Determination of the lunar body tide from global laser altimetry data. Journal of Geodesy, 95(1), 4. Retrieved from https://www.hou.usra.edu/meetings/lpsc2022/pdf/1576.pdf doi: 10.1007/s00190-020-01455-8 Tobie, G., Grasset, O., Lunine, J. I., Mocquet, A., & Sotin, S. (2005). Tidal dissipation within large icy satellites: Applications to Europa and Titan. Icarus, 175(2), 496-502. doi: 10.1016/j.icarus.2004.12.007
1594 1595 1596 1597 1598 1600 1601 1602 1603 1604 1605 1606	 Takeuchi, H., & Saito, M. (1972). Seismic Surface Waves. Methods in Computational Physics: Advances in Research and Applications, 11, 217-295. doi: 10.1016/B978-0-12-460811-5.50010-6 Tan, Y., & Harada, Y. (2021, September). Tidal constraints on the low-viscosity zone of the Moon. Icarus, 365, 114361. doi: 10.1016/j.icarus.2021.114361 Thor, R. N., Kallenbach, R., Christensen, U. R., Gläser, P., Stark, A., Steinbrügge, G., & Oberst, J. (2021, January). Determination of the lunar body tide from global laser altimetry data. Journal of Geodesy, 95(1), 4. Retrieved from https://www.hou.usra.edu/meetings/lpsc2022/pdf/1576.pdf doi: 10.1007/s00190-020-01455-8 Tobie, G., Grasset, O., Lunine, J. I., Mocquet, A., & Sotin, S. (2005). Tidal dissipation within large icy satellites: Applications to Europa and Titan. Icarus, 175(2), 496-502. doi: 10.1016/j.icarus.2004.12.007 van Kan Parker, M., Sanloup, C., Sator, N., Guillot, B., Tronche, E. J., Perrillat,
1594 1595 1596 1597 1598 1600 1601 1602 1603 1604 1605 1606 1607 1608	 Takeuchi, H., & Saito, M. (1972). Seismic Surface Waves. Methods in Computational Physics: Advances in Research and Applications, 11, 217-295. doi: 10.1016/B978-0-12-460811-5.50010-6 Tan, Y., & Harada, Y. (2021, September). Tidal constraints on the low-viscosity zone of the Moon. Icarus, 365, 114361. doi: 10.1016/j.icarus.2021.114361 Thor, R. N., Kallenbach, R., Christensen, U. R., Gläser, P., Stark, A., Steinbrügge, G., & Oberst, J. (2021, January). Determination of the lunar body tide from global laser altimetry data. Journal of Geodesy, 95(1), 4. Retrieved from https://www.hou.usra.edu/meetings/lpsc2022/pdf/1576.pdf doi: 10.1007/s00190-020-01455-8 Tobie, G., Grasset, O., Lunine, J. I., Mocquet, A., & Sotin, S. (2005). Tidal dissipation within large icy satellites: Applications to Europa and Titan. Icarus, 175(2), 496-502. doi: 10.1016/j.icarus.2004.12.007 van Kan Parker, M., Sanloup, C., Sator, N., Guillot, B., Tronche, E. J., Perrillat, JP., van Westrenen, W. (2012, March). Neutral buoyancy of titanium-
1594 1595 1596 1597 1598 1600 1601 1602 1603 1604 1605 1606 1606 1608 1609	 Takeuchi, H., & Saito, M. (1972). Seismic Surface Waves. Methods in Computational Physics: Advances in Research and Applications, 11, 217-295. doi: 10.1016/B978-0-12-460811-5.50010-6 Tan, Y., & Harada, Y. (2021, September). Tidal constraints on the low-viscosity zone of the Moon. Icarus, 365, 114361. doi: 10.1016/j.icarus.2021.114361 Thor, R. N., Kallenbach, R., Christensen, U. R., Gläser, P., Stark, A., Steinbrügge, G., & Oberst, J. (2021, January). Determination of the lunar body tide from global laser altimetry data. Journal of Geodesy, 95(1), 4. Retrieved from https://www.hou.usra.edu/meetings/lpsc2022/pdf/1576.pdf doi: 10.1007/s00190-020-01455-8 Tobie, G., Grasset, O., Lunine, J. I., Mocquet, A., & Sotin, S. (2005). Tidal dissipation within large icy satellites: Applications to Europa and Titan. Icarus, 175(2), 496-502. doi: 10.1016/j.icarus.2004.12.007 van Kan Parker, M., Sanloup, C., Sator, N., Guillot, B., Tronche, E. J., Perrillat, JP., van Westrenen, W. (2012, March). Neutral buoyancy of titaniumrich melts in the deep lunar interior. Nature Geoscience, 5(3), 186-189. doi:
1594 1595 1596 1597 1598 1600 1601 1602 1603 1604 1605 1606 1607 1608 1609 1610	 Takeuchi, H., & Saito, M. (1972). Seismic Surface Waves. Methods in Computational Physics: Advances in Research and Applications, 11, 217-295. doi: 10.1016/B978-0-12-460811-5.50010-6 Tan, Y., & Harada, Y. (2021, September). Tidal constraints on the low-viscosity zone of the Moon. Icarus, 365, 114361. doi: 10.1016/j.icarus.2021.114361 Thor, R. N., Kallenbach, R., Christensen, U. R., Gläser, P., Stark, A., Steinbrügge, G., & Oberst, J. (2021, January). Determination of the lunar body tide from global laser altimetry data. Journal of Geodesy, 95(1), 4. Retrieved from https://www.hou.usra.edu/meetings/lpsc2022/pdf/1576.pdf doi: 10.1007/s00190-020-01455-8 Tobie, G., Grasset, O., Lunine, J. I., Mocquet, A., & Sotin, S. (2005). Tidal dissipation within large icy satellites: Applications to Europa and Titan. Icarus, 175(2), 496-502. doi: 10.1016/j.icarus.2004.12.007 van Kan Parker, M., Sanloup, C., Sator, N., Guillot, B., Tronche, E. J., Perrillat, JP., van Westrenen, W. (2012, March). Neutral buoyancy of titaniumrich melts in the deep lunar interior. Nature Geoscience, 5(3), 186-189. doi: 10.1038/ngeo1402
1594 1595 1596 1597 1598 1600 1601 1602 1603 1604 1605 1606 1607 1608 1609 1610	 Takeuchi, H., & Saito, M. (1972). Seismic Surface Waves. Methods in Computational Physics: Advances in Research and Applications, 11, 217-295. doi: 10.1016/B978-0-12-460811-5.50010-6 Tan, Y., & Harada, Y. (2021, September). Tidal constraints on the low-viscosity zone of the Moon. Icarus, 365, 114361. doi: 10.1016/j.icarus.2021.114361 Thor, R. N., Kallenbach, R., Christensen, U. R., Gläser, P., Stark, A., Steinbrügge, G., & Oberst, J. (2021, January). Determination of the lunar body tide from global laser altimetry data. Journal of Geodesy, 95(1), 4. Retrieved from https://www.hou.usra.edu/meetings/1psc2022/pdf/1576.pdf doi: 10.1007/s00190-020-01455-8 Tobie, G., Grasset, O., Lunine, J. I., Mocquet, A., & Sotin, S. (2005). Tidal dissipation within large icy satellites: Applications to Europa and Titan. Icarus, 175(2), 496-502. doi: 10.1016/j.icarus.2004.12.007 van Kan Parker, M., Sanloup, C., Sator, N., Guillot, B., Tronche, E. J., Perrillat, JP., van Westrenen, W. (2012, March). Neutral buoyancy of titaniumrich melts in the deep lunar interior. Nature Geoscience, 5(3), 186-189. doi: 10.1038/ngeo1402 Viswanathan, V., Fienga, A., Minazzoli, O., Bernus, L., Laskar, J., & Gastineau, M.
1594 1595 1596 1597 1598 1600 1601 1602 1603 1604 1605 1606 1607 1608 1609 1610 1611	 Takeuchi, H., & Saito, M. (1972). Seismic Surface Waves. Methods in Computational Physics: Advances in Research and Applications, 11, 217-295. doi: 10.1016/B978-0-12-460811-5.50010-6 Tan, Y., & Harada, Y. (2021, September). Tidal constraints on the low-viscosity zone of the Moon. Icarus, 365, 114361. doi: 10.1016/j.icarus.2021.114361 Thor, R. N., Kallenbach, R., Christensen, U. R., Gläser, P., Stark, A., Steinbrügge, G., & Oberst, J. (2021, January). Determination of the lunar body tide from global laser altimetry data. Journal of Geodesy, 95(1), 4. Retrieved from https://www.hou.usra.edu/meetings/lpsc2022/pdf/1576.pdf doi: 10.1007/s00190-020-01455-8 Tobie, G., Grasset, O., Lunine, J. I., Mocquet, A., & Sotin, S. (2005). Tidal dissipation within large icy satellites: Applications to Europa and Titan. Icarus, 175(2), 496-502. doi: 10.1016/j.icarus.2004.12.007 van Kan Parker, M., Sanloup, C., Sator, N., Guillot, B., Tronche, E. J., Perrillat, JP., van Westrenen, W. (2012, March). Neutral buoyancy of titaniumrich melts in the deep lunar interior. Nature Geoscience, 5(3), 186-189. doi: 10.1038/ngeo1402 Viswanathan, V., Fienga, A., Minazzoli, O., Bernus, L., Laskar, J., & Gastineau, M. (2018, May). The new lunar ephemeris INPOP17a and its application to fun-
1594 1595 1596 1597 1598 1600 1601 1602 1603 1604 1605 1606 1607 1608 1610 1611 1612	 Takeuchi, H., & Saito, M. (1972). Seismic Surface Waves. Methods in Computational Physics: Advances in Research and Applications, 11, 217-295. doi: 10.1016/B978-0-12-460811-5.50010-6 Tan, Y., & Harada, Y. (2021, September). Tidal constraints on the low-viscosity zone of the Moon. Icarus, 365, 114361. doi: 10.1016/j.icarus.2021.114361 Thor, R. N., Kallenbach, R., Christensen, U. R., Gläser, P., Stark, A., Steinbrügge, G., & Oberst, J. (2021, January). Determination of the lunar body tide from global laser altimetry data. Journal of Geodesy, 95(1), 4. Retrieved from https://www.hou.usra.edu/meetings/lpsc2022/pdf/1576.pdf doi: 10.1007/s00190-020-01455-8 Tobie, G., Grasset, O., Lunine, J. I., Mocquet, A., & Sotin, S. (2005). Tidal dissipation within large icy satellites: Applications to Europa and Titan. Icarus, 175(2), 496-502. doi: 10.1016/j.icarus.2004.12.007 van Kan Parker, M., Sanloup, C., Sator, N., Guillot, B., Tronche, E. J., Perrillat, JP., van Westrenen, W. (2012, March). Neutral buoyancy of titaniumrich melts in the deep lunar interior. Nature Geoscience, 5(3), 186-189. doi: 10.1038/ngeo1402 Viswanathan, V., Fienga, A., Minazzoli, O., Bernus, L., Laskar, J., & Gastineau, M. (2018, May). The new lunar ephemeris INPOP17a and its application to fundamental physics. Monthly Notices of the Royal Astronomical Society, 476(2),
1594 1595 1596 1597 1598 1600 1601 1602 1603 1604 1605 1606 1606 1607 1608 1609 1610 1611 1612 1613 1614	 Takeuchi, H., & Saito, M. (1972). Seismic Surface Waves. Methods in Computational Physics: Advances in Research and Applications, 11, 217-295. doi: 10.1016/B978-0-12-460811-5.50010-6 Tan, Y., & Harada, Y. (2021, September). Tidal constraints on the low-viscosity zone of the Moon. Icarus, 365, 114361. doi: 10.1016/j.icarus.2021.114361 Thor, R. N., Kallenbach, R., Christensen, U. R., Gläser, P., Stark, A., Steinbrügge, G., & Oberst, J. (2021, January). Determination of the lunar body tide from global laser altimetry data. Journal of Geodesy, 95(1), 4. Retrieved from https://www.hou.usra.edu/meetings/1psc2022/pdf/1576.pdf doi: 10.1007/s00190-020-01455-8 Tobie, G., Grasset, O., Lunine, J. I., Mocquet, A., & Sotin, S. (2005). Tidal dissipation within large icy satellites: Applications to Europa and Titan. Icarus, 175(2), 496-502. doi: 10.1016/j.icarus.2004.12.007 van Kan Parker, M., Sanloup, C., Sator, N., Guillot, B., Tronche, E. J., Perrillat, JP., van Westrenen, W. (2012, March). Neutral buoyancy of titaniumrich melts in the deep lunar interior. Nature Geoscience, 5(3), 186-189. doi: 10.1038/ngeo1402 Viswanathan, V., Fienga, A., Minazzoli, O., Bernus, L., Laskar, J., & Gastineau, M. (2018, May). The new lunar ephemeris INPOP17a and its application to fundamental physics. Monthly Notices of the Royal Astronomical Society, 476(2), 1877-1888. doi: 10.1093/mnras/sty096
1594 1595 1596 1597 1598 1600 1601 1602 1603 1604 1605 1606 1607 1608 1609 1610 1611 1612 1613 1614	 Takeuchi, H., & Saito, M. (1972). Seismic Surface Waves. Methods in Computational Physics: Advances in Research and Applications, 11, 217-295. doi: 10.1016/B978-0-12-460811-5.50010-6 Tan, Y., & Harada, Y. (2021, September). Tidal constraints on the low-viscosity zone of the Moon. Icarus, 365, 114361. doi: 10.1016/j.icarus.2021.114361 Thor, R. N., Kallenbach, R., Christensen, U. R., Gläser, P., Stark, A., Steinbrügge, G., & Oberst, J. (2021, January). Determination of the lunar body tide from global laser altimetry data. Journal of Geodesy, 95(1), 4. Retrieved from https://www.hou.usra.edu/meetings/lpsc2022/pdf/1576.pdf doi: 10.1007/s00190-020-01455-8 Tobie, G., Grasset, O., Lunine, J. I., Mocquet, A., & Sotin, S. (2005). Tidal dissipation within large icy satellites: Applications to Europa and Titan. Icarus, 175(2), 496-502. doi: 10.1016/j.icarus.2004.12.007 van Kan Parker, M., Sanloup, C., Sator, N., Guillot, B., Tronche, E. J., Perrillat, JP., van Westrenen, W. (2012, March). Neutral buoyancy of titaniumrich melts in the deep lunar interior. Nature Geoscience, 5(3), 186-189. doi: 10.1038/ngeo1402 Viswanathan, V., Fienga, A., Minazzoli, O., Bernus, L., Laskar, J., & Gastineau, M. (2018, May). The new lunar ephemeris INPOP17a and its application to fundamental physics. Monthly Notices of the Royal Astronomical Society, 476(2), 1877-1888. doi: 10.1093/mnras/sty096 Viswanathan, V., Rambaux, N., Fienga, A., Laskar, J., & Gastineau, M. (2019,
1594 1595 1596 1597 1598 1600 1601 1602 1603 1604 1605 1606 1607 1608 1609 1610 1611 1612 1613 1614 1615 1616	 Takeuchi, H., & Saito, M. (1972). Seismic Surface Waves. Methods in Computational Physics: Advances in Research and Applications, 11, 217-295. doi: 10.1016/B978-0-12-460811-5.50010-6 Tan, Y., & Harada, Y. (2021, September). Tidal constraints on the low-viscosity zone of the Moon. Icarus, 365, 114361. doi: 10.1016/j.icarus.2021.114361 Thor, R. N., Kallenbach, R., Christensen, U. R., Gläser, P., Stark, A., Steinbrügge, G., & Oberst, J. (2021, January). Determination of the lunar body tide from global laser altimetry data. Journal of Geodesy, 95(1), 4. Retrieved from https://www.hou.usra.edu/meetings/lpsc2022/pdf/1576.pdf doi: 10.1007/s00190-020-01455-8 Tobie, G., Grasset, O., Lunine, J. I., Mocquet, A., & Sotin, S. (2005). Tidal dissipation within large icy satellites: Applications to Europa and Titan. Icarus, 175(2), 496-502. doi: 10.1016/j.icarus.2004.12.007 van Kan Parker, M., Sanloup, C., Sator, N., Guillot, B., Tronche, E. J., Perrillat, JP., van Westrenen, W. (2012, March). Neutral buoyancy of titaniumrich melts in the deep lunar interior. Nature Geoscience, 5(3), 186-189. doi: 10.1038/ngeo1402 Viswanathan, V., Fienga, A., Minazzoli, O., Bernus, L., Laskar, J., & Gastineau, M. (2018, May). The new lunar ephemeris INPOP17a and its application to fundamental physics. Monthly Notices of the Royal Astronomical Society, 476(2), 1877-1888. doi: 10.1093/mnras/sty096 Viswanathan, V., Rambaux, N., Fienga, A., Laskar, J., & Gastineau, M. (2019, July). Observational Constraint on the Radius and Oblateness of the Lunar
1594 1595 1596 1597 1598 1600 1601 1602 1603 1604 1605 1606 1607 1610 1611 1612 1613 1614 1615 1616 1617	 Takeuchi, H., & Saito, M. (1972). Seismic Surface Waves. Methods in Computational Physics: Advances in Research and Applications, 11, 217-295. doi: 10.1016/B978-0-12-460811-5.50010-6 Tan, Y., & Harada, Y. (2021, September). Tidal constraints on the low-viscosity zone of the Moon. Icarus, 365, 114361. doi: 10.1016/j.icarus.2021.114361 Thor, R. N., Kallenbach, R., Christensen, U. R., Gläser, P., Stark, A., Steinbrügge, G., & Oberst, J. (2021, January). Determination of the lunar body tide from global laser altimetry data. Journal of Geodesy, 95(1), 4. Retrieved from https://www.hou.usra.edu/meetings/lpsc2022/pdf/1576.pdf doi: 10.1007/s00190-020-01455-8 Tobie, G., Grasset, O., Lunine, J. I., Mocquet, A., & Sotin, S. (2005). Tidal dissipation within large icy satellites: Applications to Europa and Titan. Icarus, 175(2), 496-502. doi: 10.1016/j.icarus.2004.12.007 van Kan Parker, M., Sanloup, C., Sator, N., Guillot, B., Tronche, E. J., Perrillat, JP., van Westrenen, W. (2012, March). Neutral buoyancy of titaniumrich melts in the deep lunar interior. Nature Geoscience, 5(3), 186-189. doi: 10.1038/ngeo1402 Viswanathan, V., Fienga, A., Minazzoli, O., Bernus, L., Laskar, J., & Gastineau, M. (2018, May). The new lunar ephemeris INPOP17a and its application to fundamental physics. Monthly Notices of the Royal Astronomical Society, 476(2), 1877-1888. doi: 10.1093/mnras/sty096 Viswanathan, V., Rambaux, N., Fienga, A., Laskar, J., & Gastineau, M. (2019, July). Observational Constraint on the Radius and Oblateness of the Lunar Core-Mantle Boundary. Geophysical Research Letters, 46(13), 7295-7303. doi:
1594 1595 1596 1597 1598 1600 1601 1602 1603 1604 1605 1606 1607 1610 1611 1612 1613 1614 1615 1616 1617 1618	 Takeuchi, H., & Saito, M. (1972). Seismic Surface Waves. Methods in Computational Physics: Advances in Research and Applications, 11, 217-295. doi: 10.1016/B978-0-12-460811-5.50010-6 Tan, Y., & Harada, Y. (2021, September). Tidal constraints on the low-viscosity zone of the Moon. Icarus, 365, 114361. doi: 10.1016/j.icarus.2021.114361 Thor, R. N., Kallenbach, R., Christensen, U. R., Gläser, P., Stark, A., Steinbrügge, G., & Oberst, J. (2021, January). Determination of the lunar body tide from global laser altimetry data. Journal of Geodesy, 95(1), 4. Retrieved from https://www.hou.usra.edu/meetings/1psc2022/pdf/1576.pdf doi: 10.1007/s00190-020-01455-8 Tobie, G., Grasset, O., Lunine, J. I., Mocquet, A., & Sotin, S. (2005). Tidal dissipation within large icy satellites: Applications to Europa and Titan. Icarus, 175(2), 496-502. doi: 10.1016/j.icarus.2004.12.007 van Kan Parker, M., Sanloup, C., Sator, N., Guillot, B., Tronche, E. J., Perrillat, JP., van Westrenen, W. (2012, March). Neutral buoyancy of titaniumrich melts in the deep lunar interior. Nature Geoscience, 5(3), 186-189. doi: 10.1038/ngeo1402 Viswanathan, V., Fienga, A., Minazzoli, O., Bernus, L., Laskar, J., & Gastineau, M. (2018, May). The new lunar ephemeris INPOP17a and its application to fundamental physics. Monthly Notices of the Royal Astronomical Society, 476(2), 1877-1888. doi: 10.1093/mnras/sty096 Viswanathan, V., Rambaux, N., Fienga, A., Laskar, J., & Gastineau, M. (2019, July). Observational Constraint on the Radius and Oblateness of the Lunar Core-Mantle Boundary. Geophysical Research Letters, 46(13), 7295-7303. doi: 10.1029/2019GL082677
1594 1595 1596 1597 1598 1600 1601 1602 1603 1604 1605 1606 1607 1608 1609 1610 1611 1612 1613 1614 1615 1616 1617 1618 1619	 Takeuchi, H., & Saito, M. (1972). Seismic Surface Waves. Methods in Computational Physics: Advances in Research and Applications, 11, 217-295. doi: 10.1016/B978-0-12-460811-5.50010-6 Tan, Y., & Harada, Y. (2021, September). Tidal constraints on the low-viscosity zone of the Moon. Icarus, 365, 114361. doi: 10.1016/j.icarus.2021.114361 Thor, R. N., Kallenbach, R., Christensen, U. R., Gläser, P., Stark, A., Steinbrügge, G., & Oberst, J. (2021, January). Determination of the lunar body tide from global laser altimetry data. Journal of Geodesy, 95(1), 4. Retrieved from https://www.hou.usra.edu/meetings/1psc2022/pdf/1576.pdf doi: 10.1007/s00190-020-01455-8 Tobie, G., Grasset, O., Lunine, J. I., Mocquet, A., & Sotin, S. (2005). Tidal dissipation within large icy satellites: Applications to Europa and Titan. Icarus, 175(2), 496-502. doi: 10.1016/j.icarus.2004.12.007 van Kan Parker, M., Sanloup, C., Sator, N., Guillot, B., Tronche, E. J., Perrillat, JP., van Westrenen, W. (2012, March). Neutral buoyancy of titaniumrich melts in the deep lunar interior. Nature Geoscience, 5(3), 186-189. doi: 10.1038/ngeo1402 Viswanathan, V., Fienga, A., Minazzoli, O., Bernus, L., Laskar, J., & Gastineau, M. (2018, May). The new lunar ephemeris INPOP17a and its application to fundamental physics. Monthly Notices of the Royal Astronomical Society, 476(2), 1877-1888. doi: 10.1093/mnras/sty096 Viswanathan, V., Rambaux, N., Fienga, A., Laskar, J., & Gastineau, M. (2019, July). Observational Constraint on the Radius and Oblateness of the Lunar Core-Mantle Boundary. Geophysical Research Letters, 46(13), 7295-7303. doi: 10.1029/2019GL082677 Weber, R. C., Lin, PY., Garnero, E. J., Williams, Q., & Lognonné, P. (2011, Jan-

1621	.1126/science.1199375
1622	Williams, J. G., & Boggs, D. H. (2009). Lunar Core and Mantle. What Does
1623	LLR See? In Proceedings of the 16th international workshop on laser rang-
1624	ing held on 13-17 october 2008 in poznań, poland. edited by s. schilliak. pub-
1625	lished by: Space research centre, polish academy of sciences, warsaw (p. 101-
1626	120). Retrieved from http://cddis.gsfc.nasa.gov/lw16/docs/papers/
1627	proceedings_vol2.pdf,http://cddis.gsfc.nasa.gov/lw16/docs/papers/
1628	sci_1_Williams_p.pdf
1629	Williams, J. G., & Boggs, D. H. (2015, April). Tides on the Moon: Theory and de-
1630	termination of dissipation. Journal of Geophysical Research (Planets), 120(4),
1631	689-724. doi: 10.1002/2014JE004755
1632	Williams, J. G., Boggs, D. H., & Ratcliff, J. T. (2012, March). Lunar Moment of
1633	Inertia, Love Number, and Core. In 43rd annual lunar and planetary science
1634	<i>conference</i> (p. 2230).
1635	Williams, J. G., Boggs, D. H., Yoder, C. F., Ratcliff, J. T., & Dickey, J. O.
1636	(2001, November). Lunar rotational dissipation in solid body and molten
1637	core. Journal of Geophysical Research, 106(E11), 27933-27968. doi:
1638	10.1029/2000JE001396
1639	Williams, J. G., Konopliv, A. S., Boggs, D. H., Park, R. S., Yuan, DN., Lemoine,
1640	F. G., Zuber, M. T. (2014, July). Lunar interior properties from the
1641	GRAIL mission. Journal of Geophysical Research (Planets), 119(7), 1546-1578.
1642	doi: 10.1002/2013JE004559
1643	Wu, P., & Peltier, W. R. (1982). Viscous gravitational relaxation. Geophysical Jour-
1644	nal International, 70(2), 435-485. doi: 10.1111/j.1365-246X.1982.tb04976.x
1645	Wyatt, B. A. (1977, January). The melting and crystallisation behaviour of a
1646	natural clinopyroxene-ilmenite intergrowth. Contributions to Mineralogy and
1647	Petrology, 61(1), 1-9. doi: 10.1007/BF00375941
1648	Yan, J., Goossens, S., Matsumoto, K., Ping, J., Harada, Y., Iwata, T., Kawano,
1649	N. (2012, March). CEGM02: An improved lunar gravity model using
1650	Chang'E-1 orbital tracking data. Planetary and Space Science, $62(1)$, 1-9.
1651	doi: 10.1016/j.pss.2011.11.010
1652	Yan, J., Liu, S., Xiao, C., Ye, M., Cao, J., Harada, Y., Barriot, JP. (2020,
1653	April). A degree-100 lunar gravity model from the Chang'e 5TT mission.
1654	Astronomy & Astrophysics, 636, A45. doi: 10.1051/0004-6361/201936802
1655	Zhang, N., Parmentier, E. M., & Liang, Y. (2013, September). A 3-D numerical
1656	study of the thermal evolution of the Moon after cumulate mantle overturn:
1657	The importance of rheology and core solidification. Journal of Geophysical
1658	Research. Planets, 118(9), 1789-1804. doi: 10.1002/jgre.20121
1659	Zhao, Y., de Vries, J., van den Berg, A. P., Jacobs, M. H. G., & van Westrenen,
1660	W. (2019, April). The participation of ilmenite-bearing cumulates in lu-
1661	nar mantle overturn. Earth and Planetary Science Letters, 511, 1-11. doi:
1662	10.1016 J.epsi. 2019.01.022
1663	Znarkov, V. N., & Gudkova, I. V. (2005, September). Construction of Martian In-
1664	terior Model. Solar System Research, $39(5)$, $343-373$. doi: $10.1007/s11208-005$
1665	
1666	Znu, MH., Wunnemann, K., Potter, K. W. K., Kleine, T., & Morbidelli, A. (2019,
1667	August). Are the Moon's Nearside-Farside Asymmetries the Result of a Giant Interact? Lower last Q_{12} and Q_{23} and
1668	impact: Journal of Geophysical Kesearch (Planets), 124(8), 2117-2140. doi: 10
1669	.1029/2018JE005820