

Land use analysis of the N'ZI watershed of Côte d'Ivoire using multi-date and multi-source satellite data

Koffi Léon YAO¹, Koffi Eugène KOUAKOU¹, Michel Amani KOUASSI¹, Attoungbré Jean-Philippe DEGUY², and Moussa CAMARA³

¹Institut National Polytechnique Felix Houphouët-Boigny

²NANGUI ABROGOUA University

³Institut National Polytechnique Felix Houphouët-Boigny

December 7, 2022

Abstract

The uncontrolled rapid population growth in our regions and strong industrialization are putting pressure on natural resources, accelerating climate change and desertification. This study aims to follow the evolution of land use in the N'ZI watershed. Three images from Landsat 4 & 5 (1986), Landsat 7 (2000), and Landsat 8 (2020) were used to carry out this study. Remote sensing and geographic information systems (GIS) were used to monitor land cover as a whole. Various treatments were performed using Envi 5.1 and ArcGIS 10.4.1 software programs. The results showed that changes took place during the periods of 1986-2000, 2000-2020, and 1986-2020. The results of the analysis showed the regression of water surfaces from -64.95% to -52.47% during the period of 2000-2020 and 1986-2020, on the other hand, there is a great increase in bare-ground dwellings (373.63%) and low-cover soils (10.60%). These progressions were due to the destruction of forests -86.93%, savannas -3.97%, and agricultural areas -9.30% between 1986 and 2020

Hosted file

Liste -Tableau_article.docx available at <https://authorea.com/users/521645/articles/611128-land-use-analysis-of-the-n-zi-watershed-of-c%C3%B4te-d-ivoire-using-multi-date-and-multi-source-satellite-data>

Land use analysis of the N'ZI watershed of Côte d'Ivoire using multi-date and multi-source satellite data

K.L. YAO^{1,2,3}, K.E. KOUAKOU^{2,3}, A.M. KOUASSI^{2,3}, A. J-P. DEGUY⁴, M. CAMARA¹

¹National Polytechnic Institute Felix Houphouet-Boigny (INP-HB), Polytechnic Doctoral School, Yamoussoukro, Côte d'Ivoire, BP 1093, Yamoussoukro, Côte d'Ivoire

²National Polytechnic Institute Félix Houphouet-Boigny (INP-HB); Department of Earth Sciences and Mineral Resources

³Laboratory of Civil Engineering, Geosciences and Geographic Sciences, Yamoussoukro, Côte d'Ivoire

⁴NANGUI ABROGOUA University, Abidjan, Côte d'Ivoire.

Abstract

The uncontrolled rapid population growth in our regions and strong industrialization are putting pressure on natural resources, accelerating climate change and desertification. This study aims to follow the evolution of land use in the N'ZI watershed. Three images from Landsat 4 & 5 (1986), Landsat 7 (2000), and Landsat 8 (2020) were used to carry out this study. Remote sensing and geographic information systems (GIS) were used to monitor land cover as a whole. Various treatments were performed using Envi 5.1 and ArcGIS 10.4.1 software programs. The results showed that changes took place during the periods of 1986-2000, 2000-2020, and 1986-2020. The results of the analysis showed the regression of water surfaces from -64.95% to -52.47% during the period of 2000-2020 and 1986-2020, on the other hand, there is a great increase in bare-ground dwellings (373.63%) and low-cover soils (10.60%). These progressions were due to the destruction of forests - 86.93%, savannas -3.97%, and agricultural areas of -9.30% between 1986 and 2020.

Keywords: Land use, satellite data, N'ZI watershed Côte d'Ivoire.

Introduction

The forest resource of Côte d'Ivoire was estimated at 16 million hectares in 1900, and the country's forest cover was 7,850,864 ha in 1986 and 3,401,146 ha in 2015 (BNETD and Tera, 2016). The country lost a total of 78% of its crop cover, because, of the expansion of agriculture which occupied 62% of the Ivorian territory (BNETD and Tera, 2016). Forest area has fallen to more than 2 million hectares today (Koné et al., 2021; Assoma et al., 2021). Studies by Kouassi (2010), and Noho et al. (2018) on the N'ZI watershed area indicate a real decline in clear forests and forested savannas by more than 14.95%, Dense forests probably decrease by 1.93%, and shrub and/or grassland savannah by less than 15.15%.

The forests are composed of degraded secondary forests around the N'ZI River in particular (17% degraded forest, 10% light forest, and 3% gallery forest) (Noho et al., 2018). The watershed of the N'ZI River constitutes a feeding zone between the savannah and forest areas. This was the loop of cocoa and coffee during the period from 1960 to 1980 (Brou et al., 2005). The population in the study area has grown rapidly from 900,000 to 3,030,381 inhabitants (INS, 2014). Anthropogenic causes and rural land tenure have created great pressure on resources, particularly land degradation, thus leading to the abandonment of land for other sources of income. In the search for information leading to the abandonment of land to other more productive regions, remote sensing and geographical information systems (GIS) have been chosen to monitor the evolution of land use of the N'ZI watershed to inform public opinion. Remote sensing is a powerful tool for providing quantitative spatial information to assess land use changes. This tool allows the measurement of different forms of landscape changes according to the level of land use and the temporal dynamics of these changes (Dietzel et al., 2005). However, quantitative analysis of landscape changes is fundamentally dependent on available geographic data, including spatial data (cartographic, photographic, or satellite) and spatial resolution of data used. Then, visualization of land cover and land use of the N'ZI watershed, based on satellite data, could be a primary means of understanding changes in land use and decision-making, to the extent that it has expected benefits.

Changes in soil surface classification methods and criteria vary from author to author (Hoang et al., 2009). These methods fall into three processing techniques which are: visual interpretations, pixel-based methods, and object-based approaches (Desclée et al., 2006). Today, the main classification methods used are supervised and not supervised methods (Achbun et al., 2011). In this work, the aim is to use the supervised classification method to characterize the dynamics of land use and quantify the changes in land use, to identify their evolution in the watershed of N'ZI.

1. Presentation of the study area

Figure 1 shows the geographic location of the study area. This area is the sub-watershed of Bandama. It is drained by the N'ZI River and its tributaries, located between longitudes 3°49'-5°22' West and latitudes 6°00'-9°26' North. It covers an area of 35076.4 km², about 10.88% of the national territory. N'ZI river runs about 600 km long today. The watershed of the N'ZI is composed of 80 sub-prefectures and is limited to the north by the department of Ferkessédougou, to the south by the departments of Tiassalé and Divo, to the east by the department of Daoukro, and to the west by the department of Béoumi. N'ZI watershed is generally dominated by a fairly monotonous relief (Avenard et al., 1971), its altitude h varies from 668 m in the north to less than 130 m in the south. The main soil types are medium-saturated (North) and strongly-saturated (Central and South) ferritic soils (Monnet, 1972;

Koudou *et al.*, 2016). Due to its elongated geographical configuration, the N'ZI watershed (Figure 1) is representative of the major climatic groups of Côte d'Ivoire. The central part of the basin is characterized by a humid tropical regime. The south of the basin is characterized by an equatorial regime.




Figure 1: presentation map of the N'ZI watershed

2. Material and Methods

2.1. Data used

Three multi-date satellite images were used in this work to study the evolution of the basin's land use. These are images from Landsat 4 TM (Thematic Mapper) from 1986, Landsat 7 ETM+ (Enhanced Thematic Mapper plus) from 2000, and Landsat 8 OLI (Operational Land Imager) from 2020. The study area was covered by scenes that were: 196-054, 196-55, 196-056, 197-053, 197-054, and 197-055. These scenes were downloaded on <http://earthexplorer.usgs.gov>, under 30 m spatial resolution. Cartographic data on the evolution of the vegetation of Côte d'Ivoire was taken at a scale of 1: 200,000, from the Remote Sensing Mapping Center of **BNETD** (*national technical study and development office*), were used.

2.2. Methods

The approach used to study changes in land use of the N'ZI watershed is illustrated in Figure 2. The method adopted was divided into four phases. The first Phase consisted of downloading the data. The second was devoted to pre-treatments. The third was concerned with the treatment of satellite images. The fourth was dedicated to the analysis of the changes and the discussion.




Figure 2: Organizational chart analysis of the evolution of land use between 1986, 2000 and 2020

2.3.1. Image preprocessing (pre-treatment)

The image pre-processing in this work consisted of applying radiometric correction, and atmospheric correction under the QUAC and FLAASH modules to get good readings of these spectral images (Girard et al., 1999; Caloz and Pointet, 2002; Diallo et al., 2011; Tra Bi, 2013; Jofack-Sokeng, 2016; *Chaima et al.*, 2019). The geometric correction was applied while downloading, edited with the ArcGIS 10.4.1 software (downloading on: [desktop/arcgis-desktop/arcmap/10-4-1/online](https://desktop.arcgis.com/desktop/arcmap/10-4-1/online)), and

projected into the projection coordinate system: WGS_1984_UTM_Zone_30N. Then, the assembly is applied to the strips to obtain a single multi-spectral image. Mosaicking of the scenes was done to form a larger image covering the study area. Finally, the study area was extracted from the mosaic scenes using ENVI software (downloading on: <https://idl-and-envi.software.informer.com/5.1/>) to carry out the color composition of the bands. The combination of bands is as follows: bands 2, 3, and 5 for the 1986 image, bands 6, 5, and 4 or 4, 3, and 2 for the 2000 image, and bands 5, 6, and 7 for the 2020 image. These operations facilitated the interpretation and discrimination of regions of interest.

2.3.3. Image treatment

The processing of the images started with the calculation of spectral indices on ENVI software. The indices were chosen to describe and interpret the state of evolution of land use of the N'ZI watershed. This approach offers a way to analyze the bio-Geo-physical processes of landscape occupation and the degradation of the surface condition of the area. These indices described the state of a phenomenon and were based on an empirical approach using experimental data (Caloz et al., 2003; Diallo et al., 2011; Grimene et al., 2019; Tra Bi, 2013). There are many indices, but this work has used indices such as the NDVI (Normalized Difference Vegetation Index), the IB (Gloss Index), the IR (Soil Redness Index), the IC (Coloration Index), the IBI (Built Sector Based Index). The visual interpretation of the colored compositions as well as the visualization of the maps of the calculated indices allowed a good reading of the information sought (degradation or change of the surface state). The land use classes discriminated in this study were validated by reconnaissance trips carried out in the study area to check the control points for the 2020 images (17; 18; 26 and 27 December 2020; 31 march 2021; 01; 02 and 08 April 2021). This justified the choice of the supervised classification method. For the 1986 and 2000 images, their visualization on Google Earth served as a photo interpretation to attest to the information. Finally, based on all this information, the areas of interest have made it possible to distinguish the different land use classes to opt for supervised classification. Therefore, the Maximum Likelihood algorithm has been chosen to visualize the land use maps.

2.3.3. Quantification of changes in the land use of the N'ZI watershed

Quantification of changes in land use classes over the study period from 1986 to 2020 was used to detect changes in the study area using multi-date satellite images as follows:

- **Change (Δ) in the area of the unit concerned between 1986, 2000 and 2020**

Δ_1 , the difference in the area of the units concerned between 1986 and 2000;

Δ_2 , the difference in the area of the units concerned between 2000 and 2020;

$\Delta 3$, the difference in the area of the units concerned between 1986 and 2020.

If $\Delta 1$ or $\Delta 2$ or $\Delta 3 = 0$, the area of the unit is stable in time and space;

If $\Delta 1$ or $\Delta 2$ or $\Delta 3 > 0$, the area of the unit concerned is increasing;

Si $\Delta 1$ ou $\Delta 2$ ou $\Delta 3 < 0$, the area of the unit concerned is said to be in regression.

▪ **Change in the overall rate of change (T_g) in the area between 1986, 2000 and 2020**

The overall rate of change in the area of land use classes between years is estimated from the following equation, proposed by FAO (1996), which is commonly used to measure the growth of macroeconomic aggregates between two given periods (Gbombélé Soro et al., 2014; Alexis et al., 2018; Kpedenou et al., 2016b, and 2017; Salomon et al., 2021).

$$T_g = \frac{\Delta 2 - \Delta 1}{\Delta 1} \times 100$$

Finally, the change in the land cover of the N'ZI basin between the selected periods is certified by the difference in validation by analyzing the confusion matrix through the global precision and the Kappa coefficient (Congalton, 1991; Skupinski et al., 2009). To assess the degree of degradation of soil conditions in the basin, the methods already developed upstream and previously were used.

3. Results and discussions

3.1. Image corrections

Figure 3 shows the results of the radiometric and atmospheric corrections applied to the 1986, 2000, and 2020 image scenes respectively.




Figure 3: Raw image (a, b, and c), atmospheric (d, e, and f), and Radiometric (g, h, and i) corrections of Landsat images 1986 (a, b, and c), 2000 (d, e, and f), and 2020 (g, h, and i) respectively.

3.2. Image Mosaicking

It should be noted that the fusion of the bands follows the mosaics of the images. Fusion consisted of putting all the bands to produce a new image that retains the information contained in each of the original images. The results of the different treatments have led to the mosaic stage presented by the maps (1986, 2000, and 2020) in Figure 4. The different scene mosaics are intended to create a new satellite image by combining several image scenes that cover the N'ZI watershed. This section concluded with the extraction of the study area shown on the maps in Figure 4.




Figure 4: Mosaic of scenes and colored composition of bands.

3.3. Land use changes classes and maps

The pre-treatment and treatment methods contributed to 1986, 2000, and 2020 land use maps of the N'ZI watershed (Figure 5). The validation of the classification by the confusion matrix evolves significantly with Kappa values increasing slightly between 1986, 2000, and 2020 by 0.86; 0.90, and 0.92 respectively, and overall accuracy of 94.07%, 95.81%, and 94.39%. These values mean that more than 90% of the pixels in the three images were correctly classified according to the ground-truth data. The analysis of these images made it possible to define six (6) land use classes. These were the forest class; savannah class; low cover soil class; agroforestry and fallow plantation class; bare dwelling class; and water body class.





Figure 5: Land use maps of the N'ZI watershed from 1986; 2000 and 2020

▪ Variation of different classes

The analysis of the evolution of land use classes in a GIS environment is illustrated by the histograms in figure 6. It shows a strong regression of the Forest class (18.01%; 9.42%; 3.52%) to the advantages of the Bare-Floor Housing class which rose from 8.89% in 1986 to 9.45% in 2000 and then to 18.08% in 2020. Forest decline also benefited the savannah class, which rose from 10.85% in 1986 to 13.30% in 2000 and 20.62% in 2020. While the Low Covered Soil class shows a particular increase of 15.35%

over the period 2000-2020. There is an average increase of about 6% in the Plantations-agroforestry and fallow land class from 1986 to 2000 and a sharp decrease of about 14% from 2000 to 2020. In addition, the Water Bodies class increased between 1986 and 2000 and decreased between 2000 and 2020 (Figure 6).



Fo: Forest; Sa: Savannah; SFC: Sparsely Covered Soil; Pl-Agro-Ja: Plantations-agroforestry and fallow land; Ha-SN: Dwelling-bare soil; PE: Waterbody

Figure 6: Spatial evolution of land use classes 1986, 2000 and 2020

3.4. Quantification of changes in land use

The histograms in Figure 7 show two trends in land cover classes, the periods 1986-2000; 2000-2020, and 1986-2020 have revealed the changes which have occurred. Those pointing down indicate a loss in the land use class. This means a rate of change from these classes to others. While the upwardly oriented histograms highlight a gain. This means a rate of change from these classes to others. While the upwardly oriented histograms highlight a gain (Figure 7). This is a change of landscape from the study area to other landscapes and vice versa. The results show a regression of forests (Fo), water bodies (PE), and plantations-agroforestry and fallow (Pl-Agro-Ja). Except for the period 1986-2000, there is a progression of Water Bodies, and Plantations-agroforestry and fallow. There was a large increase in Savannah, low-cover soils, and bare dwellings over the periods of 1986-2000, 2000-2020, and 1986-2020 (Figure 7).




Figure 7: variation in and use of classes in the N'ZI watershed

3.5. Structuring changes

Figures 8, 9, and 10 show the different statistical variations in the temporal change of land use classes of the N'ZI watershed. After analyzing the evolution of the different classes, there is a strong transformation of land use classes over the periods of 1986-2000, 1986-2020, and 2000-2020.

General change in land use

Figure 8 provides information on the different variations in the overall change of forest, savannah, low cover, plantation-agroforestry and fallow, dwelling-bare land; and water body classes over the periods of 1986-2000, 2000-2020, and 1986-2020 that did not change.




Figure 8: Statistical evolution of changes in land use units between 1986, 2000 and 2020

changes in the different land use classes

Figure 9 shows the changes in the different land use classes considered during the different calibration periods of 1986-2000, 2000-2020, and 1986-2020.




Figure 9: Statistical summary of changes in land use units between 1986, 2000 and 2020

▪ variability of individual class changes

Figure 10 clearly illustrates the different changes in the percentage and area of the classes considered during the periods studied: 1986-2000, 2000-2020, and 1986-2020. The forest, low soil cover, and water classes have decreased significantly over the three periods, except for an increase in water from 1986 to 2000. On the other hand, there was a strong increase in the savannah, bare dwelling and agroforestry, and fallow classes over the periods 1986-2000 and 1986-2020. Except between 2000 and 2020 that there has been a decline in plantations-agroforestry and fallow land.




Figure 10: Statistical summary of changes in land use units between 1986, 2000 and 2020

3.6. Calculation of omission and commission errors

Errors of commission and errors of omission are summarized in Table 1.

Omission errors, the highest rates are recorded in the low cover class on the 2000 and 2020 images and the 1986 image in the Bare dwellings class.

Commission errors, the highest rates are also observed in the low coverage soil class for 1986, 2000, and 2020 images (Table 1). Error rates are very low, below 15% in general, except at the level of soil classes poorly covered in 2020.

Table 1: Variation of class discrimination errors on images (1986, 2000, and 2020)

1986	commission errors (%)	Omission errors (%)
Forest	11,54	2,58
Savannah	1,49	3,84
Plantations-agroforestry and fallows	6,07	6,16
Sparsely covered soil	27,22	1,52
Dwelling-bare soil	10,98	8,71
Waterbody	3,67	6,44
2000	commission errors (%)	Omission errors (%)
Forest	12,76	4,41
Savannah	4,73	5,53
Plantations-agroforestry and fallows	3,92	2,44
Sparsely covered soil	14,94	13,53
Dwelling-bare soil	5,42	7,44
Waterbody	0,46	3,01
2020	commission errors (%)	Omission errors (%)
Forest	21,85	1,69
Savannah	12,75	5,72
Plantations-agroforestry and fallows	1,94	2,49
Sparsely cover soil	43,86	70,09
Dwelling-bare soil	3,9	5,78
Waterbody	2,42	6,56

3.7. Discussion

The classification models and the different remote sensing images of 30 m spatial resolution were used to generate the land use classes as well as the detection of changes generally in the N'ZI watershed. These means of studying land use satellite data, several researchers have made use of them in recent work by showing the contribution of Landsat images in the discrimination of the main categories of land use (Diallo and Bao, 2010; Agoualé et al., 2017; Kouassi, 2010; Lienou, 2009; Mellor et al., 2013; Pelletier, 2017; Alexis et al., 2018; Noho et al., 2018; Useni et al., 2020; Salomon et al., 2021; *Zeng Ju et al., 2021*). The land use classification resulting from the analysis of the 1986 TM images, the 2000 ETM+ images, and the 2020 OLI images gave overall accuracies (94.07%, 95.81%, and 94.39%) and Kappa coefficients of 0.86; 0.90 and 0.92 respectively for the years 1986, 2000 and 2020. According to (Landis and Koch, 1977; *Pontius, 2000*) classifications are statistically good qualities and can be used wisely for the remainder of the study. Indeed, for an image analysis whose Kappa value is higher than 0.50, the results are good and exploitable. But, if the results are acceptable, they must not lose sight of the constraints encountered when analyzing the images used. Indeed, errors were recorded during the analysis of the images from several sources when discriminating against the classes mentioned. These errors can be explained by the confusion between classes with generally similar spectral signatures. In addition, the difficulties of visual observation make the distinction of the elements of land use delicate by the structure of the landscape offering homogeneous characters. This can be explained by the fact that Savannah, forest, and low-covered soils are very heterogeneous in terms of temporal evolution. They are developed, with rubber and cocoa plantations in the south and towards the center-north with cashew plantations; Also included are agricultural clearings and areas used for food crops in the previous year after the harvest (fallow). Also, cocoa plantations are mainly grown in forests. This can make it difficult to discriminate between the different formations. The same observation was made by (Kouassi, 2019; N'Guessan, 2020). Bare-ground dwellings blend with the savannah to the agricultural grip and savannah. Analysis of the results evolution of the natural environment of the N'ZI watershed is regressive between the three dates. Looking at land use maps, it appears that this degradation appears to be general and affects almost the entire basin. The overall rates of change over the 1986-2000, 2000-2020, and 1986-2020 series show regressions for the forest, savannah, plantation-agroforestry, and fallow and dwelling-bare land classes. Natural formations (forests, savannas) are increasingly replaced by anthropogenic occupation classes (fields and fallows, plantations and agroforestry, dwellings-bare ground). The same finding is made by (Mama et al., 2013) on the territory of Benin and Togo (Koumoi et al., 2013; Kpedenou et al., 2017, 2016a). This situation of degradation of the natural landscape in the N'ZI watershed is also noted by Noho et al. (2018) and in the department of Katiola by Agoualé et al.

(2017), and in other parts of Côte d'Ivoire by studies by (Aké et al., 2012; Alexis et al., 2018). Environmental degradation, reflected in the regression of natural formations, is essentially linked to the development of socio-economic activities (e.g., slash-and-burn, timber and fuelwood, mining); as observed during the field missions (Figure 11). More significantly, there is a regression of vegetation formations and an increase in the area of anthropized landscapes in this basin. The main factors in the degradation of the vegetation cover are agriculture, the remarkable increase in the population of the N'ZI region between 2010 and 2014 (INS, 2014), and late wildfires that destroy everything in their path (granaries, plantations, and various natural plant formations). Results of the work by **Kouassi (2019)** on the impact of bushfires in the N'ZI watershed confirm our results. These same observations were made in the studies of several authors on peri-urban occupations in the world (Salomon et al., 2021, Useni et al., 2020, Aguejdad and Hubert-Moy., 2016, André et al., 2014). Charcoal production and crushing also contribute to soil and vegetation degradation. Savannas burned and left in the same state remain in the same places while increasing their surface area following the degradation of certain fallow areas. In addition to wildfires, livestock farming appears to be an activity that increases land degradation in the region (Figure 11). The study area is characterized by overgrazing aggravated by internal transhumance. Indeed, overgrazing leads to a reduction in the natural regeneration of woody plants, a decrease in herbaceous cover, and the stripping of the soil and its hardening according to Agoualé et al. (2017).




Figure 11 Illustrative photos attesting to the anthropized of the N'ZI watershed (field mission: 18; 26 and 27 /12/2020: a) Bouaké; b) M'bahiakro c) Fétékro; d) Bocanda; e) Katiola, and f) Dabakala)

Conclusion

This study shows that the N'ZI watershed has rapidly changed the environment. The discrimination of the regions of interest allowed to define six (6) groups of land use (Forests; Savannas, Soils-low-cover; Plantations- agroforestry and fallow; Dwellings-bare soils; Water body) by the treatment of satellite images multi-dates (1986, 2000 and 2020). The confusion matrix of different dates with Kappa values increases slightly between 1986, 2000, and 2020 respectively by 0.86; 0.90, and 0.92 testify to the classification of images was well done, when discriminating regions of interest. The detection of change using the area calculation, the overall rates of change, and the evolution statistics of the different class units indicate a degradation of the forest, plantation, and fallow classes, of the water areas in general. On the other hand, there are an increase in bare land (67.02%) and agglomerations (103.35%) to the detriment of forests (-80.45%) and water bodies (-52.47%). A regression of approximately (-14.87%) between 2000 and 2020 of plantations-agroforestry and fallow land to the advantage of savannas (89.98%). In the end, the basin is undergoing a significant change in all categories of its landscape. The most affected classes are groups of forests and water bodies that are renewed according to the rainfall fluctuations of the area.

Acknowledgments: The authors sincerely and gratefully acknowledge, all, the contributors and colleagues who helped improve this manuscript. Too at Tivoli Sandrine, YAO Assemian, Assoumou Aka, and Moussa Camara for assistance in the preparation of the manuscript, and Deguy Attoungbré Jean Philippe for helping in the image processing.

References

- Achbun, A., Mansour, M., Layelmam, M., Smiej, M., 2011. Comparative study of the object-oriented classification of a spot5 image for land cover mapping via Cognition 9. GEO OBSERVER, 2011, pp.13. *HAL Id: hal-00915067: <https://hal.archives-ouvertes.fr/hal-00915067>*.
- Agouale Y.J,Vei K.N, Afro M.J., 2017, Anthropogenic pressures and plant cover dynamics in the Sub-prefecture of Katiola, PhD student Alassane Ouattara University (Bouaké, Côte d'Ivoire), *Revue Ivoirienne de Géographie des Savanes*, **Numéro 1 Janvier 2017, ISSN 2521-2125**. *<https://riges-uao.net/volumes/volume1/fichiers/art6.pdf>*.
- Aguejdad, R., Hubert-Moy, L., 2016. Monitoring of the artificialization of the territory in urban areas by remote sensing and using landscape metrics. Application to a medium-sized town, Rennes Metropole, *Cybergeo: European Journal of Geography*. *<https://doi.org/10.4000/cybergeo.27465>*.

- Aké, G.É., Kouadio, B.H., Adja, M.G., Ettien, J.-B., Effebi, K.R., Biémi, J., 2012. Mapping of the multifactorial vulnerability to water erosion of soils in the Bonoua region (southeast of Côte d'Ivoire). *Physio-Geo. Physical geography and environment* p1–42. <https://doi.org/10.4000/physio-geo.2285>.
- Alexis, N., Kouadi, A., Deguy, J., Hien, A., Goula, A., Savané, I., 2018. Influence of land use dynamics on the amount of land loss south of the Sassandra watershed (Cote d'Ivoire). *International Journal of Advanced Research* 6, 830–838. <https://doi.org/10.21474/IJAR01/6918>.
- André Marie ⁽¹⁾, Mahy Grégory ⁽¹⁾, Lejeune Philippe ⁽²⁾, Bogaert Jan ⁽¹⁾, 2014. Towards a synthesis of the design and definition of areas in the urban-rural gradient. *Biotechnology, Agronomy, Society and Environment* 18(1), 61–74. <https://orbi.uliege.be/bitstream/2268/161968>.
- Assoma, V.T., Yao, N.A., Dio, J.S., Jourda, J.-P., 2021. Contribution of remote sensing and GIS to the mapping of land use changes in the Lobo watershed in Côte d'Ivoire. *Physio-Geo. Physical geography and environment*. 107–126. <https://doi.org/10.4000/physio-geo.12654>.
- Avenard, J.M., Eldln, M., Girard, G., Sircoulon, J., Touchebeuf, P., Guillaumet, J.L., Adjanohoun, E. and Perraud, A. (1971) *The natural environment of the Ivory Coast. Memoirs ORSTOM*, 50, 339. <https://www.documentation.ird.fr › hor › fdi:16368>.**
- BNETD, E., Tera, R., 2016. Qualitative analysis of the factors of deforestation and forest degradation in Côte d'Ivoire. UN-REDD Programme, Abidjan. https://www.nitidae.org/files/b24e760c/161216081210_161214.
- Brou, T., Y., O., Bigot, S., Servat, E., 2005. Deforestation risks in the permanent domain of the State in Côte d'Ivoire: what future for its last forests? *Remote sensing*, 5 (1-2-3), pp. 263-275. ISSN 1028-7736. <https://www.documentation.ird.fr › hor › fdi:010047889>.
- Caloz, R., Pointet, A., and Collet, C., 2003. A comparative approach to aerial image classification methods: a case study, *Geogr. Helv.* 58, 141-153. <https://doi.org/10.5194/gh-58-141-2003>, 2003.
- Congalton, R.G., 1991. A review of assessing the accuracy of classifications of remotely sensed data. *Remote Sensing of Environment* 37, 35–46. [https://doi.org/10.1016/0034-4257\(91\)90048-B](https://doi.org/10.1016/0034-4257(91)90048-B).
- FAO, 1996. *Rome Declaration on World Food Security and World Food Summit Plan of Action: World Food Summit, 13-17 November 1996, Rome, Italy* 43p.** <https://www.fao.org/3/w3548e/w3548e00.htm>.
- INS, 2014. Mapping non-financial poverty in Abidjan District from the 2014 General Population and Housing Census of Abidjan Côte d'Ivoire 40. <https://www.ins.ci › RGPH 2014 expo dg>.

- Diallo Boubacar Amadou, Bao Zhengyu., 2010. Land cover change assessment using remote sensing: a case study of Bamako, Mali., 2010;2(4). http://www.sciencepub.net/researcher/research0204/02_2572_research0204_7_17_01.**
- Diallo, Hady, Bamba, I., Barima, Y.S.S., Visser, M., Ballo, A., Mama, A., Vranken, I., Maiga, M., Bogaert, J., 2011. Combined effects of climate and anthropogenic pressures on the evolutionary dynamics of vegetation in a protected area of Mali (Fina Reserve, Baoulé Loop). *Science and Global Change/ Drought* 22, 97–107. <https://doi.org/10.1684/sec.2011.0306>.
- Dietzel, C., Oguz, H., Hemphill, J.J., Clarke, K.C., Gazulis, N., 2005. Diffusion and coalescence of the Houston metropolitan area: evidence supporting a new urban theory. *Environ Plann B Plann Des* 32, 231–246. <https://doi.org/10.1068/b31148>
- Desclée, B., Bogaert, P., & Defourny, P. (2006). Forest change detection by statistical object-based method. Remote sensing of environment, 102(1-2), 1-11. <https://doi.org/10.1016/j.rse.2006.01.013>.**
- Girard, M.-C., and Girard, C.-M. 1999. Processing remote sensing data. Dunod Ed. Paris, 529 p ».** http://www.banque-pdf.fr/fr_girard-m-cgirard-c-m-1999-traitements-des-donnees-deleledetection.html.
- Chaima,G, Mghirbi, O., Le Grusse, P., Louvet, S., 2019. Spatial characterization of natural resource vulnerability and management of diffuse pollution risks related to agricultural phytosanitary practice. in GFP (, F. (Ed.), 49. French Pesticide Group Congress: Pesticides and Agroecological Transition: Challenges and Advances. GFP Editions, Montpellier, France, pp. 103–105.<https://doi.org/10.13140/RG.2.2.16696.80640>.**
- Gbombélé SORO^{1*}, Ernest Kouassi AHOUSSE¹, Emmanuel Konan KOUADIO¹, Tanina Drissa SORO¹, Sékouba OULARE², Mahaman Bachir SALEY^{1,2}, Nagnin SORO¹ et Jean BIEMI^{1,2}, 2014. Contribution of remote sensing to the mapping of the spatio-temporal evolution of the dynamics of land use in the Lacs region (Central Côte d'Ivoire), Afrique SCIENCE 10(3) (2014) 146 - 160 146 ISSN 1813-548X, <http://www.afriquescience.info>.**
- Koné, K.H., Konan, K.E., Bi, Z.K., 2021. Characterization of the vegetation of the Comoé National Park (north-east of Cote d'Ivoire) 14. <http://www.revue-geotropes.com> › 2-kiyofolo-al.
- Kouassi, A.M., 2010. Characterization of a possible change in the rain-flow relationship in West Africa: the case of the N'zi (Bandama) watershed in Côte d'Ivoire, PhD thesis, Cocody-Abidjan University, Côte d'Ivoire, 210 p. *Physio-Géo. Physical geography and environment* 3–4. <https://doi.org/10.4000/physio-geo.1163>.

- Kouassi, J.-L., 2019. Climate variability, fire dynamics, and local perceptions in the N'Zi watershed (Central Ivory Coast) (Ph.D. thesis). National Polytechnic Institute Felix Houphoët Boigny - Yamoussoukro., 250p. <https://tel.archives-ouvertes.fr/tel-02183044>.
- Koudou, A., N'dri, B.E., Niamke, K.H., Adjiri, O.A., Sombo, A.P., Niangoran, K.C., 2016. Analysis of productivity and estimation of vulnerability to pollution for the optimization of the implantation of drilling in the basement aquifers of the N'zi watershed (Côte d'Ivoire). Bulletin de l'Institut Scientifique, Rabat, Section Sciences de la Terre 149–164. <http://www.israbat.ac.ma › uploads › 2017/07>.
- Koumoui, Z., Alassane, A., Djangbedja, M., Boukpepsi, T., Kouya, A.-E., 2013. Spatial and temporal dynamics of land use in central Togo. Ahoho - Revue de Géographie du LARDYMES 7, 163–172. <https://hal.archives-ouvertes.fr/hal-01564256>.
- Kpedenou, K.D., Ahe, P., Boukpepsi, T., 2016a. Spatio-temporal dynamics of land use/occupation in south-east Togo: mapping test using spatial tecton. Ahoho 131–143. <https://hal.archives-ouvertes.fr/hal-01564256>.
- Kpedenou, K.D., Boukpepsi, T., Tchamie, T.T.K., 2016b. Quantification of land use changes in the Yoto prefecture (southeast Togo) using Landsat satellite imagery. Journal of Environmental Sciences. University of Lomé, 2016, pp.137-156. [ffhal-01409418. https://hal.archives-ouvertes.fr/hal-01409418](https://hal.archives-ouvertes.fr/hal-01409418).
- Kpedenou, K.D., Drabo, O., Ouoba, A.P., Da, D.C.E., Tchamie, T.T.K., 2017. Analysis of land use for monitoring the evolution of the landscape of the Ouatchi territory in south-east Togo between 1958 and 2015. Presses of the University of Ouagadougou, 2017, XXXI (55), pp.203-228. [ffhal-01701176, https://hal.archives-ouvertes.fr/hal-01701176](https://hal.archives-ouvertes.fr/hal-01701176).
- Landis, J.R., Koch, G.G., 1977. The Measurement of Observer Agreement for Categorical Data. Biometrics 33, 159–174. <https://doi.org/10.2307/2529310>
- Lienou, M.L., 2009. Machine learning of land use classes and visual representation of satellite images (Ph.D. Thesis). Telecom ParisTech.191p. [⟨pastel-00005585⟩](https://pastel-00005585)
- Mama, A., Sinsin, B., De Cannière, C., Bogaert, J., 2013. Anthropization and landscape dynamics in the Sudanese zone in northern Benin. Tropicultura 31. <http://www.tropicultura.org/text/v31n1/78>.
- Mellor, A., Haywood, A., Stone, C., & Jones, S. (2013). The performance of random forests in an operational setting for large area sclerophyll forest classification. Remote Sensing, 5(6), 2838–2856. <https://doi.org/10.3390/rs5062838>.
- Monnet Claude. (1972). Contribution à l'étude de la dynamique de la nature des suspensions d'un fleuve intertropical, le Bandama, Côte d'Ivoire : évolution des éléments chimiques des eaux**

de son estuaire. Paris: ORSTOM, 431 p. multigr. Th. Sci. Nat.: Nice: no AO7826: 1972/12/21; fdi:06019.

- N'Guessan, F.K., 2020. spatial dynamics of land use in the bembela finage (northeast of the ivory coast) 13. <http://www.revue-geotrope.com> › 20200601 › 8.
- Noho Y.A.W., N'guessan K.F., Koli B.Z 2018, Characterization of the vegetation of the N'zi catchment area (ivory coast), (EDUCI) 2018 Journal of Tropical Geography and Environment, n°1, 2018. <http://www.revue-geotrope.com> › 20180919 › 3.
- Zeng Ju¹, Mou Leong Tan¹, Narimah Samat¹, Chun Kiat Chang² Comparison of Landsat 8, Sentinel-2 and spectral indices combinations for Google Earth Engine-based land use mapping in the Johor River Basin, Malaysia. GEOGRAFIA Online™ Malaysian Journal of Society and Space 17 issue 3 (30-46) © 2021, e-ISSN 2682-7727 <https://doi.org/10.17576/geo-2021-1703-03>.**
- Pelletier, C., 2017. Mapping land use from time series of high-resolution satellite images: identification and processing of mislabelled data (Ph.D. Thesis)., University of Toulouse III-Paul Sabatier. <https://tel.archives-ouvertes.fr/tel-01665121>.
- Pontius Jr, R.G., 2000. Comparison of categorical maps. Photogramm. Eng. Remote Sens 66, 1011–1016. <https://www.asprs.org> › 2000_aug_1011-1016.
- Salomon, W., Sikuzani, Y.U., Kouakou, A.T.M., Barima, Y.S.S., Theodat, J.M., Bogaert, J., 2021. Mapping and quantification of forest cover loss in the national parks of the Republic of Haiti from 1985 to 2018: role of local agricultural activities. International Journal of Biological and Chemical Sciences 15, 477–496. [DOI: <https://dx.doi.org/10.4314/ijbcs.v15i2.9>].
- Skupinski, G., Tran, D.-B., Weber, C., 2009. Satellite images Spot multi-dates and spatial metrics in the study of urban and suburban change – The case of the lower Bruche valley (Bas-Rhin, France). CyberGeo: European Journal of Geography. <https://doi.org/10.4000/cybergeogeo.21995>.
- Tra Bi, Z.A., 2013. Study of the impact of anthropogenic activities and climate variability on vegetation and land use, by using remote sensing and agricultural statistics, on the Bouregreg watershed (MAROC). (Ph.D. doctorate). Artois. 1992p., <http://www.theses.fr/2013ARTO0008/document> <https://www.sudoc.fr/184252598>
- Useni Sikuzani, Y., Boisson, S., Cabala Kaleba, S., Nkuku Khonde, C., Malaisse, F., Halleux, J.-M., Bogaert, J., Munyemba Kankumbi, F., 2020. Land use dynamics around mining sites along the urban-rural gradient of the city of Lubumbashi, DR Congo. Biotechnology, Agronomy, Society, and Environment. DOI : 10.25518/1780-4507.18306.