

Effect of Inorganic Fertilizer Application on Green House Gas Emissions and Microbial Activity under Coffee Agroforestry in Eastern Uganda

Esther Sebuliba¹, Majaliwa Mwanjalolo², Prossy Isubikalu¹, Nelson Turyahabwe¹, Gerald Eilu¹, and Adipala Ekwamu²

¹Makerere University, College of Agricultural and Environmental Sciences

²Regional Universities Forum for Capacity Building in Higher Education

December 8, 2022

Abstract

A study was conducted in none tilled coffee agroforestry fields of Eastern Uganda to understand the effects of application of inorganic fertilizers on soil nutrient loss in form of gas for mitigation of unsustainable agricultural practices. This study specifically i) assessed the effect of application of inorganic fertilizers on greenhouse gas emissions, ii) determined their effect on microbial carbon, nitrogen and phosphorus and iii) determined their effect on leaf litter decomposition under Albizzia-coffee growing systems of the Mount Elgon. Soil gas emissions were measured with the static chamber method for twelve months in a field experiment with five different fertilizer treatments. The effect of treatments was separated using ANOVA in Genstat discovery version 13. Microbial carbon, nitrogen and phosphorus was separated using Mann-Whitney U test. Results showed that annual emissions ranged from 19.6 to 26.1 (t C/ha/yr), 3.5 to 9 (Kg N/ha/yr) and 6.9 to 9.2 (Kg C/ha/yr) for carbon dioxide, nitrous oxide and methane respectively. Significant effects on soil emissions only occurred for nitrous oxide ($P=0.017$), microbial carbon ($p=0.001$) and microbial phosphorus ($p<0.001$) for the study period. The mixture of NPK fertilizers presented the lowest carbon dioxide loss and application of TSP presented the lowest nitrous oxide emission from soil. This study underscores the need for establishment of long-term experiments across several agro-ecological zones to confirm farmers' perceptions of their soil fertility levels and ascertain the contribution of farm practices towards the retention of nutrients in the soil with minimal emission, to inform decisions of small holder farmers, policy and development partners for sustainable production.

Hosted file

Annual GHG emissions.docx available at <https://authorea.com/users/564643/articles/611572-effect-of-inorganic-fertilizer-application-on-green-house-gas-emissions-and-microbial-activity-under-coffee-agroforestry-in-eastern-uganda>

Hosted file

Study Area SSSA.docx available at <https://authorea.com/users/564643/articles/611572-effect-of-inorganic-fertilizer-application-on-green-house-gas-emissions-and-microbial-activity-under-coffee-agroforestry-in-eastern-uganda>

Hosted file

Monthly GHG emissions.docx available at <https://authorea.com/users/564643/articles/611572-effect-of-inorganic-fertilizer-application-on-green-house-gas-emissions-and-microbial-activity-under-coffee-agroforestry-in-eastern-uganda>

Hosted file

seasonal GHG emissions.docx available at <https://authorea.com/users/564643/articles/611572-effect-of-inorganic-fertilizer-application-on-green-house-gas-emissions-and-microbial-activity-under-coffee-agroforestry-in-eastern-uganda>

Hosted file

essoar.10512959.1.docx available at <https://authorea.com/users/564643/articles/611572-effect-of-inorganic-fertilizer-application-on-green-house-gas-emissions-and-microbial-activity-under-coffee-agroforestry-in-eastern-uganda>

Effect of Inorganic Fertilizer Application on Green House Gas Emissions and Microbial Activity under Coffee Agroforestry in Eastern Uganda

- The application of inorganic fertilizers increased annual soil nitrogen losses in form of soil N₂O emissions.
- Peaks of N₂O emissions were observed immediately after application of inorganic fertilizers but levelled afterwards.
- The application of inorganic fertilizers increased microbial carbon and microbial phosphorus in soil after one year.
- The application of inorganic fertilizers did not affect leaf litter decomposition rates.
- The mixture of NPK fertilizers released the lowest nutrients (CO₂ emissions) and TSP application released the lowest nitrogen (N₂O emissions) from soil.

Effect of Inorganic Fertilizer Application on Green House Gas Emissions and Microbial Activity Under Coffee Agroforestry in Eastern Uganda

Abstract

A study was conducted in none tilled coffee agroforestry fields of Eastern Uganda to understand the effects of application of inorganic fertilizers on soil nutrient loss in form of gas for mitigation of unsustainable agricultural practices. This study specifically i) assessed the effect of application of inorganic fertilizers on greenhouse gas emissions, ii) determined their effect on microbial carbon, nitrogen and phosphorus and iii) determined their effect on leaf litter decomposition under *Albizia*-coffee growing systems of the Mount Elgon. Soil gas emissions were measured with the static chamber method for twelve months in a field experiment with five different fertilizer treatments. The effect of treatments was separated using ANOVA in Genstat discovery version 13. Microbial carbon, nitrogen and phosphorus was separated using Mann-Whitney U test. Results showed that annual emissions ranged from 19.6 to 26.1 (t C/ha/yr), 3.5 to 9 (Kg N/ha/yr) and 6.9 to 9.2 (Kg C/ha/yr) for carbon dioxide, nitrous oxide and methane respectively. Significant effects on soil emissions only occurred for nitrous oxide (P=0.017), microbial carbon (p=0.001) and microbial phosphorus (p<0.001) for the study period. The mixture of NPK fertilizers presented the lowest carbon dioxide loss and application of TSP presented the lowest nitrous oxide emission from soil. This study underscores the need for establishment of long-term experiments across several agro-ecological zones to confirm farmers' perceptions of their soil fertility levels and ascertain the contribution of farm practices towards the retention of nutrients in the soil with minimal emission, to inform decisions of small holder farmers, policy and development partners for sustainable production.

Key words: Inorganic fertilizers, agroforestry, carbon dioxide, methane, nitrous oxide, *Coffea arabica*, East African highlands

Introduction

The need to increase agricultural production to satisfy the demand associated with population growth has resulted into horizontal expansion of cultivation even on marginal lands in SSA (Bekunda and Woomer, 1996; Kopittke et al., 2019). Subsequently, greenhouse gases (GHG) emission (Rosenzweig et al., 2020) and other forms of environmental degradation (FAO, 2002) including soil fertility decline (Kopittke et al., 2019) have increased. In the effort to boost soil productivity, in order to feed the growing population, various soil fertility management practices have been promoted across Africa including fertilization (Liniger et al., 2011). However, this is likely to increase the greenhouse gas emissions (GHG) from agricultural land in SSA (Nisbet et al., 2014; Musafiri et al., 2020). It is therefore imperative to identify soil fertility management technologies that increase food production and play a fundamental role in GHG fluxes control since only a few integrated studies have tried to quantify gas released or to characterize the mechanisms involved in their release.

Carbon dioxide is the most important GHG, however, methane (CH_4) and nitrous oxide (N_2O) emissions also play a substantial role in global warming (Smith et al., 2018). The net soil CO_2 emissions are produced by soil respiration (aerobic and anaerobic microbial and roots respiration) and decomposition of organic matter (Oorts et al., 2007; Sun et al., 2019; Almagro et al., 2021). Net methane fluxes are the balance between two contending microbial processes that are methane production by methanogens, and methane oxidation by methanotrophs under anaerobic and aerobic conditions, respectively (Knief, 2019). Net soil N_2O fluxes occur as a result of heterotrophic and autotrophic nitrification, chemo-denitrification, nitrifier- denitrification, and co-denitrification (Gander et al., 2012; Butterbach-Bahl et al., 2013). The strength of these GHG emissions is affected by soil properties (soil organic carbon, soil nitrogen, texture, pH), land cover changes, vegetation type, environmental factors (soil temperature, soil moisture, drought, precipitation) and farm management practices (crop residue application, tillage, manure, agroforestry, fertilizer use) (Luo et al. 2010; Powlson et al. 2011; Tongwane et al., 2016; Pelster et al., 2017; Wang et al., 2019; Dimitriou et al. 2021). Soil organic carbon (SOC) accumulation generally occurs in areas of low decomposition that thrives under low temperature, acid parent materials and anaerobic conditions. It enhances both unstable and stable macro aggregate formation (Denef et al., 2013) vital for carbon sequestration (Six et al, 2004; Plante & McGill, 2002).

Also, increasing soil N content generally leads to higher soil respiration facilitating higher net ecosystem exchange, if carbon is not limiting (Niu et al., 2010; Peng et al., 2011). According to Pilegaard et al. (2006), N_2O emissions are negatively correlated with the C/N-ratio (with N_2O emissions being lowest at C/N-ratios 30 and highest at a C/N-value of 11 (Christiansen et al., 2012). Acidic soil conditions lead to lower soil emissions, with an optimal pH-value for methanogenesis (CH_4 production) that lies between pH 4 and 7 (Dalal and Allen, 2008). Carbon dioxide emissions are observed to be highest at neutral pH-

values (Čuhel et al., 2011). Methane emissions decrease only under acidic soil conditions while nitrification increases with higher pH-values, since the equilibrium between NH_3 and NO_3 shifts to ammonia (Nugroho et al., 2007). However, no significant correlations are found between N_2O emissions and pH-value (Pilegaard et al., 2006). Saiz et al. (2006) found that young trees have higher respiration compared to old ones. Soil respiration decreases with stand age, caused by a lower fine root biomass. They added that in old forest ecosystems, the decrease levels out with stand age since lower root respiration rates are partly compensated for by higher microbial respiration due to higher organic inputs.

An increase of soil temperature leads to higher emissions and to higher soil respiration rates as a positive feedback response of increased microbial metabolism. Soil temperature and soil moisture explain 86% of the variations of N_2O emissions (Schindlbacher et al., 2004). Methane emissions are additionally forced by increasing soil respiration rates with increasing soil temperatures, leading to decreasing O_2 concentrations in the soil (Butterbach-Bahl et al., 2013). The positive temperature effect may be overlain by soil water stress, since water is needed as a transport medium for nutrients required by microbes (Fowler et al., 2013). CO_2 emissions increase exponentially with temperature (Ludwig et al., 2001; Tang et al., 2003). Soil moisture is the single most important soil parameter for soil gas emissions, since it controls microbial activity and all related processes. Nitrifying bacteria require oxygen residing in soil pores. Therefore, soils with less water-filled pore space (WFPS) have higher emissions by nitrification, with a maximum at 20% WFPS (Ludwig et al., 2001). Nitrification yields a higher potential for NO production than for N_2O production (Fowler et al., 2013). In contrast, CH_4 and N_2O producing bacteria require anaerobic conditions. N_2O production is optimal around 60% WFPS and lowest when WFPS is below 30% (Gao et al., 2014). Even an increase of WFPS above 80% can still lead to an exponential increase of N_2O emissions (Keller and Reinert, 1994). CH_4 production requires strictly anaerobic conditions and correlates positively with soil moisture (Gao et al., 2014; Smith et al., 2003). Long periods of drought can significantly reduce soil emissions and soils may then turn into a net sink for N_2O (Goldberg and Gebauer, 2009). Soils with a high proportion of large pores retain less water and therefore foster the emission of gases produced under aerobic conditions (van der Weerden et al., 2010). Soils with dominant fine pores support the formation of CH_4 and N_2O produced under anaerobic conditions (Dutaur and Verchot, 2007; Gu et al., 2013).

Soil texture and structure also influence GHGs indirectly through soil moisture. Higher CO_2 emissions are encountered with fine textured soils, especially compared to sandy soils during warm dry periods (Dilustro et al., 2005). Stable soil aggregates (concretions, crusts) lead to lower soil emissions since C and N are less available for soil microbes (Wu et al., 2012). Precipitation after extended dry periods causes the pulsing or Birch effect (Birch, 1958). Emissions increase within some minutes or hours after the onset of precipitation and return to background levels within a few days (Sponseller, 2007; Lado-Monserrat et al.,

2014). This is driven by the renewed mineralization and the availability of easily decomposable material (Borken and Matzner, 2009) for the metabolism of reactivated microbes (Ludwig et al., 2001). The Birch effect decreases with higher frequencies of wet-dry cycles (Borken and Matzner, 2009).

Also, tillage practices influence particulate organic matter fraction (Hussain et al., 1999; Liu et al., 2014). Generally, the rate of SOC storage under no tillage is relatively higher compared to conventional tillage (Johnson et al., 2005), particularly for the top soil. Several authors (Curtin et al., 2000; Al-Kaisi and Yin, 2005; Bauer et al., 2006; Ussiri and Lal, 2009) have reported higher soil CO₂ emissions under conventional tillage compared to no-tillage. This is because no-tillage reduces the diffusion and content of air-filled pores in the soil, by which soil CO₂ emissions are very low or non-existing (Bilandzija et al., 2016).

Agroforestry systems have similarly been seen as one of the promising management practices to increase soil C stocks, reduce soil degradation and mitigate greenhouse gas emissions (e.g., Frouz et al., 2013; Ehrenbergerová et al., 2016; Dollinger and Jose, 2018; Justine et al., 2019; Solis et al., 2020). Nitrogen fixing leguminous trees (such as *Albizzia sp.*, *Inga sp* and *Erythrina sp*) have been commonly used to bring N and organic matter to the system in addition to other benefits (Vaast et al., 2008; Verchot et al., 2008). However, suspicion on N fixing leguminous species to increase soil N₂O emissions (Rochette and Janzen 2005; Verchot et al., 2008) and reduce the soil CH₄ sink (Palm et al., 2002) is a growing concern in the sustainable development framework. For example, studies such as Verchot et al. (2008) and Hergoulouch et al. (2008) showed shaded coffee increased N₂O emissions by 34.8% compared to coffee monocrop. Such conflicting results compel us to question the response of soil microbial activity and production of GHGs to fertilization, since little is known about the how the type of the fertilizers applied affect soil emissions of CH₄, CO₂, and N₂O and the decomposition rate in agricultural systems (Amos et al., 2005; Mosier et al., 2006; Sainju et al., 2008).

Various scholars (e.g. Bouwman et al., 2002a&b; Phillips, 2007; Phillips et al., 2009) further found that the effects of fertilization on GHG fluxes at the soil surface tend to occur within the initial 8 to 10 weeks following N application. Since N₂O and CO₂ emissions tend to increase and CH₄ uptake tends to decline during the first few weeks following fertilization, it is generally accepted that GHG emissions tend to increase with additions of N (Bouwman et al., 2002; Mosier et al., 2006; Sainju et al., 2008). Also, the N-fertilization tends to increase CO₂ emissions (Raposo et al., 2020). Effects of Fertilizer N, P and K addition on fluxes of GHGs, however, are not consistent across studies. Sometimes fertilization of arable soil does not affect the strength of soil as a source of N₂O (Amos et al., 2005) and CO₂ (Amos et al., 2005) or the strength of the soil as a sink for CH₄ (Koga et al., 2017). CO₂ released from soil to the atmosphere, referred to as soil respiration, is a combined activity of roots, micro and macro organisms decomposing litter and organic matter in soil (heterotrophic respiration) (Högberg et al., 2020; Hanson et al., 2003) and is influenced by tem-

perature. Nevertheless, inorganic farming is reported to sequester less carbon than organic farming (Gattinger et al., 2012).

This information is very important particularly in the coffee-based farming systems where farmers have resorted to increased application of inorganic fertilizers to enhance coffee production. Coffee being one of the highest contributor to GDP in Uganda (De Beenhouwer et al., 2016), it is important to identify best management practices which sustain its production and protect soil, water and air quality under fertilized shaded conditions. This study therefore i) assessed the effect of inorganic fertilizer application on greenhouse gas emission, ii) determined the effect of fertilizer application on microbial CNP and iii) determined the effect of leaf litter decomposition under *Albizia*-coffee growing systems of the Mount Elgon.

Materials and Methods

Header 2

Description of the study area

This study was conducted in Manafwa District, on Mount Elgon of Eastern Uganda. Mount Elgon lies on the border of eastern Uganda and western Kenya. Manafwa district is bordered by Bududa District to the north, Kenya to the east and south, Tororo District to the south-west, and Mbale District to the west (Figure 1). It has an elevation of 1,354 metres.




Figure 1: The study site around Mount Elgon, in Eastern Uganda showing Manafwa and neighboring districts

The landscape of Manafwa mainly constitutes smallholder farms (< 4 acres) with intensive and mixed coffee (*C. arabica*) agricultural systems and is characterized by a relatively high population density of approximately 250–300 inhabitants per km² (Gram *et al.*, 2018). Coffee productivity in this district has been reported to have greatly declined below its potential due to low soil fertility and poor land and coffee tree management practices (Wang *et al.*, 2015). The geographical location, climate conditions, topography, geology, soils, vegetation and population are well described in Sebuliba *et al.* (2021).

Manafwa district was selected because it offered adequate land for setting up the experiment under a homogenous environment (similar slope, vegetation, climate, age of coffee) and with minimum likelihood of mud or landslides events occurring on the site. The farmers in the identified experimental site were willing to host the experiment.

Experimental design, treatment and replication

The magnitude of the effect of application of inorganic fertilizers on actual fertility of soil followed soil fertility measurement in two ways; 1) by what the farmers were saying; and 2) the experiment to understand and verify farmers' perceptions of the levels of fertility for their soil. The on-farm experiment was established on three independent farmer fields to ascertain the ability of soil to retain nutrients or facilitate escape of nutrients for coffee use through the inorganic fertilizer application. The loss of fertility of the soil was measured from the magnitude of gases lost from the soil in order to identify which treatment contributed more to rejuvenating the soil by minimizing nutrient loss by gas. Gas emission was inversely proportional to soil fertility; in that the higher the gas emitted, the lower the fertility of soil.

The experimental design was a completely randomized since coffee farmer's fields were relatively similar and showing more uniform characteristics. The assumptions were that; there was either no or minimal use of inorganic fertilizers, trees ages were above 5 years even up to 20 years, *Albizzia coriaria* (commonly known as mugavu) dominated the coffee fields, they had a similar soil type (ferralsol), same gentle slope, same climate (between 1000 to 1300mm of rain) within Manafwa. The experiment was carried out in Manafwa because its farmers rarely apply inorganic fertilizers in their coffee fields, with a moderate altitude resulting into moderate fertility levels and drier requiring rejuvenation from the application of inorganic fertilizers.

Four treatments (50P, 250N, 250K, 250N-50P-250K) $\text{Kg ha}^{-1} \text{ yr}^{-1}$ and a control) were considered and replicated four times. A total of 20 field plots, of 20m-by-20m size, was laid out randomly on the experimental farmer sites. Plots were located on same landscape positions, soil type, slope and within 300m radius from each other to ensure homogeneity of the experimental site. The properties of the experimental site in terms of texture (sandy loam), pH, soil organic matter (SOM), soil organic carbon (SOC), Total nitrogen (TN), Available phosphorus (Av. P), extractable potassium (K) and sodium (Na) are shown in Table 1. Urea, Tri-Super phosphate (TSP) and Muriate of Potash (MOP), NH_4NO_3 and NaH_2PO_4 solutions were annually applied to each plot at the beginning (August) of the short rain season.

Table 1: Initial selected soil characteristics of the farmer fields used for the experiment (N=3)

Soil depth (cm)	pH	N	Av P	K	Na	SOC	SOM	Textural class
	pH units	%	mg/kg	cmol/kg	cmol/kg	%	%	
0-15	6.42	0.19	13.86	0.68	0.10	1.90	3.27	Sandy loam
15-30	6.36	0.13	9.12	0.53	0.11	1.41	2.42	Sandy loam

Each of the experimental plots was sub-divided into four cells/grids (10×10 m). Each cell was also considered as a replicate and permanent chamber bases were installed. Chamber covers were placed on the bases and gas samples were collected 4 times at (10) minute intervals for determination of the trace soil gas fluxes (Collier *et al.*, 2014). Soil gas fluxes (CO_2 , CH_4 and N_2O) were measured monthly for twelve months.

The samples were stored in pre-evacuated glass containers with Teflon-coated stopcocks and taken to University of Gottingen for CO_2 , CH_4 and N_2O concentrations determination. Samples were analyzed using a gas chromatograph (Shimadzu GC-14B; Columbia, MD, USA) equipped with a flame ionization detector (FID), an electron capture detector (ECD) and an auto sampler (Koehler *et al.*, 2009, 2012; Corre *et al.*, 2014). The detection limits of this instrument were 50 ppm for CO_2 , 43 ppb for N_2O and 45 ppb for CH_4 . Annual and seasonal soil CO_2 , CH_4 and N_2O fluxes were calculated as a sum of the twelve-monthly fluxes and the respective monthly fluxes for each gas; respectively. Relative change to the control for monthly fluxes was also computed using;

$$R = (F_k - F_c) \frac{100}{F_c}$$

where R is the relative change of the flux in the given month for the treatment K, F_k is the flux of a given month for the treatment K, F_c is the flux for the same month for the control.

Determination of leaf litter decomposition rate

Leaf litter of *Albizia coriaria* was collected using the litterbag trapping method by placing traps on the floor of each plot at the beginning of MAM 2020. About 40 to 60g of collected leaf litter was placed in litter bags and buried in the soil at 10cm depth (Zhou *et al.*, 2008). Each plot received about 10 litter bags evenly distributed. Every three months, a litterbag was collected randomly, gently cleaned and leaf litter gently weighed using adhoc methods as described by Okalebo *et al.* (2002).

Incubation

A 72 hours' soils incubation was carried out in 50 ml plastic beakers with drainage holes at the bottom, lined with a glass filter and filled with field-moist soil (approximately 6 g dry weight) to allow inoculation (Okalebo *et al.*, 2002). A 16° C growth chamber containing 1 L glass Mason Ball jars was used for storage and transportation of soil to the laboratory under field capacity conditions with sporadic additions of de-ionized water (Davis *et al.*, 2005).

Extractable and Microbial C, N, P

The chloroform fumigation–extraction method (CFEM) on sieved, undried splits of each sample was considered in measuring microbial C, N and P within a week

of collection. A darkened vacuum desiccator fitted with fumigated soils was exposed to chloroform for 36–48h. Extraction of approximately 6 g fumigated or unfumigated (control) soil was done with Brays solution for extractable P (Bray and Kurtz, 1945), using 50 ml or 75 ml extractant while 0.5 M K_2SO_4 was used for C and N (Brookes *et al.*, 1985; Vance *et al.*, 1987; Beck *et al.*, 1997). Beforehand filtering, solutions of soil were shaken for 1h followed by refrigeration overnight. Extracts were analyzed for total N by Kjeldahl-digestion and total P by an Alpkem analyzer. A Shimadzu TOC-5050A was used to measure extracted carbon content. Unfumigated soils (control) measure represented the extractable C, N and P while microbial C, N, and P were calculated as the difference between the amount of unfumigated and fumigated soils.

Data Analysis

The effect of treatments on CO_2 , N_2O and CH_4 fluxes (annual, seasonal and monthly) and leaf litter decomposition rate was separated using ANOVA in Genstat discovery version 13. The microbial C, N and P were separated using Mann-Whitney U test. The least significant difference (lsd) for $P < 0.05$ was considered for this study.

Results and Discussion

Effect of inorganic fertilizers on annual fluxes of GHG under Albizzia shaded coffee in Manafwa District

Figure 2 shows the annual fluxes of CO_2 , N_2O and CH_4 . CO_2 emissions ranged from 19.6 to 26.1 (t C/ha/yr). N_2O emissions ranged from 3.5 to 9 (Kg N/ha/yr) and CH_4 emissions ranged from 6.9 to 9.2 (Kg C/ha/yr). Fertilizer applications had significant effect only on N_2O ($P=0.017$; One-way ANOVA) emissions. The effect of the application of NPK on emissions was significantly lower than that of the control and P but similar to that of K and N applications ($P<0.05$). The average annual emission of CO_2 was 22.8 t C/ha/yr while that of CH_4 was 8.34 Kg C/ha/yr.

[CHART]

Figure 2: Annual GHG emissions from the different fertilizer applications

Effect of inorganic fertilizers on seasonal fluxes of GHG under Albizzia shaded coffee

Figure 3 presents the seasonal GHG emissions. GHG emissions varied with season ($P<0.001$; One-way ANOVA) except for CO_2 . SON (September, October, November) and DJF (December, January, February) had significantly higher values than MAM (March, April, May) and JJA (June, July, August) for CH_4 emissions (Figure 3a) while only SON was significantly higher than the rest of the seasons for N_2O emissions (Figure 3b). There was no significant seasonal effect for CO_2 emissions with a mean of 5.68 t C/ha/yr (Figure 3c). The monthly values for CH_4 emissions ranged from 1.38 to 2.88 Kg C/ha/yr for the control,

0.93 to 3.34 Kg C/ha/yr for K applications, 0.93 to 2.84 Kg C/ha/yr for N applications, 1.07 to 3.3 Kg C/ha/yr for NPK applications and 1.4 to 2.64 Kg C/ha/yr for P applications. The monthly values for N₂O ((Figure 3b) emissions ranged from 0.44 to 1.19 Kg N/ha/yr for the control, 0.61 to 2.36 Kg N/ha/yr for K applications, 0.71 to 1.96 Kg N/ha/yr for N applications, 0.63 to 5 Kg N/ha/yr for NPK applications and 0.45 to 1.07 Kg N/ha/yr for P applications. For CO₂ emissions (Figure 3c), monthly values ranged emissions ranged from 0.44 to 7.27 t C/ha/yr for the control, 0.61 to 6.63 t C/ha/yr for K applications, 0.71 to 7.22 t C/ha/yr for N applications, 0.63 to 5.24 t C/ha/yr for NPK applications and 0.45 to 6.93 t C/ha/yr for P applications.

@ >p(- 0) * @ [CHART]

[CHART]

[CHART]

Figure 3: Seasonal GHG emissions from the different fertilizer applications in Mount Elgon

Effect of inorganic fertilizers on monthly fluxes of GHG under *Albizzia* shaded coffee

Figure 4 depicts the monthly GHG emissions. GHG emissions varied with month (P<0.001; One-way ANOVA) for N₂O and CH₄ and at P=0.014 for CO₂. The monthly values for N₂O emissions ranged from 0.14 to 0.62 Kg N/ha/yr for the control, 0.11 to 1.78 Kg N/ha/yr for K applications, 0.2 to 1.05 Kg N/ha/yr for N applications, 0.11 to 2.5 Kg N/ha/yr for NPK applications and 0.09 to 0.58 Kg N/ha/yr for P applications.

[CHART]

[CHART]

[CHART]

Figure 4: Monthly GHG emissions under different fertilizer applications in Mount Elgon

The monthly values for CH₄ emissions ranged from 0.2 to 1.1 Kg C/ha/yr for the control, 0.22 to 1.7 Kg C/ha/yr for K applications, 0.22 to 1.41 Kg C/ha/yr for N applications, 0.2 to 1.54 Kg C/ha/yr for NPK applications and 0.23 to 1.09 Kg C/ha/yr for P applications. For N₂O, October emissions were significantly higher than the rest of the months. September, March and December emissions had similar emissions but higher than May, June, July, August and November emissions. For CO₂ emissions, monthly values of emissions ranged from 0.88 to 3.48 t C/ha/yr for the control, 1.14 to 4.05 t C/ha/yr for K applications, 1.21 to 3.86 t C/ha/yr for N applications, 1.12 to 2.11 t C/ha/yr for NPK applications and 1.02 to 3.38 t C/ha/yr for P applications.

For CH₄, the months of December, November and February had significantly higher emission values compared to October, September, August, June, July, May and April emissions. Likewise, February, January, March, May, November and December had similar emission values. For CO₂, October emissions were significantly higher than those of April and February but similar to March, May, June, July, August, September, November and December emissions. Also, April emissions were significantly lower than those of March, May, July and August. Peaks in CH₄ were observed June, October and November for the control; July, December and February for P applications and May, November and February for NPK, K and N applications. Peaks in N₂O emissions were observed in May, October and December for the control; October and December for P applications; May, September, October, and December for NPK; May, October, January and March for application of K and N. Peaks in CO₂ emissions were observed May, August, October, January and March for the N, P, K applications; May, August and November for the control; May, October and December for NPK applications.

For CH₄, the months of December, November and February had significantly higher emission values compared to October, September, August, June, July, May and April emissions. Likewise, February, January, March, May, November and December had similar emission values. For CO₂, October emissions were significantly higher than those of April and February but similar to March, May, June, July, August, September, November and December emissions. Also, April emissions were significantly lower than those of March, May, July and August. Peaks in CH₄ were observed in the months of June, October and November for the control; July, December and February for P applications and May, November and February for NPK, K and N applications. Peaks in N₂O emissions were observed in May, October and December for the control; October and December for P applications; May, September, October, and December for NPK; May, October, January and March for application of K and N. Peaks in CO₂ emissions were observed May, August, October, January and March for the applications of N, P & K; May, August and November for the control; May, October and December for NPK applications.

Relative change of fluxes of GHG under *Albizzia* shaded coffee

Figure 5 shows the relative change of GHG emissions. The Emissions from other treatments were higher than the control in the months of July, November, December, February and March for CH₄. Emissions from applications of N, P, K and the mixture were higher than the control in September, October, February and March for N₂O. CO₂ emissions from other treatments were higher than the control in April, July, October, January, February and March. For CH₄ and CO₂, the emissions from the control exceeded those where fertilizers were applied in June, August, September and January and November respectively. The relative change for CH₄ emissions ranged from -60 to 82% for application of K, -74 to 133% for application of N, -46 to 99% for application of NPK, -70

to 407% for application of P. The relative change for N₂O emissions ranged from -35 to 493% for application of K, -12 to 163% for application of N, -44.9 to 558% for application of NPK, -54 to 153% for application of P. Also, the relative change for CO₂ emissions ranged from -63 to 106% for application of K, -58 to 108% for application of N, -62 to 28% for application of NPK, -69 to 153% for application of P.

[CHART]

[CHART]

[CHART]

Figure 5: Relative change in the monthly GHG fluxes under different fertilizer applications in Mount Elgon

Effect of inorganic fertilizers on microbial carbon, nitrogen and phosphorus

Table 2 shows the microbial C, N and P under different treatments. Microbial C ranged from 0.022 to 0.028, microbial P ranged from 0.006 to 0.009 while microbial N ranged from 0.017 to 0.050. Statistical analysis shows that no treatment effect was detected ($P < 0.05$; One-way ANOVA). The average microbial C, N and P were 0.024, 0.029 and 0.007 respectively.

Table 2 Effect of fertilizer application on microbial C, N and P

@ >p(- 6) * >p(- 6) * >p(- 6) * >p(- 6) * @ **Nutrient application & Microbial C & Microbial P & Microbial N**
Urea-N &

0.022

&

0.006

&

0.028

TSP- P &

0.028

&

0.007

&

0.028

Muriate of Potash-K &

0.022

&

0.007

&

0.050

NPK &

0.026

&

0.007

&

0.017

Control &

0.022

&

0.009

&

0.021

Mean &

0.024

&

0.007

&

0.028

Table 3 shows the change in microbial Carbon, Nitrogen and Phosphorus after one year of experimentation. Significant changes were observed for microbial C ($p=0.001$; One-way ANOVA) and microbial P ($p<0.001$; One-way ANOVA)

while no changes in microbial N were seen after one year of nutrient application (Appendix 1). The relative change in Microbial C was 66.7% and 1200% for microbial P.

Table 3 Change in microbial Carbon, Nitrogen and Phosphorus after one year of experimentation

@ >p(- 6) * >p(- 6) * >p(- 6) * >p(- 6) * @ **Period & Microbial C & Microbial P & Microbial N**
Start of experiment &

0.018±0.002

&

0.001±0.002

&

0.037±0.009

End of experiment &

0.030±0.002

&

0.013±0.002

&

0.020±0.007

Effect of inorganic fertilizers on decomposition rates

The decomposition rates of *Albizia* leaves varied from 0.064 to 0.079 (Figure 6). However, no statistical difference was found between treatments (Mann Whitney U test). The average decomposition rate was 0.072 per day.

[CHART]

Figure 6: Leaf litter decomposition rate under different fertilizer treatments in Mount Elgon

Discussion

Effect of fertilizer application on GHG emission in the CAF, Mount Elgon

Fertilizer applications increased N_2O emissions corroborating Owino *et al.* (2020) findings on smallholder rice paddy fields in Anyiko Wetland of Kenya. GHG emissions in the different fertilized shaded coffee of this study are comparable to the values observed by other scholars under coffee in the region for N_2O and CH_4 (eg. Capa *et al.*, 2015; Hergoualc'h *et al.*, 2008; Verchot *et al.*, 2006) except for CO_2 emissions whose values are almost triple the values observed by Hergoualc'h *et al.* (2008) in monoculture coffee systems of Costa Rica and four times that of Urzedo *et al.* (2013) under reforestation conditions in Brazil. Verchot *et al.* (2006) found N_2O emissions of approximately 7 kg N/ha/year in 3-year-old coffee plantations fertilized with 100 kg N/ha/year in southern Sumatra, India. The relatively higher magnitude of CO_2 and N_2O emissions could also be attributed to no till conditions (Bilandžija *et al.*, 2016).

The implementation of no-tillage can increase soil GHG emissions due to the maintenance of higher water content and fewer air filled pore spaces in the soil surface layer that also favors greater soil biological activity (Bilandžija *et al.*, 2016). In fact, Linn and Doran (1984) reported 3.4- and 9.4-times greater CO_2 and N_2O production from surface no till soils as compared to plowed soils at sites in Illinois, Kentucky, Minnesota and Nebraska. The low N_2O emissions under P application is in line with Sundareswar *et al.* (2003) report. It attributed the reduction of soil N_2O emissions to stimulated N immobilization. Mori *et al.* (2013) also reported that P application reduced N_2O emissions under *Acacia mangium* plantation due to the enhancement of root uptake of soil N and water. Zhang *et al.* (2014) also observed that P applications with N together significantly decreased N_2O emission. The relatively high amount of N in the studied soils then explains N_2O emissions reduction with P application observed in this study.

More so, Capa *et al.* (2015) also found that N_2O emissions increased with the fertilizer application. Largely, the small variation in the observed emissions compared to other locations is attributed to, in addition to no till, other environmental and agricultural factors such as the uniform soil water content, availability of mineral nutrients in the soil, soil temperature, and climatic conditions (Hergoualc'h *et al.* 2008; Butterfly *et al.*, 2010; Rigon *et al.*, 2018; Hiel *et al.*, 2018). Besides, Kostyanovsky *et al.* (2019) observed that emissions of CO_2 only increased with an increase in moisture and temperature but decreased under fertilizer application. Accordingly, the lack of a difference in CO_2 emissions per season in this study indicates little variation in temperature and soil water content for those seasons (Rigon *et al.*, 2018; Kostyanovsky *et al.*, 2019) that minimized the photosynthetic activity by plants and microbial activity.

Also, the increased N_2O , CH_4 and CO_2 emissions observed upon soil rewetting in the wet months after the long dry spell is in collaboration with other studies carried out in East Africa and other regions (Vilain *et al.*, 2010; Ortiz-Gonzalo *et al.*, 2018; Macharia *et al.*, 2020; Musafiri *et al.*, 2020). The pulse of the emission following a rainfall event at the onset of the season could be attributed to the Birch effect (increased decomposition and mineralization of organic matter),

since soil rewetting or high water content activates micro-organisms activity and enhances substrates supply and mineralization increasing emissions (Jarvis *et al.*, 2007; Butterfly *et al.*, 2010; Musafiri *et al.*, 2020) and contribution root respiration associated with plant growth especially for the increment in CO₂. But also, in addition to high water holding capacity, the moderate textured soils in the study area supported the higher emissions (Hiel *et al.*, 2018). However, low soil moisture associated with high soil temperatures in the severe dry spell could have restricted the flow of GHG particularly CO₂, possibly due to moderate soil microbial activity (Rigon *et al.*, 2018).

Furthermore, the highest N₂O emissions in October following fertilizer application corroborated observations of other authors (Hergoualc'h *et al.* 2008; Koehler *et al.* 2009), who reported that N₂O emissions are mainly produced within a few days after the addition of mineral fertilizers. Also, the relatively higher N₂O fluxes could be also attributed to the no till conditions because of the higher level of labile organic matter (Kostyanovsky *et al.*, 2019). Moreover, the low average soil C/N ratio of 10 also favored high emissions supporting findings of Gunderson *et al.* (2012) who reported that highest emissions occurred at the C/N ratio of 11.

Contrary to observations of Gu *et al.* (2013) that higher emissions were observed under fine textured soils, this study observed this in the sandy loam textured soils. Even an increase of soil water content or WFPS above 80% due to heavy rains especially in the last months of the experiments could have supported an exponential increase of N₂O emissions (Wu *et al.*, 2021; Dencsó *et al.*, 2020)

Effect of inorganic fertilizer application on leaf litter decomposition rate in the CAF, Mount Elgon

The average pure shade tree (*Albizia coriaria*) leaf litter decomposition rate (k) of 0.072 is comparable to findings of Cissé *et al.* (2021) for pure *D. microcarpum* (k = 0.075 week⁻¹) and *V. paradoxa* (k = 0.071 week⁻¹) leaf litters. These rates were slower compared to a mixture of shade tree leaf litters. No variation in leaf litter decomposition rate in fertilized shaded coffee supports reports by Schmitt and Perfecto (2020) who observed faster and easier decomposition with flower petals compared with leaf litter. This contradicts Chen *et al.* (2013) findings that N addition significantly decreased the decomposition of litter due to N saturation and suppression of microbial P mining within an old-growth tropical forest of China. Moreover, limited variation in temperature (mainly dry conditions) and moisture of the soil characterized by flushes of heavy rains nearing the end of the experiment hardly supported microbial activity (Zhou *et al.*, 2008; Chen *et al.*, 2020).

Effect of inorganic fertilizer application on Microbial C, N and Pin the CAF, Mount Elgon

Significant changes were observed for microbial C and microbial P while no changes in microbial N were seen after one year of nutrient application. The relative change in Microbial C was 66.7% and 1200% for microbial P. This

substantiates Su *et al.* (2014) findings that chemical fertilization significantly increased soil microbial activity involved in C, N, P and S cycling, especially for the treatments NK and NPK in paddy rice. Several authors (Crecchio *et al.* 2001; Marschner *et al.* 2001) have however reported that short term fertilizer experiments had no significant effect on the microbial activity like in long-term experiments where fertilizer additions significantly affect function, community structure, and population of soil microorganisms (Cinnadurai *et al.* 2013; Luo *et al.* 2013) for brown soil in China. Besides, this could have been attributed to the low soil moisture at the beginning of the experiment that decreased microbial activity by reducing diffusion of soluble substrates, microbial mobility and intracellular water potential (Zhou *et al.*, 2002) prompting a decrease in rates of organic matter decomposition. Likewise, heavy rains experienced towards the end of the experiment, could have steered high soil water content reducing oxygen supply to the microbes favoring inactivity (Schjønning *et al.*, 2011).

Acknowledgments

Supplemental Material

Optional Sections

Appendix 1: Effect of fertilizer application on microbial C, N and P in CAF, Mount Elgon

[CHART]

References

Al-Kaisi, M. M., & Yin, X. (2005). Tillage and crop residue effects on soil carbon and carbon dioxide emission in corn–soybean rotations. *Journal of environmental quality*, 34(2), 437-445.

Almagro, M., Ruiz-Navarro, A., Díaz-Pereira, E., Albaladejo, J., & Martínez-Mena, M. (2021). Plant residue chemical quality modulates the soil microbial response related to decomposition and soil organic carbon and nitrogen stabilization in a rainfed Mediterranean agroecosystem. *Soil Biology and Biochemistry*, 156, 108198. <https://doi.org/10.1016/j.soilbio.2021.108198>

Amos, B., Arkebauer, T. J., & Doran, J. W. (2005). Soil surface fluxes of greenhouse gases in an irrigated maize-based agroecosystem. *Agronomy--Faculty Publications*, 321. <https://digitalcommons.unl.edu/agronomyfacpub/321>

Bauer, P. J., Frederick, J. R., Novak, J. M., & Hunt, P. G. (2006). Soil CO₂ flux from a Norfolk loamy sand after 25 years of conventional and conservation tillage. *Soil and Tillage Research*, 90(1-2), 205-211.

Bayer C, Gomes J, Zanatta JA, Vieira FCB, Dieckow J. Mitigating greenhouse gas emissions from a subtropical Ultisol by using long-term no-tillage in combination with legume cover crops. *Soil Tillage Res.* 2016; 161:86–94.

Beck T, Joergensen RG, Kandeler E, Makeschin F, Nuss E, Oberholzer HR,

- Scheu S (1997). An inter-laboratory comparison of ten different ways of measuring soil microbial biomass C. *Soil Biology and Biochemistry*, 29(7):1023-32.
- Bekunda, M. A., & Woomer, P. L. (1996). Organic resource management in banana-based cropping systems of the Lake Victoria Basin, Uganda. *Agriculture, ecosystems & environment*, 59(3), 171-180.
- Bilandžija, D., Zgorelec, Ž., & Kisić, I. (2016). Influence of tillage practices and crop type on soil CO₂ emissions. *Sustainability*, 8(1), 90. doi:10.3390/su8010090
- Birch, H. F. (1958). The effect of soil drying on humus decomposition and nitrogen availability. *Plant and soil*, 10(1), 9-31. <https://doi.org/10.1007/BF01343734>
- Birkás, M., Jolánkai, M., & Schmidt, R. (2008). Environmentally-sound adaptable tillage—Solutions from Hungary, 191-194. <http://real.mtak.hu/id/eprint/5253>
- Borken, W., & Matzner, E. (2009). Reappraisal of drying and wetting effects on C and N mineralization and fluxes in soils. *Global change biology*, 15(4), 808-824. <https://doi.org/10.1111/j.1365-2486.2008.01681.x>
- Bouwman, A. F., Boumans, L. J. M., & Batjes, N. H. (2002). Modeling global annual N₂O and NO emissions from fertilized fields. *Global Biogeochemical Cycles*, 16(4), 28-1. <https://doi.org/10.1029/2001GB001812>
- Brookes, P.C., Landman, A., Pruden, G., Jenkinson, D.S. (1985). Chloroform fumigation and the release of soil nitrogen: a rapid direct extraction method to measure microbial biomass nitrogen in soil. *Soil biology and biochemistry*. 17(6):837-42.
- Butterbach-Bahl, K., Baggs, E. M., Dannenmann, M., Kiese, R., & Zechmeister-Boltenstern, S. (2013). Nitrous oxide emissions from soils: how well do we understand the processes and their controls? *Philosophical Transactions of the Royal Society B: Biological Sciences*, 368(1621), 20130122.
- Butterly, C.R., Marschner, P., McNeill, A.M., & Baldock, J.A. (2010). Rewetting CO₂ pulses in Australian agricultural soils and the influence of soil properties. *Biol Fertil Soils*. 46:739–53. <https://doi.org/10.1007/s00374-010-0481-9>.
- Capa, D., Pérez-Esteban, J., & Masaguer, A. (2015). Unsustainability of recommended fertilization rates for coffee monoculture due to high N₂O emissions. *Agronomy for sustainable development*, 35(4), 1551-1559. <https://doi.org/10.1007/s13593-015-0316-z>
- Chen, H., Li, X., Hu, F., & Shi, W. (2013). Soil nitrous oxide emissions following crop residue addition: A meta-analysis. *Global change biology*, 19(10), 2956-2964. <https://doi.org/10.1111/gcb.12274>
- Chen, Q., Niu, B., Hu, Y., Luo, T., & Zhang, G. (2020). Warming and increased precipitation indirectly affect the composition and turnover of labile-fraction soil organic matter by directly affecting vegetation and microorganisms. *Science of The Total Environment*, 714, 136787. <https://doi.org/10.1016/j.scitotenv.2020.136787>

- Christiansen, J. R., Gundersen, P., Frederiksen, P., & Vesterdal, L. (2012). Influence of hydromorphic soil conditions on greenhouse gas emissions and soil carbon stocks in a Danish temperate forest. *Forest ecology and management*, 284, 185-195. <https://doi.org/10.1016/j.foreco.2012.07.048>
- Cinnadurai, C., Gopalaswamy, G., & Balachandar, D. (2013). Diversity of cultivable *Azotobacter* in the semi-arid alfisol receiving long-term organic and inorganic nutrient amendments. *Annals of microbiology*, 63(4), 1397-1404. <https://doi.org/10.1007/s13213-013-0600-6>
- Cissé, M., Traoré, S., & Bationo, B.A. (2021). Decomposition and nutrient release from the mixed leaf litter of three agroforestry species in the Sudanian zone of West Africa. *SN Applied Sciences*, 3(2), 1-12. <https://doi.org/10.1007/s42452-021-04242-y>
- Collier, S.M., Ruark, M.D., Oates, L.G., Jokela, W.E., & Dell, C.J. (2014). Measurement of greenhouse gas flux from agricultural soils using static chambers. *J Vis Exp*. Aug 3;(90): e52110. doi: 10.3791/52110. PMID: 25146426; PMCID: PMC4827936.
- Corre, M. D., Sueta, J. P., & Veldkamp, E. (2014). Nitrogen-oxide emissions from tropical forest soils exposed to elevated nitrogen input strongly interact with rainfall quantity and seasonality. *Biogeochemistry*, 118(1), 103-120. <https://doi.org/10.1007/s10533-013-9908-3>
- Crecchio, C., Curci, M., Mininni, R., Ricciuti, P., & Ruggiero, P. (2001). Short-term effects of municipal solid waste compost amendments on soil carbon and nitrogen content, some enzyme activities and genetic diversity. *Biology and fertility of soils*, 34(5), 311-318. <https://doi.org/10.1007/s003740100413>
- Čuhel, J., & Šimek, M. (2011). Effect of pH on the denitrifying enzyme activity in pasture soils in relation to the intrinsic differences in denitrifier communities. *Folia microbiologica*, 56(3), 230-235. <https://doi.org/10.1007/s12223-011-0045-x>
- Curtin, D., Wang, H., Selles, F., McConkey, B. G., & Campbell, C. A. (2000). Tillage effects on carbon fluxes in continuous wheat and fallow-wheat rotations. *Soil Science Society of America Journal*, 64(6), 2080-2086. <https://doi.org/10.2136/sssaj2000.6462080x>
- Dalal, R. C., & Allen, D. E. (2008). Greenhouse gas fluxes from natural ecosystems. *Australian Journal of Botany*, 56(5), 369-407. <https://doi.org/10.1071/BT07128>
- Davis, K. E., Joseph, S. J., & Janssen, P. H. (2005). Effects of growth medium, inoculum size, and incubation time on culturability and isolation of soil bacteria. *Applied and environmental microbiology*, 71(2), 826-834.
- De Beenhouwer, M., Geeraert, L., Mertens, J., Van Geel, M., Aerts, R., Vanderhaegen, K., & Honnay, O. (2016). Biodiversity and carbon storage co-benefits of coffee agroforestry across a gradient of increasing management intensity in the

- SW Ethiopian highlands. *Agriculture, Ecosystems & Environment*, 222, 193-199. <http://dx.doi.org/10.1016/j.agee.2016.02.017>.
- de Urzedo, D. I., Franco, M. P., Pitombo, L. M., & do Carmo, J. B. (2013). Effects of organic and inorganic fertilizers on greenhouse gas (GHG) emissions in tropical forestry. *Forest Ecology and Management*, 310, 37-44. <https://doi.org/10.1016/j.foreco.2013.08.018>
- Dencső, M., Horel, Á., Bogunovic, I., & Tóth, E. (2021). Effects of Environmental Drivers and Agricultural Management on Soil CO₂ and N₂O Emissions. *Agronomy*, 11(1), 54. <http://doi.org/10.3390/agronomy11010054>
- Denef, K., Del Galdo, I., Venturi, A., & Cotrufo, M. F. (2013). Assessment of soil C and N stocks and fractions across 11 European soils under varying land uses. *Open Journal of Soil Science*, 3(7), 297-313. <https://doi.org/10.4236/ojss.2013.37035>
- Dilustro, J.J., Collins, B., Duncan, L., & Crawford, C. (2005). Moisture and soil texture effects on soil CO₂ efflux components in southeastern mixed pine forests. *For. Ecol. Manage.* 204, 87e97. <https://doi.org/10.1016/j.foreco.2004.09.001>.
- Dimitriou, K., Bougiatioti, A., Ramonet, M., Pierros, F., Michalopoulos, P., Liakakou, E., ... & Mihalopoulos, N. (2021). Greenhouse gases (CO₂ and CH₄) at an urban background site in Athens, Greece: Levels, sources and impact of atmospheric circulation. *Atmospheric Environment*, 253, 118372. <https://doi.org/10.1016/j.atmosenv.2021.118372>
- Dollinger, J., & Jose, S. (2018). Agroforestry for soil health. *Agroforestry Systems*, 92(2), 213-219. <https://doi.org/10.1007/s10457-018-0223-9>
- Dutaur, L., Verchot, L.V., 2007. A global inventory of the soil CH₄ sink. *Global Biogeochem. Cycles* 21, 1e9. <https://doi.org/10.1029/2006GB002734>.
- Ehrenbergerova, L., Cienciala, E., Kučera, A., Guy, L., & Habrová, H. (2016). Carbon stock in agroforestry coffee plantations with different shade trees in Villa Rica, Peru. *Agroforestry systems*, 90(3), 433-445. <https://doi.org/10.1007/s10457-015-9865-z>
- Farage, P., Pretty, J., & Ball, A. (2003). Biophysical aspects of carbon sequestration in drylands. University of Essex.
- Fowler, D., Coyle, M., Skiba, U., Sutton, M. A., Cape, J. N., Reis, S., ... & Voss, M. (2013). The global nitrogen cycle in the twenty-first century. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 368(1621), 20130164. <https://doi.org/10.1098/rstb.2013.0164>
- Frouz, J., Livečková, M., Albrechtová, J., Chronáková, A., Cajthaml, T., Pižl, V., ... & Cepáková, Š. (2013). Is the effect of trees on soil properties mediated by soil fauna? A case study from post-mining sites. *Forest Ecology and Management*, 309, 87-95. <https://doi.org/10.1016/j.foreco.2013.02.013>

- Gander, F., Proyer, R. T., Ruch, W., & Wyss, T. (2012). The good character at work: An initial study on the contribution of character strengths in identifying healthy and unhealthy work-related behavior and experience patterns. *International archives of occupational and environmental health*, 85(8), 895-904. <https://doi.org/10.1007/s00420-012-0736-x>
- Gao, X., Tenuta, M., Buckley, K.E., Zvomuya, F., Ominski, K., 2014. Greenhouse gas emissions from pig slurry applied to forage legumes on a loamy sand soil in south central Manitoba. *Can. J. Soil Sci.* 94, 149e155. <https://doi.org/10.4141/CJSS2013-117>.
- Gattinger, A., Muller, A., Haeni, M., Skinner, C., Fliessbach, A., Buchmann, N., ... & Niggli, U. (2012). Enhanced top soil carbon stocks under organic farming. *Proceedings of the National Academy of Sciences*, 109(44), 18226-18231. <https://doi.org/10.1073/pnas.1209429109>
- Goldberg, S. D., & Gebauer, G. (2009). N₂O and NO fluxes between a Norway spruce forest soil and atmosphere as affected by prolonged summer drought. *Soil Biology and Biochemistry*, 41(9), 1986-1995. <https://doi.org/10.1016/j.soilbio.2009.07.001>
- Gu, J., Nicoulaud, B., Rochette, P., Grossel, A., Hénault, C., Cellier, P., & Richard, G. (2013). A regional experiment suggests that soil texture is a major control of N₂O emissions from tile-drained winter wheat fields during the fertilization period. *Soil Biology and Biochemistry*, 60, 134-141. <https://doi.org/10.1016/j.soilbio.2013.01.029>
- Hanson, P. C., Bade, D. L., Carpenter, S. R., & Kratz, T. K. (2003). Lake metabolism: Relationships with dissolved organic carbon and phosphorus. *Limnology and Oceanography*, 48(3), 1112-1119. <https://doi.org/10.4319/lo.2003.48.3.1112>
- Hergoualc'h, K., Skiba, U., Harmand, J. M., & Hénault, C. (2008). Fluxes of greenhouse gases from Andosols under coffee in monoculture or shaded by *Inga densiflora* in Costa Rica. *Biogeochemistry*, 89(3), 329-345. <https://doi.org/10.1007/s10533-008-9222-7>
- Hiel, M.P., Barbieux, S., Pierreux, J., Olivier, C., Lobet, G., Roisin, C, ..., Dumont, B. (2018). Impact of crop residue management on crop production and soil chemistry after seven years of crop rotation in temperate climate, loamy soils. *PeerJ*, 6, 1-23. e4836. <https://doi.org/10.7717/peerj.4836>
- Högberg, M. N., Skjellberg, U., Högberg, P., & Knicker, H. (2020). Does ectomycorrhiza have a universal key role in the formation of soil organic matter in boreal forests? *Soil Biology and Biochemistry*, 140, 107635. <https://doi.org/10.1016/j.soilbio.2019.107635>
- Hussain, I., Olson, K. R., & Ebelhar, S. A. (1999). Impacts of tillage and no-till on production of maize and soybean on an eroded Illinois silt loam soil. *Soil and Tillage Research*, 52(1-2), 37-49. [https://doi.org/10.1016/S0167-1987\(99\)00054-9](https://doi.org/10.1016/S0167-1987(99)00054-9)

- Jarvis, P., Rey, A., Petsikos, C., Wingate, L., Rayment, M., Pereira, J., Banza, J., David, J., Miglietta, F., Borghetti, M., & Manca, G. (2007). Drying and wetting of Mediterranean soils stimulates decomposition and carbon dioxide emission: the "birch effect". *Tree Physiol.* 27 (7), 929–940.
- Johnson, J. M. F., Reicosky, D. C., Allmaras, R. R., Sauer, T. J., Venterea, R. T., & Dell, C. J. (2005). Greenhouse gas contributions and mitigation potential of agriculture in the central USA. *Soil and Tillage Research*, 83(1), 73-94. <https://doi.org/10.1016/j.still.2005.02.010>
- Joint FAO/WHO Expert Committee on Food Additives, Joint FAO/WHO Expert Committee on Food Additives. Meeting, & World Health Organization. (2002). *Evaluation of Certain Food Additives and Contaminants: Fifty-seventh Report of the Joint FAO/WHO Expert Committee on Food Additives* (Vol. 57). World Health Organization.
- Justine N, Tumwebaze BS, Ritah K, Nabanoga G (2019) Aboveground Species Diversity and Carbon Stocks in Smallholder Coffee Agroforestry in the Highlands of Uganda. In: Bamutaze Y., Kyamanywa S., Singh B., Nabanoga G., Lal R. (eds) *Agriculture and Ecosystem Resilience in Sub Saharan Africa. Climate Change Management*. Springer, Cham. https://doi.org/10.1007/978-3-030-12974-3_18
- Keller, M., & Reiners, W. A. (1994). Soil-atmosphere exchange of nitrous oxide, nitric oxide, and methane under secondary succession of pasture to forest in the Atlantic lowlands of Costa Rica. *Global Biogeochemical Cycles*, 8(4), 399-409. <https://doi.org/10.1029/94GB01660>
- Knief, C. (2019). Diversity of Methane-cycling Microorganisms in Soils and Their Relation to Oxygen. *Current issues in molecular biology*, 33(1), 23-56.
- Koehler, B., Corre, M. D., Veldkamp, E., Wullaert, H., & Wright, S. J. (2009). Immediate and long-term nitrogen oxide emissions from tropical forest soils exposed to elevated nitrogen input. *Global Change Biology*, 15(8), 2049-2066. <https://doi.org/10.1111/j.1365-2486.2008.01826.x>
- Koehler, B., Von Wachenfeldt, E., Kothawala, D., & Tranvik, L. J. (2012). Reactivity continuum of dissolved organic carbon decomposition in lake water. *Journal of Geophysical Research: Biogeosciences*, 117(G1). doi:10.1029/2011JG001793
- Koga, N., Shimoda, S., & Iwata, Y. (2017). Biochar impacts on crop productivity and greenhouse gas emissions from an Andosol. *Journal of environmental quality*, 46(1), 27-35. <https://doi.org/10.2134/jeq2016.04.0156>
- Kopittke, P. M., Menzies, N. W., Wang, P., McKenna, B. A., & Lombi, E. (2019). Soil and the intensification of agriculture for global food security. *Environment international*, 132, 105078. <https://doi.org/10.1016/j.envint.2019.105078>.
- Kostyanovsky, K. I., Huggins, D. R., Stockle, C. O., Morrow, J. G., & Madsen, I. J. (2019). Emissions of N₂O and CO₂ following short-term

water and n fer<https://doi.org/10.3389/fevo.2019.00063>tilization events in wheat-based cropping systems. *Frontiers in Ecology and Evolution*, 7, 63. <https://doi.org/10.3389/fevo.2019.00063>.

Lado-Monserrat, L., Lull, C., Bautista, I., Lidón, A., & Herrera, R. (2014). Soil moisture increment as a controlling variable of the “Birch effect”. Interactio<https://doi.org/10.1007/s11104-014-2037-5>ns with the pre-wetting soil moisture and litter addition. *Plant and Soil*, 379(1), 21-34. <https://doi.org/10.1007/s11104-014-2037-5>.

Lal, R. (2003). Global potential of soil carbon sequestration to mitigate the greenhouse effect. *Critical reviews in plant sciences*, 22(2), 151-184. <http://dx.doi.org/10.1080/713610854>.

Liniger, H.P., Mekdaschi Studer, R., Hauert, C., & Gurtner, M. (2011). Sustainable Land Management in Practice – Guidelines and Best Practices for Sub-Saharan Africa. TerrAfrica, World Overview of Conservation Approaches and Technologies (WOCAT) and Food and Agriculture Organization of the United Nations (FAO)

Liu, E., Teclemariam, S. G., Yan, C., Yu, J., Gu, R., Liu, S., ... & Liu, Q. (2014). Long-term effects of no-tillage management practice on soil organic carbon and its fractions in the northern China. *Geoderma*, 213, 379-384. <https://doi.org/10.1016/j.geoderma.2013.08.021>.

Ludwig, J., Meixner, F.X., Vogel, B., & Foerstner, J. (2001). Processes, influencing factors, and modelling of nitric oxide surface exchange—An overview. *Biogeochemistry* 52:225–257. <https://doi.org/10.1023/A:1006424330555>

Luo P, Han X, Wang Y, Han M, Shi H, Liu N, Bai H (2015). Influence of long-term fertilization on soil microbial biomass, dehydrogenase activity, and bacterial and fungal community structure in a brown soil of northeast China. *Annals of microbiology*. 2015 Mar;65(1):533-42.

Luo, Z., Wang, E., & Sun, O. J. (2010). Can no-tillage stimulate carbon sequestration in agricultural soils? A meta-analysis of paired experiments. *Agriculture, ecosystems & environment*, 139(1-2), 224-231. <https://doi.org/10.1016/j.agee.2010.08.006>

Macharia, M.J., Pelster, E.D., Ngetich, K.F., Shisanya, A.C., Mucheru-Muna, M., Mugendi, N.D., 2020. Soil greenhouse gas fluxes from maize production under different soil fertility management practices in East Africa. *J. Geophys. Res. Biogeosci.* <https://doi.org/10.1029/2019JG005427>.

Marschner, P., Yang, C.H., Lieberei, R., & Crowley, D.E. (2001). Soil and plant specific effects on bacterial community composition in the rhizosphere. *Soil biology and biochemistry*, 33(11):1437-45.

Mbow, C., Rosenzweig, C., Barioni, L.G., Benton, T.G., Herrero, M., Krishnapillai, M., ...& Liwenga, E. "Food security." In *Climate Change and Land*, pp. 437-550. 2019.

- Mori, T., Ohta, S., Ishizuka, S., Konda, R., Wicaksono, A., Heriyanto, J., Hamotani, Y., Gobara, Y., Kawabata, C., Kuwashima, K., Nakayama, Y., & Hardjono, A. (2013). Soil greenhouse gas fluxes and C stocks as affected by phosphorus addition in a newly established *Acacia mangium* plantation in Indonesia. *Forest Ecology and Management*, **310**, 643–651. <https://doi.org/10.1016/j.foreco.2013.08.010>
- Mosier, A. R., Halvorson, A. D., Reule, C. A., & Liu, X. J. (2006). Net global warming potential and greenhouse gas intensity in irrigated cropping systems in northeastern Colorado. *Journal of environmental quality*, *35*(4), 1584–1598. <https://doi.org/10.1007/BF00748937>
- Musafiri, C.M., Macharia, J.M., Kiboi, M.N., Ng’etich, O.K., Shisanya, C.A., Okeyo, J.M., Mugendi, D.N., Okwuosa, E.A., & Ngetich, F.K. (2020). Soil greenhouse gas fluxes from maize cropping system under different soil fertility management technologies in Kenya. *Agriculture, Ecosystems & Environment*, *301*, p.107064. <https://doi.org/10.1016/j.agee.2020.107064>
- Nisbet, E. G., Dlugokencky, E. J., & Bousquet, P. (2014). Methane on the rise—again. *Science*, *343*(6170), 493–495. <https://doi.org/10.1126/science.1247828>
- Niu, S., Sherry, R. A., Zhou, X., Wan, S., & Luo, Y. (2010). Nitrogen regulation of the climate–carbon feedback: evidence from a long-term global change experiment. *Ecology*, *91*(11), 3261–3273. <https://doi.org/10.1890/09-1634.1>
- Nugroho, R. A., Röling, W. F., Laverman, A. M., & Verhoef, H. A. (2007). Low nitrification rates in acid Scots pine forest soils are due to pH-related factors. *Microbial Ecology*, *53*(1), 89–97. <https://doi.org/10.1007/s00248-006-9142-9>
- Okalebo, J. R., Gathua, K. W., & Woomer, P. L. (1993). Fractionation of organic matter by particle size. <http://hdl.handle.net/11295/85244>
- Oorts, K., Merckx, R., Gréhan, E., Labreuche, J., & Nicolardot, B. (2007). Determinants of annual fluxes of CO₂ and N₂O in long-term no-tillage and conventional tillage systems in northern France. *Soil and Tillage Research*, *95*(1–2), 133–148. <https://doi.org/10.1016/j.still.2006.12.002>
- Ortiz-Gonzalo, D., de Neergaard, A., Vaast, P., Suárez-Villanueva, V., Oelofse, M., & Rosenstock, T. S. (2018). Multi-scale measurements show limited soil greenhouse gas emissions in Kenyan smallholder coffee-dairy systems. *Science of the Total Environment*, *626*, 328–339. <https://doi.org/10.1016/j.scitotenv.2017.12.247>
- Owino, C. N., Kitaka, N., Kipkemboi, J., & Ajwang’Ondiek, R. (2020). Assessment of Greenhouse Gases Emission in Smallholder Rice Paddies Converted from Anyiko Wetland, Kenya. *Frontiers in Environmental Science*. <https://doi.org/10.3389/fenvs.2020.00080>
- Palm, C.A., Alegre, J.C., Arevalo, L., Mutuo, P.K., Mosier, A.R., & Coe, R. (2002) Nitrous oxide and methane fluxes in six different land use sys-

- tems in the Peruvian Amazon. *Global Biogeochemical Cycles*, 16(4), 21-1. <https://doi.org/10.1029/2001GB001855>
- Paustian, K. A. O. J. H., Andren, O., Janzen, H. H., Lal, R., Smith, P., Tian, G., ... & Woomer, P. L. (1997). Agricultural soils as a sink to mitigate CO₂ emissions. *Soil use and management*, 13, 230-244. <https://doi.org/10.1111/j.1475-2743.1997.tb00594.x>
- Pelster, D., Rufino, M., Rosenstock, T., Mango, J., Saiz, G., Diaz-Pines, E., ... & Butterbach-Bahl, K. (2017). Smallholder farms in eastern African tropical highlands have low soil greenhouse gas fluxes. *Biogeosciences*, 14(1), 187-202. <https://doi.org/10.5194/bg-14-187-2017>
- Peng, Q., Dong, Y., Qi, Y., Xiao, S., He, Y., & Ma, T. (2011). Effects of nitrogen fertilization on soil respiration in temperate grassland in Inner Mongolia, China. *Environmental Earth Sciences*, 62(6), 1163-1171. <https://doi.org/10.1007/s12665-010-0605-4>
- Phillips, R. L. (2007). Organic agriculture and nitrous oxide emissions at sub-zero soil temperatures. *Journal of environmental quality*, 36(1), 23-30. <https://doi.org/10.2134/jeq2006.0205>
- Phillips, R. L., Tanaka, D. L., Archer, D. W., & Hanson, J. D. (2009). Fertilizer application timing influences greenhouse gas fluxes over a growing season. *Journal of environmental quality*, 38(4), 1569-1579. <https://doi.org/10.2134/jeq2008.0483>
- Pilegaard, K., Skiba, U., Ambus, P., Beier, C., Brüggemann, N., Butterbach-Bahl, K., ... & Zechmeister-Boltenstern, S. (2006). Factors controlling regional differences in forest soil emission of nitrogen oxides (NO and N₂O). *Biogeosciences*, 3(4), 651-661. <https://doi.org/10.5194/bg-3-651-2006>
- Plante AF, McGill WB (2002) Soil aggregate dynamics and the retention of organic matter in laboratory-incubated soil with differing simulated tillage frequencies. *Soil and Tillage research*, 66(1), 79-92. <https://doi.org/10.5194/bg-3-651-2006>
- Plante, A. F., & McGill, W. B. (2002). Soil aggregate dynamics and the retention of organic matter in laboratory-incubated soil with differing simulated tillage frequencies. *Soil and Tillage research*, 66(1), 79-92. [https://doi.org/10.1016/S0167-1987\(02\)00015-6](https://doi.org/10.1016/S0167-1987(02)00015-6)
- Powlson, D. S., Whitmore, A. P., & Goulding, K. W. (2011). Soil carbon sequestration to mitigate climate change: a critical re-examination to identify the true and the false. *European Journal of Soil Science*, 62(1), 42-55. <https://doi.org/10.1111/j.1365-2389.2010.01342.x>
- Raposo, E., Brito, L. F., Januszkiewicz, E. R., Oliveira, L. F., Versuti, J., Assumpção, F. M., ... Ruggieri, A. C. (2020). Greenhouse gases emissions from tropical grasslands affected by nitrogen fertilizer management. *Agronomy Journal*. doi:10.1002/agj2.20385

- Rastogi, M., Singh, S., & Pathak, H. (2002). Emission of carbon dioxide from soil. *Current science*, 82(5), 510-517.
- Rigon, J. P. G., Calonego, J. C., Rosolem, C. A., & Scala, N. L. (2018). Cover crop rotations in no-till system: short-term CO₂ emissions and soybean yield. *Scientia Agricola*, 75, 18-26. DOI: <http://dx.doi.org/10.1590/1678-992X-2016-0286>
- Rochette, P., & Janzen, H.H. (2005). Towards a revised coefficient for estimating N₂O emissions from legumes. *Nutrient Cycling in Agroecosystems*, 73(2), 171-179. <https://doi.org/10.1007/s10705-005-0357-9>
- Rosenzweig, C., Mbow, C., Barioni, L.G., Benton, T.G., Herrero, M., Krishnapillai, M., ... & Portugal-Pereira, J. (2020). Climate change responses benefit from a global food system approach. *Nature Food*, 1(2), 94-97. <https://doi.org/10.1038/s43016-020-0031-z>
- Sainju, U. M., Jabro, J. D., & Stevens, W. B. (2008). Soil carbon dioxide emission and carbon content as affected by irrigation, tillage, cropping system, and nitrogen fertilization. <https://doi.org/10.2134/jeq2006.0392>
- Saiz, G., Byrne, K. A., Butterbach-Bahl, K. L. A. U. S., Kiese, R., Blujdea, V., & Farrell, E. P. (2006). Stand age-related effects on soil respiration in a first rotation Sitka spruce chronosequence in central Ireland. *Global Change Biology*, 12(6), 1007-1020. <https://doi.org/10.1111/j.1365-2486.2006.01145.x>
- Schindlbacher A, Zechmeister-Boltenstern S, Butterbach-Bahl K (2004) Effects of soil moisture and temperature on NO, NO₂, and N₂O emissions from European forest soils. *Journal of Geophysical Research: Atmospheres*, 109(D17). <https://doi.org/10.1029/2004JD004590>
- Schindlbacher, A., Zechmeister-Boltenstern, S., & Butterbach-Bahl, K. (2004). Effects of soil moisture and temperature on NO, NO₂, and N₂O emissions from European forest soils. *Journal of Geophysical Research: Atmospheres*, 109(D17). <https://doi.org/10.1029/2004JD004590>
- Schjønning, P., Thomsen, I. K., Petersen, S. O., Kristensen, K., & Christensen, B. T. (2011). Relating soil microbial activity to water content and tillage-induced differences in soil structure. *Geoderma*, 163(3-4), 256-264. <https://doi.org/10.1016/j.geoderma.2011.04.022>
- Schmitt, L., & Perfecto, I. (2020). Who gives a flux? Synchronous flowering of *Coffea arabica* accelerates leaf litter decomposition. *Ecosphere*, 11(7), e03186. doi:10.1002/ecs2.3186
- Sebuliba, E., Majaliwa, J. G., Isubikalu, P., Turyahabwe, N., Eilu, G., & Ekwaru, A. (2021). Characteristics of shade trees used under Arabica coffee agroforestry systems in Mount Elgon Region, Eastern Uganda. *Agroforestry Systems*, 1-13. <https://doi.org/10.1007/s10457-021-00688-6>
- Six, J., Ogle, S. M., Jay Breidt, F., Conant, R. T., Mosier, A. R., & Paustian, K.


- (2004). The potential to mitigate global warming with no-tillage management is only realized when practised in the long term. *Global change biology*, 10(2), 155-160. <https://doi.org/10.1111/j.1529-8817.2003.00730.x>
- Skiba, U., Fowler, D., & Smith, K. A. (1997). Nitric oxide emissions from agricultural soils in temperate and tropical climates: sources, controls and mitigation options. *Nutrient Cycling in Agroecosystems*, 48(1), 139-153. <https://doi.org/10.1023/A:1009734514983>
- Smith, C. J., Forster, P. M., Allen, M., Leach, N., Millar, R. J., Passerello, G. A., & Regayre, L. A. (2018). FAIR v1. 3: a simple emissions-based impulse response and carbon cycle model. *Geoscientific Model Development*, 11(6), 2273-2297. <https://doi.org/10.5194/gmd-11-2273-2018>
- Smith, K. A., Ball, T., Conen, F., Dobbie, K. E., Massheder, J., & Rey, A. (2003). Exchange of greenhouse gases between soil and atmosphere: interactions of soil physical factors and biological processes. *European journal of soil science*, 54(4), 779-791. <https://doi.org/10.1046/j.1351-0754.2003.0567.x>
- Solis, R., Vallejos-Torres, G., Arévalo, L., Marín-Díaz, J., Ñique-Alvarez, M., Engedal, T., & Bruun, T. (2020). Carbon stocks and the use of shade trees in different coffee growing systems in the Peruvian Amazon. *The Journal of Agricultural Science*, 158(6), 450-460. <https://doi.org/10.1017/S002185962000074X>
- Sponseller, R. A. (2007). Precipitation pulses and soil CO₂ flux in a Sonoran Desert ecosystem. *Global Change Biology*, 13(2), 426-436. <https://doi.org/10.1111/j.1365-2486.2006.01307.x>
- Sun, S., Liu, J., Zhang, M., & He, S. (2019). Simultaneous improving nitrogen removal and decreasing greenhouse gas emission with biofilm carriers' addition in ecological floating bed. *Bioresource technology*, 292, 121944. <https://doi.org/10.1016/j.biortech.2019.121944>
- Sundareshwar, P. V., Morris, J. T., & Koepflex, E. K. (2003). Phosphorus limitation for coastal ecosystem processes. *Science*, 299, 563-565. <https://doi.org/10.1126/science.1079100>
- Tang, J., Baldocchi, D. D., Qi, Y., & Xu, L. (2003). Assessing soil CO₂ efflux using continuous measurements of CO₂ profiles in soils with small solid-state sensors. *Agricultural and Forest Meteorology*, 118(3-4), 207-220. [https://doi.org/10.1016/S0168-1923\(03\)00112-6](https://doi.org/10.1016/S0168-1923(03)00112-6)
- Tongwane, M., Mdlambuzi, T., Moeletsi, M., Tsubo, M., Mliswa, V., & Grootboom, L. (2016). Greenhouse gas emissions from different crop production and management practices in South Africa. *Environmental Development*, 19, 23-35. <https://doi.org/10.1016/j.envdev.2016.06.004>
- Ussiri, D. A., & Lal, R. (2009). Long-term tillage effects on soil carbon storage and carbon dioxide emissions in continuous corn cropping system from an alfisol in Ohio. *Soil and Tillage Research*, 104(1), 39-47. <https://doi.org/10.1016/j.still.2008.11.008>

- Vaast, P., Van Kanten, R., Siles, P., Angrand, J., & Aguilar, A. (2008). Bio-physical interactions between timber trees and Arabica coffee in suboptimal conditions of Central America. In *Toward Agroforestry Design* (pp. 133-146). Springer, Dordrecht. <https://doi.org/10.1007/s10457-018-0223-9>
- van der Weerden, T. J., Kelliher, F. M., & de Klein, C. A. (2012). Influence of pore size distribution and soil water content on nitrous oxide emissions. *Soil Research*, 50(2), 125-135. <https://doi.org/10.1071/SR11112>
- Vance, ED, Brookes PC, Jenkinson DS (1987). An extraction method for measuring soil microbial biomass C. *Soil biology and Biochemistry*, 19(6), 703-7.
- Verchot, L. V., Hutabarat, L., Hairiah, K., & Van Noordwijk, M. (2006). Nitrogen availability and soil N₂O emissions following conversion of forests to coffee in southern Sumatra. *Global biogeochemical cycles*, 20(4). doi:10.1029/2005GB002469
- Verchot, L. V., Júnior, S. B., de Oliveira, V. C., Mutegi, J. K., Cattânio, J. H., & Davidson, E. A. (2008). Fluxes of CH₄, CO₂, NO, and N₂O in an improved fallow agroforestry system in eastern Amazonia. *Agriculture, ecosystems & environment*, 126(1-2), 113-121. doi:10.1016/j.agee.2008.01.012
- Verchot, L. V., Júnior, S. B., de Oliveira, V. C., Mutegi, J. K., Cattânio, J. H., & Davidson, E. A. (2008). Fluxes of CH₄, CO₂, NO, and N₂O in an improved fallow agroforestry system in eastern Amazonia. *Agriculture, ecosystems & environment*, 126(1-2), 113-121. <https://doi.org/10.1016/j.agee.2008.01.012>
- Vilain, G., Garnier, J., Tallec, G., & Cellier, P. (2010). Effect of slope position and land use on nitrous oxide (N₂O) emissions (Seine Basin, France). *Agricultural and Forest Meteorology*, 150(9), 1192-1202. <https://doi.org/10.1016/j.agrformet.2010.05.004>
- Wang, Y., Hao, Y., Cui, X.Y., Zhao, H., Xu, C., Zhou, X., & Xu, Z. (2014). Responses of soil respiration and its components to drought stress. *Journal of Soils and Sediments*, 14(1):99-109.
- Wang, H., Shen, M., Hui, D., Chen, J., Sun, G., Wang, X., ... & Zhang, Y. (2019). Straw incorporation influences soil organic carbon sequestration, greenhouse gas emission, and crop yields in a Chinese rice (*Oryza sativa* L.)–wheat (*Triticum aestivum* L.) cropping system. *Soil and Tillage Research*, 195, 104377. <https://doi.org/10.1016/j.still.2019.104377>
- Wu, H., Wiesmeier, M., Yu, Q., Steffens, M., Han, X., & Kögel-Knabner, I. (2012). Labile organic C and N mineralization of soil aggregate size classes in semiarid grasslands as affected by grazing management. *Biology and Fertility of Soils*, 48(3), 305-313. <https://doi.org/10.1007/s00374-011-0627-4>
- Wu, Y., Gao, X., Cao, D., Li, L., Li, X., & Zeng, F. (2021). Nitrous Oxide Emissions from an Alpine Grassland as Affected by Nitrogen Addition. *Atmosphere*, 12(8), 976. <https://doi.org/10.3390/atmos12080976>

Zhang W, Zhu X, Luo Y, Rafique R, Chen H, Huang J, Mo J (2014) Responses of nitrous oxide emissions to nitrogen and phosphorus additions in two tropical plantations with N-fixing *vs.* non-N-fixing tree species. *Biogeosciences Discussions*, **11**, 1413–1442. <https://doi.org/10.5194/bg-11-4941-2014>

Zhang W, Zhu X, Luo Y, Rafique R, Chen H, Huang J, Mo J. (2014). Responses of nitrous oxide emissions to nitrogen and phosphorus additions in two tropical plantations with N-fixing *vs.* non-N-fixing tree species. *Biogeosciences Discussions*, **11**, 1413–1442. <https://doi.org/10.5194/bg-11-4941-2014>

Figures and Tables



[CHART]

Figure 2: Annual GHG emissions from the different fertilizer applications

[CHART]
[CHART]

Figure 3. Seasonal GHG emissions from the different fertilizer applications in Mount Elgon

[CHART]
[CHART]

Figure 7. Monthly GHG emissions under different fertilizer applications in Mount Elgon

[CHART]
[CHART]

Figure 5. Relative change in the monthly GHG fluxes under different fertilizer applications in Mount Elgon

[CHART]

Figure 6. Leaf litter decomposition rate under different fertilizer treatments in Mount Elgon