

# Major modes of climate variability dominate nonlinear Antarctic ice-sheet elevation changes 2002-2020

Matt A King<sup>1</sup> and Poul Christoffersen<sup>2</sup>

<sup>1</sup>University of Tasmania

<sup>2</sup>University of Cambridge

March 05, 2024

## Abstract

We explore the links between elevation variability of the Antarctic Ice Sheet (AIS) and large-scale climate modes. Using multiple linear regression, we quantify the cumulative effects of El Nino Southern Oscillation (ENSO) and the Southern Annular Mode (SAM) on gridded AIS elevations. Cumulative ENSO and SAM explain a median of 29% of the partial variance and up to 85% in some coastal areas. After spatial smoothing, these signals have high spatial correlation with those from GRACE gravimetry ( $r \sim 0.65$  each). Much of the signal is removed by a model of firn densification but inter-model differences exist especially for ENSO. At the lower parts of the Thwaites and Pine Island glaciers, near their grounding line, we find the Amundsen Sea Low (ASL) explains  $\sim 90\%$  of the observed elevation variability. There, firn effects explain only a small fraction of the variability, suggesting significant height changes have a climatological ice-dynamic response.

**Major modes of climate variability dominate nonlinear Antarctic ice-sheet elevation changes 2002-2020**

**Matt A. King<sup>1,2</sup>, Poul Christoffersen<sup>2,3</sup>**

<sup>1</sup>Surveying and Spatial Sciences, School of Geography, Planning, and Spatial Sciences, University of Tasmania, Hobart, Tasmania 7001, Australia

<sup>2</sup>The Australian Centre for Excellence in Antarctic Science, University of Tasmania, Hobart, Tasmania 7001, Australia

<sup>3</sup>Institute for Marine and Antarctic Studies, University of Tasmania, Hobart, Tasmania 7001, Australia

Corresponding author: Matt King ([Matt.King@utas.edu.au](mailto:Matt.King@utas.edu.au))

**Key Points:**

- Cumulative effects of large-scale climate modes dominate detrended altimeter time series of Antarctic ice elevation 2002-2020
- These decadal signals have the same spatial pattern in altimeter ice height and GRACE mass time series.
- These decadal signals are largely due to surface mass balance, but ice dynamic changes may play a role in the Amundsen Sea Embayment

## Abstract

We explore the links between elevation variability of the Antarctic Ice Sheet (AIS) and large-scale climate modes. Using multiple linear regression, we quantify the cumulative effects of El Niño Southern Oscillation (ENSO) and the Southern Annular Mode (SAM) on gridded AIS elevations. Cumulative ENSO and SAM explain a median of 29% of the partial variance and up to 85% in some coastal areas. After spatial smoothing, these signals have high spatial correlation with those from GRACE gravimetry ( $r \sim 0.65$  each). Much of the signal is removed by a model of firn densification but inter-model differences exist especially for ENSO. At the lower parts of the Thwaites and Pine Island glaciers, near their grounding line, we find the Amundsen Sea Low (ASL) explains  $\sim 90\%$  of the observed elevation variability. There, firn effects explain only a small fraction of the variability, suggesting significant height changes have a climatological ice-dynamic response.

## Plain Language Summary

This study investigates how variations in the height of the Antarctic Ice Sheet (AIS) are connected to large-scale climate patterns. We used a statistical method to measure the effects of two climate phenomena: El Niño Southern Oscillation (ENSO) and the Southern Annular Mode (SAM). We found that the cumulative effects of these phenomena account for about 29% of the variations in AIS height on average, and up to 85% in some coastal areas. These patterns match well with independent data from the GRACE satellites over the same period. Applying a model that considers the compacting of snow into ice (firn densification) removes much of this signal, suggesting much, but not all, of the signals are related to snowfall variations. At the fronts of the rapidly changing Thwaites and Pine Island glaciers, the dominant climate phenomenon is the Amundsen Sea Low (ASL), which varies in strength and location. Here, the cumulative effects of the ASL changes explain about 90% of the variations in height of these glaciers, with only a small part explained by firn effects. We suggest the unexplained variability is at least partly due to changes in ice flow.

## 1 Introduction

Observations of the changing volume of the Antarctic Ice Sheet play a major role in understanding ice-sheet change (e.g., Otonari et al., 2023; Shepherd et al., 2012) from the whole-of-ice-sheet down to individual glaciers (e.g., Smith et al., 2020; Wingham et al., 2009). The now three-decade record of continuous ice volume change captures the variability and longer-term change of both surface mass balance (SMB), and related firn processes, and elevation effects of changing ice dynamics. These changes are, respectively, related to atmospheric and oceanic processes (Horwath et al., 2012; Smith et al., 2020). Several studies have examined the relationship between ice height changes and modes of climate variability, in particular linking them to both El Niño - Southern Oscillation (ENSO) and the Antarctic Circumpolar Wave (Kaitheri et al., 2021; Mémin et al., 2015; Mémin et al., 2014).

Strangely, less studied in this context is the role of the dominant mode of climate variability in the Southern Hemisphere, the Southern Annular Mode (SAM). Despite SAM driving variability and trends in SMB across a wide range of timescales (Diener et al., 2021; Medley & Thomas,

2019; van den Broeke & van Lipzig, 2017), SAM has yet to be linked to observations of ice sheet elevation change, with one related study reporting no correlation to estimates of ice shelf elevation change (Paolo et al., 2018). By contrast, the cumulative sum of SAM (Diener et al., 2021; Kim et al., 2020) has recently been shown to be linearly related to the dominant signal in detrended surface mass time series derived from satellite gravimetry (King et al., 2023), with large-scale spatially-coherent signal across coastal regions at decadal timescales.

The ~300 km spatial resolution of satellite gravimetry, combined with uncertainties in models of SMB (Mottram et al., 2021), meant that King et al. (2023) were not able to separate the relative contributions of SMB and ice dynamical change forced respectively by the atmosphere and ocean (Hansen et al., 2021; Kim et al., 2020; Palóczy et al., 2018; Spence et al., 2017; Thomas et al., 2017; Verfaillie et al., 2022). In particular, ice dynamical change will have a distinct spatial pattern compared to SMB that is not detectable by GRACE but could be possible with altimetry (Smith et al., 2020). Detecting (or otherwise) a response of the grounded ice sheet to large-scale climate variability via the oceans and ice shelves would provide important insights into ice-sheet sensitivity to climate change.

In this paper we analyze a recent gridded compilation of satellite altimeter data and compare these time series to cumulative climate indices. We compare the derived signals to those from space gravimetry and then, taking advantage of the high-resolution altimeter data, explore the signal over key ice streams: Thwaites, Pine Island, Totten, and Denman.

## 2 Datasets and Analysis

### 2.1 Altimeter dataset

We make use of a gridded altimeter product (Nilsson et al., 2023) at 1920 m horizontal resolution and covering the period from Apr 1985 to Dec 2020 (Nilsson et al., 2022). We spatially down-sample this to 5 km horizontal resolution. To facilitate comparison with space gravimetry data we only make use of data from 2002 to the end of the record. The dataset contains monthly ice-sheet elevation-change data derived from a range of radar and laser altimeter missions; over the study period these are ERS-2, Envisat and CryoSat-2 and ICESat and ICESat-2. The approach to accounting for differences in reflection surfaces and other systematic effects is described by Nilsson et al. (2022). To reduce spatial noise we apply a Gaussian smoother with widths specified below, with width defined at the half height of the function, consistent with the definition commonly used in GRACE data smoothing (Wahr et al., 1998).

### 2.2 Space gravimetry dataset

We use the COST-G RL01 Level-3 50 km gridded GRACE and GRACE-FO V0002 dataset obtained from <http://gravis.gfz-potsdam.de/antarctica> (Sasgen et al., 2020). We make use of data from Mar 2002 to Dec 2020, with the end point chosen to match the end of the altimetry dataset.

The data are spaced approximately monthly and with a data gap of ~12 months between GRACE and GRACE-FO from mid-2017 to mid-2018.

We note that while this product is gridded at 50 km, the intrinsic GRACE resolution is 200-300 km. Post-processing steps include replacement of low-degree GRACE coefficients and insertion of degree-1 terms using standard approaches (Dahle & Murböck, 2020; Sasgen et al., 2020).

Since we are interested in decadal variability and trends, we also lightly smooth the altimetry and GRACE data with a Gaussian filter with width 7 months (Wahr et al., 1998).

### 2.3 Climate indices

We compare the altimeter and GRACE data primarily with SAM and ENSO indices, with additional comparison to Amundsen Sea Low (ASL) indices in the Amundsen Sea region. For the ASL indices, we make use of both the absolute center pressure (ASLP) and longitude (ASL $\lambda$ ) within the ASL Index version 3.20210820-era5 based on monthly ERA5 reanalysis data (Hosking et al., 2016). For SAM, we make use of the Marshall station index (Marshall, 2003). For ENSO, we make use of the Nino3.4 index based on the HadISST1 dataset (Rayner, 2003) and use a 6-month lag (King et al., 2023; Paolo et al., 2018). We normalized each index with the mean and standard deviation computed over 1971-1999 inclusive, then cumulatively summed them, limited them to the data period, and then renormalized to produce SAM $_{\Sigma}$ , ENSO $_{\Sigma}$ , ASLP $_{\Sigma}$ , and ASL $\lambda_{\Sigma}$ .

The raw indices and their cumulative sums are shown in Fig S1. Correlations above 0.7 are evident between ASLP $_{\Sigma}$  and SAM $_{\Sigma}$  and between ASL $\lambda_{\Sigma}$  and ENSO $_{\Sigma}$  (Fig S1, S2). This is due to the ASL being affected by larger-scale modes of climate variability, with SAM in particular modulating its absolute pressure and ENSO modulating the longitude of its center (Clem et al., 2017; Hosking et al., 2016; Turner et al., 2013).

### 2.4 Multi-variate Empirical Orthogonal Functions

For a data-driven analysis we make use of Multi-variate Empirical Orthogonal Functions (MEOF) (Wang, 1992). MEOFs are an extension of conventional Empirical Orthogonal Functions but allow the dominant modes across multiple variables to be identified rather than treating each variable separately. We use MEOF to analyze the elevation and mass change gridded datasets after individual normalization. We first smooth the altimetry dataset with a 50 km-wide Gaussian smoother and sub-sample the altimeter dataset to match the 50 km horizontal resolution of GRACE. Given the limited sampling of altimetry in the northern Antarctic Peninsula we truncate that region from both datasets prior to computing MEOFs.

### 2.5 Regression

Using ordinary least squares, we solved the coefficients ( $a$ ,  $b$ ,  $c$ ,  $d$ , and  $e$ ) of the functional model describing time-evolving elevation ( $h$ ) with time ( $t$ ):

$$h(t_i) = a + b(t_i - t_0) + \sum_{k=1}^2 (c_k^s \sin(2\pi f_k t_i) + c_k^c \cos(2\pi f_k t_i)) + dSAM_{\Sigma} + eENSO_{\Sigma} \quad (1)$$

Where  $f_k = [1, 2]$  cycles per year. We adopted  $t_0$  as the mid point of the altimeter series.

## 2.6 Data uncertainty

For regression parameter uncertainties, we recognize the existence of temporal correlations in the altimeter time series (Ferguson et al., 2004), in part due to SMB variation (King & Watson, 2020), and take these into account. Following King et al. (2023), we compared trend uncertainties from a linear regression using a Generalized Gauss Markov noise model to those generated using a white noise only (temporally uncorrelated) noise model using HECTOR v2.0 software (Bos et al., 2013). For regressions that included the SAM and ENSO terms, the white noise only model produced uncertainties a factor of 3 too small, taken as the median of the ratio of trend uncertainties, or factor 40 too small when not including the SAM and ENSO terms. We applied these scale factors to the uncertainties from the regression. For the GRACE uncertainties we used the scale factors of King et al. (2023).

## 3 Results

### 3.1 Ice-sheet scale analysis

Our data-driven MEOF analysis shows that ice elevation and mass time series are both dominated by decadal-scale variability (Fig. S3c, f). Together, the two leading modes explain 65% of the non-linear variance of the combined and smoothed time series. Their corresponding principal components (PCs) correlate with detrended  $\text{SAM}_\Sigma$  ( $r=0.73$ ) and 6-month lagged  $\text{ENSO}_\Sigma$  ( $r=0.89$ ). The  $\text{ASLP}_\Sigma$  and  $\text{ASL}\lambda_\Sigma$  terms are not of direct relevance at the ice-sheet scale given the limited geographical footprint of influence of the ASL, but also have high correlations with the data.

GRACE and altimetry MEOFs have a high spatial correlation (Fig. S3a-b, d-e;  $r=0.87$  for MEOF1 and  $r=0.75$  for MEOF2) suggesting they are sensing the same signal and are both dominated by coastal changes. The potential in the high-resolution altimetry record is particularly evident in MEOF1 where the spatially-diffuse signal in GRACE (Fig. S3a) is shown to be concentrated over small regions that coincide with the major ice streams of the Amundsen Sea Embayment and the coastline of the Bellingshausen Sea and Marie Byrd Land (Fig. S3b). We note that while MEOF3 (Fig. S4) is partly affected by striping in the GRACE field, characteristic of GRACE systematic error, coherent signal is evident between GRACE and altimetry along the coastlines of the Bellingshausen Sea, Marie Byrd Land and Wilkes Land, suggesting the signal is robust in those regions, although the variance explained (5%) is much smaller than MEOFs 1 and 2. A similar signal to PC3, with periodicities of ~4-7 years, has also been identified in analysis of GRACE data (King et al., 2023; Mémmin et al., 2015). Beyond MEOF3, the modes explain little variance (<4%) and are dominated by noise, at least for GRACE (Fig. S4d).

To quantify the SAM and ENSO contribution to ice sheet elevation change we regress the altimetry time series against  $\text{SAM}_\Sigma$  and  $\text{ENSO}_\Sigma$  and the other parameters in Eq. 1. Here we use the gridded data after applying a 10 km Gaussian spatial filter. The 5 km gridded altimeter regression analysis shown in Fig. 1a,b reveals large-scale spatially coherent signal relating to SAM and ENSO around the coasts of Antarctica. Together, these two terms often explain more than 40% of the partial variance of the timeseries around the coast and into the interior, with the


partial variance controlling for the other regression terms. The median partial-variance explained across the ice sheet is 29% (Fig. 1c). The  $\text{SAM}_\Sigma$  coefficient is strongest in the Amundsen Sea Embayment where it centers on the Pine Island, Thwaites, Smith, and Pope Glaciers (Fig. S5a). The negative elevation signal in this region is linked to periods where positive SAM dominates negative SAM (positive  $\text{SAM}_\Sigma$ ). Other strong signal exists along the coastal zone of the Bellingshausen Sea, Marie Byrd Land, and parts of coastal East Antarctica. A more diffuse signal is evident in the interior of West Antarctica and parts of East Antarctica (Fig. S6a). The  $\text{ENSO}_\Sigma$  coefficient has particularly high positive values, indicating elevation increase associated with sustained El Niño, along the coast of the Bellingshausen Sea and well upstream into Pine Island Glacier (Fig. S5b)

Applying a 200 km Gaussian smoother to the altimeter data and rerunning the regression (Fig. 1d-e) produces coefficients with large-scale spatial coherence and larger partial variances explained, often exceeding 60% in key coastal regions but extending well into the interior of the ice sheet (Fig. 1f). Comparing them to results of a regression with GRACE data (Fig. 1g-h) (King et al., 2023) shows high agreement in the signs and spatial distribution of the signal. We note that there are insufficient altimeter data in the Northern Antarctic Peninsula to analyze the signal in this region. Computing spatial correlations between the smoothed altimetry regression and the GRACE regression gives  $r=0.65$  for  $\text{SAM}_\Sigma$  and  $r=0.68$  for  $\text{ENSO}_\Sigma$ .

We next examine the role of SMB variability on the estimated coefficients from the altimetry regression. To do this we subtract the IMAU Firn Densification Model (IMAU FDM) v1.2A (Veldhuijsen et al., 2023) from the altimetry time series and repeat the regression. The results are shown in Figure 2. Comparing Fig. 2a with Fig. 1a shows that IMAU FDM effectively removes all the SAM-related signal in East Antarctic Ice Sheet (EAIS) but much of the SAM signal remains in West Antarctic Ice Sheet (WAIS). Much of the coastal EAIS ENSO-related signal is removed by IMAU FDM but with small over-correction evident for much of the ice sheet, including signal reversing sign in George V Land and WAIS. Repeating the regression but instead using the GSFC FDM v1.2.1 (Medley et al., 2022) shows that there is significant sensitivity to the choice of FDM (Fig. 2d-f), with GSFC FDM apparently over-correcting ENSO-related signal in the Totten Glacier region but in much better agreement with the altimetry in WAIS. Given the decadal timescales of the signals, these inter-model differences are likely to have contributions from both the FDMs themselves and their underlying SMB models (Medley et al., 2022).


The combination of coefficients estimated from each of GRACE and altimetry allows the density of these terms to be estimated. Given the GRACE resolution half-width is about 100km, we computed densities and their at locations 100 km upstream of the grounding lines of the Thwaites, Pine Island, Totten, and Denman glacier. These computed densities are sensitive to the radius of the Gaussian smoother applied to the altimetry data, and we adopted a 200 km smoother to approximate the GRACE resolution. ENSO-related results are highly uncertain in the Denman and Totten glacier regions due to limited signal, but the other densities (SAM and ENSO related) suggest the observed changes have a density between snow and ice, clustering around  $600 \text{ kg/m}^3$  (Fig. S6). While the estimates are uncertain, they suggest that some of the signal could originate in ice dynamics rather than SMB.

223



224

**Figure 1.** Results of regression analysis of gridded data. Shown are the  $SAM_{\Sigma}$  and  $ENSO_{\Sigma}$  coefficients and variances explained for the altimetry (top row), altimetry after 200 km Gaussian smoothing (middle row), and GRACE (bottom row). The partial variances explained by  $SAM_{\Sigma}$  and lagged  $ENSO_{\Sigma}$  are in the right column.



**Figure 2.** Results of regression analysis of FDM-corrected gridded altimeter data. Regression coefficients are shown (left and central columns) and the partial variances explained by SAM<sub>Σ</sub> and lagged ENSO<sub>Σ</sub> (right column). Shown are the coefficients and variances explained for the altimetry time series after subtracting of the IMAU FDM (top row) and GSFC FDM (bottom row).

Next, we explore the origins of these signals further on a glacier-by-glacier basis.

### 3.2 Regional scale analysis

#### 3.2.1 Thwaites and Pine Island glaciers

The partial variance explained by the SAM<sub>Σ</sub> and ENSO<sub>Σ</sub> terms (before subtracting an FDM) is above 60% for much of the Amundsen Sea Embayment (ASE; Fig. 1c, f; S5c,f). Regardless of the FDM model adopted, much SAM<sub>Σ</sub> signal remains in the ASE broadly and ENSO<sub>Σ</sub> signal is evident in the Pine Island Glacier region (Fig. 2). Closer examination of these regions in Fig. S5

(top row) indicates that the ASE signals are concentrated along low-elevation and fast flowing regions that correspond to Pine Island, Thwaites, and nearby glaciers. This is further evidenced through cross-sections near to the front of these glaciers (Fig. S7) along the yellow lines in Fig S5. It is notable that the phase of the SAM-related signal is switched in the fast-flowing region of Pine Island Glacier.


Coefficient magnitudes generally decay upstream of the grounding line (Fig. S8). Subtracting the IMAU FDM before performing the regression results in coefficients along the centerline and cross profiles that are shifted nearly uniformly but are not significantly altered in their spatial pattern (dashed lines Fig. S7-S8). Together these results suggest there may be an ice dynamic component within the  $SAM_{\Sigma}$  and  $ENSO_{\Sigma}$  coefficients in addition to residual SMB/FDM signal.

Along the coastal margin of the ASE the climatology is more directly controlled by the ASL than SAM and ENSO which modulate its depth and location (Clem et al., 2017; Turner et al., 2013). To explore this further we repeated the regression replacing  $SAM_{\Sigma}$  and  $ENSO_{\Sigma}$  in Equation 1 with  $ASLP_{\Sigma}$  and  $ASL\lambda_{\Sigma}$ . While the magnitude of the estimated coefficients differs between  $SAM_{\Sigma}$ / $ASLP_{\Sigma}$  and  $ENSO_{\Sigma}$ / $ASL\lambda_{\Sigma}$  the broader spatial pattern will be nearly identical due to the high correlations of these coefficient pairs over the data period (Fig. S1-S2) and so we just explore in detail the impact of estimating the ASL coefficients at one point location per glacier, at a centerline location about 20 km upstream of their respective grounding lines (Fig. S5 yellow crosses; Table S1).

The detrended data are shown in Fig. 3 (top row) where they reveal non-linear variability of several meters over the data period (blue plusses). Time series of estimated ASL coefficients sum to closely reproduce the data (black line). These two terms explain 84% (Thwaites) and 90% (Pine Island) of the partial variance of the altimeter time series. Interestingly, the phase of the  $ASLP_{\Sigma}$  term is opposite between Thwaites and Pine Island, while the  $ASL\lambda_{\Sigma}$  term is in phase.

Neither of the FDM models can explain the elevation variability at Thwaites or Pine Island glaciers (Fig. S9, brown lines). This could be because the SMB models are unable to reproduce the precipitation in this region, especially in ~2007 at Thwaites Glacier, but this would require a highly localized signal as this event does not occur at Pine Island Glacier. The misfit could be caused by errors in background altimeter models, however we note we obtain nearly identical results using the alternative dataset of Schröder et al. (2019). The most likely source of the unexplained height signal is ice flow dynamics responding to large-scale climate variability.

The dynamic effect of ice flow and its influence on ice sheet mass and surface elevation at a given point can be estimated from satellite-derived glacier velocities and the principle of mass conservation (Supplementary Text S1). Based on year-on-year changes in ice velocity since 2003, it is reasonable to expect several meters of dynamic elevation change in the lower parts of Pine Island and Thwaites due to a combination of advection and strain thinning (Fig. S10).



**Figure 3.** Detrended elevation time series at glacier point locations. Time series are shown for sites ~20 km upstream of the grounding line and along the centerline of flow (Fig. S5 yellow crosses; Table S1). Shown are the altimeter time series after 10 km Gaussian smoothing and subtracting the estimated trend and harmonics (blue plusses), and the two components of the model (colored lines) and their sum (black line) for each glacier. For Thwaites and Pine Island glaciers (top row), ASL coefficients are shown, while for Totten and Denman glaciers (bottom row) SAM and ENSO terms are shown. The partial variances explained by the sum of the two coefficients are listed in each panel. Grey shading is the 1-sigma uncertainty of the model. Error bars represent the 2-sigma uncertainties of the data.

### 3.2.2 Totten and Denman glaciers

The SAM and ENSO coefficients in the region of Totten and Denman glaciers have smaller magnitude and are much more diffuse than in the ASE (Fig. S5d-e). Nonetheless, these terms explain significant amounts of the partial variance (Fig. S5f) in this region. There is almost no non-linear signal to explain near the front of the Denman Glacier (Fig. 3), with the largest SAM or ENSO signal in the Denman region is west of Denman. Nonetheless, SAM contributes about 30% of the partial variance at Denman. If the underlying surface lowering trend of Denman is affected by climate variability it is not obviously associated with SAM and ENSO over this period.

Despite the modest signal near Totten there is still evidence that significant SAM and ENSO signals exist in the fast-flowing region of Totten Glacier (Fig. 3), at least in the 20-30 km above the grounding zone (Fig. S7c, Fig. S8). Unlike the ASE glaciers, there is insufficient ice velocity time series for Totten Glacier to explore the cumulative impacts of time-varying ice dynamics on ice elevation. As noted above, the FDM-corrected results are model-dependent in this region and so the origin(s) of the Totten Glacier non-linear elevation change signal is unclear but may contain a component due to ice dynamic changes.

## 4 Discussion

Our analysis reveals the spatial fingerprints of SAM and ENSO on AIS elevation over 2002-2021, patterns which are confirmed by analysis of GRACE mass change data over the same period. These patterns may not be stationary with time. Indeed, circulation patterns associated with SAM are known to vary over decades (Marshall et al., 2013; Silvestri & Vera, 2009), with effects including variable precipitation in the Antarctic Peninsula (Goodwin et al., 2016). Within this context it is therefore not unexpected that our pattern of SAM variability is different to the SMB-only SAM reconstruction of Medley and Thomas (2019) for the second half of the 20<sup>th</sup> century for instance. Differences with SMB-only reconstructions would also result if ice-dynamic effects on ice elevation and mass were non-negligible as hinted at by our data.

There are only a few previous studies exploring the relationship between ice dynamics, expressed as changes in ice mass, thickness, or elevation, and modes of climate variability, most notably in the Amundsen Sea Embayment region (Christie et al., 2023). In particular, Dutrieux et al. (2014) found reduced PIG ice shelf melt during a strong 2012 La Niña. Consistent with this, Paolo et al. (2018) found PIG ice shelf melting increased during El Niño, reducing ice shelf thickness, but that the ice shelf elevation increased overall due to increased accumulation. Our finding that PIG increases in elevation upstream of its grounding line when El Niño is sustained (or sustained westward ASL position), while the glacier is also dynamically thinning (Fig. S10), is consistent with this overall picture.

The SAM/ASLP-related signal upstream of PIG, Thwaites, and other ASE glaciers is the largest unexplained signal in Antarctica. The spatial pattern, with largest signal at lowest elevations, could be explained by both ice dynamics or unmodeled SMB or firn densification. Limited idealized study of the impacts of SAM on ASE basal melt is consistent with our observation of reduced upstream elevation with positive SAM but with melt response times that are decades longer than our analysis explores (Verfaillie et al., 2022), perhaps ruling out SAM but leaving the possibility of the localized ASLP as an source of immediate changes in buttressing.

We note that while the  $SAM_{\Sigma}$  and  $ASLP_{\Sigma}$  signals are correlated and our analysis cannot separate their different effects, they have different long-term implications for the ice sheet. As discussed by King et al. (2023),  $SAM_{\Sigma}$  has a trend due to the positive phase of SAM that has emerged since the 1940s.  $ASLP_{\Sigma}$  does not have a strong long-term trend, and so the extent to which the changes in coastal West Antarctica are related to the ASL rather than SAM will reduce the inferred contribution of SAM to ice-mass loss over recent decades (King et al., 2023).

Finally, our findings offer a simple way to remove decadal-scale variability from altimetry time series. This reduces correlated noise in the time series and will alter both the derived trends and, perhaps most significantly, the uncertainties of derived trends and other parameters if correlated noise is considered in the regression as it should (Ferguson et al., 2004; King & Watson, 2020; Williams et al., 2014; Wouters et al., 2013).

## 5 Conclusions

We analyzed gridded Antarctic ice elevation time series and show that much of the time series variance can be explained through a simple linear model based on the cumulative indices of the Southern Annular Mode and El Nino Southern Oscillation. The spatial pattern of this signal, once spatially smoothed, is in close agreement with the spatial pattern evident in GRACE data suggesting that observed ice elevation variability is robust and climatological. The Amundsen Sea Low is more directly relevant to the Amundsen Sea Embayment and we show that variations in its pressure and longitude explain ~90% of the variance over Pine Island and Thwaites glaciers.

Subtracting the effects of modeled firn densification removes much, but not all, signal, with inter-model differences evident. Residual climatological signal is particularly large at the fronts of fast-flowing glaciers in the Amundsen Sea Embayment. We suggest that ice dynamic effects may be contributing to this signal. Computing changes in elevation due to observed variation in horizontal velocity suggests the velocities are potential of the right magnitude to explain it. Further work is required to quantify the magnitude and response-times of upstream ice to changes in climatological variability in ice shelf melt.

## Acknowledgments

This work was supported by the ARC Australian Centre for Excellence in Antarctic Science (Project ID SR200100008). We thank the individuals and groups sharing altimetry and GRACE data, firn densification models, and climate indices.

## Open Research

## Data Availability Statement

All underlying data are openly available. Altimetry data were obtained from <https://doi.org/10.5067/L3LSVDZS15ZV>. GRACE time series were obtained from <http://gravis.gfz-potsdam.de/ais>. The IMAU FDM was obtained from <https://doi.org/10.5281/zenodo.5172513>. The GSFC FDM was obtained from <https://zenodo.org/record/7054574#.Y0iiTnbMJPY>. SAM index time series were obtained from <http://www.nerc-bas.ac.uk/public/icd/gjma/newsam.1957.2007.txt>. ASL data were obtained from [https://scotthosking.com/asl\\_index](https://scotthosking.com/asl_index) using version 3.20210820-era5. Nino3.4 index time series were obtained from [https://psl.noaa.gov/gcos\\_wgsp/Timeseries/Data/nino34.long.anom.data](https://psl.noaa.gov/gcos_wgsp/Timeseries/Data/nino34.long.anom.data). Data presented here will be made openly available upon acceptance.

## References

- Bos, M. S., Fernandes, R. M. S., Williams, S. D. P., & Bastos, L. (2013). Fast error analysis of continuous GNSS observations with missing data. *Journal of Geodesy*, 87(4), 351-360. doi:10.1007/s00190-012-0605-0
- Christie, F. D. W., Steig, E. J., Gourmelen, N., Tett, S. F. B., & Bingham, R. G. (2023). Inter-decadal climate variability induces differential ice response along Pacific-facing West Antarctica. *Nature Communications*, 14(1), 93. doi:10.1038/s41467-022-35471-3
- Clem, K. R., Renwick, J. A., & McGregor, J. (2017). Large-Scale Forcing of the Amundsen Sea Low and Its Influence on Sea Ice and West Antarctic Temperature. *Journal of Climate*, 30(20), 8405-8424. doi:10.1175/JCLI-D-16-0891.1
- Dahle, C., & Murböck, M. (2020). Post-processed GRACE/GRACE-FO Geopotential GSM Coefficients COST-G RL01 (Level-2B Product). doi:10.5880/GFZ.GRAVIS\_06\_L2B
- Diener, T., Sasgen, I., Agosta, C., Fürst, J. J., Braun, M. H., Konrad, H., & Fettweis, X. (2021). Acceleration of Dynamic Ice Loss in Antarctica From Satellite Gravimetry. *Frontiers in Earth Science*, 9. Original Research. doi:10.3389/feart.2021.741789
- Dutrieux, P., De Rydt, J., Jenkins, A., Holland, P. R., Ha, H. K., Lee, S. H., et al. (2014). Strong Sensitivity of Pine Island Ice-Shelf Melting to Climatic Variability. *Science*, 343(6167), 174-178. doi:10.1126/science.1244341
- Ferguson, A. C., Davis, C. H., & Cavanaugh, J. E. (2004). An autoregressive model for analysis of ice sheet elevation change time series. *Ieee Transactions on Geoscience and Remote Sensing*, 42(11), 2426-2436. Article. doi:10.1109/tgrs.2004.836788
- Goodwin, B. P., Mosley-Thompson, E., Wilson, A. B., Porter, S. E., & Sierra-Hernandez, M. R. (2016). Accumulation Variability in the Antarctic Peninsula: The Role of Large-Scale Atmospheric Oscillations and Their Interactions. *Journal of Climate*, 29(7), 2579-2596. doi:10.1175/JCLI-D-15-0354.1
- Hansen, N., Langen, P. L., Boberg, F., Forsberg, R., Simonsen, S. B., Thejll, P., et al. (2021). Downscaled surface mass balance in Antarctica: impacts of subsurface processes and large-scale atmospheric circulation. *The Cryosphere*, 15(9), 4315-4333. doi:10.5194/tc-15-4315-2021
- Horwath, M., Legresy, B., Remy, F., Blarel, F., & Lemoine, J. M. (2012). Consistent patterns of Antarctic ice sheet interannual variations from ENVISAT radar altimetry and GRACE satellite gravimetry. *Geophysical Journal International*, 189(2), 863-876. doi:10.1111/j.1365-246X.2012.05401.x
- Hosking, J. S., Orr, A., Bracegirdle, T. J., & Turner, J. (2016). Future circulation changes off West Antarctica: Sensitivity of the Amundsen Sea Low to projected anthropogenic forcing. *Geophysical Research Letters*, 43(1), 367-376. doi:10.1002/2015GL067143
- Kaitheri, A., Mémin, A., & Rémy, F. (2021). Inter-Annual Variability in the Antarctic Ice Sheets Using Geodetic Observations and a Climate Model. *Remote Sensing*, 13(11), 2199.

- Kim, B.-H., Seo, K.-W., Eom, J., Chen, J., & Wilson, C. R. (2020). Antarctic ice mass variations from 1979 to 2017 driven by anomalous precipitation accumulation. *Scientific Reports*, 10(1), 20366. doi:10.1038/s41598-020-77403-5
- King, M. A., Lyu, K., & Zhang, X. (2023). Climate variability a key driver of recent Antarctic ice-mass change. *Nature Geoscience*, 16(12), 1128-1135. doi:10.1038/s41561-023-01317-w
- King, M. A., & Watson, C. S. (2020). Antarctic Surface Mass Balance: Natural Variability, Noise, and Detecting New Trends. *Geophysical Research Letters*, 47(12), e2020GL087493. doi:10.1029/2020GL087493
- Marshall, G. J. (2003). Trends in the Southern Annular Mode from Observations and Reanalyses. *Journal of Climate*, 16(24), 4134-4143. doi:10.1175/1520-0442(2003)016<4134:Titsam>2.0.Co;2
- Marshall, G. J., Orr, A., & Turner, J. (2013). A Predominant Reversal in the Relationship between the SAM and East Antarctic Temperatures during the Twenty-First Century. *Journal of Climate*, 26(14), 5196-5204. doi:10.1175/Jcli-D-12-00671.1
- Medley, B., Neumann, T. A., Zwally, H. J., Smith, B. E., & Stevens, C. M. (2022). Simulations of firn processes over the Greenland and Antarctic ice sheets: 1980–2021. *The Cryosphere*, 16(10), 3971-4011. doi:10.5194/tc-16-3971-2022
- Medley, B., & Thomas, E. R. (2019). Increased snowfall over the Antarctic Ice Sheet mitigated twentieth-century sea-level rise. *Nature Climate Change*, 9(1), 34-39. doi:10.1038/s41558-018-0356-x
- Mémin, A., Flament, T., Alizier, B., Watson, C., & Rémy, F. (2015). Interannual variation of the Antarctic Ice Sheet from a combined analysis of satellite gravimetry and altimetry data. *Earth and Planetary Science Letters*, 422(0), 150-156. doi:10.1016/j.epsl.2015.03.045
- Mémin, A., Flament, T., Rémy, F., & Llubes, M. (2014). Snow- and ice-height change in Antarctica from satellite gravimetry and altimetry data. *Earth and Planetary Science Letters*, 404(0), 344-353. doi:10.1016/j.epsl.2014.08.008
- Mottram, R., Hansen, N., Kittel, C., van Wessem, J. M., Agosta, C., Amory, C., et al. (2021). What is the surface mass balance of Antarctica? An intercomparison of regional climate model estimates. *The Cryosphere*, 15(8), 3751-3784. doi:10.5194/tc-15-3751-2021
- Nilsson, J., Gardner, A., & Paolo, F. S. (2023). *MEaSUREs ITS LIVE Antarctic Grounded Ice Sheet Elevation Change, Version 1*. Retrieved from: <https://nsidc.org/data/NSIDC-0782/versions/1> doi:10.5067/L3LSVDZS15ZV
- Nilsson, J., Gardner, A. S., & Paolo, F. S. (2022). Elevation change of the Antarctic Ice Sheet: 1985 to 2020. *Earth Syst. Sci. Data*, 14(8), 3573-3598. doi:10.5194/essd-14-3573-2022
- Otosaka, I. N., Shepherd, A., Ivins, E. R., Schlegel, N. J., Amory, C., van den Broeke, M. R., et al. (2023). Mass balance of the Greenland and Antarctic ice sheets from 1992 to 2020. *Earth Syst. Sci. Data*, 15(4), 1597-1616. doi:10.5194/essd-15-1597-2023
- Palóczy, A., Gille, S. T., & McClean, J. L. (2018). Oceanic Heat Delivery to the Antarctic Continental Shelf: Large-Scale, Low-Frequency Variability. *Journal of Geophysical Research: Oceans*, 123(11), 7678-7701. doi:10.1029/2018JC014345
- Paolo, F. S., Padman, L., Fricker, H. A., Adusumilli, S., Howard, S., & Siegfried, M. R. (2018). Response of Pacific-sector Antarctic ice shelves to the El Niño/Southern Oscillation. *Nature Geoscience*, 11(2), 121-126. doi:10.1038/s41561-017-0033-0
- Rayner, N. A. (2003). Global analyses of sea surface temperature, sea ice, and night marine air temperature since the late nineteenth century. *Journal of Geophysical Research*, 108(D14). doi:10.1029/2002jd002670
- Sasgen, I., Groh, A., & Horwath, M. (2020). *COST-G GravIS RL01 Ice-Mass Change Products*. doi:10.5880/COST-G.GRAVIS\_01\_L3\_ICE
- Schröder, L., Horwath, M., Dietrich, R., Helm, V., van den Broeke, M. R., & Ligtenberg, S. R. M. (2019). Four decades of Antarctic surface elevation changes from multi-mission satellite altimetry. *The Cryosphere*, 13(2), 427-449. doi:10.5194/tc-13-427-2019
- Shepherd, A., Ivins, E. R., A, G., Barletta, V. R., Bentley, M. J., Bettadpur, S., et al. (2012). A Reconciled Estimate of Ice-Sheet Mass Balance. *Science*, 338(6111), 1183-1189. doi:10.1126/science.1228102
- Silvestri, G., & Vera, C. (2009). Nonstationary Impacts of the Southern Annular Mode on Southern Hemisphere Climate. *Journal of Climate*, 22(22), 6142-6148. doi:10.1175/2009JCLI3036.1
- Smith, B., Fricker, H. A., Gardner, A. S., Medley, B., Nilsson, J., Paolo, F. S., et al. (2020). Pervasive ice sheet mass loss reflects competing ocean and atmosphere processes. *Science*, 368(6496), 1239-1242. doi:10.1126/science.aaz5845

- Spence, P., Holmes, R. M., Hogg, A. M., Griffies, S. M., Stewart, K. D., & England, M. H. (2017). Localized rapid warming of West Antarctic subsurface waters by remote winds. *Nature Climate Change*, 7(8), 595-603. doi:10.1038/nclimate3335
- Thomas, E. R., van Wessem, J. M., Roberts, J., Isaksson, E., Schlosser, E., Fudge, T. J., et al. (2017). Regional Antarctic snow accumulation over the past 1000 years. *Clim. Past*, 13(11), 1491-1513. doi:10.5194/cp-13-1491-2017
- Turner, J., Phillips, T., Hosking, J. S., Marshall, G. J., & Orr, A. (2013). The Amundsen Sea low. *International Journal of Climatology*, 33(7), 1818-1829. doi:10.1002/joc.3558
- van den Broeke, M. R., & van Lipzig, N. P. M. (2017). Changes in Antarctic temperature, wind and precipitation in response to the Antarctic Oscillation. *Annals of Glaciology*, 39, 119-126. doi:10.3189/172756404781814654
- Veldhuijsen, S. B. M., van de Berg, W. J., Brils, M., Kuipers Munneke, P., & van den Broeke, M. R. (2023). Characteristics of the 1979–2020 Antarctic firm layer simulated with IMAU-FDM v1.2A. *The Cryosphere*, 17(4), 1675-1696. doi:10.5194/tc-17-1675-2023
- Verfaillie, D., Pelletier, C., Goosse, H., Jourdain, N. C., Bull, C. Y. S., Dalaiden, Q., et al. (2022). The circum-Antarctic ice-shelves respond to a more positive Southern Annular Mode with regionally varied melting. *Communications Earth & Environment*, 3(1), 139. doi:10.1038/s43247-022-00458-x
- Wahr, J., Molenaar, M., & Bryan, F. (1998). Time variability of the Earth's gravity field: Hydrological and oceanic effects and their possible detection using GRACE. *Journal of Geophysical Research: Solid Earth*, 103(B12), 30205-30229. doi:10.1029/98JB02844
- Wang, B. (1992). The Vertical Structure and Development of the Enso Anomaly Mode during 1979-1989. *Journal of the Atmospheric Sciences*, 49(8), 698-712. doi:10.1175/1520-0469(1992)049<0698:Tvsado>2.0.Co;2
- Williams, S. D. P., Moore, P., King, M. A., & Whitehouse, P. L. (2014). Revisiting GRACE Antarctic ice mass trends and accelerations considering autocorrelation. *Earth and Planetary Science Letters*, 385, 12-21. doi:10.1016/j.epsl.2013.10.016
- Wingham, D. J., Wallis, D. W., & Shepherd, A. (2009). Spatial and temporal evolution of Pine Island Glacier thinning, 1995-2006. *Geophysical Research Letters*, 36, 5. Article. doi:L1750, 10.1029/2009gl039126
- Wouters, B., Bamber, J. L., van den Broeke, M. R., Lenaerts, J. T. M., & Sasgen, I. (2013). Limits in detecting acceleration of ice sheet mass loss due to climate variability. *Nature Geoscience*, 6(8), 613-616. doi:10.1038/ngeo1874

## References From the Supporting Information

- Haran, T., Bohlander, J., Scambos, T., & Fahnestock, M. (2014, updated 2019). *MODIS mosaic of Antarctica 2008-2009 (MOA2009) Image Map, Version 1*. doi:10.7265/N5KP8037
- Howat, I. M. (2022). *The Reference Elevation Model of Antarctica – Mosaics, Version 2*. doi:10.7910/DVN/EBW8UC
- Rignot, E., Mouginot, J., & Scheuchl, B. (2017). *MEaSURES InSAR-Based Antarctica Ice Velocity Map, Version 2*. doi:10.5067/D7GK8F5J8M8R

**Major modes of climate variability dominate nonlinear Antarctic ice-sheet elevation changes 2002-2020**

**Matt A. King<sup>1,2</sup>, Poul Christoffersen<sup>2,3</sup>**

<sup>1</sup>Surveying and Spatial Sciences, School of Geography, Planning, and Spatial Sciences, University of Tasmania, Hobart, Tasmania 7001, Australia

<sup>2</sup>The Australian Centre for Excellence in Antarctic Science, University of Tasmania, Hobart, Tasmania 7001, Australia

<sup>3</sup>Institute for Marine and Antarctic Studies, University of Tasmania, Hobart, Tasmania 7001, Australia

Corresponding author: Matt King ([Matt.King@utas.edu.au](mailto:Matt.King@utas.edu.au))

**Key Points:**

- Cumulative effects of large-scale climate modes dominate detrended altimeter time series of Antarctic ice elevation 2002-2020
- These decadal signals have the same spatial pattern in altimeter ice height and GRACE mass time series.
- These decadal signals are largely due to surface mass balance, but ice dynamic changes may play a role in the Amundsen Sea Embayment

## Abstract

We explore the links between elevation variability of the Antarctic Ice Sheet (AIS) and large-scale climate modes. Using multiple linear regression, we quantify the cumulative effects of El Niño Southern Oscillation (ENSO) and the Southern Annular Mode (SAM) on gridded AIS elevations. Cumulative ENSO and SAM explain a median of 29% of the partial variance and up to 85% in some coastal areas. After spatial smoothing, these signals have high spatial correlation with those from GRACE gravimetry ( $r \sim 0.65$  each). Much of the signal is removed by a model of firn densification but inter-model differences exist especially for ENSO. At the lower parts of the Thwaites and Pine Island glaciers, near their grounding line, we find the Amundsen Sea Low (ASL) explains  $\sim 90\%$  of the observed elevation variability. There, firn effects explain only a small fraction of the variability, suggesting significant height changes have a climatological ice-dynamic response.

## Plain Language Summary

This study investigates how variations in the height of the Antarctic Ice Sheet (AIS) are connected to large-scale climate patterns. We used a statistical method to measure the effects of two climate phenomena: El Niño Southern Oscillation (ENSO) and the Southern Annular Mode (SAM). We found that the cumulative effects of these phenomena account for about 29% of the variations in AIS height on average, and up to 85% in some coastal areas. These patterns match well with independent data from the GRACE satellites over the same period. Applying a model that considers the compacting of snow into ice (firn densification) removes much of this signal, suggesting much, but not all, of the signals are related to snowfall variations. At the fronts of the rapidly changing Thwaites and Pine Island glaciers, the dominant climate phenomenon is the Amundsen Sea Low (ASL), which varies in strength and location. Here, the cumulative effects of the ASL changes explain about 90% of the variations in height of these glaciers, with only a small part explained by firn effects. We suggest the unexplained variability is at least partly due to changes in ice flow.

## 1 Introduction

Observations of the changing volume of the Antarctic Ice Sheet play a major role in understanding ice-sheet change (e.g., Otonari et al., 2023; Shepherd et al., 2012) from the whole-of-ice-sheet down to individual glaciers (e.g., Smith et al., 2020; Wingham et al., 2009). The now three-decade record of continuous ice volume change captures the variability and longer-term change of both surface mass balance (SMB), and related firn processes, and elevation effects of changing ice dynamics. These changes are, respectively, related to atmospheric and oceanic processes (Horwath et al., 2012; Smith et al., 2020). Several studies have examined the relationship between ice height changes and modes of climate variability, in particular linking them to both El Niño - Southern Oscillation (ENSO) and the Antarctic Circumpolar Wave (Kaitheri et al., 2021; Mémin et al., 2015; Mémin et al., 2014).

Strangely, less studied in this context is the role of the dominant mode of climate variability in the Southern Hemisphere, the Southern Annular Mode (SAM). Despite SAM driving variability and trends in SMB across a wide range of timescales (Diener et al., 2021; Medley & Thomas,

2019; van den Broeke & van Lipzig, 2017), SAM has yet to be linked to observations of ice sheet elevation change, with one related study reporting no correlation to estimates of ice shelf elevation change (Paolo et al., 2018). By contrast, the cumulative sum of SAM (Diener et al., 2021; Kim et al., 2020) has recently been shown to be linearly related to the dominant signal in detrended surface mass time series derived from satellite gravimetry (King et al., 2023), with large-scale spatially-coherent signal across coastal regions at decadal timescales.

The ~300 km spatial resolution of satellite gravimetry, combined with uncertainties in models of SMB (Mottram et al., 2021), meant that King et al. (2023) were not able to separate the relative contributions of SMB and ice dynamical change forced respectively by the atmosphere and ocean (Hansen et al., 2021; Kim et al., 2020; Palóczy et al., 2018; Spence et al., 2017; Thomas et al., 2017; Verfaillie et al., 2022). In particular, ice dynamical change will have a distinct spatial pattern compared to SMB that is not detectable by GRACE but could be possible with altimetry (Smith et al., 2020). Detecting (or otherwise) a response of the grounded ice sheet to large-scale climate variability via the oceans and ice shelves would provide important insights into ice-sheet sensitivity to climate change.

In this paper we analyze a recent gridded compilation of satellite altimeter data and compare these time series to cumulative climate indices. We compare the derived signals to those from space gravimetry and then, taking advantage of the high-resolution altimeter data, explore the signal over key ice streams: Thwaites, Pine Island, Totten, and Denman.

## 2 Datasets and Analysis

### 2.1 Altimeter dataset

We make use of a gridded altimeter product (Nilsson et al., 2023) at 1920 m horizontal resolution and covering the period from Apr 1985 to Dec 2020 (Nilsson et al., 2022). We spatially down-sample this to 5 km horizontal resolution. To facilitate comparison with space gravimetry data we only make use of data from 2002 to the end of the record. The dataset contains monthly ice-sheet elevation-change data derived from a range of radar and laser altimeter missions; over the study period these are ERS-2, Envisat and CryoSat-2 and ICESat and ICESat-2. The approach to accounting for differences in reflection surfaces and other systematic effects is described by Nilsson et al. (2022). To reduce spatial noise we apply a Gaussian smoother with widths specified below, with width defined at the half height of the function, consistent with the definition commonly used in GRACE data smoothing (Wahr et al., 1998).

### 2.2 Space gravimetry dataset

We use the COST-G RL01 Level-3 50 km gridded GRACE and GRACE-FO V0002 dataset obtained from <http://gravis.gfz-potsdam.de/antarctica> (Sasgen et al., 2020). We make use of data from Mar 2002 to Dec 2020, with the end point chosen to match the end of the altimetry dataset.

The data are spaced approximately monthly and with a data gap of ~12 months between GRACE and GRACE-FO from mid-2017 to mid-2018.

We note that while this product is gridded at 50 km, the intrinsic GRACE resolution is 200-300 km. Post-processing steps include replacement of low-degree GRACE coefficients and insertion of degree-1 terms using standard approaches (Dahle & Murböck, 2020; Sasgen et al., 2020).

Since we are interested in decadal variability and trends, we also lightly smooth the altimetry and GRACE data with a Gaussian filter with width 7 months (Wahr et al., 1998).

### 2.3 Climate indices

We compare the altimeter and GRACE data primarily with SAM and ENSO indices, with additional comparison to Amundsen Sea Low (ASL) indices in the Amundsen Sea region. For the ASL indices, we make use of both the absolute center pressure (ASLP) and longitude (ASL $\lambda$ ) within the ASL Index version 3.20210820-era5 based on monthly ERA5 reanalysis data (Hosking et al., 2016). For SAM, we make use of the Marshall station index (Marshall, 2003). For ENSO, we make use of the Nino3.4 index based on the HadISST1 dataset (Rayner, 2003) and use a 6-month lag (King et al., 2023; Paolo et al., 2018). We normalized each index with the mean and standard deviation computed over 1971-1999 inclusive, then cumulatively summed them, limited them to the data period, and then renormalized to produce SAM $_{\Sigma}$ , ENSO $_{\Sigma}$ , ASLP $_{\Sigma}$ , and ASL $\lambda_{\Sigma}$ .

The raw indices and their cumulative sums are shown in Fig S1. Correlations above 0.7 are evident between ASLP $_{\Sigma}$  and SAM $_{\Sigma}$  and between ASL $\lambda_{\Sigma}$  and ENSO $_{\Sigma}$  (Fig S1, S2). This is due to the ASL being affected by larger-scale modes of climate variability, with SAM in particular modulating its absolute pressure and ENSO modulating the longitude of its center (Clem et al., 2017; Hosking et al., 2016; Turner et al., 2013).

### 2.4 Multi-variate Empirical Orthogonal Functions

For a data-driven analysis we make use of Multi-variate Empirical Orthogonal Functions (MEOF) (Wang, 1992). MEOFs are an extension of conventional Empirical Orthogonal Functions but allow the dominant modes across multiple variables to be identified rather than treating each variable separately. We use MEOF to analyze the elevation and mass change gridded datasets after individual normalization. We first smooth the altimetry dataset with a 50 km-wide Gaussian smoother and sub-sample the altimeter dataset to match the 50 km horizontal resolution of GRACE. Given the limited sampling of altimetry in the northern Antarctic Peninsula we truncate that region from both datasets prior to computing MEOFs.

### 2.5 Regression

Using ordinary least squares, we solved the coefficients ( $a$ ,  $b$ ,  $c$ ,  $d$ , and  $e$ ) of the functional model describing time-evolving elevation ( $h$ ) with time ( $t$ ):

$$h(t_i) = a + b(t_i - t_0) + \sum_{k=1}^2 (c_k^s \sin(2\pi f_k t_i) + c_k^c \cos(2\pi f_k t_i)) + dSAM_{\Sigma} + eENSO_{\Sigma} \quad (1)$$

Where  $f_k = [1, 2]$  cycles per year. We adopted  $t_0$  as the mid point of the altimeter series.

## 2.6 Data uncertainty

For regression parameter uncertainties, we recognize the existence of temporal correlations in the altimeter time series (Ferguson et al., 2004), in part due to SMB variation (King & Watson, 2020), and take these into account. Following King et al. (2023), we compared trend uncertainties from a linear regression using a Generalized Gauss Markov noise model to those generated using a white noise only (temporally uncorrelated) noise model using HECTOR v2.0 software (Bos et al., 2013). For regressions that included the SAM and ENSO terms, the white noise only model produced uncertainties a factor of 3 too small, taken as the median of the ratio of trend uncertainties, or factor 40 too small when not including the SAM and ENSO terms. We applied these scale factors to the uncertainties from the regression. For the GRACE uncertainties we used the scale factors of King et al. (2023).

## 3 Results

### 3.1 Ice-sheet scale analysis

Our data-driven MEOF analysis shows that ice elevation and mass time series are both dominated by decadal-scale variability (Fig. S3c, f). Together, the two leading modes explain 65% of the non-linear variance of the combined and smoothed time series. Their corresponding principal components (PCs) correlate with detrended  $\text{SAM}_\Sigma$  ( $r=0.73$ ) and 6-month lagged  $\text{ENSO}_\Sigma$  ( $r=0.89$ ). The  $\text{ASLP}_\Sigma$  and  $\text{ASL}\lambda_\Sigma$  terms are not of direct relevance at the ice-sheet scale given the limited geographical footprint of influence of the ASL, but also have high correlations with the data.

GRACE and altimetry MEOFs have a high spatial correlation (Fig. S3a-b, d-e;  $r=0.87$  for MEOF1 and  $r=0.75$  for MEOF2) suggesting they are sensing the same signal and are both dominated by coastal changes. The potential in the high-resolution altimetry record is particularly evident in MEOF1 where the spatially-diffuse signal in GRACE (Fig. S3a) is shown to be concentrated over small regions that coincide with the major ice streams of the Amundsen Sea Embayment and the coastline of the Bellingshausen Sea and Marie Byrd Land (Fig. S3b). We note that while MEOF3 (Fig. S4) is partly affected by striping in the GRACE field, characteristic of GRACE systematic error, coherent signal is evident between GRACE and altimetry along the coastlines of the Bellingshausen Sea, Marie Byrd Land and Wilkes Land, suggesting the signal is robust in those regions, although the variance explained (5%) is much smaller than MEOFs 1 and 2. A similar signal to PC3, with periodicities of ~4-7 years, has also been identified in analysis of GRACE data (King et al., 2023; Mémmin et al., 2015). Beyond MEOF3, the modes explain little variance (<4%) and are dominated by noise, at least for GRACE (Fig. S4d).

To quantify the SAM and ENSO contribution to ice sheet elevation change we regress the altimetry time series against  $\text{SAM}_\Sigma$  and  $\text{ENSO}_\Sigma$  and the other parameters in Eq. 1. Here we use the gridded data after applying a 10 km Gaussian spatial filter. The 5 km gridded altimeter regression analysis shown in Fig. 1a,b reveals large-scale spatially coherent signal relating to SAM and ENSO around the coasts of Antarctica. Together, these two terms often explain more than 40% of the partial variance of the timeseries around the coast and into the interior, with the


partial variance controlling for the other regression terms. The median partial-variance explained across the ice sheet is 29% (Fig. 1c). The  $\text{SAM}_{\Sigma}$  coefficient is strongest in the Amundsen Sea Embayment where it centers on the Pine Island, Thwaites, Smith, and Pope Glaciers (Fig. S5a). The negative elevation signal in this region is linked to periods where positive SAM dominates negative SAM (positive  $\text{SAM}_{\Sigma}$ ). Other strong signal exists along the coastal zone of the Bellingshausen Sea, Marie Byrd Land, and parts of coastal East Antarctica. A more diffuse signal is evident in the interior of West Antarctica and parts of East Antarctica (Fig. S6a). The  $\text{ENSO}_{\Sigma}$  coefficient has particularly high positive values, indicating elevation increase associated with sustained El Niño, along the coast of the Bellingshausen Sea and well upstream into Pine Island Glacier (Fig. S5b)

Applying a 200 km Gaussian smoother to the altimeter data and rerunning the regression (Fig. 1d-e) produces coefficients with large-scale spatial coherence and larger partial variances explained, often exceeding 60% in key coastal regions but extending well into the interior of the ice sheet (Fig. 1f). Comparing them to results of a regression with GRACE data (Fig. 1g-h) (King et al., 2023) shows high agreement in the signs and spatial distribution of the signal. We note that there are insufficient altimeter data in the Northern Antarctic Peninsula to analyze the signal in this region. Computing spatial correlations between the smoothed altimetry regression and the GRACE regression gives  $r=0.65$  for  $\text{SAM}_{\Sigma}$  and  $r=0.68$  for  $\text{ENSO}_{\Sigma}$ .

We next examine the role of SMB variability on the estimated coefficients from the altimetry regression. To do this we subtract the IMAU Firn Densification Model (IMAU FDM) v1.2A (Veldhuijsen et al., 2023) from the altimetry time series and repeat the regression. The results are shown in Figure 2. Comparing Fig. 2a with Fig. 1a shows that IMAU FDM effectively removes all the SAM-related signal in East Antarctic Ice Sheet (EAIS) but much of the SAM signal remains in West Antarctic Ice Sheet (WAIS). Much of the coastal EAIS ENSO-related signal is removed by IMAU FDM but with small over-correction evident for much of the ice sheet, including signal reversing sign in George V Land and WAIS. Repeating the regression but instead using the GSFC FDM v1.2.1 (Medley et al., 2022) shows that there is significant sensitivity to the choice of FDM (Fig. 2d-f), with GSFC FDM apparently over-correcting ENSO-related signal in the Totten Glacier region but in much better agreement with the altimetry in WAIS. Given the decadal timescales of the signals, these inter-model differences are likely to have contributions from both the FDMs themselves and their underlying SMB models (Medley et al., 2022).


The combination of coefficients estimated from each of GRACE and altimetry allows the density of these terms to be estimated. Given the GRACE resolution half-width is about 100km, we computed densities and their at locations 100 km upstream of the grounding lines of the Thwaites, Pine Island, Totten, and Denman glacier. These computed densities are sensitive to the radius of the Gaussian smoother applied to the altimetry data, and we adopted a 200 km smoother to approximate the GRACE resolution. ENSO-related results are highly uncertain in the Denman and Totten glacier regions due to limited signal, but the other densities (SAM and ENSO related) suggest the observed changes have a density between snow and ice, clustering around  $600 \text{ kg/m}^3$  (Fig. S6). While the estimates are uncertain, they suggest that some of the signal could originate in ice dynamics rather than SMB.

223



224

**Figure 1.** Results of regression analysis of gridded data. Shown are the  $SAM_{\Sigma}$  and  $ENSO_{\Sigma}$  coefficients and variances explained for the altimetry (top row), altimetry after 200 km Gaussian smoothing (middle row), and GRACE (bottom row). The partial variances explained by  $SAM_{\Sigma}$  and lagged  $ENSO_{\Sigma}$  are in the right column.



**Figure 2.** Results of regression analysis of FDM-corrected gridded altimeter data. Regression coefficients are shown (left and central columns) and the partial variances explained by SAM<sub>Σ</sub> and lagged ENSO<sub>Σ</sub> (right column). Shown are the coefficients and variances explained for the altimetry time series after subtracting of the IMAU FDM (top row) and GSFC FDM (bottom row).

Next, we explore the origins of these signals further on a glacier-by-glacier basis.

### 3.2 Regional scale analysis

#### 3.2.1 Thwaites and Pine Island glaciers

The partial variance explained by the SAM<sub>Σ</sub> and ENSO<sub>Σ</sub> terms (before subtracting an FDM) is above 60% for much of the Amundsen Sea Embayment (ASE; Fig. 1c, f; S5c,f). Regardless of the FDM model adopted, much SAM<sub>Σ</sub> signal remains in the ASE broadly and ENSO<sub>Σ</sub> signal is evident in the Pine Island Glacier region (Fig. 2). Closer examination of these regions in Fig. S5

(top row) indicates that the ASE signals are concentrated along low-elevation and fast flowing regions that correspond to Pine Island, Thwaites, and nearby glaciers. This is further evidenced through cross-sections near to the front of these glaciers (Fig. S7) along the yellow lines in Fig S5. It is notable that the phase of the SAM-related signal is switched in the fast-flowing region of Pine Island Glacier.


Coefficient magnitudes generally decay upstream of the grounding line (Fig. S8). Subtracting the IMAU FDM before performing the regression results in coefficients along the centerline and cross profiles that are shifted nearly uniformly but are not significantly altered in their spatial pattern (dashed lines Fig. S7-S8). Together these results suggest there may be an ice dynamic component within the  $SAM_{\Sigma}$  and  $ENSO_{\Sigma}$  coefficients in addition to residual SMB/FDM signal.

Along the coastal margin of the ASE the climatology is more directly controlled by the ASL than SAM and ENSO which modulate its depth and location (Clem et al., 2017; Turner et al., 2013). To explore this further we repeated the regression replacing  $SAM_{\Sigma}$  and  $ENSO_{\Sigma}$  in Equation 1 with  $ASLP_{\Sigma}$  and  $ASL\lambda_{\Sigma}$ . While the magnitude of the estimated coefficients differs between  $SAM_{\Sigma}$ / $ASLP_{\Sigma}$  and  $ENSO_{\Sigma}$ / $ASL\lambda_{\Sigma}$  the broader spatial pattern will be nearly identical due to the high correlations of these coefficient pairs over the data period (Fig. S1-S2) and so we just explore in detail the impact of estimating the ASL coefficients at one point location per glacier, at a centerline location about 20 km upstream of their respective grounding lines (Fig. S5 yellow crosses; Table S1).

The detrended data are shown in Fig. 3 (top row) where they reveal non-linear variability of several meters over the data period (blue plusses). Time series of estimated ASL coefficients sum to closely reproduce the data (black line). These two terms explain 84% (Thwaites) and 90% (Pine Island) of the partial variance of the altimeter time series. Interestingly, the phase of the  $ASLP_{\Sigma}$  term is opposite between Thwaites and Pine Island, while the  $ASL\lambda_{\Sigma}$  term is in phase.

Neither of the FDM models can explain the elevation variability at Thwaites or Pine Island glaciers (Fig. S9, brown lines). This could be because the SMB models are unable to reproduce the precipitation in this region, especially in ~2007 at Thwaites Glacier, but this would require a highly localized signal as this event does not occur at Pine Island Glacier. The misfit could be caused by errors in background altimeter models, however we note we obtain nearly identical results using the alternative dataset of Schröder et al. (2019). The most likely source of the unexplained height signal is ice flow dynamics responding to large-scale climate variability.

The dynamic effect of ice flow and its influence on ice sheet mass and surface elevation at a given point can be estimated from satellite-derived glacier velocities and the principle of mass conservation (Supplementary Text S1). Based on year-on-year changes in ice velocity since 2003, it is reasonable to expect several meters of dynamic elevation change in the lower parts of Pine Island and Thwaites due to a combination of advection and strain thinning (Fig. S10).



**Figure 3.** Detrended elevation time series at glacier point locations. Time series are shown for sites ~20 km upstream of the grounding line and along the centerline of flow (Fig. S5 yellow crosses; Table S1). Shown are the altimeter time series after 10 km Gaussian smoothing and subtracting the estimated trend and harmonics (blue plusses), and the two components of the model (colored lines) and their sum (black line) for each glacier. For Thwaites and Pine Island glaciers (top row), ASL coefficients are shown, while for Totten and Denman glaciers (bottom row) SAM and ENSO terms are shown. The partial variances explained by the sum of the two coefficients are listed in each panel. Grey shading is the 1-sigma uncertainty of the model. Error bars represent the 2-sigma uncertainties of the data.

### 3.2.2 Totten and Denman glaciers

The SAM and ENSO coefficients in the region of Totten and Denman glaciers have smaller magnitude and are much more diffuse than in the ASE (Fig. S5d-e). Nonetheless, these terms explain significant amounts of the partial variance (Fig. S5f) in this region. There is almost no non-linear signal to explain near the front of the Denman Glacier (Fig. 3), with the largest SAM or ENSO signal in the Denman region is west of Denman. Nonetheless, SAM contributes about 30% of the partial variance at Denman. If the underlying surface lowering trend of Denman is affected by climate variability it is not obviously associated with SAM and ENSO over this period.

Despite the modest signal near Totten there is still evidence that significant SAM and ENSO signals exist in the fast-flowing region of Totten Glacier (Fig. 3), at least in the 20-30 km above the grounding zone (Fig. S7c, Fig. S8). Unlike the ASE glaciers, there is insufficient ice velocity time series for Totten Glacier to explore the cumulative impacts of time-varying ice dynamics on ice elevation. As noted above, the FDM-corrected results are model-dependent in this region and so the origin(s) of the Totten Glacier non-linear elevation change signal is unclear but may contain a component due to ice dynamic changes.

## 4 Discussion

Our analysis reveals the spatial fingerprints of SAM and ENSO on AIS elevation over 2002-2021, patterns which are confirmed by analysis of GRACE mass change data over the same period. These patterns may not be stationary with time. Indeed, circulation patterns associated with SAM are known to vary over decades (Marshall et al., 2013; Silvestri & Vera, 2009), with effects including variable precipitation in the Antarctic Peninsula (Goodwin et al., 2016). Within this context it is therefore not unexpected that our pattern of SAM variability is different to the SMB-only SAM reconstruction of Medley and Thomas (2019) for the second half of the 20<sup>th</sup> century for instance. Differences with SMB-only reconstructions would also result if ice-dynamic effects on ice elevation and mass were non-negligible as hinted at by our data.

There are only a few previous studies exploring the relationship between ice dynamics, expressed as changes in ice mass, thickness, or elevation, and modes of climate variability, most notably in the Amundsen Sea Embayment region (Christie et al., 2023). In particular, Dutrieux et al. (2014) found reduced PIG ice shelf melt during a strong 2012 La Niña. Consistent with this, Paolo et al. (2018) found PIG ice shelf melting increased during El Niño, reducing ice shelf thickness, but that the ice shelf elevation increased overall due to increased accumulation. Our finding that PIG increases in elevation upstream of its grounding line when El Niño is sustained (or sustained westward ASL position), while the glacier is also dynamically thinning (Fig. S10), is consistent with this overall picture.

The SAM/ASLP-related signal upstream of PIG, Thwaites, and other ASE glaciers is the largest unexplained signal in Antarctica. The spatial pattern, with largest signal at lowest elevations, could be explained by both ice dynamics or unmodeled SMB or firn densification. Limited idealized study of the impacts of SAM on ASE basal melt is consistent with our observation of reduced upstream elevation with positive SAM but with melt response times that are decades longer than our analysis explores (Verfaillie et al., 2022), perhaps ruling out SAM but leaving the possibility of the localized ASLP as an source of immediate changes in buttressing.

We note that while the  $SAM_{\Sigma}$  and  $ASLP_{\Sigma}$  signals are correlated and our analysis cannot separate their different effects, they have different long-term implications for the ice sheet. As discussed by King et al. (2023),  $SAM_{\Sigma}$  has a trend due to the positive phase of SAM that has emerged since the 1940s.  $ASLP_{\Sigma}$  does not have a strong long-term trend, and so the extent to which the changes in coastal West Antarctica are related to the ASL rather than SAM will reduce the inferred contribution of SAM to ice-mass loss over recent decades (King et al., 2023).

Finally, our findings offer a simple way to remove decadal-scale variability from altimetry time series. This reduces correlated noise in the time series and will alter both the derived trends and, perhaps most significantly, the uncertainties of derived trends and other parameters if correlated noise is considered in the regression as it should (Ferguson et al., 2004; King & Watson, 2020; Williams et al., 2014; Wouters et al., 2013).

## 5 Conclusions

We analyzed gridded Antarctic ice elevation time series and show that much of the time series variance can be explained through a simple linear model based on the cumulative indices of the Southern Annular Mode and El Nino Southern Oscillation. The spatial pattern of this signal, once spatially smoothed, is in close agreement with the spatial pattern evident in GRACE data suggesting that observed ice elevation variability is robust and climatological. The Amundsen Sea Low is more directly relevant to the Amundsen Sea Embayment and we show that variations in its pressure and longitude explain ~90% of the variance over Pine Island and Thwaites glaciers.

Subtracting the effects of modeled firn densification removes much, but not all, signal, with inter-model differences evident. Residual climatological signal is particularly large at the fronts of fast-flowing glaciers in the Amundsen Sea Embayment. We suggest that ice dynamic effects may be contributing to this signal. Computing changes in elevation due to observed variation in horizontal velocity suggests the velocities are potential of the right magnitude to explain it. Further work is required to quantify the magnitude and response-times of upstream ice to changes in climatological variability in ice shelf melt.

## Acknowledgments

This work was supported by the ARC Australian Centre for Excellence in Antarctic Science (Project ID SR200100008). We thank the individuals and groups sharing altimetry and GRACE data, firn densification models, and climate indices.

## Open Research

## Data Availability Statement

All underlying data are openly available. Altimetry data were obtained from <https://doi.org/10.5067/L3LSVDZS15ZV>. GRACE time series were obtained from <http://gravis.gfz-potsdam.de/ais>. The IMAU FDM was obtained from <https://doi.org/10.5281/zenodo.5172513>. The GSFC FDM was obtained from <https://zenodo.org/record/7054574#.Y0iiTnbMJPY>. SAM index time series were obtained from <http://www.nerc-bas.ac.uk/public/icd/gjma/newsam.1957.2007.txt>. ASL data were obtained from [https://scotthosking.com/asl\\_index](https://scotthosking.com/asl_index) using version 3.20210820-era5. Nino3.4 index time series were obtained from [https://psl.noaa.gov/gcos\\_wgsp/Timeseries/Data/nino34.long.anom.data](https://psl.noaa.gov/gcos_wgsp/Timeseries/Data/nino34.long.anom.data). Data presented here will be made openly available upon acceptance.

## References

- Bos, M. S., Fernandes, R. M. S., Williams, S. D. P., & Bastos, L. (2013). Fast error analysis of continuous GNSS observations with missing data. *Journal of Geodesy*, 87(4), 351-360. doi:10.1007/s00190-012-0605-0
- Christie, F. D. W., Steig, E. J., Gourmelen, N., Tett, S. F. B., & Bingham, R. G. (2023). Inter-decadal climate variability induces differential ice response along Pacific-facing West Antarctica. *Nature Communications*, 14(1), 93. doi:10.1038/s41467-022-35471-3
- Clem, K. R., Renwick, J. A., & McGregor, J. (2017). Large-Scale Forcing of the Amundsen Sea Low and Its Influence on Sea Ice and West Antarctic Temperature. *Journal of Climate*, 30(20), 8405-8424. doi:10.1175/Jcli-D-16-0891.1
- Dahle, C., & Murböck, M. (2020). Post-processed GRACE/GRACE-FO Geopotential GSM Coefficients COST-G RL01 (Level-2B Product). doi:10.5880/GFZ.GRAVIS\_06\_L2B
- Diener, T., Sasgen, I., Agosta, C., Fürst, J. J., Braun, M. H., Konrad, H., & Fettweis, X. (2021). Acceleration of Dynamic Ice Loss in Antarctica From Satellite Gravimetry. *Frontiers in Earth Science*, 9. Original Research. doi:10.3389/feart.2021.741789
- Dutrieux, P., De Rydt, J., Jenkins, A., Holland, P. R., Ha, H. K., Lee, S. H., et al. (2014). Strong Sensitivity of Pine Island Ice-Shelf Melting to Climatic Variability. *Science*, 343(6167), 174-178. doi:10.1126/science.1244341
- Ferguson, A. C., Davis, C. H., & Cavanaugh, J. E. (2004). An autoregressive model for analysis of ice sheet elevation change time series. *Ieee Transactions on Geoscience and Remote Sensing*, 42(11), 2426-2436. Article. doi:10.1109/tgrs.2004.836788
- Goodwin, B. P., Mosley-Thompson, E., Wilson, A. B., Porter, S. E., & Sierra-Hernandez, M. R. (2016). Accumulation Variability in the Antarctic Peninsula: The Role of Large-Scale Atmospheric Oscillations and Their Interactions. *Journal of Climate*, 29(7), 2579-2596. doi:10.1175/JCLI-D-15-0354.1
- Hansen, N., Langen, P. L., Boberg, F., Forsberg, R., Simonsen, S. B., Thejll, P., et al. (2021). Downscaled surface mass balance in Antarctica: impacts of subsurface processes and large-scale atmospheric circulation. *The Cryosphere*, 15(9), 4315-4333. doi:10.5194/tc-15-4315-2021
- Horwath, M., Legresy, B., Remy, F., Blarel, F., & Lemoine, J. M. (2012). Consistent patterns of Antarctic ice sheet interannual variations from ENVISAT radar altimetry and GRACE satellite gravimetry. *Geophysical Journal International*, 189(2), 863-876. doi:10.1111/j.1365-246X.2012.05401.x
- Hosking, J. S., Orr, A., Bracegirdle, T. J., & Turner, J. (2016). Future circulation changes off West Antarctica: Sensitivity of the Amundsen Sea Low to projected anthropogenic forcing. *Geophysical Research Letters*, 43(1), 367-376. doi:10.1002/2015GL067143
- Kaitheri, A., Mémin, A., & Rémy, F. (2021). Inter-Annual Variability in the Antarctic Ice Sheets Using Geodetic Observations and a Climate Model. *Remote Sensing*, 13(11), 2199.

- Kim, B.-H., Seo, K.-W., Eom, J., Chen, J., & Wilson, C. R. (2020). Antarctic ice mass variations from 1979 to 2017 driven by anomalous precipitation accumulation. *Scientific Reports*, 10(1), 20366. doi:10.1038/s41598-020-77403-5
- King, M. A., Lyu, K., & Zhang, X. (2023). Climate variability a key driver of recent Antarctic ice-mass change. *Nature Geoscience*, 16(12), 1128-1135. doi:10.1038/s41561-023-01317-w
- King, M. A., & Watson, C. S. (2020). Antarctic Surface Mass Balance: Natural Variability, Noise, and Detecting New Trends. *Geophysical Research Letters*, 47(12), e2020GL087493. doi:10.1029/2020GL087493
- Marshall, G. J. (2003). Trends in the Southern Annular Mode from Observations and Reanalyses. *Journal of Climate*, 16(24), 4134-4143. doi:10.1175/1520-0442(2003)016<4134:Titsam>2.0.Co;2
- Marshall, G. J., Orr, A., & Turner, J. (2013). A Predominant Reversal in the Relationship between the SAM and East Antarctic Temperatures during the Twenty-First Century. *Journal of Climate*, 26(14), 5196-5204. doi:10.1175/Jcli-D-12-00671.1
- Medley, B., Neumann, T. A., Zwally, H. J., Smith, B. E., & Stevens, C. M. (2022). Simulations of firn processes over the Greenland and Antarctic ice sheets: 1980–2021. *The Cryosphere*, 16(10), 3971-4011. doi:10.5194/tc-16-3971-2022
- Medley, B., & Thomas, E. R. (2019). Increased snowfall over the Antarctic Ice Sheet mitigated twentieth-century sea-level rise. *Nature Climate Change*, 9(1), 34-39. doi:10.1038/s41558-018-0356-x
- Mémin, A., Flament, T., Alizier, B., Watson, C., & Rémy, F. (2015). Interannual variation of the Antarctic Ice Sheet from a combined analysis of satellite gravimetry and altimetry data. *Earth and Planetary Science Letters*, 422(0), 150-156. doi:10.1016/j.epsl.2015.03.045
- Mémin, A., Flament, T., Rémy, F., & Llubes, M. (2014). Snow- and ice-height change in Antarctica from satellite gravimetry and altimetry data. *Earth and Planetary Science Letters*, 404(0), 344-353. doi:10.1016/j.epsl.2014.08.008
- Mottram, R., Hansen, N., Kittel, C., van Wessem, J. M., Agosta, C., Amory, C., et al. (2021). What is the surface mass balance of Antarctica? An intercomparison of regional climate model estimates. *The Cryosphere*, 15(8), 3751-3784. doi:10.5194/tc-15-3751-2021
- Nilsson, J., Gardner, A., & Paolo, F. S. (2023). *MEaSUREs ITS LIVE Antarctic Grounded Ice Sheet Elevation Change, Version 1*. Retrieved from: <https://nsidc.org/data/NSIDC-0782/versions/1> doi:10.5067/L3LSVDZS15ZV
- Nilsson, J., Gardner, A. S., & Paolo, F. S. (2022). Elevation change of the Antarctic Ice Sheet: 1985 to 2020. *Earth Syst. Sci. Data*, 14(8), 3573-3598. doi:10.5194/essd-14-3573-2022
- Otosaka, I. N., Shepherd, A., Ivins, E. R., Schlegel, N. J., Amory, C., van den Broeke, M. R., et al. (2023). Mass balance of the Greenland and Antarctic ice sheets from 1992 to 2020. *Earth Syst. Sci. Data*, 15(4), 1597-1616. doi:10.5194/essd-15-1597-2023
- Palóczy, A., Gille, S. T., & McClean, J. L. (2018). Oceanic Heat Delivery to the Antarctic Continental Shelf: Large-Scale, Low-Frequency Variability. *Journal of Geophysical Research: Oceans*, 123(11), 7678-7701. doi:10.1029/2018JC014345
- Paolo, F. S., Padman, L., Fricker, H. A., Adusumilli, S., Howard, S., & Siegfried, M. R. (2018). Response of Pacific-sector Antarctic ice shelves to the El Niño/Southern Oscillation. *Nature Geoscience*, 11(2), 121-126. doi:10.1038/s41561-017-0033-0
- Rayner, N. A. (2003). Global analyses of sea surface temperature, sea ice, and night marine air temperature since the late nineteenth century. *Journal of Geophysical Research*, 108(D14). doi:10.1029/2002jd002670
- Sasgen, I., Groh, A., & Horwath, M. (2020). *COST-G GravIS RL01 Ice-Mass Change Products*. doi:10.5880/COST-G.GRAVIS\_01\_L3\_ICE
- Schröder, L., Horwath, M., Dietrich, R., Helm, V., van den Broeke, M. R., & Ligtenberg, S. R. M. (2019). Four decades of Antarctic surface elevation changes from multi-mission satellite altimetry. *The Cryosphere*, 13(2), 427-449. doi:10.5194/tc-13-427-2019
- Shepherd, A., Ivins, E. R., A, G., Barletta, V. R., Bentley, M. J., Bettadpur, S., et al. (2012). A Reconciled Estimate of Ice-Sheet Mass Balance. *Science*, 338(6111), 1183-1189. doi:10.1126/science.1228102
- Silvestri, G., & Vera, C. (2009). Nonstationary Impacts of the Southern Annular Mode on Southern Hemisphere Climate. *Journal of Climate*, 22(22), 6142-6148. doi:10.1175/2009JCLI3036.1
- Smith, B., Fricker, H. A., Gardner, A. S., Medley, B., Nilsson, J., Paolo, F. S., et al. (2020). Pervasive ice sheet mass loss reflects competing ocean and atmosphere processes. *Science*, 368(6496), 1239-1242. doi:10.1126/science.aaz5845

- Spence, P., Holmes, R. M., Hogg, A. M., Griffies, S. M., Stewart, K. D., & England, M. H. (2017). Localized rapid warming of West Antarctic subsurface waters by remote winds. *Nature Climate Change*, 7(8), 595-603. doi:10.1038/nclimate3335
- Thomas, E. R., van Wessem, J. M., Roberts, J., Isaksson, E., Schlosser, E., Fudge, T. J., et al. (2017). Regional Antarctic snow accumulation over the past 1000 years. *Clim. Past*, 13(11), 1491-1513. doi:10.5194/cp-13-1491-2017
- Turner, J., Phillips, T., Hosking, J. S., Marshall, G. J., & Orr, A. (2013). The Amundsen Sea low. *International Journal of Climatology*, 33(7), 1818-1829. doi:10.1002/joc.3558
- van den Broeke, M. R., & van Lipzig, N. P. M. (2017). Changes in Antarctic temperature, wind and precipitation in response to the Antarctic Oscillation. *Annals of Glaciology*, 39, 119-126. doi:10.3189/172756404781814654
- Veldhuijsen, S. B. M., van de Berg, W. J., Brils, M., Kuipers Munneke, P., & van den Broeke, M. R. (2023). Characteristics of the 1979–2020 Antarctic firm layer simulated with IMAU-FDM v1.2A. *The Cryosphere*, 17(4), 1675-1696. doi:10.5194/tc-17-1675-2023
- Verfaillie, D., Pelletier, C., Goosse, H., Jourdain, N. C., Bull, C. Y. S., Dalaiden, Q., et al. (2022). The circum-Antarctic ice-shelves respond to a more positive Southern Annular Mode with regionally varied melting. *Communications Earth & Environment*, 3(1), 139. doi:10.1038/s43247-022-00458-x
- Wahr, J., Molenaar, M., & Bryan, F. (1998). Time variability of the Earth's gravity field: Hydrological and oceanic effects and their possible detection using GRACE. *Journal of Geophysical Research: Solid Earth*, 103(B12), 30205-30229. doi:10.1029/98JB02844
- Wang, B. (1992). The Vertical Structure and Development of the Enso Anomaly Mode during 1979-1989. *Journal of the Atmospheric Sciences*, 49(8), 698-712. doi:10.1175/1520-0469(1992)049<0698:Tvsado>2.0.Co;2
- Williams, S. D. P., Moore, P., King, M. A., & Whitehouse, P. L. (2014). Revisiting GRACE Antarctic ice mass trends and accelerations considering autocorrelation. *Earth and Planetary Science Letters*, 385, 12-21. doi:10.1016/j.epsl.2013.10.016
- Wingham, D. J., Wallis, D. W., & Shepherd, A. (2009). Spatial and temporal evolution of Pine Island Glacier thinning, 1995-2006. *Geophysical Research Letters*, 36, 5. Article. doi:L1750, 10.1029/2009gl039126
- Wouters, B., Bamber, J. L., van den Broeke, M. R., Lenaerts, J. T. M., & Sasgen, I. (2013). Limits in detecting acceleration of ice sheet mass loss due to climate variability. *Nature Geoscience*, 6(8), 613-616. doi:10.1038/ngeo1874

## References From the Supporting Information

- Haran, T., Bohlander, J., Scambos, T., & Fahnestock, M. (2014, updated 2019). *MODIS mosaic of Antarctica 2008-2009 (MOA2009) Image Map, Version 1*. doi:10.7265/N5KP8037
- Howat, I. M. (2022). *The Reference Elevation Model of Antarctica – Mosaics, Version 2*. doi:10.7910/DVN/EBW8UC
- Rignot, E., Mouginot, J., & Scheuchl, B. (2017). *MEaSURES InSAR-Based Antarctica Ice Velocity Map, Version 2*. doi:10.5067/D7GK8F5J8M8R

520

521

*Geophysical Research Letters*

522

Supporting Information for

523

**Major modes of climate variability dominate nonlinear Antarctic ice-sheet elevation changes 2002-  
2020**

524

525

Matt A. King<sup>1,2</sup>, Poul Christoffersen<sup>2,3</sup>

526

<sup>1</sup>School of Geography, Planning, and Spatial Sciences, University of Tasmania, Hobart, Tasmania  
7001, Australia

527

528

<sup>2</sup>The Australian Centre for Excellence in Antarctic Science, University of Tasmania, Hobart,  
Tasmania 7001, Australia

529

530

<sup>3</sup>Institute for Marine and Antarctic Studies, University of Tasmania, Hobart, Tasmania 7001,  
Australia

531

532

533

534

**Contents of this file**

535

536

Text S1

537

Figures S1 to S9

538

Table S1

539

540


## Text S1

To make a first-order estimate of plausible velocity-driven changes on surface elevation on Pine Island and Thwaites Glacier, we used annual Measures v2 ice velocities (Rignot et al., 2017) to derive year-on-year velocity changes at the glacier point positions where we extracted time series of surface elevation changes (shown in Fig. 3). The annual velocity changes were extracted in polar stereographic coordinates ( $x, y$ ) and averaged over a  $25 \times 25$  km box centred around each of the two geographical locations. The corresponding changes in surface height were estimated from the conservation of mass:

$$\frac{\partial h}{\partial t} = \dot{a} + \dot{b} + \bar{v} \cdot \nabla H + H \cdot \dot{\epsilon}_z$$


where  $\dot{a}$  is the surface accumulation from snowfall,  $\dot{b}$  is the basal accumulation when water at the bed freezes on (negative for melting),  $\bar{v}$  is the velocity vector,  $\nabla$  is the gradient operator,  $H$  is ice thickness and  $\dot{\epsilon}_z$  is the vertical strain rate averaged for the ice column. We used the third and fourth term on the RHS to derive estimates of the elevation change stemming specifically from the change in velocity, ignoring the accumulation terms  $\dot{a}$  and  $\dot{b}$ . If the velocity change is  $\delta \bar{v}$ , the change in height is  $-\delta \bar{v} \cdot \nabla H + H \cdot \delta \dot{\epsilon}_z$ , where the first term denotes elevation gain from advection of thicker ice (or the opposite) and the second term is elevation loss due to dynamic thinning when the ice velocity increases (or the opposite). The change in velocity,  $\delta \bar{v} = (\Delta v_x, \Delta v_y)$  was the annual difference from one year to the next. The advection term was calculated as  $\Delta v_x dh/dx + \Delta v_y dh/dy$ , while strain thinning was calculated as  $H(-\Delta(dv_x/dx) - \Delta(dv_y/dy))$  with the assumption that ice is incompressible ( $\delta \dot{\epsilon}_z = -\delta \dot{\epsilon}_x - \delta \dot{\epsilon}_y$ ) and vertical strain in the ice column is uniform. Fig. S10 shows the estimated elevation change in terms of advection and strain and their total.

## Supplementary Figures



**Fig S1.** Cumulatively summed and detrended climate indices. The panels show the normalized climate indices (a), their cumulative sum renormalized (b), after further detrending and renormalization (c). ENSO and ASL $\lambda$  terms are shown offset by 4 units for clarity.

576




577


578

579 **Fig S2.** Correlation coefficients of summed and detrended climate indices.


580




**Fig S3.** The leading two modes of the detrended altimeter and GRACE data based on MVEOF. The EOFs are shown after scaling to reverse the effects of normalization. GRACE modes (a, d) are shown in units of meters of water equivalent (w.e.), and altimeter modes (b, e) shown in units of meters of ice elevation. Overlain on PC1 is the detrended SAM<sub>Σ</sub> and -ASLP<sub>Σ</sub> indices and overlain on PC2 is the lagged ENSO<sub>Σ</sub> and -ASLλ<sub>Σ</sub>. The variances explained by each mode are indicated in the legends of panels c and f.




**Fig S4.** Modes 3 and 4 of the MVEOF. Note the changes in color scale from Fig S3.




**Fig S5.** Detail of Figure 1 in the Amundsen Sea Embayment (top) and Denman-Totten glacier (bottom) regions. Thick black lines show the grounding line and thin black lines the ice shelf limits (Haran et al., 2014, updated 2019). Pink lines define the 100 m/yr speed contour (Rignot et al., 2017). Yellow lines mark the locations of ice stream profiles and yellow cross marks the location of the ice stream time series.




**Fig S6.** Density of SAM<sub>Σ</sub> and ENSO<sub>Σ</sub> variability from the ratio of GRACE and smoothed-altimetry coefficients. Error bars reflect 1-sigma uncertainties. ENSO<sub>Σ</sub> values are not shown for Denman and Totten as they are too uncertain.




**Fig S7.** Cross-section profiles of SAM<sub>Σ</sub> (a) and ENSO<sub>Σ</sub> (b) coefficients as a function of distance across Thwaites (a), Pine Island (b), Totten (c) and Denman (d) glaciers. Coefficients are shown as estimated before (colored solid line) and after (colored dashed line) subtraction of the IMAU FDM. The ice elevation extracted from the REMA v2.0 1 km mosaic (Howat, 2022) is shown as a black line (central axes). Colored shading indicates the 1-sigma confidence limits. The vertical grey box indicates the 100 m/yr limits of ice flow based on Measures v2. The locations of the profiles are shown in Figure S5 as yellow lines.



**Fig S8.** Centre-line profiles of  $SAM_{\Sigma}$  (a) and  $ENSO_{\Sigma}$  (b) coefficients as a function of distance upstream of the grounding line of major ice streams. Coefficients are shown before (solid line) and after (dashed line) subtraction of the IMAU FDM. Shading indicates the 1-sigma confidence limits. The locations of the profiles are shown in Figure S5 as yellow lines.



**Figure S9.** As for Figure 3 but showing only the Thwaites and Pine Island time series. Panels show (left axes) the detrended data and their 2-sigma uncertainties (blue pluses and error bars), outputs of two models of firn densification (FDM; brown lines), the best fitting models based on ASL (black solid line with 2-sigma uncertainty in gray shading) and SAM+ENSO (black dashed lines) terms.



**Figure S10.** Dynamic elevation change anomalies (positive for gain) estimated from annual changes in satellite derived velocities for Pine Island Glacier (top) and Thwaites Glacier (bottom). These elevation change anomalies are calculated from the advective thickness change (blue circles) and strain (magenta triangle) tied to the observed change in velocity from one year to the next. The total elevation change (black squares) is the sum. Estimates are based on gridded surface velocities ( $v_x, v_y$ ) in a 25 x 25 km box centered around glacier points used to show observed elevation changes in Figure 3 of main text. Other sources of elevation change are ignored. See Text S1 for details.

Table S1

Table S1. Location in Antarctic Polar Stereographic coordinates (EPSG:3031) of sites in Fig. 3 and Fig S9 and discussed in the main text.

| <b>Glacier</b> | <b>X(m)</b> | <b>Y(m)</b> |
|----------------|-------------|-------------|
| Thwaites       | -1511702    | -463473     |
| Pine Island    | -1583860    | -232513     |
| Totten         | 2274795     | -988293     |
| Denman         | 2496950     | -423352     |