

Modern air-sea flux distributions reduce uncertainty in the future ocean carbon sink

Galen A McKinley¹, Galen A Mckinley², Val Bennington^{2,3,4}, Malte Meinshausen¹, and Zebedee Nicholls⁵

¹Affiliation not available

²Columbia University and Lamont-Doherty Earth Observatory New York

³Makai Ocean Engineering, Inc

⁴School of Geography, Earth and Atmospheric Sciences, Melbourne Climate Futures, The University of Melbourne

⁵Energy, Climate and Environment (ECE) Program, International Institute for Applied Systems Analysis (IIASA)

January 24, 2023

Modern air-sea flux distributions reduce uncertainty in the future ocean carbon sink

Galen A McKinley[†], Val Bennington^{†,‡}, Malte Meinshausen[★], Zebedee Nicholls^{★,*}

[†] Columbia University and Lamont-Doherty Earth Observatory New York, USA
[‡] Makai Ocean Engineering, Inc., Hawaii, USA
[★] Melbourne Climate Futures, School of Geography, Earth and Atmospheric Sciences, The University of Melbourne, Melbourne, Victoria, Australia
^{*} Energy, Climate and Environment (ECE) Program, International Institute for Applied Systems Analysis (IIASA), Laxenburg, Austria

Abstract.

The ocean has absorbed about 25% of the carbon emitted by humans to date. To better predict how much climate will change, it is critical to understand how this ocean carbon sink will respond to future emissions. Here, we examine the ocean carbon sink response to low emission (SSP1-1.9, SSP1-2.6), intermediate emission (SSP2-4.5, SSP5-3.4-OS), and high emission (SSP5-8.5) scenarios in CMIP6 Earth System Models and in MAGICC7, a reduced-complexity climate carbon system model. From 2020-2100, the trajectory of the global-mean sink approximately parallels the trajectory of anthropogenic emissions. Until emission growth becomes negative, the cumulative ocean carbon sink absorbs 20-30% of cumulative emissions since 2015. In scenarios where emissions decline and become negative, the ocean remains a sink and absorbs more carbon than emitted (up to 120% of cumulative emissions since 2015). Despite similar responses in all models, there remains substantial quantitative spread in estimates of the cumulative sink through 2100 within each scenario, up to 50 PgC in CMIP6 and 120 PgC in the MAGICC7 ensemble. We demonstrate that for all but SSP1-2.6, approximately half of this future spread can be eliminated if models are brought into agreement with modern best-estimates. Considering the spatial distribution of air-sea CO₂ fluxes in CMIP6, we find significant zonal-mean divergence from newly-available observation-based constraints. We conclude that a significant portion of future ocean carbon sink uncertainty is attributable to modern-day errors in the mean state of air-sea CO₂ fluxes, which in turn are associated with model representations of ocean physics and biogeochemistry. Bringing models into agreement with modern observation-based estimates at regional to global scales can substantially reduce uncertainty in future role of the ocean in absorbing anthropogenic CO₂ from the atmosphere and mitigating climate change.

1. Introduction

Since the industrial revolution, emissions due to industrial and land use activities have dramatically increased atmospheric carbon dioxide concentrations. Due to sinks in the ocean and terrestrial biosphere, less than half of these emissions remain in the atmosphere. The ocean has absorbed approximately 25% of anthropogenic emissions both in recent decades and cumulatively since 1750 [Canadell et al.2021, Friedlingstein et al.2022]. This means that as atmospheric CO₂ has increased approximately exponentially, there has also been a comparable increase in the magnitude of ocean carbon sink [Raupach et al.2014, Ridge and McKinley2021]. On millennial timescales, the ocean will absorb more than 80% of the total anthropogenic perturbation [Archer et al.2009, Cao and Caldeira2010].

Detailed assessment of the ocean carbon sink’s future state have largely focused on CMIP5 Earth System Model (ESM) projections under the RCP8.5 scenario of steadily increasing emissions through 2100 [Schwinger et al.2014, Hoffman et al.2014, Randerson et al.2015, McKinley et al.2016, Wang et al.2016, Fassbender et al.2017, Goris et al.2018]. Without significant mitigation, the ocean will remain a strong sink for anthropogenic carbon through 2100, despite modest negative feedbacks due to ocean carbon chemistry and circulation changes. The response is similar in CMIP6 ESMs [Arora et al.2020].

Nearly 200 countries have signed the UNFCCC Paris Agreement, indicating a serious intent to rapidly mitigate emissions. These reductions, as well as the changed economics of renewable energy systems, are starting to shift the plausible future emission range away from the highest scenarios [Hausfather and Peters2020]. Aside from climate targets for 2030, many countries also put forward

long-term targets - often net-zero CO₂ or net-zero greenhouse gas targets. Given that the ocean will continue to strongly influence the global-mean warming trajectory in the long-term, it is important to assess the future ocean carbon sink under scenarios of mitigation.

[Raupach et al.2014] discussed the ocean sink in the context of the atmospheric fraction (AF), which is the fraction of annual fossil and land use emissions that remain in the atmosphere. They demonstrated that the near-constant AF of 0.44 from 1959-2013 cannot be explained by underlying mechanisms of the ocean and terrestrial biosphere sinks, but instead is attributable to the approximately exponential atmospheric pCO₂ growth rate. When the atmospheric pCO₂ growth rate slows, AF will be reduced, meaning that less of the emitted carbon will remain in the atmosphere (see Figure SPM.7 in [IPCC2021]). For mitigation scenarios, the ratio of cumulative atmospheric load to cumulative emissions, or the Cumulative Atmospheric Fraction (CAF) is preferable [Jones et al.2016] because it remains numerically well behaved as emissions approach zero. The CAF is also most directly relevant to climate outcomes that depend on the reservoir of atmospheric CO₂, not its annual fluxes [Matthews et al.2020].

The global-mean ocean carbon sink will weaken in absolute magnitude as atmospheric pCO₂ growth rates slow [Cao and Caldeira2010, Jones et al.2016, Zickfeld et al.2016, Schwinger and Tjiputra2018, McKinley et al.2020, Ridge and McKinley2021, Canadell et al.2021]. Despite a reduced absolute magnitude, as long as the ocean overturning circulation continues to expose waters with additional carbon uptake capacity to the surface, the sink will continue [Zickfeld et al.2016]. Under mitigation scenarios, CAF estimates for the 21st century should be substantially lower with mitigation than without [Jones et al.2016, IPCC2021]. In other words, the ocean will be able to do proportionally more to reduce climate warming as humans increasingly mitigate emissions.

Studies to date have focused on the global-mean ocean carbon sink estimated by reduced complexity models or ESMs. Quantitative uncertainty in the magnitude of the future ocean carbon sink under mitigation scenarios has not received much attention. The global-mean ocean carbon sink is the integrated result of the anthropogenic perturbation superimposed on a background of vigorous natural fluxes [McKinley et al.2017, Crisp et al.2022]. To understand why models differ, local to regional fluxes must be considered since these are the scales on which physical and biogeochemical mechanisms of natural and anthropogenic carbon fluxes occur [McKinley et al.2016, Fay and McKinley2021]. For anthropogenic fluxes specifically, advection and water mass transformation at regional scales dominate fluxes into (reemergence) and out of (subduction) the surface mixed layer, which is critical to the movement of anthropogenic carbon to and from the deep ocean [Bopp et al.2015, Iudicone et al.2016, Toyama et al.2017, Ridge and McKinley2020]. Mechanisms of future sink will also occur locally, and thus understanding why ESMs quantitatively differ in their projections requires consideration of the spatial distribution.

[Ridge and McKinley2021] studied the three-dimensional response of the ocean carbon sink to mitigation scenarios in one ESM, the Community Earth System Model Large Ensemble [Kay et al.2015]. They found that the large-scale spatial distribution of the sink is largely conserved through 2100. The primary change is an increased or decreased amplitude of the sink in high or low emission scenarios. With strong mitigation (1.5°C scenario), some regions that were previous sinks become sources to the atmosphere as thermocline waters with high anthropogenic carbon content are recirculated to the surface. This study investigates whether these estimates of future flux distributions are consistent across a range of ESMs.

Ocean hindcast models have been critical to the annual Global Carbon Budget (GCB) since its inception [Hauck et al.2020, Friedlingstein et al.2022]. Hindcast models are forced with observed meteorology to estimate the actual evolution of ocean physics and biogeochemistry in recent years. Yet their underlying structures and parameterizations are very similar to the ocean component models in their cousin ESMs. Thus, skill assessments for hindcast models likely provide some information about ESM fidelity. A recent assessment of long-term mean and average seasonal fluxes in the nine hindcast models used in the GCB 2020 [Friedlingstein et al.2020] revealed significant regional discrepancies from observation-based estimates [Fay and McKinley2021].

A robust suite of full global coverage, monthly timescale observation-based products for surface ocean carbon content, from which air-sea CO₂ fluxes can be derived, have only recently become available [Rödenbeck et al.2015, Fay et al.2021]. Though these are based on sparse data, they have high fidelity for the long-term mean and the average seasonal cycle of air-sea CO₂ fluxes

[Gloege et al.2022a]. Compared to independent data, modern observation-based products represent surface ocean carbon concentrations much better than hindcast models [Bennington et al.2022b]. Given this new observational constraint, it is important to assess how the state-of-the-art ESMs from the sixth phase of the Coupled Model Intercomparison Project (CMIP6) suite represent modern-day mean fluxes. Do the CMIP6 ESMs demonstrate similar biases as identified for hindcast models by [Fay and McKinley2021]? What are the implications for uncertainty in projections of the future sink?

To address these issues, we analyze the ocean carbon sink response to emission scenarios with varying degrees of mitigation (SSP1-1.9, SSP1-2.6, SSP2-4.5, SSP5-3.4-OS, SSP5-8.5) from CMIP6. We also examine ocean carbon sink estimates from the reduced-complexity climate carbon cycle model, MAGICC7, that was used in the creation of the SSP scenarios [Meinshausen et al.2020].

2. Methods

2.1. Emissions Scenarios

We study the ocean carbon sink for 5 Shared Socioeconomic Pathway (SSP) scenarios with low emissions (SSP1-1.9, SSP1-2.6), intermediate emissions (SSP2-4.5, SSP5-3.4-OS), and high emissions (SSP5-8.5) over 2015-2100 [Meinshausen et al.2020].

2.2. CMIP6 Simulations

This study utilizes concentration-driven simulations of Earth System Models (ESMs) performed as part of the sixth phase of the Coupled Model Intercomparison Project (CMIP6). Monthly average simulated air-sea carbon dioxide fluxes (variable name: fgco2) for the five emission scenarios were obtained from the CMIP6 archive <https://esgf-node.llnl.gov/search/cmip6/>, (World Climate Research Programme, 2021) (Table 1). For analyses, all modeled fluxes were conservatively regridded to a regular 1° latitude by 1° longitude grid using the xesmf module in Python. Simulated fluxes are integrated in space to obtain annual air-sea carbon fluxes for each ensemble member; ensemble means are used in analysis. Ensemble-means are used to damp internal variability. For CNRM, a carbon efflux of 0.78 Pg C /yr that is due to natural inputs from rivers is removed [Séférian et al.2019]; other Earth System Models do not include natural river carbon fluxes.

2.3. MAGICC7

MAGICC7 is a reduced-complexity climate carbon system model calibrated to emulate CMIP5 ESMs [Meinshausen et al.2011, Meinshausen et al.2020] and for the results shown here, the range of CMIP6 ESMs used for IPCC AR6 WG1 assessments (see Cross-Chapter Box 7.1 in IPCC AR6 WG1 [Forster et al.2021]). MAGICC7 simulates the land and global-mean ocean carbon sinks that occur in response on emissions. It was used to develop atmospheric carbon dioxide concentrations for the SSP scenarios [Meinshausen et al.2020], which were then input to the concentration-driven CMIP6 runs that we analyze here. For the comparison between the CMIP6 ensemble and MAGICC7, the probabilistic version of MAGICC7 has been calibrated to comprehensively reflect climate system response uncertainties [Forster et al.2021]. Our comparisons are a consistency check between the ocean sink estimates from MAGICC7 and the ocean sink estimates resulting from the state-of-the-art CMIP6 suite driven with the atmospheric concentrations derived from MAGICC7.

2.4. Observation-based products

For the years 2010-2019, we compare the CMIP6 models to the currently-available ensemble of observation based products (Table 2). These products are produced using machine learning and other statistical methods to extrapolate sparse pCO₂ observations to global coverage. All products estimate monthly fluxes for the 1980s to the 2020s, while two of the newest products start in 1959 [Bennington et al.2022b, Rödenbeck et al.2022]. A suite of wind speed products is then used to estimate CO₂ flux [Fay et al.2021]. These products have been shown to offer robust estimates of long-term mean CO₂ fluxes from global to regional scales [Gloege et al.2022b, Fay and McKinley2021, Bennington et al.2022b].

Observation-based products provide an estimate of the total air-sea CO₂ flux, the sum of the anthropogenic fluxes and outgassing due to the import of natural carbon from rivers [Crisp et al.2022]. Since CMIP6 and MAGICC7 models estimate only anthropogenic fluxes, we remove the natural efflux due to rivers in each latitude band, based on the spatial distribution from [Lacroix et al.2020] and the global total flux of +0.65 PgC/yr [Regnier et al.2022] (positive to the atmosphere).

2.5. Ratio of cumulative sink to cumulative emissions (CSF_{ocean})

The atmospheric fraction (AF) is the fraction of annual fossil and land use emissions that remain in the atmosphere [Raupach et al.2014]. The cumulative atmospheric fraction (CAF) is the ratio of cumulative atmospheric load to cumulative emissions [Jones et al.2016]. Here, we evaluate the role of the ocean sink in setting the CAF. Thus, we define a cumulative sink fraction as the ratio of cumulative land and ocean sinks to cumulative emissions ($CSF = CSF_{land} + CSF_{ocean}$; and $CSF = 1 - CAF$). In this study, the ocean component (CSF_{ocean}) is the ratio of the cumulative ocean sink since 2015 to cumulative emissions since 2015.

3. Results

As emissions accumulate through 2100 (Figure 1a, S1), the cumulative ocean sink (Figure 1b) follows a similar trajectory. With higher emissions, the ocean accumulates more carbon, and with lesser emissions, less carbon is absorbed by the ocean. However, the ratio of cumulative sink to cumulative emissions (CSF_{ocean}) has the opposite response ([IPCC2021]). With accelerating emissions (SSP5-8.5 and SSP2-4.5), the ocean accumulates only 20 to 30% of the cumulative emissions after 2015. If emission rates decline, the ocean takes up a greater portion of emitted carbon, between 40% and 110% under SSP5-3.4-OS, SSP1-2.6 and SSP1-1.9 (Figure 1c, Table 3).

For each scenario, there are significant differences between the maximum and minimum predictions (Figure 1b), with a spread of 30-50 PgC in the cumulative uptake by 2100 ocean sink in CMIP6 models and range of 120 PgC in the MAGICC7 ensemble (Table 4). In all scenarios except SSP1-2.6 for both CMIP6 and MAGICC7, this spread can be reduced by 50% or more if all models estimates are adjusted to have the same sink magnitude in 2020 (3 PgC/yr, [Friedlingstein et al.2021]) (Figures 1d, S1, S2). This adjustment is accomplished by adding to all years the difference from 3 PgC/yr in 2020 for that CMIP6 or MAGICC7 model. Consistent with ESMs tending to underestimate the modern day sink [Hoffman et al.2014], uptake estimates here are increased in most cases. The reduced spread occurs because maximum uptake estimates are increased less than minimum estimates (Table 4, 5). For SSP1-2.6 in CMIP6, the maximum and minimum are increased by about the same amount.

With this adjustment, the spread in ratios of sink to emissions (CSF_{ocean}) for CMIP6 is also reduced by more than 50% in all scenarios except SSP1-2.6 (Figure 1e; Table 4). For SSP1-2.6 in CMIP6, there is a bimodal distribution of the projected cumulative sink and CSF_{ocean} after the adjustment of the 2020 sink (Figure 1d,e). This substantial reduction in spread, from homogenizing the present-day sink estimates, indicates that under most scenarios, 50% or more of future uncertainty is attributable to modern mean-state errors.

Newly available observation-based products offer the best-available constraint on the long-term mean and seasonality of air-sea CO₂ fluxes [Gloege et al.2022b, Bennington et al.2022b]. To better understand the performance of the CMIP6 models suite, we compare them to the currently-available suite of products (Table 2) for the 2010s (2010-2019). For this period, the large scale pattern of air-sea flux in the CMIP6 models is broadly comparable to the products (Figure 2, left; Figure S3). Outgassing occurs across much of the equatorial band, while uptake is greatest in the high northern latitudes and around 40°S.

Looking to the future, we choose one model that is representative of the CMIP6 suite, IPSL-CM6A-LR, to visualize expected air-sea flux changes. Changes in other models are similar (not shown). If emissions continue on a very high emission trajectory (SSP5-8.5), the ocean will take up more carbon in the 2050s than in the 2010s (Figure 2, right). By the 2090s, across the oceans south of 30°N, this pattern of greater uptake intensifies. However, there is reduced uptake compared to the 2010s north of 30°N. With intermediate emissions (SSP2-4.5) carbon uptake increases in the Southern Ocean and declines in the North Atlantic and North Pacific by the 2050s. By 2090, there

is substantially reduced uptake almost everywhere, except the high latitude Southern Ocean. Under the lowest emission scenario (SSP1-1.9) by the 2050s, carbon uptake declines compared to the 2010s except in the high latitude Southern Ocean. By the 2090s, uptake further declines everywhere. In summary, change in the ocean carbon sink under future scenarios is rather spatially homogeneous, except in the Southern Ocean where the carbon uptake generally persists, and in the North Atlantic and North Pacific where slowing uptake occurs under all scenarios.

Considering air-sea fluxes in zonal average allows for comparison of the CMIP6 suite to the modern spatial distribution (Figure 3, left), as well as the future change predicted by each ESMs (Figure 3, right). At latitudes north of 55°S, the CMIP6 ESMs capture the basic features of the observed flux distribution. However, the magnitude of modeled fluxes frequently lies outside the 2 σ spread of the 8 observation-based products. In the high latitude Southern Ocean, the observation-based products indicate a slight sink for 2010-2019, but the ESMs simulate from a large sink to a modest source.

In the future under both high, intermediate and low scenarios, the CMIP6 models all suggest a very similar response relative to their modern flux distribution. Under SSP5-8.5, the sink grows at all latitudes, with the strongest increase in the Southern Ocean. Under SSP2-4.5, most latitudes have a modest increase by 2050 and then a decline through 2090, while the Southern Ocean sink grows. Under SSP1-1.9, the sink reduces at all latitudes, with enhanced magnitude of reduction between 40°S and 60°S. The key point is here is that despite the significant spread across the CMIP6 models for the 2010s (Figure 3, left), there is strong agreement with respect to the magnitude of future sink change (Figure 3, right). This finding applies to all four scenarios considered here. In other words, as for the global mean (Figure 1), the modern state of the CMIP6 ESMs holds a significant portion of the uncertainty with respect to the magnitude of the future ocean sink across emission scenarios.

4. Discussion

The future ocean carbon sink will grow if emissions grown and decline if emissions decline. Yet, there remains substantial disagreement as to the magnitude of the projected ocean carbon sink within each scenario for both the CMIP6 Earth System Model suite and the MAGICC7 ensemble (Figure 1). Significant reduction in the spread of the future global-mean cumulative ocean sink can be achieved by adjusting the models to capture modern best-estimates (Figure 1). Under most scenarios, 50% or more of uncertainty in the magnitude of the ocean carbon sink at 2100 can be eliminated (Table 4, 5). This indicates a strong correlation between present-day and future mean-state biases. Under SSP1-2.6 in CMIP6, this adjustment leads to a tightening of projections into a bimodal distribution, but no reduction in total spread.

Global mean results from CMIP6 to MAGICC7 are in substantial agreement (Figure 1), aside from the larger spread across the MAGICC7 ensemble. Homogenizing the global-mean sink to the same 2020 value also substantially reduces the future spread of MAGICC7 projections and increases the agreement with CMIP6 (Figure 1d). This provides another line of evidence that present-day bias is a major contributor to our future uncertainty. One notable difference between CMIP6 and MAGICC7 is that in the very high emission scenario (SSP5-8.5), the CMIP6 models suggest a slightly earlier weakening of the ocean carbon sinks around the 2080s, whereas MAGICC7 ensemble members tend to plateau at this time (Figure S1). This difference may be due to reduction in the ocean overturning circulation that occurs in CMIP6 at this time [Liu et al.2022].

As noted in previous analyses [Jones et al.2016, IPCC2021], the percentage of future emissions that will be stored in the ocean is strongly dependent on the emissions trajectory (Figure 1c). We show the uncertainty in estimates of CSF_{ocean} can also be substantially reduced by addressing discrepancies in estimates of present-day fluxes (Figure 1e). Under SSP1-1.9 by 2100, cumulative emissions since 2015 and until the point of the net-zero emissions are around 209 PgC until 2055 and are then reduced again to 124 PgC by the end of the century due to net CO₂ removals (i.e. negative emissions) after 2055. With our proposed adjustment to the modern best-estimate, we would predict that all net cumulative emissions by 2100 would go into the ocean (128-147 PgC). Without adjustment, the predicted uptake has a much broader range (93 to 148 PgC) that would leave up to 30% of cumulative emissions in the atmosphere (Figure 1b,c; Table 4). The difference in these predictions illustrates the value of reducing prediction uncertainty.

For CMIP6, there are substantial regional biases against modern observation-based data products. Zonal-mean fluxes from CMIP6 frequently lie outside the 2σ spread of the observation-based products (Figure 3), a finding is consistent with recent regional assessment of related ocean hindcast models [Fay and McKinley2021]. The drivers of modeled mean-state biases are likely attributable to both the representations of ocean circulation and biogeochemistry in the models [Mongwe et al.2018, Gruber et al.2019, Fay and McKinley2021, Crisp et al.2022]. In-depth analysis of individual models will be required to fully understand biases and identify needed remedies.

Spatial patterns of air-sea CO_2 exchange do not change substantially through the 21st century (Figure 2). While emissions increase, model regions of influx increase and regions of efflux decrease in magnitude (Figure 2). There is some disagreement across models as to change in the northern middle to high latitudes under SSP5-8.5 (Figure 3). Prior studies suggest modeled physical processes that could cause some of this spread. In the previous version of CESM under high emissions (RCP8.5), the future North Atlantic experiences large freshwater fluxes that increasingly limit carbon uptake [Ridge2020, Ridge and McKinley2020]. Differential responses of the Atlantic Meridional Overturning Circulation and deep southward transport of anthropogenic carbon likely also contribute to these different predictions [Yool et al.2021, Liu et al.2022].

We demonstrate clear links between the future and the modern mean state of the ocean carbon sink in CMIP6 models and MAGICC7, and the future and the modern zonal distribution of the sink in CMIP6. This finding is consistent with [Terhaar et al.2022] who propose to constrain CMIP6 ocean sink estimates using three observational constraints related to the modern mean ocean circulation and carbon chemistry. Previous analyses of CMIP5 models under high emission scenarios also pointed to the forward propagation of modern mean state errors [Hoffman et al.2014, Wang et al.2016]. This strongly implies that improving models of the modern ocean carbon sink can lead to substantial reductions in uncertainty of future projections of the ocean carbon sink.

We propose to better constrain models to current best estimates to improve predictions. The best estimates for the global ocean sink, in fact, have substantial uncertainty. This is true if the sink is quantified for a single year or for a decade: $3.0 \pm 0.4 \text{ PgC/yr}$ (1σ) for 2020 [Friedlingstein et al.2021]; $2.5 \pm 0.6 \text{ PgC/yr}$ (1σ) for 2010-2019 (Figure 5.12 [Canadell et al.2021] [Friedlingstein et al.2020]). Uncertainties at regional scales are even larger proportions of the mean. The sparsity of ocean carbon observations drives much of the uncertainty in observation-based estimates [Bushinsky et al.2019, Gloege et al.2021]. At the same time as models are improved toward the observational constraints that are now available, tighter observational constraints should also be developed to further guide ocean and climate model development. Observations can also provide the basis for a post-simulation correction [Gloege et al.2022a, Bennington et al.2022b]. Improving both observation-based estimates and models will better constrain the global carbon cycle, which in turn will better support the climate policymaking process [Peters et al.2017].

5. Conclusion

The future ocean carbon sink will grow in absolute magnitude as future carbon emissions grow, and decline in magnitude as emissions decline. However, the proportion of future cumulative emissions that get absorbed by the ocean will be much greater if mitigation occurs. As emissions decline, the proportion of cumulative emissions absorbed by the ocean could be as high as 120%. But as long as emissions continue to rise, the ocean will accumulate only 20-30%.

While there is consensus across models as to these qualitative behaviors of the future ocean sink, significant quantitative uncertainties remain across all emission scenarios. These uncertainties can be substantially reduced by refining ocean and climate models to better represent long-term mean and seasonal flux patterns that observation-based products can now constrain. Going forward, improving modeled representations of the modern ocean carbon sink offers a tractable path forward to improving ocean and global carbon cycle projections, and thus the trajectory of future climate change.

6. Data Availability

The data used here are from the CMIP6 simulations performed by various modeling groups and publicly available from the CMIP6 archive: <https://esgf-node.llnl.gov/search/cmip6/>,

[WCRP2021], last access: 20 July 2021. The carbon emissions scenarios within this article were obtained from the SSP database hosted by the IIASA Energy Program at <https://tntcat.iiasa.ac.at/SspDb>. MAGICC7 is available for download at <https://magicc.org/download/magicc7> with the data used in this study being available at [to be made public after the review process, anonymous access can be provided to reviewers if desired].

Acknowledgments

GAM conceived of the study and drafted the manuscript, VB performed the analysis, MM and ZB contributed MAGICC data. All authors revised the manuscript. GAM and VB acknowledge support from National Oceanic and Atmospheric Administration (NA20OAR4310340); National Science Foundation through the LEAP Science Technology Center (Award 2019625) and OCE-1948624; and National Aeronautic and Space Administration (80NSSC22K0150). ZN acknowledges the European Union’s Horizon 2020 research and innovation program ESM2025 - Earth System Models for the Future (grant agreement no. 101003536). We thank Amanda Fay for assistance with the observation-based products.

7. References

- [Archer et al.2009] Archer, D., Eby, M., Brovkin, V., Ridgwell, A., Cao, L., Mikolajewicz, U., Caldeira, K., Matsumoto, K., Munhoven, G., Montenegro, A., and Tokos, K. (2009). Atmospheric Lifetime of Fossil Fuel Carbon Dioxide. *Annual Review Of Earth And Planetary Sciences*, 37:117–134.
- [Arora et al.2020] Arora, V. K., Katavouta, A., Williams, R. G., Jones, C. D., Brovkin, V., Friedlingstein, P., Schwinger, J., Bopp, L., Boucher, O., Cadule, P., Chamberlain, M. A., Christian, J. R., Delire, C., Fisher, R. A., Hajima, T., Ilyina, T., Joetzjer, E., Kawamiya, M., Koven, C. D., Krasting, J. P., Law, R. M., Lawrence, D. M., Lenton, A., Lindsay, K., Pongratz, J., Raddatz, T., Séférian, R., Tachiiri, K., Tjiputra, J. F., Wiltshire, A., Wu, T., and Ziehn, T. (2020). Carbon-concentration and carbon-climate feedbacks in CMIP6 models and their comparison to CMIP5 models. *Biogeosciences*, 17(16):4173–4222.
- [Bennington et al.2022a] Bennington, V., Galjanic, T., and McKinley, G. A. (2022a). Explicit Physical Knowledge in Machine Learning for Ocean Carbon Flux Reconstruction: The pCO₂-Residual Method. *Journal of Advances in Modeling Earth Systems*, 14(10).
- [Bennington et al.2022b] Bennington, V., Gloege, L., and McKinley, G. A. (2022b). Variability in the Global Ocean Carbon Sink From 1959 to 2020 by Correcting Models With Observations. *Geophysical Research Letters*, 49(14).
- [Bopp et al.2015] Bopp, L., Lévy, M., Resplandy, L., and Sallée, J. B. (2015). Pathways of anthropogenic carbon subduction in the global ocean. *Geophysical Research Letters*, 42(15):6416–6423.
- [Boucher et al.2020] Boucher, O., Servonnat, J., Albright, A. L., Aumont, O., Balkanski, Y., Bastrikov, V., Bekki, S., Bonnet, R., Bony, S., Bopp, L., Braconnot, P., Brockmann, P., Cadule, P., Caubel, A., Cheruy, F., Codron, F., Cozic, A., Cugnet, D., D’Andrea, F., Davini, P., de Lavergne, C., Denvil, S., Deshayes, J., Devilliers, M., Ducharne, A., Dufresne, J.-L., Dupont, E., Éthé, C., Fairhead, L., Falletti, L., Flavoni, S., Foujols, M.-A., Gardoll, S., Gastineau, G., Ghattas, J., Grandpeix, J.-Y., Guenet, B., Guez, Lionel, E., Guilyardi, E., Guimberteau, M., Hauglustaine, D., Hourdin, F., Idelkadi, A., Joussaume, S., Kageyama, M., Khodri, M., Krinner, G., Lebas, N., Levavasseur, G., Lévy, C., Li, L., Lott, F., Lurton, T., Luyssaert, S., Madec, G., Madeleine, J.-B., Maignan, F., Marchand, M., Marti, O., Mellul, L., Meurdesoif, Y., Mignot, J., Musat, I., Ottlé, C., Peylin, P., Planton, Y., Polcher, J., Rio, C., Rochetin, N., Rousset, C., Sepulchre, P., Sima, A., Swingedouw, D., Thiéblemont, R., Traore, A. K., Vancoppenolle, M., Vial, J., Vialard, J., Viovy, N., and Vuichard, N. (2020). Presentation and Evaluation of the IPSL-CM6A-LR Climate Model. *Journal of Advances in Modeling Earth Systems*, 12(7):e2019MS002010. e2019MS002010 10.1029/2019MS002010.
- [Bushinsky et al.2019] Bushinsky, S. M., Landschützer, P., Rödenbeck, C., Gray, A. R., Baker, D., Mazloff, M. R., Resplandy, L., Johnson, K. S., and Sarmiento, J. L. (2019). Reassessing Southern Ocean air-sea CO₂ flux estimates with the addition of biogeochemical float observations. *Global Biogeochemical Cycles*, 28(9):927.
- [Canadell et al.2021] Canadell, J., Monteiro, P., Costa, M., da Cunha, L. C., Cox, P., Eliseev, A., Henson, S., Ishii, M., Jaccard, S., Koven, C., Lohila, A., Patra, P., Piao, S., Rogelj, J., Syampungani, S., Zaehle, S., and Zickfeld, K. (2021). Global Carbon and other Biogeochemical Cycles and Feedbacks. In Masson-Delmotte, V., Zhai, P., Pirani, A., Connors, S., Péan, C., Berger, S., Caud, N., Chen, Y., Goldfarb, L., Gomis, M., Huang, M., Leitzell, K., Lonnoy, E., Matthews, J., Maycock, T., Waterfield, T., Yelekçi, O., Yu, R., and Zhou, B., editors, *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*, chapter 5. Cambridge University Press, Cambridge, UK and New York, NY, USA.
- [Cao and Caldeira2010] Cao, L. and Caldeira, K. (2010). Atmospheric carbon dioxide removal: long-term consequences and commitment. *Environmental Research Letters*, 5(2):024011.
- [Crisp et al.2022] Crisp, D., Dolman, H., Bastos, A., Sitch, S., Tanhua, T., McKinley, G., Hauck, J., Eggleston, S., and Aich, V. (2022). How well do we understand the land-ocean-atmosphere carbon cycle? *Reviews of Geophysics*, 60.
- [Denvil-Sommer et al.2019] Denvil-Sommer, A., Gehlen, M., Vrac, M., and Mejia, C. (2019). LSCE-FFNN-v1: A two-step neural network model for the reconstruction of surface ocean pCO₂ over the global ocean. *Geoscientific Model Development*, 12:2091–2105.
- [Fassbender et al.2017] Fassbender, A. J., Sabine, C. L., and Palevsky, H. I. (2017). Nonuniform ocean acidification and attenuation of the ocean carbon sink. *Geophysical Research Letters*, 44(16):8404–8413.
- [Fay et al.2021] Fay, A. R., Gregor, L., Landschützer, P., McKinley, G. A., Gruber, N., Gehlen, M., Iida, Y., Laruelle, G. G., Rödenbeck, C., Roobaert, A., and Zeng, J. (2021). SeaFlux: harmonization of air-sea CO₂ fluxes from surface pCO₂ data products using a standardized approach. *Earth System Science Data*, 13(10):4693–4710.
- [Fay and McKinley2021] Fay, A. R. and McKinley, G. A. (2021). Observed regional fluxes to constrain modeled estimates of the ocean carbon sink. *Geophysical Research Letters*, page e2021GL095325.
- [Forster et al.2021] Forster, P., Storelvmo, T., Armour, K., Collins, W., Dufresne, J.-L., Frame, D., Lunt, D., Mauritsen, T., Palmer, M., Watanabe, M., Wild, M., and Zhang, H. (2021). The Earth’s Energy Budget, Climate Feedbacks, and Climate Sensitivity. In Masson-Delmotte, V., Zhai, P., Pirani, A., Connors, S., Péan, C., Berger, S., Caud, N., Chen, Y., Goldfarb, L., Gomis, M., Huang, M., Leitzell, K., Lonnoy, E., Matthews, J., Maycock, T., Waterfield, T., Yelekçi, O., Yu, R., and Zhou, B., editors, *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*, chapter 7. Cambridge University Press, Cambridge, UK and New York, NY, USA.
- [Friedlingstein et al.2021] Friedlingstein, P., Jones, M. W., O’Sullivan, M., Andrew, R. M., Bakker, D. C., Hauck, J., Le Quéré, C., Peters, G. P., Peters, W., Pongratz, J., et al. (2021). Global carbon budget 2021. *Earth System Science Data Discussions*, pages 1–191.
- [Friedlingstein et al.2022] Friedlingstein, P., O’Sullivan, M., Jones, M. W., Andrew, R. M., Gregor, L., Hauck, J., Quéré, C. L., Luijkx, I. T., Olsen, A., Peters, G. P., Peters, W., Pongratz, J., Schwingshackl, C., Sitch, S., Canadell, J. G., Ciais, P., Jackson, R. B., Alin, S. R., Alkama, R., Arneth, A., Arora, V. K., Bates, N. R.,

1
2
3
4
5
6
7
8
9
10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Becker, M., Bellouin, N., Bittig, H. C., Bopp, L., Chevallier, F., Chini, L. P., Cronin, M., Evans, W., Falk, S., Feely, R. A., Gasser, T., Gehlen, M., Gkritzalis, T., Gloege, L., Grassi, G., Gruber, N., Gürses, , Harris, I., Hefner, M., Houghton, R. A., Hurtt, G. C., Iida, Y., Ilyina, T., Jain, A. K., Jersild, A., Kadono, K., Kato, E., Kennedy, D., Goldewijk, K. K., Knauer, J., Korsbakken, J. I., Landschützer, P., Lefèvre, N., Lindsay, K., Liu, J., Liu, Z., Marland, G., Mayot, N., McGrath, M. J., Metzl, N., Monacchi, N. M., Munro, D. R., Nakaoka, S.-I., Niwa, Y., O'Brien, K., Ono, T., Palmer, P. I., Pan, N., Pierrot, D., Pocock, K., Poulter, B., Resplandy, L., Robertson, E., Rödenbeck, C., Rodriguez, C., Rosan, T. M., Schwinger, J., Séférian, R., Shutler, J. D., Skjelvan, I., Steinhoff, T., Sun, Q., Sutton, A. J., Sweeney, C., Takao, S., Tanhua, T., Tans, P. P., Tian, X., Tian, H., Tilbrook, B., Tsujino, H., Tubiello, F., Werf, G. R. v. d., Walker, A. P., Wanninkhof, R., Whitehead, C., Wranne, A. W., Wright, R., Yuan, W., Yue, C., Yue, X., Zaehle, S., Zeng, J., and Zheng, B. (2022). Global Carbon Budget 2022. *Earth System Science Data*, 14(11):4811–4900.

[Friedlingstein et al.2020] Friedlingstein, P., O'Sullivan, M., Jones, M. W., Andrew, R. M., Hauck, J., Olsen, A., Peters, G. P., Peters, W., Pongratz, J., Sitch, S., Le Quéré, C., Canadell, J. G., Ciais, P., Jackson, R. B., Alin, S., Aragão, L. E. O. C., Arneth, A., Arora, V., Bates, N. R., Becker, M., Benoit-Cattin, A., Bittig, H. C., Bopp, L., Bultan, S., Chandra, N., Chevallier, F., Chini, L. P., Evans, W., Florentie, L., Forster, P. M., Gasser, T., Gehlen, M., Gilfillan, D., Gkritzalis, T., Gregor, L., Gruber, N., Harris, I., Hartung, K., Haverd, V., Houghton, R. A., Ilyina, T., Jain, A. K., Joetzjer, E., Kadono, K., Kato, E., Kitidis, V., Korsbakken, J. I., Landschützer, P., Lenton, A., Lienert, S., Liu, Z., Lombardozzi, D., Marland, G., Metzl, N., Munro, D. R., Nabel, J. E. M. S., Nakaoka, S.-i., Niwa, Y., O'Brien, K., Ono, T., Palmer, P. I., Pierrot, D., Poulter, B., Resplandy, L., Robertson, E., Rödenbeck, C., Schwinger, J., Séférian, R., Skjelvan, I., Smith, A. J. P., Sutton, A. J., Tanhua, T., Tans, P. P., Tian, H., Tilbrook, B., Van Der Werf, G., Vuichard, N., Walker, A. P., Wanninkhof, R., Watson, A. J., Willis, D., Wiltshire, A. J., Yuan, W., Yue, X., and Zaehle, S. (2020). Global Carbon Budget 2020. *Earth System Science Data*, 12:3269 – 3340.

[Gloege et al.2021] Gloege, L., McKinley, G. A., Landschützer, P., Fay, A. R., Frölicher, T. L., Fyfe, J. C., Ilyina, T., Jones, S., Lovenduski, N. S., Rodgers, K. B., Schlunegger, S., and Takano, Y. (2021). Quantifying errors in observationally based estimates of ocean carbon sink variability. *Global Biogeochemical Cycles*, 35(4):e2020GB006788. e2020GB006788 2020GB006788.

[Gloege et al.2022a] Gloege, L., Yan, M., Zheng, T., and McKinley, G. A. (2022a). Improved quantification of ocean carbon uptake by using machine learning to merge global models and pCO₂ data. *Journal of Advances in Modeling Earth Systems*, 14.

[Gloege et al.2022b] Gloege, L., Yan, M., Zheng, T., and McKinley, G. A. (2022b). Improved quantification of ocean carbon uptake by using machine learning to merge global models and pCO₂ data. *Journal of Advances in Modeling Earth Systems*, 14.

[Goris et al.2018] Goris, N., Tjiputra, J. F., Olsen, A., Schwinger, J., Lauvset, S. K., and Jeansson, E. (2018). Constraining projection-based estimates of the future North Atlantic carbon uptake. *Journal of Climate*.

[Gregor et al.2019] Gregor, L., Lebehot, A. D., Kok, S., and Monteiro, P. M. S. (2019). A comparative assessment of the uncertainties of global surface ocean CO₂ estimates using a machine-learning ensemble (CSIR-ML6 version 2019a) – have we hit the wall? *Geoscientific Model Development*, 12:5113–5136.

[Gruber et al.2019] Gruber, N., Landschützer, P., and Lovenduski, N. S. (2019). The variable southern ocean carbon sink. *Annual Review of Marine Science*, 11(1):159–186.

[Hajima et al.2020] Hajima, T., Watanabe, M., Yamamoto, A., Tatebe, H., Noguchi, M. A., Abe, M., Ohgaito, R., Ito, A., Yamazaki, D., Okajima, H., Ito, A., Takata, K., Ogochi, K., Watanabe, S., and Kawamiya, M. (2020). Development of the MIROC-ES2L Earth system model and the evaluation of biogeochemical processes and feedbacks. *Geoscientific Model Development*, 13(5):2197–2244.

[Hauck et al.2020] Hauck, J., Zeising, M., Le Quéré, C., Gruber, N., Bakker, D. C. E., Bopp, L., Chau, T. T. T., Gürses, Ö., Ilyina, T., Landschützer, P., Lenton, A., Resplandy, L., Rödenbeck, C., Schwinger, J., and Séférian, R. (2020). Consistency and Challenges in the Ocean Carbon Sink Estimate for the Global Carbon Budget . *Frontiers in Marine Science*, 7:3167.

[Hausfather and Peters2020] Hausfather, Z. and Peters, G. P. (2020). Emissions - the 'business as usual' story is misleading. *Nature*, 577(7792):618–620.

[Hoffman et al.2014] Hoffman, F. M., Randerson, J. T., Arora, V. K., Bao, Q., Cadule, P., Ji, D., Jones, C. D., Kawamiya, M., Khatiwala, S., Lindsay, K., Obata, A., Shevliakova, E., Six, K. D., Tjiputra, J. F., Volodin, E. M., and Wu, T. (2014). Causes and implications of persistent atmospheric carbon dioxide biases in Earth System Models. *Journal Of Geophysical Research-Biogeosciences*, 119(2):141 – 162.

[Iida et al.2021] Iida, Y., Takatani, Y., Kojima, A., and Ishii, M. (2021). Global trends of ocean CO₂ sink and ocean acidification: an observation-based reconstruction of surface ocean inorganic carbon variables. *Journal of Oceanography*, 77(2):323–358.

[IPCC2021] IPCC (2021). Summary for Policymakers. In *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*. Cambridge University Press, Cambridge, UK and New York, NY, USA.

[Iudicone et al.2016] Iudicone, D., Rodgers, K. B., Plancherel, Y., Aumont, O., Ito, T., Key, R. M., Madec, G., and Ishii, M. (2016). The formation of the ocean's anthropogenic carbon reservoir. *Scientific Reports*, 6:35473.

[Jones et al.2016] Jones, C. D., Ciais, P., Davis, S. J., Friedlingstein, P., Gasser, T., Peters, G. P., Rogelj, J., Vuuren, D. P. v., Canadell, J. G., Cowie, A., Jackson, R. B., Jonas, M., Kriegler, E., Littleton, E., Lowe, J. A., Milne, J., Shrestha, G., Smith, P., Torvanger, A., and Wiltshire, A. (2016). Simulating the Earth system response to negative emissions. *Environmental Research Letters*, 11(9):095012.

[Kay et al.2015] Kay, J. E., Deser, C., Phillips, A., Mai, A., Hannay, C., Strand, G., Arblaster, J. M., Bates, S. C., Danabasoglu, G., Edwards, J., Holland, M., Kushner, P., Lamarque, J.-F., Lawrence, D., Lindsay, K., Middleton, A., Munoz, E., Neale, R., Oleson, K., Polvani, L., and Vertenstein, M. (2015). The Community Earth System

- Model (CESM) Large Ensemble Project: A Community Resource for Studying Climate Change in the Presence of Internal Climate Variability. *Bulletin of the American Meteorological Society*, 96(8):1333–1349.
- [Lacroix et al.2020] Lacroix, F., Ilyina, T., and Hartmann, J. (2020). Oceanic CO₂ outgassing and biological production hotspots induced by pre-industrial river loads of nutrients and carbon in a global modeling approach. *Biogeosciences*, 17(1):55–88.
- [Landschützer et al.2020] Landschützer, P., Gruber, N., and Bakker, D. C. (2020). An observation-based global monthly gridded sea surface pco₂ product from 1982 onward and its monthly climatology. Technical Report Version 5.5, doi:10.7289/V5Z899N6, NOAA National Centers for Environmental Information.
- [Liu et al.2022] Liu, Y., Moore, J. K., Primeau, F., and Wang, W. L. (2022). Reduced CO₂ uptake and growing nutrient sequestration from slowing overturning circulation. *Nature Climate Change*, pages 1–8.
- [Matthews et al.2020] Matthews, H. D., Tokarska, K. B., Nicholls, Z. R. J., Rogelj, J., Canadell, J. G., Friedlingstein, P., Frölicher, T. L., Forster, P. M., Gillett, N. P., Ilyina, T., Jackson, R. B., Jones, C. D., Koven, C., Knutti, R., MacDougall, A. H., Meinshausen, M., Mengis, N., Séférian, R., and Zickfeld, K. (2020). Opportunities and challenges in using remaining carbon budgets to guide climate policy. *Nature Geoscience*, pages 1 – 11.
- [Mauritsen et al.2019] Mauritsen, T., Bader, J., Becker, T., Behrens, J., Bittner, M., Brokopf, R., Brovkin, V., Claussen, M., Crueger, T., Esch, M., Fast, I., Fiedler, S., Fläschner, D., Gayler, V., Giorgetta, M., Goll, D. S., Haak, H., Hagemann, S., Hedemann, C., Hohenegger, C., Ilyina, T., Jahns, T., Jimenéz-de-la Cuesta, D., Jungclaus, J., Kleinen, T., Kloster, S., Kracher, D., Kinne, S., Kleberg, D., Lasslop, G., Kornblueh, L., Marotzke, J., Matei, D., Meraner, K., Mikolajewicz, U., Modali, K., Möbis, B., Müller, W. A., Nabel, J. E. M. S., Nam, C. C. W., Notz, D., Nyawira, S.-S., Paulsen, H., Peters, K., Pincus, R., Pohlmann, H., Pongratz, J., Popp, M., Raddatz, T. J., Rast, S., Redler, R., Reick, C. H., Rohrschneider, T., Schemann, V., Schmidt, H., Schnur, R., Schulzweida, U., Six, K. D., Stein, L., Stemmler, I., Stevens, B., von Storch, J.-S., Tian, F., Voigt, A., Vrese, P., Wieners, K.-H., Wilkenskjaeld, S., Winkler, A., and Roeckner, E. (2019). Developments in the MPI-M Earth System Model version 1.2 (MPI-ESM1.2) and Its Response to Increasing CO₂. *Journal of Advances in Modeling Earth Systems*, 11(4):998–1038.
- [McKinley et al.2020] McKinley, G. A., Fay, A. R., Eddebbar, Y. A., Gloege, L., and Lovenduski, N. S. (2020). External forcing explains recent decadal variability of the ocean carbon sink. *AGU Advances*, 1(2):e2019AV000149.
- [McKinley et al.2017] McKinley, G. A., Fay, A. R., Lovenduski, N. S., and Pilcher, D. J. (2017). Natural variability and anthropogenic trends in the ocean carbon sink. *Annual Review of Marine Science*, 9(1):125–150.
- [McKinley et al.2016] McKinley, G. A., Pilcher, D. J., Fay, A. R., Lindsay, K., Long, M. C., and Lovenduski, N. S. (2016). Timescales for detection of trends in the ocean carbon sink. *Nature*, 530(7591):469–472.
- [Meinshausen et al.2020] Meinshausen, M., Nicholls, Z. R. J., Lewis, J., Gidden, M. J., Vogel, E., Freund, M., Beyerle, U., Gessner, C., Nauels, A., Bauer, N., Canadell, J. G., Daniel, J. S., John, A., Krummel, P. B., Luderer, G., Meinshausen, N., Montzka, S. A., Rayner, P. J., Reimann, S., Smith, S. J., van den Berg, M., Velders, G. J. M., Vollmer, M. K., and Wang, R. H. J. (2020). The shared socio-economic pathway (SSP) greenhouse gas concentrations and their extensions to 2500. *Geoscientific Model Development*, 13(8):3571–3605.
- [Meinshausen et al.2011] Meinshausen, M., Raper, S. C. B., and Wigley, T. M. L. (2011). Emulating coupled atmosphere-ocean and carbon cycle models with a simpler model, MAGICC6 - Part 1: Model description and calibration. *Atmospheric Chemistry And Physics*, 11(4):1417 – 1456.
- [Mongwe et al.2018] Mongwe, N., Vichi, M., and Monteiro, P. (2018). The seasonal cycle of pco₂ and co₂ fluxes in the southern ocean: diagnosing anomalies in cmip5 earth system models. *Biogeosciences*, 15(9):2851–2872.
- [Peters et al.2017] Peters, G. P., Le Quéré, C., Andrew, R. M., Canadell, J. G., Friedlingstein, P., Ilyina, T., Jackson, R. B., Joos, F., Korsbakken, J. I., McKinley, G. A., Sitch, S., and Tans, P. (2017). Towards real-time verification of CO₂ emissions. *Nature Climate Change*, 7(12):848–850.
- [Randerson et al.2015] Randerson, J., Lindsay, K., Munoz, E., Fu, W., Moore, J., Hoffman, F., Mahowald, N., and Doney, S. (2015). Multicentury changes in ocean and land contributions to the climate-carbon feedback: Carbon cycle feedbacks to 2300 in CESM. *Global Biogeochemical Cycles*, 29.
- [Raupach et al.2014] Raupach, M. R., Davis, S. J., Peters, G. P., Andrew, R. M., Canadell, J. G., Ciais, P., Friedlingstein, P., Jotzo, F., Van Vuuren, D. P., and Le Quere, C. (2014). Sharing a quota on cumulative carbon emissions. *Nature Climate Change*, 4(10):873–879.
- [Regnier et al.2022] Regnier, P., Resplandy, L., Najjar, R. G., and Ciais, P. (2022). The land-to-ocean loops of the global carbon cycle. *Nature*, 603(7901):401–410.
- [Ridge2020] Ridge, S. (2020). *Effects of Ocean Circulation on Ocean Anthropogenic Carbon Uptake*. PhD thesis, Columbia University.
- [Ridge and McKinley2020] Ridge, S. M. and McKinley, G. A. (2020). Advective Controls on the North Atlantic Anthropogenic Carbon Sink. *Global Biogeochemical Cycles*, 34(7):1138.
- [Ridge and McKinley2021] Ridge, S. M. and McKinley, G. A. (2021). Ocean carbon uptake under aggressive emission mitigation. *Biogeosciences*, 18(8):2711–2725.
- [Rödenbeck et al.2015] Rödenbeck, C., Bakker, D. C. E., Gruber, N., Iida, Y., Jacobson, A. R., Jones, S., Landschützer, P., Metzl, N., Nakaoka, S., Olsen, A., G-H, P., Peylin, P., Rodgers, K. B., Sasse, T. P., Schuster, U., Shutler, J. D., Valsala, V., Wanninkhof, R., and Zeng, J. (2015). Data-based estimates of the ocean carbon sink variability – first results of the surface ocean pco₂ mapping intercomparison (SOCOM). *Biogeosciences*, 12(23):7251–7278.
- [Rödenbeck et al.2022] Rödenbeck, C., DeVries, T., Hauck, J., Quéré, C. L., and Keeling, R. (2022). Data-based estimates of interannual sea-air CO₂ flux variations 1957-2020 and their relation to environmental drivers. *Biogeosciences*, 2022:1–43.
- [Schwinger and Tjiputra2018] Schwinger, J. and Tjiputra, J. (2018). Ocean Carbon Cycle Feedbacks Under Negative Emissions. *Geophysical Research Letters*, 45(10):5062–5070.

[Schwinger et al.2014] Schwinger, J., Tjiputra, J. F., Heinze, C., Bopp, L., Christian, J. R., Gehlen, M., Ilyina, T., Jones, C. D., Salas-Mélia, D., Segschneider, J., Séférian, R., and Totterdell, I. (2014). Nonlinearity of Ocean Carbon Cycle Feedbacks in CMIP5 Earth System Models. *Journal of Climate*, 27(11):3869 – 3888.

[Sellar et al.2019] Sellar, A. A., Jones, C. G., Mulcahy, J. P., Tang, Y., Yool, A., Wiltshire, A., O’Connor, F. M., Stringer, M., Hill, R., Palmieri, J., Woodward, S., de Mora, L., Kuhlbrodt, T., Rumbold, S. T., Kelley, D. I., Ellis, R., Johnson, C. E., Walton, J., Abraham, N. L., Andrews, M. B., Andrews, T., Archibald, A. T., Berthou, S., Burke, E., Blockley, E., Carslaw, K., Dalvi, M., Edwards, J., Folberth, G. A., Gedney, N., Griffiths, P. T., Harper, A. B., Hendry, M. A., Hewitt, A. J., Johnson, B., Jones, A., Jones, C. D., Keeble, J., Liddicoat, S., Morgenstern, O., Parker, R. J., Predoi, V., Robertson, E., Siahann, A., Smith, R. S., Swaminathan, R., Woodhouse, M. T., Zeng, G., and Zerroukat, M. (2019). UKESM1: Description and Evaluation of the U.K. Earth System Model. *Journal of Advances in Modeling Earth Systems*, 11(12):4513–4558.

[Swart et al.2019] Swart, N. C., Cole, J. N. S., Kharin, V. V., Lazare, M., Scinocca, J. F., Gillett, N. P., Anstey, J., Arora, V., Christian, J. R., Hanna, S., Jiao, Y., Lee, W. G., Majaess, F., Saenko, O. A., Seiler, C., Seinen, C., Shao, A., Sigmond, M., Solheim, L., von Salzen, K., Yang, D., and Winter, B. (2019). The Canadian Earth System Model version 5 (CanESM5.0.3). *Geoscientific Model Development*, 12(11):4823–4873.

[Séférian et al.2019] Séférian, R., Nabat, P., Michou, M., Saint-Martin, D., Voldoire, A., Colin, J., Decharme, B., Delire, C., Berthet, S., Chevallier, M., Sénési, S., Franchisteguy, L., Vial, J., Mallet, M., Joetzjer, E., Geoffroy, O., Guérémy, J.-F., Moine, M.-P., Msadek, R., Ribes, A., Rocher, M., Roehrig, R., Salas-y Mélia, D., Sanchez, E., Terray, L., Valeke, S., Waldman, R., Aumont, O., Bopp, L., Deshayes, J., Éthé, C., and Madec, G. (2019). Evaluation of CNRM Earth System Model, CNRM-ESM2-1: Role of Earth System Processes in Present-Day and Future Climate. *Journal of Advances in Modeling Earth Systems*, 11(12):4182–4227.

[Terhaar et al.2022] Terhaar, J., Frölicher, T. L., and Joos, F. (2022). Observation-constrained estimates of the global ocean carbon sink from Earth system models. *Biogeosciences*, 19(18):4431–4457.

[Toyama et al.2017] Toyama, K., Rodgers, K. B., Blanke, B., Iudicone, D., Ishii, M., Aumont, O., and Sarmiento, J. L. (2017). Large Reemergence of Anthropogenic Carbon into the Ocean’s Surface Mixed Layer Sustained by the Ocean’s Overturning Circulation. *Journal of Climate*, 30(21):8615–8631.

[Wang et al.2016] Wang, L., Huang, J., Luo, Y., and Zhao, Z. (2016). Narrowing the spread in CMIP5 model projections of air-sea CO2 fluxes. *Scientific Reports*, 6(1):37548.

[WCRP2021] WCRP (2021). CMIP6, <https://esgf-node.llnl.gov/search/cmip6>.

[Yool et al.2021] Yool, A., Palmieri, J., Jones, C. G., de Mora, L., Kuhlbrodt, T., Popova, E. E., Nurser, A. J. G., Hirschi, J., Blaker, A. T., Coward, A. C., Blockley, E. W., and Sellar, A. A. (2021). Evaluating the physical and biogeochemical state of the global ocean component of UKESM1 in CMIP6 historical simulations. *Geoscientific Model Development*, 14(6):3437–3472.

[Zeng et al.2015] Zeng, J., Nojiri, Y., Nakaoka, S.-i., Nakajima, H., and Shirai, T. (2015). Surface ocean CO2 in 1990–2011 modelled using a feed-forward neural network. *Geoscience Data Journal*, 2(1):47 – 51.

[Zickfeld et al.2016] Zickfeld, K., Macdougall, A. H., and Matthews, H. D. (2016). On the proportionality between global temperature change and cumulative CO2 emissions during periods of net negative CO2 emissions. *Environmental Research Letters*, 11:055006.

[Ziehn et al.2020] Ziehn, T., Chamberlain, M., Law, R., Lenton, A., Bodman, R., Dix, M., Stevens, L., Wang, Y.-P., and Jhan, J. S. (2020). The Australian Earth System Model: ACCESS-ESM1.5. *Journal of Southern Hemisphere Earth Systems Science*, 70:193–214.

8. Tables

Table 1. Earth System Models used, references, number of ensemble members for each SSP scenario and color in Figure 3.

Earth System Model	Reference	SSP1 1.9	SSP1 2.6	SSP2 4.5	SSP5 3.4	SSP5 8.5	Color in Fig 3
ACCESS-ESM1-5	[Ziehn et al.2020]	-	9	30	-	10	purple
CanESM5	[Swart et al.2019]	6	-	-	-	-	orange
CNRM-ESM2-1	[Boucher et al.2020]	5	5	10	5	5	red
IPSL-CM6A-LR	[Boucher et al.2020]	6	6	11	2	6	blue
MIROC-ES2L	[Hajima et al.2020]	5	3	-	-	-	pink
MPI-ESM1-2-LR	[Mauritsen et al.2019]	-	10	10	-	10	light blue
UKESM1-0-LL	[Sellar et al.2019]	5	16	16	5	6	green

Table 2. Observation-based products used in this study

	Reference
CMEMS-FFNN	[Denvil-Sommer et al.2019]
CSIR-ML6	[Gregor et al.2019]
JENA-MLS	[Rödenbeck et al.2022]
JMA-MLR	[Iida et al.2021]
LDEO-HPD	[Gloege et al.2022a, Bennington et al.2022b]
MPI-SOMFFN	[Landschützer et al.2020]
NIES-FNN	[Zeng et al.2015]
pCO ₂ -Residual	[Bennington et al.2022a]

Table 3. 2100 Ocean Carbon Ratio (Sink/Emissions). CMIP6 Ensemble Means. Adjustment for each model ensemble is accomplished by adding to all years the difference from 3 PgC/yr in 2020.

CMIP6 Ratio	SSP1 1.9	SSP1 2.6	SSP2 4.5	SSP5 3.4	SSP5 8.5
Unadjusted Max	1.12	0.52	0.33	0.46	0.20
Unadjusted Min	0.75	0.44	0.29	0.39	0.17
spread	0.37	0.08	0.04	0.07	0.03
Adjusted Max	1.19	0.61	0.34	0.47	0.20
Adjusted Min	1.04	0.52	0.33	0.44	0.19
spread	0.15	0.09	0.01	0.03	0.01

Table 4. 2100 Ocean Carbon Sink Spread Across CMIP6 Ensemble Means. Adjustment for each model ensemble is accomplished by adding to all years the difference from 3 PgC/yr in 2020. Spread is the maximum minus minimum at 2100.

CMIP6 Models	SSP1 1.9	SSP1 2.6	SSP2 4.5	SSP5 3.4	SSP5 8.5
Unadjusted Max (PgC)	139	171	264	232	428
Unadjusted Min (PgC)	93	144	233	196	377
spread (PgC)	46	27	31	35	51
Adjusted Max (PgC)	147	197	278	238	431
Adjusted Min (PgC)	128	170	267	223	403
spread (PgC)	19	27	11	15	27
Change of Spread (PgC)	-27	0	-20	-20	-24
Change of Spread (%)	-59%	0%	-66%	-58%	-46%

Table 5. 2100 Ocean Carbon Sink Spread Across MAGICC7.0 Ensemble Members. Adjustment for each model ensemble is accomplished by adding to all years the difference from 3 PgC/yr in 2020. Spread is the maximum minus minimum at 2100.

MAGICC7.0	SSP1	SSP1	SSP2	SSP5	SSP5
	1.9	2.6	4.5	3.4	8.5
Unadjusted Max (PgC)	170	119	268	232	435
Unadjusted Min (PgC)	115	76	189	159	314
spread (PgC)	55	43	79	73	121
Adjusted Max (PgC)	211	169	294	262	456
Adjusted Min (PgC)	182	135	261	227	387
spread (PgC)	29	34	33	35	69
Change of Spread (PgC)	-26	-9	-46	-38	-52
Change of Spread (%)	-47%	-21%	-58%	-52%	-43%

9. Figures






Figure 1. Cumulative CO₂ emissions and ocean sink 2015-2100 for CMIP6 (color) and MAGICC7.0 (gray). (a) Cumulative emissions from 2015, (b) Cumulative ocean carbon sink, (c) Adjusted ocean sink, with 2020 sink set to 3 PgC/yr, (d) CSF_{ocean} unadjusted, (e) CSF_{ocean} adjusted. Emission and sink rates are shown in Figure S1. For clarity, MAGICC7.0 for SSP5-3.4 and SSP1-2.6 are shown in Figure S2. Adjustment for each model ensemble is accomplished by adding to all years the difference from 3 PgC/yr in 2020.



26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60

Figure 2. Mapped air-sea CO₂ fluxes, present and future difference. (a) Mean flux from 8 observation-based products (Table 2) for the 2010s (2010-2019), (b) IPSL-CM6A-LR for 2010s, (c) flux change from the 2010s to the 2050s in IPSL under SSP1-1.9, SSP2-4.5 and SSP5-8.5, (d) as (c) for change to the 2090s. Means for other models in the 2010s are compared to the observation-based products in Figure S3.




Figure 3. Zonal mean fluxes. (a) Fluxes in 2010s for 7 CMIP6 models (colors) compared to observation-based products (black) with 2σ spread (gray), (b) change to 2050s under SSP5-8.5 (note shifted zero line), (c) change to 2050s under SSP2-4.5, (d) change to 2050s under SSP1-1.9, (e) as (b) for 2090s, (f) as (c) for 2090s, (g) as (d) for 2090s. The x-axis has the same absolute scale on all panels to allow direct comparison of magnitudes.

10. Supplementary Figures




Figure S1. Emission and sink rates. (a) Emissions under the 5 SSP scenarios considered here, (b) Ocean carbon sink rate for CMIP6 ensemble means (color) and MAGICC7.0 (gray), (c) Ocean carbon sink rate after adjustment of all models to 2020 sink rate of 3 PgC/yr [Friedlingstein et al.2021].




Figure S2. Same as Figure 1(a,b,c), but showing MAGICC7.0 for SSP5-3.4 (light gray) and SSP1-2.6 (dark gray) in panels (b) and (c).




Figure S3. Comparison of 2010s mean air-sea CO₂ flux. (a) Mean of eight observation-based products (Table 2) and (b-f) CMIP6 models (Table 1).