The association of cusp-aligned arcs with plasma in the magnetotail implies a closed magnetosphere

Stephen E. Milan¹, Michaela K Mooney¹, Gemma E. Bower¹, Matthew G. G. T. Taylor², Larry J. Paxton³, Iannis S Dandouras⁴, Andrew Fazakerley⁵, Christopher M. Carr⁶, Brian J. Anderson⁷, and Sarah Kimberly Vines⁸

¹University of Leicester
²European Space Agency
³Johns Hopkins University
⁴Institut de Recherche en Astrophysique et Planétologie, UPS-CNRS Toulouse, France
⁵University College London, UK
⁶Imperial College
⁷John Hopkins Univ.
⁸Johns Hopkins University Applied Physics Laboratory

February 20, 2023

Abstract

We investigate a fifteen-day period in October 2011. Auroral observations by the SSUSI instrument onboard the DMSP F16, F17, and F18 spacecraft indicate that the polar regions were covered by weak cusp-aligned arc emissions whenever the IMF clock angle was small, $|\vartheta| < 45^{\circ}$, which amounted to 30% of the time. Simultaneous observations of ions and electrons in the tail by the Cluster C4 and Geotail spacecraft showed that during these intervals dense (1 cm-3) plasma was observed, even as far from the equatorial plane of the tail as |ZGSE| = 13 RE. The ions had a pitch angle distribution peaking parallel and antiparallel to the magnetic field and the electrons had pitch angles that peaked perpendicular to the field. We interpret the counter-streaming ions and double loss-cone electrons as evidence that the plasma was trapped on closed field lines, and acted as a source for the cusp-aligned arc emission across the polar regions. This suggests that the magnetosphere was almost entirely closed during these periods. We further argue that the closure occured as a consequence of dual-lobe reconnection at the dayside magnetopause. Our finding forces a significant re-evaluation of the magnetic topology of the magnetosphere during periods of northwards IMF.

The association of cusp-aligned arcs with plasma in the magnetotail implies a closed magnetosphere

S. E. Milan^{1*}, M. K. Mooney¹, G. E. Bower¹, M. G. G. T. Taylor², L. J. Paxton³, I. Dandouras⁴, A. N. Fazakerley⁵, C. M. Carr⁶, B. J. Anderson⁷, and S. K. Vines⁷

6	¹ School of Physics and Astronomy, University of Leicester, Leicester, UK.
7	² ESA/ESTEC, Noordwijk, The Netherlands.
8	³ William B. Hanson Center for Space Sciences, University of Texas at Dallas, USA.
9	⁴ Institut de Recherche en Astrophysique et Planétologie (IRAP), CNRS and Université Paul Sabatier,
10	Toulouse, France.
11	⁵ Mullard Space Science Laboratory, University College, London, UK.
12	⁶ Imperial College, London, UK.
13	⁷ Johns Hopkins University, Applied Physics Laboratory, USA.

Key Points:

1

2

3

4

14

15	•	Cusp-aligned arcs observed by the DMSP spacecraft occur frequently for north-
16		ward IMF
17	•	Cluster and Geotail observations show that the arcs are accompanied by trapped
18		plasma at high latitudes in the magnetotail

We interpret cusp-aligned arcs as a signature of a magnetosphere almost entirely
 closed by dual-lobe reconnection

 $^{^{*}\}mbox{School}$ of Physics and Astronomy, University of Leicester, Leicester LE1 7RH, UK

Corresponding author: Steve Milan, steve.milan@le.ac.uk

21 Abstract

We investigate a fifteen-day period in October 2011. Auroral observations by the SSUSI 22 instrument onboard the DMSP F16, F17, and F18 spacecraft indicate that the polar re-23 gions were covered by weak cusp-aligned arc emissions whenever the IMF clock angle was 24 small, $|\theta| < 45^{\circ}$, which amounted to 30% of the time. Simultaneous observations of ions 25 and electrons in the tail by the Cluster C4 and Geotail spacecraft showed that during 26 these intervals dense ($\approx 1 \text{ cm}^{-3}$) plasma was observed, even as far from the equatorial 27 plane of the tail as $|Z_{GSE}| \approx 13 R_E$. The ions had a pitch angle distribution peaking 28 parallel and antiparallel to the magnetic field and the electrons had pitch angles that peaked 29 perpendicular to the field. We interpret the counter-streaming ions and double loss-cone 30 electrons as evidence that the plasma was trapped on closed field lines, and acted as a 31 source for the cusp-aligned arc emission across the polar regions. This suggests that the 32 magnetosphere was almost entirely closed during these periods. We further argue that 33 the closure occured as a consequence of dual-lobe reconnection at the dayside magne-34 topause. Our finding forces a significant re-evaluation of the magnetic topology of the 35 magnetosphere during periods of northwards IMF. 36

³⁷ Plain Language Summary

The magnetosphere is usually assumed to contain both open and closed magnetic 38 flux. Closed magnetic field lines have both ends connected to the Earth; open field lines 39 connect to the Earth at one end and into the interplanetary medium at the other. There 40 tends to be little plasma on open field lines as the particles escape down the magneto-41 tail, whereas plasma on closed field lines is trapped. Open flux near the poles naturally 42 explains the oval configuration of Earth's auroras, with a lack of auroras at very high 43 latitudes where there is no plasma to cause emissions. Somewhat unexpectedly, we show 44 that auroral emission near the poles is common and that at these times there is signif-45 icant plasma in the magnetotail, indicating that the magnetosphere contains only closed 46 flux. We propose that this magnetic configuration is formed by a process known as dual-47 lobe magnetic reconnection which occurs when the interplanetary magnetic field within 48 the solar wind points northwards. We must re-evaluate the standard picture of magne-49 tospheric structure during these periods of northwards interplanetary magnetic field. 50

51 **1** Introduction

In this study we investigate repeated occurrences of cusp-aligned arcs (CAAs), the poorly understood situation in which weak auroral emissions fill the polar regions, during a fifteen-day interval in 2011. We conclude that CAAs are a common occurrence during periods of northward interplanetary magnetic field (IMF), and that they are a signature of a nearly- or entirely-closed magnetosphere. This forces a significant re-evaluation of the magnetic topology of the magnetosphere during northward IMF.

The coupling between the solar wind and the magnetosphere is relatively well un-58 derstood for southward-directed IMF: as first proposed by Dungey (1961), magnetic re-59 connection near the subsolar magnetopause opens previously-closed magnetic flux which 60 then forms the magnetotail lobes; thereafter, magnetic reconnection in the central plane 61 of the tail recloses flux which returns to the dayside, resulting in the Dungey cycle of mag-62 netospheric convection. The northern and southern ionospheric polar caps, the dim re-63 gions encircled by the auroral ovals, are the ionospheric projection of the open flux form-64 ing the lobes, and their size can be used to quantify the open flux content of the mag-65 netosphere, F_{PC} (e.g., Milan et al., 2003). This naturally explains many aspects of mag-66 netospheric structure and dynamics, including the formation of the magnetotail, the twin-67 cell ionospheric convection pattern, the morphology of the auroral oval, the plasma pop-68 ulations within the magnetosphere, and the evacuated magnetotail lobes. When time-69 varying magnetic reconnection rates are considered, the expanding/contracting nature 70

of the polar cap and the substorm cycle can be understood (Siscoe & Huang, 1985; Cow-71 ley & Lockwood, 1992; Lockwood & Cowley, 1992). Typically, F_{PC} varies between 0.3 72 and 0.9 GWb (Milan et al., 2003, 2007, 2021). However, it can become as low as 0.2 GWb 73 $(\sim 2.5\%)$ of the 8 GWb associated with the terrestrial dipole) during particularly extreme 74 nightside reconnection events, accompanied by a near-total in-filling of the polar regions 75 by bright auroral emission (e.g., Milan et al., 2004). In this paper we show that in fact 76 the polar cap closes entirely, frequently, and with much less fanfare, during periods of 77 northwards IMF. 78

79 The coupling during northward-directed IMF is still poorly understood. Although Dungey (1963) correctly proposed that magnetic reconnection would take place at the 80 high latitude magnetopause, tailwards of the cusps, the ramifications are still not fully 81 resolved. The list of northward-IMF (IMF $B_Z > 0$ or NBZ) phenomena includes: single-82 and dual-lobe reconnection (e.g., Cowley, 1981; Fuselier et al., 2012), reverse convection 83 in the polar cap (e.g., Huang et al., 2000), NBZ field-aligned currents poleward of the 84 noon auroral oval (e.g. Iijima et al., 1984), cusp auroral spots (e.g., Milan et al., 2000; 85 Frey et al., 2002; Frey, 2007; Carter et al., 2018, 2020), transpolar or polar cap arcs (e.g., 86 Frank et al., 1982; Cumnock et al., 2002; Kullen et al., 2002; Milan et al., 2005; Fear et 87 al., 2014), horse-collar auroras (e.g., Hones Jr et al., 1989; Milan, Carter, Bower, et al., 88 2020; Bower, Milan, Paxton, & Anderson, 2022), and cusp-aligned arcs (e.g., Y. Zhang 89 et al., 2016; Q.-H. Zhang et al., 2020; Milan et al., 2022; Wang et al., 2023). The inter-90 ested reader is directed to the recent reviews of NBZ phenomena by Hosokawa et al. (2020) 91 and Fear (2021). 92

A key question regarding periods of NBZ is the degree to which the magnetosphere 93 loses open magnetic flux, and the resulting distribution of open and closed flux in the 94 polar regions. It has been suggested that polar cap arcs may be associated with both 95 open and closed magnetic flux (e.g., Carlson & Cowley, 2005; Reidy et al., 2020; Bower, 96 Milan, Paxton, & Imber, 2022). Evidence suggests that the most prominent of polar cap 97 arcs, those also known as theta auroras, are likely associated with closed magnetic flux 98 (Fear et al., 2014; Fryer et al., 2021; Coxon et al., 2021), proposed to be produced by 99 magnetic reconnection in the magnetotail (Milan et al., 2005; Fear & Milan, 2012a, 2012b). 100 The situation regarding less prominent sun-aligned or cusp-aligned arcs, in which mul-101 tiple weak arcs fill the polar regions (Y. Zhang et al., 2016), is as yet unresolved. If these 102 form on the open flux of the largely-evacuated magnetotail lobes, then the source of the 103 auroral precipitation is called into question, though Carlson and Cowley (2005) proposed 104 that polar rain could provide sufficient plasma to be accelerated in flow shears in the iono-105 spheric convection pattern to produce such arcs. 106

During prolonged NBZ, the polar cap can contract and become teardrop-shaped, 107 leading to the horse-collar auroras (HCAs) configuration (Hones Jr et al., 1989). Milan, 108 Carter, Bower, et al. (2020) and Bower, Milan, Paxton, and Anderson (2022) have sug-109 gested that this is formed by the closure of magnetic flux by dual-lobe reconnection (DLR) 110 for near-zero IMF clock angle, supported by numerical simulations (Wang et al., 2023). 111 Dual-lobe reconnection should be an efficient mechanism by which the magnetosphere 112 can capture solar wind plasma (e.g., Imber et al., 2006), and it has been proposed that 113 over time this can lead to the formation of a cold, dense plasma sheet (CDPS) (Øieroset 114 et al., 2005). However, this is not the only means by which solar wind plasma is thought 115 to enter the magnetosphere during periods of NBZ, with inward diffusion at the mag-116 netotail flanks (see discussion in Taylor et al., 2008) or direct entry through (single) lobe 117 reconnection (Shi et al., 2013; Mailyan et al., 2015) being other commonly discussed mech-118 anisms. It is unclear, however, how efficient diffusion can be, how quickly captured so-119 lar wind plasma can be redistributed throughout the magnetosphere, and why the plasma 120 remains near the Earth and does not escape back to the solar wind down the open field 121 lines of the lobes, as it does for southwards IMF. 122



Figure 1. (Left) The orbit of Cluster C4 (red) and Geotail (green) for the period of study, in the GSE X - Z plane. The T96 magnetic field configuration on DOY 280, 2011, at 13 UT, for $P_{dyn} = 2$ nPa, $D_{st} = -10$ nT, $B_Y = 0$ and $B_Z = 5$ nT is superimposed in grey. (Middle) The orbit of Cluster and Geotail for the period of study, in the GSE Y - Z plane; the location of the magnetopause at X = -10 R_E is shown for reference. (Right) Field-line tracing from the location of Cluster and Geotail to the northern (orange and dark green) and southern (red and light green) hemisphere ionospheres for the period of interest, presented on a geomagnetic latitude and MLT grid. An average auroral oval for $K_P = 1$ is overlaid for reference.

The cusp-aligned arcs (hereafter CAAs) auroral configuration is perhaps the most 123 poorly-studied NBZ auroral phenomenon. Partially, this is because the emissions tend 124 to be weak, so are better observed from the ground (e.g., Ismail et al., 1977; Hosokawa 125 et al., 2011), with limited geographical coverage, rather than from space. Recently, Q.-126 H. Zhang et al. (2020) and Wang et al. (2023) suggested that CAAs are produced by plasma 127 flow shears introduced into the magnetosphere by Kelvin-Helmholtz surface waves ex-128 cited on the magnetotail flanks by the flow of the solar wind. However, this does not re-129 solve the source of the precipitating plasma. On the other hand, Milan et al. (2022) pro-130 posed that if DLR continues for a prolonged period, then the magnetosphere can become 131 almost entirely closed and the horse-collar auroral configuration can develop to the point 132 where the polar slot (distorted polar cap) can almost disappear. In this scenario, the closed 133 magnetosphere will be filled with trapped solar wind plasma and, according to Milan et 134 al. (2022), flow shears produced by lobe reconnection can then accelerate this trapped 135 plasma into the atmosphere to form the CAAs. Whereas Carlson and Cowley (2005) sug-136 gested that the source of plasma to produce weak polar cap arcs was polar rain that had 137 fled to the distant magnetotail along open field lines, instead Milan et al. (2022) proposed 138 that the source is solar wind plasma captured by DLR and trapped closer to the Earth 139 on closed field lines. 140

In this study we examine plasma populations observed in the near-Earth tail by 141 the C4 spacecraft of the Cluster constellation (Escoubet et al., 2001) and the Geotail (Nishida, 142 1994) spacecraft during a prolonged period in October 2011 when auroral observations 143 from F16, F17, and F18 of the Defense Meteorological Spacecraft Program detected mul-144 tiple instances of CAAs. Dense plasma is repeatedly found in regions that would nor-145 mally be occupied by the evacuated open field lines of the northern and southern lobes. 146 and it is concluded that the magnetosphere is almost entirely closed and the plasma is 147 trapped, providing a source for the high latitude auroral emission. 148

¹⁴⁹ 2 Observations

The period under investigation is the 3 to 17 October 2011 (days-of-year 276 to 290), 150 which encompasses six orbits of the Cluster constellation during its tail season and three 151 orbits of Geotail; the orbits are shown in Figure 1. At this time, the orbit of C4 was in-152 clined such that it was southwards of the neutral sheet for most of the time, except near 153 perigee. During the study interval, C4 reached $Z_{GSE} \approx -13 R_E, X_{GSE} \approx -14 R_E$ 154 at apogee, and it was located relatively centrally in the tail, $0 < Y_{GSE} < 10 R_E$, in 155 the first half of each orbit, but closer to the dusk flank in the second. The orbit of Geo-156 157 tail was such that it sampled the magnetotail northwards of the neutral sheet, up to $Z_{GSE} \approx$ 8 R_E at $X_{GSE} \approx -19 R_E$. A representative magnetic field tracing in the T96 model 158 (Tsyganenko, 1995), with input parameters $P_{dyn} = 2 \text{ nT}$, $D_{st} = -10 \text{ nT}$, and IMF 159 $B_Y = 0$ and $B_Z = 5$ nT, is overlaid for reference. These parameters were chosen to 160 represent a quiet NBZ magnetosphere, and will be used throughout the rest of the study 161 to compare with magnetic field observations. As the plasma sheet is usually confined to 162 $|Z| < 5 R_E$, for much of their orbits Geotail and C4 are expected to be within the north-163 ern and southern lobes of the magnetotail, respectively. Field line tracings from the lo-164 cations of the two spacecraft to the northern and southern ionospheres are also presented, 165 with a $K_P = 1$ average auroral oval superimposed for reference. This indicates that near 166 apogee C4 would normally be expected to map to the central polar cap in the southern 167 hemisphere, and that Geotail would frequently map to the nightside polar cap in the north-168 ern hemisphere. 169

Auroral observations during this period are provided by the Special Sensor Ultra-170 violet Spectrographic Imager or SSUSI experiment (Paxton et al., 1992) onboard the DMSP-171 F16, -F17, and -F18. The DMSP spacecraft have sun-synchronous orbits near an alti-172 tude of 850 km. SSUSI measured auroral luminosity in a swath either side of the space-173 craft orbit in the Lyman-Birge-Hopfield short (LBHs) band, 140 to 150 nm (see Paxton 174 and Zhang (2016); Paxton et al. (2017, 2021) and the references cited therein for further 175 description of the instrument and data products). Measurements of the distribution of 176 magnetosphere-ionosphere field aligned currents (FACs) were provided by the Active Mag-177 netosphere and Planetary Electrodynamics Response Experiment (AMPERE) technique 178 (Anderson et al., 2000; Coxon et al., 2018; Waters et al., 2001). We also make use of iono-179 spheric flow observations provided by the Super Dual Auroral Radar Network (Super-180 DARN (Chisham et al., 2007)) and the Ion Driftmeter (IDM) component of the Special 181 Sensors–Ions, Electrons, and Scintillation thermal plasma analysis package or SSIES in-182 strument onboard the Defense Meteorological Satellite Program spacecraft (DMSP/IDM 183 (Rich & Hairston, 1994)). Solar wind parameters were taken from the OMNI data-set 184 (King & Papitashvili, 2005). 185

Figure 2 shows 12 snapshots of the auroral morphology from DOYs 278 to 280. Be-186 low this, a keogram of the observations along the dawn-dusk meridian made by DMSP-187 F16/SSUSI in the northern hemisphere is shown, along with the IMF clock angle from 188 OMNI. Around 10:20 UT on DOY 278 (panels a and b), the IMF had a southwards com-189 ponent and typical twin-cell ionospheric flows were observed by SuperDARN and DMSP/IDM 190 (not shown for brevity). The auroral morphology was also typical, showing an oval sur-191 rounding a dim polar cap, with evidence for substorm activity on the nightside. Between 192 approximately 16:00 UT, DOY 278, and 13:00 UT, DOY 279, the IMF turned northwards. 193 During this period the auroras dimmed, contracted to higher latitudes and acquired a 194 horse-collar auroral configuration, before the polar regions filled with auroral emission 195 mainly in the form of cusp-aligned arcs or CAAs (c to f). CAAs are seen in both north-196 ern and southern hemispheres, panels c and e being from the north and d and f being 197 from the south. Then the IMF turned southwards again, twin-cell convection resumed. 198 and the polar cap reopened (g and h). There followed another period of NBZ, during 199 which CAAs reformed (i and j), before again a southward turning and a reopening of 200 the polar cap (k to l). The lower panels clearly show the expansions and contractions 201



Figure 2. Snapshots of the LBHs auroral configuration in the northern and southern hemispheres observed by DMSP/SSUSI onboard DMSP-F16, -F17, and -F18 between 5 and 7 October 2011. Each panel is presented in a geomagnetic latitude and local time format, with noon towards the top and dawn towards the right. Grey circles indicate geomagnetic latitude in steps of 10°. Of the two lower panels, the top one shows a keogram of auroral emissions observed by DMSP-F16/SSUSI along the dawn-dusk meridian of the northern hemisphere; grey vertical bars indicate where data is missing. Red horizontal bars indicate the times that cusp-aligned arcs were observed. The bottom panel is clock angle, highlighted in red when $|\theta| < 45^{\circ}$.

of the auroral oval with changes in clock angle, and the presence of auroral emissions at high latitudes during periods when the clock angle is near zero, $|\theta| < 45^{\circ}$, highlighted in red.

Over the rest of the period considered several other intervals of CAAs were found, 205 as will be indicated in later figures. The start and end times of these intervals are ap-206 proximate due to the relatively coarse cadence of the DMSP orbits; there are also pe-207 riods when it was not possible to positively identify whether CAAs were present or not, 208 due to only partial coverage of the polar regions by the SSUSI field-of-view. During pe-209 riods when HCAs or CAAs were present, the ionospheric flows measured by SuperDARN 210 and DMSP/IDM and the field-aligned currents observed by AMPERE were consistent 211 with the observations reported by Milan, Carter, Bower, et al. (2020) and Milan et al. 212 (2022) – reverse lobe convection, NBZ FACs at noon – suggesting that dual-lobe recon-213 nection was responsible for closing the magnetosphere. The occurrence of CAAs when 214 $|\theta| < 45^{\circ}$ is also consistent with the statistical occurrence of horse-collar auroras (Bower, 215 Milan, Paxton, & Anderson, 2022), thought to be the auroral precursor to CAAs. 216

We use Cluster C4 observations from the Composition and Distribution Function 217 analyser of the Cluster Ion Spectrometry instrument (C4/CIS-CODIF (Rème et al., 1997)), 218 the Plasma Electron And Current Experiment (C4/PEACE (Johnstone et al., 1997)), 219 and the Fluxgate Magnetometer (C4/FGM (Balogh et al., 1997)). Figure 3 covers the 220 period 3 to 9 October, spanning the first three C4 orbits considered and encompassing 221 the interval shown in Figure 2. The panels are presented in the following order. (a) The 222 GSE X, Y, and Z (R_E) position of C4. (b) The B_X , B_Y , B_Z components of the mag-223 netic field measured by C4/FGM in GSE coordinates. Dots indicate predications of the 224 magnetic field measurements by the T96 (Tsyganenko, 1995) model, with fixed inputs 225 (as before). (c) The proton density, n_i , observed by C4/CIS-CODIF. Vertical red bars 226 indicate when CAAs were observed by DMSP/SSUSI. (d) The proton differential energy 227 flux spectrogram measured by C4/CIS-CODIF. (e) The electron differential energy flux 228 spectrogram measured by C4/PEACE. (f) A dawn-dusk keogram of auroral emissions 229 observed by DMSP-F16/SSUSI in the northern hemisphere. (g) A dawn-dusk keogram 230 of FAC density measured by AMPERE in the northern hemisphere, with red/blue in-231 dicating up/down currents. (h) The clock angle of the IMF, θ , highlighted in red when 232 $|\theta| < 45^{\circ}$. (i) The GSM B_X, B_Y, B_Z components of the IMF from OMNI. (j) The so-233 lar wind speed and density from OMNI. 234

The SSUSI keogram (f) reveals that at most times there is little auroral emission inside the auroral oval, consistent with an open polar cap, except at the times of CAAs (red bars in (c)) when $|\theta| < 45^{\circ}$. The region 1/2 FACs (Iijima & Potemra, 1976) observed by AMPERE (g) are enhanced when the polar cap is open, indicating Dungey cycle driving of the magnetosphere by subolar reconnection, especially during periods with IMF $B_Z < 0$ nT, $|\theta| > 90^{\circ}$. When CAAs are present the R1/R2 FACs are weak, though NBZ FACs tend to be observed at noon (not shown).

Near the perigee of each orbit (marked PG) when $|Z| < 5 R_E$, C4 passed through 242 the plasma sheet and ring current regions and enhanced proton densities, n_i , were ob-243 served (c). During the first orbit of Figure 3 almost no plasma was observed when Z <244 $-5 R_E$, consistent with the open lobe. In contrast, while C4 was near perigee on DOY 245 278 the IMF turned northwards and CAAs were observed, and throughout this period 246 C4 was engulfed in protons with $n_i \approx 1 \text{ cm}^{-3}$ up to near apogee (AG) on DOY 279. 247 As the IMF turned southwards around 13:00 UT on DOY 279, the density returned to 248 typical lobe values, $n_i < 0.1 \text{ cm}^{-3}$. Around 00 UT on DOY 280 CAAs were once again 249 observed, and n_i rose to near 1 cm⁻³, only to drop again at 04 UT as the IMF turned 250 southwards; the IMF remained predominantly southwards for the duration of the third 251 orbit, and n_i remained low throughout (except for a short period near apogee, see be-252 low). 253



Figure 3. Observations from 3 to 10 October 2011, which encompasses three orbits by Cluster. (a) The GSE position of C4. Apogees and perigees of the orbit are indicated by AG and PG. Orange highlighting indicates the time of the first Geotail orbit shown in Figure 6. (b) C4/FGM observations of the magnetic field. T96 model predictions for fixed input parameters are indicated as dots. (c) C4/CIS-CODIF ion density measurements. Red bars indicate periods of observation of cusp-aligned arcs by DMSP/SSUSI. (d) C4/CIS-CODIF ion spectrogram. (e) C4/PEACE electron spectrogram. (f) A dawn-dusk keogram of auroral observations by DMSP-F16/SSUSI in the northern hemisphere. (g) A dawn-dusk keogram of field-aligned currents measured by AMPERE, red and blue being upward and downward FACs, respectively. (h) The clock angle of the IMF from OMNI. The clock angle is highlighted in red when $|\theta| < 45^{\circ}$. (i) The components of the IMF from OMNI. (j) The solar wind speed and density from OMNI.

The C4/CIS-CODIF spectrogram (d) indicates that the CAA-related ions had energies between several 100s and several 1000s eV. (We note that at the apogees of the second and third orbits a brief interval of cold, < 100 eV, ions was also observed.) The C4/PEACE spectrogram (e) indicates that the ions were accompanied by electrons, with energies below about 1 keV. In the first CAA interval of DOY 279 the ions, and especially the electrons, reduced in energy with time, indicating a cooling of the plasma.

Figure 4 presents the next three orbits of C4. During this period, the IMF $|\theta| <$ 45° for much of the time, and there were several intervals of CAAs observed by DMSP/SSUSI. Accompanying these intervals, C4/CIS-CODIF saw elevated n_i , where otherwise lobe conditions might have been expected. During DOYs 287 to 290, there were repeated swings between $|\theta| < 45^{\circ}$ and $|\theta| > 45^{\circ}$, and CAAs and protons came and went in tandem.

Figure 5 focusses on the two intervals of CAAs presented in Figures 2 and 3. In 265 addition to the spectrograms, pitch angle distributions of the ions (d) and electrons (f)266 are shown. The proton pitch angle fluxes are integrated across the full energy range of 267 the CIS-CODIF instrument; the electron fluxes are calculated from the low energy elec-268 tron analyser head of the PEACE instrument, limited to energies above 100 eV to re-269 move the contribution of photoelectrons. During periods of CAAs, and when C4 was not 270 too close to the Earth, ion pitch angles were concentrated at 0° and 180° , indicating two 271 counter-streaming populations. The electron pitch angles, conversely, peaked at 90° , with 272 a distinct lack of electrons near 0° and 180° , indicating a double loss-cone distribution. 273 The double loss-cone electron distribution is similar to that observed by Fear et al. (2014)274 above a transpolar arc, and is indicative of plasma trapped on closed field lines. The counter-275 streaming ions support this conclusion. We note that the ion and electron densities dur-276 ing the periods of CAAs were quite variable, indicating that the trapped plasma was not 277 uniformly distributed but was present continuously. The solar wind density was variable 278 during this period, but it is not clear if variations in ion/electron density and energy ob-279 served by C4 are correlated with enhancements in n_{SW} . As noted previously, the ion and 280 electron energies decreased overall with time, which might indicate the progressive mix-281 ing of plasma sheet and solar wind populations, leading to the formation of a cold, dense 282 plasma sheet (e.g., Øieroset et al., 2005). 283

Just prior to the second period of CAAs, CIS-CODIF detected a beam of low energy ions observed only near 0° pitch angles (i.e. flowing tailward). A similar beam was observed around 00 UT near apogee on DOY 282 (see Figure 3). As there are no counterstreaming ions, we suggest that at these are the signature of plasma escaping to the solar wind along open field lines.

Superimposed on the C4/FGM observations (a) is a model prediction from T96 for fixed NBZ input parameters (as before). The model tends to match the observations well during the periods of CAAs, but the B_X component was enhanced by ≈ 10 nT during the periods when CAAs were not present, indicating that the tail was more inflated at these times. This also supports the conclusion that the magnetotail open flux was significantly reduced when CAAs were observed.

Figure 6 presents two time intervals when Geotail was located in the magnetotail; 295 Figures 3 and 4 show that these roughly correspond to orbits 3 and 5 of C4. Panels a296 and b show that in each case Geotail entered the dusk flank of the magnetotail at the 297 start of the interval, near $Z \approx 5 R_E$, and rose to $Z \approx 8 R_E$ at $X \approx -19 R_E$, $Y \approx$ 298 $0 R_E$. It later exited the dayside magnetopause in the pre-noon sector. Panels e to l present 299 ion and electron spectrograms from the Low Energy Particle (LEP) instrument (Mukai 300 et al., 1994), showing fluxes in the sunward and tailward directions. During the first or-301 bit, the plasma sheet was seen as the spacecraft entered and exited the magnetosphere, 302 but the evacuated lobe was encountered from 16 UT on DOY 281 to 14 UT on DOY 282. 303 During this period, IMF $|\theta| > 45^{\circ}$ at all times (m). During the second orbit, the plasma 304 sheet was also seen at the start and the end, with periods of lobe between 23 UT, DOY 305



Figure 4. Observations from 10 to 17 October, presented in the same format as Figure 3.



Figure 5. Observations from 5 to 7 October 2011. (a) C4/FGM observations of the magnetic field. T96 model predictions for fixed input parameters are indicated as dots. (b) C4/CIS-CODIF ion density measurements. Red bars indicate periods of observation of cusp-aligned arcs by DMSP/SSUSI. (c, d) C4/CIS-CODIF ion spectrogram and pitch angle distribution. (e, f) C4/PEACE spectrogram and pitch angle distribution. (g) A dawn-dusk keogram of auroral observations by DMSP-F16/SSUSI. (h) The clock angle of the IMF from OMNI. The clock angle is highlighted in red when $|\theta| < 45^{\circ}$. (i) The solar wind speed and density from OMNI.



Figure 6. Two time periods during which Geotail was located within the magnetotail. (a, b) The GSE location of Geotail. (c, d) The magnetic field components measured by MGF, with a T96 prediction superimposed. Red bars indicate when cusp-aligned arcs were observed by DMSP/SSUSI. (e, f) Ion spectrograms of fluxes in the sunwards direction. (g, h) Ion spectrograms of fluxes in the tailwards direction. (i, j) Electron spectrograms of fluxes in the sunwards direction. (m, n) IMF clock angle, highlighted in red when $|\theta| < 45^{\circ}$.

286, and 18 UT, DOY 287. However, three intervals of plasma, with fluxes in both the 306 sunwards and tailwards directions, were detected at the same time as CAAs were ob-307 served by DMSP/SSUSI, associated with excursions to $|\theta| < 45^{\circ}$ (n). Plasma was also 308 seen by C4 in the southern tail at these times (Figure 4), and the plasma characteris-309 tics were similar. The presence of fluxes in both the sunwards and tailwards directions. 310 indicate that the plasma is trapped on closed field lines. During the second orbit, the 311 magnetic field observed by the Magnetic Field Experiment (MGF) instrument (Kokubun 312 et al., 1994), panel d, matched closely the prediction by T96. However, during the first 313 orbit, around the time Geotail was in the lobe the B_X component was elevated above 314 the prediction (c) indicating an inflated magnetotail. 315

316 **3 Discussion**

The occurrence of cusp-aligned arcs filling the polar regions is frequent. We find 317 that CAAs have a high probability of appearing if $|\theta| < 45^{\circ}$ for an appreciable length 318 of time (one to two hours or more). During our period of study, CAAs were observed 319 for approximately 30% of the time. Although polar cap arcs received attention in the 320 past (e.g., Ismail et al., 1977), CAAs are too dim to have been detected by previous gen-321 erations of global auroral imagers (e.g., IMAGE FUV), so their importance has been over-322 looked. We find that CAAs are accompanied by dense plasma of energies from several 323 eV to several 10s keV in regions of the magnetosphere (up to $|Z| \approx 13 R_E$ in our ob-324 servations) that would normally be occupied by the evacuated tail lobes. Shi et al. (2013) 325 interpreted similar observations of such plasma in the "lobes" as an indication of direct 326 ingress of solar wind plasma via (single) lobe reconnection. However, it seems unlikely 327 that this plasma should reside in the near-Earth tail for long, rather than disappearing 328 down the tail along open field lines, as occurs during periods of southwards IMF. Instead, 329 we suggest that this plasma must be trapped on closed field lines. This interpretation 330 is supported by the presence of a double loss-cone in the C4 electron pitch angle distri-331 butions, similar to the plasma characteristics seen by Fear et al. (2014) at high Z over 332 a transpolar arc. In that case, the plasma was only observed by Cluster on field lines that 333 mapped to the arc, and evacuated lobe was seen to either side; in our case the plasma 334 is observed for prolonged periods (sometimes many hours) wherever C4 is located in the 335 tail. The double loss-cone indicates that the plasma has interacted significantly with the 336 atmosphere in both the northern and southern hemispheres over multiple bounces. The 337 presence of counter-streaming ions observed by C4 is also consistent with closed field lines 338 and trapped plasma, as are the sunwards/tailwards fluxes observed by Geotail: if the mag-339 netic field was open, only tailward fluxes would be expected. This trapped plasma is ob-340 served in both the northern and southern portions of the magnetotail, at $Z \approx 8 R_E$ 341 and $Z \approx -13 R_E$, simultaneously (see Figures 4 and 6). 342

Several mechanisms have been proposed by which plasma enters the tail during NBZ 343 conditions, and whether or not it is trapped (e.g., Taylor et al., 2008; Shi et al., 2013). 344 Milan, Carter, Bower, et al. (2020) argued that dual-lobe reconnection explains both the 345 capture and trapping of plasma, and the ionospheric flow pattern and auroral evolution 346 observed during the formation of horse-collar auroras (HCAs), and by extension CAAs 347 (Milan et al., 2022). It also explains the necessity for near-zero clock angle, $|\theta| < 45^{\circ}$, 348 for the appearance of HCAs (Bower, Milan, Paxton, & Anderson, 2022) and CAAs (this 349 study). We suggest now that this also explains why the polar cap closes and the tail loses 350 its northern and southern lobes, explaining trapped plasma on field lines that map to 351 what would normally be the central polar cap. 352

We note that between 22 UT, DOY 285, and 08 UT, DOY 286, there were two brief swings of the clock angle to $|\theta| < 45^{\circ}$, but no CAAs were detected and open lobe was observed by C4 (see Figure 4). This suggests that there is a minimum duration of DLR of one to two hours for plasma to be trapped. This minimum will be related to the reconnection rate, that is the rate at which flux is closed, such that open lobe is replaced by closed flux containing trapped solar wind plasma. Detailed measurement of the rate
of closure, difficult with low cadence DMSP/SSUSI images, will be required to further
understand this. However, ionospheric convection measurements (e.g., Chisham et al.,
2004, 2008) should help quantify the reconnection rate.

The plasma trapped by DLR then acts as a source for precipitation to produce au-362 roral emission in the polar regions. As shown by Q.-H. Zhang et al. (2020) and Milan 363 et al. (2022), CAAs are produced by inverted-V precipitation, electrons accelerated to 364 energies of a few keV, at shears in the ionospheric convection flow which are associated 365 with upwards field-aligned currents due to converging ionospheric electric fields. This 366 is consistent with the trapped electron population found in this study: in the magneto-367 tail the electrons have energies primarily below one keV and the electron pitch angle dis-368 tribution has empty loss cones. These electrons will not precipitate without the accel-369 eration provided by flow shears. Once accelerated, the electrons are seen to have ener-370 gies of several keV in the ionosphere. Precipitating ions with energies of one to several 371 keV are also observed in the ionosphere when CAAs are present, especially towards the 372 nightside, consistent with the trapped ions found in this study. Whereas Carlson and 373 Cowley (2005) proposed that weak polar cap arcs, perhaps indeed CAAs, are produced 374 by accelerated polar rain on open field lines, in our scenario trapped solar wind plasma 375 on closed field lines is the more readily available source population. 376

On the large scale the ionospheric flow pattern has reverse lobe convection cells, 377 consistent with dual-lobe reconnection (Milan et al., 2022), though on smaller scales mul-378 tiple flow shears are seen, associated with the CAAs. Q.-H. Zhang et al. (2020) and Wang 379 et al. (2023) suggested that these flow shears are produced by Kelvin-Helmholtz waves 380 on the magnetospheric flanks propagating into the magnetotail. However, the cusp-aligned 381 nature of the arcs shows that the flow shears are also cusp-aligned, which is not neces-382 sarily predicted by the KHI mechanism. Instead, Milan et al. (2022) proposed that the 383 flow shears are excited by temporally- and spatially-varying lobe reconnection rates, which 384 explains why the flows and shears naturally radiate from the cusp region. We prefer this 385 latter explanation: lobe reconnection explains the closure of magnetic flux and the trap-386 ping of solar wind plasma, it explains the "reverse" flow pattern observed in the iono-387 sphere, it explains the structuring of the auroral precipitation into multiple arcs, and it 388 also explains why the arcs are "cusp-aligned". That the field is closed at high latitudes 389 also explains why CAAs are generally seen in both hemispheres simultaneously (see Fig-390 ure 2 and Milan et al. (2022), whereas other polar cap auroral phenomena are not al-391 ways conjugate (e.g., Reidy et al., 2020; Bower, Milan, Paxton, & Imber, 2022). 392

That auroral emission is observed across the polar regions suggests that the mag-393 netosphere is almost entirely closed, though it might be expected that open and closed 394 flux is interspersed (Milan et al., 2022). As proposed by Milan et al. (2022), horse-collar 395 auroras are the preliminary stage in the development of CAAs, forming when dual-lobe 396 reconnection first commences (Milan, Carter, Bower, et al., 2020). In other words, it is 397 possible that HCAs form frequently (Bower, Milan, Paxton, & Anderson, 2022) but do 398 not necessarily fully evolve into CAAs if the IMF turns away from near-zero clock an-399 gles. The high-latitude arcs which sit at the dawn and dusk edges of the polar slot of 400 the HCA configuration are dimmer than the main auroral oval (Bower, Milan, Paxton, 401 & Anderson, 2022), but are still bright enough to be seen with global auroral imagers 402 (Cumnock & Blomberg, 2004) and can be misinterpreted as transpolar arcs (Milan, Carter, 403 Bower, et al., 2020). However, these HCA arcs are in general brighter than CAAs and 404 so may be detected more frequently, certainly with previous global auroral imagers. As 405 mentioned previously, the occurrence of CAAs is under-studied and a statistical survey 406 is required, especially as this will provide new insights into the occurrence of DLR. 407

As an aside, we note that the magnitude of the B_X component of the IMF was significant at times, e.g. ≈ 10 nT during the first period of CAAs in Figure 3, and was nearzero at others, e.g. the last period of CAAs in Figure 4. This suggests that B_X does not ⁴¹¹ play a role in modulating the occurrence of (dual-) lobe reconnection and hence the oc-⁴¹² currence of CAAs. This tallies with a lack of B_X control of the occurrence of HCAs re-⁴¹³ ported by Bower, Milan, Paxton, and Anderson (2022).

⁴¹⁴ Once DLR ceases the magnetosphere loses the trapped plasma and the polar cap ⁴¹⁵ reforms promptly, indicating the rapid opening of magnetic flux by magnetopause re-⁴¹⁶ connection. This opening will occur most rapidly if the IMF is directed southwards and ⁴¹⁷ subsolar reconnection occurs, which will be accompanied by twin-cell convection in the ⁴¹⁸ polar ionosphere. However, as noted by Milan et al. (2022), it will also occur if the IMF ⁴¹⁹ is directed northwards with $|\theta| > 45^{\circ}$ as single lobe reconnection will open a closed mag-⁴²⁰ netosphere. Hence, CAAs are only seen for $|\theta| < 45^{\circ}$.

Empirical magnetic field models, such as T96, do not reproduce a closed magne-421 tosphere well, though closed NBZ magnetospheres are readily formed in simulations (e.g., 422 Song et al., 1999; Siscoe et al., 2011; Wang et al., 2023). The magnetic field measure-423 ments made by C4/FGM and Geotail/MGF indicated that the B_X component was re-424 duced when CAAs were present, with respect to when open lobe was observed. This in-425 dicates that the tail is somewhat deflated when it is closed (though it could also be in 426 part because plasma pressure contributes to stress balance in the closed tail, rather than 427 just magnetic pressure in the lobes). However, otherwise the field did not deviate markedly 428 from the T96 predictions, indicating that if the magnetosphere is indeed closed, the mag-429 netic structure near-Earth is not significantly modified, suggesting that the closed field 430 lines stretch considerably further down-tail than the locations of C4 and Geotail. This 431 is in contradiction to simulations which suggest that a closed magnetotail might be less 432 than 20 R_E in length (Wang et al., 2023). Indeed, Milan, Carter, and Hubert (2020) es-433 timated that closed field lines associated with transpolar arcs could extend as far as X <434 $-100 R_E$. More work needs to be conducted in the distant magnetotail and with mag-435 netospheric simulations to understand the magnetic topology of the closed magnetosphere. 436

437 4 Conclusions

We have investigated a fifteen-day period during which Cluster and Geotail sam-438 pled the magnetotail. Observations of the auroras by DMSP/SSUSI indicates that cusp-439 aligned arcs were present in the high latitude polar regions whenever the IMF clock an-440 gle was small, $|\theta| < 45^{\circ}$, which during the study interval amounted to approximately 441 30% of the time. Simultaneous observations of ions and electrons by Cluster and Geo-442 tail show significant plasma densities $(n_i \approx n_e \approx 1 \text{ cm}^{-3})$ in the tail during these in-443 tervals, even as far from the equatorial plane as $|Z| \approx 13 R_E$. This region of the mag-444 netotail would normally be devoid of plasma, being occupied by the open flux of the mag-445 netotail lobe. The presence of counter-streaming ions and double loss-cone electrons sug-446 gest, instead, that the plasma was trapped, i.e., that the magnetic field was closed. This 447 trapped plasma will provide a source for the CAA auroral emission, and as the auroral 448 emission covered the polar regions, we further suggest that the magnetosphere was al-449 most entirely closed. Coxon et al. (2021) recently showed that hot plasma consistent with 450 trapping on closed field lines is frequently seen at $|Z| > 5 R_E$ during northward IMF; 451 here we have shown that CAAs are the auroral signature of this closed flux, and that at 452 such times the magnetosphere is likely (almost) entirely closed. We believe that this clo-453 sure is achieved by dual-lobe reconnection, as proposed by Milan, Carter, Bower, et al. 454 (2020) and Milan et al. (2022). Our observations indicate that closure of the magneto-455 sphere is a common occurrence. 456

457 **5 Open Research**

⁴⁵⁸ Data from the Cluster Ion Spectrometry (CIS) instrument, the Plasma Electron
 ⁴⁵⁹ And Current Experiment (PEACE), and the Fluxgate Magnetometer (FGM) were ac ⁴⁶⁰ cessed through the Cluster Science Archive (CSA, formerly Cluster Active Archive (Laakso

et al., 2010)) at https://www.cosmos.esa.int/web/csa, maintained by ESA/ESTEC. 461 Geotail Magnetic Field Experiment (MGF) and Low Energy Particle (LEP) data were 462 accessed through the DARTS Solar-Terrestrial Physics data portal (https://www.darts 463 .isas.jaxa.jp/stp/geotail/data.html) maintained by ISAS/JAXA. The Defense Meteorological Satellite Program (DMSP) Special Sensor Ultraviolet Spectrographic Instru-465 ment (SSUSI) file type EDR-AUR data were obtained from JHU/APL (http://ssusi 466 . jhuapl.edu, data version 0106, software version 7.0.0, calibration period version E0018). 467 Advanced Magnetosphere and Planetary Electrodynamics Response Experiment (AM-468 PERE) data were obtained from JHU/APL (http://ampere.jhuapl.edu/dataget/index 469 .html) and processed using software provided (http://ampere.jhuapl.edu/). The high 470 resolution (1-min) OMNI data used in this study were obtained from the NASA God-471

dard Space Flight Center Space Physics Data Facility OMNIWeb portal (https://omniweb

.gsfc.nasa.gov/form/om_filt_min.html).

474 Acknowledgments

SEM and MKM were supported by the Science and Technology Facilities Council (STFC), 475 UK, grant no. ST/S000429/1. SEM and GEB were supported by the Natural Environ-476 ment Research Council (NERC), UK, grant no. NE/W006766/1. We are grateful to the 477 Cluster CIS, PEACE, and FGM instrument teams for providing data. We acknowledge 478 use of NASA/GSFC's Space Physics Data Facility's CDAWeb service (at http://cdaweb 479 .gsfc.nasa.gov), and OMNI data, and the Cluster Science Archive (at https://www 480 .cosmos.esa.int/web/csa), maintained by ESA/ESTEC. Geotail magnetic field and 481 plasma data were provided by T. Nagai and Y. Saito through DARTS at the Institute 482 of Space and Astronomical Science, JAXA, in Japan. 483

484 References

- Anderson, B., Takahashi, K., & Toth, B. (2000). Sensing global Birkeland currents
 with Iridium® engineering magnetometer data. *Geophysical Research Letters*, 27(24), 4045–4048. doi: https://doi.org/10.1029/2000GL000094
- Balogh, A., Dunlop, M., Cowley, S., Southwood, D., Thomlinson, J., Glassmeier, K.,
 ... others (1997). The Cluster magnetic field investigation. Space Science Reviews, 79, 65–91. doi: https://doi.org/10.1023/A:1004970907748
- Bower, G., Milan, S., Paxton, L., & Anderson, B. (2022). Occurrence statis tics of horse collar aurora. Journal of Geophysical Research: Space Physics,
 e2022JA030385. doi: https://doi.org/10.1029/2022JA030385
- Bower, G., Milan, S., Paxton, L., & Imber, S. (2022). Transpolar arcs: Seasonal de pendence identified by an automated detection algorithm. Journal of Geophys *ical Research: Space Physics*, 127(1), e2021JA029743. doi: https://doi.org/10
 .1029/2021JA029743
- Carlson, H., & Cowley, S. (2005). Accelerated polar rain electrons as the source of
 sun-aligned arcs in the polar cap during northward interplanetary magnetic
 field conditions. Journal of Geophysical Research: Space Physics, 110(A5).
- Carter, J., Milan, S., Fogg, A., Paxton, L., & Anderson, B. (2018). The association of high-latitude dayside aurora with NBZ field-aligned currents. *Journal* of Geophysical Research: Space Physics, 123(5), 3637–3645. doi: https://doi .org/10.1029/2017JA025082
- Carter, J., Milan, S., Fogg, A., Sangha, H., Lester, M., Paxton, L., & Anderson,
 B. (2020). The evolution of long-duration cusp spot emission during lobe
 reconnection with respect to field-aligned currents. *Journal of Geophysical Research: Space Physics*, 125, e2020JA027922. doi: https://doi.org/10.1029/
 2020JA027922

⁵¹⁰ Chisham, G., Freeman, M., Abel, G., Lam, M., Pinnock, M., Coleman, I., ... Vil ⁵¹¹ lain, J.-P. (2008). Remote sensing of the spatial and temporal structure of

512	magnetopause and magnetotail reconnection from the ionosphere. Reviews of Geophysics $46(1)$ doi: https://doi.org/10.1029/2007BG000223
515	Chicham C. Freeman M. Coleman I. Pinnock M. Hairston M. Lester M. &
514	Sofko G (2004) Measuring the dayside reconnection rate during an interval
515	of due northward interplanetary magnetic field Annales Geonhusicae 22(12)
510	4243-4258 doi: https://doi.org/10.5194/angeo-22-4243-2004
517	Chicham C. Lester M. Milan S. E. Freeman M. Bristow W. Crocott A
518	others (2007) A decade of the Super Dual Auroral Radar Network (Super-
519	DARN): Scientific achievements, new techniques and future directions. Surveys
520	in Geonhysics 28(1) 33-109 doi: https://doi.org/10.1007/s10712-007-9017-8
521	Cowley S (1981) Magnetospheric and ionospheric flow and the interplanetary mag-
522	netic field The Physical Basis of the Jonosphere in the Solar-Terrestrial Sus-
523	tem $ACARD_CP_{-}205 (A_{-}1)_{-}(A_{-}1A)$
524	Cowley S & Lackwood M (1002) Excitation and decay of solar wind driven flows
525	in the magnetosphere ionosphere system Annales Geophysicae 10, 103–115
526	Covon I Foar B Boidy I Fryer I & Dlank I (2021) Hot plasma in the
527	magnetetail lobes shows characteristics consistent with closed field lines
528	trapped in the lobes — <i>Lowral of Coophysical Passarch: Space Physica</i> , 196(0)
529	e2021 IA020516 doi: https://doi.org/10.1020/2021 IA020516
530	Covon I Milan S & Anderson B (2018) A review of Birkeland current research
531	using AMPERE Electric Currents in Coorners and Bound 257–278 doi:
532	$\frac{1}{1000}$ https://doi.org/10.1002/0781110324522.ch16
533	Cumpeck I & Blomberg I (2004) Transpolar are evolution and associated no
534	tential patterns Annales Geonhusicae $20(4)$ 1213–1231 doi: https://doi.org/
535	10 5194/angeo. 22.1213.2004
530	Cumpock I Sharber I Healis B Blomberg I. C. Cermany C. Spann I. &
537	Coley W (2002) Interplanetary magnetic field control of theta aurora devel-
530	opment Journal of Geophysical Research: Space Physics 107(A7) SIA-4 doi:
540	https://doi.org/10.1029/2001.IA009126
5/1	Dungey J (1961) Interplanetary magnetic field and the auroral zones <i>Physical Re</i> -
542	$p_{\text{angle}}(1) = p_{\text{angle}}(1) + p_{angle$
542	Dungev J (1963) The structure of the exosphere or adventures in velocity space
544	Geophysics. The Earth's Environment.
545	Escoubet, C., Fehringer, M., & Goldstein, M. (2001). Introduction the cluster mis-
546	sion. In Annales geophysicae (Vol. 19, pp. 1197–1200), doi: https://doi.org/10
547	.5194/angeo-19-1197-2001
548	Fear, R. (2021). The northward IMF magnetosphere. <i>Magnetospheres in the Solar</i>
549	Sustem, 293–309. doi: https://doi.org/10.1002/9781119815624.ch19
550	Fear, R., & Milan, S. (2012a). The IMF dependence of the local time of transpolar
551	arcs: Implications for formation mechanism. Journal of Geophysical Research:
552	Space Physics, 117(A3).
553	Fear, R., & Milan, S. (2012b). Ionospheric flows relating to transpolar arc formation.
554	Journal of Geophysical Research: Space Physics, 117(A9).
555	Fear, R., Milan, S., Maggiolo, R., Fazakerley, A., Dandouras, L. & Mende, S. (2014).
556	Direct observation of closed magnetic flux trapped in the high-latitude mag-
557	netosphere. Science, 346 (6216), 1506–1510. doi: https://doi.org/10.1126/
558	science.1257377
559	Frank, L., Craven, J., Burch, J., & Winningham, J. (1982). Polar views of the
560	Earth's aurora with Dynamics Explorer. Geophysical Research Letters. 9(9).
561	1001–1004. doi: https://doi.org/10.1029/GL009i009p01001
562	Frey, H. (2007). Localized aurora beyond the auroral oval. <i>Reviews of Geonhusics</i> .
563	45(1). doi: https://doi.org/10.1029/2005RG000174
564	Frey, H., Mende, S., Immel, T., Fuselier, S., Claffin, E., Gérard, JC., & Hubert. B.
565	(2002). Proton aurora in the cusp. Journal of Geophysical Research: Space
566	Physics, 107(A7), SMP-2. doi: https://doi.org/10.1029/2001JA900161

567 568	Fryer, L., Fear, R., Coxon, J., & Gingell, I. (2021). Observations of closed magnetic flux embedded in the lobes during periods of northward imf. <i>Journal of Comparison Process Physics</i> , 196(6), e20211A020281, doi: https://doi.
569 570	<i>Geophysical Research: Space Physics</i> , 126(6), e2021JA029281. doi: https://doi .org/10.1029/2021JA029281
571	Fuselier, S., Trattner, K., Petrinec, S., & Lavraud, B. (2012). Davside magnetic
572	topology at the Earth's magnetopause for northward IMF. Journal of Geo-
573	physical Research: Space Physics, 117(A8). doi: https://doi.org/10.1029/
574	2012JA017852
575	Hones Jr, E., Craven, J., Frank, L., Evans, D., & Newell, P. (1989). The horse-collar
576	aurora: A frequent pattern of the aurora in quiet times. <i>Geophysical Research</i>
577	Letters, 16(1), 37–40. doi: https://doi.org/10.1029/GL016i001p00037
578	Hosokawa, K., Kullen, A., Milan, S., Reidy, J., Zou, Y., Frey, H. U., Fear, R.
579	(2020). Aurora in the polar cap: A review. Space Science Reviews, 216(1), 15.
580	doi: https://doi.org/10.1007/s11214-020-0637-3
581	Hosokawa, K., Moen, J., Shiokawa, K., & Otsuka, Y. (2011). Motion of polar cap
582	arcs. Journal of Geophysical Research: Space Physics, 116(A1). doi: https://
583	doi.org/10.1029/2010JA015906
584	Huang, CS., Sofko, G., Koustov, A., Andre, D., Ruohoniemi, J., Greenwald,
585	R., & Hairston, M. (2000). Evolution of ionospheric multicell convection
586	during northward interplanetary magnetic field with $ B_z/B_u > 1$. Jour-
587	nal of Geophysical Research: Space Physics, 105(A12), 27095–27107. doi:
588	https://doi.org/10.1029/2000JA000163
589	Iijima, T., & Potemra, T. (1976). The amplitude distribution of field-aligned cur-
590	rents at northern high latitudes observed by Triad. Journal of Geophysical Re-
591	search, 81(13), 2165–2174. doi: https://doi.org/10.1029/JA081i013p02165
592	Iijima, T., Potemra, T., Zanetti, L., & Bythrow, P. (1984). Large-scale Birkeland
593	currents in the dayside polar region during strongly northward IMF: A new
594	Birkeland current system. Journal of Geophysical Research: Space Physics,
595	89(A9), 7441–7452. doi: https://doi.org/10.1029/JA089iA09p07441
596	Imber, S., Milan, S., & Hubert, B. (2006). The auroral and ionospheric flow signa-
597	tures of dual lobe reconnection. Annales Geophyiscae, $24(11)$, $3115-3129$. doi:
598	https://doi.org/10.5194/angeo-24-3115-2006
599	Ismail, S., Wallis, D., & Cogger, L. (1977). Characteristics of polar cap Sun-aligned
600	arcs. Journal of Geophysical Research, 82(29), 4741–4749. doi: https://doi
601	.org/10.1029/JA082i029p04741
602	Johnstone, A., Alsop, C., Burge, S., Carter, P., Coates, A., Coker, A., others
603	(1997). Peace: A plasma electron and current experiment. Space Science
604	Reviews, 351-398. doi: https://doi.org/10.1023/A:1004938001388
605	King, J., & Papitashvili, N. (2005). Solar wind spatial scales in and comparisons
606	of hourly Wind and ACE plasma and magnetic field data. Journal of Geo-
607	physical Research: Space Physics, 110(A2). doi: https://doi.org/10.1029/
608	2004JA010649
609	Kokubun, S., Yamamoto, T., Acuña, M., Hayashi, K., Shiokawa, K., & Kawano, H.
610	(1994). The GEOTAIL magnetic field experiment. Journal of geomagnetism
611	and geoelectricity, 46(1), 7–21. doi: https://doi.org/10.5636/jgg.46.7
612	Kullen, A., Brittnacher, M., Cumnock, J., & Blomberg, L. (2002). Solar wind
613	dependence of the occurrence and motion of polar auroral arcs: A statistical
614	study. Journal of Geophysical Research: Space Physics, 107(A11), 13–1. doi:
615	nttps://doi.org/10.1029/2002JA009245
616	Laakso, H., Perry, C., McCaffrey, S., Herment, D., Allen, A., Harvey, C., Turner,
617	R. (2010). Cluster active archive: Overview. Springer.
618	Lockwood, M., & Cowley, S. (1992). Ionospheric convection and the substorm cycle.
619	Proceedings of the International Conference on Substorms (ICS-1), 99–109.
620	Mailyan, B., Shi, Q., Kullen, A., Maggiolo, R., Zhang, Y., Fear, R., others
621	(2015). Transpolar arc observation after solar wind entry into the high-latitude

622	magnetosphere. Journal of Geophysical Research: Space Physics, 120(5),
623	3525-3534.
624	Milan, S., Bower, G., Carter, J., Paxton, L., Anderson, B., & Hairston, M. (2022).
625	Lobe reconnection and cusp-aligned auroral arcs. Journal of Geophysical Re-
626	search: Space Physics, 127(6), e2021JA030089. doi: https://doi.org/10.1029/
627	2021JA030089
628	Milan, S., Carter, J., Bower, G., Imber, S., Paxton, L., Anderson, B., Hu-
629	bert, B. (2020). Dual-lobe reconnection and horse-collar auroras. Jour-
630	nal of Geophysical Research: Space Physics, $125(10)$, $e2020JA028567$. doi:
631	https://doi.org/10.1029/2020JA028567
632	Milan, S., Carter, J., & Hubert, B. (2020). Probing the magnetic structure of a pair
633	of transpolar arcs with a solar wind pressure step. Journal of Geophysical Re- $l_{1} = C$ $R_{1} = 105(0)$ $R_{2} = 00101 A_{1}00710C$
634	search: Space Physics, $125(2)$, $e2019JA027196$.
635	Milan, S., Carter, J., Sangha, H., Bower, G., & Anderson, B. (2021). Magne-
636	to spheric flux throughput in the dungey cycle: Identification of convection $f(2)$
637	state during 2010. Journal of Geophysical Research: Space Physics, 120(2),
638	$e_{2020JA028437}$ doi: https://doi.org/10.1029/2020JA028437
639	Milan, S., Cowley, S., Lester, M., Wright, D., Slavin, J., Fillingim, M., Singer, H.
640	(2004). Response of the magnetotal to changes in the open flux content of the magnetotal $P_{\text{magnetotal}}$ $P_{\text{magnetotal}$ $P_{\text{magnetotal}}$ $P_{\text{magnetotal}$ $P_{\text{magnetotal}}$ $P_{\text{magnetotal}$ $P_{\text{magnetotal}}$ $P_{\text{magnetotal}$ $P_{\text{magnetotal}}$ $P_{\text{magnetotal}$
641	magnetosphere. Journal of Geophysical Research: Space Physics, 109(A4). doi: https://doi.org/10.1020/20021A010250
642	Milen S. Hubert D. & Creastt A. (2005). Formation and mation of a transmo
643	Milan, S., Hubert, B., & Grocott, A. (2005). Formation and motion of a transpo-
644	har arc in response to dayside and ingliside reconnection. $Journal of Geo-$
645	physical Research. Space Physics, 110(A1). doi: https://doi.org/10.1029/20041A010825
646	Milen S. Loster, M. Cowley, S. & Brittnacher, M. (2000) Develde con
647	westion and auroral morphology during an interval of northward inter
648	planetary magnetic field $Annales Ceonhusicae 18(A) A36-AAA doi:$
650	https://doi org/10/1007/s00585-000-0436-9
651	Milan S. Lester M. Cowley S. Oksavik K. Brittnacher M. Greenwald B.
652	Villain J-P (2003) Variations in the polar cap area during two substorm
653	cycles Annales Geonhusicae 21(5) 1121–1140
654	Milan, S., Provan, G., & Hubert, B. (2007). Magnetic flux transport in the Dungev
655	cycle: A survey of dayside and nightside reconnection rates. Journal of Geo-
656	physical Research: Space Physics, 112(A1). doi: https://doi.org/10.1029/
657	2006JA011642
658	Mukai, T., Machida, S., Saito, Y., Hirahara, M., Terasawa, T., Kaya, N.,
659	Nishida, A. (1994). The low energy particle (LEP) experiment onboard the
660	Geotail satellite. Journal of geomagnetism and geoelectricity, 46(8), 669–692.
661	doi: https://doi.org/10.5636/jgg.46.669
662	Nishida, A. (1994). The geotail mission (Vol. 21) (No. 25). Wiley Online Library.
663	Øieroset, M., Raeder, J., Phan, T., Wing, S., McFadden, J., Li, W., Balogh, A.
664	(2005). Global cooling and densification of the plasma sheet during an ex-
665	tended period of purely northward IMF on October 22–24, 2003. Geophysical
666	Research Letters, $32(12)$.
667	Paxton, L., Meng, CI., Fountain, G., Ogorzalek, B., Darlington, E., Gary, S.,
668	others (1992). Special sensor ultraviolet spectrographic imager: An instru-
669	ment description. In Instrumentation for planetary and terrestrial atmospheric
670	remote sensing (Vol. 1745, pp. 2–15). doi: https://doi.org/10.1117/12.60595
671	Paxton, L., Schaefer, R., Zhang, Y., & Kil, H. (2017). Far ultraviolet instrument
672	technology. Journal of Geophysical Research: Space Physics, 122(2), 2706–
673	2733. doi: https://doi.org/10.1002/2016JA023578
674	Paxton, L., & Zhang, Y. (2016). Far ultraviolet imaging of the aurora. In Space
675	weather fundamentals (pp. 213–244). CRC Press.
676	Paxton, L., Zhang, Y., Kil, H., & Schaefer, R. (2021). Exploring the upper atmo-

677	sphere: Using optical remote sensing. Upper Atmosphere Dynamics and Ener-
678	getics, 487–522. doi: https://doi.org/10.1002/9781119815631.ch23
679	Reidy, J., Fear, R., Whiter, D., Lanchester, B., Kavanagh, A., Price, D. J., Pax-
680	ton, L. (2020). Multiscale observation of two polar cap arcs occurring on
681	different magnetic field topologies. Journal of Geophysical Research: Space
682	Physics, 125(8), e2019JA027611.
683	Rème, H., Bosqued, J., Sauvaud, J., Cros, A., Dandouras, J., Aoustin, C., others
684	(1997). The cluster ion spectrometry (cis) experiment. The cluster and phoenix
685	missions, $303-350$. doi: https://doi.org/10.1007/978-94-011-5666-0_12
686	Rich, F., & Hairston, M. (1994). Large-scale convection patterns observed by
687	DMSP. Journal of Geophysical Research: Space Physics, 99(A3), 3827–3844.
688	doi: https://doi.org/10.1029/93JA03296
689	Shi, Q., Zong, QG., Fu, S., Dunlop, M., Pu, Z., Parks, G., others (2013). Solar
690	wind entry into the high-latitude terrestrial magnetosphere during geomagneti-
691	cally quiet times. Nature communications, 4, 1466.
692	Siscoe, G., Farrugia, C., & Sandholt, P. (2011). Comparison between the two basic
693	modes of magnetospheric convection. Journal of Geophysical Research: Space
694	<i>Physics</i> , 116(A5). doi: https://doi.org/10.1029/2010JA015842
695	Siscoe, G., & Huang, T. (1985). Polar cap inflation and deflation. <i>Journal of Geo</i> -
696	physical Research: Space Physics, 90(A1), 543–547.
697	Song, P., DeZeeuw, D., Gombosi, T., Groth, C., & Powell, K. (1999). A numerical
698	study of solar wind-magnetosphere interaction for northward interplanetary
699	magnetic field. Journal of Geophysical Research: Space Physics, 104 (A12),
700	28301-28378. doi: https://doi.org/10.1029/1999JA900378
701	(2008) The plagma short and boundary layors under northward IME: A multi-
702	(2008). The plasma sheet and boundary layers under northward hvir. A multi- point and multi instrument perspective $Advances in Space Becauch (1(10))$
703	1610–1620
704	1019^{-1029} . Tsympton N (1005) Modeling the earth's magnetospheric magnetic field con
705	fined within a realistic magnetonause <u>Journal of Geonhusical Research</u> : Snace
700	Physics $100(A4)$ 5599–5612 doi: https://doi.org/10.1029/941A03193
707	Wang X - Y Zhang O - H Wang C Zhang Y - L Tang B - B Xing Z - Y
709	others (2023). Unusual shrinkage and reshaping of earth's magnetosphere un-
710	der a strong northward interplanetary magnetic field. <i>Communications Earth</i>
711	<i>Estimate a morphalical formation of the second sec</i>
712	Waters, C., Anderson, B., & Liou, K. (2001). Estimation of global field aligned
713	currents using the Iridium $(\widehat{\mathbf{R}})$ system magnetometer data. Geophysical Research
714	Letters, 28(11), 2165–2168. doi: https://doi.org/10.1029/2000GL012725
715	Zhang, QH., Zhang, YL., Wang, C., Lockwood, M., Yang, HG., Tang, BB.,
716	others (2020). Multiple transpolar auroral arcs reveal insight about
717	coupling processes in the Earth's magnetotail. Proceedings of the National
718	Academy of Sciences, 117(28), 16193–16198. doi: https://doi.org/10.1073/
719	pnas.2000614117
720	Zhang, Y., Paxton, L., Zhang, QH., & Xing, Z. (2016). Polar cap arcs: Sun-aligned
721	or cusp-aligned? Journal of Atmospheric and Solar-Terrestrial Physics, 146,
722	123–128. doi: https://doi.org/10.1016/j.jastp.2016.06.001

The association of cusp-aligned arcs with plasma in the magnetotail implies a closed magnetosphere

S. E. Milan^{1*}, M. K. Mooney¹, G. E. Bower¹, M. G. G. T. Taylor², L. J. Paxton³, I. Dandouras⁴, A. N. Fazakerley⁵, C. M. Carr⁶, B. J. Anderson⁷, and S. K. Vines⁷

6	¹ School of Physics and Astronomy, University of Leicester, Leicester, UK.
7	² ESA/ESTEC, Noordwijk, The Netherlands.
8	³ William B. Hanson Center for Space Sciences, University of Texas at Dallas, USA.
9	⁴ Institut de Recherche en Astrophysique et Planétologie (IRAP), CNRS and Université Paul Sabatier,
10	Toulouse, France.
11	⁵ Mullard Space Science Laboratory, University College, London, UK.
12	⁶ Imperial College, London, UK.
13	⁷ Johns Hopkins University, Applied Physics Laboratory, USA.

Key Points:

1

2

3

4

14

15	•	Cusp-aligned arcs observed by the DMSP spacecraft occur frequently for north-
16		ward IMF
17	•	Cluster and Geotail observations show that the arcs are accompanied by trapped
18		plasma at high latitudes in the magnetotail

We interpret cusp-aligned arcs as a signature of a magnetosphere almost entirely
 closed by dual-lobe reconnection

 $^{^{*}\}mbox{School}$ of Physics and Astronomy, University of Leicester, Leicester LE1 7RH, UK

Corresponding author: Steve Milan, steve.milan@le.ac.uk

21 Abstract

We investigate a fifteen-day period in October 2011. Auroral observations by the SSUSI 22 instrument onboard the DMSP F16, F17, and F18 spacecraft indicate that the polar re-23 gions were covered by weak cusp-aligned arc emissions whenever the IMF clock angle was 24 small, $|\theta| < 45^{\circ}$, which amounted to 30% of the time. Simultaneous observations of ions 25 and electrons in the tail by the Cluster C4 and Geotail spacecraft showed that during 26 these intervals dense ($\approx 1 \text{ cm}^{-3}$) plasma was observed, even as far from the equatorial 27 plane of the tail as $|Z_{GSE}| \approx 13 R_E$. The ions had a pitch angle distribution peaking 28 parallel and antiparallel to the magnetic field and the electrons had pitch angles that peaked 29 perpendicular to the field. We interpret the counter-streaming ions and double loss-cone 30 electrons as evidence that the plasma was trapped on closed field lines, and acted as a 31 source for the cusp-aligned arc emission across the polar regions. This suggests that the 32 magnetosphere was almost entirely closed during these periods. We further argue that 33 the closure occured as a consequence of dual-lobe reconnection at the dayside magne-34 topause. Our finding forces a significant re-evaluation of the magnetic topology of the 35 magnetosphere during periods of northwards IMF. 36

³⁷ Plain Language Summary

The magnetosphere is usually assumed to contain both open and closed magnetic 38 flux. Closed magnetic field lines have both ends connected to the Earth; open field lines 39 connect to the Earth at one end and into the interplanetary medium at the other. There 40 tends to be little plasma on open field lines as the particles escape down the magneto-41 tail, whereas plasma on closed field lines is trapped. Open flux near the poles naturally 42 explains the oval configuration of Earth's auroras, with a lack of auroras at very high 43 latitudes where there is no plasma to cause emissions. Somewhat unexpectedly, we show 44 that auroral emission near the poles is common and that at these times there is signif-45 icant plasma in the magnetotail, indicating that the magnetosphere contains only closed 46 flux. We propose that this magnetic configuration is formed by a process known as dual-47 lobe magnetic reconnection which occurs when the interplanetary magnetic field within 48 the solar wind points northwards. We must re-evaluate the standard picture of magne-49 tospheric structure during these periods of northwards interplanetary magnetic field. 50

51 **1** Introduction

In this study we investigate repeated occurrences of cusp-aligned arcs (CAAs), the poorly understood situation in which weak auroral emissions fill the polar regions, during a fifteen-day interval in 2011. We conclude that CAAs are a common occurrence during periods of northward interplanetary magnetic field (IMF), and that they are a signature of a nearly- or entirely-closed magnetosphere. This forces a significant re-evaluation of the magnetic topology of the magnetosphere during northward IMF.

The coupling between the solar wind and the magnetosphere is relatively well un-58 derstood for southward-directed IMF: as first proposed by Dungey (1961), magnetic re-59 connection near the subsolar magnetopause opens previously-closed magnetic flux which 60 then forms the magnetotail lobes; thereafter, magnetic reconnection in the central plane 61 of the tail recloses flux which returns to the dayside, resulting in the Dungey cycle of mag-62 netospheric convection. The northern and southern ionospheric polar caps, the dim re-63 gions encircled by the auroral ovals, are the ionospheric projection of the open flux form-64 ing the lobes, and their size can be used to quantify the open flux content of the mag-65 netosphere, F_{PC} (e.g., Milan et al., 2003). This naturally explains many aspects of mag-66 netospheric structure and dynamics, including the formation of the magnetotail, the twin-67 cell ionospheric convection pattern, the morphology of the auroral oval, the plasma pop-68 ulations within the magnetosphere, and the evacuated magnetotail lobes. When time-69 varying magnetic reconnection rates are considered, the expanding/contracting nature 70

of the polar cap and the substorm cycle can be understood (Siscoe & Huang, 1985; Cow-71 ley & Lockwood, 1992; Lockwood & Cowley, 1992). Typically, F_{PC} varies between 0.3 72 and 0.9 GWb (Milan et al., 2003, 2007, 2021). However, it can become as low as 0.2 GWb 73 $(\sim 2.5\%)$ of the 8 GWb associated with the terrestrial dipole) during particularly extreme 74 nightside reconnection events, accompanied by a near-total in-filling of the polar regions 75 by bright auroral emission (e.g., Milan et al., 2004). In this paper we show that in fact 76 the polar cap closes entirely, frequently, and with much less fanfare, during periods of 77 northwards IMF. 78

79 The coupling during northward-directed IMF is still poorly understood. Although Dungey (1963) correctly proposed that magnetic reconnection would take place at the 80 high latitude magnetopause, tailwards of the cusps, the ramifications are still not fully 81 resolved. The list of northward-IMF (IMF $B_Z > 0$ or NBZ) phenomena includes: single-82 and dual-lobe reconnection (e.g., Cowley, 1981; Fuselier et al., 2012), reverse convection 83 in the polar cap (e.g., Huang et al., 2000), NBZ field-aligned currents poleward of the 84 noon auroral oval (e.g. Iijima et al., 1984), cusp auroral spots (e.g., Milan et al., 2000; 85 Frey et al., 2002; Frey, 2007; Carter et al., 2018, 2020), transpolar or polar cap arcs (e.g., 86 Frank et al., 1982; Cumnock et al., 2002; Kullen et al., 2002; Milan et al., 2005; Fear et 87 al., 2014), horse-collar auroras (e.g., Hones Jr et al., 1989; Milan, Carter, Bower, et al., 88 2020; Bower, Milan, Paxton, & Anderson, 2022), and cusp-aligned arcs (e.g., Y. Zhang 89 et al., 2016; Q.-H. Zhang et al., 2020; Milan et al., 2022; Wang et al., 2023). The inter-90 ested reader is directed to the recent reviews of NBZ phenomena by Hosokawa et al. (2020) 91 and Fear (2021). 92

A key question regarding periods of NBZ is the degree to which the magnetosphere 93 loses open magnetic flux, and the resulting distribution of open and closed flux in the 94 polar regions. It has been suggested that polar cap arcs may be associated with both 95 open and closed magnetic flux (e.g., Carlson & Cowley, 2005; Reidy et al., 2020; Bower, 96 Milan, Paxton, & Imber, 2022). Evidence suggests that the most prominent of polar cap 97 arcs, those also known as theta auroras, are likely associated with closed magnetic flux 98 (Fear et al., 2014; Fryer et al., 2021; Coxon et al., 2021), proposed to be produced by 99 magnetic reconnection in the magnetotail (Milan et al., 2005; Fear & Milan, 2012a, 2012b). 100 The situation regarding less prominent sun-aligned or cusp-aligned arcs, in which mul-101 tiple weak arcs fill the polar regions (Y. Zhang et al., 2016), is as yet unresolved. If these 102 form on the open flux of the largely-evacuated magnetotail lobes, then the source of the 103 auroral precipitation is called into question, though Carlson and Cowley (2005) proposed 104 that polar rain could provide sufficient plasma to be accelerated in flow shears in the iono-105 spheric convection pattern to produce such arcs. 106

During prolonged NBZ, the polar cap can contract and become teardrop-shaped, 107 leading to the horse-collar auroras (HCAs) configuration (Hones Jr et al., 1989). Milan, 108 Carter, Bower, et al. (2020) and Bower, Milan, Paxton, and Anderson (2022) have sug-109 gested that this is formed by the closure of magnetic flux by dual-lobe reconnection (DLR) 110 for near-zero IMF clock angle, supported by numerical simulations (Wang et al., 2023). 111 Dual-lobe reconnection should be an efficient mechanism by which the magnetosphere 112 can capture solar wind plasma (e.g., Imber et al., 2006), and it has been proposed that 113 over time this can lead to the formation of a cold, dense plasma sheet (CDPS) (Øieroset 114 et al., 2005). However, this is not the only means by which solar wind plasma is thought 115 to enter the magnetosphere during periods of NBZ, with inward diffusion at the mag-116 netotail flanks (see discussion in Taylor et al., 2008) or direct entry through (single) lobe 117 reconnection (Shi et al., 2013; Mailyan et al., 2015) being other commonly discussed mech-118 anisms. It is unclear, however, how efficient diffusion can be, how quickly captured so-119 lar wind plasma can be redistributed throughout the magnetosphere, and why the plasma 120 remains near the Earth and does not escape back to the solar wind down the open field 121 lines of the lobes, as it does for southwards IMF. 122



Figure 1. (Left) The orbit of Cluster C4 (red) and Geotail (green) for the period of study, in the GSE X - Z plane. The T96 magnetic field configuration on DOY 280, 2011, at 13 UT, for $P_{dyn} = 2$ nPa, $D_{st} = -10$ nT, $B_Y = 0$ and $B_Z = 5$ nT is superimposed in grey. (Middle) The orbit of Cluster and Geotail for the period of study, in the GSE Y - Z plane; the location of the magnetopause at X = -10 R_E is shown for reference. (Right) Field-line tracing from the location of Cluster and Geotail to the northern (orange and dark green) and southern (red and light green) hemisphere ionospheres for the period of interest, presented on a geomagnetic latitude and MLT grid. An average auroral oval for $K_P = 1$ is overlaid for reference.

The cusp-aligned arcs (hereafter CAAs) auroral configuration is perhaps the most 123 poorly-studied NBZ auroral phenomenon. Partially, this is because the emissions tend 124 to be weak, so are better observed from the ground (e.g., Ismail et al., 1977; Hosokawa 125 et al., 2011), with limited geographical coverage, rather than from space. Recently, Q.-126 H. Zhang et al. (2020) and Wang et al. (2023) suggested that CAAs are produced by plasma 127 flow shears introduced into the magnetosphere by Kelvin-Helmholtz surface waves ex-128 cited on the magnetotail flanks by the flow of the solar wind. However, this does not re-129 solve the source of the precipitating plasma. On the other hand, Milan et al. (2022) pro-130 posed that if DLR continues for a prolonged period, then the magnetosphere can become 131 almost entirely closed and the horse-collar auroral configuration can develop to the point 132 where the polar slot (distorted polar cap) can almost disappear. In this scenario, the closed 133 magnetosphere will be filled with trapped solar wind plasma and, according to Milan et 134 al. (2022), flow shears produced by lobe reconnection can then accelerate this trapped 135 plasma into the atmosphere to form the CAAs. Whereas Carlson and Cowley (2005) sug-136 gested that the source of plasma to produce weak polar cap arcs was polar rain that had 137 fled to the distant magnetotail along open field lines, instead Milan et al. (2022) proposed 138 that the source is solar wind plasma captured by DLR and trapped closer to the Earth 139 on closed field lines. 140

In this study we examine plasma populations observed in the near-Earth tail by 141 the C4 spacecraft of the Cluster constellation (Escoubet et al., 2001) and the Geotail (Nishida, 142 1994) spacecraft during a prolonged period in October 2011 when auroral observations 143 from F16, F17, and F18 of the Defense Meteorological Spacecraft Program detected mul-144 tiple instances of CAAs. Dense plasma is repeatedly found in regions that would nor-145 mally be occupied by the evacuated open field lines of the northern and southern lobes. 146 and it is concluded that the magnetosphere is almost entirely closed and the plasma is 147 trapped, providing a source for the high latitude auroral emission. 148

¹⁴⁹ 2 Observations

The period under investigation is the 3 to 17 October 2011 (days-of-year 276 to 290), 150 which encompasses six orbits of the Cluster constellation during its tail season and three 151 orbits of Geotail; the orbits are shown in Figure 1. At this time, the orbit of C4 was in-152 clined such that it was southwards of the neutral sheet for most of the time, except near 153 perigee. During the study interval, C4 reached $Z_{GSE} \approx -13 R_E, X_{GSE} \approx -14 R_E$ 154 at apogee, and it was located relatively centrally in the tail, $0 < Y_{GSE} < 10 R_E$, in 155 the first half of each orbit, but closer to the dusk flank in the second. The orbit of Geo-156 157 tail was such that it sampled the magnetotail northwards of the neutral sheet, up to $Z_{GSE} \approx$ 8 R_E at $X_{GSE} \approx -19 R_E$. A representative magnetic field tracing in the T96 model 158 (Tsyganenko, 1995), with input parameters $P_{dyn} = 2 \text{ nT}$, $D_{st} = -10 \text{ nT}$, and IMF 159 $B_Y = 0$ and $B_Z = 5$ nT, is overlaid for reference. These parameters were chosen to 160 represent a quiet NBZ magnetosphere, and will be used throughout the rest of the study 161 to compare with magnetic field observations. As the plasma sheet is usually confined to 162 $|Z| < 5 R_E$, for much of their orbits Geotail and C4 are expected to be within the north-163 ern and southern lobes of the magnetotail, respectively. Field line tracings from the lo-164 cations of the two spacecraft to the northern and southern ionospheres are also presented, 165 with a $K_P = 1$ average auroral oval superimposed for reference. This indicates that near 166 apogee C4 would normally be expected to map to the central polar cap in the southern 167 hemisphere, and that Geotail would frequently map to the nightside polar cap in the north-168 ern hemisphere. 169

Auroral observations during this period are provided by the Special Sensor Ultra-170 violet Spectrographic Imager or SSUSI experiment (Paxton et al., 1992) onboard the DMSP-171 F16, -F17, and -F18. The DMSP spacecraft have sun-synchronous orbits near an alti-172 tude of 850 km. SSUSI measured auroral luminosity in a swath either side of the space-173 craft orbit in the Lyman-Birge-Hopfield short (LBHs) band, 140 to 150 nm (see Paxton 174 and Zhang (2016); Paxton et al. (2017, 2021) and the references cited therein for further 175 description of the instrument and data products). Measurements of the distribution of 176 magnetosphere-ionosphere field aligned currents (FACs) were provided by the Active Mag-177 netosphere and Planetary Electrodynamics Response Experiment (AMPERE) technique 178 (Anderson et al., 2000; Coxon et al., 2018; Waters et al., 2001). We also make use of iono-179 spheric flow observations provided by the Super Dual Auroral Radar Network (Super-180 DARN (Chisham et al., 2007)) and the Ion Driftmeter (IDM) component of the Special 181 Sensors–Ions, Electrons, and Scintillation thermal plasma analysis package or SSIES in-182 strument onboard the Defense Meteorological Satellite Program spacecraft (DMSP/IDM 183 (Rich & Hairston, 1994)). Solar wind parameters were taken from the OMNI data-set 184 (King & Papitashvili, 2005). 185

Figure 2 shows 12 snapshots of the auroral morphology from DOYs 278 to 280. Be-186 low this, a keogram of the observations along the dawn-dusk meridian made by DMSP-187 F16/SSUSI in the northern hemisphere is shown, along with the IMF clock angle from 188 OMNI. Around 10:20 UT on DOY 278 (panels a and b), the IMF had a southwards com-189 ponent and typical twin-cell ionospheric flows were observed by SuperDARN and DMSP/IDM 190 (not shown for brevity). The auroral morphology was also typical, showing an oval sur-191 rounding a dim polar cap, with evidence for substorm activity on the nightside. Between 192 approximately 16:00 UT, DOY 278, and 13:00 UT, DOY 279, the IMF turned northwards. 193 During this period the auroras dimmed, contracted to higher latitudes and acquired a 194 horse-collar auroral configuration, before the polar regions filled with auroral emission 195 mainly in the form of cusp-aligned arcs or CAAs (c to f). CAAs are seen in both north-196 ern and southern hemispheres, panels c and e being from the north and d and f being 197 from the south. Then the IMF turned southwards again, twin-cell convection resumed. 198 and the polar cap reopened (g and h). There followed another period of NBZ, during 199 which CAAs reformed (i and j), before again a southward turning and a reopening of 200 the polar cap (k to l). The lower panels clearly show the expansions and contractions 201



Figure 2. Snapshots of the LBHs auroral configuration in the northern and southern hemispheres observed by DMSP/SSUSI onboard DMSP-F16, -F17, and -F18 between 5 and 7 October 2011. Each panel is presented in a geomagnetic latitude and local time format, with noon towards the top and dawn towards the right. Grey circles indicate geomagnetic latitude in steps of 10°. Of the two lower panels, the top one shows a keogram of auroral emissions observed by DMSP-F16/SSUSI along the dawn-dusk meridian of the northern hemisphere; grey vertical bars indicate where data is missing. Red horizontal bars indicate the times that cusp-aligned arcs were observed. The bottom panel is clock angle, highlighted in red when $|\theta| < 45^{\circ}$.

of the auroral oval with changes in clock angle, and the presence of auroral emissions at high latitudes during periods when the clock angle is near zero, $|\theta| < 45^{\circ}$, highlighted in red.

Over the rest of the period considered several other intervals of CAAs were found, 205 as will be indicated in later figures. The start and end times of these intervals are ap-206 proximate due to the relatively coarse cadence of the DMSP orbits; there are also pe-207 riods when it was not possible to positively identify whether CAAs were present or not, 208 due to only partial coverage of the polar regions by the SSUSI field-of-view. During pe-209 riods when HCAs or CAAs were present, the ionospheric flows measured by SuperDARN 210 and DMSP/IDM and the field-aligned currents observed by AMPERE were consistent 211 with the observations reported by Milan, Carter, Bower, et al. (2020) and Milan et al. 212 (2022) – reverse lobe convection, NBZ FACs at noon – suggesting that dual-lobe recon-213 nection was responsible for closing the magnetosphere. The occurrence of CAAs when 214 $|\theta| < 45^{\circ}$ is also consistent with the statistical occurrence of horse-collar auroras (Bower, 215 Milan, Paxton, & Anderson, 2022), thought to be the auroral precursor to CAAs. 216

We use Cluster C4 observations from the Composition and Distribution Function 217 analyser of the Cluster Ion Spectrometry instrument (C4/CIS-CODIF (Rème et al., 1997)), 218 the Plasma Electron And Current Experiment (C4/PEACE (Johnstone et al., 1997)), 219 and the Fluxgate Magnetometer (C4/FGM (Balogh et al., 1997)). Figure 3 covers the 220 period 3 to 9 October, spanning the first three C4 orbits considered and encompassing 221 the interval shown in Figure 2. The panels are presented in the following order. (a) The 222 GSE X, Y, and Z (R_E) position of C4. (b) The B_X , B_Y , B_Z components of the mag-223 netic field measured by C4/FGM in GSE coordinates. Dots indicate predications of the 224 magnetic field measurements by the T96 (Tsyganenko, 1995) model, with fixed inputs 225 (as before). (c) The proton density, n_i , observed by C4/CIS-CODIF. Vertical red bars 226 indicate when CAAs were observed by DMSP/SSUSI. (d) The proton differential energy 227 flux spectrogram measured by C4/CIS-CODIF. (e) The electron differential energy flux 228 spectrogram measured by C4/PEACE. (f) A dawn-dusk keogram of auroral emissions 229 observed by DMSP-F16/SSUSI in the northern hemisphere. (g) A dawn-dusk keogram 230 of FAC density measured by AMPERE in the northern hemisphere, with red/blue in-231 dicating up/down currents. (h) The clock angle of the IMF, θ , highlighted in red when 232 $|\theta| < 45^{\circ}$. (i) The GSM B_X, B_Y, B_Z components of the IMF from OMNI. (j) The so-233 lar wind speed and density from OMNI. 234

The SSUSI keogram (f) reveals that at most times there is little auroral emission inside the auroral oval, consistent with an open polar cap, except at the times of CAAs (red bars in (c)) when $|\theta| < 45^{\circ}$. The region 1/2 FACs (Iijima & Potemra, 1976) observed by AMPERE (g) are enhanced when the polar cap is open, indicating Dungey cycle driving of the magnetosphere by subolar reconnection, especially during periods with IMF $B_Z < 0$ nT, $|\theta| > 90^{\circ}$. When CAAs are present the R1/R2 FACs are weak, though NBZ FACs tend to be observed at noon (not shown).

Near the perigee of each orbit (marked PG) when $|Z| < 5 R_E$, C4 passed through 242 the plasma sheet and ring current regions and enhanced proton densities, n_i , were ob-243 served (c). During the first orbit of Figure 3 almost no plasma was observed when Z <244 $-5 R_E$, consistent with the open lobe. In contrast, while C4 was near perigee on DOY 245 278 the IMF turned northwards and CAAs were observed, and throughout this period 246 C4 was engulfed in protons with $n_i \approx 1 \text{ cm}^{-3}$ up to near apogee (AG) on DOY 279. 247 As the IMF turned southwards around 13:00 UT on DOY 279, the density returned to 248 typical lobe values, $n_i < 0.1 \text{ cm}^{-3}$. Around 00 UT on DOY 280 CAAs were once again 249 observed, and n_i rose to near 1 cm⁻³, only to drop again at 04 UT as the IMF turned 250 southwards; the IMF remained predominantly southwards for the duration of the third 251 orbit, and n_i remained low throughout (except for a short period near apogee, see be-252 low). 253



Figure 3. Observations from 3 to 10 October 2011, which encompasses three orbits by Cluster. (a) The GSE position of C4. Apogees and perigees of the orbit are indicated by AG and PG. Orange highlighting indicates the time of the first Geotail orbit shown in Figure 6. (b) C4/FGM observations of the magnetic field. T96 model predictions for fixed input parameters are indicated as dots. (c) C4/CIS-CODIF ion density measurements. Red bars indicate periods of observation of cusp-aligned arcs by DMSP/SSUSI. (d) C4/CIS-CODIF ion spectrogram. (e) C4/PEACE electron spectrogram. (f) A dawn-dusk keogram of auroral observations by DMSP-F16/SSUSI in the northern hemisphere. (g) A dawn-dusk keogram of field-aligned currents measured by AMPERE, red and blue being upward and downward FACs, respectively. (h) The clock angle of the IMF from OMNI. The clock angle is highlighted in red when $|\theta| < 45^{\circ}$. (i) The components of the IMF from OMNI. (j) The solar wind speed and density from OMNI.

The C4/CIS-CODIF spectrogram (d) indicates that the CAA-related ions had energies between several 100s and several 1000s eV. (We note that at the apogees of the second and third orbits a brief interval of cold, < 100 eV, ions was also observed.) The C4/PEACE spectrogram (e) indicates that the ions were accompanied by electrons, with energies below about 1 keV. In the first CAA interval of DOY 279 the ions, and especially the electrons, reduced in energy with time, indicating a cooling of the plasma.

Figure 4 presents the next three orbits of C4. During this period, the IMF $|\theta| <$ 45° for much of the time, and there were several intervals of CAAs observed by DMSP/SSUSI. Accompanying these intervals, C4/CIS-CODIF saw elevated n_i , where otherwise lobe conditions might have been expected. During DOYs 287 to 290, there were repeated swings between $|\theta| < 45^{\circ}$ and $|\theta| > 45^{\circ}$, and CAAs and protons came and went in tandem.

Figure 5 focusses on the two intervals of CAAs presented in Figures 2 and 3. In 265 addition to the spectrograms, pitch angle distributions of the ions (d) and electrons (f)266 are shown. The proton pitch angle fluxes are integrated across the full energy range of 267 the CIS-CODIF instrument; the electron fluxes are calculated from the low energy elec-268 tron analyser head of the PEACE instrument, limited to energies above 100 eV to re-269 move the contribution of photoelectrons. During periods of CAAs, and when C4 was not 270 too close to the Earth, ion pitch angles were concentrated at 0° and 180° , indicating two 271 counter-streaming populations. The electron pitch angles, conversely, peaked at 90° , with 272 a distinct lack of electrons near 0° and 180° , indicating a double loss-cone distribution. 273 The double loss-cone electron distribution is similar to that observed by Fear et al. (2014)274 above a transpolar arc, and is indicative of plasma trapped on closed field lines. The counter-275 streaming ions support this conclusion. We note that the ion and electron densities dur-276 ing the periods of CAAs were quite variable, indicating that the trapped plasma was not 277 uniformly distributed but was present continuously. The solar wind density was variable 278 during this period, but it is not clear if variations in ion/electron density and energy ob-279 served by C4 are correlated with enhancements in n_{SW} . As noted previously, the ion and 280 electron energies decreased overall with time, which might indicate the progressive mix-281 ing of plasma sheet and solar wind populations, leading to the formation of a cold, dense 282 plasma sheet (e.g., Øieroset et al., 2005). 283

Just prior to the second period of CAAs, CIS-CODIF detected a beam of low energy ions observed only near 0° pitch angles (i.e. flowing tailward). A similar beam was observed around 00 UT near apogee on DOY 282 (see Figure 3). As there are no counterstreaming ions, we suggest that at these are the signature of plasma escaping to the solar wind along open field lines.

Superimposed on the C4/FGM observations (a) is a model prediction from T96 for fixed NBZ input parameters (as before). The model tends to match the observations well during the periods of CAAs, but the B_X component was enhanced by ≈ 10 nT during the periods when CAAs were not present, indicating that the tail was more inflated at these times. This also supports the conclusion that the magnetotail open flux was significantly reduced when CAAs were observed.

Figure 6 presents two time intervals when Geotail was located in the magnetotail; 295 Figures 3 and 4 show that these roughly correspond to orbits 3 and 5 of C4. Panels a296 and b show that in each case Geotail entered the dusk flank of the magnetotail at the 297 start of the interval, near $Z \approx 5 R_E$, and rose to $Z \approx 8 R_E$ at $X \approx -19 R_E$, $Y \approx$ 298 $0 R_E$. It later exited the dayside magnetopause in the pre-noon sector. Panels e to l present 299 ion and electron spectrograms from the Low Energy Particle (LEP) instrument (Mukai 300 et al., 1994), showing fluxes in the sunward and tailward directions. During the first or-301 bit, the plasma sheet was seen as the spacecraft entered and exited the magnetosphere, 302 but the evacuated lobe was encountered from 16 UT on DOY 281 to 14 UT on DOY 282. 303 During this period, IMF $|\theta| > 45^{\circ}$ at all times (m). During the second orbit, the plasma 304 sheet was also seen at the start and the end, with periods of lobe between 23 UT, DOY 305



Figure 4. Observations from 10 to 17 October, presented in the same format as Figure 3.



Figure 5. Observations from 5 to 7 October 2011. (a) C4/FGM observations of the magnetic field. T96 model predictions for fixed input parameters are indicated as dots. (b) C4/CIS-CODIF ion density measurements. Red bars indicate periods of observation of cusp-aligned arcs by DMSP/SSUSI. (c, d) C4/CIS-CODIF ion spectrogram and pitch angle distribution. (e, f) C4/PEACE spectrogram and pitch angle distribution. (g) A dawn-dusk keogram of auroral observations by DMSP-F16/SSUSI. (h) The clock angle of the IMF from OMNI. The clock angle is highlighted in red when $|\theta| < 45^{\circ}$. (i) The solar wind speed and density from OMNI.



Figure 6. Two time periods during which Geotail was located within the magnetotail. (a, b) The GSE location of Geotail. (c, d) The magnetic field components measured by MGF, with a T96 prediction superimposed. Red bars indicate when cusp-aligned arcs were observed by DMSP/SSUSI. (e, f) Ion spectrograms of fluxes in the sunwards direction. (g, h) Ion spectrograms of fluxes in the tailwards direction. (i, j) Electron spectrograms of fluxes in the sunwards direction. (m, n) IMF clock angle, highlighted in red when $|\theta| < 45^{\circ}$.

286, and 18 UT, DOY 287. However, three intervals of plasma, with fluxes in both the 306 sunwards and tailwards directions, were detected at the same time as CAAs were ob-307 served by DMSP/SSUSI, associated with excursions to $|\theta| < 45^{\circ}$ (n). Plasma was also 308 seen by C4 in the southern tail at these times (Figure 4), and the plasma characteris-309 tics were similar. The presence of fluxes in both the sunwards and tailwards directions. 310 indicate that the plasma is trapped on closed field lines. During the second orbit, the 311 magnetic field observed by the Magnetic Field Experiment (MGF) instrument (Kokubun 312 et al., 1994), panel d, matched closely the prediction by T96. However, during the first 313 orbit, around the time Geotail was in the lobe the B_X component was elevated above 314 the prediction (c) indicating an inflated magnetotail. 315

316 **3 Discussion**

The occurrence of cusp-aligned arcs filling the polar regions is frequent. We find 317 that CAAs have a high probability of appearing if $|\theta| < 45^{\circ}$ for an appreciable length 318 of time (one to two hours or more). During our period of study, CAAs were observed 319 for approximately 30% of the time. Although polar cap arcs received attention in the 320 past (e.g., Ismail et al., 1977), CAAs are too dim to have been detected by previous gen-321 erations of global auroral imagers (e.g., IMAGE FUV), so their importance has been over-322 looked. We find that CAAs are accompanied by dense plasma of energies from several 323 eV to several 10s keV in regions of the magnetosphere (up to $|Z| \approx 13 R_E$ in our ob-324 servations) that would normally be occupied by the evacuated tail lobes. Shi et al. (2013) 325 interpreted similar observations of such plasma in the "lobes" as an indication of direct 326 ingress of solar wind plasma via (single) lobe reconnection. However, it seems unlikely 327 that this plasma should reside in the near-Earth tail for long, rather than disappearing 328 down the tail along open field lines, as occurs during periods of southwards IMF. Instead, 329 we suggest that this plasma must be trapped on closed field lines. This interpretation 330 is supported by the presence of a double loss-cone in the C4 electron pitch angle distri-331 butions, similar to the plasma characteristics seen by Fear et al. (2014) at high Z over 332 a transpolar arc. In that case, the plasma was only observed by Cluster on field lines that 333 mapped to the arc, and evacuated lobe was seen to either side; in our case the plasma 334 is observed for prolonged periods (sometimes many hours) wherever C4 is located in the 335 tail. The double loss-cone indicates that the plasma has interacted significantly with the 336 atmosphere in both the northern and southern hemispheres over multiple bounces. The 337 presence of counter-streaming ions observed by C4 is also consistent with closed field lines 338 and trapped plasma, as are the sunwards/tailwards fluxes observed by Geotail: if the mag-339 netic field was open, only tailward fluxes would be expected. This trapped plasma is ob-340 served in both the northern and southern portions of the magnetotail, at $Z \approx 8 R_E$ 341 and $Z \approx -13 R_E$, simultaneously (see Figures 4 and 6). 342

Several mechanisms have been proposed by which plasma enters the tail during NBZ 343 conditions, and whether or not it is trapped (e.g., Taylor et al., 2008; Shi et al., 2013). 344 Milan, Carter, Bower, et al. (2020) argued that dual-lobe reconnection explains both the 345 capture and trapping of plasma, and the ionospheric flow pattern and auroral evolution 346 observed during the formation of horse-collar auroras (HCAs), and by extension CAAs 347 (Milan et al., 2022). It also explains the necessity for near-zero clock angle, $|\theta| < 45^{\circ}$, 348 for the appearance of HCAs (Bower, Milan, Paxton, & Anderson, 2022) and CAAs (this 349 study). We suggest now that this also explains why the polar cap closes and the tail loses 350 its northern and southern lobes, explaining trapped plasma on field lines that map to 351 what would normally be the central polar cap. 352

We note that between 22 UT, DOY 285, and 08 UT, DOY 286, there were two brief swings of the clock angle to $|\theta| < 45^{\circ}$, but no CAAs were detected and open lobe was observed by C4 (see Figure 4). This suggests that there is a minimum duration of DLR of one to two hours for plasma to be trapped. This minimum will be related to the reconnection rate, that is the rate at which flux is closed, such that open lobe is replaced by closed flux containing trapped solar wind plasma. Detailed measurement of the rate
of closure, difficult with low cadence DMSP/SSUSI images, will be required to further
understand this. However, ionospheric convection measurements (e.g., Chisham et al.,
2004, 2008) should help quantify the reconnection rate.

The plasma trapped by DLR then acts as a source for precipitation to produce au-362 roral emission in the polar regions. As shown by Q.-H. Zhang et al. (2020) and Milan 363 et al. (2022), CAAs are produced by inverted-V precipitation, electrons accelerated to 364 energies of a few keV, at shears in the ionospheric convection flow which are associated 365 with upwards field-aligned currents due to converging ionospheric electric fields. This 366 is consistent with the trapped electron population found in this study: in the magneto-367 tail the electrons have energies primarily below one keV and the electron pitch angle dis-368 tribution has empty loss cones. These electrons will not precipitate without the accel-369 eration provided by flow shears. Once accelerated, the electrons are seen to have ener-370 gies of several keV in the ionosphere. Precipitating ions with energies of one to several 371 keV are also observed in the ionosphere when CAAs are present, especially towards the 372 nightside, consistent with the trapped ions found in this study. Whereas Carlson and 373 Cowley (2005) proposed that weak polar cap arcs, perhaps indeed CAAs, are produced 374 by accelerated polar rain on open field lines, in our scenario trapped solar wind plasma 375 on closed field lines is the more readily available source population. 376

On the large scale the ionospheric flow pattern has reverse lobe convection cells, 377 consistent with dual-lobe reconnection (Milan et al., 2022), though on smaller scales mul-378 tiple flow shears are seen, associated with the CAAs. Q.-H. Zhang et al. (2020) and Wang 379 et al. (2023) suggested that these flow shears are produced by Kelvin-Helmholtz waves 380 on the magnetospheric flanks propagating into the magnetotail. However, the cusp-aligned 381 nature of the arcs shows that the flow shears are also cusp-aligned, which is not neces-382 sarily predicted by the KHI mechanism. Instead, Milan et al. (2022) proposed that the 383 flow shears are excited by temporally- and spatially-varying lobe reconnection rates, which 384 explains why the flows and shears naturally radiate from the cusp region. We prefer this 385 latter explanation: lobe reconnection explains the closure of magnetic flux and the trap-386 ping of solar wind plasma, it explains the "reverse" flow pattern observed in the iono-387 sphere, it explains the structuring of the auroral precipitation into multiple arcs, and it 388 also explains why the arcs are "cusp-aligned". That the field is closed at high latitudes 389 also explains why CAAs are generally seen in both hemispheres simultaneously (see Fig-390 ure 2 and Milan et al. (2022), whereas other polar cap auroral phenomena are not al-391 ways conjugate (e.g., Reidy et al., 2020; Bower, Milan, Paxton, & Imber, 2022). 392

That auroral emission is observed across the polar regions suggests that the mag-393 netosphere is almost entirely closed, though it might be expected that open and closed 394 flux is interspersed (Milan et al., 2022). As proposed by Milan et al. (2022), horse-collar 395 auroras are the preliminary stage in the development of CAAs, forming when dual-lobe 396 reconnection first commences (Milan, Carter, Bower, et al., 2020). In other words, it is 397 possible that HCAs form frequently (Bower, Milan, Paxton, & Anderson, 2022) but do 398 not necessarily fully evolve into CAAs if the IMF turns away from near-zero clock an-399 gles. The high-latitude arcs which sit at the dawn and dusk edges of the polar slot of 400 the HCA configuration are dimmer than the main auroral oval (Bower, Milan, Paxton, 401 & Anderson, 2022), but are still bright enough to be seen with global auroral imagers 402 (Cumnock & Blomberg, 2004) and can be misinterpreted as transpolar arcs (Milan, Carter, 403 Bower, et al., 2020). However, these HCA arcs are in general brighter than CAAs and 404 so may be detected more frequently, certainly with previous global auroral imagers. As 405 mentioned previously, the occurrence of CAAs is under-studied and a statistical survey 406 is required, especially as this will provide new insights into the occurrence of DLR. 407

As an aside, we note that the magnitude of the B_X component of the IMF was significant at times, e.g. ≈ 10 nT during the first period of CAAs in Figure 3, and was nearzero at others, e.g. the last period of CAAs in Figure 4. This suggests that B_X does not ⁴¹¹ play a role in modulating the occurrence of (dual-) lobe reconnection and hence the oc-⁴¹² currence of CAAs. This tallies with a lack of B_X control of the occurrence of HCAs re-⁴¹³ ported by Bower, Milan, Paxton, and Anderson (2022).

⁴¹⁴ Once DLR ceases the magnetosphere loses the trapped plasma and the polar cap ⁴¹⁵ reforms promptly, indicating the rapid opening of magnetic flux by magnetopause re-⁴¹⁶ connection. This opening will occur most rapidly if the IMF is directed southwards and ⁴¹⁷ subsolar reconnection occurs, which will be accompanied by twin-cell convection in the ⁴¹⁸ polar ionosphere. However, as noted by Milan et al. (2022), it will also occur if the IMF ⁴¹⁹ is directed northwards with $|\theta| > 45^{\circ}$ as single lobe reconnection will open a closed mag-⁴²⁰ netosphere. Hence, CAAs are only seen for $|\theta| < 45^{\circ}$.

Empirical magnetic field models, such as T96, do not reproduce a closed magne-421 tosphere well, though closed NBZ magnetospheres are readily formed in simulations (e.g., 422 Song et al., 1999; Siscoe et al., 2011; Wang et al., 2023). The magnetic field measure-423 ments made by C4/FGM and Geotail/MGF indicated that the B_X component was re-424 duced when CAAs were present, with respect to when open lobe was observed. This in-425 dicates that the tail is somewhat deflated when it is closed (though it could also be in 426 part because plasma pressure contributes to stress balance in the closed tail, rather than 427 just magnetic pressure in the lobes). However, otherwise the field did not deviate markedly 428 from the T96 predictions, indicating that if the magnetosphere is indeed closed, the mag-429 netic structure near-Earth is not significantly modified, suggesting that the closed field 430 lines stretch considerably further down-tail than the locations of C4 and Geotail. This 431 is in contradiction to simulations which suggest that a closed magnetotail might be less 432 than 20 R_E in length (Wang et al., 2023). Indeed, Milan, Carter, and Hubert (2020) es-433 timated that closed field lines associated with transpolar arcs could extend as far as X <434 $-100 R_E$. More work needs to be conducted in the distant magnetotail and with mag-435 netospheric simulations to understand the magnetic topology of the closed magnetosphere. 436

437 4 Conclusions

We have investigated a fifteen-day period during which Cluster and Geotail sam-438 pled the magnetotail. Observations of the auroras by DMSP/SSUSI indicates that cusp-439 aligned arcs were present in the high latitude polar regions whenever the IMF clock an-440 gle was small, $|\theta| < 45^{\circ}$, which during the study interval amounted to approximately 441 30% of the time. Simultaneous observations of ions and electrons by Cluster and Geo-442 tail show significant plasma densities $(n_i \approx n_e \approx 1 \text{ cm}^{-3})$ in the tail during these in-443 tervals, even as far from the equatorial plane as $|Z| \approx 13 R_E$. This region of the mag-444 netotail would normally be devoid of plasma, being occupied by the open flux of the mag-445 netotail lobe. The presence of counter-streaming ions and double loss-cone electrons sug-446 gest, instead, that the plasma was trapped, i.e., that the magnetic field was closed. This 447 trapped plasma will provide a source for the CAA auroral emission, and as the auroral 448 emission covered the polar regions, we further suggest that the magnetosphere was al-449 most entirely closed. Coxon et al. (2021) recently showed that hot plasma consistent with 450 trapping on closed field lines is frequently seen at $|Z| > 5 R_E$ during northward IMF; 451 here we have shown that CAAs are the auroral signature of this closed flux, and that at 452 such times the magnetosphere is likely (almost) entirely closed. We believe that this clo-453 sure is achieved by dual-lobe reconnection, as proposed by Milan, Carter, Bower, et al. 454 (2020) and Milan et al. (2022). Our observations indicate that closure of the magneto-455 sphere is a common occurrence. 456

457 **5 Open Research**

⁴⁵⁸ Data from the Cluster Ion Spectrometry (CIS) instrument, the Plasma Electron
 ⁴⁵⁹ And Current Experiment (PEACE), and the Fluxgate Magnetometer (FGM) were ac ⁴⁶⁰ cessed through the Cluster Science Archive (CSA, formerly Cluster Active Archive (Laakso

et al., 2010)) at https://www.cosmos.esa.int/web/csa, maintained by ESA/ESTEC. 461 Geotail Magnetic Field Experiment (MGF) and Low Energy Particle (LEP) data were 462 accessed through the DARTS Solar-Terrestrial Physics data portal (https://www.darts 463 .isas.jaxa.jp/stp/geotail/data.html) maintained by ISAS/JAXA. The Defense Meteorological Satellite Program (DMSP) Special Sensor Ultraviolet Spectrographic Instru-465 ment (SSUSI) file type EDR-AUR data were obtained from JHU/APL (http://ssusi 466 . jhuapl.edu, data version 0106, software version 7.0.0, calibration period version E0018). 467 Advanced Magnetosphere and Planetary Electrodynamics Response Experiment (AM-468 PERE) data were obtained from JHU/APL (http://ampere.jhuapl.edu/dataget/index 469 .html) and processed using software provided (http://ampere.jhuapl.edu/). The high 470 resolution (1-min) OMNI data used in this study were obtained from the NASA God-471

dard Space Flight Center Space Physics Data Facility OMNIWeb portal (https://omniweb

.gsfc.nasa.gov/form/om_filt_min.html).

474 Acknowledgments

SEM and MKM were supported by the Science and Technology Facilities Council (STFC), 475 UK, grant no. ST/S000429/1. SEM and GEB were supported by the Natural Environ-476 ment Research Council (NERC), UK, grant no. NE/W006766/1. We are grateful to the 477 Cluster CIS, PEACE, and FGM instrument teams for providing data. We acknowledge 478 use of NASA/GSFC's Space Physics Data Facility's CDAWeb service (at http://cdaweb 479 .gsfc.nasa.gov), and OMNI data, and the Cluster Science Archive (at https://www 480 .cosmos.esa.int/web/csa), maintained by ESA/ESTEC. Geotail magnetic field and 481 plasma data were provided by T. Nagai and Y. Saito through DARTS at the Institute 482 of Space and Astronomical Science, JAXA, in Japan. 483

484 References

- Anderson, B., Takahashi, K., & Toth, B. (2000). Sensing global Birkeland currents
 with Iridium® engineering magnetometer data. *Geophysical Research Letters*, 27(24), 4045–4048. doi: https://doi.org/10.1029/2000GL000094
- Balogh, A., Dunlop, M., Cowley, S., Southwood, D., Thomlinson, J., Glassmeier, K.,
 ... others (1997). The Cluster magnetic field investigation. Space Science Reviews, 79, 65–91. doi: https://doi.org/10.1023/A:1004970907748
- Bower, G., Milan, S., Paxton, L., & Anderson, B. (2022). Occurrence statis tics of horse collar aurora. Journal of Geophysical Research: Space Physics,
 e2022JA030385. doi: https://doi.org/10.1029/2022JA030385
- Bower, G., Milan, S., Paxton, L., & Imber, S. (2022). Transpolar arcs: Seasonal de pendence identified by an automated detection algorithm. Journal of Geophys *ical Research: Space Physics*, 127(1), e2021JA029743. doi: https://doi.org/10
 .1029/2021JA029743
- Carlson, H., & Cowley, S. (2005). Accelerated polar rain electrons as the source of
 sun-aligned arcs in the polar cap during northward interplanetary magnetic
 field conditions. Journal of Geophysical Research: Space Physics, 110(A5).
- Carter, J., Milan, S., Fogg, A., Paxton, L., & Anderson, B. (2018). The association of high-latitude dayside aurora with NBZ field-aligned currents. *Journal* of Geophysical Research: Space Physics, 123(5), 3637–3645. doi: https://doi .org/10.1029/2017JA025082
- Carter, J., Milan, S., Fogg, A., Sangha, H., Lester, M., Paxton, L., & Anderson,
 B. (2020). The evolution of long-duration cusp spot emission during lobe
 reconnection with respect to field-aligned currents. *Journal of Geophysical Research: Space Physics*, 125, e2020JA027922. doi: https://doi.org/10.1029/
 2020JA027922

⁵¹⁰ Chisham, G., Freeman, M., Abel, G., Lam, M., Pinnock, M., Coleman, I., ... Vil ⁵¹¹ lain, J.-P. (2008). Remote sensing of the spatial and temporal structure of

512	magnetopause and magnetotail reconnection from the ionosphere. Reviews of Geophysics $46(1)$ doi: https://doi.org/10.1029/2007BG000223
515	Chicham C. Freeman M. Coleman I. Pinnock M. Hairston M. Lester M. &
514	Sofko G (2004) Measuring the dayside reconnection rate during an interval
515	of due northward interplanetary magnetic field Annales Geonhusicae 22(12)
510	4243-4258 doi: https://doi.org/10.5194/angeo-22-4243-2004
517	Chicham C. Lester M. Milan S. E. Freeman M. Bristow W. Crocott A
518	others (2007) A decade of the Super Dual Auroral Radar Network (Super-
519	DARN): Scientific achievements, new techniques and future directions. Surveys
520	in Geonhysics 28(1) 33-109 doi: https://doi.org/10.1007/s10712-007-9017-8
521	Cowley S (1981) Magnetospheric and ionospheric flow and the interplanetary mag-
522	netic field The Physical Basis of the Jonosphere in the Solar-Terrestrial Sus-
523	tem $ACARD_CP_{-}205 (A_{-}1)_{-}(A_{-}1A)$
524	Cowley S & Lackwood M (1002) Excitation and decay of solar wind driven flows
525	in the magnetosphere ionosphere system Annales Geophysicae 10, 103–115
526	Covon I Foar B Boidy I Fryer I & Dlank I (2021) Hot plasma in the
527	magnetetail lobes shows characteristics consistent with closed field lines
528	trapped in the lobes — <i>Lowral of Coophysical Passarch: Space Physica</i> , 196(0)
529	$_{2021}$ IA020516 doi: https://doi.org/10.1020/2021 IA020516
530	Covon I Milan S & Anderson B (2018) A review of Birkeland current research
531	using AMPERE Electric Currents in Coorners and Bound 257–278 doi:
532	$\frac{1}{1000}$ https://doi.org/10.1002/0781110324522.ch16
533	Cumpeck I & Blomberg I (2004) Transpolar are evolution and associated no
534	tential patterns Annales Geonhusicae $20(4)$ 1213–1231 doi: https://doi.org/
535	10 5194/angeo. 22.1213.2004
530	Cumpock I Sharber I Healis B Blomberg I. C. Cermany C. Spann I. &
537	Coley W (2002) Interplanetary magnetic field control of theta aurora devel-
530	opment Journal of Geophysical Research: Space Physics 107(A7) SIA-4 doi:
540	https://doi.org/10.1029/2001.IA009126
5/1	Dungey J (1961) Interplanetary magnetic field and the auroral zones <i>Physical Re</i> -
542	$p_{\text{angle}}(1) = p_{\text{angle}}(1) + p_{angle$
542	Dungev J (1963) The structure of the exosphere or adventures in velocity space
544	Geophysics. The Earth's Environment.
545	Escoubet, C., Fehringer, M., & Goldstein, M. (2001). Introduction the cluster mis-
546	sion. In Annales geophysicae (Vol. 19, pp. 1197–1200), doi: https://doi.org/10
547	.5194/angeo-19-1197-2001
548	Fear, R. (2021). The northward IMF magnetosphere. <i>Magnetospheres in the Solar</i>
549	Sustem, 293–309. doi: https://doi.org/10.1002/9781119815624.ch19
550	Fear, R., & Milan, S. (2012a). The IMF dependence of the local time of transpolar
551	arcs: Implications for formation mechanism. Journal of Geophysical Research:
552	Space Physics, 117(A3).
553	Fear, R., & Milan, S. (2012b). Ionospheric flows relating to transpolar arc formation.
554	Journal of Geophysical Research: Space Physics, 117(A9).
555	Fear, R., Milan, S., Maggiolo, R., Fazakerley, A., Dandouras, L. & Mende, S. (2014).
556	Direct observation of closed magnetic flux trapped in the high-latitude mag-
557	netosphere. Science, 346 (6216), 1506–1510. doi: https://doi.org/10.1126/
558	science.1257377
559	Frank, L., Craven, J., Burch, J., & Winningham, J. (1982). Polar views of the
560	Earth's aurora with Dynamics Explorer. Geophysical Research Letters. 9(9).
561	1001–1004. doi: https://doi.org/10.1029/GL009i009p01001
562	Frey, H. (2007). Localized aurora beyond the auroral oval. <i>Reviews of Geonhusics</i> .
563	45(1). doi: https://doi.org/10.1029/2005RG000174
564	Frey, H., Mende, S., Immel, T., Fuselier, S., Claffin, E., Gérard, JC., & Hubert. B.
565	(2002). Proton aurora in the cusp. Journal of Geophysical Research: Space
566	Physics, 107(A7), SMP-2. doi: https://doi.org/10.1029/2001JA900161

567 568	Fryer, L., Fear, R., Coxon, J., & Gingell, I. (2021). Observations of closed magnetic flux embedded in the lobes during periods of northward imf. <i>Journal of Comparison Process Physics</i> , 196(6), e20211A020281, doi: https://doi.
569 570	<i>Geophysical Research: Space Physics</i> , 126(6), e2021JA029281. doi: https://doi .org/10.1029/2021JA029281
571	Fuselier, S., Trattner, K., Petrinec, S., & Lavraud, B. (2012). Davside magnetic
572	topology at the Earth's magnetopause for northward IMF. Journal of Geo-
573	physical Research: Space Physics, 117(A8). doi: https://doi.org/10.1029/
574	2012JA017852
575	Hones Jr, E., Craven, J., Frank, L., Evans, D., & Newell, P. (1989). The horse-collar
576	aurora: A frequent pattern of the aurora in quiet times. <i>Geophysical Research</i>
577	Letters, 16(1), 37–40. doi: https://doi.org/10.1029/GL016i001p00037
578	Hosokawa, K., Kullen, A., Milan, S., Reidy, J., Zou, Y., Frey, H. U., Fear, R.
579	(2020). Aurora in the polar cap: A review. Space Science Reviews, 216(1), 15.
580	doi: https://doi.org/10.1007/s11214-020-0637-3
581	Hosokawa, K., Moen, J., Shiokawa, K., & Otsuka, Y. (2011). Motion of polar cap
582	arcs. Journal of Geophysical Research: Space Physics, 116(A1). doi: https://
583	doi.org/10.1029/2010JA015906
584	Huang, CS., Sofko, G., Koustov, A., Andre, D., Ruohoniemi, J., Greenwald,
585	R., & Hairston, M. (2000). Evolution of ionospheric multicell convection
586	during northward interplanetary magnetic field with $ B_z/B_u > 1$. Jour-
587	nal of Geophysical Research: Space Physics, 105(A12), 27095–27107. doi:
588	https://doi.org/10.1029/2000JA000163
589	Iijima, T., & Potemra, T. (1976). The amplitude distribution of field-aligned cur-
590	rents at northern high latitudes observed by Triad. Journal of Geophysical Re-
591	search, 81(13), 2165–2174. doi: https://doi.org/10.1029/JA081i013p02165
592	Iijima, T., Potemra, T., Zanetti, L., & Bythrow, P. (1984). Large-scale Birkeland
593	currents in the dayside polar region during strongly northward IMF: A new
594	Birkeland current system. Journal of Geophysical Research: Space Physics,
595	89(A9), 7441–7452. doi: https://doi.org/10.1029/JA089iA09p07441
596	Imber, S., Milan, S., & Hubert, B. (2006). The auroral and ionospheric flow signa-
597	tures of dual lobe reconnection. Annales Geophyiscae, $24(11)$, $3115-3129$. doi:
598	https://doi.org/10.5194/angeo-24-3115-2006
599	Ismail, S., Wallis, D., & Cogger, L. (1977). Characteristics of polar cap Sun-aligned
600	arcs. Journal of Geophysical Research, 82(29), 4741–4749. doi: https://doi
601	.org/10.1029/JA082i029p04741
602	Johnstone, A., Alsop, C., Burge, S., Carter, P., Coates, A., Coker, A., others
603	(1997). Peace: A plasma electron and current experiment. Space Science
604	Reviews, 351-398. doi: https://doi.org/10.1023/A:1004938001388
605	King, J., & Papitashvili, N. (2005). Solar wind spatial scales in and comparisons
606	of hourly Wind and ACE plasma and magnetic field data. Journal of Geo-
607	physical Research: Space Physics, 110(A2). doi: https://doi.org/10.1029/
608	2004JA010649
609	Kokubun, S., Yamamoto, T., Acuña, M., Hayashi, K., Shiokawa, K., & Kawano, H.
610	(1994). The GEOTAIL magnetic field experiment. Journal of geomagnetism
611	and geoelectricity, 46(1), 7–21. doi: https://doi.org/10.5636/jgg.46.7
612	Kullen, A., Brittnacher, M., Cumnock, J., & Blomberg, L. (2002). Solar wind
613	dependence of the occurrence and motion of polar auroral arcs: A statistical
614	study. Journal of Geophysical Research: Space Physics, 107(A11), 13–1. doi:
615	nttps://doi.org/10.1029/2002JA009245
616	Laakso, H., Perry, C., McCaffrey, S., Herment, D., Allen, A., Harvey, C., Turner,
617	R. (2010). Cluster active archive: Overview. Springer.
618	Lockwood, M., & Cowley, S. (1992). Ionospheric convection and the substorm cycle.
619	Proceedings of the International Conference on Substorms (ICS-1), 99–109.
620	Mailyan, B., Shi, Q., Kullen, A., Maggiolo, R., Zhang, Y., Fear, R., others
621	(2015). Transpolar arc observation after solar wind entry into the high-latitude

622	magnetosphere. Journal of Geophysical Research: Space Physics, 120(5),
623	3525-3534.
624	Milan, S., Bower, G., Carter, J., Paxton, L., Anderson, B., & Hairston, M. (2022).
625	Lobe reconnection and cusp-aligned auroral arcs. Journal of Geophysical Re-
626	search: Space Physics, 127(6), e2021JA030089. doi: https://doi.org/10.1029/
627	2021JA030089
628	Milan, S., Carter, J., Bower, G., Imber, S., Paxton, L., Anderson, B., Hu-
629	bert, B. (2020). Dual-lobe reconnection and horse-collar auroras. Jour-
630	nal of Geophysical Research: Space Physics, $125(10)$, $e2020JA028567$. doi:
631	https://doi.org/10.1029/2020JA028567
632	Milan, S., Carter, J., & Hubert, B. (2020). Probing the magnetic structure of a pair
633	of transpolar arcs with a solar wind pressure step. Journal of Geophysical Re- $l_{1} = C$ $R_{1} = 105(0)$ $R_{2} = 00101 A_{1}00710C$
634	search: Space Physics, $125(2)$, $e2019JA027196$.
635	Milan, S., Carter, J., Sangha, H., Bower, G., & Anderson, B. (2021). Magne-
636	to spheric flux throughput in the dungey cycle: Identification of convection $f(2)$
637	state during 2010. Journal of Geophysical Research: Space Physics, 120(2),
638	$e_{2020JA028437}$ doi: https://doi.org/10.1029/2020JA028437
639	Milan, S., Cowley, S., Lester, M., Wright, D., Slavin, J., Fillingim, M., Singer, H.
640	(2004). Response of the magnetotal to changes in the open flux content of the magnetotal $P_{\text{magnetotal}}$ and $P_{\text{magnetotal}}$ $P_{\text{magnetotal}$ $P_{\text{magnetotal}}$ $P_{\text{magnetotal}$ $P_{\text{magnetotal}}$ $P_{\text{magnetotal}$ $P_{\text{magnetotal}}$ $P_{\text{magnetotal}$ $P_{\text{magnetotal}$ $P_{\text{magnetotal}$ $P_{\text{magnetotal}$ $P_{\text{magnetotal}}$ $P_{\text{magnetotal}$ $P_{\text{magnetotal}}$
641	magnetosphere. Journal of Geophysical Research: Space Physics, 109(A4). doi: https://doi.org/10.1020/20021A010250
642	Milen S. Hubert D. & Creastt A. (2005). Formation and mation of a transmo
643	Milan, S., Hubert, B., & Grocott, A. (2005). Formation and motion of a transpo-
644	har arc in response to dayside and ingliside reconnection. $Journal of Geo-$
645	physical Research. Space Physics, 110(A1). doi: https://doi.org/10.1029/20041A010825
646	Milen S. Loster, M. Cowley, S. & Brittnacher, M. (2000) Develde con
647	westion and auroral morphology during an interval of northward inter
648	planetary magnetic field $Annales Ceonhusicae 18(A) A36-AAA doi:$
650	https://doi org/10/1007/s00585-000-0436-9
651	Milan S. Lester M. Cowley S. Oksavik K. Brittnacher M. Greenwald B.
652	Villain J-P (2003) Variations in the polar cap area during two substorm
653	cycles Annales Geonhusicae 21(5) 1121–1140
654	Milan, S., Provan, G., & Hubert, B. (2007). Magnetic flux transport in the Dungev
655	cycle: A survey of dayside and nightside reconnection rates. Journal of Geo-
656	physical Research: Space Physics, 112(A1). doi: https://doi.org/10.1029/
657	2006JA011642
658	Mukai, T., Machida, S., Saito, Y., Hirahara, M., Terasawa, T., Kaya, N.,
659	Nishida, A. (1994). The low energy particle (LEP) experiment onboard the
660	Geotail satellite. Journal of geomagnetism and geoelectricity, 46(8), 669–692.
661	doi: https://doi.org/10.5636/jgg.46.669
662	Nishida, A. (1994). The geotail mission (Vol. 21) (No. 25). Wiley Online Library.
663	Øieroset, M., Raeder, J., Phan, T., Wing, S., McFadden, J., Li, W., Balogh, A.
664	(2005). Global cooling and densification of the plasma sheet during an ex-
665	tended period of purely northward IMF on October 22–24, 2003. Geophysical
666	Research Letters, $32(12)$.
667	Paxton, L., Meng, CI., Fountain, G., Ogorzalek, B., Darlington, E., Gary, S.,
668	others (1992). Special sensor ultraviolet spectrographic imager: An instru-
669	ment description. In Instrumentation for planetary and terrestrial atmospheric
670	remote sensing (Vol. 1745, pp. 2–15). doi: https://doi.org/10.1117/12.60595
671	Paxton, L., Schaefer, R., Zhang, Y., & Kil, H. (2017). Far ultraviolet instrument
672	technology. Journal of Geophysical Research: Space Physics, 122(2), 2706–
673	2733. doi: https://doi.org/10.1002/2016JA023578
674	Paxton, L., & Zhang, Y. (2016). Far ultraviolet imaging of the aurora. In Space
675	weather fundamentals (pp. 213–244). CRC Press.
676	Paxton, L., Zhang, Y., Kil, H., & Schaefer, R. (2021). Exploring the upper atmo-

677	sphere: Using optical remote sensing. Upper Atmosphere Dynamics and Ener-
678	getics, 487–522. doi: https://doi.org/10.1002/9781119815631.ch23
679	Reidy, J., Fear, R., Whiter, D., Lanchester, B., Kavanagh, A., Price, D. J., Pax-
680	ton, L. (2020). Multiscale observation of two polar cap arcs occurring on
681	different magnetic field topologies. Journal of Geophysical Research: Space
682	Physics, 125(8), e2019JA027611.
683	Rème, H., Bosqued, J., Sauvaud, J., Cros, A., Dandouras, J., Aoustin, C., others
684	(1997). The cluster ion spectrometry (cis) experiment. The cluster and phoenix
685	missions, $303-350$. doi: https://doi.org/10.1007/978-94-011-5666-0_12
686	Rich, F., & Hairston, M. (1994). Large-scale convection patterns observed by
687	DMSP. Journal of Geophysical Research: Space Physics, 99(A3), 3827–3844.
688	doi: https://doi.org/10.1029/93JA03296
689	Shi, Q., Zong, QG., Fu, S., Dunlop, M., Pu, Z., Parks, G., others (2013). Solar
690	wind entry into the high-latitude terrestrial magnetosphere during geomagneti-
691	cally quiet times. Nature communications, 4, 1466.
692	Siscoe, G., Farrugia, C., & Sandholt, P. (2011). Comparison between the two basic
693	modes of magnetospheric convection. Journal of Geophysical Research: Space
694	<i>Physics</i> , 116(A5). doi: https://doi.org/10.1029/2010JA015842
695	Siscoe, G., & Huang, T. (1985). Polar cap inflation and deflation. <i>Journal of Geo</i> -
696	physical Research: Space Physics, 90(A1), 543–547.
697	Song, P., DeZeeuw, D., Gombosi, T., Groth, C., & Powell, K. (1999). A numerical
698	study of solar wind-magnetosphere interaction for northward interplanetary
699	magnetic field. Journal of Geophysical Research: Space Physics, 104 (A12),
700	28301-28378. doi: https://doi.org/10.1029/1999JA900378
701	(2008) The plagma short and boundary layors under northward IME: A multi-
702	(2008). The plasma sheet and boundary layers under northward hvir. A multi- point and multi instrument perspective $Advances in Space Becauch (1(10))$
703	1610–1620
704	1019^{-1029} . Tsympton N (1005) Modeling the earth's magnetospheric magnetic field con
705	fined within a realistic magnetonause <u>Journal of Geonhusical Research</u> : Snace
700	Physics $100(A4)$ 5599–5612 doi: https://doi.org/10.1029/941A03193
707	Wang X - Y Zhang O - H Wang C Zhang Y - L Tang B - B Xing Z - Y
709	others (2023). Unusual shrinkage and reshaping of earth's magnetosphere un-
710	der a strong northward interplanetary magnetic field. <i>Communications Earth</i>
711	<i>Estimate a morphalical formation of the second sec</i>
712	Waters, C., Anderson, B., & Liou, K. (2001). Estimation of global field aligned
713	currents using the Iridium $(\widehat{\mathbf{R}})$ system magnetometer data. Geophysical Research
714	Letters, 28(11), 2165–2168. doi: https://doi.org/10.1029/2000GL012725
715	Zhang, QH., Zhang, YL., Wang, C., Lockwood, M., Yang, HG., Tang, BB.,
716	others (2020). Multiple transpolar auroral arcs reveal insight about
717	coupling processes in the Earth's magnetotail. Proceedings of the National
718	Academy of Sciences, 117(28), 16193–16198. doi: https://doi.org/10.1073/
719	pnas.2000614117
720	Zhang, Y., Paxton, L., Zhang, QH., & Xing, Z. (2016). Polar cap arcs: Sun-aligned
721	or cusp-aligned? Journal of Atmospheric and Solar-Terrestrial Physics, 146,
722	123–128. doi: https://doi.org/10.1016/j.jastp.2016.06.001