

# Is equilibrium climate sensitivity clouding our judgement?

Jonny Williams<sup>1</sup>

<sup>1</sup>NIWA

March 6, 2023

## Abstract

Equilibrium climate sensitivity - ECS - is easily-understood, has been studied for over 150 years and is therefore appealing as a metric for communication of climate model results. In this work I argue that ECS is not a good metric for comparing different climate models. Via brief examples concerning the Pliocene epoch and the Paleocene-Eocene Thermal Maximum , I further argue that models which produce temperatures towards the higher end of model intercomparisons are useful in spite of recent studies concluding that these models are ‘too hot’. I hope that this brief manuscript generates discussion on how to prioritise the consideration of more useful, and potentially novel, ways of comparing climate models going forward.

# Is equilibrium climate sensitivity clouding our judgement?

Jonny Williams<sup>1,\*</sup>

<sup>1</sup>*NIWA, Wellington, New Zealand\**

*\*Corresponding author: Jonny Williams,  
jonny.williams@niwa.co.nz*

March 2023

## Abstract

Equilibrium climate sensitivity – ECS – is easily-understood, has been studied for over 150 years and is therefore appealing as a metric for communication of climate model results. In this work I argue that ECS is not a good metric for comparing different climate models. Via brief examples concerning the Pliocene epoch and the Paleocene–Eocene Thermal Maximum, I further argue that models which produce temperatures towards the higher end of model intercomparisons are useful in spite of recent studies concluding that these models are ‘too hot’. I hope that this brief manuscript generates discussion on how to prioritise the consideration of more useful, and potentially novel, ways of comparing climate models going forward.

## 1 Introduction

The climate is warming due to anthropogenic forcing [1]; so much so in fact that we have entered a new geological period called the Anthropocene [2]. Quantifying the amount of warming that is expected for an additional, marginal amount of greenhouse gases in the atmosphere is – in general – known as the climate sensitivity. In particular, the *equilibrium* climate sensitivity – ECS – is defined as the global mean atmospheric warming for a doubling of carbon dioxide – CO<sub>2</sub> – with respect to preindustrial conditions.

The ECS has a long history and – due to its simple definition – is often used as a zeroth-order, comparative measure of different climate models’ ability to reproduce observed warming. These difference models have, since the early 1990s, been compared in several generations of Model Intercomparison Projects

---

\*jonny.williams@niwa.co.nz




Figure 1: Figure TS.16(a) in IPCC, 2021: Technical Summary. In: Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change.

or MIPs and the range of the predicted ECS has remained about the same, see Figure 1.

Ostensibly, this could be seen as a failure of the models to converge on the ‘correct’ value of the ECS. This however, obscures the fact that the application of the concept of ECS is somewhat misleading when applied to today’s complex climate models.

## 2 History

Estimates of ECS date back to the late 19th century when Arrhenius estimated values between approximately 4-6° [3]. Ever since, and in spite of spectacular advances in fundamental scientific understanding and computational resources, the ECS remains elusive.

The main reason for this stubborn refusal to converge lies in the continual ‘moving of the goal posts’ with respect to the system being studied. The first studies were the best part of a century before the first computational estimates of ECS were published in 1967 by Manabe and Wetherald and computer simulations have steadily increased in complexity and resource requirements ever since.

Computational studies have moved from considerations of the atmosphere alone <sup>1</sup> to inclusion of land-surface processes, dynamic ocean models, complex atmospheric chemistry, sea-ice, ice-sheets and ocean biogeochemistry. The addition of each of these components – while undoubtedly beneficial

<sup>1</sup>Even within atmosphere-only models there is a hierarchy of complexity; with and without water vapour feedbacks for example [4]

to the understanding in fundamental science – have each introduced additional uncertainties in the response of the systems to increased greenhouse gas forcing. Indeed given the level of complexity that the current state of the art models represent – bearing in mind that all climate models necessarily differ in terms of the processes they represent – it is surely not surprising that the estimates of ECS differ so much;  $2 \lesssim \text{ECS} \lesssim 5.5$  to the nearest  $0.5^\circ$  [5].

### 3 Feedbacks

Key to the argument that ECS is ‘under-defined’ is the concept of feedbacks, such as the relationship between ice amount and albedo (reflectivity). As high-reflectivity ice melts and the underlying land or ocean is exposed, less sunlight is reflected directly back into space, thus causing more warming, and so on. In atmosphere-only models, this feedback is inherently absent – i.e there is no dynamic ice model – and so when ECS is predicted using an atmosphere coupled to a sea-ice model, the estimate of ECS will change. The same is true for considerations of grounded ice sheets, the inclusion of which is one of the most recent additions to state-of-the-art climate models [6].

Even recent advances in ocean biogeochemistry [7] introduce feedbacks, for example by allowing surface gas exchange to planktonic species, which are themselves bound by predictions of ocean circulation. Following this, ocean circulation is far from constant on timescales longer than a century or so. Indeed consideration of the Gulf Stream which keeps northern Europe warmer than other similar-latitude locations make it clear that changes to ocean circulation could potentially have enormous repercussions [8]. All of this is without any discussion *at all* of the longer-term implications gleaned from paleoclimate studies, e.g. [9].

The above discussion – I hope – shows that use of the ECS as a useful metric for our ability to ‘correctly’ predict the amount of warming expected for a given amount of greenhouse gas forcing is far from ideal. Note that I am deliberately not considering the related metric of transient climate response, or TCR, here for brevity and the interested reader is referred elsewhere [10].

With the initial publication of climate model data from CMIP6, it quickly became apparent that some models were producing climate sensitivities which were noticeably higher than before [11]. Because of this, there has been some hesitancy to ‘believe’ the results produced by the models. This leads me to discuss the ‘what if’ situation of very high ECS.

### 4 Overheated arguments?

Values at the high end of CMIP6 predictions of ECS are arguably unphysical. For example Zhu et al. argue that the high values obtained in CMIP6 are not supported by paleoclimate evidence [12]. However, in [12] the period studied is approximately 50 million years ago and when – in common with essentially

the entire climatic history of the Earth – climate shifted from quasi-equilibrium (QE) state to QE state via changes in paleogeography [13] and orbital cycles [14] for example. Indeed, arguably the most-studied ‘abrupt’ climate change events known – the Paleocene-Eocene thermal maximum PETM – occurred over a period of over 100,000 years, e.g. [15].

The current situation with respect to anthropogenic climate change is that the climate system is no longer moving in a sequence of QEs and hence we have scant idea about the impact of multi-century-timescale changes in ice-sheet dynamics, for example. This is well illustrated by examining the last time that atmospheric CO<sub>2</sub> concentrations exceeded 400 parts per million; the value first exceeded in the Anthropocene in 2013 [16]. The Pliocene epoch, between about 2 and about 5 million years ago, had comparable CO<sub>2</sub> concentrations to today yet was characterised by higher sea-levels [17], extreme polar amplification of warming [18] and a significantly smaller Antarctic ice-sheet compared to today, as shown in Figure 2.

At this point it is important to stress that I am not arguing for the ‘correctness’ of very high values of ECS. However, if the ‘true’ ECS did turn out to be, say, 5° or higher the results would be societally devastating and it seems to me that to ignore this based on studies of an unknowable quantity is – at best – dangerous.

Another potentially important feedback in the climate system as the world warms in the 21st century – although this argument holds for any currently not-quantified feedback – is that of methane release. Although many climate models do include representations of land-atmosphere exchange, there are potentially large and unaccounted-for climate impacts from the release of methane from tropical wetlands [19] and permafrost melt [20].

These possible, additional methane-induced climate impacts are just one example of a process which could increase the amount of warming for a given amount of increased greenhouse gas forcing. Therefore, the study of relatively extreme warming from high-sensitivity models is surely warranted given that if these models are ‘right for the wrong reason’ about a high temperature future, then at least we are collectively better prepared for it.

## 5 Where do we go from here?

As a community of researchers, we have learnt a vast amount about our climate system, how it has changed in the past and how we think it may change in the future. In summary, I feel it is very important not to over-rely on an idealised quantity, the value of which we will arguably never know for sure.

The ECS is undoubtedly a convenient and simple way of distilling future temperature change projections into easily communicable information. However, the over-reliance on its utility as a useful comparative measure of climate models can give the false impression of a lack of progress in understanding fundamental climate processes.

In terms of the ECS’ uncertainty bounds remaining approximately constant




Figure 2: Cross-Chapter Box 2.4, Figure 1 in IPCC, 2021: Chapter 2. In: Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change

over the past few decades – e.g. Figure 1 – it seems to me that there is at least a passing similarity with the fallacy of a 50% probability of rainfall in a weather forecast being interpreted as *we don't know whether it will rain or not*.

Communicating uncertainty in weather and climate studies is clearly one of the most ‘wicked’ problems of our time [21], but over-simplification is not the answer that future generations need.

## References

- [1] IPCC. “Summary for Policymakers”. In: *Climate Change 2021: The Physical Science Basis. Contribution of Working Group I to the Sixth Assessment Report of the Intergovernmental Panel on Climate Change*. Ed. by V. Masson-Delmotte et al. Cambridge, United Kingdom and New York, NY, USA: Cambridge University Press, 2021. URL: [https://www.ipcc.ch/report/ar6/wg1/downloads/report/IPCC\\_AR6\\_WGI\\_SPM.pdf](https://www.ipcc.ch/report/ar6/wg1/downloads/report/IPCC_AR6_WGI_SPM.pdf).
- [2] Simon L Lewis and Mark A Maslin. “Defining the anthropocene”. In: *Nature* 519.7542 (2015), pp. 171–180.
- [3] Andrei G Lapenis. “Arrhenius and the intergovernmental panel on climate change”. In: *Eos, Transactions American Geophysical Union* 79.23 (1998), pp. 271–271.
- [4] William Ingram. “A very simple model for the water vapour feedback on climate change”. In: *Quarterly Journal of the Royal Meteorological Society: A journal of the atmospheric sciences, applied meteorology and physical oceanography* 136.646 (2010), pp. 30–40.
- [5] Laura A McBride et al. “Comparison of CMIP6 historical climate simulations and future projected warming to an empirical model of global climate”. In: *Earth System Dynamics* 12.2 (2021), pp. 545–579.
- [6] Robin S Smith et al. “Coupling the UK Earth System Model to dynamic models of the Greenland and Antarctic ice sheets”. In: *Journal of Advances in Modeling Earth Systems* 13.10 (2021), e2021MS002520.
- [7] Andrew Yool et al. “Evaluating the physical and biogeochemical state of the global ocean component of UKESM1 in CMIP6 historical simulations”. In: *Geoscientific Model Development* 14.6 (2021), pp. 3437–3472.
- [8] Jon Robson et al. “The role of anthropogenic aerosol forcing in the 1850–1985 strengthening of the AMOC in CMIP6 historical simulations”. In: *Journal of Climate* 35.20 (2022), pp. 3243–3263.
- [9] Eelco J Rohling et al. “Comparing climate sensitivity, past and present”. In: *Annual Review of Marine Science* 10 (2018), pp. 261–288.
- [10] JM Gregory and PM Forster. “Transient climate response estimated from radiative forcing and observed temperature change”. In: *Journal of Geophysical Research: Atmospheres* 113.D23 (2008).

- [11] Gerald A Meehl et al. “Context for interpreting equilibrium climate sensitivity and transient climate response from the CMIP6 Earth system models”. In: *Science Advances* 6.26 (2020), eaba1981.
- [12] Jiang Zhu, Christopher J Poulsen, and Bette L Otto-Bliesner. “High climate sensitivity in CMIP6 model not supported by paleoclimate”. In: *Nature Climate Change* 10.5 (2020), pp. 378–379.
- [13] Alex Farnsworth et al. “Past East Asian monsoon evolution controlled by paleogeography, not CO<sub>2</sub>”. In: *Science Advances* 5.10 (2019), eaax1697.
- [14] André Berger. “Milankovitch theory and climate”. In: *Reviews of geophysics* 26.4 (1988), pp. 624–657.
- [15] Ursula Röhl et al. “On the duration of the Paleocene-Eocene thermal maximum (PETM)”. In: *Geochemistry, Geophysics, Geosystems* 8.12 (2007).
- [16] Randy Showstack. “Carbon dioxide tops 400 ppm at Mauna Loa, Hawaii”. In: *Eos, Transactions American Geophysical Union* 94.21 (2013), pp. 192–192. DOI: <https://doi.org/10.1002/2013E0210004>.
- [17] Lorraine E Lisiecki and Maureen E Raymo. “A Pliocene-Pleistocene stack of 57 globally distributed benthic  $\delta^{18}\text{O}$  records”. In: *Paleoceanography* 20.1 (2005).
- [18] Timothy Naish et al. “Obliquity-paced Pliocene West Antarctic ice sheet oscillations”. In: *Nature* 458.7236 (2009), pp. 322–328.
- [19] Paul Voosen. “Ominous feedback loop may be accelerating methane emissions”. In: *Science* 377.6603 (2022), pp. 250–251.
- [20] Edward AG Schuur et al. “Climate change and the permafrost carbon feedback”. In: *Nature* 520.7546 (2015), pp. 171–179.
- [21] Horst WJ Rittel and Melvin M Webber. “Dilemmas in a general theory of planning”. In: *Policy sciences* 4.2 (1973), pp. 155–169.