Unsupervised clustering of oceanic Lagrangian particles: identification of the main pathways of the Labrador Current

Mathilde Jutras¹, Noémie Planat¹, Carolina Dufour¹, and Lauryn C Talbot¹

¹McGill University

July 23, 2023

Abstract

Modelled geospatial Lagrangian trajectories are widely used in Earth Science, including in oceanography, atmospheric science and marine biology. The typically large size of these dataset makes them arduous to analyze, and their underlying pathways challenging to identify. Here, we show that a Machine Learning unsupervised k-means++ clustering method can successfully identify the pathways of the Labrador Current from a large set of modelled Lagrangian trajectories. The presented method requires simple pre-processing of the data, including a Cartesian correction on longitudes and a PCA reduction. The clustering is performed in a kernalized space and uses a larger number of clusters than the number of expected pathways. During postprocessing, similar clusters are grouped into pathway categories by experts in the circulation of the region of interest. We find that the Labrador Current mainly follows a westward-flowing and an eastward retroflecting pathway (20% and 50% of the flow, respectively) that compensate each other through time in a see-saw behaviour. These pathways experience a strong variability of up to 96\%. We find that two thirds of the retroflection occurs at the tip of the Grand Banks, and one quarter at Flemish Cap. The westward pathway is mostly fed by the on-shelf branch of the Labrador Current, and the eastward pathway by the shelf-break branch. Pathways of secondary importance feed the Labrador Sea, the Gulf of St. Lawrence through the Belle Isle Strait, and the subtropics across the Gulf Stream.

































Unsupervised clustering of oceanic Lagrangian particles: identification of the main pathways of the Labrador Current

M. Jutras¹, N. Planat², C. O. Dufour², L. C. Talbot²

 $^{1}\mathrm{Department}$ of Earth Sciences, McGill University $^{2}\mathrm{Department}$ of Atmospheric and Oceanic Sciences, McGill University

Key Points:

1

2

3

4

5 6

7

8	•	Unsupervised clustering can identify the main pathways in geospatial Lagrangian
9		trajectories.
10	•	The clusters provide information on the properties and origin of the pathways.
11	•	The Labrador Current breaks in an east-west see-saw at the tip of the Grand Banks.

 $Corresponding \ author: \ Mathilde \ Jutras, \texttt{mathilde.jutras@mail.mcgill.ca}$

12 Abstract

Modelled geospatial Lagrangian trajectories are widely used in Earth Science, includ-13 ing in oceanography, atmospheric science and marine biology. The typically large size 14 of these dataset makes them arduous to analyze, and their underlying pathways chal-15 lenging to identify. Here, we show that a Machine Learning unsupervised k-means++ 16 clustering method can successfully identify the pathways of the Labrador Current from 17 a large set of modelled Lagrangian trajectories. The presented method requires simple 18 pre-processing of the data, including a Cartesian correction on longitudes and a PCA 19 reduction. The clustering is performed in a kernalized space and uses a larger number 20 of clusters than the number of expected pathways. During post-processing, similar clus-21 ters are grouped into pathway categories by experts in the circulation of the region of 22 interest. We find that the Labrador Current mainly follows a westward-flowing and an 23 eastward retroflecting pathway (20% and 50% of the flow, respectively) that compen-24 sate each other through time in a see-saw behaviour. These pathways experience a strong 25 variability of up to 96%. We find that two thirds of the retroflection occurs at the tip 26 of the Grand Banks, and one quarter at Flemish Cap. The westward pathway is mostly 27 fed by the on-shelf branch of the Labrador Current, and the eastward pathway by the 28 shelf-break branch. Pathways of secondary importance feed the Labrador Sea, the Gulf 29 of St. Lawrence through the Belle Isle Strait, and the subtropics across the Gulf Stream. 30

³¹ Plain language summary

Lagrangian trajectories, in which we follow a parcel of water or air parcel as it is 32 moved around by currents, are widely used in Earth Science, including in oceanography, 33 atmospheric science and marine biology. They typically come in very large and chaotic 34 dataset, from which it is difficult to identify the main pathways of a flow. Here, we use 35 a Machine Learning based algorithm, more specifically an unsupervised clustering algo-36 rithm, to identify the main pathways of the Labrador Current based on a large set of La-37 grangian trajectories obtained from an ocean model. This study shows the power of such 38 a method to help analyze this type of data, and provides a detailed description of the applied recipe so it can be used by people in the field. We find that, when it reached the 40 Grand Banks of Newfoundland, most of the Labrador Current flows either westward to-41 wards the Slope Sea or eastward towards the North Atlantic Ocean, in a see-saw behaviour. 42 We also identify a previously unknown minor pathway that brings Labrador Current wa-43 ters south of the Gulf Stream front. 44

45 **1** Introduction

Lagrangian trajectories are diagnostics that are widely used across climate sciences. 46 Such trajectories are obtained from the positioning of observational platforms such as 47 drifting floats in the ocean (e.g. Argo floats, surface drifters, RAFOS floats) and balloons 48 in the atmosphere, as well as from the advection of virtual particles derived from veloc-49 ity fields reconstructed from satellite altimetry or output from numerical simulations (among 50 others, A. F. Thompson & Sallée, 2012; van Sebille et al., 2018). Lagrangian trajecto-51 ries are used to study ocean and atmospheric circulations (e.g., Schulze Chretien & Frajka-52 Williams, 2018; Gillard et al., 2016; Bower et al., 2011; Fischer & Schott, 2002) and sea 53 ice drift (e.g., Williams et al., 2016; Brunette et al., 2019), to identify the origin and fate 54 of water masses (e.g., Kawasaki et al., 2022; Kelly et al., 2019), to assess connectivity 55 timescales (e.g., Jönsson & Watson, 2016), and to study the fate of atmospheric and oceanic 56 pollutants (e.g., Hertwig et al., 2015; Viikmäe et al., 2013), plastic (e.g., Lebreton et al., 57 2012), larvae (e.g., Avata et al., 2010; Cetina-Heredia et al., 2015; Phelps et al., 2015; 58 Simons et al., 2013), icebergs (e.g., Marson et al., 2018; Merino et al., 2016), and debris 59 or people during search and rescue (e.g., Hart-Davis & Backeberg, 2021). Yet, sets of 60 Lagrangian trajectories are challenging to analyze. It is often not possible to clearly dis-61

tinguish pathways given the chaotic nature of geophysical flows, which generally prevents 62 the use of simple and objective criteria to produce classification. In oceanography, tra-63 ditional classification methods of Lagrangian trajectories are based on counting parti-64 cles crossing sections based on hydrography (Jutras et al., 2023; Daher et al., 2020; Merino 65 et al., 2016; Bower et al., 2011), topography, or dynamic water properties (e.g. on fronts, 66 Roach & Speer, 2019; Schulze Chretien & Frajka-Williams, 2018). Apart from passages, 67 straits or other clearly defined topographic features that provide non-ambiguous phys-68 ical boundaries for the flow, criteria used for classification of trajectories often appear 69 adhoc or subjective (Fig. 1b). Besides, in modelling studies, dataset typically reach up 70 to millions of trajectories, making visual inspection overwhelming and non-efficient. 71

Machine Learning (ML) offers several algorithms that can help analyze (extremely) 72 large and complex Lagrangian datasets. Here, we consider clustering algorithms, which 73 automatically classify objects into "clusters", or groups of elements with similar prop-74 erties. Supervised clustering is trained on a pre-classified dataset, which can be obtained, 75 e.g., based on visual inspection. These types of methods are useful when the classifica-76 tion is already known or obvious to the human eye. On the other hand, *unsupervised* clus-77 tering lets the algorithm identify the clusters itself, removing potential biases in the choice 78 of classes. Unsupervised clustering has already successfully been applied to vertical pro-79 files from Conductivity-Temperature-Depth sensors (CTD; Boehme & Rosso, 2021) and 80 Argo floats (Houghton & Wilson, 2020; Rosso et al., 2020; Jones et al., 2019), to radar 81 data (Tiira & Moisseev, 2020), to cyclones tracks (Kremer et al., 2020) and to air pol-82 lutants (Brankov et al., 1998), as well as to identify mean flows (Koszalka & Lacasce, 83 2010), ocean fronts (Thomas et al., 2021), and finite-time (couple of days) coherent struc-84 tures in a flow (Filippi, Hadjighasem, et al., 2021; Wichmann et al., 2021; Schneide et 85 al., 2018), to name a few. The above-mentioned studies use various ML clustering meth-86 ods, including Gaussian-Mixture Models (Boehme & Rosso, 2021; Rosso et al., 2020; Jones 87 et al., 2019; Thomas et al., 2021), k-means (Houghton & Wilson, 2020; Kremer et al., 88 2020; Schneide et al., 2018; Koszalka & Lacasce, 2010), optimized-parameter spectral meth-89 ods based on k-means (Filippi, Hadjighasem, et al., 2021; Filippi, Rypina, et al., 2021), 90 hierarchical clustering, and density-oriented clustering like DBSCAN (Wichmann et al., 91 2021). Yet, to our knowledge, no study has applied unsupervised clustering to large-scale 92 (more than a couple of days) geophysical Lagrangian trajectories, nor used such a method 93 to identify the main pathways of a geophysical flow. This technique appears especially 94 suitable to identify and characterize the pathways of an ocean current, removing the sub-95 jectivity inherent to more traditional methods mentioned earlier. 96

In this study, we use such a method to study the Labrador Current (Fig. 1a). The 97 Labrador Current is a western boundary current. It forms the western limb of the sub-98 polar gyre, and as such is a critical component of the North Atlantic circulation. The 99 Labrador Current is composed of an inshore and a shelf-break branch that flow south 100 on and along the Labrador shelf, respectively (Florindo-López et al., 2020; Loder et al., 101 1998; Lazier & Wright, 1993), until the tip of the Grand Banks. Eventually, most of the 102 Labrador Current Water is entrained into the subpolar North Atlantic by the North At-103 lantic Current (NAC) and the remainder follows the continental shelf southwestward (Fig. 1a; 104 Townsend et al., 2015; Fratantoni & McCartney, 2010; Pérez-Brunius et al., 2004). By 105 doing so, the Labrador Current carries cold, relatively fresh and well-oxygenated waters 106 from the subarctic to both the subpolar North Atlantic and to the Slope Sea and east-107 ern American continental shelf. Variability in the strength and exact path of the Labrador 108 Current therefore affects the water properties in both regions (e.g., Jutras et al., 2023; 109 Gonçalves Neto et al., 2021; Chen et al., 2020; Holliday et al., 2020; Claret et al., 2018; 110 B. D. Petrie & Drinkwater, 1993) and in connected bodies of water such as the Gulf of 111 St. Lawrence Estuary (Jutras et al., 2020; Gilbert et al., 2005; Han et al., 1999) and the 112 Gulf of Maine (Whitney et al., 2022; Pershing et al., 2016), with direct consequences on 113 marine ecosystems (Poitevin et al., 2019; Chabot & Dutil, 1999) and fisheries (Pershing 114 et al., 2016; Mills et al., 2013). Some Labrador Current Waters also leak into the Labrador 115



Figure 1. (a) Region of the Labrador Current. The arrows show the approximate location of the main currents of the region. The main topographic and oceanographic features are labelled. FC indicates Flemish Cap. NAC indicates the North Atlantic Current. (b) Example subset of Lagrangian trajectories launched in the Labrador Current. The colour indicates the distance from the shore at initialization. The thick black line indicates the launch section. The blue line indicates the entry point to the Labrador Sea, used in section 3.2.1. (c) Probability density plot of the complete dataset of Lagrangian trajectories. We stop tracking particles east of 50E.

Sea (Schulze Chretien & Frajka-Williams, 2018; Howatt et al., 2018; Palter et al., 2008;
Myers, 2005), possibly affecting stratification and modulating deep water formation and
the Atlantic Meridional Overturning Circulation (AMOC; New et al., 2021). Still, little is known about the spatio-temporal characteristics, magnitude and drivers of these
pathways, or about other possible pathways (Jutras et al., 2023; Fratantoni & McCartney, 2010).

This paper uses the Labrador Current as a case study to demonstrate that unsu-122 pervised clustering can be used to identify pathways in geophysical Lagrangian tracks. 123 To do so, we implement an unsupervised kmeans++ clustering method on a large set 124 of Lagrangian trajectories within the Labrador Current (Fig. 1b), to identify and char-125 acterize the main and secondary pathways of this current. The trajectories are almost 126 127 impossible to distinguish through traditional methods, being continuously distributed in the western North Atlantic (Fig. 1b,c). Section 2 presents a step-by-step description 128 of our method intended for non-experts, hoping that this case study can inspire appli-129 cations in other Earth system contexts. Section 3.1 presents the results of the cluster-130 ing, including the identification of the pathways, a characterization of their properties, 131 and quantitative comparisons with the literature. Section 3.2 uses the classification of 132 the complete dataset to look at the variability of each pathway of the Labrador Current 133 on seasonal and decadal time scales. Section 4 offers some concluding remarks on the 134 method and the results. 135

136 2 Methods

137

2.1 Lagrangian trajectories

The Lagrangian trajectories are generated from virtual particles advected offline by 3D velocity from the GLORYS12V1 ocean reanalysis (Lellouche et al., 2018). GLO-RYS12V1 is based on the NEMO3.1 modelling platform (Madec et al., 2019). It has a spatial resolution of 1/12° on an ORCA grid and 50 levels in the vertical, with thicknesses ranging from 0.5 m at the surface to 160 m at a 1 km depth and with 18 levels ¹⁴³ in the top 50 m. The simulation covers the 1993 to 2018 period and is forced with the ¹⁴⁴ ERA-Interim atmospheric reanalysis (ECMWF Re-Analysis, Dee et al., 2011).

The virtual particles are tracked with the OceanParcels tool for Python (Probably 145 A Really Computationally Efficient Lagrangian Simulator; Delandmeter & Van Sebille, 146 2019). We use the daily horizontal velocity outputs on a longitude-latitude grid provided 147 on the Copernicus Marine Service (CMS) website. The vertical velocities are reconstructed 148 from sea surface height. Particles are seeded every $1/12^{\circ}$ along the $(53^{\circ}N, 56.7^{\circ}W) - (54.3^{\circ}N, 52.0^{\circ}W)$ 149 line (Fig. 1b) and every 10 m in the vertical, in waters with a salinity lower than 34.8, 150 for a total of 966 particles per seeding event. The salinity cut-off is used to delineate the 151 Labrador Current from the Labrador Sea (Myers, P., personal communication; Loder et 152 al., 1998). Particles are released every week from January 1^{st} 1993 to January 1^{st} 2015 153 and are tracked with a 10-minute time step. The complete data set contains 1.2 millions 154 trajectories. The seeding temporal and spatial frequencies are chosen so that increas-155 ing the number of particles does not change their general distribution downstream, op-156 timizing the use of computational resources (van Sebille et al., 2018). We stop tracking 157 the particles when they hit topography or the boundaries of the domain (Fig. 1c) or af-158 ter 550 days, time after which they have left the Labrador Shelf and reached their final 159 export zone. These experiments are also described in Jutras et al. (2023), who look at 160 the variability in the retroflection of the Labrador Current. In addition to the position 161 and depth of the particles, we track their temperature, salinity and age since release. We 162 use a purely advective scheme. Tamsitt et al. (2017) showed that the addition of tur-163 bulent diffusion did non affect Lagrangian trajectories significantly in eddy-resolving mod-164 els. In addition, there is no consensus on a realistic value for diffusive coefficients, espe-165 cially when covering both coastal and open ocean areas (van Sebille et al., 2018). 166

167

2.2 Observational dataset

We compare the trajectories of the virtual particles with those from actual obser-168 vational platforms, namely surface drifters, Argo floats and RAFOS/SOFAR floats. A 169 direct comparison is not possible because virtual particles can move vertically, while floats 170 and drifters flow at a fixed depth. In addition, most of the Argo and RAFOS/SOFAR 171 floats drift deeper than the virtual particles, more specifically into the Deep Western Bound-172 ary Current. We therefore expect the trajectories to differ, in particular where the Labrador 173 Current waters dive as they interact with the Gulf Stream – NAC front. Still, we use the 174 observations to validate qualitatively the simulated pathways, as well as to offer a rough 175 comparison of the magnitude of each pathway. 176

We use surface drifters deployed as part of the Global Drifter Program. These satellitetracked buoys drift at the surface of the ocean and are equipped with 15 m or 1 m drogues. We consider the floats that are carried by the Labrador Current by selecting the ones that cross the virtual particles seeding line and that enter the Grand Banks area, as defined by the (55°W; 41°W) – (45°N; 50°N) box (Fig. 6). Based on this criterion, we identify 79 drifters from 2000 to 2018.

Argo floats are autonomous profilers that drift passively with ocean currents at a 183 parking depth (typically 1 km) and profile temperature, salinity and pressure down to 184 approximately 2 km every 10 days. RAFOS/SOFAR floats are autonomous platforms 185 that drift at a fixed depth between 500 m and 1 km. We select the floats based on the 186 187 same criteria as for the surface drifters, except that we extend the seeding line and the box offshore by two degrees (Fig. 6) to account for the fact that floats drift deeper over 188 the continental slope. We identify 64 Argo floats fitting these criteria between 2001 and 189 2019 and 50 RAFOS/SOFAR floats between 2003 and 2007. 190

A visual inspection suggests that the pathways of observational platforms and of virtual particles generally agree (Fig. 1b and 6). The small number of drifters and floats rules out applying a clustering algorithm to their trajectories. Hence, we manually classify the platforms into pathways using the following hydrographic sections (Fig. 6):

- Westward-flowing: crosses the 54^{th} meridian south of the Grand Banks;
- Westward then retroflected: crosses the 54th meridian south of the Grand Banks and eventually drifts eastward;
 - Retroflecting: enters the zone from 0°W to 60°W and from 47°N to 65°N;
 - Southward-flowing: enters the zone from 54°W to 35°W and from 35°N to 47°N.
- 200 2.3 Clustering algorithm, step by step
- 201 **2.3.1 Overview**

195

198

199

Machine Learning unsupervised clustering algorithms build a classification model 202 that attributes each object (here, trajectories) to a cluster. The model is characterized 203 by parameters called *hyperparameters* that can include, for instance, the number of trans-204 formations applied to the data, the number of clusters, or criteria on the within-cluster 205 maximal distance. Three independent data subsets are used to feed the model, namely the training, validation and test sets. These sets must be large (at least hundreds of ob-207 jects) and of high quality (e.g. evenly sampled or without missing values). The train-208 ing set is used to train the model, which is validated with the validation set for a range 209 of hyperparameter values. By comparing the results with performance metrics, the most 210 performant hyperparameters values are determined. Once the model is ready, its per-211 formance is validated with the test set. To avoid overfitting the model to the subsets, 212 the test set must be used only once, to validate the final results. Overfitting would lead 213 to a model that offers a good classification of the training subset, but not of new data. 214 Finally, once the model is ready, it can be applied to the complete dataset or to new dataset. 215 An overview of the method is presented in Figure 2. 216

2.3.2 Pre-processing

Before building this model, we need to prepare the data. Since the goal of the study 218 is to identify the various pathways of the Labrador Current as it flows over and along 219 the Labrador Shelf, we are interested in the shape of the Lagrangian trajectories. We 220 therefore base our classification on latitude and longitude coordinates. Additional vari-221 ables (temperature, salinity and depth) were also considered to be used in the clustering algorithm, but showed no significant improvement on the classification results. We build the clustering model with a subset of 100 000 trajectories out of a total 1.2 mil-224 lions (Fig. 2). These trajectories are selected randomly every four years, as preliminary 225 analyses showed no periodicity in the preferred pathways over that timescale. This sub-226 set is further separated into an 80 000 particle training set, a 10 000 particle validation 227 set, and a 10 000 particle test set. While there is no universal rule on the number of ob-228 jects required in each set, an 0.8-0.1-0.1 ratio is commonly used. 229

- ²³⁰ We apply the following pre-processing to each set (Fig. 2):
- 231 232 233

234

235

217

- To avoid a bias by which the particles would be clustered based on their initialization location, we translate all the particles to the same starting point. This *translation* step increases the efficiency of the clustering (not shown).
- Trajectories shorter than 550 days e.g. due to the particles reaching the bottom of the ocean, the shore, or the boundaries of the domain are filled with zeros.
- To account for the sphericity of the Earth, we apply a longitudinal correction. The 236 particles flow approximately from 54°N, where one degree of latitude represents 237 65 km, to 30°N , where one degree of latitude represents 96 km. Because we are 238 interested in the shape of the trajectory in a Cartesian space (km) but operate 239 the classification in latitude-longitude space, we apply a " $\cos \lambda$ " weight to the lon-240 gitudes, where λ represents the latitude. The resulting Euclidian distance in mod-241 ified latitude-longitude space offers a good approximation of the real (physical) 242 Cartesian distance at the surface of the ocean. 243



Figure 2. Schematic diagram of the unsupervised clustering method. A detailed description of the method is provided in Section 2.3. The 30 clusters are presented in Fig. 4 and the seven pathway categories are presented in Fig. 5.

• To lower the computational cost of the analysis (the training set has a size of 80 000 244 \times 550 locations \times 2 coordinate variables), it is common practice to reduce the num-245 ber of features by implementing a Principal Component Analysis (PCA). By keep-246 ing the features responsible for 99.99% of the total variance, we reduce the dataset 247 to 12% of its original size. We implement the PCA and the k-mean clustering (see 248 Section 2.3.3) in a kernalized space, i.e. a transformed variable space. A kernel 249 helps convergence when using linear classifiers on non-linear data, by implicitly 250 adding non-linearities to the algorithm (Hofmann et al., 2008). Among the tested 251 kernels (sigmoid, polynomial, cosinus), the cosinus transformation, defined as fol-252 lows, led to the most efficient clustering: 253

$$k(x,y) = \frac{xy^{T}}{||x|| \cdot ||y||}$$
(1)

where x and y are the vectors containing the variables (here, the coordinates of the trajectories).

Computing the kernalized k-means and kernalized PCA requires high RAM, as large matrices need to be temporarily loaded. In our case, the computation takes about one day
on a HPC system with 186 GB of RAM. In a non-kernalized space, the clustering algorithm could run on a regular work station.

260 2.3.3 Processing

We here apply a k-means++ clustering algorithm (Fig. 2), which is common, easy 261 to implement, and requires only one hyperparameter: the number of clusters. The k-means 262 method classifies the data by minimizing the within-cluster variance of the Euclidean dis-263 tance between each object. More specifically, each cluster is characterized by a centroid. 264 or mean vector, to which the distance with each object belonging to that cluster is min-265 imized. In the k-means++, the spread between the initial centroids is maximized by test-266 ing multiple initializations and keeping the one offering the best classification, signifi-267 cantly improving the convergence and speed compared to the traditional k-means method. 268 We here implement 20 random initializations. To accelerate the convergence of the clas-269 sification itself, we then fold it 15 times: we randomly split the dataset in 15 pieces, it-270 eratively apply the classification to 14 pieces and evaluate the results on the 15th. The 271 results are not sensitive to a higher number of folds or initializations. These steps are 272 implemented using the k-means++ functions of the Python scikit-learn package (scikit 273 -learn.org/). 274

For the value of the hyperparameter, namely the number of clusters, prior knowl-275 edge of the circulation of the Labrador Current suggests two major pathways plus some 276 minor ones (see Section 1). The k-means method has difficulty converging in the pres-277 ence of clusters of unequal sizes (i.e. containing unequal number of objects). Using a large 278 number of clusters and grouping them afterwards has been shown to improve the per-279 formance of the classification (Echols et al., 2020) and helps reveal secondary pathways. 280 To find the optimal number of clusters, we use two performance metrics: the silhouette 281 score (Rousseeuw, 1987), and a physics-based metrics that is adapted to our scientific question. The silhouette score measures the overall performance of the clustering algo-283 rithm based on the intra and inter cluster distances. This metric is expected to monotonously 284 decrease with the number of clusters, since a higher number of clusters necessarily im-285 proves the performance on average (i.e. the intra cluster spread decreases as the num-286 ber of objects per clusters decreases). The number of clusters can therefore be chosen 287 based on a stabilization of this score (Fig. 3a). We then define a physics-based metric 288 that evaluates the spatial and temporal coherence of the particles. We first define five 289 regions that the particles are likely to visit (Fig. 3b). For each cluster, we identify the 290 most popular region at each time step, and compute the fraction of particles found in 291 that region. This provides a score for each cluster, between 0 and 1. We then average 292



Figure 3. (a) Algorithm's performances for a varying number of clusters, based on silhouette and a physics-based scores. (b) Regions used for the physics-based performance metric.

the scores of all clusters to obtain a global score. The model's performance is highest when this metric is maximized, indicating that a high number of particles simultaneously visit the same region. Both the numerical and physics base metrics show a plateau around 30 clusters (Fig. 3a).

297 2.3.4 Post-processing

As expected, a visual inspection of the obtained clusters reveals some redundan-298 cies in the pathways represented in some clusters (Fig. 4). While, as we will show in Sec-299 tion 3.2.3, most of the clusters which look alike actually present differences that are not 300 visible in the particle trajectories, it is useful to group the clusters identifying similar path-301 ways. Based on the shape of the trajectories and on their export location, we visually 302 identified six pathway categories which are described in detail in Section 3.1.1. The iden-303 tification of these categories was nourished by prior knowledge of the circulation discussed 304 in the literature (Section 3.1.3). To avoid biases in the categorization, we invited eight 305 experts of the northwestern Atlantic circulation to sort the 30 clusters into the six iden-306 tified pathway categories or to new ones they would discern. The experts overall agree 307 on the classification (see Appendix B for details). 308

309 **3 Results**

310

3.1 Pathways

The unsupervised clustering method successfully classifies the trajectories into 30 clusters showing similar trajectories (Fig. 4), which were combined into six pathway categories (Table 1 and Fig. 5). Note that throughout this section, we display results from the test set. We first describe how we identified the pathway categories (section 3.1.1) before assessing the qualitative agreement with the scarce observations (section 3.1.2) and with the literature (section 3.1.3 and 3.1.4).

317 3.1.1 Definition of the pathway categories

From the 30 clusters, we identify two main pathway categories: particles retroflected towards the subpolar North Atlantic (henceforth referred to as *retroflected*), and particles following the shelf westward into the Slope Sea and along the eastern American continental shelf (henceforth referred to as *westward-flowing*, Fig. 4 and 5). These pathways account in total for respectively 48% and 21% of all the trajectories (Table 1), mean-

ing that close to 70% of the water from the Labrador Current feeds either the subpo-323 lar North Atlantic or the Slope Sea and eastern American continental shelf. Less than 324 1% of the particles first enter the Slope Sea before retroflecting towards the subpolar North 325 Atlantic (cluster #22 on Fig. 4). We include these particles in the westward-flowing cat-326 egory, as they first affect the water properties of the Slope Sea and have lost most of their 327 Labrador Current water signature once they retroflect. Another significant pathway cat-328 egory comprises the particles that are killed on the Labrador Shelf as they hit the seafloor 329 (22% of the trajectories; referred to as Labrador Shelf). This category does not repre-330 sent a real pathway, and is rather an artifact of the virtual Lagrangian tracking. We also 331 identify three secondary pathway categories: the particles that travel southward from 332 the tip of the Grand Banks (8%; referred to as *southward-flowing*), the ones that enter 333 the Gulf of St. Lawrence through the Belle Isle Strait (1%; referred to as *Belle Isle*), and 334 the ones that feed the Labrador Sea (<1%; referred to as Labrador Sea, Fig. 4 and 5). 335

The uncertainty on this classification comes from two sources: the clustering al-336 gorithm itself (algorithm uncertainty), and the categorization of clusters (human-induced 337 uncertainty). The two are not independent, since a large algorithm error will lead to dis-338 agreement in the experts' classification. First, the algorithm error manifests as trajec-339 tories that are classified into a cluster even if, from a visual inspection, they would have 340 fitted better in another. For instance, cluster #17 belongs to the Labrador Shelf path-341 way category, but a few particles still reach the Scotian Shelf, and should have been clas-342 sified in a cluster belonging to the westward-flowing pathway category. There currently 343 exists no widely accepted method to evaluate the error from unsupervised clustering al-344 gorithms (e.g., Abdar et al., 2021; Kläs & Vollmer, 2018). We cannot use the within-cluster 345 spread to assess the algorithm's error, because particles can end up quite far from each 346 other but still belong to the same cluster (e.g., particles retroflecting eastward can reach 347 from 30N to 55N). Hence, we simply report the algorithm's performance based on the 348 physical metric presented in Section 2.3. We find that the score is high for all (> 0.7)349 but some Labrador Shelf clusters, in which a few particles enter the Belle Isle Strait, and 350 for the Belle Isle cluster, in which a few particles flow along the Scotian Shelf (Fig. A3). 351

Second, for the errors in the categorization of clusters, we find that the experts are almost unanimous in classifying the clusters in the Belle Isle, southward-flowing and Labrador Sea pathway categories. For other categories, the error ranges between 7 and 10% (Table 1, see also Appendix B and Table B1). Overall, the errors appear sufficiently small to go forward with the analysis of the results.

357

3.1.2 Comparison against trajectories of observational platforms

We perform a visual comparison between the obtained pathways and that of Argo 358 floats, RAFOS/SOFAR floats, and surface drifters (see Section 2.2). We find that the 359 retroflected, westward-flowing and southward-flowing pathways clearly appear in the tra-360 jectories of autonomous platforms (Fig. 6). There is also a significant amount of plat-361 forms going westward and then retroflecting, more than in the virtual particles. We do 362 not expect any observational platforms to follow the Labrador Sea pathway because our 363 selection criteria filter out these platforms (section 2.2). The same holds for the Labrador 364 Shelf pathway, which is an artifact of the virtual Lagrangian tracking, and for the Belle 365 Isle pathway, since no autonomous platforms have been launched within the coastal cur-366 rent that feeds this strait. The agreement in the pathways provides confidence in the clus-367 tering. The bulk sizes of the observed and modelled pathway categories generally agree 368 (Table 1), keeping in mind that (i) there are too few observational platforms to allow 369 a statistically robust comparison and (ii) observational platforms drift at a fixed depth 370 while Lagrangian particles can move vertically (see Section 2.2), and (iii) the on-shelf 371 category, composing >20% of the virtual particles, is an artifact of the Lagrangian track-372 ing and is absent from the observations. Note that most of the surface drifters retroflect 373 eastward (Fig. 6). 374



Figure 4. Density map of the trajectories for each of the 30 clusters identified by the kmeans++ model for the test set (see Section 2.3.3). The numbers on the top right indicate the number of particles (or trajectories) in each cluster, while the numbers on the bottom left correspond to the cluster identification number. The dark grey line shows the 350 m isobath. The coloured frames indicate in which pathway category the cluster is classified by the experts: retroflected (red); westward-flowing (green); Labrador Shelf (purple); Labrador Sea (blue); southward-flowing (pink); Belle Isle (cyan). The westward-flowing cluster with a dashed contour contains particles that go westward first and are then retroflected. See Section 3.1.1 for a description of each pathway category.

Pathway cate- gory	Cluster ID	Percentage: mean (min- max)	Exp- ert's error	% of ob- serva- tions
Retroflected	$\begin{vmatrix} 2, 4, 7^*, 8^*, 9, \\ 10, \\ 11^*, 19^*, 20, 24, \\ 28^*, 30 \end{vmatrix}$	47.6 % (24.9-73.7)	10 %	74 %
Westward- flowing Westward then retroflected	3, 18, 22, 23, 26 22	21.0 % (5.4-42.2) 0.6 % (0.0-2.2)	6 % 7 %	10 % 4 %
Labrador Sea	27	$ \begin{array}{ c c c c c c c c c c c c c c c c c c c$	0 %	-
Labrador Shelf	$\left \begin{array}{c}1,5,6,13,14,\\15,16,17,21\end{array}\right $	$ \begin{vmatrix} 21.8 \% \\ (12.7-31.3) \end{vmatrix} $	9 %	-
Southward- flowing	12, 29	$\begin{array}{ c c c c c c c c c c c c c c c c c c c$	0 %	16 %
Belle Isle	25	$ \begin{vmatrix} 1.4 \% \\ (0.0-6.5) \end{vmatrix} $	0 %	0 %

Table 1. Classification of the 30 clusters into the six pathway categories (see Section 3.1.1 for a description of each pathway category). The first column indicates the name of the pathway category; the second column indicates the identity number (ID) of the clusters classified within that category (see Fig. 4 for the IDs); the third column indicates the mean percentage of trajectories classified into a given category, computed from the complete dataset, as well as the lowest and highest percentage over the 1993 to 2018 period; the fourth column indicates the error coming from the disagreement in the experts' categorization; the last column indicates the percentage of observational platforms corresponding to each category (see Section 2.2). In the retroflected category, the clusters marked with an asterisk retroflect at Flemish Cap while the others retroflect at the tip of the Grand Banks.



Figure 5. Density maps for each of the six pathway categories of the Labrador Current (see Section 2.3.4). See Figure 4 for a detailed description of the plot. The percentage provides the average magnitude of each pathway category.

3.1.3 Validation of pathway categories against the literature

375

We compare the relative importance of each pathway category with results from 376 previous studies. First, many studies also report that the retroflected and westward-flowing 377 pathways are, respectively, the main and secondary pathways for the Labrador Current 378 (e.g., Gonçalves Neto et al., 2023; Holliday et al., 2020; Han et al., 2014; Fratantoni & 379 McCartney, 2010). Fox et al. (2022) also observed that modelled Lagrangian trajecto-380 ries can retroflect after having flowed westward. Our estimate of the Labrador Current 381 export towards the Labrador Sea (0 - 3%, Table 1) is in good agreement with observation-382 based studies (0 - 3%; Howatt et al., 2018; Schmidt & Send, 2007) and with model-based 383 studies, (6 - 8%; Myers, 2005). We expect an underestimation, because the above-mentioned 384 studies focus on the shelf-break branch of the Labrador Current, while we also consider 385 the inshore, on-shelf branch of the current ($\sim 15\%$ of the volume transport). The inflow of water through the Belle Isle strait has been estimated to range from 0.1 Sy in the spring 387 to 0.4 Sv in the winter, based on observations (Shaw & Galbraith, 2023; B. Petrie et al., 388 1988), and from 0.15 Sv to 1 Sv during winter storms, based on a model (Saucier et al., 380 2003). Relative to the mean 8.1 Sv Labrador Current volume transport found in GLO-390 RYS12V1, this represents 1 - 12% of the current, in broad agreement with the results 391 of the clustering (0 - 7%). The southward-flowing pathway has not been explicitly de-392 scribed as a Labrador Current pathway in the literature. However, such a pathway has 303 been described for the Deep Western Boundary Current (DWBC), which exports Labrador 394 Sea Waters equatorward below ~ 1500 m (Bower et al., 2009). The virtual particles that 395 follow the southward-flowing pathway first sink to a depth of ~ 1000 m (see Fig. 11), thus 396 reaching the upper limb of the DWBC (Handmann et al., 2018). This pathway could there-397 fore emerge from interactions between the two currents. Overall, the relative importance 398 of each pathway obtained from the clustering agrees well with previously model-based 399 and observation-based estimates, further supporting the method. 400



Figure 6. Observational platforms drifting within the Labrador Current between year 2000 and 2018 – a time period that overlaps with the GLORYS12V1 reanalysis period – sorted into four of the pathway categories identified from the clustering algorithm (see Section 2.3.4). The name of the pathway category is indicated at the bottom right, along with the percentage of platforms classified in this category. The colours of the trajectories refers to the drifting depth of the platform. For each panel, a pie chart represents the fraction of platforms in depth classes, with the corresponding number of platforms indicated within each class. The grey contour delineates the 350 m isobath. The black straight line and the yellow boxes indicate the criteria used to select the platforms of interest (see section 2.2 for further details).



Figure 7. (a-f) Potential temperature – salinity diagrams of all the virtual particles classified according to pathway categories. The color shading represents the density of particles for a given T-S combination.(g) Potential temperature – salinity diagram of the two main water masses interacting in the region of interest: the North Atlantic Central Waters (NACW) originating from the Gulf Stream, and the Labrador Current Waters (LCW) formed by the Labrador Current flowing southward on the Labrador Shelf. Data for these signatures come from the World Ocean Circulation Experiment (WOCE) climatology (Gouretski, 2018). The LCW is defined as the waters lying between 150 m and the seafloor on the Labrador Shelf and slope, and the NACW as the waters lying between 250 and 1300 m within the Gulf Stream jet (see Jutras et al., 2020). The dashed grey lines show isopycnals (+1000 kg m⁻³).

3.1.4 Thermohaline signature across pathways

We further verify the ability of the clustering algorithm to properly classify the tra-402 jectories by comparing the thermohaline properties of each category with what is expected 403 for these pathways. The Labrador Sea category clearly and almost exclusively shows the signature of Labrador Current Waters (Fig. 7), which makes sense since these waters leave 405 the continental shelf before any contamination can occur (see section 3.2.2). All the other 406 categories show the signature of the LCW getting fresher as they receive river outflow 407 along the Labrador coast. The Belle Isle category contains only the LCW. In addition to the signature of the LCW, the retroflected, westward-flowing and southward-flowing 409 categories show that of the warm and salty North Atlantic Central Waters (NACW), which 410 progressively mix with the LCW along the Labrador Current – NAC front. In the westward-411 flowing category, we only find the signature of the coldest, freshest NACW, as the con-412 tact time with the NAC is shorter than for the retroflected category. Most of the path-413 ways show the additional weak signature (few trajectories) of warm and fresh waters formed 414 on the Labrador Shelf during the summer. The fact that each pathway category has a 415 thermohaline signature that fits with what can be expected from the circulation supports 416 the algorithm and our choice of categories. 417

418

401

3.2 Spatio-temporal characteristics of the pathway categories

⁴¹⁹ Now that the different pathways of the Labrador Current are identified and val-⁴²⁰ idated, we document their temporal evolution (section 3.2.1) and spatial characteristics (sections 3.2.2, 3.2.3 and 3.2.4). To do so, we use the classification of the complete dataset (1.2 millions trajectories) from 1993 to 2018.

423

3.2.1 Temporal variability in the pathways

The classification provides time series of the relative importance of each identified pathway (Fig. 8). The relative importance of the westward-flowing pathway from 1996 to 1998 fits with a salinification of the subpolar North Atlantic (Holliday et al., 2020), while that of the retroflected pathway since 2011 coincides with a strong freshening of the subpolar North Atlantic (Holliday et al., 2020), a warming of the eastern American continental shelf (Chen et al., 2020), and a deoxygenation of the western North Atlantic and adjacent basins (Jutras et al., 2020; Claret et al., 2018).

We find that the two main pathways display a strong interannual and seasonal variability (Fig. 8). Their magnitude varies largely: between 24% and 73% of the particles are retroflected and between 4% and 42% flow westward, respectively a 77% and 96% variability (variance/mean × 100). The retroflected pathway is always dominant, while the westward-flowing pathway can be almost shut down when the retroflection is strong. The southward-flowing pathway is the most stable pathway, with a variability of 13%.

The retroflected and westward-flowing pathway categories are strongly anti-correlated 437 (correlation coefficient (c.c.) = -0.97, p-value < 0.001, top panel of Fig. 8a). This agrees 438 well with a see-saw behaviour of the Labrador Current at the tip of the Grand Banks 439 (Jutras et al., 2023; Han et al., 2019): when the retroflection towards the subpolar North 440 Atlantic is strong, little water reaches the Slope Sea or the continental shelf break, and 441 442 vice-versa. The Labrador Sea category is the opposite, being anti-correlated with the westwardflowing one (c.c. = -0.64, p < 0.001) and hence correlated with the retroflected one (c.c. 443 = 0.55, p < 0.001). These correlations support the idea that the westward-flowing path-444 way is mainly associated with similar branches of the Labrador Current, namely the in-445 shore branch, while the Labrador Sea and retroflected pathways are associated with the 446 offshore branch of the current (see Section 3.2.2). The Belle Isle and southward-flowing 447 pathway categories show no correlation with other pathways, suggesting that they are 448 forced by different mechanisms. 449

There is a significant seasonal cycle in the time at which the particles veer west-450 ward or are retroflected at the tip of the Grand Banks (Fig. 8b). The retroflection is strongest 451 in late summer (Aug.-Sep.) and generally weakest in the winter (Jan.), although it is strong 452 in some winters, while the westward-flowing pathway is greatest in the winter (Feb.) and 453 weakest in the summer (Jun.-Aug.). The opposite seasonal cycles between the two main 454 pathways suggest that the see-saw behaviour also occurs at a seasonal scale. Seasonal 455 variations in the circulation patterns near the Grand Banks are discussed in the liter-456 ature, and are suggested to be driven by seasonal variations in the water temperature 457 and salinity (advection of meltwater; Fratantoni & McCartney, 2010; Lazier & Wright, 458 1993) affecting stratification (Fratantoni & McCartney, 2010), in the density gradients 459 across the shelf-break (Schneider et al., 2015), in the winds (Holliday et al., 2020; Han, 460 2005; K. R. Thompson et al., 1986), and to a southern drift of the Gulf Stream in the 461 summer (Seidov et al., 2021). 462

In addition to the two main pathway categories, there is a marked seasonality in the trajectory of the particles leaving the Labrador Shelf towards the Labrador Sea (crossing the blue line on Fig. 1b). More particles do so in the early summer (Jun.-Jul.) compared to other seasons (not shown). This behavior agrees with the observations of Howatt et al. (2018), who suggest that northward winds, which are only present in the summer, drive an offshore Ekman transport that supports the export of freshwater to the Labrador Sea.



Figure 8. Temporal variability of the pathway categories. (a) Time series of the percentage of the total number of particles for each of the six pathway categories, with the time recorded at the seeding time. The time series are smoothed over one year and are presented in three different panels for readability. Note the differences between y-axes. (b) Seasonal cycle of the retroflected and westward-flowing pathways, for the time at which the particles reach the tip of the Grand Banks (i.e. when they cross the 49°N line, see Fig. 1a). This way of recording the time gives a better estimate of the local seasonal export variability, given the wide range of propagation times for the particles (Fig. 10; Fox et al., 2022). The amplitude is normalized. The shaded area shows the interannual spread in the seasonal cycle, computed from the squared sum of each year difference.

3.2.2 Spatial characteristics of the pathways

We can use the characteristics of the different pathway categories to deduce infor-471 mation about their origin. In the retroflected pathway category, the virtual particles have 472 slightly higher velocities on the Labrador Shelf compared to the other pathways (Fig. 9iii). 473 and most (though not all) particles originate from and flow within the offshore portion 474 of the shelf (Fig. 9i and ii). The retroflected pathway thus seems to be mostly fed by the 475 offshore, or shelf-break, faster branch of the Labrador Current. In contrast, the virtual 476 particles in the westward-flowing and in the Labrador Shelf pathway categories originate 477 equally from across the seeding line (Fig. 9iv), and then most converge towards the in-478 shore section of the shelf (Fig. 9v). Still, more than 5% of particles drifting within the 479 shelf-break branch of the Labrador Current join the westward-flowing pathway (Fig. 9v). 480 The westward-flowing and Labrador Shelf pathways are also associated with generally 481 slower velocities (Fig. 9vi and xii). The Labrador Shelf category overall seems to be fed 482 predominantly by the offshore branch of the Labrador Current (Fig. 9xi). The particles 483 that end up in the Labrador Sea originate from close to the shelf-break, and are carried 484 by the offshore-most and fastest portion of the Labrador Current (Fig. 9viii and ix). In 485 contrast, the particles entering the Belle Isle Strait are very slow (Fig. 9xviii). They can 486 originate from any location across the shelf, although not from the offshore-most por-487 tion of the current (Fig. 9xvi), and more specifically from shallower depths (< 50 m) 488 than the other particles (Supplementary fig. A1). The depth distribution of the particles at initialization does not play a role for the other pathways. Finally, similarly to the 490 westward-flowing category, the particles associated with the southward-flowing pathway 491 do not appear to have a preferred origin and travel across the whole shelf (Fig. 9xiii and 492 xiv). They also show a wider range of velocities than the other pathway categories (Fig. 9xv). 493 This suggests that the westward-flowing and southward-flowing categories are not fed 494 by a particular branch of the Labrador Current. We find that the particles in the southward-495 flowing category show turbulent motion soon after they leave the Grand Banks (Fig. 1b) 496 and 6). Since that region is located in the transition zone between the Gulf Stream and 497 the more stable NAC, we suggest that the southward-flowing pathway emerges as par-498 ticles get caught in small-scale features such as eddies, common in that region (Rossby, 499 1999; Brooks, 1987), explaining why these particles do not follow the average circulation 500 of the Gulf Stream/NAC, directed northeastward (Bower et al., 2011). 501

502

470

3.2.3 Specific circulation patterns

Grouping the different clusters into pathway categories is useful to concentrate on 503 the general properties of the pathways of the Labrador Current. Yet, within a pathway 504 category, individual clusters often show distinct characteristics. These characteristics re-505 veal important details of the circulation that can refine our view of the Labrador Cur-506 rent pathways. For instance, the particles in different retroflecting clusters veer at dif-507 ferent locations. About one third of the retroflecting particles do so near Flemish Cap, 508 and the remainder at the tip of the Grand Banks (Table 1 and Fig. 4). These propor-509 tions do not vary significantly with time. We notice that depending on where the par-510 ticles retroflect, they reach slightly different regions of the North Atlantic. The parti-511 cles that retroflect at Flemish Cap feed the north of the subpolar gyre (\sim 52–57°N) and 512 the particles that retroflect further downstream, at the tip of the Grand Banks, feed the 513 center of the subpolar gyre ($\sim 45-52^{\circ}$ N). 514

A more detailed look at the westward-flowing category provides information on the specific pathways of Labrador Current Waters. Some of the waters reaching the Slope Sea do so through the Avalon Channel (cluster #23), while others flow over the Grand Banks (cluster #18). We find that the waters entering the Laurentian Channel and reaching further south along the Scotian Shelf mostly go through the Avalon Channel (cluster #3 and 26). We also mentioned in Section 3.1.1 how cluster #22 contains particles that first go westward and are then retroflected (Fig. 4).



Figure 9. Left: Histograms of the longitudes of origin (along the seeding line), for each pathway. Middle: Histograms of the longitudes covered during the first 50 days, for each pathway. The first 50 days represent the portion of the trajectory north of Flemish Cap, where the current spreads longitudinally because of the presence of the Grand Banks. Right: For each pathway category, histograms of the average velocity of the particles (degrees of latitude travelled per day; zero indicates zonal displacement) over their first 100 days, the average time that particles spend north of the tip of the Grand Banks. The black vertical line indicates the peak of the histogram. The vertical grey lines in the left and middle panels show the location of the shelf-break. A cut-off at 0.20 was chosen for clarity, as the focus is on the bulk of the particles.

Finally, we find that the two clusters that belong to the southward-flowing pathway category are actually associated with two slightly different pathways (#12 and 29; Fig. 4). A first pathway flows *along* the Grand Banks shelf-break and, once it detaches from the shelf, veers slightly east and reaches deeper depths (maximal depths of 1500 m, cluster #12). Another pathway flows *over* the Grand Banks and, once it detaches from the shelf, continues straight to the south and reaches maximal depths of 700 m (cluster #29).

When looking at individual clusters, we also notice that the algorithm classified 529 particles based on their propagation time (Fig. 10). This is not surprising, since it clas-530 sified them based on their location at every time step, which implicitly contains infor-531 mation on the velocity. For instance, the particles classified in the westward-flowing path-532 way category take about 8 months to reach the Slope Sea from their seeding position, 533 except for the particles classified in cluster #18, which take about one year (Fig. 10). 534 Similarly, the particles classified in the retroflected pathway category take 2-3 months 535 to reach the tip of the Grand Banks, except for the particle classified in clusters #2, 9536 and 24, which take 4-9 months. While, here, our interest is on the various pathways of 537 the Labrador Current, information on the propagation time is useful to evaluate how long 538 it takes for anomalies carried by the Labrador Current to reach different export zones. 539

540

3.2.4 Characteristic depths of the pathways

Each identified pathway category has a distinct signature in depth (Fig. 11). The 541 particles moving eastward stay at shallow depths, while the particles moving southward 542 and westward reach deeper. The southward-flowing particles reach the deepest depths, 543 diving on average to maximal depths of ~ 1200 m or below 2000 m for 10% of them. These 544 particles cross the Gulf Stream front, which acts as a barrier to cross-front flow down 545 to 700 m, and as a stirrer below (Palter et al., 2013; Bower et al., 1985). Hence, it is ex-546 pected that the southward-flowing particles travel at such great depths, since only the 547 particles that subduct can cross the front and follow this pathway (Fig. 12). The westward-548 flowing particles dive on average to maximal depths of ~ 900 m, or even down to 1800 m 549 for 10% of them. Cluster #18 however remains above 700 m, gathering the particles that 550 are not entrained below the front. Within the retroflected pathway category, particles 551 descend once they quit the shelf, near the 50th meridian (Fig. 11b), reaching on aver-552 age depths of ~ 500 m, or 1000 m for 10% of them. The particles retroflecting at the tip 553 of the Grand Banks (clusters #2 and 10) reach greater depths (~1500 m) compared to particles retroflecting at Flemish Cap (\sim 700 m). Hence, the latter feed the core of the 555 North Atlantic Current (down to ~ 800 m, Gouretski, 2018), while the former feed the 556 deep ocean. Finally, the particles of the Labrador Sea pathway category remain relatively 557 close to the surface (above ~ 150 m). This is probably due to the low salinity (33-34.5) 558 of these waters compared to the open ocean (34.5-36.5), and suggests that the weak fresh-559 water export from the Labrador Current contributes to increasing the stratification in 560 the Labrador Sea (Howatt et al., 2018). 561

⁵⁶² 4 Discussion and conclusion

In this study, we present a method to classify geophysical Lagrangian trajectories 563 using unsupervised clustering. Our results demonstrate that this method is useful and 564 efficient to (i) classify Lagrangian tracks that are challenging to classify with more tra-565 ditional methods (e.g. counting particles crossing hydrographic sections), (ii) assist in 566 the treatment of huge Lagrangian tracks datasets, (iii) identify the main pathways of an 567 ocean current, and (iv) analyze the variability in the magnitude of these pathways. The 568 method was applied to 1.2 millions modelled trajectories along the Labrador Current and 569 was successful in identifying the different pathways of the Labrador Current, including 570 a previously unknown pathway directed southward from the tip of the Grand Banks. The 571



Figure 10. (Histograms, right) For each cluster in four pathway categories, histogram of the propagation time (in days) of the virtual particles, from the initialization line (thickest black on the maps on the left) to the entry point to the export zones associated with each category (medium-thickness black line, maps on the left). The cluster ID is indicated in the top right of the panel. We do not show the propagation time for the Labrador Shelf pathway category because they are not exported, and for the Southward-flowing pathway category because the export zone is not clearly definable. The thin black contour on the maps corresponds to the 350 m isobath.



Figure 11. (a) Average depth of the particles in each pathway category along the distance travelled by the particles, using 100 km bins. The size of the dots represent the number of particles used in the average for each distance bin. This number tends to decrease with the travelled distance because particles can exit the domain or hit bathymetry. We stop displaying the data when less than 25% of the particles remain. The vertical black line indicates the approximate location of the Tip of the Grand Banks. (b) For each pathway category, averaged longitude of the particles in a given depth bin. In both plots, the shaded areas show the zone encompassing the 10% and 90% longitude percentile.



Figure 12. Particle distribution along a transect at 50° W that runs across the Gulf Stream front, from the tip of the Grand Banks (43°N, left) to 32.5°N (right). The inset shows the location of the transect (red line). Colours provide the number of particles passing by. White lines show isopycnals of potential density (+1000 kg m⁻³). The strong tilt in the isopycnals around 39° N is due to the Gulf Stream front.

pre-processing applied here is relatively straightforward to implement, and the k-means++ 572 algorithm is simple to use and converges well (Section 2.3). The use of a larger number 573 of clusters than the expected number of pathways proves adequate, as it leads to a good 574 performance of the clustering and to the identification of details of the circulation that 575 we were initially not hoping to resolve. The choice of the number of clusters relies on two 576 metrics: the silhouette score and an ad hoc "physics-based" metrics tailored to our sci-577 entific question. Overall, the algorithm is relatively cheap to run, except for the kernal-578 ized PCA step (see Section 2.3), which requires a lot of computational resources (here, 579 186 GB of RAM for one day, on a HPC system). 580

The results of the clustering confirm that the Labrador Current splits into two main 581 branches: a branch retroflecting east towards the subpolar North Atlantic, representing 582 $\sim 50\%$ of the Labrador Current water, and a branch flowing west along the eastern Amer-583 ican continental shelf-break and into the Slope Sea, representing $\sim 20\%$ of it (Fig. 13). 584 Two-thirds of the eastward retroflection occurs at the tip of the Grand Banks, and a quar-585 ter at Flemish Cap. The waters retroflecting at the tip of the Grand Banks reach deeper 586 and get close to the Northeast corner, while the water retroflecting at Flemish Cap stay shallower and reach higher north. Secondary pathways of the Labrador Current include 588 one exporting water from the Labrador Shelf to the Labrador Sea, one entering the Gulf 589 of St. Lawrence through the Strait of Belle Isle, and one retroflecting after vising the Slope 590 Sea (Fig. 13). In addition to these pathways, which were already documented in the lit-591 erature, the clustering reveals a pathway bringing waters southwards from the tip of the 592 Grand Banks and representing on average 8% of the Lagrangian trajectories. This path-593 way has been described for the underlying Deep Western Boundary Current (Bower et 594 al., 2011, 2009) and suggests a connection between the two currents. 595

The variability of the two main branches is strong, reaching up to 96% of the mean state. These two branches strongly compensate each other through time, which is characteristic of a see-saw system. The time series of the magnitude of each pathway can be very useful to study what drives their variability. For instance, Jutras et al. (2023) in-

troduce an index for the retroflection of the Labrador Current and use it to study the 600 drivers of the retroflection and its impact on the physical and biogeochemical proper-601 ties of the northwestern Atlantic. That index is based on counting the number of par-602 ticles reaching a hydrographic line south of the Grand Banks, therefore not discriminat-603 ing the particles going southward from the tip of the Grand Banks, or reaching the Labrador 604 Sea, from those actually retroflected towards the subpolar North Atlantic. The time se-605 ries of the magnitude of the retroflected pathway obtained in the present study through 606 unsupervised clustering represent a more precise estimate of the magnitude of the retroflec-607 tion of the Labrador Current. 608

Finally, by analyzing the origin of each pathway and the associated water masses 609 (section 3.1.1), we can discuss what influences water properties in the export regions of 610 each pathway. The two main pathways, retroflected and westward-flowing, are fed by both 611 the inshore and shelf-break branches of the Labrador Current, but the shelf-break branch 612 contributes slightly more to the retroflected pathway (about 30%) and the inshore branch 613 slightly more to the westward-flowing pathway (about 90%, Fig. 13). The inshore branch 614 is fed by Arctic outflow through Davis Strait and by river outflow along the Labrador 615 Shelf, while the shelf-break branch is mostly fed by the Greenland Current (section 1, 616 Florindo-López et al., 2020). Hence, variations in the magnitude of rivers outflow along 617 the Labrador Shelf likely affect salinity in the Slope Sea slightly more than in the sub-618 polar North Atlantic. In contrast, variations in salinity in the Greenland Current, due 619 to changes in the Arctic freshwater outflow (de Steur et al., 2018) or to Greenland ice 620 sheet melt (Marson et al., 2021), will affect salinity in the subpolar North Atlantic slightly 621 more than in the Slope Sea. These variations would also likely affect salinity in the Labrador 622 Sea through the Labrador Sea pathway, fed exclusively by this pathway. The southward-623 flowing pathway has a weak variability and is not associated with a particular branch 624 of the Labrador Current. Hence, its variability does not contribute to that in western 625 North Atlantic Ocean water properties, but changes in its water properties could. Fi-626 nally, we also find that the pathways exporting water to the Labrador Sea and to the 627 subpolar North Atlantic supply the surface ocean (Fig. 11 and 13). Since the Labrador 628 Current carries freshwater, variations in these exports would likely affect the stratifica-629 tion in these regions, including the occurrence and intensity of deep convection, with po-630 tential effects on the Atlantic Meridional Overturning Circulation (AMOC; Lozier, 2012). 631 carbon uptake (Fontela et al., 2016), and on oxygen repletion of the deep North Atlantic 632 waters (Koelling et al., 2022; Atamanchuk et al., 2021). 633

To finish, this paper offers first and foremost methodological advancements for the geophysical community. The method, extensively described in this paper, could be applied to other oceanic currents or other types of geophysical Lagrangian trajectories.

5 Open Research

The Lagrangian tracking experiments can be reproduced by downloading the publicly available GLORYS12V1 outputs from the Copernicus Marine Environment Monitoring Service (CMS) website: resources.marine.copernicus.eu/product-detail/ GLOBAL_MULTIYEAR_PHY_001_030/INFORMATION. Information about the OceanParcels tool for Python is available at oceanparcels.org. The scripts used to run the Lagrangian tracking experiments can be found as a supplementary material to Jutras et al. (2023).

The ML tools are available through the Python scikit-learn package (scikit-learn .org/). The scripts of the unsupervised clustering method are available at https://github .com/noemieplanat/Clustering_Lagragian_particles.

The data from the Global drifter program was obtained from the Atlantic Oceanographic and Meteorological Laboratory of the National Oceanic and Atmospheric Administration (AOML/NOAA, ftp.aoml.noaa.gov/phod/pub/buoydata). The RAFOS/SOFAR subsurface float trajectories are compiled from 52 experiments by the WOCE Subsurface Float Data Assembly Center (WFDAC, www.aoml.noaa.gov/phod/float_traj/).



Figure 13. Summary of the pathways of the Labrador Current as identified by the unsupervised clustering of Lagrangian trajectories. The dashed arrows indicate the shelf-branch (green) and the shelf-break (blue) branches of the Labrador Current. The full arrows indicate the different pathways, and are identified with a letter: (A) Labrador Sea, (B) Labrador Shelf, (C) Belle Isle, (D) westward-flowing, (E) southward-flowing, (F) retroflected eastward, (D₂) westwardflowing and then retroflected. Pathway (B) ends with a bar because it contains particles that die on the shelf after they hit bathymetry. For each pathway, the width of the arrow indicates its averaged magnitude (corresponding to the fraction of the particles classified in that pathway category), and the color of the arrows indicates which of the Labrador Current branches mainly feeds the pathway. Colored triangles indicate the depth reached by the particles. The month labels indicate the average transit time from initialization to the export zone. The insert in the bottom-right illustrates the see-saw behaviour of the two main pathways (westward-flowing and retroflected): one weakens as the other strengthens, and vice-versa. The thin gray line indicates the 250 m isobath.

- ⁶⁵² The Argo data were collected and made freely available by the International Argo Pro-
- gram and the national programs that contribute to it (argo.ucsd.edu, https://www.ocean
- -ops.org). The Argo Program is part of the Global Ocean Observing System.
655 Appendix A Supplementary material



Figure A1. Histograms of the initialization depth of the particles associated with the different pathways.



Figure A2. Trajectories in each cluster, for the test set. The color indicates the depth of the particles.



Figure A3. Same as Figure 4, but indicating in the top right the score of the physical metrics (P.S.) defined in section 2.3 and the internal spread (Spr).

1	67%	11	90%	21	100%
2	70%	12	100%	22	66%
3	90%	13	90%	23	90%
4	90%	14	100%	24	60%
5	90%	15	100%	25	100%
6	90%	16	100%	26	50%
7	100%	17	55%	27	90%
8	100%	18	90%	28	100%
9	90%	19	55%	29	100%
10	90%	20	100%	30	100%

Table B1. Agreement rate during the expert's classification into the different pathway categories, for each cluster (section 2.3.4).

656 Appendix B Clusters grouping exercise

The agreement rate between the consulted experts is of 100% in 12 clusters, all agree 657 but one or two in 10 clusters, and is above 60% in 5 clusters (Table B1). The agreement 658 rate is of 55% for the westward-flowing cluster #17, that 45% of the experts assign to 659 the Labrador Shelf category. We assign this cluster to the Labrador Shelf category, be-660 cause the strong majority of the particles remain on the Labrador Shelf. Disagreement 661 regarding the Labrador Sea pathway is probably due to a lack of clear distinction be-662 tween the Labrador Sea and the north of the subpolar North Atlantic. For the retroflected 663 cluster #19, 45% of the experts assign to the Labrador Sea category. We assign cluster 664 it to the retroflected category because, even if some of the particles in this cluster cross 665 the southern portion of the Labrador Sea, they eventually feed the subpolar North At-666 lantic, where they will affect the water properties. For similar reasons, we go against the 667 expert agreement on cluster #11 (which was categorized as belonging to the Labrador 668 Sea category), because the particles are retroflected at Flemish Cap before aiming North. 669 Finally, there is equality in the vote for cluster #26, between the Belle Isle and westward-670 flowing categories. An analysis of the individual trajectories, as opposed to a density view, 671 reveals that while a fair amount of particles enter Belle Isle Strait, most go around New-672 foundland and some around the Grand Banks (not shown). We therefore assign this clus-673 ter to the westward-flowing category. 674

675 Acknowledgments

The authors would like to thank Pr. Daniel Bourgault, Frédéric Cyr, Pr. Juliana Marson, Pr. Alfonso Mucci, Pr. Paul Myers, Pr. Jaime Palter, Nicolai von Oppeln-Bronikowski, Mariona Claret and Peter S. Galbraith for acting as experts in classifying the clusters into pathway categories, as well as Jan Klaus Rieck, David Straub and Bruno Tremblay for testing the classification protocol. We also want to thank Pr. David Rolnick for an informal conversation on Machine Learning applications to geosciences.

This research was enabled in part by support provided by Calcul Québec (calculquebec 682 .ca) and Compute Canada (compute canada.ca). MJ was supported by the The Nat-683 ural Sciences and Engineering Research Council of Canada (NSERC), the Fonds de recherche 684 du Québec - Nature et technologie (FRQNT), Ouranos; NP by the FRQNT; and LCT 685 by McGill University Science Undergraduate Research Award. This research was also 686 funded by NSERC Accelerator Supplements (grant no. RGPAS/2018-522502) and the 687 Canada Research Chair program awarded to COD, as well as by a NSERC Discover Grant 688 awarded to Pr. Bruno Tremblay (grant no. RGPIN/2018-04838). Finally, we acknowl-689 edge the support of Québec-Océan. 690

MJ and NP designed and conducted the study with input from COD. MJ did the 691 preprocessing of the data, and NP developed the ML model, with the assistance of MJ. 692 LCT processed and analyzed the drifter and float dataset. MJ and NP analyzed the re-693 sults and COD contributed to the interpretation. MJ wrote the manuscript, with con-694 tribution from NP for the method section, from LCT for the observation section and from 695 COD for editing. 696

References 697

715

- Abdar, M., Pourpanah, F., Hussain, S., Rezazadegan, D., Liu, L., Ghavamzadeh, 698 M., ... Nahavandi, S. (2021, December). A review of uncertainty quan-699 tification in deep learning: Techniques, applications and challenges. In-700 formation Fusion, 76, 243–297. Retrieved 2023-03-02, from https:// 701 www.sciencedirect.com/science/article/pii/S1566253521001081 doi: 702 10.1016/j.inffus.2021.05.008 703 Atamanchuk, D., Palter, J., Palevsky, H., Le Bras, I., Koelling, J., & Nicholson, 704
- D. (2021, December). Linking Oxygen and Carbon Uptake with the Merid-705 ional Overturning Circulation Using a Transport Mooring Array. In Fron-706 tiers in Ocean Observing: Documenting Ecosystems, Understanding Envi-707 ronmental Changes, Forecasting Hazards. (E.S. Kappel, S.K. Juniper, S. 708
- Seevave, E. Smith, and M. Visbeck ed., p. 9). Retrieved 2023-03-17, from 709 https://doi.org/10.5670/oceanog.2021.supplement.02-03 710
- Ayata, S. D., Lazure, P., & Thiébaut, (2010).How does the connectivity be-711 tween populations mediate range limits of marine invertebrates? A case 712 study of larval dispersal between the Bay of Biscay and the English Chan-713 nel (North-East Atlantic). Progress in Oceanography, 87(1-4), 18–36. doi: 714 10.1016/j.pocean.2010.09.022
- Boehme, L., & Rosso, I. (2021).Classifying Oceanographic Structures in the 716 Amundsen Sea, Antarctica. Geophysical Research Letters, 48(5), 1–8. doi: 717 10.1029/2020GL089412 718
- Bower, A. S., Lozier, M. S., Gary, S. F., & Böning, C. W. (2009). Interior pathways 719 of the North Atlantic meridional overturning circulation. Nature, 459(7244), 720 243–247. doi: 10.1038/nature07979 721
- Bower, A. S., Lozier, S., & Gary, S. (2011). Export of Labrador Sea Water from the 722 Deep-Sea Research Part subpolar North Atlantic: A Lagrangian perspective. 723 II: Topical Studies in Oceanography, 58(17-18), 1798–1818. Retrieved from 724 http://dx.doi.org/10.1016/j.dsr2.2010.10.060 (Publisher: Elsevier) 725 doi: 10.1016/j.dsr2.2010.10.060 726
- Bower, A. S., Rossby, H. T., & Lillibridge, J. L. (1985). The Gulf Stream-Barrier 727 or Blender? Journal of Physical Oceanography, 15(1), 24–32. doi: 10.1175/ 728 1520-0485(1985)015(0024:tgsob)2.0.co;2 729
- Brankov, E., Trivikrama Rao, S., & Steven Porter, P. (1998).A Trajectory-730 Clustering-Correlation Methodology for Examining the Long-Range Transport 731 of Ari Pollutants. Atmospheric Environment, 32(9), 1525–1534. 732
- Brooks, D. A. (1987). The Influence of Warm-Core Rings on Slope Water Entering 733 the Gulf of Maine. Journal of Geophysical Research, 92(C8), 8183–8196. 734
- Brunette, C., Tremblay, B., & Newton, R. (2019, February). Winter Coastal 735 Divergence as a Predictor for the Minimum Sea Ice Extent in the Laptev 736 Journal of Climate, 32(4), 1063–1080. Sea. Retrieved 2022-11-25, from https://journals.ametsoc.org/doi/10.1175/JCLI-D-18-0169.1 doi: 738 10.1175/JCLI-D-18-0169.1 739
- Cetina-Heredia, P., Roughan, M., van Sebille, E., Feng, M., & Coleman, M. A. 740 (2015).Strengthened currents override the effect of warming on lobster lar-741 val dispersal and survival. Global Change Biology, 21(12), 4377–4386. doi: 742 10.1111/gcb.13063 743

744	Chabot, D., & Dutil, J. D. (1999). Reduced growth of Atlantic cod in non-lethal hy-
745	poxic conditions. Journal of Fish Biology, 55(3), 472–491. doi: 10.1006/jfbi
746	.1999.1005
747	Chen, Z., Kwon, Y. O., Chen, K., Fratantoni, P. S., Gawarkiewicz, G., & Joyce,
748	T. M. (2020). Long-Term SST Variability on the Northwest Atlantic Con-
749	tinental Shelf and Slope. Geophysical Research Letters, $47(1)$, 1–11. doi:
750	10.1029/2019GL085455
751	Claret, M., Galbraith, E. D., Palter, J. B., Bianchi, D., Fennel, K., Gilbert, D.,
752	& Dunne, J. P. (2018). Rapid coastal deoxygenation due to ocean cir-
753	culation shift in the northwest Atlantic. Nature Climate Change, 8. Re-
754	trieved from http://www.nature.com/articles/s41558-018-0263-1 doi:
755	10.1038/s41558-018-0263-1
756	Daher, H., Beal, L. M., & Schwarzkopf, F. U. (2020). A New Improved Estimation
757	of Agulhas Leakage Using Observations and Simulations of Lagrangian Floats
758	and Drifters. Journal of Geophysical Research: Oceans, 125(4), 1–17. doi:
759	10.1029/2019JC015753
760	Dee, D. P., Uppala, S. M., Simmons, A. J., Berrisford, P., Poli, P., Kobavashi, S.,
761	Vitart, F. (2011). The ERA-Interim reanalysis: Configuration and performance
762	of the data assimilation system. Quarterly Journal of the Royal Meteorological
763	Society, 137(656), 553–597. doi: 10.1002/gj.828
764	Delandmeter, P., & Van Sebille, E. (2019). The Parcels v2.0 Lagrangian framework:
765	New field interpolation schemes. Geoscientific Model Development, 12(8).
766	3571–3584. doi: 10.5194/gmd-12-3571-2019
767	de Steur, L., Peralta-Ferriz, C., & Paylova, O. (2018). Freshwater Export
768	in the East Greenland Current Freshens the North Atlantic. Geonhusi-
769	cal Research Letters, 45(24), 13,359–13,366. Retrieved 2023-06-22, from
770	https://onlinelibrary.wiley.com/doi/abs/10.1029/2018GL080207
771	(_eprint: https://onlinelibrary.wiley.com/doi/pdf/10.1029/2018GL080207)
772	doi: 10.1029/2018GL080207
773	Echols, R., Rocap, G., & Riser, S. (2020). Exploring biogeographical provinces with
774	machine learning and Argo float data. In AGU 2020 fall meeting (pp. OS022–
775	06).
776	Filippi, M., Hadiighasem, A., Rayson, M., Rypina, I. L. Ivey, G., Lowe, R.,, Pea-
777	cock, T. (2021). Investigating transport in a tidally driven coral atoll flow
778	using Lagrangian coherent structures. Limnology and Oceanography, 66(11),
779	4017–4027. doi: 10.1002/lno.11939
780	Filippi, M., Rypina, I. I., Hadiighasem, A., & Peacock, T. (2021). An optimized-
781	parameter spectral clustering approach to coherent structure detection in
782	geophysical flows. Fluids, $6(1)$, 1–34. doi: 10.3390/fluids6010039
783	Fischer, J., & Schott, F. A. (2002). Labrador Sea Water tracked by profiling floats -
784	From the boundary current into the Open North Atlantic. Journal of Physical
785	Oceanography, 32(2), 573-584.
786	Florindo-López, C., Bacon, S., Aksenov, Y., Chafik, L., Colbourne, E., Penny Hol-
787	liday, N., & Holliday, N. P. (2020). Arctic Ocean and Hudson Bay Fresh-
788	water Exports : New Estimates from 7 Decades of Hydrographic Sur-
789	vevs on the Labrador Shelf. Journal of Climate, 33(20), 1–62. doi:
790	10.1175/jcli-d-19-0083.1
701	Fontela M García-Ibáñez M I Hansell D A Mercier H & Pérez F F (2016)
792	Dissolved Organic Carbon in the North Atlantic Meridional Overturning Cir-
793	culation. Scientific Reports 6 1–9 Retrieved from http://dx doi_org/
79/	10.1038/srep26931 (Publisher: Nature Publishing Group ISBN: 2045-2322
795	(Electronic) 2045-2322 (Linking)) doi: 10.1038/srep26931
796	Fox, A. D., Biastoch, A., Cunningham, S. A. Garv, S. F. Handmann, P. Penny, N.
797	Schmidt, C. (2022). How reduced Labrador Sea surface heat loss caused
798	exceptional freshening and cooling in the eastern subpolar North Atlantic

799	Ocean Science, in prep.(April), 1–35.
800	Fratantoni, P. S., & McCartney, M. S. (2010). Freshwater export from the Labrador
801	Current to the North Atlantic Current at the Tail of the Grand Banks of New-
802	foundland. Deep-Sea Research Part I: Oceanographic Research Papers, 57(2).
803	258–283. Retrieved from http://dx.doi.org/10.1016/j.dsr.2009.11.006
804	doi: 10.1016/i.dsr 2009.11.006
004	Cilbert D Sundhy B B Cobeil C Mucci A & Tremblay C H (2005) A
805	seventy two year record of diminishing doop water ovygen in the St. Lawrence
806	ostupry: The northwest Atlantic connection Limneleeu and Oceanography
807	50(5) 1654 1666
808	Cillard I C Hu X Muora D C l_2 Dombor I I (2016) Moltwater nothward
809	Ginard, L. C., Hu, A., Myers, P. G., & Damber, J. L. (2010). Mentwater pathways
810	from marine terminating glaciers of the Greemand ice sneet. Geophysical Re-
811	search Letters, $43(20)$, $10,873-10,882$. doi: $10.1002/2010GL070909$
812	Gonçalves Neto, A., Langan, J. A., & Palter, J. B. (2021). Changes in the Gui
813	Stream precede rapid warming of the Northwest Atlantic Shelf. Nature Com-
814	munications Earth & Environment, 2(74). Retrieved from http://dx.doi
815	.org/10.1038/s4324/-021-00143-5 doi: 10.1038/s4324/-021-00143-5
816	Gonçalves Neto, A., Palter, J. B., Xu, X., & Fratantoni, P. (2023). Temporal
817	Variability of the Labrador Current Pathways Around the Tail of the Grand
818	Banks at Intermediate Depths in a High-Resolution Ocean Circulation Model.
819	Journal of Geophysical Research: Oceans, 128(3), e2022JC018756. Retrieved
820	2023-03-13, from https://onlinelibrary.wiley.com/doi/abs/10.1029/
821	2022JC018756 doi: 10.1029/2022JC018756
822	Gouretski, V. (2018). WOCE-Argo Global Hydrographic Climatology (WAGHC
823	<i>Version 1.0).</i> (Publication Title: World Data Center for Climate (WDCC) at
824	DKRZ)
825	Han, G. (2005). Wind-driven barotropic circulation off Newfoundland and Labrador.
826	Continental Shelf Research, 25(17), 2084–2106. doi: 10.1016/j.csr.2005.04.015
827	Han, G., Chen, N., & Ma, Z. (2014). Is there a north-south phase shift in the sur-
828	face Labrador Current transport on the interannual-to-decadal scale? Journal
829	of Geophysical Research: Oceans, $119(1)$, 276–287. (ISBN: 2169-9275) doi: 10
830	.1002/2013 JC009102
831	Han, G., Loder, J. W., & Smith, P. C. (1999). Seasonal-Mean Hydrography and
832	Circulation in the Gulf of St. Lawrence and on the Eastern Scotian and South-
833	ern Newfoundland Shelves. Journal of Physical Oceanography, $29(6)$, 1279–
834	1301. Retrieved from http://journals.ametsoc.org/doi/abs/10.1175/
835	1520-0485%281999%29029%3C1279%3ASMHACI%3E2.0.C0%3B2 (ISBN: 0022-
836	3670) doi: $10.1175/1520-0485(1999)029(1279:SMHACI)2.0.CO;2$
837	Han, G., Ma, Z., & Chen, N. (2019). Ocean climate variability off Newfound-
838	land and Labrador over 1979–2010: A modelling approach. Ocean Mod-
839	elling, 144 (October), 101505. Retrieved from https://doi.org/10.1016/
840	j.ocemod.2019.101505 doi: 10.1016/j.ocemod.2019.101505
841	Handmann, P., Fischer, J., Visbeck, M., Karstensen, J., Biastoch, A., Böning,
842	C. W., & Patara, L. (2018). The Deep Western Boundary Current in
843	the Labrador Sea From Observations and a High-Resolution Model. Jour-
844	nal of Geophysical Research: Oceans, 123(4), 2829–2850. doi: 10.1002/
845	2017JC013702
846	Hart-Davis, M., & Backeberg, B. (2021, April). Towards a particle trajectory mod-
847	elling approach in support of South African search and rescue operations at
848	sea. Journal of Operational Oceanography, 1–9. Retrieved 2022-11-25, from
849	https://www.tandfonline.com/doi/full/10.1080/1755876X.2021.1911485
850	doi: 10.1080/1755876X.2021.1911485
851	Hertwig, D., Burgin, L., Gan, C., Hort, M., Jones, A., Shaw, F., Zhang, K.
852	(2015). Development and demonstration of a Lagrangian dispersion modeling
853	system for real-time prediction of smoke haze pollution frombiomass burning in

854	Southeast Asia. Journal of Geophysical Research, 120(24), 12,605–12,630. doi: 10.1002/2015JD023422
856	Hofmann, T., Schölkopf, B., & Smola, A. J. (2008, June). Kernel methods in
857	machine learning. The Annals of Statistics, 36(3), 1171–1220. Retrieved 2022-
858	11-28. from https://projecteuclid.org/journals/annals-of-statistics/
859	volume-36/issue-3/Kernel-methods-in-machine-learning/10.1214/
860	009053607000000677.full (Publisher: Institute of Mathematical Statistics)
861	doi: 10.1214/00905360700000677
862	Holliday N P Bersch M Bers B Chafik L Cunningham S Florindo-López
863	C Yashavaev, I. (2020). Ocean circulation causes the largest freshening
864	event for 120 years in eastern subpolar North Atlantic Nature Communica-
865	tions, $11(1)$, doi: 10.1038/s41467-020-14474-v
866	Houghton I A & Wilson J D (2020) El Niño Detection Via Unsupervised Clus-
867	tering of Argo Temperature Profiles Journal of Geophysical Research: Oceans
969	125(9) 1–12 doi: 10.1029/2019.IC015947
000	Howatt T M Palter I B Matthews I B B deVoung B Bachmayer B k
809	Claus B (2018) Ekman and addy exchange of freshwater and ovygen across
870	the Labrador Shelf break Iournal of Physical Oceanography 18(5) 1015-
871	1031 doi: 10.1175/IPO-D-17-0148.1
872	Jones D. C. Holt H. J. Maijars A. J. & Shuckhurgh F. (2010) Unsupervised
8/3	Clustering of Southern Ocean Argo Float Temperature Profiles Lowrad of
874	Combusical Research: Oceans 12/(1) 300-402 doi: 10.1020/2018 IC014620
875	$\begin{array}{c} \text{Geophysical Rescaled. Occurs, 124(1), 390-402. \text{ doi: 10.1029/20103C014029} \\ \text{Intrag. M. Dufour, C. O. Muggi A. Cur, F. & Cilbert, D. (2020) Temporal. \end{array}$
876	Changes in the Causes of the Observed Ovygen Decline in the St. Lawrence
877	Estuary Lowrad of Coonhusical Research: Oceans 125(12) 1 20 doi:
878	10 1020 /2020 IC016577
879	10.1029/2020JC010377 Jutras M. Dufaun Carolina O. Mussi Alfanza & Talbat Laurum (2022) Demote
880	Jutras, M., Dulour, Carolina O., Mucci, Alfonso, & Talbot, Lauryn. (2023). Remote
881	Lingson D. E. & Wetson J. D. (2016 April) The timescales of global surface
882	Joinsson, D. F., & Watson, J. R. (2010, April). The timescales of global surface-
883	11.25 from https://www.networe.com/orticles/neerme11220 (Number 1
884	Dublishow Nature Dublishing Crown) doi: 10.1029/ncommo11220
885	Verregeli T. Metarreure V. & Hearrei H. (2022) Deep meter petbruore in
886	the North Decife Ocean revealed by Legrangian particle tracking
887	the North Fachic Ocean revealed by Lagrangian particle tracking. $Scientific Perpendicular (10, 1) = 10$
888	c/1502 022 10020 2 (Dublicher: Noture Dubliching Croup IIV ISDN:
889	(1 ubishing Group OX 15DN, 1150802210080) doi: 10.1038/s41508.022.10080 8
890	
891	the Origin of Water Massag in the Boaufort Cure Lowroad of Coonhusical Re
892	the Origin of water masses in the beautoft Gyre. Journal of Geophysical Re- search: O_{acana} , $12/(7)$, 4606 , 4700 , doi: 10.1020/2010IC015022
893	$V_{122}^{122} = M_{122}^{122} (1), 4050-4109. $ doi: 10.1029/20190001022
894	actions: A Prostice Driven Classification of Uncertainty in Machine Learning Appli-
895	A Strayburg E Schoitach & E Ditach (Edg.) Computer Safety Deliabil
896	A. Skavnaug, E. Scholtsch, & F. Ditsch (Eds.), Computer Sujety, Retadou-
897	$10 \ 1007 \ /078 \ 3 \ 210 \ 00220 \ 7 \ 36$
898	Valling L Atomorphyle D Konstonson L Handmann D & Wallace D W D
899	(2022 January) — Owners around to the deep accentally in Tahuadan Carl
900	(2022, January). Oxygen export to the deep ocean following LaDrador Sea Water formation $Bioaccession and 10(2) 427 454 Detwiewed 2022 06 22$
901	water formation. Diogeosciences, $19(2)$, $437-454$. Refrieved 2025-00-22, from https://bg.copporpieus.org/orticl.og/10/427/2022/ (Dublisher)
902	Constraints CmbH) doi: 10.5104/bg.10.427.2022 (PUDISITER:
903	Copermicus Gimbil) uci. 10.0194/ 0g-19-401-2022
904	ROSZAIKA, I. IVI., & Lacasce, J. R. (2010, August). Lagrangian analysis by clustering.
905	$U_{\text{remov}} = \frac{1}{2} \int \frac{1}{2} \int$
906	tailored to structure sharps of tropical suslance wing a very large sur-
907	tanored to structure change of tropical cyclones using a very large num-
908	before the periodic trajectories. Moning weather Review, $14\delta(10)$, $4209-4229$. doi:

909	10.1175/MWR-D-19-0408.1
910	Lazier, J. R. N., & Wright, D. G. (1993). Annual velocity variations in the Labrador
911	Current. Journal of Physical Oceanography, 23(4), 659–678. doi: 10.1175/1520
912	-0485(1993)023(0659:AVVITL)2.0.CO:2
913	Lebreton, L. C., Greer, S. D., & Borrero, J. C. (2012). Numerical modelling of
914	floating debris in the world's oceans. Marine Pollution Bulletin, 64 (3), 653–
915	661. Retrieved from http://dx.doi.org/10.1016/j.marpolbul.2011.10.027
916	(Publisher: Elsevier Ltd) doi: 10.1016/i.marpolbul.2011.10.027
017	Lellouche J-M Greiner E Le Galloudec O Garric G Begnier C Drevillon
018	M Le Traon P Y (2018) Recent updates to the Copernicus Marine
010	Service global ocean monitoring and forecasting real-time $1/12^{\circ}$ high-resolution
919	system Ocean Science $1/(5)$ 1093–1126 doi: 10.5194/os-14-1093-2018
021	Loder I W Petrie B D & Gawarkiewicz G (1998) The coastal ocean off north-
921	eastern North America: a large-scale view. In <i>The Sea</i> (Vol. 11, pp. 105–133)
922	Lozier M S (2012) Overturning in the North Atlantic Annual Re-
923	view of Marine Science /(1) 201-315 Retrieved 2023-02-07 from
924	t = 120710 - 100740
925	https://doi.org/10.1146/annurey_marine_120710_100740
920	annurey_marine_120710_100740
927	Madae G. Bourdallé Badia B. Chanut I. Clomonti F. Coward A. Ethé C.
928	(2010) NFMQ accan engine (Version $u(0)$) Notes Du Pôlo Do
929	Modélisation Do L'institut Pierre simon Laplace (IPSL) Zenodo [Dataset]
930	Marson I M Cillard I C & Myors P C (2021) Distinct Ocean Responses
931	to Croopland's Liquid Bunoff and Johorg Molt
932	Research: Oceans, 196(12), 2001 IC017542 Botrioved 2023 06 22 from
933	https://onlinelibrary.wiley.com/doi/abs/10.1020/2021 $IC017542$
934	(eprint: https://onlinelibrary.wiley.com/doi/pdf/10.1029/2021JC017542)
935	doi: 10.1029/2021 IC017542
930	Marson I M Myers P C Hu X & Le Sommer I (2018) Using Verti-
937	cally Integrated Ocean Fields to Characterize Greenland Icebergs' Distribu-
930	tion and Lifetime Geonbusical Research Letters (5(9), 4208–4217 doi:
939	10 1029/2018GL077676
940	Merino N Le Sommer I Durand G Jourdain N C Madec G Mathiot P
941	& Tournadre I (2016) Antarctic icebergs melt over the Southern Ocean :
942	Climatology and impact on sea ice Ocean Modelling 10/ 99–110 Betrieved
943	from http://dx.doj.org/10_1016/j_ocemod_2016_05_001 (Publisher:
945	Elsevier Ltd) doi: 10.1016/i.ocemod.2016.05.001
046	Mills K E Pershing A J Brown C J Chen Y Chiang F S Holland D S
947	Wahle R A (2013) Fisheries management in a changing climate: Lessons
948	from the 2012 ocean heat wave in the Northwest Atlantic. Oceanography.
949	26(2), doi: 10.5670/oceanog.2013.27
050	Myers P G (2005) Impact of freshwater from the Canadian Arctic Archinelago on
950	Labrador Sea Water formation Geophysical Research Letters 32(6) 1–4 doi:
951	10 1029/2004GL022082
053	New A Smeed D A Czaja A Blaker A T Mecking J V Mathews J P
954	Sanchez-Franks, A. (2021). Labrador Slope Water Connects the Subarctic with
955	the Gulf Stream. Environmental Research Letters, 16(8), 084019. (Publisher
956	IOP Publishing) doi: 10.1088/1748-9326/ac1293
957	Palter, J. B., Lozier, M. S., & Lavender, K. L. (2008) How does Labrador sea wa-
958	ter enter the deep western boundary current? Journal of Physical Oceanoara-
950	<i>phy.</i> 38(5), 968–983, doi: 10.1175/2007.IPO3807.1
960	Palter, J. B., Marinov, L., Sarmiento, J. L. & Gruber N. (2013) Large-Scale Per-
961	sistent Nutrient Fronts of the World Ocean: Impacts on Biogeochemistry In
962	(pp. 1–38). Berlin, Heidelberg: Springer. Retrieved 2023-03-03, from https://
963	doi.org/10.1007/698_2013_241 doi: 10.1007/698_2013_241

964	Pershing, A. J., Kerr, L. A., Bris, A. L., Mills, K. E., Sherwood, G. D., Alexander,
965	M. A., Thomas, A. C. (2016). Slow adaptation in the face of rapid warm-
966	ing leads to collapse of the Gulf of Maine cod fishery. Science, 352(6284), 423.
967	doi: 10.1126/science.aae0463
968	Petrie, B., Toulany, B., & Garrett, C. (1988). The transport of water, heat and salt
969	through the strait of Belle Isle. Atmosphere - $Ocean$, $26(2)$.
970	Petrie, B. D., & Drinkwater, K. F. (1993). Temperature and Salinity Variability on
971	the Scotian Shelf and in the Gulf of Maine 1945-1990. Journal of Geophysical
972	Research, 98(C11), 20079–20089.
973	Phelps, J. J., Polton, J. A., Souza, A. J., & Robinson, L. A. (2015). Behaviour
974	influences larval dispersal in shelf sea gyres: Nephrops norvegicus in the Irish
975	Sea. Marine Ecology Progress Series, 518, 177–191. doi: 10.3354/meps11040
976	Poitevin, P., Thebault, J., Siebert, V., Donnet, S., Archambault, P., Doré, J.,
977	Lazure, P. (2019). Growth response of Arctica islandica to North Atlantic
978	oceanographic conditions since 1850. Frontiers in Marine Science, 6, 1–14.
979	doi: 10.3389/fmars.2019.00483
980	Pérez-Brunius, P., Rossby, T., & Watts, D. R. (2004). Absolute transports of mass
981	and temperature for the North Atlantic Current-subpolar front system. Jour-
982	nal of Physical Oceanography, 34(8), 1870–1883.
092	Roach C. I. & Speer K (2019) Exchange of Water Between the Boss Gyre
984	and ACC Assessed by Lagrangian Particle Tracking Journal of Geophysical
985	Research: Oceans, 124(7), 4631–4643, doi: 10.1029/2018JC014845
986	Rossby T (1999) On gyre interactions Deen-Sea Research Part II: Tonical Studies
987	in Oceanoaranhy, 46(1-2), 139–164, doi: 10.1016/S0967-0645(98)00095-2
088	Rosso I Mazloff M B Talley L D Purkey S G Freeman N M & Maze
900	G = (2020) Water Mass and Biogeochemical Variability in the Kerguelen
909	Sector of the Southern Ocean: A Machine Learning Approach for a Mixing
991	Hot Spot. Journal of Geophysical Research: Oceans. 125(3), 1–23. doi:
992	10.1029/2019JC015877
003	Rousseeuw P. J. (1987) Silhouettes: A graphical aid to the interpretation and vali-
995	dation of cluster analysis Journal of Computational and Annlied Mathematics
995	20(C), 53–65. doi: 10.1016/0377-0427(87)90125-7
996	Saucier, F. J., Roy, F., Gilbert, D., Pellerin, P., & Ritchie, H. (2003). Modeling the
997	formation and circulation processes of water masses and sea ice in the Gulf
998	of St. Lawrence, Canada. Journal of Geophysical Research C: Oceans, 108(8),
999	25-1. Retrieved from http://doi.wiley.com/10.1029/2000JC000686 (ISBN:
1000	2156-2202) doi: 10.1029/2000jc000686
1001	Schmidt, S., & Send, U. (2007). Origin and composition of seasonal Labrador Sea
1002	freshwater. Journal of Physical Oceanography, 37(6), 1445–1454. doi: 10.1175/
1003	JPO3065.1
1004	Schneide, C., Pandey, A., Padberg-Gehle, K., & Schumacher, J. (2018). Probing tur-
1005	bulent superstructures in Rayleigh-Bénard convection by Lagrangian trajectory
1006	clusters. <i>Physical Review Fluids</i> , $3(11)$, 1–15. (arXiv: 1810.07396 Publisher:
1007	American Physical Society) doi: 10.1103/PhysRevFluids.3.113501
1008	Schneider, L., Kieke, D., Jochumsen, K., Colbourne, E., Yashayaev, I., Steinfeldt, R.,
1009	Rhein, M. (2015). Variability of Labrador Sea Water transported through
1010	Flemish Pass during 1993–2013. Journal of Geophysical Research: Oceans,
1011	120, 5514–5533. doi: 10.1002/2015JC010939
1012	Schulze Chretien, L. M., & Frajka-Williams, E. (2018). Wind-driven transport
1013	of fresh shelf water into the upper 30m of the Labrador Sea. Ocean Science,
1014	14(5), 1247-1264. doi: $10.5194/os-14-1247-2018$
1015	Seidov, D., Mishonov, A., & Parsons, R. (2021). Recent warming and
1016	decadal variability of Gulf of Maine and Slope Water. Limnology and
1017	<i>Oceanography</i> , 66(9), 3472–3488. Retrieved 2022-11-29, from https://
1018	onlinelibrary.wiley.com/doi/abs/10.1002/lno.11892 (_eprint:

1019	https://onlinelibrary.wiley.com/doi/pdf/10.1002/lno.11892) doi: 10.1002/
1020	lno.11892
1021	Shaw, JL., & Galbraith, P. S. (2023). Climatology of Transport in
1022	the Strait of Belle Isle. Journal of Geophysical Research: Oceans,
1023	128(2), e2022JC019084. Retrieved 2023-03-15, from https://
1024	onlinelibrary.wiley.com/doi/abs/10.1029/2022JC019084 (_eprint:
1025	https://onlinelibrary.wiley.com/doi/pdf/10.1029/2022JC019084) doi:
1026	10.1029/2022JC019084
1027	Simons, R. D., Siegel, D. A., & Brown, K. S. (2013). Model sensitivity and ro-
1028	bustness in the estimation of larval transport: A study of particle track-
1029	ing parameters. Journal of Marine Systems, 119-120, 19–29. Retrieved
1030	from http://dx.doi.org/10.1016/j.jmarsys.2013.03.004 doi:
1031	10.1016/j.jmarsys.2013.03.004
1032	Tamsitt, V., Drake, H. F., Morrison, A. K., Talley, L. D., Dufour, C. O., Gray,
1033	A. R., Weijer, W. (2017). Spiraling pathways of global deep waters to
1034	the surface of the Southern Ocean. Nature Communications, $8(1)$, 1–10. doi:
1035	10.1038/s41467-017-00197-0
1036	Thomas, S. D. A., Jones, D. C., Faul, A., Mackie, E., & Pauthenet, E. (2021)
1037	November). Defining Southern Ocean fronts using unsupervised classi-
1038	fication. Ocean Science, 17(6), 1545–1562. Retrieved 2023-01-16, from
1039	https://os.copernicus.org/articles/17/1545/2021/ doi: 10.5194/
1040	os-17-1545-2021
10/1	Thompson A F & Sallée J-B (2012 June) Jets and Topography: Jet Tran-
1041	sitions and the Impact on Transport in the Antarctic Circumpolar Current
1042	Journal of Physical Oceanoaranhy 42(6) 956–972 Betrieved 2023-06-27
1045	from http://journals_ametsoc_org/doi/10_1175/IPO-D-11-0135_1doi:
1045	10 1175/JPO-D-11-0135 1
1045	Thompson K B Lazier I B N & Taylor B (1086) Wind-forced changes in
1040	Labrador Current transport <i>Journal of Geophysical Research</i> 91(C12) 14261
1047	doi: 10.1029/ic091ic12p14261
1040	Tiira I la Moisseev D (2020) Unsupervised classification of vertical profiles
1049	of dual polarization radar variables Atmospheric Measurement Techniques
1050	13(3) 1227-1241 doi: 10.5104/2mt-12-1227-2020
1051	Townsond D W Pottigrow N P Thomas M A Neary M C Magilliauddy
1052	D. I. & Donnell I. O. (2015). Water masses and nutrient fluxes to the Culf of
1053	Maine Lowmal of Marine Basarah 72, 03, 122, doi: 10.1038/141548a0
1054	Malle E Criffica C M Abamathan D Adama T D Darlaff D Diag
1055	tach A Zilta I D (2018) Lagrangian according to the fundament
1056	toli, A., Zika, J. D. (2018). Lagrangian ocean analysis: Fundamen-
1057	tals and practices. $Ocean Model(My, 121 (November 2017), 49-75.$ doi: 10.1016/j. accmod.2017.11.008
1058	Wilmäe D. Tengrik T. & Seeman T. (2012). Impact of herizoptal oddy diffugiy
1059	ity on Lormon statistics for coastal pollution from a major maning fairman
1060	Try on Lagrangian statistics for coastal ponution from a major marine fairway
1061	Madalling Concern (IONCMOD) in Breat France 21 22 Mar 2012
1062	Modelling Group (JONSMOD) in Brest, France 21-23 May 2012. Ocean Dy -
1063	namics, b3(5), 589-597. doi: 10.1007/s10236-013-0615-3
1064	Whitney, N. M., Wanamaker, A. D., Ummenhofer, C. C., Cresswell-clay, N., Kreutz,
1065	K. J., & Johnson, B. J. (2022). Rapid 20th century warming reverses 900-year
1066	cooling in the Gulf of Maine. Nature Communications Earth & Environment,
1067	3(179), 1-15. doi: $10.1038/s43247-022-00504-8$
1068	Wichmann, D., Kehl, C., Dijkstra, H. A., & Van Sebille, E. (2021). Ordering of
1069	trajectories reveals hierarchical finite-time coherent sets in Lagrangian par-
1070	ticle data: Detecting Agulhas rings in the South Atlantic Ocean. Nonlinear
1071	Processes in Geophysics, 28(1), 43–59. doi: 10.5194/npg-28-43-2021
1072	Williams, J., Tremblay, B., Newton, R., & Allard, R. (2016). Dynamic precondi-
1073	tioning of the minimum september sea-ice extent. Journal of Climate, $29(16)$,

¹⁰⁷⁴ 5879–5891. doi: 10.1175/JCLI-D-15-0515.1

Figure 1.



Figure 2.

DATASET PREPARATION



PRE-PROCESSING

PROCESSING

30 clusters

POST-PROCESSING

7 pathway categories



Figure 3.



Figure 4.



Retroflected Westward Labrador Sea

Labrador Shelf Southward Belle Isle Number of particles

10²

Figure 5.



Figure 6.



Figure 7.



Figure 8.





Figure 9.





Figure 10.



Figure 11.



Figure 12.

Transect at 50W



Figure 13.


Figure A1.



Figure A2.



















































Figure A3.

