

Investigating Exhumation of the High Tatra Mountains: Implications for the Western Carpathians, Slovakia by Zircon and Apatite (U-Th)/He Thermochronometry

Daniel Campos¹, Elizabeth Catlos², Milan Kohut³, Igor Broska³, and Daniel F Stockli¹

¹Dept. Geological Sciences, University of Texas at Austin

²Affiliation not available

³Earth Science Institute of the Slovak Academy of Sciences, Bratislava, Slovakia

August 4, 2023

Abstract

The Carpathian Mountains form the large collisional orocline stretching from Vienna, Austria to Bucharest, Romania. The Western and Inner Carpathians include the High Tatra mountains, which exhibit the highest elevation peaks of the entire mountain belt. Here we studied the exhumation history of an area near Gerlachovský štít, the topographically highest point of the High Tatras. Granitoid samples from different elevations were collected and analyzed for apatite (U-Th)/He (n=12; 5-6 aliquots) and zircon (U-Th)/He ages (n=22; 2-4 aliquots). In addition, apatite U-Pb dating by LA-ICPMS was conducted to complement existing zircon U-Pb dates to track the evolution of the High Tatra Mountains from the onset of magmatism during the Variscan orogeny. The (U-Th)/He apatite ages show a general increase from 9.6 ± 0.6 Ma to 31.9 ± 2.0 Ma to from lower to higher elevations. The zircon (U-Th)/He ages are more scattered and range from 13.5 ± 1.1 Ma to 47.8 ± 3.9 Ma. These reported ages agree with published low-temperature thermochronometric results. However, the apparent average exhumation rates for zircon and apatite (U-Th)/He data derived from the age-to-elevation profile near Gerlachovský štít are inconsistent with a proposed rapid early Miocene exhumation pulse. Apatite U-Pb ages obtained in this study are between 337.61 ± 2.21 Ma and 372.74 ± 3.09 Ma. These ages agree with previously reported zircon dates from the same or nearby samples. This observation is indicative of rapid cooling of the granitoids following crystallization. However, the greatest variance in both data sets were observed from samples collected near the sub-Tatra fault and along the Ružbachy fault. This observation was used to confine regions about these major structures that have distinct exhumation records. The results of the (U-Th)/He ages captures both pre- and post-Miocene slow cooling interrupted by early Miocene tectonic unroofing. Overall, these results are used to outline the earliest tectonic history of the High Tatra Mountains until the onset of more recent exhumation and impacts our understanding of the origin and development of this section of the arcuate mountain belt.

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/368536814>

Investigating Exhumation of the High Tatra Mountains: Implications for the Western Carpathians, Slovakia by Zircon and Apatite (U–Th)/He Thermochronometry

Conference Paper · September 2022

CITATIONS

3

READS

34

5 authors, including:



Daniel Campos

University of Texas at Austin

8 PUBLICATIONS 5 CITATIONS

[SEE PROFILE](#)



Elizabeth J. Catlos

University of Texas at Austin

150 PUBLICATIONS 3,878 CITATIONS

[SEE PROFILE](#)



Milan Kohut

Earth Science Institute, Slovak Academy Sciences, Bratislava, Slovakia

124 PUBLICATIONS 1,532 CITATIONS

[SEE PROFILE](#)



Daniel F. Stockli

University of Texas at Austin

727 PUBLICATIONS 16,866 CITATIONS

[SEE PROFILE](#)

Some of the authors of this publication are also working on these related projects:



UHP metamorphic rocks e.g. Himalaya [View project](#)



Arctic Tectonics [View project](#)

Investigating Exhumation of the High Tatra Mountains: Implications for the Western Carpathians, Slovakia by Zircon and Apatite (U-Th)/He Thermochronometry

Daniel S. Campos¹, Elizabeth J. Catlos¹, Milan Kohút², Igor Broska², Daniel F. Stockli¹

¹ Department of Geological Sciences, Jackson School of Geosciences, The University of Texas at Austin
Austin, Texas, United States of America

² Earth Science Institute of the Slovak Academy of Sciences, Dúbravská cesta 9, 840 05
Bratislava, Slovak Republic

The Carpathian Mountains form the large collisional orocline stretching from Vienna, Austria to Bucharest, Romania. The Western and Inner Carpathians include the High Tatra mountains, which exhibit the highest elevation peaks of the entire mountain belt. Here we studied the exhumation history of an area near Gerlachovský štít, the topographically highest point of the High Tatras. Granitoid samples from different elevations were collected and analyzed for apatite (U-Th)/He (n=12; 5-6 aliquots) and zircon (U-Th)/He ages (n=22; 2-4 aliquots). In addition, apatite U-Pb dating by LA-ICPMS was conducted to complement existing zircon U-Pb dates to track the evolution of the High Tatra Mountains from the onset of magmatism during the Variscan orogeny. The (U-Th)/He apatite ages show a general increase from 9.6 ± 0.6 Ma to 31.9 ± 2.0 Ma to from lower to higher elevations. The zircon (U-Th)/He ages are more scattered and range from 13.5 ± 1.1 Ma to 47.8 ± 3.9 Ma. These reported ages agree with published low-temperature thermochronometric results. However, the apparent average exhumation rates for zircon and apatite (U-Th)/He data derived from the age-to-elevation profile near Gerlachovský štít are inconsistent with a proposed rapid early Miocene exhumation pulse. Apatite U-Pb ages obtained in this study are between 337.61 ± 2.21 Ma and 372.74 ± 3.09 Ma. These ages agree with previously reported zircon dates from the same or nearby samples. This observation is indicative of rapid cooling of the granitoids following crystallization. However, the greatest variance in both data sets were observed from samples collected near the sub-Tatra fault and along the Ružbachy fault. This observation was used to confine regions about these major structures that have distinct exhumation records. The results of the (U-Th)/He ages captures both pre- and post-Miocene slow cooling interrupted by early Miocene tectonic unroofing. Overall, these results are used to outline the earliest tectonic history of the High Tatra Mountains until the onset of more recent exhumation and impacts our understanding of the origin and development of this section of the arcuate mountain belt.

Acknowledgements. This work was supported by the US National Science Foundation #1460050 and Slovak Research and Development Agency APVV-18-0107 and VEGA 2/0075/20. Thank you to the University of Texas Chronology Laboratory, the Stockli Group, GeoFORCE Texas, and the Center for Russian and Eastern European Studies.