

Preliminary Analysis of a Retrogressive Thaw Slump Multi-Source Database for the West Siberian Arctic

Nina Nesterova^{1,2}, Ilya Tarasevich^{3,4}, Marina Leibman^{3,4}, Alexander Kizyakov⁵, Ingmar Nitze¹, and Guido Grosse^{1,2}

¹Alfred-Wegener-Institute for Polar and Marine Research

²University of Potsdam, Institute of Geosciences

³University of Tyumen

⁴Earth Cryosphere Institute

⁵Lomonosov Moscow State University

December 27, 2023

Preliminary Analysis of a Retrogressive Thaw Slump Multi-Source Database for the West Siberian Arctic

Nina Nesterova^{1,2}, Ilya Tarasevich^{3,4}, Marina Leibman^{3,4}, Alexander Kizyakov⁵,
Ingmar Nitze¹, Guido Grosse^{1,2}

- 1. Alfred-Wegener-Institute for Polar and Marine Research, Potsdam, Germany
- 2. University of Potsdam, Institute of Geosciences, Potsdam, Germany
- 3. University of Tyumen, Tyumen, Russia
- 4. Earth Cryosphere Institute, Tyumen, Russia
- 5. Lomonosov Moscow State University, Moscow, Russia



nina.nesterova@awi.de

Background & Study area

Retrogressive thaw slump (RTS) is a cryogenic landform due to ice-rich permafrost thawing or massive ground ice melting:

- Significantly altering the environment: vegetation, topography and carbon emissions
- Vulnerable to the temperature fluctuations
- Polycyclic nature: active or stabilized (or ancient)
- Complex features: multiage zones within one landform

West Siberian Arctic: Continuous permafrost + tabular massive ground ice close to the surface → widespread RTS occurrence




The aim

Further detailed investigation of RTS occurrence and environmental factors

Materials & Methods

- Mapped RTS points by Nesterova et al. 2021, Nitze et al., 2018 and Yang et al., 2023
- Manual collection of RTS points based on: ESRI satellite base map, Google map satellite, Yandex map satellite
- Classification

How does RTS look in West Siberia?



Results

- 4390 points collected
- 3150 are morphologically Thermocirques
 - 828 combined morphologies and only 160 Thermoterraces
 - 2139 complex and 1999 single forms

Most common RTS: single thermocirque at the lakeshore (with thermokarst subsidence and thermoerosion)



AI thoughts on RTS in West Siberia

Outlook

- In relation to data on:
- Geology
 - Ground ice content
 - Climate
 - Landcover
 - Other environmental parameters

More figures, statistics & References



Source: OpenAI