

# The Past and Future of the Fisheries and Marine Ecosystem Model Intercomparison Project

Camilla Novaglio<sup>1</sup>, Andrea Bryndum-Buchholz<sup>2</sup>, Derek Tittensor<sup>3</sup>, Tyler Eddy<sup>4</sup>, Heike K Lotze<sup>3</sup>, Cheryl Shannon Harrison<sup>5</sup>, Ryan Heneghan<sup>4</sup>, Olivier Maury<sup>6</sup>, Kelly Ortega-Cisneros<sup>7</sup>, Colleen M Petrik<sup>8</sup>, Kelsey E Roberts<sup>9</sup>, and Julia L. Blanchard<sup>1</sup>

<sup>1</sup>University of Tasmania

<sup>2</sup>Centre for Fisheries Ecosystems Research, Fisheries and Marine Institute, Memorial University of Newfoundland

<sup>3</sup>Department of Biology, Dalhousie University, 1355 Oxford Street, Halifax, NS B3H 4R2, Canada

<sup>4</sup>Unknown

<sup>5</sup>Louisiana State University

<sup>6</sup>Institut de Recherche pour le Développement

<sup>7</sup>Department of Biological Sciences, University of Cape Town, Cape Town, South Africa

<sup>8</sup>UC San Diego

<sup>9</sup>Department of Ocean and Coastal Science and Center for Computation and Technology, Louisiana State University, Baton Rouge, LA, USA

January 16, 2024

## Abstract

Climate-driven ecosystem changes are increasingly affecting the world's ocean ecosystems, necessitating urgent guidance on adaptation strategies to limit or prevent catastrophic impacts. The Fisheries and Marine Ecosystem Model Intercomparison Project (FishMIP) is a network and framework that provides standardised ensemble projections of the impacts of climate change and fisheries on ocean life and the benefits that it provides to people through fisheries. Since its official launch in 2013 as a small, self-organised project within the larger Inter-Sectoral Impact Model Intercomparison Project, the FishMIP community has grown substantially and contributed to key international policy processes, such as the IPCC AR5 and AR6, and the IPBES Global Biodiversity Assessment. While not without challenges, particularly around comparing heterogeneous ecosystem models, integrating fisheries scenarios, and standardising regional-scale ecosystem models, FishMIP outputs are now being used across a variety of applications (e.g., climate change targets, fisheries management, marine conservation, Sustainable Development Goals). Over the next decade, FishMIP will focus on improving ecosystem model ensembles to provide more robust and policy-relevant projections for different regions of the world under multiple climate and societal change scenarios, and continue to be open to a broad spectrum of marine ecosystem models and modellers. FishMIP also intends to enhance leadership diversity and capacity-building to improve representation of early- and mid-career researchers from under-represented countries and ocean regions. As we look ahead, FishMIP aims to continue enhancing our understanding of how marine life and its contributions to people may change over the coming century at both global and regional scales.

## Hosted file

982925\_0\_art\_file\_11715640\_s61444.docx available at <https://authorea.com/users/364325/articles/697849-the-past-and-future-of-the-fisheries-and-marine-ecosystem-model->

[intercomparison-project](#)

**Hosted file**

982925\_0\_supp\_11715641\_s61444.docx available at <https://authorea.com/users/364325/articles/697849-the-past-and-future-of-the-fisheries-and-marine-ecosystem-model-intercomparison-project>

**Hosted file**

982925\_0\_supp\_11715643\_s61444.docx available at <https://authorea.com/users/364325/articles/697849-the-past-and-future-of-the-fisheries-and-marine-ecosystem-model-intercomparison-project>

# The Past and Future of the Fisheries and Marine Ecosystem Model Intercomparison Project

Camilla Novaglio<sup>1,2\*</sup>, Andrea Bryndum-Buchholz<sup>3\*</sup>, Derek P. Tittensor<sup>4</sup>, Tyler D. Eddy<sup>3</sup>, Heike K. Lotze<sup>4</sup>, Cheryl S. Harrison<sup>5</sup>, Ryan F. Heneghan<sup>6</sup>, Olivier Maury<sup>7</sup>, Kelly Ortega-Cisneros<sup>8</sup>, Colleen M. Petrik<sup>9</sup>, Kelsey E. Roberts<sup>5</sup>, Julia L. Blanchard<sup>1,2</sup>

<sup>1</sup>Institute for Marine and Antarctic Studies, University of Tasmania, Australia.

<sup>2</sup>Centre for Marine Socioecology, University of Tasmania, Australia.

<sup>3</sup>Centre for Fisheries Ecosystems Research, Fisheries and Marine Institute, Memorial University of Newfoundland, St. John's, NL, Canada

<sup>4</sup>Department of Biology, Dalhousie University, 1355 Oxford Street, Halifax, NS B3H 4R2, Canada

<sup>5</sup>Department of Ocean and Coastal Science and Center for Computation and Technology, Louisiana State University, Baton Rouge, LA, USA

<sup>6</sup>School of Science, Technology and Engineering, University of the Sunshine Coast, Petrie, Australia

<sup>7</sup>IRD, Univ. Montpellier, Ifremer, CNRS, INRAE, MARBEC, Sète France

<sup>8</sup>Department of Biological Sciences, University of Cape Town, Cape Town, South Africa

<sup>9</sup>Scripps Institution of Oceanography, University of California San Diego, CA, USA

\*Corresponding authors: Camilla Novaglio ([camilla.novaglio@gmail.com](mailto:camilla.novaglio@gmail.com); ORCID 0000-0003-3681-1377) & Andrea Bryndum-Buchholz ([andrea.buchholz@mi.mun.ca](mailto:andrea.buchholz@mi.mun.ca); ORCID 0000-0002-7635-7845)

<sup>∞</sup>Contributed equally to this manuscript

## Key Points:

- There is an urgent need for policy to develop strategies to adapt to the impacts of climate change on ecosystems and their services
- The Fisheries and Marine Ecosystem Model Intercomparison Project has contributed understanding of climate impacts on marine ecosystems
- The next 10 years will see the FishMIP improved ensemble model pushing the boundaries of the field and increasing outputs policy-relevance

## Abstract

Climate-driven ecosystem changes are increasingly affecting the world's ocean ecosystems, necessitating urgent guidance on adaptation strategies to limit or prevent catastrophic impacts.

The Fisheries and Marine Ecosystem Model Intercomparison Project (FishMIP) is a network and framework that provides standardised ensemble projections of the impacts of climate change and

36 fisheries on ocean life and the benefits that it provides to people through fisheries. Since its  
37 official launch in 2013 as a small, self-organised project within the larger Inter-Sectoral Impact  
38 Model Intercomparison Project, the FishMIP community has grown substantially and contributed  
39 to key international policy processes, such as the IPCC AR5 and AR6, and the IPBES Global  
40 Biodiversity Assessment. While not without challenges, particularly around comparing  
41 heterogeneous ecosystem models, integrating fisheries scenarios, and standardising regional-  
42 scale ecosystem models, FishMIP outputs are now being used across a variety of applications  
43 (e.g., climate change targets, fisheries management, marine conservation, Sustainable  
44 Development Goals). Over the next decade, FishMIP will focus on improving ecosystem model  
45 ensembles to provide more robust and policy-relevant projections for different regions of the  
46 world under multiple climate and societal change scenarios, and continue to be open to a broad  
47 spectrum of marine ecosystem models and modellers. FishMIP also intends to enhance  
48 leadership diversity and capacity-building to improve representation of early- and mid-career  
49 researchers from under-represented countries and ocean regions. As we look ahead, FishMIP  
50 aims to continue enhancing our understanding of how marine life and its contributions to people  
51 may change over the coming century at both global and regional scales.

## 52 **1 Introduction**

53 In 2013, the Fisheries and Marine Ecosystem Model Intercomparison Project (Fish-MIP -  
54 [www.fishmip.org](http://www.fishmip.org)) was officially launched at the 1<sup>st</sup> Inter-Sectoral Impact Model  
55 Intercomparison Project (ISIMIP) cross-sectoral workshop in Potsdam, Germany. This launch  
56 filled a crucial gap in ISIMIP's mission "*to improve global and regional risk management by*  
57 *advancing knowledge of the risks of climate change through integrating climate impacts across*

58 *sectors and scales in a multi-impact model framework*” (ISIMIP, n.d.), which until then had only  
59 included terrestrial sectors and lacked contributions from the marine realm.

60 FishMIP tackled this gap through the development of an ensemble modelling framework to  
61 quantitatively assess uncertainties across marine ecosystem models and, further, to contribute to  
62 a multi-sectoral, multi-scale assessment of climate change impacts. This was and is a novel  
63 approach within the marine ecosystem modelling world, as most global studies of climate  
64 impacts on marine ecosystems were using single marine ecosystem model approaches of  
65 coupled-biophysical models (e.g., Cheung et al, 2011, Blanchard et al. 2012, Barange et al. 2014,  
66 Lefort et al. 2015). Using single marine ecosystem models, even if forced by multiple earth-  
67 system models (ESMs), limits our ability to quantify and understand the sources and range of  
68 uncertainty associated with the different ways ecosystems and fisheries have been  
69 conceptualised, mathematically formulated and computationally implemented in models, as there  
70 are very different interpretations of how best to integrate ecological processes into modelling  
71 frameworks (Tittensor et al. 2018, Lotze et al. 2019, Heneghan et al. 2021). Today, FishMIP  
72 comprises 100+ marine ecosystem and climate-impact modellers and contributors from around  
73 the world, aiming to collectively fulfil its (still unchanged) mission: *“to bring together diverse  
74 marine ecosystem models to help better understand and project long-term impacts of climate  
75 change on fisheries and marine ecosystems, and to use our findings to help inform policy”*  
76 ([www.fishmip.org](http://www.fishmip.org)). Specifically, FishMIP aims to answer questions around the future of fish and  
77 fisheries, food security, marine biodiversity and marine ecosystem functioning.


78 FishMIP’s ensemble modelling framework combines simulations from multiple marine  
79 ecosystem models, all forced by standardised inputs and scenarios, as defined by a specific  
80 project protocol (Tittensor et al. 2018). Such a standardised approach is necessary to make the

81 results comparable and to be able to calculate an ensemble mean and the variation around it. To  
82 develop a simulation protocol that works for many different models, and doing so in a  
83 collaborative, open way, involves many painstaking steps, discussions, and coordination among  
84 climate and ecosystem modellers. The first steps towards the FishMIP 1.0 protocol were taken in  
85 2014 and published in 2018 (Tittensor et al. 2018), followed by publishing the first ensemble  
86 results of six global marine ecosystem models in 2019 (Lotze et al. 2019).

87 The FishMIP 1.0 protocol established the foundational framework for FishMIP modelling  
88 efforts (Tittensor et al. 2018). It was specifically designed to support the Intergovernmental  
89 Panel on Climate Change (IPCC) and focussed on assessing climate change impacts on marine  
90 ecosystems by using heterogeneous marine ecosystem models forced by standardised outputs  
91 from two Earth System Models (ESMs), GFDL-ESM2M and IPSL-CM5A-LR, and four  
92 Representative Concentration Pathways (RCPs) provided by the Coupled Model Intercomparison  
93 Project Phase 5 and 6 (CMIP5 and CMIP6; <https://esgf-node.llnl.gov/search/cmip5/>; Bopp et al.  
94 2013), and standardised to a common 1° resolution global grid defined by ISIMIP. This protocol  
95 will be further developed into the FishMIP 2.0 protocol, which focuses on both climate change  
96 and fishing impacts, and extended to consider topics, such as food security, of interest to other  
97 policy bodies, including the Food and Agriculture Organization (FAO). Currently, FishMIP has  
98 achieved several major milestones and developments along the way that have contributed to  
99 scientific understanding and policy applications (Figure 1).

100 In this review, we chart the development, progress, and applications of FishMIP over the  
101 past decade, with the overarching aim of asking how far we have come in meeting the above  
102 mission and what future directions are needed to better deliver policy support at a time where  
103 rapid and robust answers are needed to guide strategies to reduce the impacts of climate-driven


104 changes in marine life and the resources it provides. We synthesise how FishMIP results have  
105 helped address key policy questions both within the marine ecosystems and fisheries sector and  
106 in cross-sectoral studies, as well as tracking the research impact of selected key papers. Finally,  
107 we highlight the path ahead over the next decade of FishMIP 2.0 (Blanchard et al., *this issue*).



**Figure 1.** Timeline of FishMIP development and milestones since 2013.

110 **2 Growing applications and impact**


111 Several key FishMIP papers have shown a growing level of research impact through time, in  
112 terms of number of citations and policy uptake. They have helped answer questions about key  
113 topics in the FishMIP mission: fish and fisheries, seafood supply, marine biodiversity and marine  
114 ecosystem functioning, thus supporting policies related to these key areas. By analysing the  
115 number of citations since 2016, derived from the FishMIP Google Scholar  
116 (<https://tinyurl.com/usw9e92p>), we identified six key FishMIP papers with >100 cumulative  
117 citations each (Figure 2).



118  
119 **Figure 2.** Number of citations per year since 2017 for six key FishMIP papers (cumulative  
120 citations >100), derived from the FishMIP Google Scholar <https://tinyurl.com/usw9e92p>.

121 Based on a cross-checked Altmetric analysis, we determined the policy uptake of FishMIP  
122 publications and which policy themes were addressed. Our cross-check removed duplicates in  
123 the Altmetric records and added missing reports that were not captured by Altmetric at the time

124 of analysis (Table S1). Over time, FishMIP publications have addressed three main, often  
 125 intersecting, policy categories: 1) climate change targets, and the importance of meeting these  
 126 targets for fish, fisheries and their management; 2) cross sectoral trade-offs in Sustainable  
 127 Development Goals (SDGs), including future challenges for seafood supply and food security;  
 128 and 3) marine biodiversity conservation and ecosystem functioning (Figure 3). The category of  
 129 “Other”, included policy reports that did not fit into the main categories identified. FishMIP  
 130 publications have led to the use and the uptake of FishMIP output into national and international  
 131 policy documents, including reports by the IPCC, IPBES, FAO, World Bank and UN (Table S2).



132 **Figure 3.** Number of policy categories addressed in policy reports that use FishMIP data or cite  
 133 published FishMIP and ISIMIP publications between 2017-2021. Note that some policy reports  
 134 address multiple themes. See the number and title of policy reports citing FishMIP and ISIMIP  
 135 publications in the Supplemental Material, Table S1, S2.  
 136

## 137 **2.1 Climate change targets**

138 A primary objective has been, and continues to be, to provide future projections to the IPCC  
139 as evidence of the benefits of limiting climate change. This includes the first synthesis and inter-  
140 comparison of standardised climate-change driven marine ecosystem model outputs on global  
141 (Lotze et al. 2019) and regional scales (Bryndum-Buchholz et al. 2019), which were integrated  
142 into the IPCC Special Report for the Ocean and Cryosphere in a Changing Climate (IPCC  
143 2019b). These works showed consistent and exacerbated future declines in ocean biomass under  
144 the high emissions scenario (RCP8.5) compared to the low emissions scenario (RCP2.6), with  
145 substantial geographical variation, and thus highlighted the benefits of climate mitigation  
146 actions.

147 Large uncertainties across the ensemble projections within the global FishMIP CMIP5  
148 results (Lotze et al. 2019) and for key regions (Bryndum-Buchholz et al. 2019) led to a targeted  
149 simulation experiment in 2018-2019 to tease apart the effects of different climate drivers on  
150 individual marine ecosystem model projections of future fish biomass change (Heneghan et al.  
151 2021). This analysis revealed that the nature of lower trophic level coupling in models was a key  
152 source of inter-model uncertainty. Despite CMIP6-forced projections revealing a narrower  
153 spread of ecosystem projections, with a more pessimistic outlook relative to CMIP5 (due to both  
154 higher climate sensitivity of FishMIP models and higher warming in CMIP6) that was integrated  
155 into the IPCC's 6th assessment report (Tittensor et al. 2021; IPCC 2022), All four FishMIP  
156 papers led to recommendations for improved understanding of uncertainty and a need for more  
157 robust regional-scale projections.

158 A further investigation revealed that large uncertainties and limited quantitative and  
159 standardised ensemble model validation hamper the robustness of modelling outputs and may

160 have restricted the use of these outputs in management and policy contexts (Steenbeek et al.  
161 2021). Among the clear gaps in improving confidence in and robustness of marine ecosystem  
162 model projections are: 1) the limited cross-ecosystem model validation against historical data  
163 (Heneghan et al. 2021, Rynne et al. *this issue*); and 2) the high computational cost of marine  
164 ecosystem model simulations, which makes it difficult to develop large ensemble simulations to  
165 conduct systematic sensitivity analysis and parameter estimation (Steenbeek et al. 2021).  
166 Solutions to these challenges are currently being developed by the FishMIP community,  
167 including the collaborative development of a model validation framework (Rynne et al. *this*  
168 *issue*), distributed computation to tackle the high computational cost of marine ecosystem model  
169 simulations (Steenbeek et al.2021), and collaboration with the Earth System Grid Federation  
170 (ESGF2) to better validate marine ecosystem drivers in CMIP6 (Fu et al. 2022).

## 171 **2.2 Climate-resilient fisheries management**

172 Global projections of ocean biomass have been used to assess fisheries management  
173 challenges in the Northwest Atlantic under different climate change scenarios (Bryndum-  
174 Buchholz et al. 2020a, Lotze et al. 2022). Here, the FishMIP ensemble revealed regionally  
175 disparate biomass changes, with high projected decreases overlapping with historical and current  
176 areas of high fisheries landings, while areas with lower historical landings, such as Arctic and  
177 sub-Arctic areas, showed large biomass increases, albeit with large uncertainties (Bryndum-  
178 Buchholz et al. 2020a). For Australia, more specifically, Pethybridge et al. (2020) explored  
179 contrasting futures for national, large-scale fisheries stocks, using an ensemble of regional and  
180 two of the global FishMIP models (DBEM, DBPM). Across Australia, demersal fisheries were  
181 projected to experience larger climate-related impacts than pelagic fisheries, notably Australia's  
182 invertebrate fisheries. Using the ensemble approach, this study highlighted priorities for fisheries

183 specific, proactive, and flexible management systems that effectively account for climate change  
184 impacts.

185 The call by Pethybridge et al. (2020) for proactive and flexible fisheries management  
186 adaptation in Australia was complemented by a comprehensive review of the implementation of  
187 climate change adaptation in fisheries management policy and legislation (Bryndum-Buchholz et  
188 al. 2021). Here, global FishMIP projections of ocean biomass changes within 11 Exclusive  
189 Economic Zones (EEZs) were used to illustrate the impact of climate change on marine  
190 ecosystems and fisheries, and highlighted the urgency of developing adaptation plans by the  
191 respective nations responsible for these EEZs (Bryndum-Buchholz et al. 2021). This was critical  
192 information, since across these EEZs, none of the reviewed fisheries management policies and  
193 legislation explicitly addressed climate change impacts or mandated the integration of those  
194 impacts into stock assessments or decision making. The findings of this study were highlighted  
195 in the IPCC 6th assessment report (IPCC 2022). Understanding climate change impacts on  
196 marine ecosystems and fisheries governance is imperative for ensuring that they remain  
197 productive and sustainable in a changing ocean.

198 While these FishMIP studies address questions around climate change impacts on marine  
199 ecosystems and fisheries management, projections of climate impacts did not account for  
200 impacts due to fishing. At the time of these analyses, a standardised, spatially and temporally  
201 explicit, representation of future fishing scenarios (i.e. future evolution of fishing effort,  
202 mortality or exploitation rate) was unavailable. Developing the Ocean System Pathways (OSPs),  
203 a set of standardised fishing scenarios and the associated modelling framework required to  
204 simulate them (Maury et al, *this issue*), has been one of FishMIP's key challenges and has led to  
205 the development of the FishMIP 2.0 protocol (Blanchard et al., *this issue*). This process

206 commenced in 2021 following extensive community consultation on research and policy  
207 priorities. It involves defining an expanded set of simulations directly targeted at specific policy  
208 processes that includes dynamic fisheries and socioeconomic scenarios (Maury et al., *this issue*)  
209 using models that have gone through a validation and benchmarking stage (Blanchard et al., *this*  
210 *issue*).

### 211 **2.3 Food security and cross sectoral trade-offs**

212 A preliminary synthesis of CMIP5 projections from FishMIP and the Agriculture Model  
213 Intercomparison Project (AgMIP) provided the first cross-sectoral ensemble assessment of joint  
214 climate change impacts across land and sea (Blanchard et al. 2017), highlighting trade-offs for  
215 food security in the Special Report on Climate Change and Land (IPCC 2019a). This assessment  
216 showed that the ceiling of food production from both land and sea is expected to face declines in  
217 most countries (Blanchard et al. 2017). Other studies using projections to assess cross-sectoral  
218 impacts have followed and shown that, under a high emissions scenario, most countries are  
219 expected to face losses in both agriculture and fisheries production, while under a low emissions  
220 scenario they would experience gains in both sectors, thus advocating for prompt climate  
221 mitigation actions (Thiault et al. 2019).

222 Building on this work and using detailed household surveys enabled an assessment of future  
223 local impacts on agriculture and fisheries production for coastal communities of the Indo-Pacific  
224 (Cinner et al. 2022). This study showed that potential losses to fisheries are generally higher than  
225 those to agriculture, and that strong climate change mitigation could drastically reduce the risk of  
226 simultaneous losses in agriculture and fisheries production for most of the coastal communities  
227 considered. The socio-economic benefits of meeting climate mitigation targets were also  
228 highlighted in another study linking marine ecosystem model projections to a range of socio-

229 economic indicators, at national to global scales (Boyce et al. 2020). These benefits include  
230 preventing the widening of existing climate-driven equity gaps, in particular for nations that are  
231 heavily dependent on decreasing food resources and face increasingly poor nutritional status,  
232 wealth and ocean health (Boyce et al. 2020). This work was integrated into the 6th IPCC  
233 assessment report, emphasising socio-economic consequences in the context of climate change  
234 impacts and adaptation (IPCC 2022).

235 Looking ahead, as extreme climate events increase, the frequency and intensity of sudden  
236 losses to food production on land and in the sea are expected to increase, with negative  
237 repercussions for food supply and security, livelihoods and human well-being (Cottrell et al.  
238 2019). To prepare for such losses, we need marine ecosystem models that are capable of  
239 estimating the full extent of impacts of extreme climate events, and cross-sectoral assessments  
240 that make use of such modelling outputs to account for the complex linkages across food  
241 systems. A test of how well the current suite of models from multiple sectors can capture the  
242 impacts of an extreme climate event revealed that severe impacts on agriculture and ecosystem  
243 productivity are largely underestimated, and further highlighted the need to improve such  
244 models' ability and to work on cross-sectoral aspects (Schewe et al. 2019).

## 245 **2.4 Marine biodiversity conservation and ecosystem functioning**

246 The FishMIP-CMIP5 ensemble contributed to the 2019 Intergovernmental Science-Policy  
247 Platform on Biodiversity and Ecosystem Services (IPBES) global assessment of biodiversity and  
248 ecosystem services, by providing insights into future climate change impacts on marine  
249 biodiversity and ecosystem functioning, as well as the potential consequences for ecosystem  
250 services (Blanchard et al. 2017, Tittensor et al. 2018, Lotze et al. 2019, 2022, IPBES, 2019).

251 The enhanced FishMIP-CMIP6 model ensemble was applied to regional marine  
252 conservation questions (Bryndum-Buchholz et al. 2023a), ultimately generating knowledge for  
253 the United Nations Sustainable Development Goals (SDGs), particularly SDG 13 (climate  
254 action) and SDGs 14 (life below water). Here, projections of ocean biomass and key physical  
255 variables for Atlantic Canada were combined to identify climate refugia and hotspots and to  
256 evaluate the future ability of marine conservation areas to protect biodiversity. Results provide  
257 important long-term context for adaptation and for building climate-resilience into spatial marine  
258 conservation planning in the region.

259 The effects of climate change on ecosystem structure were recently assessed using the  
260 enhanced global FishMIP-CMIP6 ensemble (Guibourd de Luzinai et al. 2023). By analysing  
261 changes within marine food webs in the model ensemble, the study quantified the extent of  
262 trophic amplification (the larger decline in the biomass of higher trophic level organisms relative  
263 to the decrease in primary producer biomass) and highlighted a more complex response within  
264 food webs than anticipated due to disparate parameterization and structure of the individual  
265 FishMIP models. Rather than a consistent amplification, results showed how temperature  
266 changes can attenuate or even offset trophic amplification across marine ecosystems (Guibourd  
267 de Luzinai et al. 2023).

### 268 **3 Looking ahead with FishMIP 2.0**

269 With the Paris agreement climate change target, the associated Global Stocktake revision to  
270 come after the UNFCCC COP28 (<https://www.cop28.com/en/>) and the 2030 agenda for  
271 sustainable development (SDGs) imminent, delivering policy-relevant climate impact and  
272 adaptation science is of paramount importance across sectors and disciplines. For the ocean and  
273 its living marine resources, FishMIP model ensemble projections are used to support

274 vulnerability and climate risk studies and adaptation plans for different countries and regions of  
275 the world. To date, FishMIP has informed climate, biodiversity, fisheries, food security, and  
276 marine conservation sectors. FishMIP results can increasingly be used to inform current and  
277 future progress towards United Nations Sustainable Development Goals  
278 (<https://sdgs.un.org/goals>), specially: Climate Action (SDG 13), Life Below Water (SDG 14),  
279 Zero Hunger (SDG 2), Decent Work and Economic Growth (SDG 8), and Responsible  
280 Consumption & Production (SDG 12), and a range of Ocean Decade Challenges  
281 (<https://oceandecade.org/challenges/>) (Figure 4, Table S3). However, to successfully guide  
282 potential actions for informing progress towards the above goals, we highlight several focal areas  
283 for FishMIP 2.0 to focus on over the next decade.

284 We have developed five key areas for focal working groups (WGs), described below, that  
285 are designed to contribute to a range of SDGs and to provide scientific information needed to  
286 strengthen resilience and adaptive capacity to climate-related hazards, integrate climate change  
287 measures into policies and planning, and sustainably manage marine resources (Figure 4, Table  
288 S3). Each working group also addresses specific Ocean Decade Challenges, FAO Blue  
289 Transformation goals, as well as targets and goals of the Convention of Biological Diversity  
290 (CBD). Together, they will create research capacity to:

### 291 **3.1 WG1: Marine ecosystem model improvement to enhance policy relevance**

292 Previous studies have highlighted that projections of climate change impacts on marine  
293 ecosystems and fisheries have large uncertainties, particularly in regions where rapid changes are  
294 occurring, e.g., polar waters (Lotze et al. 2019; Tittensor et al. 2021). However, robust model  
295 ensembles of climate impact projections and their associated inter-model variability, on which  
296 reliable future sustainable pathways can be based, are needed to inform decision making. To be

297 reliable, these projections must be accompanied by appropriate estimates of uncertainty, where  
298 lower uncertainty means higher confidence in the direction, magnitude, and geographical pattern  
299 of change (Payne et al. 2016).

300 Improving model ensemble skills through appropriate calibration and validation of  
301 individual marine ecosystem models is one of the primary goals of current and future FishMIP  
302 efforts. This requires a stronger assessment of the ability of individual marine ecosystem models  
303 to capture past ecosystem states and environmental- and exploitation-driven changes, including  
304 the development of standardised datasets against which to evaluate historical model simulations  
305 (Blanchard et al. *this issue*) and standardised methodological frameworks for model skill  
306 evaluation (Rynne et al., *this issue*). The FishMIP Model Improvement Working Group (WG 1;  
307 Figure. 4) is developing tools to assess the reliability of marine ecosystem model ensembles  
308 through advanced skill assessment techniques (Novaglio et al., *this issue*; Rynne et al., *this issue*)  
309 using global and regional observational datasets, ranging from fisheries catches and effort to fish  
310 abundance and biomass surveys, including novel data streams (e.g., satellite data, eDNA). This  
311 working group will also develop tools to help build regional marine ecosystem model ensembles  
312 spanning a gradient of data-rich to data-poor regions.

313 In addition, confronting marine ecosystem models with observations helps to identify the  
314 processes that affect simulation outputs and the parameterisation that needs to be refined,  
315 ultimately leading to the improvement of marine ecosystem models (Heneghan et al. 2021).  
316 FishMIP 2.0 will assess the ability of the FishMIP ensemble to detect past ecosystem and fishery  
317 changes, including the attribution of changes to specific stressors. As the majority of fisheries  
318 catches are made in coastal waters, this is likely to require improved representation of coastal  
319 physical and biogeochemical processes in ESMs and increased reliability of projections for

320 coastal regions. This will in particular involve the use of higher-resolution ESM input data that  
321 better represent coastal enrichment processes such as upwelling and nutrient runoff from land-  
322 based human activities, sediment dynamics (Liu et al. 2021; Frieler et al. 2023), as well as  
323 regional downscaling of global ESMs (Drenkard et al. 2021, Jacox et al. 2020, Holt et al. 2017),  
324 and the use of standardised fishing effort for global and regional models (Rousseau et al., in  
325 press).

326 An improved FishMIP model ensemble, forced by improved ESMs, will contribute  
327 knowledge to major science-policy efforts (e.g. IPCC, IPBES), policy processes (e.g. FAO  
328 Committee on Fisheries, and Convention on Biological Diversity) and international targets, such  
329 as SDG 13 (Climate Action) and 14 (Life Below Water), as well as SDG targets focussing on  
330 increasing scientific knowledge and capacity to improve ocean health and effective climate  
331 change-related planning and managing (e.g., 13.3, 14.B, 14.A;  
332 [https://sdgs.un.org/goals/goal13#targets\\_and\\_indicators](https://sdgs.un.org/goals/goal13#targets_and_indicators),  
333 [https://sdgs.un.org/goals/goal14#targets\\_and\\_indicators](https://sdgs.un.org/goals/goal14#targets_and_indicators)). Likewise, improved marine ecosystem  
334 models and ESMs will help address Ocean Decade Challenges 8 (Create a digital representation  
335 of the Ocean), and 9 (Skill, knowledge and technologies for all; Table S3).

### 336 **3.2 WG2: Socio-economic scenarios to foster sustainable fisheries**

337 Billions of people depend on living marine resources and sustainable fisheries that are at risk  
338 due to climate change. The Socio-economic Scenarios Working Group (Fig. 4) will lead the  
339 development, implementation in models, testing and simulation of the OSPs, a set of SSP-  
340 consistent scenarios of the future of fisheries, including a modelling framework for embedding  
341 market and fleet dynamics into ecosystem models (Maury et al. *this issue*), that will, amongst

342 other objectives, help to address where adaptive fisheries management measures are most  
343 needed, and which measures can successfully lead to climate resilient fisheries.

344 Policymakers across multiple sectors need projections of marine biomass, fisheries catches  
345 and seafood availability under scenarios of climate and socio-economic changes that take into  
346 account human behaviour and choices in order to inform management decisions on scales  
347 ranging from national to global. In addition, these projections are needed to test new  
348 conservation and adaptation strategies that will be developed during the next decade of climate-  
349 impact research and implementation. With better-validated models that can simultaneously  
350 capture human impacts and impacts on humans, FishMIP will be in a position to explore  
351 spatially and temporally explicit socio-economic fisheries scenarios that are consistent with and  
352 extend the climate scenarios considered thus far (Maury et al., *this issue*). This requires the  
353 implementation of dynamic fishing, which simultaneously depends on economic, governance  
354 and management drivers as well as changing climatic conditions, into all FishMIP models, the  
355 translation of qualitative OSP storylines into quantitative driver pathways, and the design of  
356 modelling experiments aimed at exploring the potential range of climate and socio-economic  
357 impacts on the marine realm along with adaptation and mitigation options (Maury et al. *this*  
358 *issue*).

359 Following extensive community consultation, the Socio-economic Scenarios Working  
360 Group has already made progress on the implementation of the features described above, with  
361 work on informing climate resilient fisheries planned to continue over the next decade, with the  
362 goal of contributing to the knowledge base required for major, international sustainability goals  
363 (Figure 4, Table S3).

### 364 **3.3 WG3: Promote climate-resilient food-security**

365 Climate change and resource use have affected and will continue to affect the substantial  
366 contributions that aquatic food already makes to the diet and livelihoods of many nations (FAO.  
367 2020), highlighting the urgent need to strengthen the climate-resilience of aquatic food. To  
368 increase resilience and to meet the ever increasing demand for food (van Dijk et al. 2021,  
369 Costello et al. 2020), we need a better understanding of how marine ecosystems respond to  
370 perturbations, now and in the future, and of the linkages between land and sea food production  
371 systems. The Food Security Working Group (Figure. 4) will work on such topics. For instance,  
372 through well established cross-sectoral links with agriculture modellers (AgMIP), FishMIP will  
373 carry out the first ensemble model projection for aquaculture and combine agriculture,  
374 aquaculture, and fisheries projections to evaluate land-sea interactions and food security-  
375 biodiversity trade-offs. A larger set of indicators will be developed to simplify the integration of  
376 FishMIP results with those from other sectors, including AgMIP and the Food Model  
377 Intercomparison Project (FoodNut). Such integration would allow for answering questions on the  
378 sustainability of interconnected food systems of fisheries, aquaculture, and agriculture and their  
379 respective impacts on and vulnerabilities to changes in biodiversity and ecosystem functions and  
380 services, from regional to global scales.

381 A second core objective of this working group is to develop tools for improving the  
382 representation of coastal fisheries in key regions of the world and developing countries. Indeed, a  
383 clear policy and decision-making request to FishMIP is to strengthen the focus on regional and  
384 local scales, where projections are lacking (e.g. Tittensor et al. 2021). This means increasing the  
385 currently patchy coverage of regional ecosystem models, with the global south particularly  
386 underrepresented; focusing on the development of regional marine ecosystem model ensembles

387 that are often difficult to assemble for regional institutions; promoting the development of  
388 dynamically downscaled climate projections, as well as less labour-intensive statistically  
389 downscaled projections; and increasing efforts on the analysis of global model outputs by  
390 Exclusive Economic Zones (plus the High Seas). To increase the coverage of regional models  
391 and the diversity of ecosystem modellers, resources such as the Global South Climate Database  
392 that lists climate experts and their skills will be explored ([https://www.carbonbrief.org/global-](https://www.carbonbrief.org/global-south-climate-database/)  
393 [south-climate-database/](https://www.carbonbrief.org/global-south-climate-database/)) and a higher number of postgraduate students and postdoctoral fellows  
394 from underrepresented regions will be engaged. Focusing on improving spatial and temporal  
395 resolutions of regional modelling outputs is an ongoing exercise, which will gather momentum  
396 throughout the coming decade of FishMIP, actively addressing multiple SDGs, Ocean Decade  
397 Challenges, and Blue Transformation goals, such as SDG 2 (Zero Hunger), Ocean Challenge 3  
398 (Sustainably feed the global population), and Better Nutrition, Programme Priority Areas 1 to 4  
399 (Figure 4, Table S3).

#### 400 **3.4 WG4: Protect ecosystems and biodiversity**

401 Climate-adaptive marine ecosystem management has the potential to mitigate long-term  
402 effects of climate change and reduce biodiversity loss by restricting fisheries exploitation by  
403 space, time, species, and size. The Biodiversity Conservation Working Group (Figure. 4) will  
404 work to evaluate the extent to which Marine Protected Areas (MPAs) and other spatial  
405 conservation measures such as Other Effective Area-Based Conservation Measures (OECMs)  
406 can and will be able to contribute to marine ecosystem protection and restoration, in addition to  
407 exploring the impact of fisheries management decisions on marine biodiversity and ecosystem  
408 health. In particular, these modelling experiments will explore the role of MPAs and other area-

409 based fisheries management strategies in enhancing biodiversity, and strengthening ecosystems’  
410 recreational value.


411 To quantify the benefits of good management and trade-offs between climate mitigation and  
412 adaptation pathways that emphasise different ecological and socio-economic values of marine  
413 ecosystems, the range of biodiversity metrics that can be captured with FishMIP models will be  
414 extended. Because most global FishMIP models do not include species but rather focus on size  
415 classes or functional groups, and hence cannot easily provide species-specific biodiversity  
416 indicators, a possible way forward is the use of traits or functional diversity or empirical  
417 relationships to convert modelled biomass to species richness and thus to indirectly translate  
418 modelling outputs to biodiversity. An additional aspect of the work will be aligning biodiversity-  
419 focussed scenarios with those used by IPBES, in particular the Nature Futures Framework  
420 (Pereira et al. 2020; Maury et al. *this issue*). Outputs of the biodiversity-focussed scenarios can  
421 also be used to inform the newly adopted Biodiversity Beyond National Jurisdiction (BBNJ;  
422 Tiller & Mendenhall, 2023) treaty by exploring changing biodiversity on the high seas.

423 The Biodiversity Protection Working Group aims at contributing to the knowledge base of  
424 SDGs 13 and 14, in particular to targets for sustainable management and protection of marine  
425 and coastal ecosystems and the conservation of coastal and marine areas, consistent with national  
426 and international law (SDG 14.2, SDG 14.5; Figure 4, Tabel S3). This work is also consistent  
427 with Ocean Decade Challenge 2 (Protect and restore ecosystems and biodiversity) and 5 (Unlock  
428 ocean-based solutions to climate change; Figure 4, Table S3).

### 429 **3.5 WG 5: Assess intervention scenarios for climate change remediation**

430 Reducing emissions and limiting global warming to 1.5 degrees Celsius has been repeatedly  
431 shown to improve prospects for future global fisheries and ecosystems (e.g., Boyce et al. 2020).

432 Additionally, as climate targets are slow to be reached, a wide range of climate intervention  
433 approaches are being debated to modify the Earth’s radiation budget and remove greenhouse  
434 gases from the atmosphere (National Academies of Sciences, Engineering, and Medicine 2021,  
435 2022). The many consequences of these interventions on fisheries and marine ecosystems are  
436 largely unknown, potentially detrimental, and therefore crucial to understand given their  
437 importance to the global food supply. The Climate Change Mitigation Working Group (Figure 4)  
438 will explore the potential impacts of climate interventions on marine ecosystems, in an effort to  
439 add to the knowledge base around geoengineering impacts and hence inform decision- and  
440 policymaking with due consideration of potential impacts on ecosystem services, biodiversity,  
441 and food security in future climate scenarios. While FishMIP is already contributing to climate  
442 risk and vulnerability assessments (Cinner et al. 2022), this focused set of policy-relevant and  
443 management-driven analyses will allow a more in-depth consideration of climate intervention  
444 strategies, and increase scientific knowledge, develop research capacity and transfer marine  
445 technology to improve ocean health (SDG target 14.A) and increase community resilience to  
446 ocean hazards (Ocean Decade Challenge 6; Figure 4, Table S3).



447  
448  
449  
450  
451  
452  
453  
454  
455

**Figure 4.** Overview of FishMIP goals and their respective policy categories, the objectives of current and future FishMIP working groups (WGs), and how both address targets and goals of the Sustainable Development Goals (SDGs), the Ocean Decade Challenges, the Food and Agriculture Organization (FAO) Strategic Framework for Blue Transformation, and the targets (numbers) and goals (letters) of the Convention of Biological Diversity (CBDs). Please note that SDGs, Ocean Decade Challenge, FAO and CBDs goals for WG5 and WG2 are grouped as both WGs fall within the broader Fish & Fisheries topic. Specific explanation of each target and goal are in Table S3.

#### 456 **4 Key challenges and solutions**

457 Since its inception in 2013, FishMIP has faced and addressed important challenges.  
458 Discussion around these challenges and proposed solutions have been opportunities for learning  
459 and the lessons gained have often informed other modelling projects and teams, thus advancing  
460 marine ecosystem modelling more broadly. Most of these challenges are long-standing and  
461 require continuous work, while others have appeared only recently. Here is an overview of the  
462 main challenges:

463 **Relevant climate models that simulate and save all the necessary data for the marine**  
464 **ecosystem models:** Through the collaboration of FishMIP and CMIP (Ruane et al. 2016) we  
465 received improved representation of biochemical parameters in CMIP6 compared to CMIP5  
466 which helped advance FishMIP projections (Tittensor et al. 2021). However, the number of  
467 ESMs that simulate, save and can provide all the physical and biogeochemical variables that are  
468 required to drive marine ecosystem models is still limited. Thus, this collaborative process  
469 between climate and marine modellers needs to continue and deepen.

470 **Empirical data for model calibration and validation:** the ever-increasing expansion of  
471 global, observational datasets will provide more opportunities to better calibrate, constrain and  
472 validate both climate and marine ecosystem models.

473 **Improved ecosystem models:** Each marine ecosystem model has its own idiosyncrasies and  
474 biases, and is necessarily an incomplete representation of the complexity of marine ecosystems.  
475 Thus, incorporating into the ensemble and comparing the predictions of more marine ecosystem  
476 models that are based on different paradigms or reflect different ecosystem structures and  
477 processes is highly informative (Tittensor et al. 2018) and can help identify gaps in individual  
478 models and make progress in addressing them. There are plenty of ways to improve, for instance,

479 including marine ecosystem models that incorporate the potential for species acclimatisation and  
480 adaptation into the FishMIP ensemble would enable assessment of how this would affect  
481 projected future changes in distribution and abundance of marine animal biomass.

482       **Compare global with regional projections:** Regional ecosystem models may more  
483 accurately capture processes at management-relevant scales, both due to better resolution of  
484 physical and biogeochemical processes with finer regional ocean model resolution and use of  
485 regionally-tuned plankton models, and due to better representation of local ecosystem processes  
486 in regional marine ecosystem models. To address whether or why regional marine ecosystem  
487 models are more accurate and/or different from global models, systematic comparisons between  
488 global and regional model ensembles are needed, along with continued effort at regional  
489 downscaling of climate projections sensitive to the needs of ecosystem managers.

490       **Optimal level of complexity:** As FishMIP models evolve and new processes are being  
491 integrated to answer more complex questions, the ability to understand the effect of a change in a  
492 driver becomes more difficult. In the future, this difficulty will further increase when dynamic  
493 fishing is integrated in all FishMIP models. Increased validation of regional and global models  
494 and the implementation of detection and attribution analyses are expected to help in this regard.

495       **Work-flow preceding modelling experiments:** Efforts around the preparation of climate  
496 and fishing inputs have increased at every protocol iteration. This is due to the need to meet the  
497 requirements of an increasing set of models and to accommodate the different requests of global  
498 and regional models, with the latter often being forced, calibrated, and validated using high-  
499 resolution ocean model and fishing inputs that come from alternative sources and that are likely  
500 more appropriate for regional modelling than the standardised, global-scale FishMIP inputs.  
501 Solutions to this growing challenge include agreement on a downscaling method that takes

502 global-scale inputs and provides more meaningful regional products. To reduce bottlenecks, the  
503 FishMIP team is developing a series of tools that ease access, downloading and formatting of  
504 climate model input data and to support global and regional modelling ([www.fishmip.org](http://www.fishmip.org);  
505 <https://github.com/Fish-MIP>).

506 **Computational needs and quality:** Computational needs have become a limiting factor for  
507 some modelling teams as inputs increase in resolution, hence in volume, and experiments expand  
508 to answer multiple questions. Model improvement is not only related to model structure and the  
509 choice of parameters but also to its computational quality - i.e., how efficient the code is.  
510 Improving the model computational quality depends on the economic, personnel and  
511 computational resources available to each modelling team (Steenbeek et al. 2021, Steenbeek et  
512 al. *this issue*).

513 **Voluntary commitment:** Currently FishMIP relies on voluntary commitments from  
514 modellers and scientists. Ensuring the long-term longevity of this ever growing project requires  
515 setting up mechanisms that guarantee consistent funding similar to those supporting other  
516 relevant projects in this field, e.g., CMIP. As a first step to increase its visibility to potential  
517 funders, FishMIP applied to become a UN Decade Action, making FishMIP a UN-endorsed  
518 project directly contributing to the Ocean Decade vision of '*the science we need for the ocean we*  
519 *want*' (<https://oceandecade.org/decade-actions/>)

520 **Dissemination of outputs:** FishMIP outputs are becoming harder to analyse and make  
521 widely available as they grow in complexity and number. To ensure a continuous and increasing  
522 use of these outputs, it is paramount that interactive tools where outputs can be stored,  
523 summarised and downloaded continue to be developed and innovative solutions to be explored  
524 and adopted.

## 525 **5 Conclusion**

526 Over the past decade, FishMIP has brought together and engaged an international  
527 community of marine climate, fisheries and ecosystem scientists and modellers, who have  
528 contributed substantially to the field of climate-impact science in the marine realm, and informed  
529 key international climate and biodiversity science-policy efforts. Along the way, FishMIP has  
530 faced important challenges, most of them requiring continuous work, such as improving the  
531 robustness of future projections through standardised marine ecosystem model evaluation, the  
532 development of fishery scenarios accounting for the socio-economic factors that drive the  
533 evolution of fishing effort and markets in interaction with ecosystems, and providing regional  
534 and local model-based projections to inform policy-making. By addressing these challenges over  
535 the next decade, FishMIP can play an important role in the development of marine ecosystem  
536 models at all scales, and accelerate their progress. It can also be useful in providing the  
537 information that fisheries management and marine conservation need to address the issue of  
538 resilience to climate change and develop effective adaptation strategies. Likewise, it can  
539 contribute to the knowledge base to achieve the UN Sustainable Development Goals, particularly  
540 those addressing hunger and life below water. Over the next decade, FishMIP will address long-  
541 standing scientific and policy challenges through the remarkable efforts of a large, diverse,  
542 collaborative, and growing scientific community around the world.

## 543 **Data Availability Statement**

544 No new data was produced as part of this manuscript. All data analysed is publicly available and  
545 cited in the manuscript.

546 **Acknowledgments**

547 We acknowledge all participants of the September 2022 and June 2023 FishMIP workshops, as  
548 well as the August 2023 NOAA-FishMIP joint workshop. The workshops gave the authors of  
549 this manuscript access to a broad range of information that were developed into the main themes  
550 and key challenges presented in this manuscript. We also thank past and present FishMIP  
551 coordinators for their impact on the success of FishMIP. JLB and CN were supported by  
552 Australian Research Council FT210100798. ABB acknowledges funding from Ocean Frontier  
553 Institute Module H funded by the Canada First Research Excellence Fund. HKL acknowledges  
554 funding from the Natural Sciences and Engineering Research Council (NSERC) of Canada. DPT  
555 acknowledges funding from the Jarislowsky Foundation and NSERC. TDE acknowledges  
556 funding from Fisheries and Oceans Canada (DFO) Atlantic Fisheries Fund and NSERC. KOC  
557 acknowledges support from the National Research Foundation of South Africa (grant 136481).  
558 CSH and KR were supported by the US National Science Foundation (NSF) and NOAA climate  
559 program office. OM acknowledges support from the European Union's Horizon 2020 research  
560 and innovation program under grant agreement N° 817806.

561 **References**

- 562 Barange, M., Merino, G., Blanchard, J.L., Scholtens, J., Harle, J., Allison, E.H., Allen, J.I., Holt,  
563 J. and Jennings, S., 2014. Impacts of climate change on marine ecosystem production in  
564 societies dependent on fisheries. *Nature Climate Change*, 4(3), 211-216.  
565 <https://doi.org/10.1038/NCLIMATE2119>
- 566 Barrier, N., Lengaigne, M., Rault, J., Person, R., Ethé, C., Aumont, O. and Maury, O.,  
567 2023. Mechanisms underlying the epipelagic ecosystem response to ENSO in the  
568 equatorial Pacific ocean. *Progress in Oceanography*, 213, 103002.  
569 <https://doi.org/10.1016/j.pocean.2023.103002>
- 570 Blanchard, J.L., Jennings, S., Holmes, R., Harle, J., Merino, G., Allen, J.I., Holt, J., Dulvy, N.K.  
571 and Barange, M., 2012. Potential consequences of climate change for primary production  
572 and fish production in large marine ecosystems. *Philosophical Transactions of the Royal  
573 Society B: Biological Sciences*, 367(1605), 2979-2989.  
574 <https://doi.org/10.1098/rstb.2012.0231>

575 Blanchard, J.L., Watson, R.A., Fulton, E.A., Cottrell, R.S., Nash, K.L., Bryndum-  
576 Buchholz, A., Büchner, M., Carozza, D.A., Cheung, W.W., Elliott, J. and Davidson,  
577 L.N., 2017. Linked sustainability challenges and trade-offs among fisheries, aquaculture  
578 and agriculture. *Nature ecology & evolution*, 1(9), 1240-1249.  
579 <https://doi.org/10.1038/s41559-017-0258-8>

580 Blanchard, J.L., Novaglio, C., Maury, O., Harrison, C., Petrik, C., Ortega Cisneros, L., Eddy,  
581 T.D., Heneghan, R., Liu, X., Stock, C., Rousseau, Y., Buechner, M., Roberts, K.,  
582 Bryndum-Buchholz, A., Adekoya, E., Andersen, K., Bianchi, D., Bulman, C., Chueng,  
583 W., Christensen, V., Coll, M., Capitani, L., Datta, S., Fierro Arcos, D., Fulton, B.,  
584 Garza, V., Guiet, J., Murphy, K., Oliveros-Ramos, R., Palacios-Abrantes, J., Reum, J.,  
585 Schewe, J., Scherrer, K., Shin, Y., Steenbeck, J, van Denderen, D., Woodworth-Jefcoats,  
586 P., Tittensor, D.P. (This issue). Detecting, attributing, and projecting global marine  
587 ecosystem and fisheries change: FishMIP 2.0.

588 Bianchi, D., Carozza, D.A., Galbraith, E.D., Guiet, J. and DeVries, T., 2021. Estimating  
589 global biomass and biogeochemical cycling of marine fish with and without fishing.  
590 *Science advances*, 7(41), eabd7554.  
591 <https://www.science.org/doi/full/10.1126/sciadv.abd7554>

592 Bopp, L., Resplandy, L., Orr, J. C., Doney, S. C., Dunne, J. P., Gehlen, M., Halloran, P.,  
593 Heinze, C., Ilyina, T., Séférian, R., Tjiputra, J., & Vichi, M. (2013). Multiple stressors of  
594 ocean ecosystems in the 21st century: Projections with CMIP5 models. *Biogeosciences*,  
595 10(10), 6225–6245. <https://doi.org/10.5194/bg-10-6225-2013>

596 Bopp, L., Aumont, O., Kwiatkowski, L., Le Mezo, P., Maury, O., Séférian, R., and  
597 Tagliabue, A.: Projecting net primary production in a sea of uncertainty: next steps and  
598 why should we care?, 2022. EGU General Assembly 2022, Vienna, Austria, 23–27 May  
599 2022, EGU22-8687, <https://doi.org/10.5194/egusphere-egu22-8687>

600 Boyce, D.G., Lotze, H.K., Tittensor, D.P., Carozza, D.A. and Worm, B., 2020. Future  
601 ocean biomass losses may widen socioeconomic equity gaps. *Nature Communications*,  
602 11(1), 2235. <https://doi.org/10.1038/s41467-020-15708-9>

603 Boyce, D.G., Tittensor, D.P., Garilao, C., Henson, S., Kaschner, K., Kesner-Reyes, K.,  
604 Pigot, A., Reyes Jr, R.B., Reygondeau, G., Schleit, K.E. and Shackell, N.L., 2022. A  
605 climate risk index for marine life. *Nature Climate Change*, 12(9), 854-862.  
606 <https://doi.org/10.1038/s41558-022-01437-y>

607 Bryndum-Buchholz, A., Tittensor, D.P., Blanchard, J.L., Cheung, W.W., Coll, M.,  
608 Galbraith, E.D., Jennings, S., Maury, O. and Lotze, H.K., 2019. Twenty-first-century  
609 climate change impacts on marine animal biomass and ecosystem structure across ocean  
610 basins. *Global change biology*, 25(2), 459-472. <https://doi.org/10.1111/gcb.14512>

611 Bryndum-Buchholz, A., Boyce, D.G., Tittensor, D.P., Christensen, V., Bianchi, D. and  
612 Lotze, H.K., 2020a. Climate-change impacts and fisheries management challenges in the  
613 North Atlantic Ocean. *Marine Ecology Progress Series*, 648, 1-17.  
614 <https://doi.org/10.3354/meps13438>

615 Bryndum-Buchholz, A., Prentice, F., Tittensor, D.P., Blanchard, J.L., Cheung, W.W.L.,  
616 Christensen, V., Galbraith, E.D., Maury, O. and Lotze, H.K., 2020b. Differing marine  
617 animal biomass shifts under 21st century climate change between Canada’s three oceans.  
618 *Facets* 5, 105–122. <https://doi.org/10.1139/facets-2019-0035>

619 Bryndum-Buchholz, A., Tittensor, D.P. and Lotze, H.K., 2021. The status of climate change  
620 adaptation in fisheries management: Policy, legislation and implementation. *Fish and*  
621 *Fisheries*, 22(6), 1248-1273. <https://doi.org/10.1111/faf.12586>

622 Bryndum-Buchholz, A., 2022. Keeping up hope as an early career climate-impact  
623 scientist. *ICES Journal of Marine Science*, 79(9), 2345-2350.  
624 <https://doi.org/10.1093/icesjms/fsac180>

625 Bryndum-Buchholz, A., Blanchard, J.L., Coll, M., Pontavice, H.D., Everett, J.D., Guiet, J.,  
626 Heneghan, R.F., Maury, O., Novaglio, C., Palacios-Abrantes, J. and Petrik, C.M., 2023a.  
627 Applying ensemble ecosystem model projections to future-proof marine conservation  
628 planning in the Northwest Atlantic Ocean. *Facets*, 8, 1-16. [https://doi.org/10.1139/facets-](https://doi.org/10.1139/facets-2023-0024)  
629 [2023-0024](https://doi.org/10.1139/facets-2023-0024)

630 Bryndum-Buchholz, A., Lotze, H.K., Novaglio, C. and Eddy, T.D. 2023b. Ocean  
631 Biomass and Climate Change. Elsevir. [https://doi.org/10.1016/B978-0-323-90798-](https://doi.org/10.1016/B978-0-323-90798-9.00010-X)  
632 [9.00010-X](https://doi.org/10.1016/B978-0-323-90798-9.00010-X)

633 Cheung, W.W., Dunne, J., Sarmiento, J.L. and Pauly, D., 2011. Integrating ecophysiology and  
634 plankton dynamics into projected maximum fisheries catch potential under climate  
635 change in the Northeast Atlantic. *ICES Journal of Marine Science*, 68(6), 1008-1018.  
636 <https://doi.org/10.1093/icesjms/fsr012>

637 Cinner, J.E., Caldwell, I.R., Thiault, L., Ben, J., Blanchard, J.L., Coll, M., Diedrich, A.,  
638 Eddy, T.D., Everett, J.D., Folberth, C. and Gascuel, D., 2022. Potential impacts of  
639 climate change on agriculture and fisheries production in 72 tropical coastal  
640 communities. *Nature communications*, 13(1), 3530. [https://doi.org/10.1038/s41467-022-](https://doi.org/10.1038/s41467-022-30991-4)  
641 [30991-4](https://doi.org/10.1038/s41467-022-30991-4)

642 Coll, M., Steenbeek, J., Pennino, M.G., Buszowski, J., Kaschner, K., Lotze, H.K.,  
643 Rousseau, Y., Tittensor, D.P., Walters, C., Watson, R.A. and Christensen, V., 2020.  
644 Advancing global ecological modeling capabilities to simulate future trajectories of  
645 change in marine ecosystems. *Frontiers in Marine Science*, 7, 567877.  
646 <https://doi.org/10.3389/fmars.2020.567877>

647 Costello, C., Cao, L., Gelcich, S., Cisneros-Mata, M.Á., Free, C.M., Froehlich, H.E.,  
648 Golden, C.D., Ishimura, G., Maier, J., Macadam-Somer, I. and Mangin, T., 2020. The  
649 future of food from the sea. *Nature*, 588(7836), 95-100. [https://doi.org/10.1038/s41586-](https://doi.org/10.1038/s41586-020-2616-y)  
650 [020-2616-y](https://doi.org/10.1038/s41586-020-2616-y)

651 Cottrell, R.S., Nash, K.L., Halpern, B.S., Remenyi, T.A., Corney, S.P., Fleming, A., Fulton,  
652 E.A., Hornborg, S., Johne, A., Watson, R.A. and Blanchard, J.L., 2019. Food production  
653 shocks across land and sea. *Nature Sustainability*, 2(2), 130-137.  
654 <https://doi.org/10.1038/s41893-018-0210-1>

655 Drenkard, E. J., Stock, C., Ross, A. C., Dixon, K. W., Adcroft, A., Alexander, M., ... & Wang,  
656 M. 2021. Next-generation regional ocean projections for living marine resource  
657 management in a changing climate. *ICES Journal of Marine Science*, 78(6), 1969-1987.  
658 <https://doi.org/10.1093/icesjms/fsab100>

659 Dupont, L., Le Mézo, P., Aumont, O., Bopp, L., Clerc, C., Ethé, C. and Maury, O., 2023.  
660 High trophic level feedbacks on global ocean carbon uptake and marine ecosystem  
661 dynamics under climate change. *Global Change Biology*, 29(6), 1545-1556.  
662 <https://doi.org/10.1111/gcb.16558>

663 du Pontavice, H., 2019. Changing biomass flows in marine ecosystems: from the past to  
664 the future. In *Predicting Future Oceans* (pp. 121-128). Elsevier.  
665 <https://doi.org/10.1016/B978-0-12-817945-1.00012-5>

666 du Pontavice, H., Gascuel, D., Reygondeau, G., Stock, C. and Cheung, W.W., 2021.  
667 Climate-induced decrease in biomass flow in marine food webs may severely affect  
668 predators and ecosystem production. *Global Change Biology*, 27(11), pp.2608-2622.  
669 <https://doi.org/10.1111/gcb.15576>

670 Eddy, T.D., 2019. Building confidence in projections of future ocean capacity. In  
671 *Predicting Future Oceans* (pp. 69-76). Elsevier. [https://doi.org/10.1016/B978-0-12-](https://doi.org/10.1016/B978-0-12-817945-1.00007-1)  
672 [817945-1.00007-1](https://doi.org/10.1016/B978-0-12-817945-1.00007-1)

673 Erauskin-Extramiana, M., Chust, G., Arrizabalaga, H., Cheung, W.W., Santiago, J.,  
674 Merino, G. and Fernandes-Salvador, J.A., 2023. Implications for the global tuna fishing  
675 industry of climate change-driven alterations in productivity and body sizes. *Global and*  
676 *Planetary Change*, 222, 104055. <https://doi.org/10.1016/j.gloplacha.2023.104055>

677 FAO. 2020. *The State of World Fisheries and Aquaculture 2020. Sustainability in action*. Rome.  
678 <https://doi.org/10.4060/ca9229en>

679 Fu, W., Moore, J. K., Primeau, F., Collier, N., Ogunro, O. O., Hoffman, F. M., & Randerson, J.  
680 T. 2022. Evaluation of Ocean Biogeochemistry and Carbon Cycling in CMIP Earth  
681 System Models With the International Ocean Model Benchmarking (IOMB) Software  
682 System. *Journal of Geophysical Research: Oceans*, 127(10), e2022JC018965.  
683 <https://doi.org/10.1029/2022JC018965>

684 Frieler, K., Lange, S., Piontek, F., Reyer, C.P., Schewe, J., Warszawski, L., Zhao, F., Chini, L.,  
685 Denvil, S., Emanuel, K. and Geiger, T., 2017. Assessing the impacts of 1.5 C global  
686 warming—simulation protocol of the Inter-Sectoral Impact Model Intercomparison Project  
687 (ISIMIP2b). *Geoscientific Model Development*, 10(12), 4321-4345.  
688 <https://doi.org/10.5194/gmd-10-4321-2017>

689 Frieler, K., Volkholz, J., Lange, S., Schewe, J., Mengel, M., Rivas López, M.D.R., Otto,  
690 C., Reyer, C.P., Karger, D.N., Malle, J.T. and Treu, S., 2023. Scenario set-up and forcing  
691 data for impact model evaluation and impact attribution within the third round of the  
692 Inter-Sectoral Model Intercomparison Project (ISIMIP3a). *EGUsphere*, 1-83.  
693 <https://doi.org/10.5194/egusphere-2023-281>

694 Gómara, I., Rodríguez-Fonseca, B., Mohino, E., Losada, T., Polo, I. and Coll, M., 2021.  
695 Skillful prediction of tropical Pacific fisheries provided by Atlantic Niños. *Environmental*  
696 *Research Letters*, 16(5), 054066. <https://doi.org/10.1088/1748-9326/abfa4d>

697 Guibourd de Luziniais, V., Du Pontavice, H., Reygondeau, G., Barrier, N., Blanchard,  
698 J.L., Bornarel, V., Büchner, M., Cheung, W.W., Eddy, T.D., Everett, J.D. and Guiet, J.,  
699 2023. Trophic amplification: A model intercomparison of climate driven changes in  
700 marine food webs. *PloS one*, 18(8), e0287570.  
701 <https://doi.org/10.1371/journal.pone.0287570>

702 Heneghan, R.F., Hatton, I.A. and Galbraith, E.D., 2019. Climate change impacts on  
703 marine ecosystems through the lens of the size spectrum. *Emerging Topics in Life*  
704 *Sciences*, 3(2), 233-243. <https://doi.org/10.1042/ETLS20190042>

705 Heneghan, R.F., Galbraith, E., Blanchard, J.L., Harrison, C., Barrier, N., Bulman, C., Cheung,  
706 W., Coll, M., Eddy, T.D., Erauskin-Extramiana, M. and Everett, J.D., 2021.  
707 Disentangling diverse responses to climate change among global marine ecosystem

708 models. *Progress in Oceanography*, 198, 102659.  
709 <https://doi.org/10.1016/j.pocean.2021.102659>

710 Heneghan, R.F., Everett, J.D., Blanchard, J.L., Sykes, P. and Richardson, A.J., 2023.  
711 Climate-driven zooplankton shifts cause large-scale declines in food quality for fish.  
712 *Nature Climate Change*, 13(5), 470-477. <https://doi.org/10.1038/s41558-023-01630-7>

713 Holt, J., Hyder, P., Ashworth, M., Harle, J., Hewitt, H.T., Liu, H., New, A.L., Pickles, S.,  
714 Porter, A., Popova, E. and Allen, J., 2017. Prospects for improving the representation of  
715 coastal and shelf seas in global ocean models. *Geoscientific Model Development*, 10(1),  
716 499-523. <https://doi.org/10.5194/gmd-10-499-2017>

717 IPBES (2019) Global assessment report on biodiversity and ecosystem services of the  
718 Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services.  
719 Díaz, S., Settele, J., Brondizio E.S., Ngo, H. T. , Guèze, M., Agard, J., Arneth, A.,  
720 Balvanera, P., Brauman, K. A., Butchart, S. H. M., Chan, K. M. A., Garibaldi, L. A.,  
721 Ichii, K., Liu, J., Subramanian, S. M., Midgley, G. F., Miloslavich, P., Molnár, Z., Obura,  
722 D., Pfaff, A., Polasky, S., Purvis, A., Razzaque, J., Reyers, B., Chowdhury, R. R., Shin,  
723 Y.-J., Visseren-Hamakers, I. J., Willis, K. J., Zayas, C. N. (eds.). IPBES secretariat,  
724 Bonn, Germany.

725 IPCC, (2019a): Climate Change and Land: an IPCC special report on climate change,  
726 desertification, land degradation, sustainable land management, food security, and  
727 greenhouse gas fluxes in terrestrial ecosystems[P.R. Shukla, J. Skea, E. Calvo Buendia,  
728 V. Masson-Delmotte, H.-O. Pörtner, D. C. Roberts, P. Zhai, R. Slade, S. Connors, R. van  
729 Diemen, M. Ferrat, E. Haughey, S. Luz, S. Neogi, M. Pathak, J. Petzold, J. Portugal  
730 Pereira, P. Vyas, E. Huntley, K. Kissick, M. Belkacemi, J. Malley, (eds.)]. IPCC, Geneva,  
731 Switzerland.

732 IPCC 2019b. IPCC Special Report on the Ocean and Cryosphere in a Changing Climate.  
733 Pörtner, H. O., Roberts, D. C., Masson-Delmotte, V., Zhai, P., Tignor, M., Poloczanska,  
734 E., Mintenbeck, K., Nicolai, M., Okem, A., Petzold, J., Rama, B., Weyer, N. (eds.).  
735 IPCC, Geneva, Switzerland.

736 IPCC, 2022. Climate Change 2022: Impacts, Adaptation, and Vulnerability. Contribution of  
737 Working Group II to the Sixth Assessment Report of the Intergovernmental Panel on  
738 Climate Change [H.-O. Pörtner, D.C. Roberts, M. Tignor, E.S. Poloczanska, K.  
739 Mintenbeck, A. Alegría, M. Craig, S. Langsdorf, S. Lösschke, V. Möller, A. Okem, B.  
740 Rama (eds.)]. Cambridge University Press. Cambridge University Press, Cambridge, UK  
741 and New York, NY, USA, 3056 pp.,

742 ISIMIP, (n.d.). The Inter-sectoral Impacts Model Intercomparison Project (ISIMIP). Mission &  
743 Implementation.  
744 [https://www.isimip.org/documents/646/MissionAndImplementation\\_12Sep2018\\_5Hlvj2](https://www.isimip.org/documents/646/MissionAndImplementation_12Sep2018_5Hlvj2N.pdf)  
745 [N.pdf](https://www.isimip.org/documents/646/MissionAndImplementation_12Sep2018_5Hlvj2N.pdf)

746 Jacox, M. G., Alexander, M. A., Siedlecki, S., Chen, K., Kwon, Y.-O., Brodie, S., Ortiz, I.,  
747 Tommasi, D., Widlansky, M. J., Barrie, D., Capotondi, A., Cheng, W., Di Lorenzo, E.,  
748 Edwards, C., Fiechter, J., Fratantoni, P., Hazen, E. L., Hermann, A. J., Kumar, A., ...  
749 Rykaczewski, R. (2020). Seasonal-to-interannual prediction of North American coastal  
750 marine ecosystems: Forecast methods, mechanisms of predictability, and priority  
751 developments. *Progress in Oceanography*, 183, 102307.  
752 <https://doi.org/10.1016/j.pocean.2020.102307>

753 Lefort, S., Aumont, O., Bopp, L., Arsouze, T., Gehlen, M. and Maury, O., 2015. Spatial  
754 and body-size dependent response of marine pelagic communities to projected global  
755 climate change. *Global change biology*, 21(1), 154-164.  
756 <https://doi.org/10.1111/gcb.12679>

757 Liu, X., Stock, C. A., Dunne, J. P., Lee, M., Shevliakova, E., Malyshev, S., & Milly, P. C. D.  
758 2021. Simulated Global Coastal Ecosystem Responses to a Half-Century Increase in  
759 River Nitrogen Loads. *Geophysical Research Letters*, 48(17), e2021GL094367.  
760 <https://doi.org/10.1029/2021GL094367>

761 Lotze, H.K., Tittensor, D.P., Bryndum-Buchholz, A., Eddy, T.D., Cheung, W.W., Galbraith,  
762 E.D., Barange, M., Barrier, N., Bianchi, D., Blanchard, J.L. and Bopp, L., 2019. Global  
763 ensemble projections reveal trophic amplification of ocean biomass declines with climate  
764 change. *Proceedings of the National Academy of Sciences*, 116(26), 12907-12912.  
765 <https://doi.org/10.1073/pnas.1900194116>

766 Lotze, H.K., Bryndum-Buchholz, A. and Boyce, D.G., 2021. Effects of climate change  
767 on food production (fishing). In *The Impacts of Climate Change*, pp. 205-231. Elsevier.  
768 <https://doi.org/10.1016/B978-0-12-822373-4.00017-3>

769 Lotze, H.K., Mellon, S., Coyne, J., Betts, M., Burchell, M., Fennel, K., Dusseault, M.A.,  
770 Fuller, S.D., Galbraith, E., Suarez, L.G. and de Gelleke, L., 2022. Long-term ocean and  
771 resource dynamics in a hotspot of climate change. *Facets*. 7, 1142-1184.  
772 <https://doi.org/10.1139/facets-2021-0197>

773 Maury, O., Campling, L., Arrizabalaga, H., Aumont, O., Bopp, L., Merino, G., Squires, D.,  
774 Cheung, W., Goujon, M., Guivarch, C. and Lefort, S., 2017. From shared socio-economic  
775 pathways (SSPs) to oceanic system pathways (OSPs): Building policy-relevant scenarios  
776 for global oceanic ecosystems and fisheries. *Global Environmental Change*, 45, 203-216.  
777 <https://doi.org/10.1016/j.gloenvcha.2017.06.007>

778 Maury, O., Tittensor, D.P., Eddy, T.D., Allison, E.H., Fulton, E.A., Heneghan, R.F.,...Cheung,  
779 W.W.L. The Ocean System Pathways (OSPs): a new FishMIP scenario framework to  
780 investigate the future of oceans. Pre-print in this issue.  
781 National Academies of Sciences, Engineering, and Medicine. 2021. Reflecting Sunlight:  
782 Recommendations for Solar Geoengineering Research and Research Governance.  
783 Washington, DC: The National Academies Press. <https://doi.org/10.17226/25762>.

784 National Academies of Sciences, Engineering, and Medicine. 2022. A Research Strategy for  
785 Ocean-based Carbon Dioxide Removal and Sequestration. Washington, DC: The  
786 National Academies Press.

787 Novaglio C., Tittensor D.P., Blanchard J.L.. Using ecological theory to evaluate and  
788 reduce uncertainties of a marine ecosystem model ensemble. pre-print in this issue.

789 Payne, M.R., Barange, M., Cheung, W.W., MacKenzie, B.R., Batchelder, H.P., Cormon,  
790 X., Eddy, T.D., Fernandes, J.A., Hollowed, A.B., Jones, M.C. and Link, J.S., 2016.  
791 Uncertainties in projecting climate-change impacts in marine ecosystems. *ICES Journal*  
792 *of Marine Science*, 73(5), 1272-1282. <https://doi.org/10.1093/icesjms/fsv231>

793 Pereira, L.M., Davies, K.K., den Belder, E., Ferrier, S., Karlsson-Vinkhuyzen, S., Kim,  
794 H., Kuiper, J.J., Okayasu, S., Palomo, M.G., Pereira, H.M. and Peterson, G., 2020.  
795 Developing multiscale and integrative nature–people scenarios using the Nature Futures  
796 Framework. *People and Nature*, 2(4), 1172-1195. <https://doi.org/10.1002/pan3.10146>

797 Perryman, H.A., Kaplan, I.C., Blanchard, J.L., Fay, G., Gaichas, S.K., McGregor, V.L.,  
798 Morzaria-Luna, H.N., Porobic, J., Townsend, H. and Fulton, E.A., 2023. Atlantis

799 Ecosystem Model Summit 2022: Report from a workshop. *Ecological Modelling*, 483,  
800 110442. <https://doi.org/10.1016/j.ecolmodel.2023.110442>

801 Pethybridge, H.R., Fulton, E.A., Hobday, A.J., Blanchard, J., Bulman, C.M., Butler, I.R.,  
802 Cheung, W.W., Dutra, L.X., Gorton, R., Hutton, T. and Matear, R., 2020. Contrasting  
803 futures for Australia’s fisheries stocks under IPCC RCP8.5 emissions—a multi-ecosystem  
804 model approach. *Frontiers in Marine Science*, 7, 577964.  
805 <https://doi.org/10.3389/fmars.2020.577964>

806 Petrik, C.M., Luo, J.Y., Heneghan, R.F., Everett, J.D., Harrison, C.S. and Richardson,  
807 A.J., 2022. Assessment and constraint of mesozooplankton in CMIP6 earth system  
808 models. *Global Biogeochemical Cycles*, 36(11), e2022GB007367.  
809 <https://doi.org/10.1029/2022GB007367>

810 Rosenzweig, C., Arnell, N.W., Ebi, K.L., Lotze-Campen, H., Raes, F., Rapley, C., Smith,  
811 M.S., Cramer, W., Frieler, K., Reyer, C.P. and Schewe, J., 2017. Assessing inter-sectoral  
812 climate change risks: the role of ISIMIP. *Environmental Research Letters*, 12(1), 010301.  
813 <https://doi.org/10.1088/1748-9326/12/1/010301>

814 Rousseau Y., Blanchard, J.L., Novaglio, C., Pinnell, K.A., Tittensor, D.T., Watson, R.A., Ye,  
815 Y. (in press). A database of mapped global fishing activity, 1950- 2017’ Scientific Data.  
816 Ruane, A.C., Teichmann, C., Arnell, N.W., Carter, T.R., Ebi, K.L., Frieler, K., Goodess,  
817 C.M., Hewitson, B., Horton, R., Kovats, R.S. and Lotze, H.K., 2016. The vulnerability,  
818 impacts, adaptation and climate services advisory board (VIACS AB v1.0) contribution  
819 to CMIP6. *Geoscientific Model Development*, 9(9), pp.3493-3515.  
820 <https://doi.org/10.5194/gmd-9-3493-2016>

821 Rynne N., Novaglio C., Heneghan R., Bianchi D., Christensen V., Coll M., Guet J.,  
822 Steenbeek J., Bryndum-Buchholza A., Eddy T.D., Maury O., Petrik C.M., Ortega-  
823 Cisneros K., Tittensor D.P., Blanchard J. A skill assessment framework for the Fisheries  
824 and Marine Ecosystem Model Intercomparison Project. pre-print in this issue.

825 Schewe, J., Gosling, S.N., Reyer, C., Zhao, F., Ciais, P., Elliott, J., Francois, L., Huber,  
826 V., Lotze, H.K., Seneviratne, S.I. and Van Vliet, M.T., 2019. State-of-the-art global  
827 models underestimate impacts from climate extremes. *Nature communications*, 10(1),  
828 1005. <https://doi.org/10.1038/s41467-019-08745-6>

829 Steenbeek, J., Buszowski, J., Chagaris, D., Christensen, V., Coll, M., Fulton, E.A., Katsanevakis,  
830 S., Lewis, K.A., Mazaris, A.D., Macias, D. and De Mutsert, K., 2021. Making spatial-  
831 temporal marine ecosystem modelling better—a perspective. *Environmental Modelling &*  
832 *Software*, 145, 105209. <https://doi.org/10.1016/j.envsoft.2021.105209>

833 Stock, C. A., Alexander, M. A., Bond, N. A., Brander, K. M., Cheung, W. W. L., Curchitser, E.  
834 N., Delworth, T. L., Dunne, J. P., Griffies, S. M., Haltuch, M. A., Hare, J. A., Hollowed,  
835 A. B., Lehodey, P., Levin, S. A., Link, J. S., Rose, K. A., Rykaczewski, R. R., Sarmiento,  
836 J. L., Stouffer, R. J., ... Werner, F. E. (2011). On the use of IPCC-class models to assess  
837 the impact of climate on Living Marine Resources. *Progress in Oceanography*, 88(1), 1–  
838 27. <https://doi.org/10.1016/j.pocean.2010.09.001>

839 Thiault, L., Mora, C., Cinner, J.E., Cheung, W.W., Graham, N.A., Januchowski-Hartley, F.A.,  
840 Mouillot, D., Sumaila, U.R. and Claudet, J., 2019. Escaping the perfect storm of  
841 simultaneous climate change impacts on agriculture and marine fisheries. *Science*  
842 *Advances*, 5(11), eaaw9976. [10.1126/sciadv.aaw9976](https://doi.org/10.1126/sciadv.aaw9976)

843 Tiller, R. and Mendenhall, E., 2023. And so it begins–The adoption of the ‘Biodiversity  
844 Beyond National Jurisdiction’ treaty. *Marine Policy*, 157, 105836.  
845 <https://doi.org/10.1016/j.marpol.2023.105836>

846 Tittensor, D.P., Eddy, T.D., Lotze, H.K., Galbraith, E.D., Cheung, W., Barange, M., Blanchard,  
847 J.L., Bopp, L., Bryndum-Buchholz, A., Büchner, M. and Bulman, C., 2018. A protocol  
848 for the intercomparison of marine fishery and ecosystem models: Fish-MIP v1. 0.  
849 *Geoscientific Model Development*, 11(4), 1421-1442. [https://doi.org/10.5194/gmd-11-](https://doi.org/10.5194/gmd-11-1421-2018)  
850 [1421-2018](https://doi.org/10.5194/gmd-11-1421-2018)

851 Tittensor, D.P., Novaglio, C., Harrison, C.S., Heneghan, R.F., Barrier, N., Bianchi, D., Bopp, L.,  
852 Bryndum-Buchholz, A., Britten, G.L., Büchner, M. and Cheung, W.W., 2021. Next-  
853 generation ensemble projections reveal higher climate risks for marine ecosystems.  
854 *Nature Climate Change*, 11(11), 973-981. <https://doi.org/10.1038/s41558-021-01173-9>

855 Van Dijk, M., Morley, T., Rau, M.L. and Saghai, Y., 2021. A meta-analysis of projected global  
856 food demand and population at risk of hunger for the period 2010–2050. *Nature Food*,  
857 2(7), 494-501. <https://doi.org/10.1038/s43016-021-00322-9>