INVESTIGATING THE RESILIENCE OF SALT MARSHES TO EXTERNAL DISTURBANCE

Natascia Pannozzo¹, Rachel K Smedley¹, Iacopo Carnacina², and Nicoletta Leonardi¹

¹Department of Geography and Planning, University of Liverpool ²Department of Civil Engineering -Liverpool, John Mores University

March 11, 2024

Abstract

Salt marshes are valuable ecosystems that provide numerous services and act as natural coastal defences by buffering storm waves and stabilising sediments. However, it is not clear whether they will be able to retain their resilience with accelerating rate in sea-level rise, possible increases in storm intensity and increasing land reclamation. The current paradigm is that a positive sediment budget supports the survival and accretion of salt marshes while a negative sediment budget causes marsh degradation. Here we present the results of a series of studies that used an integration of modelling and paleoenvironmental analysis and a sediment budget approach to investigate the resilience of estuaries and salt marshes to projected rise in sea-level, possible increases in storm activity and existing anthropogenic disturbance. The studies were conducted using the Ribble Estuary-NorthWest England-as a test case, the hydrodynamic model Delft3D to simulate the estuary morpho-dynamics under selected scenarios, and optically stimulated luminescence (OSL), geochemistry and particle size distribution analysis to reconstruct the past evolution and adaptation of the estuary morphology. Results showed that sea-level rise threatens estuary and marsh stability by promoting ebb dominance and triggering a net export of sediment. Conversely, storm surges aid the resilience of the system by promoting flood dominance and triggering a net import of sediment and have the potential to counteract the negative impact of sea-level rise by masking its effects on the sediment budget. The addition of embankments, on the other hand, can further promote ebb dominance in the system and intensify sediment export, further threatening marsh stability.

References

Leonardi, N., Ganju, N.K. and Fagherazzi, S. (2016). A linear relationship between wave power and erosion determines saltmarsh resilience to violent storms and hurricanes. Proceedings of the National Academy of Sciences 113(1), 64-68.

Ganju, N.K., Kirwan, M.L., Dickhudt, P.J., Guntenspergen, G.R., Cahoon, D.R. and Kroeger, K.D. (2015). Sediment transportbased metrics of wetland stability. Geophysical Research Letters, 42(19), 7992-8000.





INVESTIGATING THE RESILIENCE OF SALT MARSHES TO EXTERNAL DISTURBANCE

N. Pannozzo^{1*}, R.K. Smedley¹, I. Carnacina², N. Leonardi¹

¹Department of Geography and Planning – University of Liverpool, Liverpool, Merseyside, United Kingdom

²Department of Civil Engineering – Liverpool John Mores University, Liverpool, Merseyside, United Kingdom

*sgnpanno@liverpool.ac.uk



Paper 1

'Saltmarsh resilience to sea-level rise and increased storm intensity' + follow-up work

Paper 2

'Influence of sediment availability and embankment construction on salt marsh resilience to sea-level rise'



- Salt marshes act as natural, long-term, low-cost coastal defences by buffering storm waves and stabilising sediments.
- Marshes are found in intertidal areas and their growth happens by vertical and lateral accretion, which occurs by delivery of fine sediments from rivers and the sea to accommodation spaces.
- Marsh resilience can be monitored analysing the sediment budget = balance between sediment added to and removed from the marsh, when more material is added than it is removed, there is a surplus of sediment and the shore builds seaward, and vice versa.

Paper 1 - 'Saltmarsh resilience to sea-level rise and increased storm intensity'

Aim and objectives


To investigate marsh and estuary resilience under various storm surge and sea-level scenarios by using a sediment budget approach and the hydrodynamic model Delft3D:

- Change in sea-level only
- Change in surge intensity only
 - timing of the storm surge relative to high or low tide
 - duration of the surge
 - change in tidal range
- Combination of storm surge and sea-level scenarios

One month of simulation.


Case study - Ribble Estuary (North West England):

- Funnel shaped, hypertidal.
- One of the largest tidal flat salt marsh complexes in Europe, part of which has been recently restored through managed realignment.





Results: sediment budget

- > Deposit increases when the surge height increases.
- > Timing of the surge with respect to HT and LT and presence of vegetation can affect the trends.
- > With longer storm duration or lower tidal range these effects decrease and become negligible.



Results: sediment budget





Sea-level

> Deposition decreases with an increase in sea-level.

Sea-level + storm surges (48h spring)

- The trends in deposition caused by the increase in surge height remain similar in all sea-level scenarios.
- There is an overall decrease in the magnitude of deposition with the increase in sea-level.
- The effects of storm surges mask the effects of sealevel rise on the sediment budget; this is especially true for surges with the highest intensities (>3 m).



Storm surges could contribute to salt marsh and estuarine resilience

Key points

- Sea-level rise will threaten the stability of the marsh
- Storm surges might contribute to increase salt marsh resilience to sea-level rise

References

Pannozzo N., Leonardi N., Carnacina I., Smedley R. (2021). Salt marsh resilience to sea-level rise and increased storm intensity. *Geomorphology*, 389 (4): 107825. Pannozzo N., Leonardi N., Carnacina I., Smedley R. (2021). Dataset of results from numerical simulations of increased storm intensity in an estuarine salt marsh system. *Data in Brief*, 38 (6): 107336.

Follow-up work

'Investigating spatial distribution of storm deposit on the marsh platform and its transport pathway'

- Investigate parameters that affect spatial variability of storm sedimentation on the marsh platform
- Investigate provenance of sediment that are transported on the marsh platform by storms
- 1. Analysis performed on simulated surge scenarios (the ones just discussed)
 - + machine learning tools used to
 - classify significance of the various hydrodynamic and morphological parameters impact on the different tidal flat/salt marsh areas + generate predictor for calculating sediment budget for the different areas given certain hydrodynamic conditions
 - investigate sediment connectivity to understand the provenance and pathways of storm sediment from ocean to marsh
- 2. Analysis performed on sediment collected in the field using sediment traps
 - sediments collected from the marsh platform on a monthly basis as well as sediments collected from possible sources

Paper 2 - 'Influence of sediment availability and embankment construction on salt marsh resilience to sea-level rise'


Case study - Ribble Estuary (North West England):

- Widespread anthropogenic interventions including embankment construction since 1810.
- Marsh accreting at a fast rate, previously thought to be linked to embankment presence.

Aim and objectives


Investigate salt marsh resilience to sea-level rise in relation to sediment supply and embankment construction using a combination of paleoenvironmental analysis and Delft 3D:


- Investigate provenance of sediments performing geochemical analysis on sediments from the marsh and possible sources (i.e. river, ocean);
- Detect changes in sediment accretion rate using OSL alongside down-core changes in particle size distribution, geochemistry and organic content to detect changes or shifts in sedimentation processes and depositional environment;
- Compare changes to a record of estuary management to investigate any correlation with embankment construction;
- Use Delft3D to investigate sediment budget calculated from numerical simulations modelling scenarios with and without embankment.



Results: paleoenvironmental reconstruction

- ➤ Rapid accretion followed by a slower deposition.
- The accretion is currently a magnitude higher than sea-level rise (c.a. 2 mm/yr) meaning that it allows the marsh to survive.
- The rapid accretion precedes the beginning of embankment construction (c.a. 1810).
- > Embankment cannot be responsible for a faster accretion.
- > Could it be the natural evolution of the system?







Results: paleoenvironmental reconstruction

- Geochemistry correlates sediment supply to the marsh platform with sediments from the bed of the Irish Sea.
- The Irish Sea is characterised by high quantity of mobile sediment, legacy of the Irish Sea Glacier.
- Could the fast accretion be naturally driven by a high sediment supply to the system and the trend resemble the natural evolution of a marsh system?

Results: paleoenvironmental reconstruction


- Multi-proxy down-core analysis shows gradual evolution of the system from sandflat to mudflat and salt marsh.
- Correlation between evolution of the marsh system and shift in accretion rate.





Results: modelled scenarios

- After one year of simulation, the increase in sediment budget on Hesketh Out Marsh platform is higher for the scenario where no embankments are present.
- When embankments are added, about 10% of the domain is affected by a change in asymmetry. More than 50% of this change involves a shift from flood dominated to ebb dominated cells, while a shift from ebb dominated to flood dominated cells involves less than 50% of this change.
- An overall intensification of ebb dominance and weakening of flood dominance favours sediment export.





High sedimentation rates support mudflat and salt marsh resilience to sea-level rise

Key points

- > The construction of embankments can promote sediment export out of estuarine systems
- > Natural sediment supply from the ocean can be fundamental for the survival of estuarine marshes

References

Pannozzo N., Smedley R., Chiverrell R., Carnacina I., Leonardi N. (2021). Influence of sediment availability and embankment construction on salt marsh resilience to sea-level rise. Journal of Geophysical Research: Earth Surface, In review.

