# Estimation of spectral solar irradiance in the ecliptic plane using synthetic solar surface magnetograms

Isabela de Oliveira<sup>1</sup>, K Sowmya<sup>2</sup>, N.-E Nèmec<sup>3</sup>, and A I Shapiro<sup>2</sup>

<sup>1</sup>Affiliation not available

<sup>2</sup>Max Planck Institute for Solar System Research <sup>3</sup>Institute for Astrophysics and Geophysics, Georg-August-Universität Göttingen

April 26, 2024

#### Abstract

The primary external energy input to planetary atmospheres comes from solar radiation. Accurate estimates of solar irradiance are needed in order to understand atmospheric conditions at other planets. Many studies have interpolated Earth-based irradiance measurements at other planets, disregarding the evolution of solar features, which are responsible for the irradiance variability. We present an approach for assessing solar irradiance at other planets by using synthetic full surface magnetograms of the Sun, generated by the Surface Flux Transport Model, and then converted into sunspots and faculae areas. Following the Spectral And Total Irradiance REconstruction approach, we compute the irradiance as the sum of contributions from sunspots, faculae, and the quiet Sun. We calculate the S-index, which is a proxy for near-ultraviolet irradiance, and the total solar irradiance, which is the wavelength-integrated value. We demonstrate that a simple phase shift of the Earth-observed total solar irradiance does not produce reliable estimates in the ecliptic. By comparing and correlating our results with the interpolated measurements, we find that the two models agree in the S-index variability, because faculae have a long lifetime, so the effect of the solar rotation becomes less important. Conversely, the methods disagree strongly in the total solar irradiance, since sunspots have a short lifetime, and the interpolation does not take into account their emergence and evolution on the far-side of the Sun, as it only uses the information of the disc visible for an Earth-bound observer.

## Estimation of spectral solar irradiance in the ecliptic plane using synthetic solar surface magnetograms

I. de Oliveira<sup>1</sup>, K. Sowmya<sup>1</sup>, N.-E. Nèmec<sup>2</sup>, and A. I. Shapiro<sup>1</sup>

<sup>1</sup>Max Planck Institute for Solar System Research, Justus-von-Liebig-Weg 3, 37077 Göttingen, Germany <sup>2</sup>Institute for Astrophysics and Geophysics, Georg-August-Universität Göttingen, Friedrich-Hund-Platz 1, 37077 Göttingen, Germany

## Key Points:

1

2

3

4 5

6

7

8	• We estimate the solar irradiance at any point in the ecliptic plane, accounting for
9	the evolution of magnetic features on the solar surface.
10	• The method agrees for values in the ultraviolet but disagrees for the TSI when com-
11	pared to the interpolation of Earth-based measurements.
12	• We calculate the correlation between our method and the phase shifting and the
13	interpolation of Earth-based measurements of irradiance.

Corresponding author: I. de Oliveira, oliveira@mps.mpg.de

#### 14 Abstract

The primary external energy input to planetary atmospheres comes from solar ra-15 diation. Accurate estimates of solar irradiance are needed in order to understand atmo-16 spheric conditions at other planets. Many studies have interpolated Earth-based irra-17 diance measurements at other planets, disregarding the evolution of solar features, which 18 are responsible for the irradiance variability. We present an approach for assessing so-19 lar irradiance at other planets by using synthetic full surface magnetograms of the Sun, 20 generated by the Surface Flux Transport Model, and then converted into sunspots and 21 22 faculae areas. Following the Spectral And Total Irradiance REconstruction approach, we compute the irradiance as the sum of contributions from sunspots, faculae, and the 23 quiet Sun. We calculate the S-index, which is a proxy for near-ultraviolet irradiance, and 24 the total solar irradiance, which is the wavelength-integrated value. We demonstrate that 25 a simple phase shift of the Earth-observed total solar irradiance does not produce reli-26 able estimates in the ecliptic. By comparing and correlating our results with the inter-27 polated measurements, we find that the two models agree in the S-index variability, be-28 cause faculae have a long lifetime, so the effect of the solar rotation becomes less impor-29 tant. Conversely, the methods disagree strongly in the total solar irradiance, since sunspots 30 have a short lifetime, and the interpolation does not take into account their emergence 31 and evolution on the far-side of the Sun, as it only uses the information of the disc vis-32 ible for an Earth-bound observer. 33

## 34 1 Introduction

Solar radiation is the main source of external energy input to planetary atmospheres, 35 playing a crucial role in the atmospheric structure and composition. In particular, the 36 ultraviolet (UV) irradiance drives the photochemistry of the Earth's middle atmosphere 37 (Gray et al., 2010), while the extreme ultraviolet (EUV) radiation is responsible for the 38 photochemistry of the upper atmospheres of the planets in the Solar System (Fox et al., 39 2008). A number of general circulation models have been employed in order to under-40 stand the mechanisms that rule the atmospheric behavior. These models need, as an in-41 put, the values of solar irradiance entering the planetary atmosphere. However, the di-42 rect measurements of such irradiance are scarce, so that it must be obtained with mod-43 elling. 44

Several studies have used measurements made at Earth to evaluate the solar irra-45 diance at other planets (e.g. Ramstad et al., 2015; Peter et al., 2014). These studies are 46 based on the assumption that irradiance from the non-rotating Sun is constant and, thus, 47 the only source of irradiance variability is the solar rotation (hereafter referred as the 48 lighthouse approximation). Then, to obtain the irradiance entering the atmosphere of 49 a planet, the measurements made from Earth are either phase-shifted (see Section 3.1) 50 or interpolated (see Section 3.2) according to the relative position of the Earth and the 51 planet and then scaled to the Sun-planet distance (Thiemann et al., 2017, 2021). 52

There have been efforts to test the accuracy of the lighthouse approximation in the 53 literature. For example, Thiemann et al. (2021) used measurements from the Extreme 54 Ultraviolet Monitor (EUVM, Eparvier et al., 2015) onboard the Mars Atmosphere and 55 Volatile EvolutioN (MAVEN) spacecraft to conclude that the error introduced by the 56 lighthouse approximation can be comparable to the amplitude of the solar rotational vari-57 ability. Quémerais et al. (2019) developed an algorithm which combines measurements 58 from the Solar Wind ANisotropies (SWAN) Solar Heliospheric Observatory (SOHO) Lyman-59  $\alpha$  photometer with Earth-based solar flux measurements to derive the irradiance at Lyman-60  $\alpha$  for any direction in the ecliptic plane. They found that the SWAN-based method is 61 superior to the interpolation at periods of high solar activity. On the contrary, the in-62 terpolation method works better than the SWAN-based method at periods of low solar 63 activity. 64

The lighthouse approximation introduces an error that depends on the angle within 65 the ecliptic plane between Earth and the planet of interest, centered on the Sun. This 66 Earth-Sun-planet angle  $(\phi)$  is shown in Figure 1. This error is a consequence of the con-67 stantly evolving distribution of magnetic fields on the solar surface. The magnetic fields 68 emerge from the interior of the Sun, forming active regions on the surface, which evolve 69 and decay on timescales from days to months (see e.g. Solanki et al., 2013, and refer-70 ences therein). In the absence of active regions, the solar irradiance is constant. In this 71 case, the solar irradiance measured at Earth can be scaled to other positions in the So-72 lar System with a simple function of distance. The active regions, however, make the so-73 lar surface heterogeneous and lead to anisotropic solar irradiance, as well as the tempo-74 ral variability of the irradiance. 75



Figure 1. Earth-Sun-planet angle  $(\phi)$ , within the ecliptic plane. The Earth and the planet of interest are depicted by the symbols  $\oplus$  and the full circle, respectively, and the star depicts the Sun. Solar rotation is anti-clockwise.

There are two main sources of solar variability on the timescales from days to months. 76 The first is brought about by solar rotation: the active regions rotate in and out of view, 77 similar to the behavior of a lighthouse. The second is attributed to the evolution of ac-78 tive regions. Nèmec et al. (2020) showed that the variability of the solar irradiance in 79 the visible spectral domain is dominated by rotation for timescales between 4-5 days and 80 the synodic solar rotation period (27.3 days). For timescales shorter than 4-5 days or 81 longer than a solar rotation, the variability is dominated by the evolution and emergence 82 pattern of magnetic features, which is not taken into consideration by the lighthouse ap-83 proximation. 84

In this study, we present a method for estimating solar irradiance at different po-85 sitions within the ecliptic plane of the Solar System, for the total solar irradiance (TSI, 86 i.e. spectrally-integrated irradiance) and for the S-index (a proxy for near-ultraviolet). 87 We take into consideration the evolution of active regions in the Sun and follow an ap-88 proach rooted in the Spectral And Total Irradiance REconstruction model (SATIRE, Fligge 89 et al., 2000; Krivova et al., 2003, 2011). Our SATIRE-inspired approach is described in 90 Section 2 and then compared to both simple shifting of irradiance measurements and the 91 interpolation method in Section 3. The conclusions of this work are drawn in Section 4. 92

## 93 2 Methods

98

In order to calculate the TSI and the S-index, we combine spectra of the quiet Sun (i.e. regions on the solar surface free from any visible manifestations of magnetic activity) and the magnetic features with their corresponding area coverages on the visible solar disc. Each of these aspects will be covered in this section.

2.1 Spectral and total solar irradiance

<sup>99</sup> The SATIRE-inspired approach makes use of the intensity of the quiet Sun and of <sup>100</sup> magnetic features, weighted by their areas on the solar disc. It takes into account the <sup>101</sup> distribution and time evolution of magnetic features on the full solar surface so that the <sup>102</sup> irradiance variability in any given direction can be directly calculated. The approach is

based on the assumption that solar irradiance variations on timescales longer than one

day can be explained by variations in the surface magnetic fields (Fligge et al., 2000; Krivova et al., 2003).

In this approach, the spectral solar irradiance (SSI) at the Earth-Sun-planet angle  $\phi$  (see Figure 1), at an orbital inclination  $\theta$ , for a certain time t and wavelength  $\lambda$  is computed as the sum of two components:

$$E^{\phi\theta}(t,\lambda) = E_{q}(\lambda) + E_{a}^{\phi\theta}(t,\lambda).$$
(1)

Then, the TSI is obtained by integrating the SSI calculated using Eq. 1 over all wavelengths.

In Eq. 1,  $E_q(\lambda)$  is the time-independent spectral irradiance from the quiet Sun.  $E_q(\lambda)$  does not depend on  $\phi$  and  $\theta$ , and is given by

$$E_{\mathbf{q}}(\lambda) = \sum_{i} I_{\mathbf{q}}(\lambda, \mu_{i}) \Delta \Omega_{i}, \qquad (2)$$

where  $I_q(\lambda, \mu_i)$  is the intensity emerging from the quiet Sun for a given  $\lambda$  at a disc position given by  $\mu_i$ , where  $\mu$  is the cosine of the heliocentric angle. The solar surface is divided into many pixels indexed with *i* (see Section 2.2) and  $\Delta\Omega_i$  is the solid angle subtended by the *i*-th pixel as seen from the planet.

The term  $E_{\rm a}^{\phi\theta}(t,\lambda)$  in Eq. 1 represents the contribution to the SSI from magnetic activity, and can be decomposed as

$$E_{a}^{\phi\theta}(t,\lambda) = \sum_{i} \alpha_{u}^{i\phi\theta}(t) [I_{u}(\lambda,\mu_{i}) - I_{q}(\lambda,\mu_{i})] \Delta\Omega_{i} + \sum_{i} \alpha_{p}^{i\phi\theta}(t) [I_{p}(\lambda,\mu_{i}) - I_{q}(\lambda,\mu_{i})] \Delta\Omega_{i} + \sum_{i} \alpha_{f}^{i\phi\theta}(t) [I_{f}(\lambda,\mu_{i}) - I_{q}(\lambda,\mu_{i})] \Delta\Omega_{i}.$$
(3)

Here,  $I(\lambda, \mu_i)$  are the intensities of the magnetic features, namely, sunspot umbrae (u), sunspot penumbrae (p), and faculae (f). Unruh et al. (1999) have shown that including only the quiet Sun, sunspots, and faculae in the computations is adequate to explain the observed SSI and TSI variations (see e.g. Solanki et al., 2013; Ermolli et al., 2013). We employ the intensities computed by Unruh et al. (1999) with the use of the ATLAS9 radiative transfer code (Kurucz, 1992; Castelli & Kurucz, 1994), under the assumption of local thermodynamic equilibrium (LTE).

The intensities are derived for 1221 wavelengths, ranging from 9.09 nm to 10020 nm, with varying spectral resolution, but only values computed above 160 nm can be trusted, due to the assumption of LTE in ATLAS9 (Tagirov et al., 2019). To account for the centreto-limb variation of the intensity, the intensities are calculated for 11 positions on the solar disc corresponding to  $\mu = 1.0$  (disc center), 0.9, 0.8, 0.7, 0.6, 0.5, 0.4, 0.3, 0.2, 0.1, and 0.05. For intermediate disc positions, i.e. when  $\mu_i$  of the pixel is different from these 11 values, the intensities are interpolated to the value of  $\mu_i$ .

In Eq. 3,  $\alpha^{i\phi\theta}(t)$  are the fractional area coverages per pixel by a given magnetic feature. Depending on  $\phi$  and  $\theta$ , an observer sees different parts of the solar surface, i.e., the visible solar disc changes. The dependence of the SSI on the observer's location comes solely from  $E_{a}^{\phi\theta}(t,\lambda)$ , because the disc area coverages of sunspots and faculae depend on  $\phi$  and  $\theta$ . Therefore, the calculation of the SSI for different  $\phi$  and  $\theta$  angles requires that the distribution of sunspots and faculae over the entire solar surface is known, in addition to their intensities.

In the following section, we describe the method used to obtain the area coverages 133 of magnetic features that are needed to compute the SSI and TSI at various locations 134 of the observer. Several studies have used this method to model solar irradiance at ar-135 bitrary inclinations (see e.g. Nèmec et al., 2020; Sowmya, Shapiro, et al., 2021). We note 136 that for any  $\phi$  and  $\theta$ , an observer sees only half of the solar surface and therefore the sum-137 mations in Eqs. 2 and 3 are carried out only over the pixels on the visible disc. Further-138 more, to compare our results with those which use measurements made exclusively on 139 Earth, we limit our analysis to the solar disc as observed from the ecliptic plane, and there-140 fore the superscript  $\theta$  will be omitted from now on. 141

## 2.2 Area coverages

142

We use the spots and faculae area coverage maps obtained by Nèmec et al. (2020), who utilised the Surface Flux Transport Model (SFTM) in the form presented in Cameron et al. (2010). We summarise their approach here briefly, but refer to Nèmec et al. (2020) for details.

In the SFTM, magnetic flux emerges as bipolar magnetic regions and, once emerged, 147 the flux evolves due to advection, diffusion and large scale flows (i.e. the meridional flow 148 and differential rotation). The emergence of the bipolar magnetic regions is based on the 149 semi-empirical input record of sunspot groups, as described in Jiang et al. (2011), with 150 the addition that the emergence longitude of the bipolar magnetic regions is randomised 151 in order to get the full surface distribution of the regions, as needed for our purpose. We 152 note that the statistical input record preserves the latitudes of emergence (i.e. the so-153 lar butterfly diagram), but due to the nature of the semi-empirical record, the actual sizes 154 and emergence of the active regions cannot be directly compared on a day-to-day ba-155 sis to the observed distribution on the Sun. While the record by Jiang et al. (2011) dates 156 back to 1700, we only utilise the synthesized solar cycle 22, which represents a cycle of 157 intermediate activity level. 158

Subsequently, the flux distribution is converted into spot and faculae areas. Fol-159 lowing the input record, sunspots are immediately emerged at their largest size, i.e. we 160 neglect the growing phase. On the day of emergence, Nèmec et al. (2020) calculated the 161 size of the spots using the given areas in the input record, but needed to add a simple 162 model to take into account the growth and decay of the spots. Penumbral area cover-163 ages are taken as 80% of the spot areas, while the remaining 20% is attributed to um-164 brae. Using the spatial and temporal distribution of spots, the magnetic flux in each pixel 165 of the full surface map is corrected for the spot flux. The remaining magnetic field in 166 the pixel is ascribed to faculae. The area coverages of faculae are determined following 167 the saturation threshold approach of the SATIRE model (Fligge et al., 2000; Krivova et 168 al., 2003). From the full surface maps of area coverages, the coverages on the visible disc 169 for any  $\phi$  can be computed by making appropriate geometrical transformations (see e.g. 170 Sowmya, Nèmec, et al., 2021). 171

Figure 2 illustrates an example of the synthetic distribution of magnetic features on the full solar surface and the visible discs corresponding to  $\phi$  angles of 0°, 90°, 180°, and 270°. The spatial distribution as well as the area coverages of spots and faculae on the visible disc are very different depending on  $\phi$ , as clearly shown by the orthographic projections in Figure 2.

## 177 **2.3 S-index**

The Mg II spectral lines around 280 nm exhibit one of the largest solar irradiance variability in the near-UV. Due to this, the Mg II index (core to wing ratio) is used as a proxy for the irradiance variability in the UV and EUV regime. The Mg II index is also shown to be a better proxy for solar EUV than the widely used F10.7 index (Viereck



Figure 2. Snapshot of a synthetic distribution of faculae (red shades) and sunspots (blue pixels). Top: full surface map. Bottom: orthographic projections centered at the  $\phi$  angle indicated above each panel. An observer placed at  $\phi = 0^{\circ}$  views the visible solar disc as in panel (a), for example.

et al., 2001). Very accurate and high resolution spectra of the Mg II lines synthesized in non-LTE as a function of  $\mu$  are needed to correctly determine the Mg II index and its dependence on  $\phi$ . Since such detailed modelling is currently unavailable and beyond the scope of this work, we instead use the Ca II H and K lines to investigate the irradiance changes in the near-UV. Such a choice is directly driven by the strong linear correlation between the irradiance in the Mg II and Ca II lines (DeLand & Marchenko, 2013).

We quantify the variability in Ca II via the well-known chromospheric activity indicator, namely the S-index (Vaughan et al., 1978). The S-index is calculated as the ratio of the summed fluxes in the Ca II H and K line cores to the summed fluxes in two spectral regions near the blue wing of the K line and the red wing of the H line. This index was established by the Mt. Wilson Observatory, and has been recorded for lower main-sequence stars since 1966 (Wilson, 1978).

The formation of Ca II H and K lines spans a wide range of heights in the solar atmosphere. The line cores are formed in the chromosphere, above the solar photosphere, where LTE assumptions are no longer valid. Sowmya, Shapiro, et al. (2021) followed a non-LTE approach to compute the emergent intensities of the quiet Sun and faculae, using the RH code (Uitenbroek, 2001). Here, again, the intensities are computed for different disc position, as explained in Section 2.1, in order to account for center-to-limb variations.

We use the spectra from Sowmya, Shapiro, et al. (2021) in Eqs. 2 and 3 to determine the near-UV SSI variability, neglecting the contribution from spots, since the variability in the Ca II H and K lines are mainly driven by faculae, as shown by Sowmya, Shapiro, et al. (2021) and Sowmya et al. (2023). The S-index is then obtained according to Eq. 1 of Sowmya, Shapiro, et al. (2021).

## <sup>206</sup> 3 Results and discussion

The synthetic calculations of TSI and S-index in this work are performed for two one-year periods within solar cycle 22, using the methods presented in Sections 2.1 and 2.3. The first year is a period of high solar activity and it is centered on the day with the maximum disc area coverage of faculae during the cycle. The second is a period of low solar activity and comprises the last 365 days of solar cycle 22. Figure 3 shows the obtained TSI for the whole solar cycle 22, as well as the chosen periods of high and low solar activity.



**Figure 3.** Synthetic TSI based on the magnetic flux of the visible disc for solar cycle 22. The one-year time periods of high and low solar activity used in this work are marked by red dashed lines.

We define the normalized variability on the time scale of a solar rotation as

$$\Delta E^{\phi}(t,\lambda) = \frac{E^{\phi}(t,\lambda) - \langle E^{\phi}(t,\lambda) \rangle_{81}}{\langle E^{\phi}(t,\lambda) \rangle_{81}},\tag{4}$$

where  $\langle E^{\phi}(t,\lambda) \rangle_{81}$  is the moving average with an 81-days time window centered on t. We use a time window of 81-days, which is the equivalent to three solar rotations, in order to remove the effects that arise due to the transit of the magnetic features.

For the S-index, we simply calculate the rotational variability as

$$\Delta S^{\phi}(t) = S^{\phi}(t) - \langle S^{\phi}(t) \rangle_{81}, \tag{5}$$

where  $S^{\phi}(t)$  is the S-index at time t and angle  $\phi$ , and  $\langle S^{\phi}(t) \rangle_{81}$  is the 81-days moving average.

#### 3.1 Phase shifts and cross-correlation analysis

In the phase shift approach, the irradiance at the planet of interest is obtained by simply phase shifting the irradiance measurements made at Earth and scaling them to the Sun-planet distance. In this approach, we neglect the evolution of active regions and assume that in the absence of solar rotation, the solar irradiance is constant.

Here, we perform an experiment to assess the accuracy of the phase shift approach at three different  $\phi$  angles in the plane of the solar equator: 90°, 180°, and 270° (see Figure 1). According to the phase shift approach, the irradiance values at these angles are

$$E^{90^{\circ}}(t,\lambda) = E^{0^{\circ}}(t-7\ d,\lambda)$$
(6)

$$E^{180^{\circ}}(t,\lambda) = E^{0^{\circ}}(t-14\ d,\lambda)$$
(7)

$$E^{270^{\circ}}(t,\lambda) = E^{0^{\circ}}(t-21\ d,\lambda),\tag{8}$$

with t in days (d). The time periods are equivalent to the angular distance, assuming the synodic solar rotation to be ~ 28 days. The irradiance at  $\phi = 90^{\circ}$  is expected to be the irradiance at  $\phi = 0^{\circ}$  a quarter of a solar rotation (seven days) earlier, and so on for the other  $\phi$  angles.

Figure 4 shows the comparison between TSI values calculated with the phase shift approach (i.e. with Eqs. 6–8, see blue, green, and red curves) and with the SATIREbased approach (i.e. with Eqs. 1–3, see black curves). The comparison is given for periods of high (left panels) and low (right panels) solar activity during solar cycle 22. We note that the dips in the time series are indicative of spot transits. Since the spots are dark, they reduce the TSI. On the contrary, the peaks are caused by faculae, which are bright regions that increase the TSI.

The variability of the S-index computed with the phase shift and with SATIREbased approaches is shown in Figure 5, following the same color scheme as in Figure 4. We note that the S-index calculations are performed by neglecting spots, since Sowmya et al. (2023) showed that the solar S-index and its variability are mainly driven by faculae. Hence, the variability seen in Figure 5 are attributed to faculae. The solar rotational cycles are clearly visible in the time series at both high and low activity periods.

In Table 1, we show the linear Pearson correlation coefficients obtained for each 244 of the cases presented in Figures 4 and 5. The correlation between the reference and the 245 shifted TSI time series is poor during both periods of solar activity. In addition, the cor-246 relation worsens with increasing  $\phi$ . This can be attributed to spots which dominate TSI 247 at shorter timescales. The spots are short-lived and evolve quickly on the solar surface. 248 Because of that, the visible disc at various  $\phi$  differs considerably and the correlation be-249 comes low when a shift of the order of a few days is applied. This suggests that a sim-250 ple phase shift of the irradiance time series is not a good solution for determining the 251 TSI and its variability at different positions within the Solar System ecliptic plane. 252

The S-index curves at  $\phi = 0^{\circ}$  and at  $\phi = 90^{\circ}$  shifted by seven days (blue curves) 253 show a very good correlation, with linear correlation coefficients above 0.9. The differ-254 ence between these two time series at  $\phi = 90^{\circ}$  and the series at  $\phi = 0^{\circ}$  arises from 255 the evolution of faculae over the course of seven days. Since the faculae survive on the 256 solar surface longer than most spots, the time series recorded seven days apart still sees 257 most of the original faculae features and thus the correlation between the time series is 258 high. With increasing  $\phi$ , the correlation between the time series generally decreases, al-259 though the correspondence is still reasonably high. This indicates that shifting the so-260



Figure 4. Synthetic time series of the normalized TSI variability for periods of high (left) and low (right) solar activity. The panels show the calculated (black) and phase shifted (colors) variability. From top to bottom, the panels show  $\phi = 90^{\circ}$  (blue),  $\phi = 180^{\circ}$  (green), and  $\phi = 270^{\circ}$  (red).



Figure 5. The same as in Figure 4, but for the S-index variability.

lar irradiance time series measured at Earth to other positions within the ecliptic planeworks relatively better for S-index than for TSI.

We also determine the cross-correlation between the time series. With this analysis, our aim is to find how many days (forward or backward) the irradiance time series at  $\phi = 90^{\circ}, 180^{\circ}$ , and 270° should be shifted in order to achieve the highest correlation

**Table 1.** Correlation Coefficients Between the Calculated and the Phase Shifted Time Series ofIrradiance.

	High a	High activity		ctivity
$\phi$	TSI	S-index	TSI	S-index
90°	0.39	0.92	0.47	0.93
$180^{\circ}$	0.10	0.84	0.17	0.75
$270^{\circ}$	-0.10	0.75	0.18	0.63

 Table 2.
 Results of the Cross-Correlation Analysis Between the Time Series.

	High activity				Low a	activity		
	TSI		S-index		TSI		S-index	
$\phi$	Shift (days)	r	Shift (days)	r	Shift (days)	r	Shift (days)	r
$90^{\circ}$	-6	0.45	-6	0.92	-7	0.47	-6	0.94
$180^{\circ}$	-22	0.19	-13	0.87	-5	0.30	+13	0.82
$270^{\circ}$	+4	0.38	+7	0.90	+7	0.41	+7	0.88

Note. Given are the maximum correlation coefficients (r) and their corresponding shifts with respect to the time series at  $\phi = 0^{\circ}$ . Negative shifts indicate preceding days, while positive shifts indicate following days.

coefficient, r, to the time series at  $\phi = 0^{\circ}$ . This is done by shifting the time series day by day, with a time window ranging from a solar rotation in the past to a solar rotation in the future, and comparing them with the time series at  $\phi = 0^{\circ}$ . Table 2 shows the results of the cross-correlation analysis. Negative shifts indicate preceding days (the same logic as in Eqs. 6–8), while positive shifts indicate following days.

As discussed previously, an angular distance of  $\phi = 90^{\circ}$  should correspond approx-271 imately to -7 days,  $\phi = 180^{\circ}$  to -14 days, and  $\phi = 270^{\circ}$  to -21 days. The cross-correlation 272 analysis gives the expected result for  $\phi = 90^{\circ}$ , even though the coefficients are relatively 273 low for the TSI. For higher  $\phi$  angles, the maximum correlation coefficients for the TSI 274 do not seem to have any physical meaning, since the shifts are very distant from what 275 is expected. As for the S-index, some of the maximum coefficients are achieved when the 276 time series are shifted one entire solar rotation ahead of what is expected (positive shifts). 277 From Figures 4 and 5 and Tables 1 and 2, we infer that a simple phase shift is not suf-278 ficient to provide good correlation results and, accordingly, irradiance estimates. 279

280

#### 3.2 Comparison with interpolated Earth-based measurements

As described by Thiemann et al. (2017), the interpolation method estimates the irradiance at a certain planet (or angle  $\phi$  within the ecliptic plane) for a day  $t_d$ , by using measurements at Earth  $(E_{\oplus})$  made before  $(t_d - \Delta t_1)$  and after  $(t_d + \Delta t_2)$  the visible solar disc rotates past the planet. The time differences  $\Delta t_1$  and  $\Delta t_2$  follow the same logic as in Eqs. 6 to 8. For example, if the Earth-Sun-planet angle  $\phi$  is 90° today, the irradiance at the planet would be estimated as the weighted sum of the irradiance at Earth  $\sim$ 7 days ago and  $\sim$ 21 days in the future. The interpolation is given by

$$E(t_d, \phi) = \frac{w_1 \ E_{\oplus}(t_d - \Delta t_1) + w_2 \ E_{\oplus}(t_d + \Delta t_2)}{r^2(t_d)},\tag{9}$$

where  $r^2(t_d)$  is the distance between Earth and the planet on day  $t_d$ , in AU. The measurements are linearly interpolated and weighted, and the weights  $w_1$  and  $w_2$  are pro-

**Table 3.** Correlation Coefficients Between the Calculated and the Interpolated Time Series ofIrradiance.

	High activity		Low a	ctivity
$\phi$	TSI	S-index	TSI	S-index
90°	0.30	0.93	0.49	0.93
$180^{\circ}$	0.17	0.93	0.21	0.87
$270^{\circ}$	0.24	0.93	0.45	0.90

<sup>290</sup> portional to the periods  $\Delta t_1$  and  $\Delta t_2$ . In the example given, the weights  $\Delta t_1$  and  $\Delta t_2$ <sup>291</sup> would be 0.75 and 0.25, respectively. For a more detailed explanation of the method, see <sup>292</sup> Section 2.1 from Thiemann et al. (2017).

We apply the interpolation method to the irradiance time series at  $\phi = 0^{\circ}$ , and determine the irradiance at  $\phi = 90^{\circ}$ , 180°, and 270°. Then, we compare these interpolated time series to the ones evaluated at  $\phi = 90^{\circ}$ , 180°, and 270° using our SATIREinspired approach, by considering the Pearson correlation coefficient of the time series at each  $\phi$ . The coefficients are shown in Table 3. Figure 6 shows such a comparison for the TSI during periods of high and low activity. The time series calculated using our SATIREinspired approach at  $\phi$  (black curves) and the values interpolated from  $\phi = 0^{\circ}$  to  $\phi =$  $90^{\circ}$ , 180°, and 270° (in colors) differ greatly.

The interpolation method misses the exact timing of most spot transits (valleys), 301 especially for periods of high solar activity. During periods of low solar activity, extrap-302 olating the data seems to overestimate the duration and/or intensity of the sunspots. For 303 all three positions around the Sun, the correlation between the calculated and the in-304 terpolated TSI variability does not surpass 0.49. This suggests further that the inter-305 polation of the TSI variability observed at Earth to other planets does not represent the 306 true variability seen at those planets. By comparing Figures 4 and 6, however, the val-307 leys and peaks characteristics of each time series are better reconstructed with the in-308 terpolation than with the phase shifting method. 309

Following the same scheme, Figure 7 shows the time series of the S-index variability at high and low activity periods. Contrary to the TSI case, the correlations between the calculated and interpolated S-index variability are high, and the curves look very similar. Again, the shapes of the interpolated irradiance time series are more in agreement to the ones obtained with the SATIRE-inspired approach than the ones estimated with the phase shifting method (Figure 5).

On one hand, the interpolation method works very well for the proxy of UV wavelengths because facular regions evolve very slowly, so that the evolution variability they induce is not significant on solar rotation timescales. On the other hand, the application of the interpolation method for the TSI or wavelengths which are dominated by sunspots would result in a very large error in the estimated irradiance.

#### <sup>321</sup> 4 Summary and conclusions

In this study, we present a method for estimating solar irradiance within the ecliptic plane. First, we use a synthetic statistical model of magnetic field transport, as well as semi-empirical information on active regions sources in order to build a synthetic model of faculae and spots area coverages observed at any position within the ecliptic plane during solar cycle 22. Then, we compute synthetic values of spectral solar irradiance at several Earth-Sun-planet angles in the Solar System using a method that is built on the SATIRE model for solar irradiance variability reconstruction.



Figure 6. Synthetic time series of the normalized TSI variability for periods of high (left) and low (right) solar activity. The panels show the variability calculated using our SATIRE-inspired approach (black) and the interpolation method (colors). From top to bottom, the panels show  $\phi = 90^{\circ}$  (blue),  $\phi = 180^{\circ}$  (green), and  $\phi = 270^{\circ}$  (red).



Figure 7. The same as in Figure 6, but for the S-index variability.

<sup>329</sup> Using our SATIRE-inspired synthetic model, we evaluate the TSI and the S-index, <sup>330</sup> which is a proxy for irradiance at near-UV wavelengths, at four Earth-Sun-planet an-<sup>331</sup> gles:  $\phi = 0^{\circ}, 90^{\circ}, 180^{\circ}$ , and 270°, for a year of high solar activity and a year of low so-<sup>332</sup> lar activity. We apply a simple phase shift to the time series at  $\phi = 90^{\circ}, 180^{\circ}$ , and 270°, <sup>333</sup> proportional to the solar rotation, and compare the results with the time series at  $\phi =$  <sup>334</sup> 0°. We conclude that a simple phase shift of the data is not enough to obtain good es-<sup>335</sup> timates of solar irradiance.

We also compared our method with the traditional interpolation method (Thiemann et al., 2017). This method consists of a "lighthouse" model, where Earth-based measurements of SSI are interpolated to other planets assuming that irradiance variability is only caused by solar rotation.

We investigate the correlation of the time series computed with our SATIRE-inspired 340 approach and the interpolated time series. The results show that there is a high corre-341 lation between the interpolation and the SATIRE-inspired approach when it comes to 342 the S-index. Due to the relatively greater lifetime of faculae, which are the sources of 343 solar UV variability, the interpolation method is able to catch slow changes in irradiance. 344 However, the correlation drops significantly for the TSI, since it is dominated by sunspots, 345 which emerge, evolve, and decay very fast. In this case, the interpolation method can-346 not properly account for the effect of spots, which are relatively short-lived and do not 347 survive a full solar rotation. The SATIRE-inspired approach directly calculates the TSI, 348 accounting for effects of faculae and spots, which is advantageous compared to the in-349 terpolation method. 350

We point out that our SATIRE-inspired approach for evaluating irradiance is still based on a synthetic model of full surface solar magnetograms and we used statistically valid emergences of magnetic regions to calculate the irradiance, and not real data. Therefore, the model is not capable of producing a nowcast estimation of solar irradiance. In the future, we will use information of far-side helioseismology to derive the actual area coverages of magnetic regions.

<sup>357</sup> Open Research Section

The area coverages data and flux intensities used for the calculations of irradiance in the study are available in De Oliveira (2024).

360 Acknowledgments

I. de O. would like to thank Laurent Gizon and Dan Yang for their contribution on re-

viewing the draft of this paper. This work was supported by the International Max Planck
 Research School for Solar System Science at the University of Göttingen.

## 364 References

- Cameron, R. H., Jiang, J., Schmitt, D., & Schüssler, M. (2010, August). Surface
   Flux Transport Modeling for Solar Cycles 15-21: Effects of Cycle-Dependent
   Tilt Angles of Sunspot Groups. *The Astrophysical Journal*, 719(1), 264-270.
   doi: 10.1088/0004-637X/719/1/264
- Castelli, F., & Kurucz, R. L. (1994, January). Model atmospheres for Vega. Astronomy & Astrophysics, 281, 817-832.
- DeLand, M., & Marchenko, S. (2013, April). The solar chromospheric Ca and Mg indices from Aura OMI. *Journal of Geophysical Research (Atmospheres)*, 118(8), 3415-3423. doi: 10.1002/jgrd.50310
- 374De Oliveira, I.(2024, February).Data from: Estimation of solar irradiance375in the ecliptic plane using synthetic solar surface magnetograms.Zen-376odo.Retrieved from https://doi.org/10.5281/zenodo.10630214doi:37710.5281/zenodo.10630214
- Eparvier, F. G., Chamberlin, P. C., Woods, T. N., & Thiemann, E. M. B. (2015, December). The Solar Extreme Ultraviolet Monitor for MAVEN. Space Science Reviews, 195(1-4), 293-301. doi: 10.1007/s11214-015-0195-2

381	Ermolli, I., Matthes, K., Dudok de Wit, T., Krivova, N. A., Tourpali, K., Weber, M.,
382	Woods, T. N. (2013). Recent variability of the solar spectral irradiance and
383	its impact on climate modelling. Atmospheric Chemistry & Physics, $13(8)$ ,
384	3945-3977. doi: $10.5194/acp-13-3945-201310.5194/acpd-12-24557-2012$
385	Fligge, M., Solanki, S. K., & Unruh, Y. C. (2000, January). Modelling irradiance
386	variations from the surface distribution of the solar magnetic field. Astronomy
387	& Astrophysics, 353, 380-388.
388	Fox, J. L., Galand, M. I., & Johnson, R. E. (2008, August). Energy Deposition in
389	Planetary Atmospheres by Charged Particles and Solar Photons. Space Science
390	Reviews, 139(1-4), 3-62. doi: 10.1007/s11214-008-9403-7
391	Gray, L. J., Beer, J., Geller, M., Haigh, J. D., Lockwood, M., Matthes, K.,
392	White, W. (2010, October). Solar Influences on Climate. Reviews of Geo-
393	pnysics, 48(4), RG4001. doi: 10.1029/2009RG000282
394	Jiang, J., Cameron, R. H., Schmitt, D., & Schussler, M. (2011, April). The so-
395	ar magnetic field since 1700. II. Physical reconstruction of total, polar and
396	open nux. Astronomy & Astrophysics, 528, A85. doi: 10.1051/0004-0301/ 201016168
397	Krivova N A Solanki S K Eliggo M & Unruh V C (2003 February)
398	Reconstruction of solar irradiance variations in cycle 23: Is solar surface
399	magnetism the cause? Astronomy & Astronbusics 399 L1-L4 doi:
401	10.1051/0004-6361:20030029
402	Krivova, N. A., Solanki, S. K., & Unruh, Y. C. (2011, February). Towards a long-
403	term record of solar total and spectral irradiance. Journal of Atmospheric and
404	Solar-Terrestrial Physics, 73(2-3), 223-234. doi: 10.1016/j.jastp.2009.11.013
405	Kurucz, R. L. (1992, March). Remaining Line Opacity Problems for the Solar Spec-
406	trum. Revista Mexicana de Astronomia y Astrofísica, 23, 187.
407	Nèmec, N. E., Shapiro, A. I., Krivova, N. A., Solanki, S. K., Tagirov, R. V.,
408	Cameron, R. H., & Dreizler, S. (2020, April). Power spectra of solar brightness
409	variations at various inclinations. Astronomy & Astrophysics, 636, A43. doi:
410	10.1051/0004- $6361/202037588$
411	Peter, K., Pätzold, M., Molina-Cuberos, G., Witasse, O., González-Galindo, F.,
412	Withers, P., Tyler, G. L. (2014). The dayside ionospheres of Mars and
413	Venus: Comparing a one-dimensional photochemical model with MaRS (Mars
414	Express) and VeRa (Venus Express) observations. <i>Icarus</i> , 233, 66-82. Re-
415	trieved from https://www.sciencedirect.com/science/article/pii/
416	S0019103514000591 doi: https://doi.org/10.1016/j.icarus.2014.01.028
417	Quemerais, E., 1 niemann, E., Snow, M., Ferron, S., & Schmidt, W. (2019).
418	ing SOHO SWAN and MAVEN FUVM Lymph & Measurements
419	and of Geophysical Research: Space Physics $12/(6)$ 3040-3060 doi:
420	https://doi.org/10.1029/2019JA026674
421	Ramstad B. Barabash S. Futaana V. Nilsson H. Wang XD. & Holmström
422	M (2015) The Martian atmospheric ion escape rate dependence on solar
423	wind and solar EUV conditions: 1. Seven years of Mars Express observations.
425	Journal of Geophysical Research: Planets, 120(7), 1298-1309. Retrieved
426	from https://agupubs.onlinelibrary.wiley.com/doi/abs/10.1002/
427	2015JE004816 doi: https://doi.org/10.1002/2015JE004816
428	Solanki, S. K., Krivova, N. A., & Haigh, J. D. (2013, August). Solar Irradiance Vari-
429	ability and Climate. Annual Review of Astronomy and Astrophysics, 51(1),
430	311-351. doi: 10.1146/annurev-astro-082812-141007
431	Sowmya, K., Nèmec, N. E., Shapiro, A. I., Işık, E., Witzke, V., Mints, A.,
432	Solanki, S. K. (2021, October). Predictions of Astrometric Jitter for Sun-like
433	Stars. II. Dependence on Inclination, Metallicity, and Active-region Nesting.
434	The Astrophysical Journal, 919(2), 94. doi: 10.3847/1538-4357/ac111b
435	Sowmya, K., Shapiro, A. I., Rouppe van der Voort, L. H. M., Krivova, N. A., &

436	Solanki, S. K. (2023, October). Modeling Stellar Ca II H and K Emis-
437	sion Variations: Spot Contribution to the S-index. $, 950(1), L10.$ doi:
438	10.3847/2041-8213/act92a
439	Sowmya, K., Snapiro, A. I., Witzke, V., Nemec, N. E., Chatzistergos, I., Yeo, K. L.,
440	Solanki, S. K. (2021, June). Modeling Stellar Ca II H and K Emission
441	Variations. I. Effect of Inclination on the S-index. The Astrophysical Journal, 01/(1) $01$ $1$ $10$ $2047/1520$ $4257/16247$
442	914(1), 21.  doi:  10.3847/1538-4357/abf247
443	Tagirov, R. V., Snapiro, A. I., Krivova, N. A., Unrun, Y. U., Yeo, K. L., & Solanki,
444	S. K. (2019, November). Readdressing the UV solar variability with $CATELEE = C$
445	SATIRE-S: non-LIE effects. Astronomy & Astrophysics, 031, A178. doi:
446	10.1051/0004-0301/201935121 Thismann E. Chambarlin D. C. Francisco F. C. Tamalanan D. Waada T. N.
447	Deurshen C. W. & Jakashur, P. M. (2017) The MAVEN EUVM model of
448	bougher, S. W., & Jakosky, D. M. (2017). The MAVEN ED VM model of
449	solar spectral irradiance variability at Mars: Algorithms and results. $Jour-$
450	from https://ammuha.enlipeliberen viles.cs/doi/abc/10.1002/
451	1000000000000000000000000000000000000
452	$ \begin{array}{c} 2010 \text{JA023512} & \text{doi. https://doi.org/10.1002/2010JA023512} \\ \text{Thiomann E Ensuring E C Knoor V Al Muhamani A & IIilia D I (2021) \\ \end{array} $
453	Solar Extreme Illtraviolet Irradiance Uncertainties for Planetary Studies
454	Journal of Geophysical Research: Space Physics $126(1)$ $_{2}0201\Lambda028184$
455	Batrieved from https://agupubs.onlinelibrary.wiley.com/doi/abs/
450	10 1029/2020 IA028184 doi: https://doi.org/10.1029/2020 IA028184
457	Uitenbroek H (2001 August) Multilevel Badiative Transfer with Partial Frequency
450	Bedistribution The Astrophysical Journal 557(1) 389-398 doi: 10.1086/
459	321659
461	Unruh, Y. C., Solanki, S. K., & Fligge, M. (1999, May). The spectral dependence
462	of facular contrast and solar irradiance variations. Astronomy & Astronhysics.
463	3/5. 635-642.
464	Vaughan, A. H., Preston, G. W., & Wilson, O. C. (1978, June). Flux measurements
465	of Ca II and K emission. Publications of the Astronomical Society of the Pa-
466	cific, 90, 267-274. doi: 10.1086/130324
467	Viereck, R., Puga, L., McMullin, D., Judge, D., Weber, M., & Tobiska, W. K.
468	(2001). The Mg II index: A proxy for solar EUV. <i>Geophysical Research</i>
469	Letters, 28(7), 1343-1346. Retrieved from https://agupubs.onlinelibrary
470	.wiley.com/doi/abs/10.1029/2000GL012551 doi: https://doi.org/10.1029/
471	2000GL012551
472	Wilson, O. C. (1978, December). Chromospheric variations in main-sequence stars.
473	226, 379-396. doi: 10.1086/156618