

<b>TABLE 1</b> Study demographics	
Variable	N = 23 No. (%)
<b>Sex</b>	
Male	12 (52.2)
Female	11 (47.8)
<b>Age at Diagnosis, years</b>	
Median [Minimum-Maximum]	9 [5-19]
Mean $\pm$ SD	9.7 $\pm$ 4.4
5-8	11 (47.8)
9-13	7 (30.4)
14-19	5 (21.7)
<b>Ethnicity</b>	
Non-Hispanic	13 (56.5)
Hispanic	10 (43.5)
<b>Race</b>	
White	12 (52.2)
Black or African American	3 (13.0)
Asian	2 (8.7)
Other	5 (21.7)
Unknown	1 (4.3)
<b>Primary Language</b>	
English	17 (73.9)
Spanish	6 (26.1)
<b>Final Risk Group</b>	
High Risk <sup>a</sup>	9 (39.1)
Intermediate Risk	3 (13.0)
Low Risk	11 (47.8)
<b>BMI at Diagnosis</b>	
Underweight (<5th percentile)	3 (13.0)
Normal weight (5-84th percentile)	13 (56.5)
Overweight (85-94th percentile)	2 (8.7)
Obese ( $\geq$ 95th percentile)	5 (21.7)
<b>Timing of Initiation of Nutrition Counseling</b>	
Study Entry	19 (82.6)
End of Induction	4 (17.4)
<sup>a</sup> High risk group includes very high risk (n=6) and high risk (n=3). Abbreviation: SD, standard deviation; BMI, body mass index.	