

## **Title**

The effect of COVID-19 stay-at-home order and campus closure on the prevalence of acute respiratory infection symptoms in college campus cohorts.

**Short running title:** COVID-19 stay-at-home order and the prevalence of Acute Respiratory Infections (ARI) symptoms.

## **Authors**

Oluwasanmi Adenaiye<sup>1</sup>, P. Jacob Bueno de Mesquita<sup>1</sup>, Qiong Wu<sup>2</sup>, Filbert Hong<sup>1</sup>, Jianyu Lai<sup>1,2</sup>, Shuo Chen<sup>3</sup>, Donald K. Milton<sup>1\*</sup> and Prometheus@UMD Consortium

<sup>1</sup>Maryland Institute for Applied Environmental Health, and

<sup>2</sup>Department of Epidemiology and Biostatistics,

University of Maryland, College Park, MD 20742, USA

<sup>3</sup>Division of Biostatistics and Bioinformatics, School of Medicine, University of Maryland, Baltimore, MD, USA.

## **Acknowledgment**

This study is part of Prometheus-UMD, which was sponsored by the Defense Advanced Research Projects Agency (DARPA) BTO under the auspices of Col. Matthew Hepburn through agreements N66001-18-2-4015, and by the Office of the Assistant Secretary for Preparedness and Response, Biomedical Advanced Research and Development Authority. The findings and conclusions in this report are those of the authors and do not necessarily represent the official position or policy of the funding agency and no official endorsement should be inferred.

## **Abstract**

Evaluation of population-based COVID-19 control measures informs strategies to quell the current pandemic and reduce the impact of those yet to come. Effective COVID-19 control measures may simultaneously reduce the incidence of other acute respiratory infections (ARIs) due to shared transmission modalities. To assess the impact of stay-at-home orders and other physical distancing measures on the prevalence of ARI-related symptoms, we compared symptoms reported by prospective college cohorts enrolled during two consecutive academic years. ARI-related symptoms declined following campus closure and implementation of stay-at-home orders, demonstrating the impact of population-based physical distancing measures on control of a broad range of respiratory infections.

## **Keywords**

COVID-19, Lockdown, Influenza, ARI, transmission mitigation.

## **Introduction**

As part of efforts to prevent the escalation of SARS-CoV-2 transmission, the US Centers for Disease Control and Prevention issued nationwide stay-at-home recommendations in the United States on March 15, 2020.<sup>1</sup> Also, states across the country issued varying degrees of physical distancing measures, including restrictions on transportation and movement, school closures, and business closures. Since then, studies have shown, using mathematical and epidemiological models, that such distancing measures likely reduced transmission of COVID-19.<sup>2,3</sup> Others have shown the effect of these measures on the incidence of other viral acute respiratory infections (ARIs) such as influenza.<sup>4,5</sup> However, only a few community-surveillance studies were ongoing before the stay-at-home orders and can show the effect of these measures on a population already under ARI surveillance. Here, we use data from a prospective college cohort under ARI surveillance to assess the impact of the physical distancing measures on the rate of ARI-related symptoms.

## **Methods**

We recruited new cohorts of college students and staff each academic year for four years and monitored them prospectively for the occurrence of ARI (2017-2020). At the beginning of the study, we asked subjects to complete an online survey on their demography and health history and to provide baseline biological specimens. Subjects were then prospectively monitored for ARI. In the most recent two years (2019 and 2020), we monitored the cohorts for ARI using symptom surveys sent daily at 10 AM via text message to all enrolled subjects. Subjects were asked to rate each of the ARI-related symptoms on the daily text survey on a severity scale of 0 to 3 (3 = most severe), based on how they felt at the time of their response (Figure 1). The daily symptom surveys were part of a larger study wherein participants with ARI-related symptoms were invited to the study clinic to provide mid-turbinate (MT) nasal swab specimens. Swabs were tested for 44 respiratory pathogens using a TaqMan Array Card® (Thermo Fisher, Waltham, MA, USA).<sup>6</sup> Physical environmental conditions in the living spaces of the subjects who tested positive and their contacts were monitored as described by Zhu et al.<sup>7</sup>

The data presented in this report are from the daily symptom surveys for two consecutive spring semesters, 2019, and 2020, when we monitored subjects daily for ARI symptoms. During the 2020 spring semester, each subject was compensated \$1 per day for completing the symptom survey, plus \$5 on a random day in a month if they completed the survey on that day. During the 2019 spring semester, we performed a daily and a weekly lottery among all respondents, and the randomly selected subject from each lottery received \$20 and \$100, respectively.

We created two indicator variables for ARI-related symptoms for each person-day observation: i.) sum of symptom scores  $>3$ , and ii.) presence of self-reported fever, with cough or sore throat. A 3-day simple-moving-average (SMA) was constructed with SMA package<sup>8</sup> from the percentages of respondents who met the criteria for each of the two categories. Plots were created with ggplot2<sup>9</sup> to compare the trends before and after stay-at-home orders. To account for differential average response rates, we calculated the proportion of enrollees who reported symptoms in each of the two symptom categories. Categorical variables were compared across cohort years with chi-squared tests and continuous variables were compared with Mann-Whitney U tests. We tested, using one-dimensional scan statistics<sup>10</sup>, to evaluate differences between the reporting rates of the two groups of ARI-related symptoms between the years 2019 and 2020 (SI).

## **Result**

For the spring semester of 2019, daily symptom surveillance commenced on January 30, 2019, and ended on May 21, 2019. For the spring semester of 2020, the daily symptom surveillance for some cohort members started in December, but the majority enrolled in January and the data in this report includes reports starting on January 27 and ending on May 21, 2020. The characteristics of respondents are summarized in Table 1. Seventy-four percent of the subjects who were monitored in 2020 lived on-campus during the period when the campus was open and residence halls were occupied. The majority recorded Maryland as their state of permanent residence. Figure 2 shows the daily trend in the number of subjects and the proportion of enrollees that responded to the daily text. Figure 3 shows the daily trend of the proportions of respondents and enrollees with ARI-related symptoms for the two academic semesters. Figures S1 and S2 show the epidemic curve of the infections that were detected weekly in the subjects who were tested in the 2019 and 2020 cohorts respectively.

Within the 2019 cohort, matching on response rate in the prior week, subjects who received a lottery payment had a 3.13-fold ( $p < 0.001$ ) greater response rate in the following week than those who did not receive payment. The average completion rate of the daily symptom survey was 80% in the 2020 semester and 50% in 2019 ( $p < 0.0001$ ). There were no observed significant differences in age ( $p > 0.79$ ) and sex ( $p > 0.37$ ) distributions between the subjects in the 2019 and 2020 cohorts. No significant differences were observed between the respondents and non-respondents in 2019 (Age:  $p > 0.89$ , sex:  $p > 0.67$ ).

The ARI-related symptoms detection rate was reduced over the course of the semester in each of the two years. The drop in ARI-related symptoms fell at a higher rate in 2020 compared to 2019, and the drop to minimal ARI-related symptoms detection rate persisted in 2020. There was a steady decrease in the rate of ARI-related symptoms starting from around March 16, 2020. When compared to 2019, we observed a significant reduction in the reporting rates of symptoms with a sum of scores greater than 3 ( $p < 0.05$ ) and reporting rates of fever with cough or sore throat ( $p < 0.0001$ ) starting from March 31 and April 1 respectively (Figure S3, S4). There was no significant difference in the reporting rates of the ARI-related symptoms between the two years before campus closure for either set of symptoms ( $p \geq 0.46$ ).

## **Discussion**

We showed variations in the prevalence of ARI-related symptoms in two college campus cohorts monitored for ARI in the spring semesters of two consecutive academic years. The members of the 2020 cohort with a low prevalence of ARI-related symptoms spent a considerable period observing stay-at-home orders with presumably reduced physical contacts. The members of the 2019 cohort spent the surveillance period living in a large university campus with high numbers of daily interactions and a much higher occupant density setting.

The period of a significant decrease in the prevalence of ARI-related symptoms in the Spring-2020 cohort started from around March 31, 2020; eighteen days after the campus was closed on March 13 and when physical distancing measures were introduced in the state of Maryland where most of the subjects live.<sup>11,12</sup> The difference between the two semesters was no longer significant after April 18 in part due to falling symptom rates in late spring during the reference period (2019) which we believe is because of the decreasing prevalence of influenza infections (Figure S1, S2). The general decrease in 2019 in ARI detection over semester could be related to the seasonality of many ARIs with higher incidence generally observed in the winter compared with the summer season in temperate climates.

We did not observe a substantial reduction in the number of daily survey completions following campus closure in the year 2020. A comparison of response rates over the two years showed that payment for

daily performance was a much more effective means of compensation than a lottery. Thus, by compensating subjects every day that they completed the survey, we positively reinforced the survey-completing behavior. Because we adjusted for the response rate in the analysis comparing ARI-related symptom detection between cohort years 2019 and 2020, we don't believe that these differences in response rate affected the main analysis.

This report offers evidence for the effectiveness of physical distancing measures and campus closure on reducing the incidence of ARI in the campus community. This may explain the large reduction in the number of influenza cases observed in the southern hemisphere in 2020 compared to the same calendar period in 2019.<sup>13</sup> Furthermore, our findings demonstrate the impact of population-based physical distancing measures on control of a broad range of respiratory infections.

## **References**

1. CDC. Communities, Schools, Workplaces, & Events. Centers for Disease Control and Prevention. Published April 30, 2020. Accessed June 7, 2020. <https://www.cdc.gov/coronavirus/2019-ncov/community/large-events/mass-gatherings-ready-for-covid-19.html>
2. Lai S, Ruktanonchai NW, Zhou L, et al. Effect of non-pharmaceutical interventions to contain COVID-19 in China. *Nature*. Published online May 4, 2020:1-7. doi:10.1038/s41586-020-2293-x
3. Chinazzi M, Davis JT, Ajelli M, et al. The effect of travel restrictions on the spread of the 2019 novel coronavirus (COVID-19) outbreak. *Science*. 2020;368(6489):395-400. doi:10.1126/science.aba9757
4. Cowling BJ, Ali ST, Ng TWY, et al. Impact assessment of non-pharmaceutical interventions against coronavirus disease 2019 and influenza in Hong Kong: an observational study. *Lancet Public Health*. 2020;5(5):e279-e288. doi:10.1016/S2468-2667(20)30090-6
5. Wu D, Lu J, Liu Y, Zhang Z, Luo L. Positive effects of COVID-19 control measures on influenza prevention. *Int J Infect Dis*. 2020;95:345-346. doi:10.1016/j.ijid.2020.04.009
6. Harvey JJ, Chester S, Burke SA, et al. Comparative analytical evaluation of the respiratory TaqMan Array Card with real-time PCR and commercial multi-pathogen assays. *J Virol Methods*. 2016;228:151-157. doi:10.1016/j.jviromet.2015.11.020
7. Zhu S, Jenkins S, Addo K, et al. Ventilation and laboratory confirmed acute respiratory infection (ARI) rates in college residence halls in College Park, Maryland. *Environ Int*. 2020;137:105537. doi:10.1016/j.envint.2020.105537
8. Svetunkov I, Petropoulos F. Old dog, new tricks: a modelling view of simple moving averages. *Int J Prod Res*. 2018;56(18):6034-6047. doi:10.1080/00207543.2017.1380326
9. Wickham H. *Ggplot2: Elegant Graphics for Data Analysis*. 2nd ed. Springer International Publishing; 2016. doi:10.1007/978-3-319-24277-4
10. Gotway CA, Waller LA. *Applied Spatial Statistics for Public Health Data*. Vol 368. John Wiley & Sons; 2004.
11. Archived Calendars | Division of Academic Affairs. Accessed June 12, 2020. <https://svp.umd.edu/archived-calendars>
12. Coronavirus timeline: Key dates as the virus spread in DC, Maryland and Virginia. WTOP. Published May 13, 2020. Accessed June 13, 2020. <https://wtop.com/coronavirus/2020/05/coronavirus-timeline-key-dates-as-the-virus-spread-in-dc-maryland-and-virginia/>
13. WHO | Influenza update - 375. WHO. Accessed September 10, 2020. [http://www.who.int/influenza/surveillance\\_monitoring/updates/latest\\_update\\_GIP\\_surveillance/en/](http://www.who.int/influenza/surveillance_monitoring/updates/latest_update_GIP_surveillance/en/)

**Table 1. Characteristics of respondents**

	<b>2020</b>	<b>2019</b>
Surveillance period	January 27 – May 21	January 30 – May 21
Number of subjects sent daily survey <sup>†</sup> Mean $\pm$ SD (Range)	230.2 $\pm$ 31.2 (132 - 249)	507.6 $\pm$ 78.4 (349 - 596)
Number of daily respondents Mean $\pm$ SD (Range)	182.7 $\pm$ 22.1 (106 - 211)	250.1 $\pm$ 15.7 (204 - 284)
Survey completion rate $\pm$ SD <sup>*</sup>	79.8% $\pm$ 5.9	50.4% $\pm$ 8.1
Age in years Mean $\pm$ SD	20.4 $\pm$ 3.3	19.9 $\pm$ 2.7
Female (%)	150 (60.2%)	339 (56.9%)
On-campus residents (%)	183 (73.5%)	476 (79.8%)
Residents of the State of Maryland (%)	225 (90.4%)	372 (82.4%)





SD, Standard Deviation

<sup>†</sup> This number represents the number of subjects that were receiving daily text survey messages. It increased as more people enrolled in the study.

<sup>\*</sup> Survey completion rate, 2020 vs 2019; p-value (<0.0001)


During the 2020 spring semester, each subject was compensated \$1 per day, plus \$5 on a random day in a month, for completing the daily symptom survey. Whereas during the 2019 spring semester, we performed a daily and a weekly lottery among all respondents, and the randomly selected subject from each lottery received \$20 and \$100, respectively.

Please rate the following symptoms for how you feel **right now**.

	0= no symptoms 	1=just noticeable 	2= bothersome 	3= awful 
<b>Runny nose:</b> * must provide value	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
				reset
<b>Sore throat:</b> * must provide value	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
				reset
<b>Cough:</b> * must provide value	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
				reset
<b>Sweats/fever/chills</b> * must provide value	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>	<input type="radio"/>
				reset

**Figure 1.** The daily symptom survey that was sent to study participants.


The symptom score is rated as follows, 0 – No symptoms, 1 – Just noticeable, 2- Bothersome, 3 – Awful.



**Figure 2.** The daily trend of the number of responses received and the proportion of the enrollees that responded to the daily text.

As more people enrolled in the study, the number of sent surveys increased.

The campus closed for spring break on March 13, 2020, and remained closed due to the COVID-19 pandemic, till the end of the surveillance period on May 21, 2020.



**Figure 3.** The daily trend of the proportions of respondents and enrollees with ARI-related symptoms for the two academic semesters. **A.** The proportion of daily respondents reporting a sum of symptoms' score greater than 3. **B.** The proportion of the enrollees reporting a sum of symptoms' score greater than 3. **C.** The proportion of respondents reporting having a fever and either a sore throat or cough, **D.** The



proportion of the enrollees that reported having a fever and either a sore throat or cough. The campus closed on March 13, 2020, and did not reopen during the study due to COVID-19.