

Table 3: Perinatal outcomes of 1061 pregnant women randomly assigned to undergo an additional ultrasound examination at 35-37 weeks' gestation (study group) versus 30-33 weeks' gestation (control group)

Variables	Control group (n = 513)	Study group (n = 548)	<i>p</i> value
Labor induction (n [%])	133 (25.9)	162 (29.6)	0.19
Gestational age at delivery (wk) (median $\pm$ IQR)	40 (39-40.5)	40 (39.1-40.5)	0.83
Male gender (n [%])	266 (51.9)	284 (51.8)	0.99
Birthweight (g) (median $\pm$ IQR)	3295 (3030-3615)	3282.5 (3035-3620)	0.99
Birthweight centile (median $\pm$ IQR)	39.6 (22.7-64.1)	39.2 (21.7-64.9)	0.64
Birthweight			
< 10th centile (n [%])	46 (9.0)	52 (9.5)	0.77
< 3rd centile (n [%])	9 (1.8)	8 (1.5)	0.81
Operative vaginal delivery (n [%])	135 (26.3)	168 (30.7)	0.12
Operative vaginal delivery for nonreassuring fetal status (n [%])	53 (39.3)	41 (24.4)	0.005
Cesarean delivery (n [%])	103 (20)	107 (19.5)	0.82
Cesarean delivery for nonreassuring fetal status (n [%])	40 (38.8)	18 (16.8)	<0.001

Abbreviations: wk, weeks; IQR, interquartile range; SD, standard deviation.