

Kong M. Wong,¹ Keith E. Duncan,¹ Sergey Komarov,² Dierdra Daniels,³
Daniela S. Floss,³ Yuan-Chuan Tai,² Christopher N. Topp¹

1. Donald Danforth Plant Science Center, St. Louis, MO, 63132 USA
2. Department of Radiology, Washington University in St. Louis, St. Louis, MO, 63110 USA
3. Valent BioSciences Corporation, Libertyville, IL, 60048 USA

The symbiosis between crops and arbuscular mycorrhizal fungi (AMF) have become an attractive route towards achieving carbon neutral agriculture and reducing the use of chemical fertilizers. Yet, our understanding of how active AMF infections influence the uptake, allocation, and exchange of carbon is limited. Here, we combine X-ray CT and PET imaging to observe and quantify the flow of carbon from leaves to roots to hyphae. Comparison of maize grown with and without AMF allow us to measure changes in the amount of $^{11}\text{CO}_2$ taken up in leaves and subsequently the amount of ^{11}C allocated to below-ground roots. Then, co-registered CT and PET images are used to identify hot spots which may indicate active AMF infection sites. Finally, analysis of ^{11}C kinetics at these hot spots are used to assess the amount of carbon exchanged between maize roots and hyphae. By combining structural and biochemical information, we begin to deepen our understanding of the different types of changes in carbon flow in Maize-AMF systems and how we can improve sustainable agriculture efforts.