

One-Dimensional Variational Ionospheric Retrieval Using Radio Occultation Bending Angles: Part 2 - Validation

S. Elvidge¹, S. B. Healy², and I. D. Culverwell³

¹Space Environment and Radio Engineering (SERENE), University of Birmingham, United Kingdom

²European Centre for Medium-Range Weather Forecasts, Reading, United Kingdom

³Met Office, Exeter, United Kingdom

Key Points:

- Improved computational performance of the 1D-Var bending angle retrieval is demonstrated with a better background model
- Extensive validation of the 1D-Var retrieval approach compared to ionosondes and COSMIC-2 retrievals has been undertaken
- The 1D-Var retrieval, using four-layers, is shown to have an ~40% reduction in root mean square error compared to COSMIC-2 retrievals

Corresponding author: Sean Elvidge, s.elvidge@bham.ac.uk

Abstract

Culverwell et al. (2023) described a new one-dimensional variational (1D-Var) retrieval approach for ionospheric GNSS radio occultation (GNSS-RO) measurements. The approach maps a one-dimensional ionospheric electron density profile, modeled with multiple “Vary-Chap” layers, to bending angle space. This paper improves the computational performance of the 1D-Var retrieval using an improved background model and validates the approach by comparing with the COSMIC-2 profile retrievals, based on an Abel Transform inversion, and co-located (within 200 km) ionosonde observations using all suitable data from 2020. A three or four layer Vary-Chap in the 1D-Var retrieval shows improved performance compared to COSMIC-2 retrievals in terms of percentage error for the F2 peak parameters (NmF2 and hmF2). Furthermore, skill in retrieval (compared to COSMIC-2 profiles) throughout the bottomside (~90 km to 300 km) has been demonstrated. With a single Vary-Chap layer the performance is similar, but this improves by approximately 40% when using four-layers.

Plain Language Summary

Culverwell et al. (2023) presented a new way of estimating ionospheric electron density using the amount of bending experienced by GNSS signals in the upper atmosphere. In this paper, as well as providing extensive validation of the technique, the computational performance is improved by using better initial conditions. The validation is done by comparing the newly described approach with that used by the COSMIC-2 satellite constellation using additional data from ground-based sensors known as ionosondes. The newly described technique is found to provide improvements of approximately 40% compared to a complementary approach using by COSMIC-2.

1 Introduction

Culverwell et al. (2023) described a one-dimensional variational (1D-Var) ionospheric retrieval that can be applied to the Metop-SG measurement geometry, which will truncate the ionospheric GNSS-RO measurements around 600 km above the surface. This new, general approach for the ionospheric GNSS-RO retrieval problem, is valid for both the truncated and standard GNSS-RO measurement geometry. A new 1D ionospheric bending angle forward operator was described that computes the L2-L1 bending angle differences as a function of impact parameter, assuming that the ionospheric electron density can be modeled with multiple Vary-Chap layers. This approach is close to how GNSS-RO data is used in neutral atmosphere applications, such as operational Numerical Weather Prediction (NWP) models (Healy & Thépaut, 2006).

Furthermore Culverwell et al. (2023) showed that gradient based minimisation techniques can be successfully applied to this bending angle retrieval problem, and the non-linearity of the Vary-Chap functions was demonstrated to not be problematic. The 1D-Var filters out measurement noise and the retrieved electron density solutions are smooth in comparison with the more traditional Abel transform retrieval. The use of bending angles may have implications for how GNSS-RO data is used in ionospheric data assimilation (DA) systems. Until now, slant TEC values have usually been assimilated in ionospheric DA systems, but these require a correction for the Differential Code Biases (DCB). Since the DCB is usually assumed to be constant during the occultation, it will not impact the bending angle values derived from the raw phase delays, and so it is not required to assimilate these measurements.

The purpose of this paper is to fully validate the 1D-Var ionospheric retrieval using COSMIC-2 bending angles. The 1D-Var and COSMIC-2 retrievals, based on the Abel Transform inversion as implemented by University Corporation for Atmospheric Research (UCAR), hereafter known simply as “COSMIC-2 retrievals”, are compared to co-located

Table 1. Background and analysis parameters for a one-layer Vary-Chap with different background conditions.

	Parameters				
	$NmF2$	$hmF2$	$H_{0,F2}$	k_{F2}	Number of iterations
Background	2.00×10^{12}	300	50.0	0.15	9
Analysis	5.66×10^{11}	244	50.1	0.14	
Background	7.00×10^{11}	300	50.0	0.15	11
Analysis	5.66×10^{11}	244	50.1	0.14	
Background	7.00×10^{11}	250	50.0	0.15	6
Analysis	5.66×10^{11}	244	49.9	0.14	
Background	2.00×10^{11}	300	50.0	0.15	14
Analysis	5.66×10^{11}	244	50.1	0.14	

(within 200 km) ionosonde observations using all suitable data from 2020, which involves a statistical comparison of over 10,000 ionospheric profiles.

2 COSMIC-2

The FORMOSAT-7/COSMIC-2 constellation is a constellation of six identical satellites in a low inclination orbit at a nominal altitude of 520-550 km with an inclination of 24° (Anthes & Schreiner, 2019). It follows the successful COSMIC-1 program, which launched in April 2006 (Ho et al., 2020). The COSMIC-2 constellation’s primary aim is to observe the atmosphere from a low latitude orbit using RO, supporting operational global weather prediction, tropical weather and climate research, space weather forecasting, and ionospheric research.

Whilst COSMIC-2 do not provide L1 and L2 bending angles, here we use the derivative of slantwise TEC S , with respect to the impact parameter a , $\partial S/\partial a$. Culverwell et al. (2023) showed that, to a good approximation, this is proportional to the difference between the L2 and L1 bending angles, plus an electron density term at the LEO satellite. Here we fit to bending angles with impact heights (impact parameter - radius of curvature) between 100 and 500 km. In contrast COSMIC-2 retrievals use data from 0 up to a height of ~ 700 km at a vertical resolution of 1 Hz (≈ 2 to 3 km).

3 Background Model

In this implementation of a 1D-Var system the *a priori* (background) values are effectively used as a first guess to start the minimisation process rather than a strong constraint on the final 1D-Var solution. For example, using COSMIC-1 measurements provided by the Institut D’Estudis Espacials de Catalunya (IEEC) in Spain using a one-layer Vary-Chap and a variety of background parameters gives the analysis results and the number of iterations until convergence shown in Table 1.

What is clear from Table 1 is that varying the background model parameters has almost no impact on the final analysis (the only difference at all is in the third block with an $H_{0,F2}$ value of 49.9 where in all other instances it is 50.1). The main difference between varying the parameters is the impact on the number of iterations required to arrive at the analysis results. This varies from 14 iterations to 6. This suggests that an im-

proved background guess may reduce the number of iterations required for the model to converge.

Noting this increased performance when using a more accurate background guess, rather than a fixed set of values, a simple model for these parameters is defined below. Whilst a more complicated background model could be used, this increases the computational cost and run times. Since the *a priori* values are not a strong constraint in this system, these cost increases provide no advantages.

This peak parameter (NmF2, NmF1, NmE, hmF2, hmF1 and hmE) model is based on Nava et al. (2008); Angling et al. (2018).

3.1 NmE

The peak density of the ionospheric E-region, NmE , is well modelled by a simple function based on the a seasonal relationship with F10.7 and solar zenith angle χ (Leitinger & Kirchengast, 1997; Nava et al., 2008):

$$NmE = \frac{a_e}{80.616} \sqrt{F_{10.7}} \cos^{0.6}(\chi_{eff}), \quad (1)$$

where a_e is a seasonal term given by (Nava et al., 2008)

$$a_e = (1.112 - 0.019s_p)^2, \quad (2)$$

$$s_p = s \cdot \tanh(0.15\phi), \quad (3)$$

where ϕ is latitude in radians, $s = -1$ if the month is January, February, November or December, $s = 0$ if March, April, September or October and $s = 1$ if May, June, July or August. Finally, χ_{eff} is the effective solar zenith angle given by

$$\chi_{eff} = \frac{\chi + [90 - 0.24 \exp(20 - 0.2\chi)] \cdot \exp(12(\chi - \chi_0))}{1 + \exp(12(\chi - \chi_0))}, \quad (4)$$

where χ_0 is the zenith angle at night-day transition, 86.23° as given in ITU-R P.2297-1 (2019).

3.2 hmE

Across a range of models including NeQuick and the IRI hmE , the height of the E-region peak density, is usually set to a fixed height. In this model

$$hmE = 110 \text{ km}, \quad (5)$$

as per the updated NeQuick model from Angling et al. (2018).

3.3 NmF2

NmF2, the peak density of the F2-layer, is dependent on a number of factors including geographical longitude, latitude, time, season/day of year and solar activity. A number of approaches can be used to described solar activity, for this simple model the sunspot number (SSN) is used. The NeQuick (Nava et al., 2008) and the Ionospheric Reference Model (IRI; e.g. (Bilitza & Reinisch, 2008)) use the Committee Consultative for Ionospheric Radiowave propagation (CCIR) files to compute NmF2 and M(3000)F2 (the ratio of the maximum usable frequency at a distance of 3000 km to the F2 layer critical frequency, foF2). The CCIR maps for the F2 parameter consist of 988 coefficients for each month. CCIR provides two sets of coefficients, one for low sunspot numbers and one for high (Haralambous et al., 2021). The coefficients for intermediate levels of solar activity are determined by linear interpolation (European Commission, 2016).

As per (Nava et al. (2008); European Commission (2016) Equations 61–68 and 77) the CCIR coefficients are interpolated for the current date and solar activity levels and then vectors of coefficients for Legendre polynomials are calculated. The NmF2 (electrons/m³) is specified in terms of foF2 (Hz):

$$NmF2 = 0.0124foF2^2, \quad (6)$$

where foF2 itself is defined as the sum of nine intermediate terms:

$$foF2 = \sum_{n=1}^9 foF2_n, \quad (7)$$

where

$$foF2_1 = \sum_{k=1}^{12} f_k m_k, \quad (8)$$

and

$$foF2_n = \cos^{n-1}(\text{lat}) \sum_{k=1}^{q_n} [f_{k_n+2k-1} \cos((n-1)\text{lon}) + f_{k_n+2k} \sin((n-1)\text{lon})] m_k \quad (9)$$

for $n = 2, \dots, 9$ where

$$q_n = \{12, 12, 9, 5, 2, 1, 1, 1, 1\}, \quad n = 1, \dots, 9 \quad (10)$$

$$k_n = \{-12, 12, 36, 54, 64, 68, 70, 72, 74\}, \quad n = 1, \dots, 9. \quad (11)$$

m_k are the “modip” coefficients (fully described in European Commission (2016) equations 6-16), lon and lat are the longitude and latitude of interest and the f_j are the 76 CCIR coefficients for the Legendre calculation (European Commission, 2016; Equations 40-50; note that f is written as $cf2$ in those equations).

3.4 hmF2

The calculation of the height of the maximum F2 density, hmF2, is based on the NmF2, NmE and the M(3000)F2 (European Commission, 2016; Equations 69–76). The M(3000)F2 is calculated similarly to NmF2 from Section 3.3 as the sum of seven intermediate terms:

$$M(3000)F2 = \sum_{n=1}^7 M(3000)F2_n, \quad (12)$$

where

$$M(3000)F2_1 = \sum_{k=1}^7 u_k m_k, \quad (13)$$

and

$$M(3000)F2_n = \cos^{n-1}(\text{lat}) \sum_{k=1}^{r_n} [u_{h_n+2k-1} \cos((n-1)\text{lon}) + u_{h_n+2k} \sin((n-1)\text{lon})] m_k \quad (14)$$

where

$$r_n = \{8, 6, 3, 2, 1, 1\} \quad \text{and} \quad (15)$$

$$h_n = \{7, 23, 35, 41, 45, 47\}, \quad (16)$$

for $n = 2, \dots, 7$. u_j are the interpolated CCIR coefficients (European Commission, 2016; Equations 40–51; note that u is written as $cm3$ in those equations).

Then the $hmF2$ is defined as (Nava et al., 2008)

$$hmF2 = \frac{1490M}{M + \Delta M} \sqrt{\frac{0.0196M^2 + 1}{1.2967M^2 - 1}} - 176 \quad (17)$$

where $M = M(3000)F2$ and

$$\Delta M = \frac{0.253}{\rho - 1.215} - 0.012, \quad (18)$$

$$\rho = N \cdot \frac{\exp(20(N - 1.75)) + 1.75}{\exp(20(N - 1.75)) + 1}, \quad (19)$$

$$N = \sqrt{\frac{NmF2}{NmE}}. \quad (20)$$

3.5 NmF1

The maximum density of the F1-layer, $NmF1$, is primarily defined in this model as in NeQuick (Nava et al., 2008), in terms of the critical density of the E-Region. Specifically

$$NmF1 = 1.96NmE. \quad (21)$$

3.6 hmF1

The height of the peak density of the F1-layer, $hmF1$, is defined here simply as the average of the height of peak density of the E and F2 layer:

$$hmF1 = \frac{hmF2 + hmE}{2} = \frac{hmF2 + 110}{2}. \quad (22)$$

3.7 Using the New Background Model

Table 1 demonstrated the impact of the background model values on the number of iterations needed to converge to a solution with the 1D-Var. Rather than using a fixed set of values for the initial guesses for the Vary-Chap layers but instead using values from the model described in the previous sections has a major impact on the iteration rate and the total number of successful convergences (a successful convergence is defined as a convergence which takes less than 50 iterations), Table 2.

The four-layer Vary-Chap model in Table 2 is an E-, F1-, F2- and topside-layer model. The D-region is excluded as it has very little impact on the overall results (see Culverwell et al. (2023)). It can be seen from Table 2 that using the new background model not only reduces the average number of iterations required for convergence in each case but also improves the overall percentage of observations which do converge. The average reduction in the number of observations is 5 iterations and the increased percentage of observation convergence means that all test versions of the model have at least a 73.4% convergence rate (improved from 58.7%).

4 Statistical Performance of the 1D-Var Retrieval

To undertake a rigorous statistical analysis of the 1D-Var retrieval technique two things are needed:

Table 2. Percentage of observations which converge within 50 iterations and statistics of the rate of convergence for up to four Vary-Chap layers using the first-guess background conditions described in Table 1 of Culverwell et al. (2023) and the background model described here.

	Background using old parameter conditions		Background using new parameter model	
	Observations which converged	Mean Iterations	Observations which converged	Mean Iterations
1 layer	98.6%	16	99.3%	11
2 layers	85.5%	34	92.7%	26
3 layers	66.7%	28	80.2%	24
4 layers	58.7%	31	73.4%	28

1. a comparison to independent observations,
2. sufficient retrievals to reduce uncertainty in the analysis.

4.1 Radio Occultation Observations

The current most abundant, and freely available, source of recent RO observations is from the COSMIC-2 satellites (see Section 2). To perform a rigorous statistical analysis of the 1D-Var retrieval technique all available Level 1B (TEC observations) and Level 2 (ionospheric profiles) products from 2020 have been used (UCAR, 2019). COSMIC-2 ionospheric retrievals use the calibrated TEC data derived from the L1 and L2 phase differences, which is then calibrated using the method described in CDAAC (Strauss et al., 2020).

In total there are 1,838,920 Level 1B occultations (each containing several hundred TEC observations) and an associated 966,358 Level 2 derived ionospheric profiles available to download in 2020.

4.2 Independent Observations

By stepping through a range of HF frequencies transmitted vertically upwards and measuring the return echoes ionosondes can image the vertical profile of the ionosphere up to the peak density. These observations are both widely assimilated by ionospheric models and also commonly used as reference observations (often incorrectly called ‘truth’) for comparative studies e.g. (Feltens et al., 2011; Elvidge et al., 2014; Scherliess et al., 2011).

Lin et al. (2020) and Cherniak et al. (2021) have previously validated the COSMIC-2 profiles, in terms of peak density/height, by comparison to ionosonde profiles at eight locations across one month in late 2019 and two months in early 2020 respectively. The Lin et al. (2020) study resulted in the comparison of 135 RO profiles and the Cherniak et al. (2021) study used ~ 2200 profiles.

In this work every ionosonde observation (profiles) from within 200 km of the location of an occultation in 2020 has been used to validate both the COSMIC-2 profile reconstruction as well as the 1D-Var retrieval. Usually ionosonde profiles are “autoscaled” to get the true height information from the observations, but this can, and does, give rise to “autoscaling errors”. Ionogram autoscaling is the process of automatically detecting the traces on a graph of time-of-flight against transmitted frequency which can then be used to infer the electron densities. The most commonly used autoscaling software is the

Table 3. Number of COSMIC-2 profiles analysed

Total occultations	1,838,920	100.0%
Number which have associated electron density profiles	966,358	52.5%
Number which have nearby ionosonde observations	10,935	0.59%
Number which do not contain obvious autoscaling errors	10,612	0.58%

Automatic Real-Time Ionogram Scaler with True height (ARTIST) (Galkin et al., 2008) which uses a hyperbolic trace fitting method. The best way to overcome these autoscaling errors is to manually scale the ionograms (as was done in Lin et al. (2020)), however that is a time consuming process. Another way to address the problem is by looking at the confidence scores that autoscaling software, such as ARITST, uses to assess the success of the inversion. Only profiles with maximum confidence (100) have been used. That results in 10,935 profiles used in this study.

However, even profiles with the maximum confidence score can still contain errors (Themens et al., 2022). To address this issue, each ionosonde profile has been examined and removed from the analysis if they contain obvious errors. Overall this results in 10,612 profiles which are used in the analysis, far more than used in previous studies. The number of observations are summarized in Table 3.

4.3 Analysis Results

Initial analysis of the 1D-Var retrieval, in a similar approach to (Lin et al., 2020), is to look at the performance of the F2 peak in terms of density and altitude compared to the ionosondes. These results can then be compared to the COSMIC-2 electron density profiles. However, before looking at the statistical performance of the two approaches, some quality control (QC) of the peak parameters is required.

An easy first approximation to find the F2 peak parameters is to take the maximum density of the electron density profile and associated altitude. Figures 1 and 2 show the resulting probability density plots for the maximum density and the associated altitudes respectively. By comparing the COSMIC-2 height of maximums to the probability distribution function of hmF2's using data from the Chilton, UK ionosonde between 2000 and 2019 (Figure 2) it is clear that the main bulk of the distributions agree closely. However the second and third peaks centred at 100 km and just greater than 0 in the COSMIC-2 data are likely not actually hmF2 values. These likely included both observations of sporadic-E and other errors in the COSMIC-2 profiles (e.g. see examples in Figure 3). In this analysis hmF2 values are defined to be between 200 km and 500 km; if a value is outside of this range it is excluded from the analysis.

4.4 Statistical Performance

To provide a statistical comparison between the F2 peak parameters the relative percentage error ($\Delta X(\%) = 100(X - X_{ionosonde})/X_{ionosonde}$) of the specification of both NmF2 and hmF2 between these ionosonde observations with both the 1D-Var retrieval with up to four-layers and COSMIC-2 retrieval is shown in Table 4.

From these results it is clear that the 1D-Var retrieval outperforms COSMIC-2 when using three or four layers. There is a 7% relative error in the COSMIC-2 retrieval of NmF2 (which is consistent with the results of Cherniak et al. (2021)) compared with -5.3% and -4.2% when using 1D-Var with three and four layers respectively. However, when only using one-layer the 1D-Var has a -14.8% error in the retrieval of the parameter. For hmF2 the COSMIC-2 error is 2.6% and all versions of the 1D-Var retrieval result in a smaller




Figure 1. Histogram of COSMIC-2 maximum densities.




Figure 2. Probability distribution function of COSMIC-2 maximum densities overlaid with hmF2 data from Chilton, UK ionosonde using data from 2000 to 2019.

Table 4. Statistical performance of retrieving NmF2 and hmF2 from the 1D-Var compared with COSMIC-2 relative to ionosondes

COSMIC-2		1D-Var			
		1 Layer	2 Layers	3 Layers	4 Layers
NmF2	7.0%	-14.8%	-10.6%	-5.3%	-4.2%
hmF2	2.6%	1.3%	0.7%	-0.1%	-0.1%




Figure 3. Sample of COSMIC-2 electron density profiles, showing occasional sporadic E-layers and spurious results, COSMIC-2 filenames are above each subplot.

error decreasing from 1.3% to just -0.1%. However it should be noted that errors in hmF2 specification from autoscaled ionosonde observations can be large ($\sim \pm 10\%$) and this should be kept in mind when comparing the parameters.

A more detailed view of how the number of layers improves the F2 specification can be seen by looking at probability density functions of the error in NmF2 specification. Figure 4a shows the probability density function for COSMIC-2 and a single F2 layer, overall the distributions are similar, albeit with a slight negative bias in the 1D-Var retrieval. Moving around in Figure 4 the 1D-Var retrieval has more layers added to the solution, from one to four. With increasing layers the bias and standard deviation is reduced, with overall the four layers in Figure 4d performing the best, with a significantly reduced standard deviation compared to the COSMIC-2 retrieval.

4.5 Bottomside Profile

Whilst the F2-peak is an important parameter and a good indicator of how well the 1D-Var retrieval is working, one reason for using the Vary-Chap layers is to reconstruct the full ionospheric profile. To assess the bottomside (the altitudes region below that of the peak ionospheric density) performance all of the ionospheric profiles (from the ionosonde, COSMIC-2 and the 1D-Var) are interpolated onto the same altitude grid, at 1 km resolution, and the root mean square error (RMSE) with respect to the ionosonde at each altitude is then calculated. The resulting RMSE altitude profile is shown in Figure 5.

This shows that the 1D-Var retrieval with one-layer (F2) performs very similarly to the COSMIC-2 profiles, and above 225 km there is no statistically significant differences in the results, however below 200 km the COSMIC-2 retrieval is clearly superior. A two-layer model (F2+F1) shows an improvement over COSMIC-2 above 200 km, which is consistent with the fact of adding a lower-altitude layer. The four-layer (F2+F1+E+Topside) 1D-Var retrieval shows an excellent performance throughout the altitude range, with an average improvement over COSMIC-2 by approximately 40%. Only above ~ 280 km do the COSMIC-2 results show comparable/improved performance relative to the four-layer




Figure 4. Panels show comparisons of probability density functions of the error in NmF2 from COSMIC-2 compared to the 1D-Var solution with (a) an F2 Layer, (b) F2+F1 layers, (c) F2+F1+E layers and (d) F2+F1+E+Topside layers. Vertical dashed line is marked at 0.




Figure 5. Root mean square error (RMSE) altitude profiles for COSMIC-2 (red) and the 1D-Var retrieval with only 1 Layer (F2; blue), 2 Layers (F2+F1; orange) and 4 Layers (F2+F1+E+Topside; green) compared to ionosonde observations.

model, although the number of data points at these altitudes is relatively small and the results are not statistically distinguishable.

5 Discussion and Conclusions

The statistical performance of the 1D-Var retrieval technique has been thoroughly evaluated in the current work, comparing it to independent observations and leveraging a substantial set of retrievals to reduce uncertainty in the analysis. The method was applied to the Level 1B and Level 2 data obtained from COSMIC-2 satellites and validated using ionosonde observations, constituting a rigorous and expansive analytical process.

From the analysis, it was clear that the 1D-Var retrieval technique demonstrates a robust performance, outperforming COSMIC-2 retrievals, particularly when employing three or four layers. The relative percentage errors in the specification of both NmF2 and hmF2 were found to be lower for the 1D-Var retrieval. This demonstrates the significant potential of the 1D-Var technique in the field of ionospheric profile reconstruction. However, it's worth noting that the 1D-Var retrieval's performance was significantly lower when only one layer was used, indicating the importance of multi-layer modeling for improved accuracy. This is an important finding and suggests that future work should focus on refining and utilizing multi-layer models to improve the retrieval of ionospheric parameters.

Furthermore, the comprehensive analysis of the full ionospheric profiles showed that the four-layer 1D-Var retrieval exhibited excellent performance throughout the altitude range. Notably, there was an average improvement over COSMIC-2 by approximately

40%, showcasing the promising potential of this technique. Although the COSMIC-2 results showed comparable performance to the four-layer model above 280 km, the number of data points at these altitudes was relatively small, and the results weren't statistically distinguishable. This suggests that further research should be dedicated to improving the modeling and retrieval at these higher altitudes.

In conclusion, the 1D-Var retrieval technique, particularly when using three or four layers, offers a significant advancement in ionospheric profile reconstruction. While there are still areas for improvement, particularly in the retrieval at higher altitudes, the findings of this work provide a strong foundation for further research in this field.

6 Open Research

COSMIC-1 data is available from <https://data.cosmic.ucar.edu/gnss-ro/cosmic1/repro2013/>. The processed files from the IEEC may be requested from Dr Hernández-Pajares. COSMIC-2 data is available from <https://data.cosmic.ucar.edu/gnss-ro/cosmic2/nrt/> and the ionosonde observations were retrieved from The National Centers for Environmental Information (NCEI) <https://www.ngdc.noaa.gov/stp/iono/ionogram.html>.

The MODIP and CCIR files required to reconstruct the *a priori* model described in this work are available from the International Telecommunications Union (<https://www.itu.int/md/R07-WP3L-C-0094/en>) or by contacting Bruno Nava (<https://t-ict4d.ictp.it/nequick2/source-code>).

The background model described in this paper is coded in Python 3.x, as is the 1DVar retrieval code that uses it. An officially supported Fortran95 implementation of the latter has been part of the Radio Occultation Processing Package (ROPP) since version 11.0 (released January 2022). ROPP is maintained, developed and supported by EUMETSAT, through the Radio Occultation Meteorology Satellite Applications Facility (ROM SAF), and freely available to download from its website (ROM SAF, 2023).

Acknowledgments

We thank Dr Haixia Lyu and Dr Hernández-Pajares for providing the COSMIC-1 data used here. Dr Riccardo Notarpietro and Dr David Themens are thanked for useful discussions during the early stages of this work.

This work was conducted as part of the Visiting Scientist program of the Radio Occultation Meteorology Satellite Applications Facility (ROM SAF) which is a decentralized operational radio occultation processing center under EUMETSAT. S. Elvidge was a ROM SAF Visiting Scientist for this project, and S. Healy and I. Culverwell are respectively current and former members of the ROM SAF.

References

- Angling, M. J., Elvidge, S., & Healy, S. B. (2018, April). Improved model for correcting the ionospheric impact on bending angle in radio occultation measurements. *Atmospheric measurement techniques*, 11(4), 2213–2224. Retrieved 2019-02-21, from <https://www.atmos-meas-tech.net/11/2213/2018/> doi: 10.5194/amt-11-2213-2018
- Anthes, R., & Schreiner, W. (2019). Six new satellites watch the atmosphere over earth's equator. *Eos*, 100. doi: <https://doi.org/10.1029/2019EO131779>
- Bilitza, D., & Reinisch, B. (2008, August). International Reference Ionosphere 2007: Improvements and new parameters. *Advances in Space Research*, 42(4), 599–609. Retrieved 2019-02-21, from <https://linkinghub.elsevier.com/retrieve/pii/S0273117708000288> doi: 10.1016/j.asr.2007.07.048

- Cherniak, I., Zakharenkova, I., Braun, J., Wu, Q., Pedatella, N., Schreiner, W., ...
Hunt, D. (2021). Accuracy assessment of the quiet-time ionospheric F2 peak
parameters as derived from COSMIC-2 multi-GNSS radio occultation mea-
surements. *Journal of Space Weather and Space Climate*, 11, 18. Retrieved
2021-09-11, from [https://www.swsc-journal.org/articles/swsc/abs/](https://www.swsc-journal.org/articles/swsc/abs/2021/01/swsc200070/swsc200070.html)
2021/01/swsc200070/swsc200070.html (Publisher: EDP Sciences) doi:
10.1051/swsc/2020080
- Culverwell, I. D., Healy, S. B., & Elvidge, S. (2023). One-dimensional variational
ionospheric retrieval using radio occultation bending angles: Part 1 - theory.
Journal of Geophysical Research: Space Physics, X(Y).
- Elvidge, S., Angling, M. J., & Nava, B. (2014, September). On the use of modified
Taylor diagrams to compare ionospheric assimilation models. *Radio Science*,
49(9), 737–745. Retrieved 2019-02-21, from [http://doi.wiley.com/10.1002/](http://doi.wiley.com/10.1002/2014RS005435)
2014RS005435 doi: 10.1002/2014RS005435
- European Commission. (2016). *Ionospheric correction algorithm for galileo single
frequency users* (Tech. Rep.). Retrieved from [https://www.gsc-europa.eu/](https://www.gsc-europa.eu/sites/default/files/sites/all/files/Galileo_Ionospheric_Model.pdf)
sites/default/files/sites/all/files/Galileo_Ionospheric_Model.pdf
- Feltens, J., Angling, M. J., Jackson-Booth, N., Jakowski, N., Hoque, M., Hernández-
Pajares, M., ... Zandbergen, R. (2011, December). Comparative testing of
four ionospheric models driven with GPS measurements. *Radio Science*, 46(6).
Retrieved 2019-02-21, from <http://doi.wiley.com/10.1029/2010RS004584>
doi: 10.1029/2010RS004584
- Galkin, I., Khmyrov, G., Kozlov, A., Reinisch, B. W., Huang, X., & Paznukhov, V.
(2008). The ARTIST 5. *Radio Sounding and Plasma Physics, AIP Proceedings*
#974.
- Haralambous, H., Leontiou, T., Petrou, V., Kumar Singh, A., Charalambides, M.,
Lithoxopoulos, N., & Agisilaou, A. (2021). Adjusting ccir maps to improve
local behaviour of ionospheric models. *Atmosphere*, 12(6). Retrieved from
<https://www.mdpi.com/2073-4433/12/6/691> doi: 10.3390/atmos12060691
- Healy, S. B., & Thépaut, J.-N. (2006). Assimilation experiments with champ gps
radio occultation measurements. *Quarterly Journal of the Royal Meteorological
Society*, 132. doi: <https://doi.org/10.1256/qj.04.182>
- Ho, S.-p., Anthes, R. A., Ao, C. O., Healy, S., Horanyi, A., Hunt, D., ... Zeng, Z.
(2020, July). The COSMIC/FORMOSAT-3 Radio Occultation Mission after
12 Years: Accomplishments, Remaining Challenges, and Potential Impacts of
COSMIC-2. *Bulletin of the American Meteorological Society*, 101(7), E1107–
E1136. Retrieved 2023-04-28, from [https://journals.ametsoc.org/view/](https://journals.ametsoc.org/view/journals/bams/101/7/bamsD180290.xml)
journals/bams/101/7/bamsD180290.xml (Publisher: American Meteoro-
logical Society Section: Bulletin of the American Meteorological Society) doi:
10.1175/BAMS-D-18-0290.1
- ITU-R P.2297-1. (2019). *Electron density models and data for transionospheric radio
propagation* (Tech. Rep.). Retrieved from [https://www.itu.int/dms_pub/itu](https://www.itu.int/dms_pub/itu-r/opb/rep/R-REP-P.2297-1-2019-PDF-E.pdf)
-r/opb/rep/R-REP-P.2297-1-2019-PDF-E.pdf
- Leitinger, R., & Kirchengast, G. (1997). Easy to Use Global and Regional Iono-
spheric Models. A Report on Approaches Used in Graz. *Acta Geod. Geoph.*
Hung, 32, 329–342. doi: <https://doi.org/10.1007/BF03325504>
- Lin, C.-Y., Lin, C. C.-H., Liu, J.-Y., Rajesh, P. K., Matsuo, T., Chou, M.-Y.,
... Yeh, W.-H. (2020, October). The Early Results and Validation of
FORMOSAT-7/COSMIC-2 Space Weather Products: Global Ionospheric Spec-
ification and Ne-Aided Abel Electron Density Profile. *Journal of Geophysical
Research: Space Physics*, 125(10), e2020JA028028. Retrieved 2021-08-01, from
<http://agupubs.onlinelibrary.wiley.com/doi/10.1029/2020JA028028>
(Publisher: John Wiley & Sons, Ltd) doi: 10.1029/2020JA028028
- Nava, B., Coisson, P., & Radicella, S. (2008). A new version of the neQuick iono-
sphere electron density model. *Journal of Atmospheric and Solar Terrestrial*

- 422 *Physics*, 70(15), 1856–1862.
- 423 ROM SAF. (2023). *ROPP Software Deliverable*. Retrieved from [https://www](https://www.romsaf.org/ropp/)
- 424 [.romsaf.org/ropp/](https://www.romsaf.org/ropp/)
- 425 Scherliess, L., Thompson, D. C., & Schunk, R. W. (2011). Data assimilation models:
426 A new tool for ionospheric science and applications. In W. Liu & M. Fujimoto
427 (Eds.), *The Dynamics Magnetosphere* (pp. 329–339). Springer, Berlin.
- 428 Strauss, P., Schreiner, W., & Santiago, J. (2020). *FORMOSAT-7/COSMIC-2*
429 *TGRS Space Weather Provisional Data Release 1* (Tech. Rep.). Retrieved
430 from [https://data.cosmic.ucar.edu/gnss-ro/cosmic2/provisional/](https://data.cosmic.ucar.edu/gnss-ro/cosmic2/provisional/spaceWeather/F7C2_SpWx_Provisional_Data_Release_1.pdf)
431 [spaceWeather/F7C2_SpWx_Provisional_Data_Release_1.pdf](https://data.cosmic.ucar.edu/gnss-ro/cosmic2/provisional/spaceWeather/F7C2_SpWx_Provisional_Data_Release_1.pdf)
- 432 Themens, D. R., Reid, B., & Elvidge, S. (2022). Artist ionogram autoscaling confi-
433 dence scores: Best practices. *Radio Science Letters - Under Review*.
- 434 UCAR. (2019). *COSMIC-2 Data Products* (Tech. Rep.). Retrieved from
435 <https://data.ucar.edu/en/dataset/cosmic-2-data-products> doi:
436 10.5065/t353-c093