

The SET HASDM density database

W. Kent Tobiska¹, Bruce R. Bowman¹, Dave Bouwer¹, Alfredo Cruz^{1,3}, Kaiya Wahl¹

Marcin Pilinski²

Piyush M. Mehta³ and Richard J. Licata³

¹Space Environment Technologies, Pacific Palisades, CA

²Freestream, Boulder, CO

³West Virginia University, Morgantown, WV

Abstract

The SET HASDM density database is available for scientific studies through a SQL database with open community access. The information in the SET HASDM density database covers the period from January 1, 2000 through December 31, 2019. Data records exist every 3 hours during solar cycles 23 and 24. The database has a grid size of $10^\circ \times 15^\circ$ (latitude, longitude) with 25 km altitude steps between 175–825 km. A description of the source of the database, its validation, its information content, and its accessibility are provided.

Current state of art for operational thermospheric densities

The United States Air Force (now United States Space Force, USSF) Combined Space Operations Center (CSpOC) creates arguably the world's most accurate current and forecast neutral thermosphere densities as part of their operation to produce the NORAD satellite catalog. To do this they use the High Accuracy Satellite Drag Model (HASDM) v2 (Storz *et al.*, 2005) as a data assimilative system working with its associated JB2008 model (Bowman *et al.*, 2008; Tobiska *et al.*, 2008; ISO 14222, 2013). HASDM continuously derives densities from several dozens of calibration satellites and revises the current epoch JB2008 densities using temperature correction coefficients to achieve very low global and regional density uncertainty at most epochs.

The JB2008 thermospheric density model (Bowman *et al.*, 2008) was developed by the U.S. Air Force Space Command and Space Environment Technologies (SET) to improve current specification and forecast of thermospheric mass densities. Its creation led to the largest single reduction in thermospheric mass density uncertainties since the early Jacchia-type models in the 1960's (Marcos *et al.*, 2006) where the 1-sigma uncertainties at 400 km for a given epoch dropped from 15% to 8% compared with previous Jacchia and MSIS models. The first improvement was to create new solar and geomagnetic indices that directly mapped solar irradiances from the Sun to the Earth's upper atmosphere. In addition to the legacy F10 solar proxy and the ap geomagnetic index, three new operational solar indices (S10, M10, and Y10) were developed for JB2008 (Tobiska *et al.*, 2008; ISO 14222, 2013) and the Dst geomagnetic index was incorporated as well.

The new solar indices represent the combined energy input from multiple solar sources: a) S10 is the 26–34 nm bandpass EUV chromospheric irradiance index; b) M10 is the 160 nm FUV Schumann-Runge photospheric irradiance proxy; and c) Y10 is the 121.6 nm Lyman-alpha chromospheric/transition region irradiance convolved with the 0.1–0.8 nm X-ray coronal irradiance to create a new solar index. S10 represents the energy for O photoabsorption in the middle and upper thermosphere (>180 km), M10 represents the energy for O₂ dissociation in the lower thermosphere (~110 km), and Y10 represents the energy for mesosphere and lower thermosphere H₂O chemistry (~90 km). The F10 was retained for legacy use to account for unmodeled solar flux.

Geomagnetic storm and sub-storm representation was accomplished using the 1-h Dst index when it was less than -75 nT. Non-storm, but unsettled, geomagnetic conditions continue to be modeled with the 3-h ap geomagnetic index, including during high speed stream events. Forecast ap is operationally provided by NOAA SWPC and forecast Dst plus the solar indices are operationally provided since 2012 by SET at <https://sol.spacenvironment.net/JB2008/>. Licata *et al.* (2020) have benchmarked the uncertainties in SET's solar and geomagnetic predictions.

JB2008 was adopted as part of the COSPAR International Reference Atmosphere (CIRA, 2014) and the ISO International Standard 14222 on Earth's upper atmosphere (ISO 14222, 2013), both of which recommend using JB2008 for calculating mass densities related to satellite drag. The densities are used in satellite orbit determination (OD) algorithms for Low Earth Orbit (LEO) to provide a solution for atmospheric drag.

The current epoch runs of the HASDM system during the course of an operational day at CSpOC are archived and contain the temperature-corrected coefficients that have been applied to the JB2008 atmosphere. The temperature corrected coefficients created at CSpOC are not released to the public. However, SET does validations of the HASDM output on a weekly basis to ensure the quality of the output product and is able to recreate the densities of the global atmosphere. This recreation is called the SET HASDM density database.

HASDM validation

Satellite drag-derived atmospheric density variations depend upon the drag coefficient, C_D , times the satellite frontal area, A , divided by mass, m . The inverse of this expression is also known as the ballistic coefficient, b . C_D is a function of satellite shape, thermospheric temperature, and thermospheric composition. Atmospheric winds cause variable velocity, v , affecting satellite drag variations at high latitudes during large geomagnetic storms. Legacy methods obtained density variations from analysis of satellite orbits using mean orbital elements or raw observational tracking data. Current methods analyze orbit decay rates using special orbit perturbations to fit satellite observations; however, those require very accurate radar availability and optical observations. The HASDM validation method uses densities derived from the use of special perturbations (SP) OD employing daily orbital energy dissipation rates (EDRs) to obtain very accurate daily variations. Radar and optical observations from the worldwide USAF Space Surveillance Network (SSN) are used to fit precise orbital elements. A key element in this method is obtaining calibrated ballistic coefficients to compute the daily-observed density variations from the drag-derived computation (DDC) method using observed daily EDRs and differential orbit correction (DOC). Once these densities are determined for the calibration satellites, the least squares fitting assimilation of these calibration satellites' data into the JB2008 current epoch density solutions is then able to drop the HASDM density uncertainties to generally less than 5% for a given epoch.

Validation of the overall HASDM system is done weekly by SET. With the Drag-Derived Computation (DDC) method Bowman *et al.* (2004) uses government data sets and computes accurate daily thermospheric density values down to 150 km derived from SSN data. A SP differential orbit correction (DOC) program fits SSN radar and optical observational data to obtain a standard 6-element state vector plus a ballistic coefficient, b . In this algorithm, the atmospheric density model used is the Modified Jacchia 1970 (J70Mod) developed originally for HASDM v1. EDRs are modeled over the observation span of 3–8 days using the J70Mod density values and the fitted b -value. Overlapping EDR values are used to compute accurate daily values. Daily temperature and density values are then calculated from the EDR using the average 30-year b -value of each satellite. The HASDM density model computes partial derivatives of temperature and EDR changes. The method was validated through comparison of daily temperature results during 2001 with the HASDM Dynamic Calibration Atmosphere (DCA) project. The compari-

sons were excellent and the computed daily densities were validated against historical daily densities for the last 30 years using many satellites. The density accuracy was quantified by comparing geographically overlapping perigee location data with over 8500 pairs of density values used for the comparisons. The density errors are <4% overall and ~2% during the latest solar maximum.

The Differential Orbit Correction (DOC) method is used to validate the HASDM orbital fit for individual orbiting objects and uses the SSN observations for the six Keplerian elements plus b . A minimum of 10 accurate radar observations spread throughout the fit span ensures accuracy; the majority use at least 30–60 observations per fit. The DOC orbit fits for an orbiting object are obtained using a weighted least squares DOC program with SP orbit integration. The geopotential used in the DOC is the EGM96 model truncated to a 48×48 field. The use of the 48×48 field results in maximum b errors <4% with a standard deviation <1% for an orbit with a perigee height close to 400 km. These are acceptable to obtain the desired density accuracy. The SP integration includes third-body gravitational effects of the Sun and Moon, solar radiation pressure, and atmospheric drag accelerations. A variable C_D is required for the processing.

A final step in the HASDM validation uses the Energy Dissipation Rate (EDR) to compute accurate daily density values for an orbiting object. DOC was modified to compute orbital EDR over an observation span using J70Mod density and fitted b values (Bowman and Hrnir, 2007). Overlapping EDR values are obtained by moving the observation span by one day and then recomputing another orbit fit with the same span length. Overlapping fits are then used to compute accurate daily EDR values using a weighting method favoring EDR values obtained from the middle of the observation span as opposed to those at the ends of the span. Finally, daily EDR values are used to compute observed daily temperature values. The daily EDR values are obtained from averaging the same day's overlapping values obtained from the orbit fits. Because of fit error, the best method for obtaining an accurate daily EDR is to constrain the average daily EDR to maintain conservation of energy over each orbit fit span. Thus, energy change over each fit must be preserved since these were the real measured quantities when the orbit fits were obtained.

SET HASDM density database extraction and availability

Under authority from USSF, SET has extracted two solar cycles of temperature-corrected coefficients and created the SET HASDM density database. This is the first-time there has been an extraction of this operational database for scientific use; there is no comparable dataset in existence. All solar cycle, geomagnetic storm and sub-storm, extended solar flare, and thermospheric cooling perturbations are embedded in the data. Because of its accuracy, time resolution, global scale, and information content, the SET HASDM database densities are suitable for use as a new space weather benchmark for atmospheric expansion against which space weather events are measured. A new benchmark refines the Phase 1 Benchmark that was released by the National Science and Technology Council (National Space Weather Strategy and Action Plan, 2019) for upper atmospheric expansion.

The statistical metrics in the SET HASDM density database are being characterized by the META-HASDM project which uses machine learning to quantify the database. That work is being developed separately for publication. Called HASDM ML, it will characterize the global density environment for rapid operational use, for quantification of uncertainty, and for improving ballistic coefficients for all LEO objects.

The SET HASDM density database now resides in a SQL database with open community access for scientific studies. The information in the SQL-based SET HASDM density database, with an example in Figure 1 for the Halloween 2003 storm period, covers the period from January 1,

2000 through December 31, 2019. Data records exist every 3 hours during solar cycles 23 and 24. The database has a grid size of $10^\circ \times 15^\circ$ (latitude, longitude) with 25 km altitude steps between 175–825 km. The database access, along with supplementary information, can be found at <https://spacewx.com/hasdm/>.

Acknowledgements

The SET HASDM density database was developed with internal R&D funding at Space Environment Technologies. The *Machine Learning Enabled Thermosphere Advanced by HASDM (META-HASDM)* project gratefully acknowledges the support of the NASA SBIR contract 80NSSC20C0292 for funding to translate the SET HASDM density database into a neural net set of coefficients.

References

- Bowman, B. R., F. A. Marcos, and M. J. Kendra (2004), A Method For Computing Accurate Daily Atmospheric Density Values From Satellite Drag Data, 14th AAS/AIAA Space Flight Mechanics Conference, AAS 04-173, February 8-12.
- Bowman, B. R. and S. Hrncir (2007), Drag Coefficient Variability at 100-300 km from the Orbit Decay Analysis of Rocket Bodies, AIAA 2007-262, AIAA Astroynamics Specialist Conference, Mackinaw Island, MI, August.
- Bowman, B., W. K. Tobiska, F. Marcos, C. Huang, C. Lin, and W. Burke (2008), A new empirical thermospheric density model JB2008 using new solar and geomagnetic indices, In *AIAA/AAS Astro-dynamics specialist conference and exhibit* (p. 6438).
- ISO 14222 (2013), Space environment (natural and artificial) - Earth upper atmosphere, Standard, International Organization for Standardization, Geneva, CH.
- Licata, R. J., W. K. Tobiska, and P. M. Mehta (2020), Benchmarking forecasting models for space weather drivers. *Space Weather*, 18, e2020SW002496. <https://doi.org/10.1029/2020SW002496>.
- Marcos, F. A., B. R. Bowman, and R. E. Sheehan (2006), Accuracy of Earth's thermospheric neutral density models, AIAA 2006-6167, AIAA/AAS Astroynamics Specialist Conference, Keystone, CO, August.
- National Science and Technology Council (2019), National Space Weather Strategy and Action Plan, Washington, DC.
- Storz, M. F., B. R. Bowman, M. J. I. Branson, S. J., Casali, and W. K. Tobiska (2005). High accuracy satellite drag model (HASDM). *Advances in Space Research*, 36(12), 2497-2505.




Fig. 1. The SET HASDM database example shown for the Halloween storm period of 2003.

Tobiska, W. K., S. D. Bouwer, and B. R. Bowman (2008), The development of new solar indices for use in thermospheric density modeling, *Journal of Atmospheric and Solar-Terrestrial Physics*, 70(5), 803 – 819, doi:<https://doi.org/10.1016/j.jastp.2007.11.001>.