

UPDATED RADIATIVE TRANSFER MODEL FOR TITAN: VALIDATION ON VIMS-CASSINI OBSERVATIONS OF THE HUYGENS LANDING SITE AND APPLICATION TO THE ANALYSIS OF THE DRAGONFLY LANDING AREA.

M. ES-SAYEH¹, S. RODRIGUEZ¹, T. CORNET², L. MALTAGLIATI³, M. COUTELIER⁴, P. RANNOU⁴, B. GRIEGER⁵, E. KARKOSCHKA⁶, B. SEIGNOVERT⁷, S. LE MOUÉLIC⁷, C. SOTIN⁷

¹Université de Paris, Institut de Physique du Globe de Paris, CNRS, France

²Aurora Technology BV for European Space Agency (ESA), European Space Astronomy Centre (ESAC), Madrid, Spain

³Nature Astronomy, Springer Nature, London, UK

⁴Groupe de Spectrométrie Moléculaire et Atmosphérique, UMR CNRS 7331, Université de Reims Champagne-Ardenne, Reims, France

⁵RHEA System for ESA, European Space Astronomy Centre, Madrid, Spain

⁶Lunar and Planetary Laboratory, Tucson, AZ, USA and

⁷Laboratoire de Planétologie et Géodynamique, UMR 6112, CNRS, Université de Nantes, France

ABSTRACT

Titan, the only moon in the solar system with a thick atmosphere and a methane-based hydrological cycle similar to the water-based cycle on Earth, is a prime target for planetary and astrobiological researches. Organic materials from atmospheric chemistry precipitate on the surface and are subject to geological processes (e.g. eolian and fluvial erosion) that lead to the formation of dune fields, river networks, lakes and seas similar to their terrestrial counterparts. The analysis of the surface reflectance in the near-infrared (NIR) allows to constrain the surface composition, which is crucial to understand these atmosphere/surface interactions. However, Titan's atmosphere prevents the surface from being probed in the NIR, except in 7 transmission windows (centered at 0.93, 1.08, 1.27, 1.59, 2.01, 2.7 and 5 μm). We use an updated version of the Radiative Transfer (RT) model of Hirtzig et al. (2013), with updated gases and aerosols opacities, in order to better simulate atmospheric absorption and scattering and retrieve surface albedos in the 7 NIR transmission windows with an enhanced accuracy. Our RT model is based on the SHDOMPP and CDISORT solvers (Evans 2007) and (Buras et al. 2011) to solve the RT equations in a plane-parallel and pseudo-spherical approximations respectively. We recently improve atmospheric inputs of the model with up-to-date gaseous CH₄, CH₃D, ¹³CH₄, C₂H₂, HCN and CO abundances profiles and absorption coefficients (Vinatier et al. 2007), (Niemann et al. 2010), (Maltagliati et al. 2015), (Serigano et al. 2016), (Rey et al. 2018), (Thelen et al. 2019), (Gautier 2021), and improved the photochemical aerosol optical properties. In particular, the optical properties of Titan's aerosols are now computed from a fractal aggregate model (Rannou et al. 2003) constrained by in situ measurements of the Descent Imager/Spectral Radiometer (DISR) onboard the Huygens probe (Tomasko et al. 2008) and (Dose et al. 2016). Our RT model is benchmarked with the help of the most recent RT model for Titan in the literature (Coutelier 2021) and validated using the in situ observations of DISR acquired during descent and once landed. Coupled with an efficient inversion scheme, our model can be applied to the Cassini's Visual and Infrared Mapping Spectrometer (VIMS) complete dataset for the retrieval of Titan's atmospheric opacity and surface albedos at regional and global scales. This will help to analyzing the near-future observations of Titan with the James Webb Space Telescope (JWST) (Nixon et al. 2016) and preparing the future exploration of Titan by the Dragonfly mission (Lorenz et al. 2018).

REFERENCES

- Buras, R., Dowling, T., and Emde, C. (2011). New secondary-scattering correction in disort with increased efficiency for forward scattering. *Journal of Quantitative Spectroscopy and Radiative Transfer*, 112(12):2028–2034.
- Coutelier, M. (2021). Distribution and intensity of water ice signature in south xanadu and tui regio.
- Dose, L. R., Karkoschka, E., Tomasko, M. G., and Anderson, C. M. (2016). Vertical structure and optical properties of titan's aerosols from radiance measurements made inside and outside the atmosphere. *Icarus*, 270:355–375.
- Evans, K. F. (2007). Shdomppda: A radiative transfer model for cloudy sky data assimilation. *Journal of the atmospheric sciences*, 64(11):3854–3864.
- Gautier, T. (2021). Reevaluation of methane vertical profile at huygens landing site from huygens/gcms measurements.
- Hirtzig, M., Bézard, B., Lellouch, E., Coustenis, A., De Bergh, C., Drossart, P., Campargue, A., Boudon, V., Tyuterev, V., Rannou, P., et al. (2013). Titan's surface and atmosphere from cassini/vims data with updated methane opacity. *Icarus*, 226(1):470–486.
- Lorenz, R. D., Turtle, E. P., Barnes, J. W., Trainer, M. G., Adams, D. S., Hibbard, K. E., Sheldon, C. Z., Zacny, K., Peplowski, P. N., Lawrence, D. J., et al. (2018). Dragonfly: A rotorcraft lander concept for scientific exploration at titan. *Johns Hopkins APL Technical Digest*, 34(3):14.
- Maltagliati, L., Bézard, B., Vinatier, S., Hedman, M. M., Lellouch, E., Nicholson, P. D., Sotin, C., de Kok, R. J., and Sicardy, B. (2015). Titan's atmosphere as observed by cassini/vims solar occultations: CH₄, CO and evidence for C₂H₆ absorption. *Icarus*, 248:1–24.
- Niemann, H., Atreya, S., Demick, J., Gautier, D., Haberman, J., Harpold, D., Kasprzak, W., Lunine, J., Owen, T., and Raulin, F. (2010). Composition of titan's lower atmosphere and simple surface volatiles as measured by the cassini-huygens probe gas chromatograph mass spectrometer experiment. *Journal of Geophysical Research: Planets*, 115(E12).

- Nixon, C. A., Achterberg, R. K., Ádámkovics, M., Bézard, B., Bjoraker, G. L., Cornet, T., Hayes, A. G., Lellouch, E., Lemmon, M. T., López-Puertas, M., et al. (2016). Titan science with the james webb space telescope. *Publications of the Astronomical Society of the Pacific*, 128(959):018007.
- Rannou, P., McKay, C., and Lorenz, R. (2003). A model of titan's haze of fractal aerosols constrained by multiple observations. *Planetary and Space Science*, 51(14-15):963–976.
- Rey, M., Nikitin, A. V., Bézard, B., Rannou, P., Coustenis, A., and Tyuterev, V. G. (2018). New accurate theoretical line lists of 12ch₄ and 13ch₄ in the 0–13400 cm⁻¹ range: Application to the modeling of methane absorption in titan's atmosphere. *Icarus*, 303:114–130.
- Serigano, J., Nixon, C. A., Cordiner, M. A., Irwin, P. G., Teanby, N. A., Charnley, S. B., and Lindberg, J. E. (2016). Isotopic ratios of carbon and oxygen in titan's co using alma. *The Astrophysical Journal Letters*, 821(1):L8.
- Thelen, A. E., Nixon, C., Chanover, N., Cordiner, M., Molter, E., Teanby, N., Irwin, P., Serigano, J., and Charnley, S. (2019). Abundance measurements of titan's stratospheric hcn, hc₃n, c₃h₄, and ch₃cn from alma observations. *Icarus*, 319:417–432.
- Tomasko, M. G., Doose, L., Engel, S., Dafoe, L., West, R., Lemmon, M., Karkoschka, E., and See, C. (2008). A model of titan's aerosols based on measurements made inside the atmosphere. *Planetary and Space Science*, 56(5):669–707.
- Vinatier, S., Bézard, B., and Nixon, C. A. (2007). The titan 14n/15n and 12c/13c isotopic ratios in hcn from cassini/cirs. *Icarus*, 191(2):712–721.