

Deciphering the Cenozoic exhumation history of the Eastern Pyrenees along a crustal-scale normal fault using low-temperature thermochronology

G. Milesi^{1,2}, P. Monié¹, R. Soliva¹, P. Münch¹, P.G. Valla³, S. Brichau⁴, M. Bonno¹, C. Martin¹, M. Bellanger⁵

¹ Géosciences Montpellier, Université de Montpellier, CNRS, Université des Antilles, Montpellier, France

² Chrono-Environnement, UMR 6249-CNRS, Université Bourgogne Franche-Comté, Besançon, France

³ University of Grenoble Alpes, University of Savoie Mont Blanc, CNRS, IRD, IFSTTAR, ISTerre, 38000 Grenoble, France

⁴ Géosciences Environnement Toulouse (GET), Université de Toulouse, UPS, Univ. Paul Sabatier, CNRS, IRD, CNES, 14 av. Edouard Belin, F-31400 Toulouse, France

⁵ TLS Geothermics, 91 chemin de Gabardie, 31200 Toulouse, France

Corresponding author: Gaétan Milesi (gaetan.milesi@univ-fcomte.fr)

Key Points:

- Tectonic evolution along a major normal fault recorded by low-temperature thermochronology
- Priabonian (35-32 Ma) switch between contractional and extensional regime in the Eastern Pyrenees
- Cenozoic westward propagation of the deformation along the Têt fault

Abstract

The timing of transition between the contractional and extensional regimes along the Pyrenean range remains debated. Compared to its central and western parts, the eastern part of the chain was significantly affected by extensional tectonics mostly related to the opening of the Gulf of Lion. The Têt normal fault is the best example of this tectonic activity, with topographic reliefs above 2,000 m in its footwall. In this study, we synthesized previous thermochronological data and performed new (U-Th)/He and fission-track dating in the Eastern Pyrenean massifs. Output apparent exhumation rate and thermal modeling in the hanging-wall of the Têt fault highlight a rapid exhumation (0.48 km/Ma) and cooling (~30°C/Ma) phase between 38 and 35 Ma, followed by slower exhumation/cooling afterwards. In the footwall, cooling subsequently propagated westward along the fault during Priabonian (35-32 Ma), upper Oligocene and lower Miocene (26-19 Ma), and Serravalian-Tortonian times (12-9 Ma). These data and modeling outcomes suggest that the exhumation of the Têt fault hanging-wall related to southward thrusting ended at 35 Ma, and was followed by different extensional stages, with a propagation of the deformation towards the West during the upper Miocene. We propose that the onset of extension in the Eastern Pyrenees occurred during the late Priabonian period, contemporaneously with the large-scale rifting episode recorded in Western Europe. After this event, the Têt fault activity and the westward propagation of the deformation appear mainly controlled by the opening of the Gulf of Lion.

Plain Language Summary

The Pyrenees result from the North-South convergence of the Eurasian and Iberian plates. The eastern part of the range experienced strong extensional tectonics mostly related to the opening of the Gulf of Lion, which timing and influence on the modern topographic relief remain unclear. To better characterize the transition timing between contractional and extensional regimes and the tectonic evolution in the Eastern Pyrenees, we used low-temperature

thermochronology and thermal modeling to reconstruct the exhumation/cooling histories of the different massifs along the Têt fault. Our data and modeling outcomes show a switch between contractional and extensional tectonics during the Priabonian (ca. 35 Ma), followed by different extensional stages recorded in the Têt fault footwall, coeval with a global westward propagation of the deformation along the fault until ca. 9 Ma.

1 Introduction

In orogenic belts, crustal-scale faults are key deformation markers that accommodate various regimes of plate tectonics during rock burial, exhumation or strike slip activity (Jones and Wesnousky, 1992; Norris and Cooper, 2001; Ratschbacher et al., 2003; Viola et al., 2004; Malusà et al., 2009). When syn- to post-orogenic sedimentary record or chronological constraints are lacking for bracketing fault activity within orogens, low-temperature (low-T) thermochronology is a powerful tool to quantify the timing and magnitude of exhumation along major faults, since it provides time constraints on the thermal evolution of rocks during their exhumation towards the Earth's surface (Farley, 2002; Ehlers and Farley, 2003; Malusà et al., 2005; Glotzbach et al., 2011; Stockli, 2005; Colgan et al., 2006; Reiners and Brandon, 2006). This situation is common within orogens for which the transition between syn- and post-orogenic periods, or the transition from contraction to extension, remains difficult to date and is often highly debated (Price and Henri, 1984; Carmignani and Kligfield, 1990; Jolivet et al., 2020, 2021a; Séranne et al., 2021). Low-T thermochronology has been widely used in large-scale extensional domains to date the activity of normal faults, as for example in the Basin and Range Province (Foster et al., 1999; Surpless et al., 2002; Armstrong et al., 2003; Colgan et al., 2008) or the Aegean domain (Coutand et al., 2014; Brichau et al., 2006). However, few studies have investigated the onset of post-orogenic extension using low-T thermochronology (i.e. Cederbom et al., 2000; Danišik et al., 2012; Fillon et al., 2021; Martín-González et al., 2012) and even less on the lateral migration of the tectonic activity along normal faults in orogenic context (Deeken et al., 2006; Krugh, 2008; Curry et al., 2016).

In the Pyrenees, previous thermochronological studies have focused mainly on the central part of the chain, which is composed of a stack of crustal nappes formed during the main Eocene - early Oligocene orogenic build up (Jolivet et al., 2007; Mouthereau et al., 2014; Bosch et al., 2016; Labaume et al., 2016; Vacherat et al., 2016; Waldner et al., 2021). In the eastern part of the Pyrenees, less studies have been carried out (Maurel et al., 2002; 2008; Gunnell et al., 2009), which do not provide a detailed view of fault activity through time. This orogen segment shows a similar nappe structure as further West (Laumonier et al., 2015, 2017; Calvet et al., 2021) but has experienced significant post-orogenic crustal thinning to 25 km of total thickness, as indicated by recent geophysical data (Nercessian et al., 2001; Lacan and Ortuño, 2012; Chevrot et al., 2018; Diaz et al., 2018). This thinning is assigned to the presence of numerous and widely distributed normal faults onshore and offshore (Jolivet et al., 2020, 2021a; Romagny et al., 2020; Calvet et al., 2021; Séranne et al., 2021; Taillefer et al., 2021). The geodynamic origin for the onset of the extension has been linked to either the initiation of the West European Rifting which formed a large intraplate feature (Mouthereau et al., 2021; Angrand and Mouthereau, 2021) or the early onset of back-arc extension leading to the formation of the Gulf of Lion (Séranne, 1999; Séranne et al., 2021). The Têt fault is the most prominent normal fault of the Eastern Pyrenees, which localizes high-relief massifs in its footwall such as the Canigou and Carança (Fig. 1). The development of these high topographic reliefs has been attributed to normal faulting during the Oligo-Miocene period (Maurel et al., 2008). However, the pre-extensional history of this area, the onset of extension and its polyphase activity along strike during the Cenozoic are still poorly understood (e.g. Huyghe et al., 2020; Jolivet et al., 2020, 2021a, 2021b; Angrand and Mouthereau, 2021; Taillefer et al., 2021).

In this study, we present a new low-T thermochronology dataset from bedrock samples collected on both sides of the Têt fault, including (U-Th)/He on apatite (AHe) and zircon (ZHe), and apatite fission track (AFT). Low-T thermochronological data from previous studies (Maurel et al., 2002; 2008; Gunnell et al., 2009; Milesi et al., 2019; 2020a, 2020b) have been also synthesized with the new dataset, and all data are used for thermal modeling to assess the exhumation history of the footwall and hanging wall massifs along the southwestern segment of the Têt fault. Based on these results, we discuss the onset, timing and spatial evolution of Cenozoic extension in the eastern part of the Pyrenees as well as the potential driving mechanisms for this evolution.

2 Geological setting

2.1 Tectonic evolution of the eastern part of the Pyrenees

The Pyrenees result from the North-South convergence of the Eurasian and Iberian plates since the late Cretaceous (Choukroune et al. 1989; Roure et al. 1989; Muñoz, 1992; Beaumont et al., 2000; Mouthereau et al., 2014; Teixell et al., 2016), and form a double-wedged mountain range of around 1,000 km long and 150 km wide (Fig. 1a). The maximum of shortening occurred during the Eocene in the central part of the range (e.g. Vergés et al., 1995; Gibson et al., 2007; Sinclair et al., 2005; Metcalf et al., 2009; Whitchurch et al., 2011; Fillon and van der Beek, 2012; Mouthereau et al., 2014; Teixell et al., 2016; Curry et al., 2019). The Pyrenees are divided into three main latitudinal tectonostratigraphic domains (Vergés et al., 2002; Grool et al., 2018). To the North, three main units are recognized: the Aquitaine foreland basin, the Sub Pyrenean Zone and the North Pyrenean Zone, the last two being separated by the North Pyrenean Frontal Thrust (Fig. 1a). Further South, the North Pyrenean Fault (NPF) separates the North Pyrenean Zone from the Axial Zone and is interpreted as the suture between the Eurasian and Iberian plates. The Axial Zone consists of a stack of south-verging nappes made of late Proterozoic and Paleozoic sedimentary, metamorphic and magmatic rocks involved in the Variscan orogeny. The South Pyrenean Zone extends to the South of the Axial Zone and is composed of a sequence of Mesozoic to Eocene sediments involved in several thrust sheets transported southward. The Ebro Basin forms the southern foreland basin of the Pyrenean orogen.

In the eastern Axial Zone, it is accepted that the mountain building occurred through the emplacement of south-verging nappes rooted in the northern part of the Axial Zone, south of the NPF (Vergés et al., 1995; Sibuet et al., 2004; Laumonier et al., 2015; Teixell et al., 2016). In the studied area, the balanced cross sections of Ternois et al. (2019) suggest an Eocene thrusting of the Aspres–Mont-Louis massifs onto the Canigou massif, in agreement with available thermochronological data (Maurel et al., 2008). The reactivation of Variscan structures during the Pyrenean orogeny has been proposed, the most significant example being the Merens fault to the North of our study area (McCaig and Miller, 1986; Burbank et al., 1992; Guitard et al., 1998; Cochelin et al., 2017; Laumonier et al., 2017). The particularity of the Eastern Pyrenees is the reactivation of compressional structures during extensional tectonic regime (Séranne et al., 1995; Séranne 1999; Jolivet et al., 2020; Calvet et al., 2021; Séranne et al., 2021). This regional scale extension is witnessed by geophysical data that show a progressive crustal thinning, with crustal thickness varying between 45 km in the eastern part of the Axial Zone (~1°E) to 25 km at the margin of the Gulf of Lion (Chevrot et al., 2018; Diaz et al., 2018). This regional extensional episode led to the (re-)activation of major structures as normal faults with different orientations (NE-SW, NW-SE and N-S, Fig. 1b), from the end of the Oligocene to the Quaternary (Taillefer et al., 2021). Some of these faults have been considered as inherited ductile Variscan faults (Guitard et al., 1992, 1998; Bouchez and Gleizes, 1995; Autran et al., 2005; Laumonier et al., 2015, 2017). Two main NE-SW trending normal faults are recognized in the studied area: the Têt and the Tech faults (Fig. 1b). The Têt fault represents the southern margin of the Cerdagne and Conflent basins, while the Tech fault is the southern bounding fault of the Roussillon basin. Noteworthy is the importance of a NW-SE trending fault network that affects particularly the Mont-Louis and Carança massifs (e.g. Fontpédrouse and Nuria faults, Fig. 1.b) and cuts the North Catalan Coastal Range further South (Fig. 1.a). Some of these faults have nearly E-W directions probably recording spatial and/or temporal changes of stress orientation and/or stress regime. Major N-S faults in the eastern part of the Pyrenees are rare, among which the Capçir fault is described as a Quaternary normal fault (Briaïs et al., 1990). In the study area, the kinematics and amount of exhumation associated to these different faults are still debated. In Figure 1b, major crustal blocks have been differentiated and delimited by the Têt fault, namely the Mont-Louis block to the North (hanging wall) and Canigou-Costabonne and Carança blocks (footwall, delimited by the Py secondary fault) to the South.

Previous multi-thermochronological studies (Maurel et al., 2002; 2008; Gunnel et al., 2009) in the Canigou (footwall of the Têt fault) and Mont-Louis (hanging wall of the Têt fault) provided insights and results guiding our study. Maurel et al. (2002, 2008) proposed that the Canigou massif was exhumed during two periods, the first one at a rate of ~0.30 km/Ma between 27 and 21 Ma, followed by a significant slowdown of exhumation (~0.10 km/Ma) until present-day. In the Mont-Louis massif, thermochronological data suggest an earlier exhumation between 50 and 35 Ma (~0.30 km/Ma) accompanied by a rapid cooling. Since 35 Ma, the Mont Louis exhumation has been relatively slow, estimated at 0.04-0.06 km/Ma (Maurel et al., 2008). These different exhumation and cooling histories between the two massifs since 35 Ma were interpreted to be related to the normal motion of the Têt fault, without erasing the thermochronological record of Eocene tectonic activity in the hanging wall. In the Carança massif, thermal modeling based on AHe data (Milesi et al., 2019a; 2020b) suggests two main cooling events that occurred in the Oligo-Miocene, a major one between 30 and 24 Ma (at a rate of 25°C/Ma) followed by a second episode between 12 and 9 Ma (at a

rate of 15°C/Ma). Despite these previous thermochronological studies, the spatio-temporal evolution of the main tectonic structures in the eastern part of the Axial Zone of the Pyrenees since the Priabonian remains still poorly constrained (see Taillefer et al., 2021).




Figure 1. a) Structural map of the Pyrenees showing the main structural domains delimited by faults (modified after Taillefer et al., 2017). The major Neogene normal faults of the Eastern Pyrenees are reported in red. The study area is outlined with an open purple-dashed box. b) Structural sketch map of the study area showing the different massifs (in bold italics) and basins (in italics) along the Southwestern (SW) and Northeastern (NE) segments of the Têt fault (modified from Taillefer et al., 2021). Secondary faults are indicated by red numbers (see legend for details).

2.2 Tectonic evolution and sedimentary record along the Têt fault

The southern segment of the Têt normal fault is a NE-SW north-dipping and 100-km long crustal-scale fault (Maurel et al., 2002; 2008; Chevrot et al., 2018; Diaz et al., 2018; Fig. 1a). It crosscuts Palaeozoic magmatic and metamorphic rocks of the Mont Louis, Canigou and Carança massifs along which Neogene sedimentary basins developed (Fig. 1b). In the Canigou massif, the main period of fault activity during the Oligo-Miocene has been well constrained using low-T thermochronology (Maurel et al., 2002; 2008). A second stage of normal motion along the entire Têt fault has been recorded between the middle-Miocene and the late Pliocene, with associated vertical displacements in the range of 150-500 m (Pous et al., 1986; Rehault et al., 1987; Cabrera et al., 1988; Roca and Desegaulx, 1992; Tassone et al.,

1994; Calvet, 1999; Carozza and Baize, 2004; Delcaillau et al., 2004; Agustí et al., 2006; Clauzon et al., 2015) to kilometric (Calvet, 1996; Mauffret et al. 2001). However, thermochronological data in the Canigou massif (Maurel et al., 2008) are apparently not consistent with an hypothesis of km-scale vertical displacements. Since the end of Miocene, a main difference is recorded along the Têt fault between the western (Cerdagne basin) and eastern (Conflent and Roussillon basins) segments. Indeed, only the western segment of the Têt fault has been active (Calvet, 1999) which led to the opening of the Cerdagne pull-apart basin accommodated by normal (Pous et al., 1986; Agustí et al., 2006) and right-lateral displacement along the Têt fault (Cabrera et al., 1988). Based on geomorphological observations, a westward propagation of the deformation along the Têt fault has also been proposed to occur during the Plio-Pleistocene (Carozza and Delcaillau, 1999; Carozza and Baize, 2004). The amplitude of Pliocene to Quaternary normal activity on the eastern segment of the Têt fault is still debated. For some authors, the presence of triangular facets along the Têt fault scarp documents a recent normal fault activity (Briais et al., 1990; Calvet, 1999). However, Petit and Mouthereau (2012) suggested these are only the morphological expression of the differential erosion within Variscan mylonites. It is important to note that facets are also observed on scarps with no apparent mylonite nor favorably-oriented Variscan foliation (western segment of the Têt fault, Py and Capcir faults, Delmas et al., 2018). Finally, over the last 6 Ma, low incision rates of maximum 25 m/Ma in the Têt valley indicate weak vertical uplift in the study area (Sartégou et al., 2018), bringing further evidence to the ongoing discussion on Late-Miocene potential uplift from paleoelevation studies (Huyghe et al., 2020; Suc and Fauquette, 2012).

The sedimentary record is not continuous along the Têt fault system, and three main depositional areas can be distinguished from East to West: (1) the Roussillon basin bounded to the North by the northern segment of the Têt fault that is antithetic to the southern segment, (2) the Conflent basin that connects to the Roussillon basin to the East and (3) the Cerdagne basin along the southwestern segment of the Têt fault (Fig. 1b). The Roussillon basin is a large graben belonging to the West European Rift system and was highly subsident during the Oligocene-Aquitania interval that corresponds to the rifting phase preluding the Liguro-Provençal Sea opening. Post-rift deposits within the Roussillon basin were deposited in a passive margin geotectonic setting with low tectonic subsidence, and were deeply incised during the Messinian salinity crisis after which the passive margin sedimentation resumed during the Pliocene (Clauzon et al., 1987; Clauzon, 1990; Calvet et al., 2015; Calvet et al., 2021). The Conflent basin is an intramontane half-graben lying along the southwestern segment of the Têt fault, at an elevation ranging from 250 to 1,000 m. Its sedimentary infill is composed of up to ~1,000 m thick continental deposits, thought to be related to the main tectonic activity of the Têt fault (Guitard et al., 1998; Calvet et al., 2014). However, the stratigraphy of this basin is debated and the main sedimentary units, peculiarly an olistostrome with km-scale olistoliths originated from the Canigou massif, may be either early Burdigalian (Guitard et al., 1998; Calvet et al., 2014) or Pliocene (Clauzon et al., 2015). Towards the southwest, the Cerdagne basin, at an elevation of 1,100 m, is interpreted as a pull-apart basin formed by dextral-strike slip along the Têt fault (Cabrera et al., 1988). It has been infilled by 400 to 1,000 m of Neogene sediments divided in two depositional units from early Miocene and late Miocene, separated by an unconformity (Pous et al., 1986; Agustí and Roca, 1987; Cabrera et al., 1988; Roca, 1996). The source area of clastic sediments switched from the North to the South between these two units, with tectonic activity strongly decreasing during the late Miocene (Roca and Santanach, 1986; Cabrera et al., 1988).

3 Methodology

3.1 Low-temperature thermochronology

3.1.1 Sampling strategy

Our main objective is to quantify the exhumation and thermal evolution of the different crustal blocks separated by main regional faults, and to provide new data on the kinematic history of these faults. In the hanging wall of the Têt fault, two main blocks, separated by the Mérens fault, have been studied: respectively the North and South Mérens blocks. The North Mérens block is composed of the Millas and Querigut granitic massifs, and the South Mérens block is formed by Mont-Louis, Campcardos and Carlit massifs (Fig. 1b). In the footwall of the Têt fault, two main blocks, separated by the NE-SW trending Py fault, have been sampled: the Canigou-Costabonne block (eastern segment of the Têt fault, Canigou and Costabonne sub-blocks separated by the NW-SE Llipodère fault) and the Carança block (western segment). New AHe, AFT and ZHe ages have been obtained mainly in the footwall of the Têt fault (Carança and Canigou-Costabonne blocks), which represents a total of 44 AHe ages, 3 AFT ages and 25 ZHe ages (Tables 1 and 2). Thermochronological data from previous studies (Maurel et al., 2008; Gunnell et al., 2009) have been synthesized and supplemented by AHe ages from our previous studies (Milesi et al., 2019, 2020b). Note that we have

excluded samples affected by hydrothermalism and Rare Earth Element mobility, therefore not relevant to define regional exhumation and thermal evolution of the studied area (Milesi et al., 2019, 2020b, Fig. 2). Sample localities and corresponding thermochronological data from literature are summarized in Supplementary Section Table S1 and shown in Figure 2.

In the hanging wall of the Têt fault, six samples at an elevation between 730 m and 2,380 m were analyzed in the North Mérens block (DON, MAD and MTB). The South Mérens block (i.e. Mont-Louis massif) provided seventeen samples (CAR, CMPC, GAL, LPCH, ML, ST) with an elevation difference of ~1800 m between the lowest sample in the Têt Valley (1,081 m) and that of the summit of Campcardos (2,900 m). In the footwall of the Têt fault, the Costabonne massif includes four samples (GUIL and POMA) from Gunnell et al. (2009) and two samples (VER) dated in this study. In the Canigou massif, Maurel et al. (2008) reported thermochronological data on seven samples (CAN) collected along a profile from the base of the massif (970 m) to the summit (2,784 m). Three apatite samples (CAN4, CAN9 and CAN12), initially dated with the AHe population method, have been redated with AHe single grain method (see section 3.1.2). Two augen gneiss blocks (OL1 and OL2) from the olistostrome formation deposited in the Conflent basin and originating from the Canigou massif (Clauzon et al., 2015) have been also dated with the AHe single-grain method. In the Carança block, five new samples have been dated with the AHe method (GAL5, ST6, ST7, ST9 and ST10) to complete the AHe dataset from Milesi et al. (2019, 2020b). AFT ages have been obtained on three samples from different sampling profiles (ST2, GAL4 and TET4). Finally, a ZHe age-elevation profile (900 to 1900 m) has been realized with six samples from the Carança block (TET1.1, TET4, TET5, GAL7, GAL3, PLA3 and ST3).




Figure 2. Location of samples projected on DEM under GMT (Wessel et al., 2019) using SRTM1s. Different crustal blocks are delimited by regional major faults. From the North to the South, the sample names are for ST profile : ST2, ST3, ST4, ST10, ST9, ST8, ST7, ST6 and for GAL profile: GAL7, GAL6, GAL5, GAL4, GAL3, GAL1. Samples ST2, ST6, ST7, ST9, ST10, GAL4 and GAL5 were dated in this study.

3.1.2 Apatite and zircon (U-Th)/He dating

Apatite and zircon (U-Th)/He analyses were conducted at the Noble Gas Laboratory of Géosciences Montpellier (France). All samples were crushed and sieved, and apatite and zircon concentrates were obtained by heavy liquid

methods. Inclusion-free crystals with no evidence of fracture were hand-picked under a binocular microscope. Each single grain was packed in Pt tubes for apatite or Nb tube for zircon, placed under vacuum, and heated with a 1,090-nm fibre laser operating at 4.0W (900°C) for apatite and 6.2W (1,100°C) for zircon. We applied a duration of heating of 5 min for apatite and 15 min for zircon. After ^3He spiking, gas purification was achieved by a cryogenic trap and two SAES AP-10-N getters, and helium content was measured on a quadrupole PrismaPlus QMG 220. The ^4He content was determined by the peak height method and was 10–10000 times above typical blank levels. A second heating run using the same analytical procedure was systematically conducted to verify that more than 99% of ^4He was extracted during the first run. After helium extraction, Pt or Nb tubes were retrieved from the sample chamber and transferred in a 2 ml polypropylene conical tube. Samples were doubly spiked (^{230}Th and ^{233}U) and dissolved using procedures previously described by Wu et al. (2016) for apatite and Gautheron et al. (2021) for zircon. The resulting solutions were diluted, and U (^{233}U and ^{238}U) and Th (^{230}Th and ^{232}Th) were measured by using isotope dilution ICPMS. For age calculation, alpha ejection correction (Farley et al., 1996) was calculated using the Ft software (Gautheron and Tassan-Got, 2010; Ketcham et al., 2011). Durango apatite and Fish Canyon Tuff (FCT) zircon replicates were analysed between four unknown grains and yielded a mean age of 31.24 ± 2.18 Ma and 29.19 ± 1.19 Ma, respectively, during the different analyses of this study. These results are consistent with the Durango reference age of 31.02 ± 1.01 Ma given by McDowell et al. (2005) and FCT reference age of 28.30 ± 2.8 Ma (Reiners and Nicolescu, 2006). Conservatively, the He Partial Retention Zone (PRZ) for the zircon system is assumed to be between 140°C and 220°C (Guenther et al., 2013) and in the range of 40°C to 80°C for apatite (Stockli et al., 2000). It is important to note that the helium retention is sensitive to the crystal chemistry (eU values, chlorine content) and cooling history of samples (see Ault et al., 2019), and also that the PRZ can spread over a larger range of temperature (see Ault et al., 2019).

3.1.3 Apatite fission tracks

Apatite grains were mounted and polished for etching to reveal the natural spontaneous fission tracks. Apatites were etched using 5.5N HNO₃ at 20°C for 20s. Etched grain mounts were packed with mica external detectors and corning glass (CN5) dosimeters and irradiated in the Chilean CCHEN nuclear reactor. Following irradiation, the external detectors were etched using 40% HF at 20°C for 40 minutes. Analyses were carried out on an Olympus BX61 microscope at a magnification of $\times 1,250$, using a dry ($\times 100$) objective in the Dating laboratory of Géosciences Environnement Toulouse (France). Confined track-length measurements were performed using a drawing tube and digitizing tablet, calibrated against a stage micrometer. Single-grain AFT ages were calculated using the external detector method and the zeta calibration approach, as recommended by the I.U.G.S. Subcommittee on Geochronology (Hurford, 1990). Track-length measurements were restricted to confined tracks parallel to the c-crystallographic axis. Fission tracks in apatite shorten or anneal with increased temperature and duration of heating. For apatite of typical Durango composition (0.4 wt% Cl), experimental and borehole data (Green et al. 1989; Ketcham et al. 1999) show that over geologic time fission tracks begin to anneal at a sufficient rate to be measurable above $\sim 60^\circ\text{C}$, with complete annealing and total resetting of the apatite fission track age occurring between 100 and 120°C. This range of temperatures is usually labelled the apatite fission-track partial annealing zone (PAZ).

3.2 Thermochronological data interpretation

3.2.1 Age-elevation relationships (AER)

For each crustal block (Fig. 2), the age-elevation relationships (AERs) between the different thermochronological data have been used to estimate first-order apparent exhumation rates and also to get information on timing for potential changes in exhumation (i.e. break-in-slope in AERs). This approach is independent from the thermal structure of the block under consideration (e.g. Wagner et al., 1977; Fitzgerald et al., 1995; Braun, 2002; Fitzgerald and Malusà, 2019), but it relies on several assumptions and simplifications. First, it only considers the measured thermochronological ages without taking into account potential sample-specific kinetics from parent element content for instance (e.g. Ault et al., 2019). The AER approach also considers a vertical distribution of investigated samples (Stüwe et al. 1994), which is rarely the case in the field, and may also be influenced by potential changes in topography (Braun, 2002) or the presence of secondary faults. A major potential problem concerning the interpretation of AERs is the complexity of the exhumation scenario (i.e. number of segments which can be defined in an age-elevation dataset), we thus used a Bayesian Information Criterion (BIC) to select the appropriate model complexity (Schwarz, 1978). In this study, we followed the approach developed by Glotzbach et al. (2011) to determine the best-fitting AER estimates for AHe, AFT and ZHe data with minimization of the BIC.

3.2.2 Inverse thermal modeling under QTQt

In order to reconstruct the thermal history of the different crustal blocks (Fig. 2), time-temperature paths were modeled with QTQt 5.7.0 software (Gallagher et al., 2009; Gallagher, 2012) using AHe and ZHe single-grain ages and parameters (eU, Rs) together with AFT single-grain ages with length distribution data. QTQt software uses a Bayesian Markov chain Monte Carlo (MCMC) sampling method to infer sample time-temperature histories (Sambridge, 1999; Charvin et al., 2009). This software is particularly efficient to model together several samples from a same elevation profile. We parametrized modeling to allow all samples of a given elevation profile to evolve under a common thermal path with a typical geothermal gradient of $30^{\circ}\text{C} \pm 10^{\circ}\text{C}$ in order to take full advantage of the multi-sample inversion approach (Vermeesch and Tian, 2014). The radiation-damage model of Gautheron et al. (2009) has been chosen for the AHe, the kinetic models of Ketcham et al. (2007) for AFT and Guenther et al. (2013) diffusion model for ZHe. For each model, 100,000 iterations have been performed and the predicted vs. observed ages graph is systematically presented with output time-temperature histories. ZHe data are modeled only for the Carança block (where we obtained a ZHe elevation profile), and are used as first-order time-temperature constraints to define the thermal histories of the other crustal blocks (no available ZHe profile, only scarce individual data obtained with the population method).

4 Results

4.1 New thermochronological ages

4.1.1 Apatite and zircon (U-Th)/He

All AHe and ZHe single-grain ages obtained in this study are reported in Table 1. We also present different graphs of ages vs. Rs, eU and Th/U in the Supporting Information (Fig. S1). For the South Mérens block, an augen gneiss (sample ST13) was collected in the footwall of the Fontpédrouse fault and provides a mean AHe age of 16.7 ± 1.0 Ma. Two apatite grains have not been considered to calculate the mean AHe age due to their anomalous high eU content compared to the other grains, possibly due to U-rich inclusions in these apatite grains (Table 2 and Fig. S1). Note that ST13 has an AHe age younger than all AHe ages (all >25 Ma) previously obtained in the South Mérens block (Maurel et al., 2008; Milesi et al., 2020b). This cannot be explained by different Rs or eU values of the dated apatite grains (Table 1, Fig. S1) and therefore sample ST13 will be considered independently of other samples from the South Mérens block due to its particular structural position in the footwall of the Fontpédrouse fault (Fig. 3).




Figure 3. Synthesis of AHe and ZHe ages in the study area. Samples with green and italic labels are new samples from this study, those with black labels are from previous literature studies (Maurel et al. 2008; Gunnell et al., 2009; Milesi et al., 2019; 2020a; 2020b). Along altitudinal profiles, samples from North to South are : ST profile - ST3, ST4, ST10, ST9, ST8, ST7, ST6; GAL profile - GAL7, GAL6, GAL5, GAL3. Samples ST6, ST7, ST9, ST10 and GAL5 were dated in this study.

In the footwall of the Têt fault, three samples from the Canigou massif, previously analyzed using multigrain AHe approach, were re-processed using a single-grain approach. Sample CAN12 from the base of the profile (970 m) shows a mean AHe age of 16.7 ± 1.3 Ma that agrees with the multigrain AHe age of 18.8 ± 1.0 Ma (Maurel et al., 2008). On top of the massif (2,784 m), sample CAN4 displays larger single-grain AHe age dispersion between 24.3 and 33.5 Ma, without any clear relationship with the apatite chemical composition (Table 1 and Fig. S.1). The mean single-grain AHe age of CAN4 (27.9 ± 4.8 Ma), despite high uncertainty, is younger than the multigrain AHe age of 34.7 ± 1.7 Ma obtained on three aliquots by Maurel et al. (2008). At an intermediate elevation (2,050 m), a single apatite grain provides an AHe age of 19.6 ± 1.0 Ma for sample CAN8. In the southern Costabonne massif, two samples VER11 (1,560 m) and VER13 (1,935 m) show low intra-sample age dispersion, except one apatite grain excluded for the mean age calculation due to its important eU content and young AHe age (Table 1 and Fig. S.1). AHe ages are respectively of 30.6 ± 1.8 Ma for VER11 and 34.9 ± 1.8 Ma for VER13. In the olistostrome of the Conflent basin, two augen gneisses (OL1 and OL2, Fig. 2) provide five AHe ages with four of them between 40.8 ± 2.3 Ma and 49.5 ± 2.2 Ma, and one at 21.7 ± 1.1 Ma. In the Carança massif, a new AHe mean age of 11.9 ± 0.9 Ma has been obtained for a granite sample (GAL5), thus confirming previous single-grain AHe ages between 10.0 ± 0.4 Ma and 14.1 ± 1.1 Ma obtained for the GAL profile (Milesi et al., 2020b). In the western part of Carança block, samples ST6, ST7, ST9 and ST10 collected at a similar elevation provide mean AHe ages of 25.5 ± 6.5 Ma, 17.1 ± 1.4 Ma, 24.3 ± 3.0 Ma and 19.6 ± 4.4 Ma, respectively (Table 1 and Fig. 3). ST samples provide quite large intra-sample variability in AHe ages, which cannot be explained by the chemical characteristics (eU, Th/U) or the grain size (Rs).

In the Carança massif, seven samples collected at different elevations (from 900 to 1,900 m) have been dated using the single-grain ZHe method. These zircon grains have an eU content mostly ranging between 500 and 1900 ppm, except sample PLA2 (1,900 m) that contains two zircons with eU values above 3000 ppm. These samples do not display important intra-sample age variation and show mean ZHe ages increasing regularly with elevation from 22.0 ± 1.7 Ma to 32.5 ± 3.3 Ma. The two samples PLA3 (1,622 m) and TET5 (1,900 m) from the top of the profile display similar ZHe ages of 36.2 ± 2.9 Ma and 37.3 ± 3.0 Ma, respectively (Table 1 and Fig. 3).

Table 1. (U-Th)/He data on apatite and zircon. Single-grain data with * (and in italics) have been considered as outliers and not considered for the mean age calculation.

Block	Sample/ grain	Rs μm	U ppm	Th ppm	eU ppm	Th/U	4He ncc/g	± s ncc/g	Ft	Corrected age Ma	Error ±1σ (Ma)
Apatite											
South Mérens block	ST13 (42.50727N 2.15867E 1,289m) Augen gneiss										
	ST13a	81.8	12.4	18.6	16.9	1.5	29023.1	1160.9	0.84	17.0	0.8
	ST13b	81.3	14.1	16.0	17.9	1.1	28315.3	849.5	0.85	15.5	0.6
	ST13c	67.0	24.2	24.4	30.0	1.0	49313.1	986.3	0.82	16.7	0.7
	ST13d	59.9	14.3	14.6	17.8	1.0	30276.5	1211.1	0.82	17.2	0.7
	ST13e	61.3	77.2	73.7	94.9	1.0	143789.9	1437.9	0.82	15.4	0.7
	ST13f	45.5	48.5	59.8	62.9	1.2	102885.4	2057.7	0.7	18.1	0.8
	ST13g*	53.3	<i>130.3</i>	<i>158.8</i>	<i>168.4</i>	<i>1.2</i>	<i>356975.6</i>	<i>3569.8</i>	<i>0.8</i>	<i>22.9</i>	<i>1.0</i>
	ST13h*	43.9	<i>94.3</i>	<i>102.8</i>	<i>119.0</i>	<i>1.1</i>	<i>227047.7</i>	<i>2951.6</i>	<i>0.7</i>	<i>21.9</i>	<i>1.1</i>
									Mean	16.7	1.0
Carançon block	GAL5 (42.51287N 2.20037E 1,147m) Granite										
	GAL5a	57.3	20.6	5.1	21.8	0.2	19915.4	597.5	0.77	9.8	0.6
	GAL5b	62.7	11.2	5.6	12.5	0.5	15730.0	471.9	0.81	12.9	0.8
	GAL5c	60.8	7.8	2.1	8.4	0.3	10375.3	415.0	0.79	13.0	0.8
									Mean	11.9	1.8
	ST10 (42.4949N 2.17104E 1,383m) Augen gneiss										
	ST10a	61.3	50.1	28.0	56.8	0.6	109832.3	2196.6	0.77	20.9	1.2
	ST10b	56.3	62.3	27.3	68.8	0.4	94741.0	1136.9	0.75	15.3	0.8
	ST10c	75.1	56.1	26.4	62.4	0.5	103165.4	1547.5	0.81	17.0	0.9
	ST10d	67.0	29.1	7.4	30.8	0.3	76314.5	1526.3	0.81	25.2	1.3
									Mean	19.6	4.4
	ST9 (42.49203N 2.17351E 1,421m) Fractured augen gneiss										
	ST9a	60.4	46.6	17.7	50.8	0.4	119148.9	1191.5	0.79	24.7	1.2
	ST9b	68.1	38.6	10.2	41.0	0.3	83334.3	1666.7	0.83	20.4	1.0
	ST9c	68.0	59.8	25.8	66.0	0.4	158824.9	1588.2	0.81	24.5	1.3
	ST9d	63.2	54.0	19.8	58.8	0.4	164830.2	1648.3	0.84	27.8	1.7
									Mean	24.3	3.0
	ST7 (42.48421N 2.17433E 1,494m) Augen gneiss										
	ST7a	61.0	11.3	3.2	12.1	0.3	18475.0	923.8	0.79	16.1	0.8
	ST7b	64.6	16.1	4.6	17.2	0.3	31125.3	1245.0	0.80	18.7	0.9
	ST7c	70.5	24.1	5.9	25.5	0.2	40022.5	800.4	0.82	15.8	0.7
	ST7d	63.5	14.2	3.7	15.1	0.3	26010.2	1040.4	0.80	17.9	0.9
									Mean	17.1	1.4
	ST6 (42.48116N 2.17433E 1,533m) Augen gneiss										
	ST6a	82.1	7.4	2.6	8.1	0.4	27252.5	817.6	0.85	33.1	1.8
	ST6b	62.3	13.1	3.3	13.9	0.3	28447.2	1137.9	0.78	21.7	1.1
	ST6c	67.9	7.2	3.3	8.0	0.5	17354.2	867.7	0.82	21.9	1.0
									Mean	25.5	6.5
Canigou block	CAN12 (42.56647N 2.48237E 970m) Augen gneiss										
	CAN12a	77.8	32.7	17.4	36.9	0.5	67808.4	678.1	0.84	18.1	0.9
	CAN12b	61.3	32.4	10.6	35.0	0.3	54050.7	540.5	0.83	15.5	0.7
	CAN12c	102.6	8.3	24.1	14.1	2.9	25356.0	253.6	0.90	16.6	0.8
									Mean	16.7	1.3
	CAN8 (42.53956N 2.46652E 2,050m) Augen gneiss										
	CAN8a	80	33.5	2.4	34.1	0.1	67878.1	678.8	0.84	19.6	1.0
	CAN4 (42.51892N 2.45676E 2,784m) Augen gneiss										
	CAN4a	42.6	8.1	17.9	12.4	2.2	25850.1	258.5	0.71	24.3	1.5
	CAN4b	45.1	16.5	43.7	27.0	2.6	62074.6	620.7	0.74	26.0	1.4
	CAN4c	61.7	11.6	35.4	20.1	3.0	64660.9	646.6	0.80	33.5	1.9
									Mean	27.9	4.8
Costabonne block	VER11 (42.477943N 2.305973E 1,560m) Highly fractured augen gneiss with chlorite										
	VER11a	50.0	237.6	17.0	241.7	0.1	639343.3	6393.4	0.75	29.2	1.4
	VER11b	47.9	228.1	19.4	232.7	0.1	684958.3	6849.6	0.74	32.7	1.6
	VER11c	49.9	153.7	13.2	156.9	0.1	429612.5	4296.1	0.76	29.9	1.5
									Mean	30.6	1.8
	VER13 (42.471203N 2.343885E 1,935m) Augen gneiss										
	VER13a*	61.9	<i>377.9</i>	<i>156.1</i>	<i>415.3</i>	<i>0.4</i>	<i>969933.9</i>	<i>9699.3</i>	<i>0.81</i>	<i>24.0</i>	<i>1.3</i>
	VER13b	52.9	275.0	61.6	289.8	0.2	920151.0	9201.5	0.77	34.3	1.8
	VER13c	57.7	237.9	65.3	253.6	0.3	860128.9	8601.3	0.79	35.4	1.8
									Mean	34.9	1.8

Olistolithes	OL2 (42.55702N 2.39468E 780m) Fractured augen gneiss										
	OL2a	81.2	15.2	9.2	17.4	0.6	76552.5	765.5	0.85	43.1	2.0
	OL2b	38.6	41.7	49.8	53.7	1.2	100135.0	1001.4	0.71	21.7	1.1
	OL2c	82.5	13.0	16.3	16.9	1.3	87438.1	874.4	0.87	49.5	2.2
	OL1 (42.53754N 2.3375E 930m) Fractured augen gneiss										
	OL1a	48.4	17.2	10.7	19.7	0.6	72121.3	721.2	0.71	42.9	2.4
	OL1b	56.7	20.9	9.7	23.2	0.5	88868.5	888.7	0.78	40.8	2.3
Zircon											
Carana block	TET1.1 (42.52611N 2.2430555E 900 m) Granite with chlorite										
	TET1.1a	47.9	1185.3	530.6	1312.6	0.4	2188733.2	43774.7	0.70	19.7	1.6
	TET1.1b	71.0	632.9	201.8	681.3	0.3	1577103.0	41004.7	0.80	23.9	1.9
	TET1.1c	53.5	525.5	215.5	577.2	0.4	1158414.3	20851.5	0.77	21.6	1.7
	TET1.1d	54.7	581.0	338.7	662.3	0.6	1397788.4	29353.6	0.77	22.7	1.8
									Mean	22.0	1.7
	TET 4 (42.51175N 2.25487E 1,390m) Augen gneiss										
	TET4a	73.7	460.9	175.7	503.0	0.4	1449839.5	21747.6	0.83	28.7	2.3
	TET4b	60.8	548.1	770.2	733.0	1.4	2244859.4	24693.5	0.76	33.5	2.7
	TET4c	56.1	869.6	320.0	946.4	0.4	3000148.0	48002.4	0.75	35.2	2.8
									Mean	32.5	3.3
	TET 5 (42.49078N 2.23036E 1,900m) Augen Gneiss										
	TET5a	63.7	564.0	149.9	600.0	0.3	2172324.1	39101.8	0.77	38.9	3.1
	TET5b	67.8	933.1	423.6	1034.8	0.5	3586410.1	71728.2	0.79	36.3	2.9
	TET5c	68.5	1192.0	315.4	1267.7	0.3	4567608.3	68514.1	0.82	36.4	2.9
	TET5d	58.9	968.6	452.4	1077.1	0.5	3262406.1	48936.1	0.76	33.2	2.7
									Mean	36.2	2.9
	GAL7 (42.51505N 2.19904E 1,025m) Fractured fine grained gneiss with quartz and calcite veins and locally oxides										
	GAL7a	62.4	1090.8	483.6	1206.8	0.4	2620687.7	39310.3	0.80	22.4	1.8
	GAL7b	54.7	950.0	407.1	1047.7	0.4	2191714.2	39450.9	0.74	23.5	1.9
	GAL7c	52.3	1264.8	556.2	1398.3	0.4	3105604.0	52795.3	0.77	24.0	1.9
									Mean	23.3	1.9
	GAL3 (42.51018N 2.20525E 1,363m) Fine grained gneiss										
	GAL3a	55.4	1678.1	328.7	1757.0	0.2	4976741.5	59720.9	0.74	31.6	2.5
	GAL3b	59.8	685.1	436.9	789.9	0.6	2120801.6	31812.0	0.78	28.6	2.3
	GAL3c	67.4	685.6	477.6	800.3	0.7	2144053.6	40737.0	0.81	27.5	2.2
	GAL3d	66.0	881.9	207.1	931.6	0.2	2581295.1	41300.7	0.78	29.4	2.3
									Mean	29.3	2.3
	ST3 (42.50001N 2.16697E 1,174m) Unaltered gneiss with biotite										
	ST3a	59.9	1316.3	557.5	1450.2	0.4	3801126.6	49414.6	0.76	28.6	2.3
	ST3b	63.4	1802.8	314.4	1878.2	0.2	4910004.5	68740.1	0.77	28.0	2.2
	ST3c	61.3	1121.4	661.1	1280.0	0.6	2993707.8	47899.3	0.77	25.3	2.0
									Mean	27.3	2.2
	PLA3 (42.49343N 2.15462E 1,622 m) Fractured leucocratic gneiss and locally oxidized										
	PLA3a	49.5	2958.1	226.3	3012.4	0.1	9898474.3	89086.3	0.75	36.2	2.9
	PLA3b	50.4	3478.6	417.7	3578.8	0.1	11888683.1	106998.1	0.75	36.8	2.9
	PLA3c	61.3	1678.8	277.0	1745.2	0.2	6282251.9	50258.0	0.77	38.9	3.1
									Mean	37.3	3.0
Note. Ft: Alpha ejection correction (Farley et al., 1996)											

4.1.2 Apatite fission tracks (AFT)

In the Carana massif, three new AFT ages have been obtained for samples TET4 (1,390 m) and GAL4 (1,221 m) and ST2 (1,217 m) (Fig. 4). They are respectively of 17.4 ± 1.7 Ma, 15.2 ± 1.4 Ma and 17.4 ± 1.7 Ma, with related mean track lengths of 12.84 ± 0.50 μm , 12.81 ± 0.70 μm and 12.39 ± 0.50 μm . AFT data and mean track lengths are summarized in Table 2 and shown with literature data on Figure 4.




Figure 4. AFT central ages for the study area. Samples ST2, GAL4 and TET4 (in green) are from this study, AFT ages in black have been extracted from Maurel et al. (2008) and Gunnell et al. (2009) (See Supplementary Table S1 for details and locations).

Sample (elevation)	No. of crystals	Track density ($\times 10^6$ tr.cm $^{-2}$)			Age dispersion		U (ppm)	Central age (Ma $\pm 1\sigma$)	Mean track length (μ m)	StD (μ m)	No. of tracks measured
		ρ_d [Nd]	ρ_s [Ns]	ρ_i [Ni]	RE (%)	P χ^2 (%)					
TET4 1,390 m	10	1.183 [10391]	0.400 [119]	4.546 [1377]	0.1	67.37	48.0	17.4 \pm 1.7	12.84 \pm 0.5	1.81	35
GAL4 1,221 m	16	1.189 [10391]	0.333 [163]	4.659 [2203]	14.7	17.14	49.0	15.20 \pm 1.4	12.81 \pm 0.7	2.53	40
ST2 1,217 m	20	1.177 [10391]	0.307 [150]	3.601 [1741]	17.6	28.01	38.3	17.4 \pm 1.7	12.39 \pm 0.5	2.46	57

Table 2. Fission-track data for the Carança massif. Analyses were determined by the external detector method using 0.5 for the $4\pi/2\pi$ geometry correction factor. Apatite fission-track ages were calculated using dosimeter glass (CN-5; Analyst Stephanie Brichau, $\xi=341.8 \pm 7.8$) calibrated by multiple analyses of IUGS apatite age standards (Hurford, 1990). P χ^2 is probability of obtaining χ^2 value for ν degrees of freedom, where ν is the amount of crystals. Central age is a modal age, weighted for different precisions of individual crystals. In track density, ρ_d is the fission track

density of the standard U-glass (CN-5); Ns (spontaneous), Ni (induced) and Nd (dosemeter) are the fission track numbers corresponding to ps, pi and pd, respectively.

4.2 AERs and apparent exhumation rates

4.2.1 Hanging wall of the Têt fault

In the hanging wall of the Têt fault, AERs are presented only for the South Mérens block (Fig. 5a). AERs based on AFT and AHe data suggest a three-stage exhumation scenario defined by the lowest BIC (Fig. 5a). Samples between 1,400 m and 2,400 m provide AFT central ages between 32.3 ± 3.4 Ma and 38.6 ± 2.4 Ma, corresponding to a mean apparent exhumation rate of 0.48 km/Ma. The uncertainty on this exhumation rate is relatively large (from 0.22 to 1.90 km/Ma) because most samples lie on an apparent vertical straight line. Samples CMPC1 and CMPC2 from the top of the profile (2,900 m), with AFT central ages ~ 50 Ma, indicate a lower apparent exhumation rate (0.04 km/Ma) that prevailed between ~ 35 and 50 Ma (Fig. 5a, upper graph); although the two ZHe ages in this block suggest potential variability in the exhumation rate during this period. CMPC1 and CMPC2 are the westernmost samples, it may also be possible that they have experienced different exhumation than other samples further East. However, these are the only thermochronological data available above 2,400 m for the South Mérens block, so we cannot assess further this potential difference.

AHe ages from samples above 1,700 m indicate an apparent negative exhumation rate between 35 and 40 Ma. Sample ML3 (2,030 m), which presents an AHe mean age older than its AFT central age has not been considered. This age inversion can find several explanations: an excess helium in the apatite grains (Green et al., 2006), the presence of inclusion inside or rich U-Th grain boundary phases (Murray et al., 2014). Sample ST13 is not presented in the Figure 5, its mean AHe age (16.7 ± 1.0 Ma) is younger than that of other samples and cannot be explained by the regional AER trend. The particular structural location of this sample in the footwall of the Fontpédrouse fault, close to the fault corner between Fontpédrouse (NW-SE) and the Têt fault (NE-SW) can explain the specific exhumation history due to the NW-SE fault activity (see Section 2.1). The negative apparent exhumation rate obtained can be due to: (1) the small number of samples (4 in total) above 1,700 m used to precisely define an exhumation rate in this block; (2) a change in AHe kinetics due to the rapid exhumation (e.g Ault et al., 2019); (3) a major decrease of relief during this period (Braun, 2002; Reiners, 2007; McDannell et al., 2014). This AER above 1,700 m is strongly influenced by AHe mean ages from CMPC1/2 samples at the top of the profile (Fig. 5a, lower graph), and can be explained only by rapid exhumation rates, consistently with the exhumation rates derived from the AFT central ages during this period (Fig. 5a, upper graph). Samples between 1,000 m and 1,700 m (Fig. 5a, lower graph) provide AHe mean ages between 24.2 ± 4.0 Ma and 40.0 ± 2.0 Ma, suggesting an important decrease in the apparent exhumation rate (0.05 km/Ma). For comparison, AFT ages in the North Mérens block support a mean apparent exhumation rate of 0.46 km/Ma between ~ 52 and 48 Ma, with high uncertainty due to the low number of AFT central ages obtained for this block (see Supporting Information Fig. S2).




Figure 5. Age-Elevation Relationships (AERs) for AFT and ZHe (first row) and AHe data (second row) for (a) the South Mérens block (the AHe mean age of sample ML3 at elevation of 2,050 m, with AHe mean age older than AFT central age has not been considered), (b) the Canigou-Costabonne massif and (c) the Carança massif.

4.2.2 Footwall of the Têt fault

In the footwall of the Têt fault, the Canigou-Costabonne (Fig. 5b) and Carança blocks (Fig. 5c) are separated by the Py fault and therefore their AERs have been considered individually. On Figure 5b (upper graph), the AER deduced from AFT data in the Canigou sub-block (between 970 m and 2,784 m), suggests a single exhumation phase between ca. 22 and 27 Ma, with an apparent exhumation rate of 0.33 km/Ma. AHe mean ages from the same block (Fig. 5b, lower graph) are between 16.7 ± 1.8 Ma and 34.7 ± 2.5 Ma, suggesting an apparent exhumation rate of 0.16 km/Ma from the Priabonian to the end of the Burdigalian. South of the Canigou massif, samples from the Costabonne massif do not show enough elevation difference to provide a reliable exhumation rate from AERs. However, it can be noted that for samples taken at similar elevations in these two massifs, the AFT and AHe ages are 1 to 10 Ma older in the Costabonne massif than in the Canigou massif (Fig. 5b).

In the Carança massif (Fig. 5c), both ZHe and AHe data have been used to constrain apparent exhumation rates from AERs. Three AFT central ages cannot be used given the limited elevation distribution (Fig. 5c, upper graph). ZHe data obtained on 7 samples show a quasi-ideal AER with an apparent exhumation rate of 0.06 km/Ma between ca. 37 and 22 Ma. AHe data suggest a similar apparent exhumation rate (0.07 km/Ma), between ca. 22 and 10 Ma, with some age variability for samples between 1,250 and 1,550 m on the ST profile, probably due to the proximity of secondary NW-SE faults that locally fragmented the massif in many sub-blocks (Fig. 5c, lower graph). We can also note that the

AER slope defined between 17 and 15 Ma by the three AFT central ages of the Carança block is in agreement with that derived from AHe mean ages from 20 to 10 Ma (Fig. 5c).

4.3 Thermal evolution

4.3.1 Hanging-wall of the Têt fault

The thermal history of the South Mérens block has been derived for all AHe (30) and AFT (12) data from 16 samples used to define AERs (Fig. 5a). For this block, the two ZHe ages of samples ML1 and ML6 (Fig. 3) have been used as time-temperature constraints for numerical modeling. Another model set-up, including AHe ages of ST13 and ML3 samples and without any ZHe constraint, has been considered and is presented in the Supporting Information (Figure S3). The output thermal evolution, depicted on Figure 6a, shows that between 50 and 38 Ma, the South Mérens block experienced a cooling rate of around 5°C/Ma, followed by an abrupt acceleration in cooling (~30°C/Ma) between 38 and 35 Ma. Then, since 35 Ma, this block was experiencing slow and continuous cooling (<1°C/Ma). Similar results have been observed in the alternative model (Supporting Information Figure S3), while AHe ages of ST13 and ML3 samples cannot be correctly reproduced (Figure S3). With the exception of two AHe ages, all predicted AHe, AFT ages and track lengths are consistent with the observed data implemented for inverse modeling (Fig. 6a).

4.3.2 Footwall of the Têt fault

For the Têt footwall, QTQt thermal modeling was conducted successively on the Canigou and Carança blocks, which are separated by the Py fault. In the Canigou block, data available in the Costabonne sub-block were not considered due to the presence of the Llipodère fault between the Canigou and Costabonne sub-blocks (Fig. 1b) and the lack of data under 2,200 m (only 2 samples with AHe method, VER11 and VER13). An alternative modeling set-up with data from Costabonne sub-block is available in the Supporting Information (Fig. S3). The Canigou thermal modeling (Fig. 6b) was designed with all the AHe (12), AFT (6) and track-length data from 7 samples available from the bottom to the top of the massif (thermal modeling output without ZHe constraint is available in the Supporting Information, Figure S3). The output thermal history suggests an important cooling event until ca. 33 Ma (onset timing not precisely constrained) at around 30°C/Ma, followed by slow cooling (<1°C/Ma) until ca. 26 Ma. A second cooling phase at ~10°C/Ma can be observed between 26 and 19 Ma, followed by slow cooling until present-day. The thermal history reproduces well AHe, AFT ages and mean track lengths, except the AHe age of sample CAN9 (2,100 m) and mean track lengths measured on samples from the Canigou summit (CAN4 and CAN5). Thermal modeling based on data from the Costabonne sub-block (Supporting Information, Figure S3) also suggests rapid cooling (30°C/Ma) for this block between 32 and 29 Ma, followed by slow cooling (<1°C/Ma); however this model output should be considered with caution due to the small amount of data (4 AFT and 2 AHe). This rapid cooling would be consistent with an early Oligocene cooling phase, before the Oligo-Miocene phase recorded between 26 and 19 Ma for the Canigou massif (Fig. 6b).

The modeled thermal history of the Carança block (Fig. 6c) is based on AHe (59), AFT (3) and ZHe (24) data from 20 samples. Output thermal history reveals slow cooling (<1°C/Ma) of the massif between 40 and 25 Ma. The main cooling phase at ~20°C/Ma occurred between 25 and 21 Ma, followed by slow cooling (<1°C/Ma) until 12 Ma. A second cooling pulse, of relatively minor magnitude, can be observed between 12 and 9 Ma with a predicted cooling rate of 10°C/Ma, and is followed by slow cooling (<1°C/Ma) since 9 Ma. Despite the important amount of data and an apparent dispersion of AHe ages (see Fig. 6c), the modeled thermal history reproduces well the AHe ages (except for samples GAL6, GAL3, ST6 and ST9), AFT ages and ZHe ages (except for sample GAL7). However, we can note that the predicted mean track lengths are not well reproduced and are generally longer than the observed ones (Fig. 6c).




Figure 6. Thermal history of (a) South Mérens block from the hanging wall of the Têt fault, (b) Canigou massif and (c) Carança block from the footwall of the Têt fault. Thermal models were computed using QTQt software (Gallagher, 2012). T-t paths for the uppermost (blue) and the lowermost (red) samples are presented (dashed lines correspond to 95 % confidence interval). Black boxes are constraints based on ZHe data from South Mérens block and Canigou massif, ZHe data are modeled for the Carança block. To the right, age-elevation profiles using predicted vs. observed ages for each block are presented as well as observed and predicted track lengths. AHe ages represented with orange error bars in the South Mérens block are not used to construct the thermal evolution model. Sample names for which several thermochronometers were used are indicated in bold. Note that mean predicted/observed data are presented

for clarity, but that thermal modeling has been using/predicting single-grain AHe/ZHe data and (U-Th)/He ages (uncorrected for alpha ejection; Farley et al., 1996).

5 Discussion

5.1 The Têt fault hanging wall: contractional stage

In the hanging wall of the Têt fault, North and South Mérens blocks were distinguished in the present study. In the North Mérens block, AHe mean ages are between 30 and 40 Ma (Fig. 3), while AFT central ages are between 45 and 54 Ma (Figs. 4 and S3). These ages are older than those obtained at similar elevations in the South Mérens block. This difference in low-T thermochronological data suggests an early exhumation of the North Mérens block during the Early Eocene, which is in agreement with McCaig and Miller (1986), who proposed on the basis of $^{40}\text{Ar}/^{39}\text{Ar}$ mica dating that the Mérens fault was reactivated southward around 50-60 Ma. The scarcity of data in the North Mérens block has not allowed to perform thermal modeling.

The thermal history of the South Mérens block (Fig. 6a), obtained using AHe and AFT data, highlights a first stage of cooling between 50 and 38 Ma ($> 5^\circ\text{C}/\text{Ma}$), that is coeval with a period of maximum shortening in the Eastern Pyrenees that has been evidenced in the Agly-Salvezines massifs to the North of our study area (Ternois et al., 2019). This cooling stage became more rapid between 38 and 35 Ma ($\sim 30^\circ\text{C}/\text{Ma}$, Fig. 6a). The fast exhumation rate that prevailed during this last cooling stage (0.45 km/Ma from AER, Fig. 5a) can be associated with the activity of the Cadi-Canigou thrust fault that emerges further South (Ternois et al., 2019). This thrust is one of the major fault accommodating the convergence between the Iberian and European plates during the Eocene (also see Fitzgerald et al., 1999; Whitchurch et al., 2011; Rushlow et al., 2013; Mouthereau et al., 2014; Bosch et al., 2016; Labaume et al., 2016; Cruset et al., 2020). This interpretation is also consistent with the general propagation and stacking of the nappes from the North to the South in the Pyrenees (Jolivet et al., 2007; Fillon and van der Beek, 2012; Cruset et al., 2020).

At around 35 Ma, our thermal model output suggests that nearly all the samples collected from 1100 m to 2900 m were above their respective PAZ and PRZ. After 35 Ma, low cooling rates are consistent with an important decrease in exhumation towards present-day in the Têt-fault hanging wall (Fig. 5a). This is in agreement with the recent exhumation model for the Axial Zone proposed by Curry et al. (2021). On the basis of a regional thermochronological data compilation and thermo-kinematic modeling (for details see Curry et al. 2011), this exhumation model suggests that rock uplift rates peak at 30-40 Ma in the Eastern Pyrenees, about 10 Ma earlier than in the western Pyrenees (see also Fillon and van der Beek, 2012 for a similar conclusion).




Figure 7. Output thermal histories for the study area: the South Mérens block (blue), the Canigou massif (red, with the associated box for ZHe constraint) and the Carança block (green). Thermal models were computed using QTQt software (Gallagher, 2012). Main cooling events are indicated by purple (hanging wall of the Têt fault) and grey (footwall of the Têt fault) bars.

5.2 The Têt fault footwall : extensional stage

In the different crustal blocks from the southern Têt fault footwall, we used a large number of ZHe, AFT and AHe data to constrain output thermal histories that emphasize multiple cooling phases since the end of the Eocene. (Fig. 7). A first fast cooling ($\sim 25^{\circ}\text{C}/\text{Ma}$), that started at an unconstrained period but ended at ca. 33 Ma, is recorded essentially by samples from the top of the Canigou massif (CAN4 and CAN5). Within these two samples, the differences between modeled and observed mean track lengths (Fig. 6b) can be explained by the the small amount of measured tracks ($n=30$ and 69 , respectively, see Fig. 4). We can note that zircon fission-track ages of Maurel et al. (2008) from the top and bottom of the Canigou massif are very similar (30.9 ± 2.5 Ma and 33.8 ± 2.1 Ma respectively, see Supporting Information Table S1). This suggests an important exhumation step of at least 2,000 m during the Priabonian-Rupelian period, which is not recorded further West in the Carança block by the ZHe data (Fig. 5c). Thermochronological data from the Costabonne massif are also consistent with an early Rupelian cooling phase in the Py fault footwall (Supporting Information Figure S3). The Py normal fault is a NW dipping master fault between the Canigou and Costabonne massifs (with numerous field evidence of substantial displacement : triangular facets, metric fault core with gouges) that branches out on the Têt fault to the North (Fig. 1 and 2).

This important exhumation signal in the Canigou and Costabonne massifs is better explained by normal faulting rather than south-verging thrusting at regional scale, such as described further South of the study area (e.g. Cruset et al., 2020). We propose that this interpretation of exhumation before 33 Ma is only relevant to the Canigou and Costabonne massifs (Figs. 1 and 6) and not to the whole Canigou-Carança range in agreement with Ternois et al. (2019). The Têt and Py faults had probably both accommodated the main exhumation of the Canigou and Costabonne blocks, the normal activity of the Py fault (or both faults) resulting in maintaining the Carança block at depth to the West. Normal activity of the western part of the Têt fault (Carança block) cannot be excluded due to the connection between the Py and the Têt faults (Fig. 1). The normal activity of the Py fault thus explains why the low-T thermochronometers used in our study do not record any cooling below PRZ nor PAZ during this period in the Carança block. In a contractional context, the diachronism between the Canigou and Carança blocks would require the presence of a master reverse back-thrust between these two blocks, which is not supported by field observations along the Py fault. Because the South Mérens block was already at shallow crustal level and thus has not recorded any significant cooling/exhumation since 35 Ma, both the Py fault and the southeasternmost segment of the Têt fault were probably active during the Priabonian-Rupelian period to allow for the exhumation of the Canigou-Costabonne massifs only.

The second major cooling event from our output thermal histories occurred between the upper Oligocene and the lower Miocene (i.e. ca. 26 to 19 Ma), and was recorded by both the Canigou and the Carança massifs (Fig. 7). During this period, the Canigou massif experienced relatively fast exhumation (0.33 km/Ma from AER, Fig. 5b). This cooling/exhumation signal can be thus associated to normal faulting all along the Têt fault. In the Canigou massif, low-T thermochronology data do not document any major cooling/exhumation since 19 Ma, suggesting that the southeastern segment of the Têt fault remained partly inactive since the Burdigalian. This is in agreement with the sedimentary record in the Conflent basin, showing that the main subsidence, associated with normal activity of the eastern segment of the Têt fault, was concentrated from the Aquitanian to the Early Burdigalian (Calvet et al., 2014). In addition, the AHe mean ages (mostly older than 40 Ma) obtained on gneiss samples from the olistotrome formation in the Conflent basin suggest that the olistolithes collapsed during this upper Oligocene–lower Miocene phase of significant exhumation. Indeed, AHe mean ages from the olistotrome formation are older than for modern bedrock samples at the top of the Canigou profile (AHe mean ages about 30 Ma, Fig. 5b). These old ages also show that the olistolithes were not buried enough to reset the AHe signal.

In the Carança block, our AHe data allow to differentiate two sub-blocks separated by the NW-SE Fontpédrouse fault (Figs. 1 and 3). AHe mean ages from the eastern sub-block (TET and GAL samples) are younger (10-15 Ma) than for the western sub-block (ST and PLA samples) collected at similar elevations (15-25 Ma, Fig. 3). This AHe age difference is obvious for samples between 1,250 and 1,550 m (Fig. 5c). In addition, the Fontpédrouse normal fault propagates in the South Mérens block, and it seems likely that the AHe mean age of 16.7 ± 1.0 Ma obtained close to this fault (sample ST13) recorded the fault activity during the Burdigalian (see also alternative thermal modeling in

Supporting Information, Figure S3). Note that despite the proximity of a huge gouge zone and evidence for fluid alteration, the Rare Earth Element distribution of this sample remains unaffected by hydrothermalism (see Supporting Information, Figure S4) compared to our previous observations along the Têt fault itself (Milesi et al., 2020b). NW-SE trending faults are frequent in this western segment of the Têt fault (see Milesi, 2020; Taillefer et al., 2021) and their activity can account for an important segmentation of the Carança massif with therefore a spatial variability in AHe data due to slightly different cooling histories within the different sub-blocks. In spite of these local perturbations by NW-SE faults in the Carança block, AHe and ZHe data are well reproduced by the QTQt model (Fig. 6c), and only mean track lengths show important differences between observed and modeled data, which can be explained by the small amount of tracks measured on the three samples (see Table 2 for details).

A third cooling event has been recorded between 12 and 9 Ma (Serravalian-Tortonian) but only for the Carança block (Fig. 7). The lack of record in the Canigou-Costabonne and South Mérens crustal blocks suggests a tectonic activity limited to the southwestern segment of the Têt fault, rather than a general exhumation of the eastern part of the Pyrenees (Huyghe et al., 2020; Calvet et al., 2021). This relatively recent activity can explain the preservation of triangular facets along the Têt fault (Petit and Mouthereau, 2012; Delmas et al., 2018) and is also consistent with the syntectonic sedimentation of late-Miocene age recorded by the lower unit in the Cerdagne basin (Pous et al., 1986; Augusti and Roca, 1987; Roca, 1996). The opening of the Cerdagne pull-apart sedimentary basins appears essentially controlled by the development of the NW-SE normal faults, facilitated by pre-existing NW-SE segments along the Têt fault (Cabrera et al., 1988).

5.3 Fault system evolution model and geodynamic implications

In the eastern part of the Pyrenees, North-South shortening has been recorded until ca. 35 Ma by our low-T thermochronological data. This is consistent with the timing for late contractional episode on the North Pyrenean Thrust Front (Grool et al., 2018) and the last main peak of pyrenean activity (Bartonian-Priabonian) recorded in Provence (Lacombe and Jolivet, 2005). On another side, new U-Pb on calcite studies suggest that shortening in the external units of the Pyrenees proceeded until the middle Miocene (Cruset et al., 2020; Hoareau et al., 2021; Parizot et al., 2021), which could be a consequence of the far field stress imposed by Africa-Europe convergence (Jolivet et al., 2021b; Mouthereau et al., 2021). Based on the sedimentary record, a recent study in the Gulf of Lion margin revealed that the shift between the pyrenean contractional and extensional tectonics occurred during the late Rupelian (~30 Ma, Séranne et al., 2021), with evidence for a rapid change in the tectonic regime. Although the timing of this shift in tectonic regime is globally consistent (see Section 5.2), our results suggest a slightly earlier onset of normal faulting along the Py and Têt faults, i.e. during the Priabonian, and an end of extensional tectonics at ca. 33 Ma (Fig. 7). We should also note that previous thermochronological studies proposed a large-scale episode of exhumation recorded in the Eastern Pyrenees between 35 Ma and 30 Ma (Morris et al., 1998) that could be regarded as a consequence of normal faulting, rather than thrusting. This first extensional event preceded a ~7 Ma long period of exhumation quiescence between 33 and 26 Ma (Fig. 7), which is synchronous to the development of back-arc extension in the Mediterranean domain (onset at 32–30 Ma, Jolivet and Faccenna, 2000). Thus the first exhumation and coeval extensional tectonic phase does not appear to be related to the rifting phase leading to the opening of the Liguro-Provençal domain, especially with regard to the specific configuration of the Py fault (Fig. 8a, i.e. oriented N030E compared to the N060E main trend of the Gulf of Lion faults). This event may rather correspond to the West European Rifting from strain geometry and age of exhumation (Ziegler, 1992; Romagny et al., 2020; Angrand and Mouthereau, 2021; Jolivet et al., 2021b; Mouthereau et al., 2021; Séranne et al., 2021). The West European Rifting is considered geodynamically independent and can lead or be immediately followed by the Gulf of Lion opening (Réhault et al., 1984; Séranne, 1999; Vignaroli et al., 2008; Jolivet et al., 2015, 2020).

A second extensional event has been recorded between the upper Oligocene and Burdigalian for the whole Canigou-Carança range, associated to a main normal faulting phase along the Têt fault (Figs. 7 and 8b). This event corresponds to the main cooling event recorded by Maurel et al. (2008), and it appears to be related to the opening of the Gulf of Lion, consistently with sedimentary records on the Catalan margin (Bartrina et al., 1992). In terms of direction of extension (NW-SE), this event clearly corresponds to the NE-SW trend of the faults observed in the Languedoc, Roussillon, Catalan and Valencia troughs, as well as offshore faults observed at the margin of the Gulf of Lion (e.g. Séranne, 1999; Mauffret et al., 2001; Maillard et al., 2020; Romagny et al., 2020; Jolivet et al., 2021a). In terms of timing, this second extensional event appears slightly younger than the onset of rifting in Languedoc (late Rupelian, Séranne, 1999), and earlier than the second stage of normal faulting on the Catalan margin (Roca and Desegaulx, 1992), probably reflecting the rift propagation towards the Southwest (Séranne, 1999).

A third extensional event (Fig. 8c) has been recorded by AHe data in the Carança and the South Mérens blocks, not in the Canigou-Costabonne block (Fig. 3). In the Carança massif, AHe data suggest a change in the direction of extension from NW-SE to NE-SW during the Lower-Miocene times (ca. 18 Ma), with normal-sense movement on the NW-SE Fontpédrouse fault. This stage evolved afterwards between 12 and 9 Ma on the southwestern segment of the Têt fault, commonly associated to a reactivation stage with moderate normal displacements between 150 and 500 m (Pous et al., 1986; Clauzon et al., 1987; Rehault et al., 1987; Roca and Desegaulx, 1992; Tassone et al., 1994; Calvet, 1999; Carozza and Baize, 2004; Delcaillau et al., 2004; Agustí et al., 2006; Clauzon et al., 2015). AHe data along the Têt fault reveal that the exhumation was probably more pronounced along the southwestern segment (>500 m). This stage, that is not recorded by low-T thermochronology data in the Canigou massif (Maurel et al., 2008, this study), marks differential exhumation along the Têt fault, more pronounced at this stage in the southwestern part, consistently with sediment infills of the Cerdagne basin (Pous et al., 1986; Agustí et al., 2006). This late activity on the southwestern segment of the Têt fault confirms the southwestward propagation of the exhumation along the Têt fault (Carozza and Delcaillau, 1999; Carozza and Baize 2004). The direction of extension is also consistent with Middle-Miocene to Pliocene normal faulting in the Emporda basin and the North-Catalan Ranges that trends globally NW-SE (Medialdea et al., 1994; Saula et al., 1994; Tassone et al., 1994; Lewis et al., 2000; Taillefer et al., 2021). Moreover, the pull-apart opening of the Cerdagne basin, accommodated by normal activity of NW-SE to E-W faults (Pous et al., 1986; Agustí et al., 2006) and right-lateral displacements on NE-SW faults (Cabrera et al., 1988), suggests that the main direction of extension was NNE-SSW, allowing the NE-SW Têt fault to be reactivated in right-lateral strike-slip movement (Fig. 8, Cabrera et al., 1988, Goula et al., 1999; Carozza and Baize, 2004, Delcaillau et al., 2004). We should also note that this trend of extension is also compatible with the stress tensors obtained in the Cerdagne area by Cruset et al. (2020). NW-SE faults could therefore have contributed to the uplift of the Cerdagne basin during Middle Miocene (Huyghe et al., 2020; Calvet et al., 2021; Tosal et al., 2021).




Figure 8. Reconstitution of the extensional tectonic evolution since the Priabonian in the eastern part of the Pyrenees. a) Priabonian-Rupelian period (35-32 Ma) is marked by the exhumation of the Canigou-Costabonne massif, linked to the Py fault normal motion. A WNW-ESE direction of extension is proposed. b) Aquitanian-Burdigalian

period (26-19 Ma) is characterized by the opening of the Gulf of Lion and normal motion of the Têt fault, more pronounced on the eastern segment of the fault. This observation is in agreement with the early formation of the Conflent basin at 23 Ma. During this period, the Têt fault normal activity is associated to a change towards the North of extensional direction c) Burdigalian to Tortonian period (19-9 Ma) reveals a propagation further West of the exhumation along the Têt fault with late cooling event recorded for the Carança massif. Local (re-)activation of the NW-SE faults can be involved in AHe dispersion for this block. To the West, the formation of the Cerdagne basin during the Serravalian (13 Ma) is consistent with a spatial migration of the tectonic activity. Normal activity of the NW-SE faults and NE-SW Têt fault is possible under NNE-SSW extension. d) Plio-Quaternary period (5-0 Ma) is marked by a N-S fault activation (Capçir f.) and E-W direction of extension (Calvet, 1999; Rigo et al., 2015).

This Lower-Miocene change in direction of extension could be related to geodynamic processes implying stress changes at the Mediterranean domain scale. Romagny et al. (2020) proposed a global change in the main direction of slab retreat at about 20 Ma, with a change in the direction of retreat from NNW-SSE to mostly E-W towards the Apennines. Although at far distance from our study area and not clearly kinematically consistent, such process involving mantle flux perturbations may have implied stress changes at far distances in the Pyrenean lithosphere. Another potential source of stress perturbation could be the mechanical interaction and linkage (e.g. Crider and Pollard, 1998; Kattenhorn et al., 2000) between the Cevennes and the Catalan lithospheric normal faults, through a very large-scale relay zone located in the Eastern Pyrenees. Such large-scale mechanical interaction could have favoured stress changes and strain distribution along multiple faults in this eastern part of the Pyrenees. Linkage had to develop with new NW-SE relay faults after the growth of the two NE-SW Cevennes and Penedes master faults in the Oligocene – Lower Miocene (e.g. Seranne et al., 1999), consistently with the timing and direction of the Upper Miocene NW-SE faults observed in the study area. Also note that both master fault segmentation at the place of the pre-existing Pyrenees and the timing of linkage are consistent with the margin development in the Roussillon and its specific orientation (NNW-SSE) in the Gulf of Lion (Mauffret et al., 2001). Finally, another hypothesis to consider is the presence of an new extensional phase due to a not well known geodynamic process in the area (e.g. stresses due to wedge collapse, erosion, or new mantle dynamic, etc.) in a larger domain since a similar cooling event has been recorded in the western Axial Zone (Fillon et al., 2021).

During the Plio-quaternary period (Fig. 8d), seismic data inversion highlight a global N-S contraction in the area, while we can note E-W extension in the Cerdagne basin (Rigo et al., 2015). This E-W extension can be responsible for the Capçir N-S normal faulting (Calvet, 1999; Baize et al., 2002), kinematically consistent with a recent return to N-S Pyrenean contraction in the study area.

6 Conclusions

Low-temperature thermochronology and inverse thermal modeling reveal successive cooling periods associated to the differential exhumation of crustal blocks along the southern Têt fault. In the hanging wall of the Têt fault, low-T thermochronological data indicate a significant exhumation/cooling period ($\sim 30^{\circ}\text{C}/\text{Ma}$) between 38 and 35 Ma, followed by an important decrease in exhumation/cooling ($< 1^{\circ}\text{C}/\text{Ma}$). This slowdown is interpreted as the result of the last Pyrenean contractional stage during the Priabonian. In the Têt fault footwall, we propose that an early exhumation stage of the Canigou-Costabonne block is recorded until 33 Ma ($\sim 30^{\circ}\text{C}/\text{Ma}$) but not in the Carança block (further West), in association to the normal activity of both the Têt and Py faults. These results suggest a rapid switch between contractional and extensional regime in the Eastern Pyrénées during the Priabonian. A second major cooling event ($\sim 20^{\circ}\text{C}/\text{Ma}$) between the Upper Oligocene and Lower Miocene (26-19 Ma) is recorded both in the Canigou and Carança massifs, associated to the major period of activity of the Têt fault linked to the opening of the Gulf of Lion. During the upper Miocene, low-T thermochronological data from solely the Carança massif suggest a third cooling event ($\sim 10^{\circ}\text{C}/\text{Ma}$) during the Serravalian-Tortonian (12-9 Ma) and its segmentation in different sub-blocks separated by NW-SE faults. Our results reveal a progressive propagation of the deformation towards the Southwest along the Têt fault, and also account for major changes in the direction of extension in the Eastern Pyrenees since the Priabonian.

Acknowledgments, Samples, and Data

Competing interests. The authors declare that they have no conflict of interest.

Data availability. All the data used in this study can be obtained in the figures, references, and supporting information. The data are deposited in Geochron database (<https://www.geochron.org>).

Acknowledgements. Thanks for technical support go to Olivier Bruguier, Lucie Koeller and Léa Causse for ICP-MS analyses, and Cyprien Astoury for apatite and zircon separation. Thanks to V. Combaudon and G. de Montbrun for help during Canigou sampling and apatite preparation for dating and Frédéric Mouthereau and Riccardo Asti for their positive remarks and improvements. P.G.V. acknowledges funding support from the Swiss National Science Foundation SNSF (Grant PP00P2_170559) and the French ANR-PIA program (ANR-18-MPGA-0006). This research received funding from the programme TelluS of the Institut National des Sciences de l'Univers, CNRS.

Financial support. This research has been supported by the THERMOFAULT, a project supported by the Region Occitanie (France) involving TLS Geothermics (main sponsor), Géosciences Montpellier, and the TelluS Program of CNRS/INSU.

References

- Agustí, J., Oms, O., Furió, M., Pérez-Vila, M.-J., & Roca, E. (2006). The Messinian terrestrial record in the Pyrenees: The case of Can Vilella (Cerdanya Basin). *Palaeogeography, Palaeoclimatology, Palaeoecology*, 238(1–4), 179–189. <https://doi.org/10.1016/j.palaeo.2006.03.024>
- Angrand, P., & Mouthereau, F. (2021). Evolution of the Alpine orogenic belts in the Western Mediterranean region as resolved by the kinematics of the Europe-Africa diffuse plate boundary. *BSGF - Earth Sciences Bulletin*, 192, 42. <https://doi.org/10.1051/bsgf/2021031>
- Armstrong, P. A., Ehlers, T. A., Chapman, D. S., Farley, K. A., & Kamp, P. J. J. (2003). Exhumation of the central Wasatch Mountains, Utah: 1. Patterns and timing of exhumation deduced from low-temperature thermochronology data. *Journal of Geophysical Research: Solid Earth*, 108(B3). <https://doi.org/10.1029/2001JB001708>
- Ault, A. K., Gautheron, C., & King, G. E. (2019). Innovations in (U–Th)/He, Fission Track, and Trapped Charge Thermochronometry with Applications to Earthquakes, Weathering, Surface-Mantle Connections, and the Growth and Decay of Mountains. *Tectonics*, 38(11), 3705–3739. <https://doi.org/10.1029/2018TC005312>
- Autran, A., Calvet, M., Delmas, M. (2005). Carte géologique de la France à 1/50 000, feuille mont-louis. BRGM.
- Bartrina, M. T., Cabrera, L., Jurado, M. J., Guimerà, J., & Roca, E. (1992). Evolution of the central Catalan margin of the Valencia trough (western Mediterranean). *Tectonophysics*, 203(1–4), 219–247. [https://doi.org/10.1016/0040-1951\(92\)90225-U](https://doi.org/10.1016/0040-1951(92)90225-U)
- Beaumont, C., Muñoz, J. A., Hamilton, J., & Fullsack, P. (2000). Factors controlling the Alpine evolution of the central Pyrenees inferred from a comparison of observations and geodynamical models. *Journal of Geophysical Research: Solid Earth*, 105(B4), 8121–8145. <https://doi.org/10.1029/1999JB900390>
- Bosch, G. V., Teixell, A., Jolivet, M., Labaume, P., Stockli, D., Domènech, M., & Monié, P. (2016a). Timing of Eocene–Miocene thrust activity in the Western Axial Zone and Chaînons Béarnais (west-central Pyrenees) revealed by multi-method thermochronology. *Comptes Rendus Geoscience*, 348(3–4), 246–256. <https://doi.org/10.1016/j.crte.2016.01.001>
- Bouchez, J. L., & Gleizes, G. (1995). Two-stage deformation of the Mont-Louis-Andorra granite pluton (Variscan Pyrenees) inferred from magnetic susceptibility anisotropy. *Journal of the Geological Society*, 152(4), 669–679. <https://doi.org/10.1144/gsjgs.152.4.0669>
- Braun, J. (2002). Quantifying the effect of recent relief changes on age–elevation relationships. *Earth and Planetary Science Letters*, 200(3–4), 331–343. [https://doi.org/10.1016/S0012-821X\(02\)00638-6](https://doi.org/10.1016/S0012-821X(02)00638-6)
- Briaies, A., Armijo, R., Winter, T., Tapponnier, P., Herbecq, A. (1990). Morphological evidence for quaternary normal faulting and seismic hazard in the eastern pyrenees. *Ann. Tecton.* 4, 19–42.
- Brichau, S., Ring, U., Ketcham, R. A., Carter, A., Stockli, D., & Brunel, M. (2006). Constraining the long-term evolution of the slip rate for a major extensional fault system in the central Aegean, Greece, using thermochronology. *Earth and Planetary Science Letters*, 241(1–2), 293–306. <https://doi.org/10.1016/j.epsl.2005.09.065>
- Burbank, D. W., Puigdefàbregas, C., & Munoz, J. A. (1992). The chronology of the Eocene tectonic and stratigraphic development of the eastern Pyrenean foreland basin, northeast Spain. *Geological Society of America Bulletin*, 20.
- Cabrera, L., Roca, E., & Santanach, P. (1988). Basin formation at the end of a strike-slip fault: the Cerdanya Basin (eastern Pyrenees). *Journal of the Geological Society*, 145(2), 261–268. <https://doi.org/10.1144/gsjgs.145.2.0261>
- Calvet, M. (1996). Morphogenèse d’une montagne méditerranéenne, les pyrénées orientales (vol. 1, 1170 p). University of Paris, Paris.

- Calvet, M. (1999). Régimes des contraintes et volumes de relief dans l'est des Pyrénées/Stress regimes and volumes of reliefs in the Eastern Pyrenees. *Géomorphologie : relief, processus, environnement*, 5(3), 253–278. <https://doi.org/10.3406/morfo.1999.991>
- Calvet, M., Gunnell, Y., & Delmas, M. (2014). The Têt River Valley: A Condensed Record of Long-Term Landscape Evolution in the Pyrenees. In M. Fort & M.-F. André (Eds.), *Landscapes and Landforms of France* (pp. 127–138). Dordrecht: Springer Netherlands. https://doi.org/10.1007/978-94-007-7022-5_13
- Calvet, M., Autran, A., Wiazemsky, M., Laumonier, B., Guitard, G. (2015). Carte géologique de la France, 1:50,000 scale, sheet Argelès-sur-Mer/Cerbère (1097). BRGM, Orléans. Handbook by Laumonier, B., Calvet, M., Barbey, P., Guennoc, P., Lambert J., Lenoble, J.L., Wiazemsky, M., 149 p.
- Calvet, M., Gunnell, Y., & Laumonier, B. (2021). Denudation history and palaeogeography of the Pyrenees and their peripheral basins: an 84-million-year geomorphological perspective. *Earth-Science Reviews*, 215, 103436. <https://doi.org/10.1016/j.earscirev.2020.103436>
- Carmignani, L., & Kligfield, R. (1990). Crustal extension in the northern Apennines: The transition from compression to extension in the Alpi Apuane Core Complex. *Tectonics*, 9(6), 1275–1303. <https://doi.org/10.1029/TC009i006p01275>
- Carozza, J.-M. & Delcaillau, B. (1999). L'enregistrement géomorphologique de la tectonique quaternaire par les nappes alluviales: l'exemple du bassin de la têt (roussillon, france). *Comptes Rendus de l'Académie des Sciences-Series IIA-Earth and Planetary Science* 329 (10), 735–740.
- Carozza, J.-M., & Baize, S. (2004). L'escarpement de faille de la Têt est-il le résultat de la tectonique active Plio-Pléistocène ou d'une exhumation Pléistocène? *Comptes Rendus Geoscience*, 336(3), 217–226. <https://doi.org/10.1016/j.crte.2003.10.026>
- Cederbom, C., Larson, S. Å., Tullborg, E.-L., & Stiberg, J.-P. (2000). Fission track thermochronology applied to Phanerozoic thermotectonic events in central and southern Sweden. *Tectonophysics*, 316(1–2), 153–167. [https://doi.org/10.1016/S0040-1951\(99\)00230-9](https://doi.org/10.1016/S0040-1951(99)00230-9)
- Charvin, K., Gallagher, K., Hampson, G. L., & Labourdette, R. (2009). A Bayesian approach to inverse modelling of stratigraphy, part 1: method. *Basin Research*, 21(1), 5–25. <https://doi.org/10.1111/j.1365-2117.2008.00369.x>
- Chevrot, S., Sylvander, M., Diaz, J., Martin, R., Mouthereau, F., Manatschal, G., et al. (2018). The non-cylindrical crustal architecture of the Pyrenees. *Scientific Reports*, 8(1). <https://doi.org/10.1038/s41598-018-27889-x>
- Choukroune, P. (1989). The Eors Pyrenean deep seismic profile reflection data and the overall structure of an orogenic belt. *Tectonics*, 8(1), 23–39. <https://doi.org/10.1029/TC008i001p00023>
- Clauzon, G., Aguilar, J.-P., Michaux, J. (1987). Le bassin pliocène du Roussillon (Pyrénées orientales, France): exemple d'évolution géodynamique d'une ría méditerranéenne consécutive à la crise de salinité messinienne. *C. R. Acad. Sci. Paris Ser. 2* (304), 585e590.
- Clauzon, G. (1990). Restitution de l'évolution géodynamique néogène du bassin du Roussillon et de l'unité adjacente des Corbières d'après les données écostratigraphiques et paléogéographiques. *Paleobiol. Cont.* 17, 125e155.
- Clauzon, G., Le Strat, P., Duvail, C., Do Couto, D., Suc, J.-P., Molliex, S., et al. (2015). The Roussillon Basin (S. France): A case-study to distinguish local and regional events between 6 and 3 Ma. *Marine and Petroleum Geology*, 66, 18–40. <https://doi.org/10.1016/j.marpetgeo.2015.03.012>
- Cochelin, B., Chardon, D., Denèle, Y., Gumiaux, C., & Le Bayon, B. (2017). Vertical strain partitioning in hot Variscan crust: Syn-convergence escape of the Pyrenees in the Iberian-Armorican syntax. *Bulletin de La Société Géologique de France*, 188(6), 39. <https://doi.org/10.1051/bsgf/2017206>

- Colgan, J. P., Dumitru, T. A., Reiners, P. W., Wooden, J. L., & Miller, E. L. (2006). Cenozoic Tectonic Evolution of the Basin and Range Province in Northwestern Nevada. *American Journal of Science*, 306(8), 616–654. <https://doi.org/10.2475/08.2006.02>
- Colgan, Joseph P., Shuster, D. L., & Reiners, P. W. (2008). Two-phase Neogene extension in the northwestern Basin and Range recorded in a single thermochronology sample. *Geology*, 36(8), 631. <https://doi.org/10.1130/G24897A.1>
- Crider, J. G., & Pollard, D. D. (1998). Fault linkage: Three-dimensional mechanical interaction between echelon normal faults. *Journal of Geophysical Research: Solid Earth*, 103(B10), 24373–24391. <https://doi.org/10.1029/98JB01353>
- Cruset, D., Vergés, J., Albert, R., Gerdes, A., Benedicto, A., Cantarero, I., & Travé, A. (2020). Quantifying deformation processes in the SE Pyrenees using U–Pb dating of fracture-filling calcites. *Journal of the Geological Society*, 177(6), 1186–1196. <https://doi.org/10.1144/jgs2020-014>
- Curry, M. A. E., Barnes, J. B., & Colgan, J. P. (2016). Testing fault growth models with low-temperature thermochronology in the northwest Basin and Range, USA: TESTING FAULT GROWTH MODELS WITH AHE. *Tectonics*, 35(10), 2467–2492. <https://doi.org/10.1002/2016TC004211>
- Curry, M. E., van der Beek, P., Huismans, R. S., Wolf, S. G., & Muñoz, J.-A. (2019). Evolving paleotopography and lithospheric flexure of the Pyrenean Orogen from 3D flexural modeling and basin analysis. *Earth and Planetary Science Letters*, 515, 26–37. <https://doi.org/10.1016/j.epsl.2019.03.009>
- Curry, M. E., van der Beek, P., Huismans, R. S., Wolf, S. G., Fillon, C., & Muñoz, J.-A. (2021). Spatio-temporal patterns of Pyrenean exhumation revealed by inverse thermo-kinematic modeling of a large thermochronologic data set. *Geology*. <https://doi.org/10.1130/G48687.1>
- Danišík, M., Štěpančíková, P., & Evans, N. J. (2012). Constraining long-term denudation and faulting history in intraplate regions by multisystem thermochronology: An example of the Sudetic Marginal Fault (Bohemian Massif, central Europe): THERMOCHRONOLOGY OF THE SUDETES. *Tectonics*, 31(2). <https://doi.org/10.1029/2011TC003012>
- Deeken, A., Sobel, E. R., Coutand, I., Haschke, M., Riller, U., & Strecker, M. R. (2006). Development of the southern Eastern Cordillera, NW Argentina, constrained by apatite fission track thermochronology: From early Cretaceous extension to middle Miocene shortening: EXHUMING THE SOUTHEASTERN ANDES. *Tectonics*, 25(6). <https://doi.org/10.1029/2005TC001894>
- Delcaillau, B., Carozza, J.-M., & Font, M. (2004). The northern segment of the Têt Fault (Pyrenees-Orientales): Neogene evolution and geomorphic implications. *Bulletin de la Société Géologique de France*, 175(3), 257–272. <https://doi.org/10.2113/175.3.257>
- Delmas, M., Calvet, M., Gunnell, Y., Voinchet, P., Manel, C., Braucher, R., et al. (2018). Terrestrial ^{10}Be and electron spin resonance dating of fluvial terraces quantifies quaternary tectonic uplift gradients in the eastern Pyrenees. *Quaternary Science Reviews*, 193, 188–211. <https://doi.org/10.1016/j.quascirev.2018.06.001>
- Diaz, J., Vergés, J., Chevrot, S., Antonio-Vigil, A., Ruiz, M., Sylvander, M., & Gallart, J. (2018). Mapping the crustal structure beneath the eastern Pyrenees. *Tectonophysics*, 744, 296–309. <https://doi.org/10.1016/j.tecto.2018.07.011>
- Ehlers, T. A., & Farley, K. A. (2003). Apatite (U–Th)/He thermochronometry: methods and applications to problems in tectonic and surface processes. *Earth and Planetary Science Letters*, 206(1–2), 1–14. [https://doi.org/10.1016/S0012-821X\(02\)01069-5](https://doi.org/10.1016/S0012-821X(02)01069-5)
- Farley, K. A., Wolf, R. A., and Silver, L. T. (1996). The effects of long alpha-stopping distances on (U–Th)/He ages, *Geochim. Cosmochim. Ac.*, 60, 4223–4229, [https://doi.org/10.1016/S0016-7037\(96\)00193-7](https://doi.org/10.1016/S0016-7037(96)00193-7)

- Farley, K. A. (2002). (U-Th)/He Dating: Techniques, Calibrations, and Applications. *Reviews in Mineralogy and Geochemistry*, 47(1), 819–844. <https://doi.org/10.2138/rmg.2002.47.18>
- Faulkner, D. R., Jackson, C. A. L., Lunn, R. J., Schlische, R. W., Shipton, Z. K., Wibberley, C. A. J., & Withjack, M. O. (2010). A review of recent developments concerning the structure, mechanics and fluid flow properties of fault zones. *Journal of Structural Geology*, 32(11), 1557–1575. <https://doi.org/10.1016/j.jsg.2010.06.009>
- Fillon, C., & van der Beek, P. (2012). Post-orogenic evolution of the southern Pyrenees: constraints from inverse thermo-kinematic modelling of low-temperature thermochronology data. *Basin Research*, 24(4), 418–436. <https://doi.org/10.1111/j.1365-2117.2011.00533.x>
- Fillon, C., Mouthereau, F., Calassou, S., Pik, R., Bellahsen, N., Gautheron, C., ... & van der Beek, P. (2021). Post-orogenic exhumation in the western Pyrenees: evidence for extension driven by pre-orogenic inheritance. *Journal of the Geological Society*, 178(2). <https://doi.org/10.1144/jgs2020-079>
- Fitzgerald, P. G., Sorkhabi, R. B., Redfield, T. F., & Stump, E. (1995). Uplift and denudation of the central Alaska Range: A case study in the use of apatite fission track thermochronology to determine absolute uplift parameters. *Journal of Geophysical Research: Solid Earth*, 100(B10), 20175–20191. <https://doi.org/10.1029/95JB02150>
- Fitzgerald, P. G., Muñoz, J. A., Coney, P. J., & Baldwin, S. L. (1999). Asymmetric exhumation across the Pyrenean orogen: implications for the tectonic evolution of a collisional orogen. *Earth and Planetary Science Letters*, 173(3), 157–170. [https://doi.org/10.1016/S0012-821X\(99\)00225-3](https://doi.org/10.1016/S0012-821X(99)00225-3)
- Fitzgerald, P. G., & Malusà, M. G. (2019). Concept of the Exhumed Partial Annealing (Retention) Zone and Age-Elevation Profiles in Thermochronology. In M. G. Malusà & P. G. Fitzgerald (Eds.), *Fission-Track Thermochronology and its Application to Geology* (pp. 165–189). Cham: Springer International Publishing. https://doi.org/10.1007/978-3-319-89421-8_9
- Foster, D. A., & John, B. E. (1999). Quantifying tectonic exhumation in an extensional orogen with thermochronology: examples from the southern Basin and Range Province. *Geological Society, London, Special Publications*, 154(1), 343–364. <https://doi.org/10.1144/GSL.SP.1999.154.01.16>
- Gallagher, K., Charvin, K., Nielsen, S., Sambridge, M., & Stephenson, J. (2009). Markov chain Monte Carlo (MCMC) sampling methods to determine optimal models, model resolution and model choice for Earth Science problems. *Marine and Petroleum Geology*, 26(4), 525–535. <https://doi.org/10.1016/j.marpetgeo.2009.01.003>
- Gallagher, K. (2012). Transdimensional inverse thermal history modeling for quantitative thermochronology. *Journal of Geophysical Research: Solid Earth*, 117(B2). <https://doi.org/10.1029/2011JB008825>
- Gautheron, C., Tassan-Got, L., Barbarand, J., & Pagel, M. (2009). Effect of alpha-damage annealing on apatite (U-Th)/He thermochronology. *Chemical Geology*, 266(3–4), 157–170. <https://doi.org/10.1016/j.chemgeo.2009.06.001>
- Gautheron, C., & Tassan-Got, L. (2010). A Monte Carlo approach to diffusion applied to noble gas/helium thermochronology. *Chemical Geology*, 273(3–4), 212–224. <https://doi.org/10.1016/j.chemgeo.2010.02.023>
- Gautheron, C., Pinna-Jamme, R., Derycke, A., Ahadi, F., Sanchez, C., Haurine, F., et al. (2021). Analytical protocols and performance for apatite and zircon (U-Th)/He analysis on quadrupole and magnetic sector mass spectrometer systems between 2007 and 2020. *Geochronology*, 3(1), 351–370. <https://doi.org/10.5194/gchron-3-351-2021>
- Gibson, M., Sinclair, H. D., Lynn, G. J., & Stuart, F. M. (2007). Late- to post-orogenic exhumation of the Central Pyrenees revealed through combined thermochronological data and modelling. *Basin Research*, 19(3), 323–334. <https://doi.org/10.1111/j.1365-2117.2007.00333.x>
- Glotsbach, C., van der Beek, P. A., & Spiegel, C. (2011). Episodic exhumation and relief growth in the Mont Blanc massif, Western Alps from numerical modelling of thermochronology data. *Earth and Planetary Science Letters*, 304(3–4), 417–430. <https://doi.org/10.1016/j.epsl.2011.02.020>

- Goula, X., Olivera, C., Fleta, J., Grellet, B., Lindo, R., Rivera, L. A., et al. (1999). Present and recent stress regime in the eastern part of the Pyrenees. *Tectonophysics*, 308(4), 487–502. [https://doi.org/10.1016/S0040-1951\(99\)00120-1](https://doi.org/10.1016/S0040-1951(99)00120-1)
- Green PF, Duddy IR, Laslett GM, Hegarty KA, Gleadow AJW, Lovering JF (1989) Thermal annealing of fission tracks in apatite, 4, quantitative modelling techniques and extension to geological timescale. *J Pet Geol* 12:111–114. [https://doi.org/10.1016/0168-9622\(89\)90018-3](https://doi.org/10.1016/0168-9622(89)90018-3)
- Green, P. F., Crowhurst, P. V., Duddy, I. R., Japsen, P., & Holford, S. P. (2006). Conflicting (U–Th)/He and fission track ages in apatite: Enhanced He retention, not anomalous annealing behaviour. *Earth and Planetary Science Letters*, 250(3–4), 407–427. <https://doi.org/10.1016/j.epsl.2006.08.022>
- Grool, A. R., Ford, M., Vergés, J., Huisman, R. S., Christophoul, F., & Dielforder, A. (2018). Insights Into the Crustal-Scale Dynamics of a Doubly Vergent Orogen From a Quantitative Analysis of Its Forelands: A Case Study of the Eastern Pyrenees. *Tectonics*, 37(2), 450–476. <https://doi.org/10.1002/2017TC004731>
- Guenther, W. R., Reiners, P. W., Ketcham, R. A., Nasdala, L., & Giester, G. (2013). Helium diffusion in natural zircon: Radiation damage, anisotropy, and the interpretation of zircon (U–Th)/He thermochronology. *American Journal of Science*, 313(3), 145–198. <https://doi.org/10.2475/03.2013.01>
- Guitard, G., Geyssant, J., Laumonier, B., Autran, A., Fontelles, M., Dalmayrac, B., Vidal, J., Bandet, Y. (1992). Carte géologique de la France (1/50 000), feuille prades (1095).
- Guitard, G., Laumonier, B., Autran, A., Bandet, Y., Berger, G. (1998). Notice explicative, carte géologique France (1/50.000), feuille prades (1095). BRGM, Orléans.
- Gunnell, Y., Calvet, M., Brichau, S., Carter, A., Aguilar, J.-P., & Zeyen, H. (2009). Low long-term erosion rates in high-energy mountain belts: Insights from thermo- and biochronology in the Eastern Pyrenees. *Earth and Planetary Science Letters*, 278(3–4), 208–218. <https://doi.org/10.1016/j.epsl.2008.12.004>
- Hoareau, G., Crognier, N., Lacroix, B., Aubourg, C., Roberts, N. M. W., Niemi, N., et al. (2021). Combination of $\Delta 47$ and U–Pb dating in tectonic calcite veins unravel the last pulses related to the Pyrenean Shortening (Spain). *Earth and Planetary Science Letters*, 553, 116636. <https://doi.org/10.1016/j.epsl.2020.116636>
- Hurford, A. J. (1990). Standardization of fission track dating calibration: Recommendation by the Fission Track Working Group of the I.U.G.S. Subcommittee on Geochronology. *Chemical Geology: Isotope Geoscience Section*, 80(2), 171–178. [https://doi.org/10.1016/0168-9622\(90\)90025-8](https://doi.org/10.1016/0168-9622(90)90025-8)
- Huyghe, D., Mouthereau, F., Ségalen, L., & Furió, M. (2020). Long-term dynamic topographic support during post-orogenic crustal thinning revealed by stable isotope ($\delta^{18}\text{O}$) paleo-altimetry in eastern Pyrenees. *Scientific Reports*, 10(1), 2267. <https://doi.org/10.1038/s41598-020-58903-w>
- Jolivet, L., & Faccenna, C. (2000). Mediterranean extension and the Africa-Eurasia collision. *Tectonics*, 19(6), 1095–1106. <https://doi.org/10.1029/2000TC900018>
- Jolivet, L., Gorini, C., Smit, J., & Leroy, S. (2015). Continental breakup and the dynamics of rifting in back-arc basins: The Gulf of Lion margin. *Tectonics*, 34(4), 662–679. <https://doi.org/10.1002/2014TC003570>
- Jolivet, L., Romagny, A., Gorini, C., Maillard, A., Thinon, I., Couëffé, R., et al. (2020). Fast dismantling of a mountain belt by mantle flow: Late-orogenic evolution of Pyrenees and Liguro-Provençal rifting. *Tectonophysics*, 776, 228312. <https://doi.org/10.1016/j.tecto.2019.228312>
- Jolivet, L., Menant, A., Roche, V., Le Pourhiet, L., Maillard, A., Augier, R., et al. (2021a). Transfer zones in Mediterranean back-arc regions and tear faults. *BSGF - Earth Sciences Bulletin*, 192, 11. <https://doi.org/10.1051/bsgf/2021006>

- Jolivet, L., Baudin, T., Calassou, S., Chevrot, S., Ford, M., Issautier, B., et al. (2021b). Geodynamic evolution of a wide plate boundary in the Western Mediterranean, near-field versus far-field interactions. *BSGF - Earth Sciences Bulletin*, 192, 48. <https://doi.org/10.1051/bsgf/2021043>
- Jolivet, M., Labaume, P., Monié, P., Brunel, M., Arnaud, N., & Campani, M. (2007). Thermochronology constraints for the propagation sequence of the south Pyrenean basement thrust system (France-Spain): PROPAGATION OF THE SOUTH PYRENEAN PRISM. *Tectonics*, 26(5). <https://doi.org/10.1029/2006TC002080>
- Jones, C. H., & Wesnousky, S. G. (1992). Variations in Strength and Slip Rate Along the San Andreas Fault System. *Science*, 256(5053), 83–86. <https://doi.org/10.1126/science.256.5053.83>
- Kattenhorn, S. A., Aydin, A., & Pollard, D. D. (2000). Joints at high angles to normal fault strike: an explanation using 3-D numerical models of fault-perturbed stress fields p. *Journal of Structural Geology*, 22(1), 1-23. [https://doi.org/10.1016/S0191-8141\(99\)00130-3](https://doi.org/10.1016/S0191-8141(99)00130-3)
- Ketcham, Richard A., Donelick, R. A., & Carlson, W. D. (1999). Variability of apatite fission-track annealing kinetics; III, Extrapolation to geological time scales. *American Mineralogist*, 84(9), 1235–1255. <https://doi.org/10.2138/am-1999-0903>
- Ketcham, R. A., Carter, A., Donelick, R. A., Barbarand, J., & Hurford, A. J. (2007). Improved modeling of fission-track annealing in apatite. *American Mineralogist*, 92(5–6), 799–810. <https://doi.org/10.2138/am.2007.2281>
- Ketcham, R. A., Gautheron, C., and Tassan-Got, L. (2011). Accounting for long alpha-particle stopping distances in (U-Th-Sm)/He geochronology: Refinement of the baseline case, *Geochim. Cosmochim. Ac.*, 75, 7779–7791, <https://doi.org/10.1016/j.gca.2011.10.011>
- Krugh, W. C. (2008). Low-temperature thermochronologic constraints on fault array evolution and patterns of range-scale denudation. *ETH Zurich*. <https://doi.org/10.3929/ETHZ-A-005772437>
- Labaume, P., Meresse, F., Jolivet, M., Teixell, A., & Lahfid, A. (2016). Tectonothermal history of an exhumed thrust-sheet-top basin: An example from the south Pyrenean thrust belt: JACA THRUST-SHEET-TOP BASIN. *Tectonics*, 35(5), 1280–1313. <https://doi.org/10.1002/2016TC004192>
- Lacan, P., & Ortuño, M. (2012). Active Tectonics of the Pyrenees: A review. *Journal of Iberian Geology*, 38(1). https://doi.org/10.5209/rev_JIGE.2012.v38.n1.39203
- Lacombe, O., & Jolivet, L. (2005). Structural and kinematic relationships between Corsica and the Pyrenees-Provence domain at the time of the Pyrenean orogeny: CORSICA AND PROVENCE IN THE LATE EOCENE. *Tectonics*, 24(1). <https://doi.org/10.1029/2004TC001673>
- Laumonier, B., Calvet, M., Le Bayon, B., Barbey, P., Lenoble, J.-L. (2015). avec la collaboration de Cocherie A., Melleton J. – Notice explicative, Carte géologique de la France à 1/50 000, feuille Prats-de-Mollo-la-Preste (1099). In: Orléans : BRGM, page 189.
- Laumonier, B., Calvet, M., Delmas, M., Barbey, P., Lenoble, J., Autran, A. (2017). Notice explicative Carte géologique de la France `a 1/50 000, feuille Mont-Louis (1094). In: Orléans : BRGM, page 139.
- Lewis, C. J., Vergés, J., & Marzo, M. (2000). High mountains in a zone of extended crust: Insights into the Neogene-Quaternary topographic development of northeastern Iberia. *Tectonics*, 19(1), 86–102. <https://doi.org/10.1029/1999TC900056>
- Malusà, M. G., Polino, R., Zattin, M., Bigazzi, G., Martin, S., & Piana, F. (2005). Miocene to Present differential exhumation in the Western Alps: Insights from fission track thermochronology. *Tectonics*, 24(3). <https://doi.org/10.1029/2004TC001782>

- Malusà, M. G., Polino, R., & Zattin, M. (2009). Strain partitioning in the axial NW Alps since the Oligocene: STRAIN PARTITIONING IN THE AXIAL ALPS. *Tectonics*, 28(3). <https://doi.org/10.1029/2008TC002370>
- Martín-González, F., Barbero, L., Capote, R., Heredia, N., & Gallastegui, G. (2012). Interaction of two successive Alpine deformation fronts: constraints from low-temperature thermochronology and structural mapping (NW Iberian Peninsula). *International Journal of Earth Sciences*, 101(5), 1331–1342. <https://doi.org/10.1007/s00531-011-0712-9>
- Mauffret, A., Durand de Grossouvre, B., Tadeu Dos Reis, A., Gorini, C., & Nercessian, A. (2001). Structural geometry in the eastern Pyrenees and western Gulf of Lion (Western Mediterranean). *Journal of Structural Geology*, 23(11), 1701–1726. [https://doi.org/10.1016/S0191-8141\(01\)00025-6](https://doi.org/10.1016/S0191-8141(01)00025-6)
- Maurel, O., Brunel, M., & Monié, P. (2002). Exhumation cénozoïque des massifs du Canigou et de Mont-Louis (Pyrénées orientales, France). *Comptes Rendus Geoscience*, 334(12), 941–948. [https://doi.org/10.1016/S1631-0713\(02\)01834-5](https://doi.org/10.1016/S1631-0713(02)01834-5)
- Maurel, O., Monié, P., Pik, R., Arnaud, N., Brunel, M., & Jolivet, M. (2008). The Meso-Cenozoic thermo-tectonic evolution of the Eastern Pyrenees: an $40\text{Ar}/39\text{Ar}$ fission track and (U–Th)/He thermochronological study of the Canigou and Mont-Louis massifs. *International Journal of Earth Sciences*, 97(3), 565–584. <https://doi.org/10.1007/s00531-007-0179-x>
- McDannell, K. T., Toro, J., Hourigan, J. K., & Harris, D. (2014). Thermochronologic constraints on Late Cretaceous to Cenozoic exhumation of the Bendeleben Mountains, Seward Peninsula, Alaska. *Geochemistry, Geophysics, Geosystems*, 15(10), 4009–4023. <https://doi.org/10.1002/2014GC005424>
- McCaig, A. M., & Miller, J. A. (1986). 40Ar - 39Ar age of mylonites along the Merens Fault, central Pyrenees. *Tectonophysics*, 129(1–4), 149–172. [https://doi.org/10.1016/0040-1951\(86\)90250-7](https://doi.org/10.1016/0040-1951(86)90250-7)
- McDowell, F. W., McIntosh, W. C., and Farley, K. A. (2005). A precise 40Ar – 39Ar reference age for the Durango apatite (U–Th)/He and fission-track dating standard, *Chem. Geol.*, 214, 249–263. <https://doi.org/10.1016/j.chemgeo.2004.10.002>
- Medialdea, T., Vázquez, J., Vegas, R. (1994). Estructura y evolución geodinámica del extremo noreste del margen continental catalán durante el neógeno. *Acta Geol. Hisp.* 29 (2–4), 39–53.
- Metcalf, J. R., Fitzgerald, P. G., Baldwin, S. L., & Muñoz, J.-A. (2009). Thermochronology of a convergent orogen: Constraints on the timing of thrust faulting and subsequent exhumation of the Maladeta Pluton in the Central Pyrenean Axial Zone. *Earth and Planetary Science Letters*, 287(3–4), 488–503. <https://doi.org/10.1016/j.epsl.2009.08.036>
- Milesi, G., Soliva, R., Monié, P., Münch, P., Bellanger, M., Bruguier, O., et al. (2019a). Mapping a geothermal anomaly using apatite (U–Th)/He thermochronology in the Têt fault damage zone, eastern Pyrenees, France. *Terra Nova*, 31(6), 569–576. <https://doi.org/10.1111/ter.12429>
- Milesi, G. (2020). Analyse thermochronologique, géochimique et structurale du système hydrothermal de la faille de la Têt (Pyrénées, France), un nouvel outil d'exploration géothermique. PhD thesis. Université de Montpellier.
- Milesi, G., Monié, P., Soliva, R., Münch, P., Taillefer, A., Bellanger, M., et al. (2020a). Imaging Geothermal Anomalies Using Low-temperature (U–Th)/He Thermochronometry: A Case Study from the Active Têt Fault Hydrothermal System (Eastern Pyrenees, France). In *Proceedings World Geothermal Congress 2020 Reykjavik, Iceland, April 26–May 2, 2020*.
- Milesi, G., Monié, P., Münch, P., Soliva, R., Taillefer, A., Bruguier, O., et al. (2020b). Tracking geothermal anomalies along a crustal fault using (U – Th)/He apatite thermochronology and rare-earth element (REE) analyses: the example of the Têt fault (Pyrenees, France). *Solid Earth*, 11(5), 1747–1771. <https://doi.org/10.5194/se-11-1747-2020>
- Morris, R.G., Sinclair, H.D. and Yelland, A.J. (1998). Exhumation of the Pyrenean orogen: implications for sediment discharge. *Basin Research*, 10, 69–86, <https://doi.org/10.1046/j.1365-2117.1998.00053.x>

- Mouthereau, F., Filleaudeau, P.-Y., Vacherat, A., Pik, R., Lacombe, O., Fellin, M. G., et al. (2014). Placing limits to shortening evolution in the Pyrenees: Role of margin architecture and implications for the Iberia/Europe convergence: Plate convergence in the Pyrenees. *Tectonics*, 33(12), 2283–2314. <https://doi.org/10.1002/2014TC003663>
- Mouthereau, F., Angrand, P., Jourdon, A., Ternois, S., Fillon, C., Calassou, S., et al. (2021). Cenozoic mountain building and topographic evolution in Western Europe: impact of billions of years of lithosphere evolution and plate kinematics. *BSGF - Earth Sciences Bulletin*, 192, 56. <https://doi.org/10.1051/bsgf/2021040>
- Muñoz, J. A. (1992). Evolution of a continental collision belt: ECORS-Pyrenees crustal balanced cross-section. In K. R. McClay (Ed.), *Thrust Tectonics* (pp. 235–246). Dordrecht: Springer Netherlands. https://doi.org/10.1007/978-94-011-3066-0_21
- Nercessian, A., Mau, A., Gallart, J., & Diaz, J. (2001). Deep reflection seismic images of the crustal thinning in the eastern Pyrenees and western Gulf of Lion. *Journal of Geodynamics*, 15. [https://doi.org/10.1016/S0264-3707\(00\)00029-6](https://doi.org/10.1016/S0264-3707(00)00029-6)
- Norris, R. J., & Cooper, A. F. (2001). Late Quaternary slip rates and slip partitioning on the Alpine Fault, New Zealand. *Journal of Structural Geology*, 23(2–3), 507–520. [https://doi.org/10.1016/S0191-8141\(00\)00122-X](https://doi.org/10.1016/S0191-8141(00)00122-X)
- Parizot, O., Missenard, Y., Haurine, F., Blaise, T., Barbarand, J., Benedicto, A., & Sarda, P. (2021). When did the Pyrenean shortening end? Insight from U–Pb geochronology of syn-faulting calcite (Corbières area, France). *Terra Nova*, 33(6), 551–559. <https://doi.org/10.1111/ter.12547>
- Petit, C., & Mouthereau, F. (2012). Steep topographic slope preservation by anisotropic diffusion: An example from the Neogene Têt fault scarp, eastern Pyrenees. *Geomorphology*, 171–172, 173–179. <https://doi.org/10.1016/j.geomorph.2012.05.016>
- Pous, J., Julià, R., Sugrañes, L.S. (1986). Cerdanya basin geometry and its implication on the neocene evolution of the eastern pyrenees. *Tectonophysics* 129 (1–4), 355–365. [https://doi.org/10.1016/0040-1951\(86\)90261-1](https://doi.org/10.1016/0040-1951(86)90261-1)
- Price, J. G., & Henry, C. D. (1984). Stress orientations during Oligocene volcanism in Trans-Pecos Texas: Timing the transition from Laramide compression to Basin and Range tension. *Geology*, 12(4), 238–241. [https://doi.org/10.1130/0091-7613\(1984\)12<238:SODOVI>2.0.CO;2](https://doi.org/10.1130/0091-7613(1984)12<238:SODOVI>2.0.CO;2)
- Ratschbacher, L., Hacker, B. R., Calvert, A., Webb, L. E., Grimmer, J. C., McWilliams, M. O., et al. (2003). Tectonics of the Qinling (Central China): tectonostratigraphy, geochronology, and deformation history. *Tectonophysics*, 366(1–2), 1–53. [https://doi.org/10.1016/S0040-1951\(03\)00053-2](https://doi.org/10.1016/S0040-1951(03)00053-2)
- Rehault, J., Moussat, E., Fabbri, A. (1987). Structural evolution of the tyrrhenian back-arc basin. *Mar. Geol.* 74 (1-2), 123–150. [https://doi.org/10.1016/0025-3227\(87\)90010-7](https://doi.org/10.1016/0025-3227(87)90010-7)
- Reiners, P.W., Farley, K.A., and Hiskes, H.J. (2002). He diffusion and (U-Th)/He thermochronology of zircon: initial results from Fish Canyon Tuff and Gold Butte. *Tectonophysics*, 349: 297–308, [https://doi.org/10.1016/S0040-1951\(02\)00058-6](https://doi.org/10.1016/S0040-1951(02)00058-6)
- Reiners, P. W., & Brandon, M. T. (2006). USING THERMOCHRONOLOGY TO UNDERSTAND OROGENIC EROSION. *Annual Review of Earth and Planetary Sciences*, 34(1), 419–466. <https://doi.org/10.1146/annurev.earth.34.031405.125202>
- Reiners, P. W. (2007). Thermochronologic Approaches to Paleotopography. *Reviews in Mineralogy and Geochemistry*, 66(1), 243–267. <https://doi.org/10.2138/rmg.2007.66.10>
- Reiners, P. W., and S. Nicolescu (2006), Measurement of parent nuclides for (U-Th)/He chronometry by solution sector ICP-MS, ARHDL Rep., 1, Ariz. Radiogenic Helium Dating Lab., Tucson. Available at https://www.geo.arizona.edu/~reiners/arhdl/U-Th-Sm_measurement_Reiners_042107.pdf

- Rigo, A., Vernant, P., Feigl, K. L., Goula, X., Khazaradze, G., Talaya, J., et al. (2015). Present-day deformation of the Pyrenees revealed by GPS surveying and earthquake focal mechanisms until 2011. *Geophysical Journal International*, 201(2), 947–964. <https://doi.org/10.1093/gji/ggv052>
- Roca, E. & Santanach, P. (1986). Génesis y evolución de la Fosa de la Cerdanya (Pirineos orientales). *Geogaceta* 1, 37–38.
- Roca, E. & Desegaulx, P. (1992). Analysis of the geological evolution and vertical movements in the valencia trough area, western mediterranean. *Marine and Petroleum Geology*, 9(2) :167–176.
- Roca, E. (1996). The Neogene Cerdanya and Seu d'Urgell intramontane basins (eastern Pyrenees). In: Friend, P., Bario, C.J. (Eds.), *Tertiary Basins of Spain. The Stratigraphic Record of Crustal Kinematics*. Cambridge University Press, Cambridge, pp. 114–118.
- Romagny, A., Jolivet, L., Menant, A., Bessière, E., Maillard, A., Canva, A., et al. (2020). Detailed tectonic reconstructions of the Western Mediterranean region for the last 35 Ma, insights on driving mechanisms. *BSGF - Earth Sciences Bulletin*, 191, 37. <https://doi.org/10.1051/bsgf/2020040>
- Roure, F., Choukroune, P., Berastegui, X., Munoz, J. A., Villien, A., Matheron, P., et al. (1989). Eors deep seismic data and balanced cross sections: Geometric constraints on the evolution of the Pyrenees. *Tectonics*, 8(1), 41–50. <https://doi.org/10.1029/TC008i001p00041>
- Rushlow, C. R., Barnes, J. B., Ehlers, T. A., & Vergés, J. (2013). Exhumation of the southern Pyrenean fold-thrust belt (Spain) from orogenic growth to decay: EXHUMATION OF THE SOUTHERN PYRENEES. *Tectonics*, 32(4), 843–860. <https://doi.org/10.1002/tect.20030>
- Sambridge, M. (1999). Geophysical inversion with a neighbourhood algorithm - II. Appraising the ensemble. *Geophysical Journal International*, 138, 727–746. <https://doi.org/10.1046/j.1365-246x.1999.00900.x>
- Sartégou, A., Bourlès, D. L., Blard, P.-H., Braucher, R., Tibari, B., Zimmermann, L., et al. (2018). Deciphering landscape evolution with karstic networks: A Pyrenean case study. *Quaternary Geochronology*, 43, 12–29. <https://doi.org/10.1016/j.quageo.2017.09.005>
- Saula, E., Picart, J., Mató, E., Llenas, M., Losantos, M., Berástegui, X., & Agustí, J. (1994). Evolución geodinámica de la fosa del Empordà y las Sierras Transversales. *Acta geológica hispánica*, 29(2-4), 55–75.
- Schmitz, M.D. & Bowring, S.A. (2001). U-Pb zircon and titanite systematics of the Fish Canyon Tuff: an assessment of high precision U-Pb geochronology and its application to young volcanic rocks. *Geochim. Cosmochim. Acta*, 65: 2571–2587, [https://doi.org/10.1016/S0016-7037\(01\)00616-0](https://doi.org/10.1016/S0016-7037(01)00616-0)
- Schwarz, G. (1978). Estimating the dimension of a model. *The annals of statistics*, 6(2), 461–464.
- Séranne, M., Benedicto, A., Labaume, P., Truffert, C., & Pascal, G. (1995). Structural style and evolution of the Gulf of Lion Oligo-Miocene rifting: Role of the Pyrenean orogeny. *Marine and Petroleum geology*, 12(8), 809–820. [https://doi.org/10.1016/0264-8172\(95\)98849-Z](https://doi.org/10.1016/0264-8172(95)98849-Z)
- Séranne, M. (1999). The Gulf of Lion continental margin (NW Mediterranean) revisited by IBS: an overview. *Geological Society, London, Special Publications*, 156(1), 15–36. <https://doi.org/10.1144/GSL.SP.1999.156.01.03>
- Séranne, M., Couëffé, R., Husson, E., Baral, C., & Villard, J. (2021). The transition from Pyrenean shortening to Gulf of Lion rifting in Languedoc (South France)—A tectonic-sedimentation analysis. *BSGF-Earth Sciences Bulletin*, 192(1), 27. <https://doi.org/10.1051/bsgf/2021017>
- Sibuet, J.-C., Srivastava, S. P., & Spakman, W. (2004). Pyrenean orogeny and plate kinematics. *Journal of Geophysical Research: Solid Earth*, 109(B8). <https://doi.org/10.1029/2003JB002514>

- Sinclair, H. D., Gibson, M., Naylor, M., & Morris, R. G. (2005). Asymmetric growth of the Pyrenees revealed through measurement and modeling of orogenic fluxes. *American Journal of Science*, 305(5), 369–406. <https://doi.org/10.2475/ajs.305.5.369>
- Stockli, D. F., Farley, K. A., & Dumitru, T. A. (2000). Calibration of the apatite (U-Th)/He thermochronometer on an exhumed fault block, White Mountains, California. *Geology*, 28(11), 983–986. [https://doi.org/10.1130/0091-7613\(2000\)28<983:COTAHT>2.0.CO;2](https://doi.org/10.1130/0091-7613(2000)28<983:COTAHT>2.0.CO;2)
- Stockli, D. F. (2005). Application of Low-Temperature Thermochronometry to Extensional Tectonic Settings. *Reviews in Mineralogy and Geochemistry*, 58(1), 411–448. <https://doi.org/10.2138/rmg.2005.58.16>
- Stüwe, K., White, L., & Brown, R. (1994). The influence of eroding topography on steady-state isotherms. Application to fission track analysis. *Earth and Planetary Science Letters*, 124(1–4), 63–74. [https://doi.org/10.1016/0012-821X\(94\)00068-9](https://doi.org/10.1016/0012-821X(94)00068-9)
- Surpless, B. E., Stockli, D. F., Dumitru, T. A., & Miller, E. L. (2002). Two-phase westward encroachment of Basin and Range extension into the northern Sierra Nevada: BASIN AND RANGE ENCROACHMENT. *Tectonics*, 21(1), 2-1-2–10. <https://doi.org/10.1029/2000TC001257>
- Taillefer, A., Soliva, R., Guillou-Frottier, L., Le Goff, E., Martin, G., & Séranne, M. (2017). Fault-Related Controls on Upward Hydrothermal Flow: An Integrated Geological Study of the Têt Fault System, Eastern Pyrénées (France). *Geofluids*, 2017, 1–19. <https://doi.org/10.1155/2017/8190109>
- Taillefer, A., Milesi, G., Soliva, R., Monnier, L., Delorme, P., Guillou-Frottier, L., & Le Goff, E. (2021). Polyphased brittle deformation around a crustal fault: A multi-scale approach based on remote sensing and field data on the mountains surrounding the Têt hydrothermal system (Eastern Pyrénées, France). *Tectonophysics*, 804, 228710. <https://doi.org/10.1016/j.tecto.2020.228710>
- Tassone, A., Roca, E., Muñoz, J. A., Cabrera, L., & Canals, M. (1994). Evolución del sector septentrional del margen continental catalán durante el Cenozoico. *Acta geológica hispánica*, 29(2-4), 3-37.
- Teixell, A., Labaume, P., & Lagabrielle, Y. (2016). The crustal evolution of the west-central Pyrenees revisited: Inferences from a new kinematic scenario. *Comptes Rendus Geoscience*, 348(3–4), 257–267. <https://doi.org/10.1016/j.crte.2015.10.010>
- Ternois, S., Odlum, M., Ford, M., Pik, R., Stockli, D., Tibari, B., et al. (2019). Thermochronological Evidence of Early Orogenesis, Eastern Pyrenees, France. *Tectonics*, 38(4), 1308–1336. <https://doi.org/10.1029/2018TC005254>
- Tosal, A., Verduzco, O., & Martín-Closas, C. (2021). CLAMP-based palaeoclimatic analysis of the late Miocene (Tortonian) flora from La Cerdanya Basin of Catalonia, Spain, and an estimation of the palaeoaltitude of the eastern Pyrenees. *Palaeogeography, Palaeoclimatology, Palaeoecology*, 564, 110186. <https://doi.org/10.1016/j.palaeo.2020.110186>
- Vacherat, A., Mouthereau, F., Pik, R., Huyghe, D., Paquette, J.-L., Christophoul, F., et al. (2016). Rift-to-collision sediment routing in the Pyrenees: A synthesis from sedimentological, geochronological and kinematic constraints. *Earth-Science Reviews*, 172, 43–74. <https://doi.org/10.1016/j.earscirev.2017.07.004>
- Vergés, J., Millán, H., Roca, E., Muñoz, J. A., Marzo, M., Cirés, J., et al. (1995). Eastern Pyrenees and related foreland basins: pre-, syn- and post-collisional crustal-scale cross-sections. *Marine and Petroleum Geology*, 12(8), 903–915. [https://doi.org/10.1016/0264-8172\(95\)98854-X](https://doi.org/10.1016/0264-8172(95)98854-X)
- Vergés, J., Fernandez, M., & Martinez, A. (2002). The Pyrenean orogen: Pre-, syn-, and post-collisional evolution. *Journal of the Virtual Explorer*, 08, 55–74. <https://doi.org/10.3809/jvirtex.2002.00058>

- Vermeesch, P., & Tian, Y. (2014). Thermal history modelling: HeFTy vs. QTQt. *Earth-Science Reviews*, 139, 279–290. <https://doi.org/10.1016/j.earscirev.2014.09.010>
- Vignaroli, G., Faccenna, C., Jolivet, L., Piromallo, C., & Rossetti, F. (2008). Subduction polarity reversal at the junction between the Western Alps and the Northern Apennines, Italy. *Tectonophysics*, 450(1–4), 34–50. <https://doi.org/10.1016/j.tecto.2007.12.012>
- Viola, G., Odonne, F., & Mancktelow, N. S. (2004). Analogue modelling of reverse fault reactivation in strike-slip and transpressive regimes: application to the Giudicarie fault system, Italian Eastern Alps. *Journal of Structural Geology*, 26(3), 401–418. <https://doi.org/10.1016/j.jsg.2003.08.014>
- Wagner, G.A., Reimer, G.M., Jäger, E. (1977). Cooling ages derived by apatite fission track, mica Rb–Sr and K–Ar dating: the uplift and cooling history of the central Alps. *Mem. Sci. Geol. Padova* 30, 1–27.
- Waldner, M., Bellahsen, N., Mouthereau, F., Bernet, M., Pik, R., Rosenberg, C. L., & Balvay, M. (2021). Central Pyrenees Mountain Building: Constraints From New LT Thermochronological Data From the Axial Zone. *Tectonics*, 40(3). <https://doi.org/10.1029/2020TC006614>
- Wessel, P., Luis, J. F., Uieda, L., Scharroo, R., Wobbe, F., Smith, W. H. F., & Tian, D. (2019). The generic mapping tools version 6. *Geochemistry, Geophysics, Geosystems*, 20(11), 5556–5564. <https://doi.org/10.1029/2019GC008515>
- Whitchurch, A. L., Carter, A., Sinclair, H. D., Duller, R. A., Whittaker, A. C., & Allen, P. A. (2011). Sediment routing system evolution within a diachronously uplifting orogen: Insights from detrital zircon thermochronological analyses from the South-Central Pyrenees. *American Journal of Science*, 311(5), 442–482. <https://doi.org/10.2475/05.2011.03>
- Wu, L., Monié, P., Wang, F., Lin, W., Ji, W., Bonno, M., Münch, P., and Wang, Q. (2016). Cenozoic exhumation history of Sulu terrane: Implications from (U–Th)/He thermochronology. *Tectonophysics*, 672–673, 1–15. <https://doi.org/10.1016/j.tecto.2016.01.035>
- Ziegler, P. A. (1992). European Cenozoic rift system. *Tectonophysics*, 208(1–3), 91–111. [https://doi.org/10.1016/0040-1951\(92\)90338-7](https://doi.org/10.1016/0040-1951(92)90338-7)

References From the Supporting Information

- Sun, S. S., and McDonough, W. F. (1989) Chemical and isotopic systematics of oceanic basalts: implications for mantle composition and processes. *Geological Society, London, Special Publications*, 42, 313–345. <https://doi.org/10.1144/GSL.SP.1989.042.01.19>