



# 1. PRODUCER PERSPECTIVE: ON-FARM LAND AND WATER USE

We measured water use for soybean, maize, rice and bean [4] and modeled water use for cattle in Mato Grosso, Brazil [5]. Results (**Tab. 1**) offer insight into strategies to improve water and land productivity with a focus on agricultural production output.

## Main conclusions

1. Irrigation can be used to enable three crop harvests per year through early soybean planting, and a dry season bean crop. Irrigated soybean showed the same yield as rain-fed soybean [4];
2. The largest water use activity for cattle production came from evaporation of farm impoundments typically used for drinking water (**Fig. 2**). Total Mato Grosso impoundment area reached 44,000 ha in 2014 [5].

**Tab. 1.** Water use for crop and cattle respectively measured [4] and modeled [5] in Mato Grosso, Brazil.

| <i><b>Product</b></i> | <i><b>Water use</b></i> |                  |
|-----------------------|-------------------------|------------------|
| Soybean               | 800-1300                | L/kg             |
| Maize                 | 320-500                 | L/kg             |
| Rice                  | 725-950                 | L/kg             |
| Bean                  | 1500-1800               | L/kg             |
| Cattle                | 251-255                 | L/kg live weight |

## Methods and Materials

### 1. Water use for cropland

Crop evapotranspiration (ET) was measured with an eddy covariance tower [4] installed at Capuaba Farm in Lucas do Rio Verde, Mato Grosso, Brazil (**Fig. 3**) (18 Sep 2015 to 4 Feb 2017). Crop water use was calculated as the sum of crop ET over the course of the crop development cycle (**Fig. 1**), to derive a crop water use per kg of soybean from the yield reported by the farmer [4].



**Fig. 3.** The eddy covariance tower installed at Capuaba Farm in Lucas do Rio Verde, Mato Grosso, Brazil Link (<http://ameriflux.lbl.gov/sites/siteinfo/BR-CMT>).

[VIDEO] <https://www.youtube.com/embed/MUFU0k3yek0?feature=oembed&fs=1&modestbranding=1&rel=0&showinfo=0>

**Vid. 1.** The soybean development cycle at Capuaba Farm in Lucas do Rio Verde, Mato Grosso, Brazil [4]

## 2. Water use for cattle

We modeled average water use for cattle in Mato Grosso, Brazil, following the model of Ridoutt *et al* [6] which combines water use with the cattle development cycle. We considered an average cattle development cycle of 48 months with both pasture or feedlot finishing in the 6 months leading up to slaughter. Evaporation from small farm impoundments ( $W_{res}$  in **Fig. 4**) were obtained using remote sensing by detecting total reservoir surface area within pasture land uses mapped by LAPIG (<http://www.lapig.iesa.ufg.br/lapig/>) (upper limit of 1 km<sup>2</sup> per reservoir).




Fig. 4. Average water consumption modeled for cattle ( $VWF_{animal}$ ) in Mato Grosso, Brazil, considering both pasture and feedlot finishing (6 months prior to slaughter): water use for feed ( $W_{feed}$ ), milk consumption ( $W_{milk}$ ), water drunk by the animal ( $W_{drink}$ ), water used to mix feed ( $W_{mix}$ ), evaporation from reservoirs ( $W_{res}$ ), and metabolic water ( $W_{met}$ ) [5].

## 2. PRODUCER PERSPECTIVE: BASIN LAND AND WATER USE

We carried out a water footprint sustainability assessment [7] in the Xingu Basin of Mato Grosso (Fig. 5) to assess conditions under which land and water sustainable limits may be exceeded [8].

### Main conclusions:

1. Current land and water uses are within sustainable limits, but these limits may be reached in 2030 and 2050 following agricultural expansion and intensification;
2. Irrigation together with cattle intensification could push the basin towards sustainable limits in the dry season;
3. Compliance to the Brazilian Forest Code, sensible intensification, and increased on-farm water use efficiency could prevent future conflicts over water use up- and downstream, and in- and outside the basin.





Fig. 5. The Xingu Basin of Mato Grosso, Brazil, separated into the Upper Xingu Basin and Xingu Headwaters [8].

### Methods and Materials



We calculated water scarcity for the basin (**Fig. 5**) in 2000, 2014, 2030 and 2050 by computing the ratio of total water use (agricultural, industrial and domestic) with water availability (as natural runoff minus demand) [7].

Land and water use in the basin were obtained following agricultural production scenarios and assumptions about livestock population and production, and domestic and industrial water uses (**Fig. 6**) [8]. Water availability was obtained using the IBIS model combined to land use information (2000, 2014), and deforestation scenarios (2030, 2050) [8].




**Fig. 6.** Initial conditions (2000, 2014) and agricultural production scenarios (2030, 2050) considered for estimating land and water use in the Xingu Basin of Mato Grosso [8].

## MAIN MESSAGE

This study aims to highlight the **specific roles of actors on land and water use for soybean and cattle production in Mato Grosso, Brazil**, by combining perspectives of producers and consumers, considering (1) agricultural products, and (2) hydro-geography (**Fig. 1**). The state of Mato Grosso (**Fig. 2**) is the largest soybean and cattle producer in Brazil with close to 30 Mtons soybean produced annually, and 30 million heads of cattle [1]. Soybean and cattle production have historically relied on deforestation of tropical forest and savanna in the Amazon and Cerrado biomes to expand.

We combined field measurements, modeling and trade analysis to map land and water use actions specific to producers and consumers (**Fig. 1**).




**Fig. 1.** Summary of actions on land and water use for soybean and cattle production in Mato Grosso, Brazil. Actions are focused on either the product, or hydro-geographic region, and are specific to producers (left) and consumers (right). Numbers 1-4 refer to the studies shown in panels 1-4.

### Approach

We carry out product- and hydro-geographic focussed assessments [2] of land and water use in Mato Grosso, Brazil (**Fig. 2**), described in detail in the four panels:

1. Producer land and water use at the field level;
2. Producer land and water use at the river basin level;
3. Consumer land use through trade;
4. Consumer land and water product impact assessment.



**Fig. 2.** The state of Mato Grosso in central western Brazil with its Amazon, Cerrado and Pantanal biomes [3]




### 3. CONSUMER PERSPECTIVE: EMBEDDED LAND USE

We obtained embedded deforestation in soybean leaving Mato Grosso, Brazil, using a combination of high resolution trade data and remote sensing shown in the Transparency for Sustainable Economies (Trase) platform [9] (**Vid. 2**). Consuming countries obtain commodities such as soybean (as bean, oil and cake) and beef which carry deforestation and greenhouse gas emissions from land use change.

#### Main conclusions

1. In 2017, Mato Grosso, Brazil, produced 30 Mtons of soybean (as bean, oil and cake), of which 42% was destined to China, 21% remained in Brazil, 15% was shipped to the EU and 22% to the rest of the world (**Fig. 7**);
2. These exports were associated with almost 15,000 ha deforestation in 2017, leading to 2.8 Mtons of CO<sub>2</sub> emission. There was greater soy deforestation in the Amazon biome (58%) than in the Cerrado (43%);
3. Roughly half of soy trade was made under a zero deforestation commitment in 2017.



**Fig. 7.** Soybean trade flows from Mato Grosso, Brazil, used in Brazil and exported internationally [9]. Trade flows are coloured by zero deforestation commitment: Soy Moratorium (blue), Company commitment (green), or None (red)

#### Methods and Materials

##### 1. Trade analysis

The Transparency for Sustainable Economies (Trase) platform (**Vid. 2**) relies on the combination of high resolution trade information, asset information (silos, ports) and tax information to link Brazilian municipalities of production to destination countries [9]. This information is then combined to production [10] and deforestation maps [11,12] to derive soybean deforestation. Greenhouse gas emissions were estimated using publicly available carbon maps [13].

[VIDEO] <https://www.youtube.com/embed/wMnAQJBtj8?feature=oembed&fs=1&modestbranding=1&rel=0&showinfo=0>

**Vid. 2:** Introducing the Transparency for Sustainable Economies (TRASE) platform [9]

##### 2. Soybean deforestation and zero deforestation commitments


Zero deforestation commitments are distinguished between "Soy Moratorium" and "Company commitments" based on whether soybean has been sourced from the Amazon biome, or whether a trader has made a specific pledge to zero deforestation (identified through company websites).

# 4. CONSUMER PERSPECTIVE: PRODUCT IMPACT ASSESSMENT

We calculated environmental impacts of agricultural production by considering impacts of intensification and extensification scenarios to the water cycle per production output (**Fig. 8, 9**) [14].

**Main conclusions**

- 1. Agricultural extensification has the potential to reduce precipitation and increase runoff in the basin;
- 2. Impacts on the water cycle were greater in the Amazon compared to the Cerrado biome;
- 3. Additional impacts must be considered such as loss of biodiversity and greenhouse gas emissions [15].



**Fig. 8.** Impacts of agricultural production in the Amazon biome (m³ per kg of crop) considering rain-fed and irrigated cropland extensification onto tropical forest (labeled as "forest"), and intensification on current pastureland (labeled as "pasture"). Cropland includes options of growing rain-fed soybean, maize, or irrigated soybean, rain-fed rice, and irrigated bean [14].




Fig. 9. Impacts of cattle production in Amazon and Cerrado biomes (m³ per kg of live weight) considering pasture expansion onto tropical forest in both biomes [14].

#### Methods

We use life cycle assessment (LCA) to estimate impacts per kg of product using water use estimated in panels 1 and 2, and using the Xingu Basin as the geographic boundary (Fig. 5 shows a portion of this basin which extends further north). The method follows the framework for including land use in LCA [16], focussing specifically on impacts of deforestation on precipitation recycling [17] and groundwater recharge [18] following the equation  $\text{Impact} = A \cdot \text{Factor}$ , where **Impact** is the impact to the water cycle (positive or negative flows per kg),  $A$  (ha (kg product)⁻¹) is the area required for 1 kg of product, and  $\text{Factor}$  (m³ ha⁻¹) is the impact factor allowing to allocate the impact per kg of product [16].

# DISCLOSURES

## Acknowledgements

This research was supported by the Natural Sciences and Engineering Research Council (NSERC) Vanier Canada Graduate Scholarship [201411DVC-347484-257696], the NSERC Michael Smith Foreign Study Supplement [488016-2015], and the Mitacs Globalink Research Award [IT05435]. The Transparency for Sustainable Economies platform (Trase, [trase.earth](http://trase.earth) (<http://trase.earth>)) is supported by the Gordon and Betty Moore Foundation, the Swedish Research Council Formas and the World Wildlife Fund.

The authors acknowledge the additional contributors to this work and its results: H.J. Dalmagro, M.S. Johnson, A.T. Black, P.H.Z. Arruda, I. Hawthorne, E.G. Couto, C. Bulle, E.J. Miranda, K. Solvik, J. Graesser, A. Castanho, M. N. Macedo, M.T. Coe, P. Löfgren, B. Ayre, and E. zu Ermgassen.

## ABSTRACT

For decades, agricultural expansion in Southern Amazonia has relied on deforestation to increase the national and global supplies of cattle and soybean products. Research on the use of land and water for increasing the region's pasture and cropland outputs has provided important insight into the role of producers in managing these resources. However, the roles of other actors on cattle and soybean production systems (e.g. traders) have been less apparent. For instance, some private initiatives, such as the Cattle Agreement or the Soybean Moratorium, have been proposed to curb deforestation through the cattle and soybean supply chains as a means to influence the production process. Here, we highlight the role of producers and consumers in the use of land and water resources for agricultural production in the state of Mato Grosso, Brazil, by combining field measurements and modelling with trade mapping. We use the Transparency for sustainable economies platform (Trase, <https://trase.earth>) to highlight the role of trade actors by combining high resolution trade information (e.g. custom declarations) with up-to-date deforestation and water scarcity maps. In 2015, up to five traders with zero deforestation commitments exported over 40% of soybean produced in the state of Mato Grosso. Within this context, producers can increase agricultural output by irrigating cropland, and/or concentrating cattle production on current pastureland, but this combined system has the potential to increase water scarcity in the dry season. Our analysis provides additional information on the drivers shaping the Brazilian agricultural frontier and attempts to bring producers and consumers closer together in agricultural supply chains.

## REFERENCES

- [1] SIDRA-IBGE 2018 Banco de dados agregados sidra.ibge.gov.br (<http://sidra.ibge.gov.br>)
- [2] Lathuillière MJ, Bulle C, Johnson MS 2018 A contribution to harmonize water footprint assessments *Global Environmental Change* **53**: 252-264 doi: 10.1016/j.gloenvcha.2018.10.004 Link (<http://www.sciencedirect.com/science/article/abs/pii/S0959378017314656>)
- [3] Lathuillière MJ, Johnson MS, Galford GL, Couto EG 2014 Environmental footprints show China and Europe's evolving resource appropriation for soybean production in Mato Grosso, Brazil *Environmental Research Letters* **9**(7): 074001 doi: 10.1088/1748-9326/9/7/074001 Link (<http://iopscience.iop.org/article/10.1088/1748-9326/9/7/074001/meta>)
- [4] Lathuillière MJ, Dalmagro HJ, Black AT, Arruda PHZ, Hawthorne I, Couto EG, Johnson MS 2018 Rain-fed and irrigated cropland-atmosphere water fluxes and their implications for agricultural production in Southern Amazonia *Agricultural and Forest Meteorology* **256-257**: 407-419 doi: 10.1016/j.agrformet.2018.03.02 Link (<http://www.sciencedirect.com/science/article/pii/S0168192318301060>)
- [5] Lathuillière MJ, Solvik K, Macedo MN, Graesser J, Miranda EJ, Couto EG, Johnson MS 2018 Cattle production in Southern Amazonia: implications for land and water management *in preparation*
- [6] Ridoutt BG, Sanguansri P, Freer M, Harper GS 2012 Water footprint of livestock: comparison of six geographically defined beef production systems *The International Journal of Life Cycle Assessment* **17**: 165-175 doi: 10.1007/s11367-011-0346-y Link (<http://link.springer.com/article/10.1007%2Fs11367-011-0346-y>)
- [7] Hoekstra AY, Chapagain AK, Aldaya MM, Mekonnen MM 2011 *The Water Footprint Assessment Manual* (Earthscan) London UK Link (<http://waterfootprint.org/en/resources/publications/water-footprint-assessment-manual/>)
- [8] Lathuillière MJ, Coe MT, Castanho A, Graesser J, Johnson MS 2018 Evaluating Water use for Agricultural Intensification in Southern Amazonia Using the Water Footprint Sustainability Assessment *Water* **10**: 349 doi: 10.3390/w10040349 Link (<http://www.mdpi.com/2073-4441/10/4/349>)
- [9] Transparency for Sustainable Economies (Trase) 2018 trase.earth (<http://trase.earth>)
- [10] Agrosatellite 2018 <http://agrosatellite.com.br> (<http://agrosatellite.com.br>)
- [11] INPE 2018 Taxas anuais de desmatamento na Amazônia Legal Brasileira (AMZ) <http://www.obt.inpe.br/prodes/dashboard/prodes-rates.html> (<http://www.obt.inpe.br/prodes/dashboard/prodes-rates.html>)
- [12] INPE 2018 Incrementos de desmatamento anuais no Cerrado Brasileiro <http://terrabrasilis.dpi.inpe.br/dashboard/deforestation/biomes/cerrado/increments/> (<http://terrabrasilis.dpi.inpe.br/dashboard/deforestation/biomes/cerrado/increments/>)
- [13] FUNCATE 2018 <https://www.funcate.org.br/pt/> (<https://www.funcate.org.br/pt/>)
- [14] Lathuillière MJ, Bulle C, Johnson MS 2018 Complementarity in mid-point impacts for water use in life cycle assessment applied to cropland and cattle production in Southern Amazonia *Journal of Cleaner Production* in review
- [15] Lathuillière MJ, Miranda EJ, Bulle C, Couto EG, Johnson MS 2017 Land occupation and transformation impacts of soybean production in Southern Amazonia *Journal of Cleaner Production* **149**: 680-689 doi: 10.1016/j.jclepro.2017.02.120 Link (<http://www.sciencedirect.com/science/article/pii/S0959652617303438>)
- [16] Koellner T, de Baan L, Beck T, Brandão M, Civit B, Margni M, Milà i Canals L, Saad R, Maia de Souza D, Müller-Wenk R 2013 UNEP-SETAC guideline on global land use impact assessment on biodiversity and ecosystem services in LCA *The International Journal of Life Cycle Assessment* **18**(6): 1185-1187 doi: 10.1007/s11367-013-0579-z Link (<http://link.springer.com/article/10.1007/s11367-013-0579-z>)
- [17] Lathuillière MJ, Bulle C, Johnson MS 2016 Land use in LCA: including regionally altered precipitation to quantify ecosystem damage *Environmental Science and Technology* **50**(21): 11769-11778 doi: 10.1021/acs.est.6b02311 Link (<http://pubs.acs.org/doi/10.1021/acs.est.6b02311>)
- [18] Saad R, Koellner T, Margni M 2013 Land use impacts on freshwater regulation, erosion regulation, and water purification: a spatial approach for a global scale level *The International Journal of Life Cycle Assessment* **18**(6): 1253-1264 doi: 10.1007/s11367-013-0577-1 Link (<http://link.springer.com/article/10.1007/s11367-013-0577-1>)



