



Authors: Ruiyang Yu<sup>1</sup>, Yunjun Yao<sup>1,\*</sup>, Ke Shang<sup>1</sup>, Junming Yang<sup>1</sup>, Xiaozheng Guo<sup>1</sup>, Zijing Xie<sup>1</sup>


1. Beijing Normal University, Beijing 100875, China; yuruiyang927@163.com (Ruiyang Yu)

\* Correspondence: boyyunjun@163.com

## Introduction

Terrestrial latent heat flux (LE), which refers to the heat flux transferred from the land surface to the atmosphere through soil evaporation, vegetation transpiration and interception, is an essential component for characterizing the global and regional hydrological budget, energy redistribution and carbon cycles. Accurate estimation of LE at high spatial resolution is of great vital importance for energy balance and water resource management, especially for agricultural production at field scale. However, there are relatively few LE products based on Chinese GaoFen-1 (GF-1) data.

## Results



The validation results show that the 16 m GF and 1 km GLASS LE products of the three countries all presented good spatial consistency, and the coefficient of determination of LE estimates based on GF-1 data against GLASS LE products were all greater than 0.6, indicating that the accuracy of LE products based on GF-1 data was high.

## Data

### Study area:

The part regions of Ethiopia, Laos and Pakistan

### Satellite data:

The 16 m Chinese GF-1 remote sensing data

### Modern Era Retrospective Analysis for Research and Applications-2 (MERRA-2) Meteorological data:

Air temperature ( $T_a$ ), the ATI derived by the diurnal air temperature range ( $DT$ ), the net radiation ( $R_n$ )

### LE product:

1 km Global LAnd Surface Satellite (GLASS) LE product

## Conclusions

The 16 m GF and 1 km GLASS LE products of the three countries all presented good spatial consistency. Validated by GLASS LE product, the accuracy of LE products based on GF-1 data was high. LE estimation based on GF-1 data is of great significance for energy balance and water resource precision management.

## Acknowledgments

This research was funded by the National Natural Science Foundation of China (Grant No. 41671331), the National Key Research and Development Program of China (Grant No. 2016YFA0600103 and No. 2016YFA0600102).

### REFERENCE

[1] Yao, Y.; Liang, S.; Cheng, J.; Liu, S.; Fisher, J.B.; Zhang, X.; Jia, K.; Zhao, X.; Qing, Q.; Zhao, B.; et al. MODIS-driven estimation of terrestrial latent heat flux in China based on a modified Priestley-Taylor algorithm. *Agric. For. Meteorol.* 2013, 171, 187–202.

## Methods

GF-1 LE product was generated based on a Modified Satellite Priestley - Taylor (MS-PT) algorithm proposed by Yao et al. [1] The terrestrial LE consists of the unsaturated soil evaporation ( $LE_s$ ), the canopy transpiration ( $LE_c$ ), the canopy interception evaporation ( $LE_{ic}$ ), and the saturated wet soil surface evaporation ( $LE_{ws}$ ). The MS-PT algorithm can be given by:

$$\begin{aligned} LE &= LE_s + LE_c + LE_{ic} + LE_{ws}, \\ LE_s &= (1 - f_{wet}) f_{sm} \alpha \frac{\Delta}{\Delta + \gamma} (R_{ns} - G), \\ LE_c &= (1 - f_{wet}) f_v f_T \alpha \frac{\Delta}{\Delta + \gamma} R_{nv}, \\ LE_{ic} &= f_{wet} \alpha \frac{\Delta}{\Delta + \gamma} R_{nv}, \quad LE_{ws} = f_{wet} \alpha \frac{\Delta}{\Delta + \gamma} (R_{ns} - G), \\ f_{sm} &= ATI^k = \left( \frac{1}{DT} \right)^{DT/DT_{max}}, \quad f_{wet} = f_{sm}^4, \\ f_v &= \frac{NDVI - NDVI_{min}}{NDVI_{max} - NDVI_{min}} \end{aligned}$$

where  $\alpha$  is the Priestly-Taylor coefficient (1.26),  $\gamma$  is the psychrometric constant (0.066 kPa  $^{\circ}\text{C}^{-1}$ ),  $\Delta$  is the slope of saturation water vapor pressure versus temperature curve,  $f_{wet}$  is the relative surface wetness,  $f_{sm}$  is soil moisture constraint,  $f_T$  represents plant temperature constraint ( $\exp(-(T_a - T_{opt})/T_{opt})^2$ ),  $DT_{max}$  describes the maximum diurnal air temperature range (40  $^{\circ}\text{C}$ ),  $T_{opt}$  is an optimum temperature (25 $^{\circ}\text{C}$ ),  $R_{ns}$  is the surface net radiation to the soil ( $R_{ns} = R_n(1 - f_v)$ ),  $G$  is soil heat flux ( $\mu R_n(1 - f_v)$ ,  $\mu = 0.18$ ),  $R_{nv}$  represents the surface net radiation to the vegetation ( $R_{nv} = R_n f_v$ ),  $f_v$  is the vegetation cover fraction, and  $NDVI_{min}$  and  $NDVI_{max}$  are the minimum and maximum normalized difference vegetation index (NDVI), respectively.