

Time, hydrologic landscape and the long-term storage of peatland carbon in sedimentary basins

David J. Large^{1*}, Chris Marshall², Malte Jochmann^{3,4}, Maria Jensen³, Baruch F. Spiro⁵,
Snorre Olaussen³

¹ Department of and Environmental Engineering, Faculty of Engineering, University of Nottingham,
University Park, Nottingham NG7 2RD, UK (0115) 82 32760

² Environmental Research Institute, University of the Highlands and Islands, North Highland College UHI,
Castle Street, Thurso, KW14 7JD, United Kingdom

³ University Centre in Svalbard (UNIS) P.O Box 156, N-9171, Longyearbyen, Norway

⁴ Store Norske Spitsbergen Kulkompani AS, PO Box 613, N-9171 Longyearbyen, Norway

⁵ Department of Mineralogy, Natural History Museum, Cromwell Road, London SW7 5BD, UK

Corresponding author: David J. Large (david.large@nottingham.ac.uk)

Key Points:

- To understand how peat enters long term carbon storage requires consideration of peat accumulation over periods of 10^5 to 10^6 years.
- Peat accumulation over these timeframes requires sustained storage of water above depositional base level
- This has profound implications for our understanding of peat in the earth system and the interpretation of the geological record

Abstract

Peatland carbon may enter long-term storage in sedimentary basins preserved as either coal or lignite. To understand the process by which this happens requires extrapolation of our understanding of peatland carbon accumulation over timescales that greatly exceed those of Holocene peat. By applying this extrapolation, we deduce that the amount of time required to account for the carbon in 1 – 10 m thick coal seams must represent 10^5 to 10^6 years, an order of magnitude more than assumed in current interpretations of stratigraphic frameworks. Extrapolating peat growth to periods of 10^6 years requires consideration of the conditions of landscape, hydrology, accommodation space and crustal deformation required to ensure the sustained growth and accommodation of peat deposits. We conclude that the generation of accommodation space at low rates (0.1 to 0.2 mm/yr) can adequately accommodate thick peat accumulation over periods $>10^5$ yrs. However, generation of accommodation space at high rates (>1 mm/yr) cannot accommodate significant peat accumulation. Key to this process is the maintenance of a saturated peat body above the level of clastic deposition. This long-term stability of the system has implications for our understanding of the processes that limit peatland growth under very low accommodation rates and the long-term validity of current peat growth models.

Plain Language Summary

This is optional but will help expand the reach of your paper. Information on writing a good plain language summary is available [here](#).

1 Introduction

Peatland carbon may enter into long-term crustal storage in sedimentary basins where it occurs as either coal or lignite. To understand the process by which this happens requires a conceptual bridge between peatland processes measured on Holocene timescales and processes over the much greater timescales required to explain features of substantial deposits of coal and lignite. Key to this conceptual bridge are the consequences of extrapolating Holocene peatland processes over long timeframes. However, these consequences are not considered by either peat scientists or coal geologists, and the absence of this analysis results, not only in flawed stratigraphic interpretation but also undermines our ability to understand peat within the Earth system on timescales that greatly exceed those of the Holocene.

Coal is one of our most significant energy resources and generations of geologists studied coal; its sedimentology, geochemistry and stratigraphy. Many approaches (Diessel 1992; Dai et al., 2020) have been applied to interpret coal stratigraphy. Currently, the most favoured approaches integrate coal into stratigraphic frameworks using the framework proposed by Bohacs and Suter (1997) and subsequently developed and refined (Diessel et al., 2000; Holz et al., 2002; Jerrett et al., 2011a; Jerrett et al., 2011b). In this stratigraphic framework, growth of the water-saturated precursor peat responds to the rate of change of the space available for sediment accumulation, known as the accommodation space. The balance between the rate at which accommodation space changes (on account of tectonic subsidence, change in sea level and sediment accumulation) and the peat production rate determines the resulting thickness, areal extent and composition of the peat. This stratigraphic framework acknowledges but does not focus on other factors (e.g. climate, groundwater, vegetation, landscape) that influence peat accumulation. By doing this, coal is integrated into stratigraphic models (Davies et al., 2005; Jerrett et al., 2011b; Michaelsen et al., 2000; Staub, 2002) where it forms an interpretative bridge between the marine and terrestrial realms (Wadsworth et al., 2010).

While there is an unquestionable need to accommodate peat to form coal via the creation of accommodation space, the assumptions that underpin the interpretation of coal within stratigraphic frameworks including that of Bohacs and Suter (1997) are not firmly grounded in the processes that determine the formation and accumulation of peat. Specifically there are two flaws: stratigraphic frameworks consider only the volumetric growth of peat (predominantly water) and subsequent compaction while ignoring the processes that determine carbon accumulation; and they also assume that peat properties, including thickness and composition, are highly responsive to the rate at which accommodation space is generated. The first flaw results in a gross order of magnitude underestimate of the time required to account for the carbon in coal; the second flaw devalues the interplay between hydrology, landscape and water storage in determining long-term hydrological stability and duration of a peat body. Consequently, stratigraphic interpretations based on the approach of Bohacs and Suter (1997) treat coal as a relatively transient component of the stratigraphic record, which in turn requires the inference of significant stratigraphic hiatuses (e.g. Jerrett et al., 2011a; Jerrett et al., 2011b; Scott and Stephens, 2015). It also results in coal seams that must have accumulated over periods of 10^5 to 10^6 yr being interpreted as would a Holocene peat with duration of 10^4 yr, effectively ignoring the extent and potential stratigraphic and palaeoenvironmental value of the record contained within the coal. Consequently, the approach also fails to recognise the capacity peat to modify sediment transport and storage within a basin and the capacity of coal to inform our long-term

understanding of basin hydrology and its evolution relative to active structures and palaeo-landscape.

The study of Holocene peat on its own is inherently constrained by the Holocene timeframe and Holocene peat scientists rarely consider the extension of Holocene processes over timeframes in which peat may enter crustal storage or become limited within the current landscape. For example, peat growth models (e.g. Baird et al., 2012; Morris et al., 2012; Frothingham et al., 2010) are never extended over timeframes that greatly exceed those of the Holocene. Consequently, their validity with respect to long-term crustal storage of peat remains untested.

In this paper, we address this issue by defining a new basis for the stratigraphic interpretation of peat, lignite and coal founded on the understanding of carbon accumulation within a hydrologic landscape. The result is a more informed model based on a wider scientific basis that encapsulates the interaction of geological structures, landscape and hydrology over a wide range of timescales and enables a greater understanding of peatland within the Earth system. Furthermore the results demonstrate that lignite and coal can provide invaluable long duration records of the conditions prevailing on the Earth surface at the time of formation.

2. Basis for a new hydrologic landscape approach

In the sections that follow, we define base level as the local graded stream profile, the level at which clastic deposition and erosion balance (Catuneanu, 2006). The rate at which accommodation space (Catuneanu, 2006) is generated is, for simplicity and to align with the use in the stratigraphic model based on Bohacs and Suter (1997), referred to as the accommodation rate.

For ease of reference, we use the term coal as the generic term to refer to both coal and lignite. In this context coal is a readily combustible rock containing more than 50 percent organic matter by weight (on a dry basis), which was formed from the compaction and alteration of plant remains (Jackson, 1997; Alpern and de Sousa, 2002; Schopf, 1956). There is no single formal definition of peat so for the purposes of this paper we use a definition modified after Schopf (1966) in which peat is an un-consolidated, hydrophilic, carbonaceous sediment, formed by accumulation of partially fragmented and decomposed, plant remains. The precursor to coal is always peat however not all materials classed as peat could form coal.

2.1 Carbon accumulation and time

As coal is predominantly carbon, a carbon basis for determining the duration of a coal seam is far more appropriate and robust (Large and Marshall, 2015) than volumetric alternatives. The reason for this is that the processes determining the rate of carbon accumulation in peat and subsequent loss of carbon during coalification are well understood and quantifiable (Large and Marshall, 2015; Clymo, 1984; Diessel, 1992). For example, starting with a range of peatland vegetation chemistry (sphagnum, sedge, shrub, wood) the relative positions of vegetation, peat

and coal on a van Krevelen diagram (Fig 1) are consistent with a continuous process of transformation of vegetation to coal. Loss of CO₂ and CH₄ in equal proportions can account for this trend (Fig 1) as can loss of more complex mixtures including dissolved organic carbon (Moore et al 2018). The positions of coal, peat and vegetation (Fig 1) are also consistent with a similar mix of peatland vegetation chemistry accounting for all coal compositions throughout geological time. This illustrates that the mass balance during the transition from peat to coal is a predictable and measureable process. By accounting for the mass lost during the transition from peat to coal it is also possible to predict (Large and Marshall, 2015) the coal thickness equivalent to 20kyrs of carbon accumulation (Fig.2) for a range of carbon accumulation rates that are typical of global rates in the Holocene.

This leads to three important conclusions. First, the duration of coal seams is about an order of magnitude greater than previously considered e.g. 1 m of bituminous coal formed under temperate palaeo-climatic conditions at a carbon accumulation rate of 20g/m²/yr would represent 100 kyrs rather than the typically assumed 10 kyrs (e.g. Diessel et al, 2000; Jerrett et al, 2011a). Second, most of the inorganic matter in coal can be accounted for by the deposition of atmospheric mineral dust at rates of 0.02 to 20 g/m²/yr (Large and Marshall, 2015; Marshall et al., 2016). Third, global variations in the composition of coal are predictable using uniformitarian assumptions of carbon accumulation rate and dust deposition (Large and Marshall, 2015; Marshall et al., 2016). This indicates that




Figure 1 A van Krevelen diagram illustrating the continuous relationship between the C-O-H composition of vegetation, peat, lignite and coal. Coal data spanning the compositional range from lignite to anthracite, ranging in age from Neogene to Carboniferous is from the USGS coal quality database ($n = 7000$). Mean peat, peatland and vegetation compositions are from Moore et al (2018) and references therein. Mean peat data is shown for UK, Latvia (Lat), Ontario (Ont), Indonesia (Ind) and Southeast USA. Projection lines are indicated assuming mass is lost as mean organic matter CH₂O equivalent to loss of CO₂ and CH₄ in equal proportions. Positions of average swamp and bog are shown after 60% and 70% loss of mass.

176

177




Figure 2 Plot showing the relationship between the coal seam stratigraphic thickness equivalent to 20 ky of carbon accumulation, carbon concentration in coal on a dry ash free basis and carbon accumulation rate in the precursor peat. Carbon accumulation rates are chosen to represent the greater part of the range of reported Holocene rates, so are equivalent to the long term rates measured over ca. 10 kyrs. Coal thicknesses were calculated using the empirical carbon loss model of Large and Marshall (2015) and a starting carbon concentration in the precursor peat of 52% by weight on a dry ash free basis.

the processes of carbon accumulation in peat have not varied markedly over geological time (Large and Marshall, 2015; Marshall et al., 2016).

Several stratigraphic consequences arise from these conclusions:

1) Thick coal seams (1 m to > 20 m) can be of sufficient duration to span, without hiatus, periods of 0.1 to > 1 Ma and may account for more time than inter-seam clastic sediments (Large and Marshall, 2015).

2) Coal is indicative of periods of sediment bypass during which fluvial clastic deposition is excluded. This conclusion arises as:

a. Silici-clastic deposition rates in coal forming environments (floodplains, estuaries and deltas) are typically in the range 0.02 to 1 mm per year (Einsele, 2000)

b. At typical bulk soil densities of 1200 to 1600 kg/m³ this corresponds to mass deposition rates of silici-clastic sediments of 24 to 1600 g/m²/yr.

c. Supply of silic-clastic matter to peat at rates greater than 35 g/m²/yr would result in a rock with greater than 50% inorganic matter that could not be classed as a coal, and even the lowest of these rates would produce a low quality coal containing at least 20% inorganic matter.

3) In a given palaeo-geographic setting, the long-term rate of carbon accumulation tends to be constant and lie within the global Holocene range. This does not mean that the rate of carbon accumulation does not vary; it only implies that given suitable hydrology the conditions of nutrient supply (via atmospheric deposition), climate and productivity will tend to produce a stable long-term average rate in a given palaeo-geographic setting. It is therefore more appropriate to assume that for coals of similar paleogeography and rank thicker coal accumulated over longer periods of time not at a faster rate.

4) The accumulation of peat over long periods requires sustained supply of fresh water, which in turn requires sustained hydrological stability.

Some of the consequences above have been inferred previously. For example, Broadhurst and France (1986) and Broadhurst and Simpson (1983) examining the influence of syndepositional faulting on coal seams observed that coal seams were more strongly influenced by active structures and associated landscape than the interseam sediments. They concluded that the duration of coal seams must be considerably greater than that of the intervening sediments. Similarly, in the UK Carboniferous Scott and Stephens (2015) concluded that a greater amount of time must be in coal rather than clastic sediments.

2.2 Water storage

With typical dry bulk densities in the region of 0.1 g/cm^3 (Clymo, 1984) by mass and by volume, peat is predominantly water. The volume of water stored in peat in low-lying areas is responsible for the exclusion of clastic sediment and therefore represents a volume of water stored above base level. In upland areas of blanket bog, landscape incision and consequent sediment bypass are primarily responsible for the exclusion of clastic sediment from the peat. Therefore, accumulation and preservation of peat is not only a matter of balancing rates of peat growth and accommodation; it is also a matter of sustaining water storage in the form of peat above the level of clastic sediment deposition. This is an important and fundamental difference between this approach and previous approaches (e.g. Bohacs and Suter 1997; Diessel et al 2000).


Evidence for either being in a state of clastic supply or clastic exclusion is the typically abrupt transition (clastic sediment to coal and vice versa) from one state to another at the boundaries of many coal seams (Diessel, 1992). Even during more gradual transitions from clastic sediment to coal, intercalated thin coal and clastic horizons occur rather than a gradual dilution of clastic sediment with organic matter (Diessel, 1992). If we accept that there is more time in the coal than the clastic sediment this intercalation of clastic sediment and thin coal indicates that clastic accommodation space was extremely limited at the point where coal seams start to establish. A finding that is consistent with the base of coal seams being unconformable (Haszeldine, 1989).

The hydrological stability required to initiate and sustain a volume of water above the graded stream profile is strongly influenced by the capacity of groundwater to maintain the water saturated state of a peat body during periods of water shortage (Glaser et al., 2004; Glaser et al., 1997). Consequently, the position of peat within a landscape is strongly related to the discharge (or seepage) of groundwater (Winter, 2000) and the response of this discharge to crustal deformation (Glaser et al., 2004). Peatland supported by discharge from large groundwater catchments will be resilient to climate change. Peatland associated with small groundwater catchments that are highly dependent on annual rainfall will be less resilient (Winter, 2000, Hokansen et al., 2016).

2.3 The hydrologic landscape

Peat, as the precursor to coal, initiates in a landscape. The base of coal seams is typically unconformable i.e. on land (Haszeldine, 1989) and based on analysis of controlling variables (eustasy, tectonics, climate, vegetation) most coal seams initiate during periods of long-term lowering of global sea level (Railsback, 1995) i.e. when land is created. A useful way of conceptualising the relationship between peat, landscape and hydrology on different scales in a variety of settings is the hydrologic landscape concept (Winter, 2001). In this approach, Winter (2001) uses the idea of a fundamental hydrologic landscape unit. This unit consists of an upland separated from a lowland by a steeper slope (Fig 3) and incorporates both geology and climatic setting. Associated with the hydrologic landscape unit, Winter (2001) defines a hydrologic system (the components of the total hydrologic budget for any point on the landscape). This consists of surface water controlled by slope, slope aspect and surface permeability; ground water controlled by the subsurface hydraulic characteristics; atmospheric water controlled by climate. Where the supply of water is sufficient, ground water will tend to flow towards the surface (or discharge) at the base of a slope. Given a sufficiently wet climate or large enough groundwater catchment this creates the fresh water saturation required for the initiation and

257 maintenance of istable wetlands or peatlands (Winter, 2000). The base of slopes will therefore
 258 be the setting in which to initiate and maintain water saturated peat deposits (Fig 3).



259 There are two important aspects to the hydrologic landscape concept. First, the
 260 fundamental hydrologic landscape unit operates on multiple scales that can range from the break
 261 in slope between river terraces, to the transition from mountain range to basin. Scale is
 262 important because the scale of catchment determines the vulnerability and resilience of the
 263 peatland. Small catchments relating to small landscape units will be highly dependent on

Figure 3 A schematic representation of the fundamental hydrologic landscape unit described by Winter (2000) illustrating the optimal position for peat accumulation.

264 precipitation to sustain associated peatlands. Whereas, large-scale units and associated large-
 265 scale ground water flow will be capable of buffering peatlands through periods of variable
 266 climate (Winter, 2000). Second, landscape evolution driven by tectonics and shaped by erosion
 267 evolves slowly over periods of 10^5 to 10^7 years, whereas climate and base level may display
 268 large fluctuations over much shorter timeframes. Hence, the long periods (10^5 to 10^6 years) of
 269 hydrologic landscape stability required to account for thick coal seams (e.g. Large et al., 2004;
 270 Large, 2007; Briggs et al., 2007; Large and Marshall, 2015) must depend on hydrologic
 271 landscape units controlled by tectonics and active structures within a basin. This is essential if
 272 hydrologic stability is to be sustained during periods of higher frequency climate and base level
 273 fluctuation.

274 The importance of hydrologic landscape is clear in the Holocene during which long-
 275 term accumulation of peat depends more on the capacity of the landscape to sustain water
 276 storage than on the capacity to accommodate peat into long-term crustal storage. This is
 277 evidenced by the widespread accumulation of peat on various scales in a range of hydrologic
 278 landscapes (e.g boreal lowlands, upland blanket bogs, high altitude plateaus and coastal plains)
 279 irrespective of base level, uplift or subsidence. The short timeframe of the Holocene relative to
 280 the timescale of significant crustal deformation, also tends to limit our understanding of the
 281 tectonic and structural influence of glacio-eustatic rebound on peat initiation and hydrology
 282 (Glaser et al., 2004).

283 An aspect not explicitly considered by Winter (2001) is that as peat accumulates it will
 284 modify the hydrologic landscape, raising the local water table and shifting the breaks in slope.
 285 An example of this phenomenon is the spring line, visible on optical satellite images,

surrounding the Hongyuan peat (Large et al., 2009) and neighbouring peatlands on the Qinghai-Tibetan Plateau.

2.3 Influence of changing of base level

A change in base level constitutes a change in the hydrologic landscape. As base level rises, surface wetness and run off increase, and new accommodation space is generated. At the level of the local graded stream profile deposition and erosion balance, and the accommodation space is filled with clastic sediment.

In a hydrologic landscape in which peat has initiated and accumulated at the breaks in slope, rising base level is accompanied by clastic deposition away from breaks in slope. At smaller scale hydrologic landscape units, groundwater discharge will decrease and the water budgets required to sustain peat accumulation above base level will become increasingly dependent on local precipitation. If the water budget is insufficient to sustain a peat body above base level, the peat will be inundated by clastic sediment. As base level continues to rise, larger scale hydrologic landscape units will maintain continuous peat.

Falling base level will reverse this trend. Water tables will fall and peat surfaces at the basin margin will become more vulnerable and collapse in response to falling base level. However, if the scale of the hydrologic landscape units is sufficient large, peat growth may be sustained even during periods of falling base level. At the same time, new breaks in slope associated with topographic features on the emerging and incised land surface will provide the necessary areas of groundwater seepage to initiate new peat growth.

Note that collapse of the peat surface does not necessarily mean that the peat becomes drier, as mechanically weak peat will initially collapse and track the falling water table. Only once the water table has reached the level at which the peat has sufficient strength to resist compaction will the water table continue to fall below the peat surface resulting in drying and oxidation of the peat.

Another means of modifying the relationship between base level and the hydrologic landscape is tectonic tilting. During this process, water storage will shift down the hydraulic gradient. Up gradient peat will collapse and possibly oxidise as water is withdrawn, while the down gradient time equivalent peat will be water saturated and tend to expand. In these circumstances time equivalent peats could have developed markedly different ecosystems. If the collapsed peat surface falls below base level then the capacity exists to rapidly infill accommodation space above the peat surface. Collapse, due to water withdrawal, could explain the burial of large in-situ tree trunks immediately above some coal seams (Broadhurst and Magraw, 1961; Falcon-Lang, 2006) and in particular in areas adjacent to incised palaeochannels (Guion, 1987).

2.4 Accommodation and peat growth models

For peat to enter long term crustal storage it needs to be accommodated and rates of accommodation need to be capable of allowing continuous peat accumulation over periods of 10^5 to 10^6 yrs. To achieve this an appropriate balance of the rate of peat growth to accommodation rate is essential (Bohacs and Suter, 1997). This is complicated, as the rate of peat accumulation

with respect to time is an inherently non-linear balance between the input of organic matter and the mass of organic matter lost due to decay (Clymo, 1984). Furthermore this non-linear property of peat growth is ignored in previous stratigraphic models that assume linear peat growth rates (Diessel et al., 2000; Bohacs and Suter, 1997).

A simple evaluation of the consequences of this non-linearity can be made by extending simple peat growth models (Clymo et al., 1998; Clymo 1992) which provide realistic estimates of carbon accumulation over the Holocene to periods that greatly exceed 10ka. The shape of the peat growth curve generated by these models is governed by the initial decay rate, a (yr^{-1}), the decay rule that determines the change in decay rate as decay progresses, and the rate of input of carbon to the peat, p ($\text{kmolC m}^{-2} \text{yr}^{-1}$). Reasonable estimates can be made for rate of input and initial decay rates for the Holocene, however the decay rule over long periods of time is unknown. We also ignore short-term ecohydrological feedback. This is not unreasonable, as over long periods the mass balance between input and decay must shape the peat growth curve. Also, the longer term effects of ecohydrological feedback, limited by the climate space in which the peat forms should be represented as oscillations on the long term trend.




Figure 4 A) The relationship between cumulative peat thickness, cumulative peat carbon, time and linear accommodation rates (dashed lines with values given in mm/yr). Peat growth curves are derived using the quadratic model of Clymo et al (1998). Values of input (p) and initial decay rate (a) are chosen to represent the current range of global values: extreme low latitude, low latitude, boreal and high latitude. Conversion of cumulative carbon to thickness is based on a peat dry bulk density of 0.1 g/cm³ and a carbon concentration of 50% on a dry ash free basis. B) Derived from 4A this illustrates the height of peat accumulation above base level for accommodation rates of 0.1 mm/yr (thin lines) and 0.2 mm/yr (thick lines). Values of p and a are those given in 4A. C) The results in figure 4A expressed as bituminous coal thickness. Coal thickness was calculated from the cumulative carbon using the empirical method of Large and Marshall 2015 to account for loss of organic matter during coalification.

345 Clymo et al (1998).

$$M_t = (p/a) \left((\sqrt{1 + 2at}) - 1 \right)$$

346 Where M_t is the cumulative dry mass of carbon per unit area after time t (yr) and p and a are
 347 described above. This model is only a fixed point on continuum of possible decay rules (Clymo,
 348 1992) and is chosen for illustrative purposes as it generates reasonable accumulation over 10^4
 349 years and over much longer timeframes. Other points on the spectrum of possible models for
 350 example the constant decay model have unreasonable stratigraphic consequences over longer
 351 periods of time e.g. using Holocene parameters could not account for any significant coal
 352 accumulation.

353 Four growth curves were generated (Fig 4A) that cover a range of inputs and decay
 354 values. Values were chosen to represent: a high latitude value ($p=0.001$, $a=10^{-6}$); a boreal value
 355 ($p=0.003$, $a=10^{-4}$, similar to Clymo et al 1998); a low latitude value (0.008 , 10^{-3} much greater
 356 than the boreal values of Clymo et al 1998); and an extreme low latitude value ($p=0.1$, $a=0.1$)
 357 equivalent to a tropical net primary production and a high decay rate in tropical soil (e.g. Yule
 358 and Gomez, 2009).

359 The peat growth curves are then analysed relative to linear accommodation rates,
 360 ineffect constant subsidence. In reality accommodation rate may also be non-linear, but this
 361 simplifying assumption helps focus our analyses on the consequences of non-linear peat growth.
 362 To provide an understanding of what might be observed in terms of long term crustal storage
 363 equivalent bituminous coal thicknesses (Fig 4C) are calculated based on the cumulative carbon
 364 and accounting for the loss of carbon during coalification (Large and Marshall, 2015).

365 Analysis of the relationship between accommodation rate and the modelled growth
 366 curves (Fig 4) indicates the following key features for a system in which clastic supply is
 367 sufficient to fill the available accommodation space.

- 368 1) Peat accumulation above base level occurs if the value of p and its
 369 volumetric equivalent exceeds the accommodation rate (Fig 4A and 4B).
- 370 2) Significant peat accommodation sufficient to generate >50 cm of
 371 bituminous coal occurs at accommodation rates <0.5mm/yr (Fig 4C). Contrary to the
 372 conclusion of Bohacs and Suter (1997) it is improbable that high accommodation rates of
 373 1 to 2 mm per year could permit thick peat accumulation.
- 374 3) Tropical peats with high values of p can grow and establish above base
 375 level more rapidly (Fig 4B).
- 376 4) Over short timeframes of <70ka apparent long-term accumulation rates
 377 appear higher in the tropics and lower at the poles whereas, over longer time frames this
 378 relationship reverses (Fig 4A). Whether or not that reversal is apparent over the
 379 timeframe of the Holocene is unclear, however currently, greater Holocene peat
 380 thicknesses are typically reported from tropical regions (e.g. Anderson, 1983; Page et al
 381 1999)

5) At accommodation rates of between 0.1 and 0.2 mm per year, which are at the extreme low end of the range used by Bohacs and Suter (1997), 100 to 400ka of continuous peat accumulation can be accommodated with the height of peat above base level not exceeding 8 m (Fig 4B).

6) Temperate peatlands offer the maximum capacity for long-term carbon storage as they have the right balance of input and output to enable establishment and long-term growth above base level (Fig 4A and 4B).

7) The ratio of the accommodation rate to the peat production rate varies throughout the period of peat accumulation and bears no necessary relationship to the condition of the peatland (Fig 4A).

8) Given sufficient time and a constant accommodation rate all peat deposits will be terminated by inundation however if the accommodation rate is too low, or zero the model effectively permits indefinite peat accumulation above base level and this process must in some way be limited by erosion (Fig 4B).

2.5 Limits to peat accumulation.

Stratigraphic models assume that erosion and/or unfavourable hydrology rapidly limit peat growth in areas with no accommodation or falling base level. Over the short term, such as the period of the Holocene, the influence of landscape overcomes this and explains why extensive peat continues to accumulate in actively eroding landscapes with no increase in accommodation space. Furthermore, reported Holocene peat thickness, for example those in the Northern Hemisphere database of Loisel et al (2014), rarely exceed a thickness of 8 m so the influence of erosion on thicker accumulations above base level (Fig 4B) remains untested over the timeframe of the Holocene.

From observation of Holocene peat we deduce that 10 ka is generally insufficient time to observe significant limits to peat growth. So what processes ultimately limit peat accumulation in settings with little or no accommodation space on timescales much greater than 10 ka? The answer to this question is significant, as considerable quantities of Holocene peat occur in uplands or areas of active post-glacial uplift with limited or no accommodation space capacity.

Three key limiting processes are:

Hydrological and Oxidative limits – without accommodation the supply of water ultimately limits peat growth in any landscape setting. As peat growth approaches the hydrological limits of the system, oxidative decay above the water table increasingly limits carbon accumulation. This intrinsic limit in which the balance between productivity and decay determine an upper limit to carbon accumulation (Clymo, 1984) has been used to predict future long term limits ($>10^4$ years) to northern peatland carbon stocks. (Alexandrov et al 2019).

This approach is problematic for several reasons:

1. By necessity it assumes constant values for hydraulic conductivity, ignoring the poro-elastic response of the multiphase (solid, liquid, gas) peat matrix. For example, peat dry

bulk density in the saturated zone can vary from 0.02 to 0.2 g/cm³ (Large et al., 2009; Page et al., 2004) and peat hydraulic conductivity may vary by several orders of magnitude (Charman, 2002). Both of which illustrate that mechanical change in the pore structure of the peat can buffer the hydrology and create large uncertainty in model outputs.

2. It ignores the role of the hydrologic landscape.
3. Assuming some level of sustained input of atmospheric mineral dust at even quite low rates, the long-term consequences of oxidative limitation would be that the peat soil would transition into a mineral soil. For example, recent rates of dust deposition over peatland in Western Siberia range from 1 to 12 g/m²/yr (Fiałkiewicz-Kozieł, 2016), approximately 5-50% of the Holocene boreal long-term carbon accumulation rate (Loisel et al, 2014) before an oxidative limit has been reached. Some of this mass will be lost as soluble elements; however, most of the mass in the form of low solubility aluminosilicates, is retained (Large and Marshall, 2015).
4. Other physical and mechanic processes, discussed below, also act to limit peat growth and the relative importance of these processes over long periods is unknown.

Mechanical Deformation - An alternative not explored fully by peat scientists is that the stability of a peat body in the absence of accommodation space is mechanically limited and that the weak peat body will eventually undergo structural failure (mass movement), drainage and subsequent erosion even under sustained ideal conditions for peat formation. The inevitability of such mass movement has been noted and observed in specific blanket bog settings (Tallis, 1985) however, it is not generally considered as a limiting process in all settings in which accommodation space is limit or unavailable. The surface of the peat body may also wrinkle as proposed by Pearsall (1956) and observed in one mechanical simulation (Briggs et al., 2007), possibly into the characteristic patterns of hummocks and hollows observed in the field (Pearsall, 1956; Morris et al, 2013) and in some coal seams (Broadhurst and Simpson, 1983; Rippon, 1998). Such wrinkles could conceivably generate water channels leading to erosion, drainage and oxidation of a raised peatland under sustained wet conditions.

Advective erosion – The accumulation and growth of peat above base level is in-effect an increase in elevation, which from the perspective of erosion is no different to tectonic uplift. Observations and theories of landscape formation (e.g. Kirby and Whipple, 2012) require that an increase in elevation of the land must be constrained by erosion, particularly so in the typically wet climates associated with peat formation. Fluvial erosion has been studied in peatland (Li et al, 2019; Watters et al, 2007; Gradzinski et al 2003) but often from a short-term perspective of land management and carbon budgets (Li et al, 2019; Cowley et al, 2018; Evans et al, 2006)) rather than the perspective of long-term limits. This lack of consideration of significant natural erosion limits probably results from two aspects of the peatland erosion. First, the fibrous nature of peat is relatively resistant to erosion creating distinctive channel morphology and limiting channel movement (Watters et al, 2007; Gradzinski et al 2003). Second, it is difficult to quantify natural processes of erosion in highly modified (e.g. from the effects of over grazing and trampling) and managed landscapes. An additional and notable contribution to peat erosion, particularly in upland areas is aeolian deflation, which can rapidly occur following fluvial

incision (Foulds and Warburton, 2007). An ideal location in which to test the relative influence of erosion, mass movement and hydrology is the Falkland Islands. Although intensively grazed, a significant proportion of Falkland Island blanket peat initiated pre-Holocene (Payne et al, 2019), is highly dissected and in optical images and in the field shows ample evidence of established erosion and mass movement that must predate human occupation of the islands.

2.5 Hiatuses within peat and coal

In many interpretations, the lateral equivalent of stratigraphic boundaries generated on account of base level fluctuations within a subsiding basin are interpreted as intra-seam hiatuses in coal and vice versa. Evidenced by change in humification without notable siliciclastic deposition these hiatuses are interpreted to represent periods on the order of 10^4 to 10^6 years (Davies et al., 2005; Holdgate et al., 1995; Jerrett et al., 2011a). This is improbable. The generation of depositional breaks in peat requires sustained erosion or non-accumulation of the peat surface without accommodating clastic input or atmospheric dust. This is difficult, as peat will continue to lose mass due to on-going decay, in effect creating its own accommodation space. Furthermore, organic rich peat surfaces are naturally attractive sites for plant growth and, even in the absence of clastic deposition re-establishment of peat accumulation is likely on periods much less than 10^4 years. Evidence to support this are the short periods (about 10^2 to 10^3 years) that account for hiatuses in Holocene peat known as recurrence surfaces (Borgmark, 2005). The short duration of these time gaps is not surprising as even the loss of the top 1 m of a typical Holocene peatland need not result in a stratigraphic gap of more than 10^3 years (Borgmark, 2005; Page et al., 2004). Consequently, in a subsiding basin, it is improbable that intra-seam discontinuities without clastic deposition could represent long periods.

2.6 Principles of the new hydrologic landscape approach to the crustal storage of peat

Based on the above it is possible to define a set of considerations and related guidelines for the interpretation of peat and coal.

1. The hydrologic landscape determines the position, volume and extent of a peat body.
2. Thick peat forms in areas of prolonged hydrological stability.
3. The geometry, scale and persistence of the hydrological landscape unit determines the hydrological stability of a peatland.
4. In a given palaeo-geographic and palaeo-climate setting thicker coal seams accumulated over longer periods, not at a different rate.
5. The formation of coal requires clastic sediment bypass and hence storage of water in the precursor peat above the level of the equilibrium stream profile.
6. The thickness of a coal seam is proportional to the time integrated carbon accumulation rate.
7. Change into and out of periods of peat formation is a response to the change in the volume of water stored above base level.

8. Relative change in base level will modify the hydrologic landscape but does not preclude the capacity of the hydrologic landscape to maintain peat growth above base level.

3. Discussion

Application of these considerations and rules to the interpretation of peat and coal produces a very different understanding of the processes leading to the crustal storage of peat. It also places greater value on peat and coal as an indicator of the processes of landscape and basin evolution.

By far the biggest change in interpretation arises from the order of magnitude increase in time required to account for the carbon in coal. Why previous interpretations did not consider a carbon basis for time in coal is quite inexplicable, as the processes on which it is based are understood. The consequence that coal seams with thickness in the range 1 to 20 m must account for periods in the range 0.1 to 1 My, a range that overlaps falls within the periods of 3rd and 4th order sea level cycles (0.5-5 and 0.1-0.5 My respectively), requires greater significance to be placed on the coal record and the time therein. It also means that in many coal-bearing systems the greatest amount of time is contained within the coal not the clastic sediments. For example, when realistic estimates of time in coal are applied to the Duckmantian-Langsettian interval of the Carboniferous the time accounted for in the clastic sediments and coal increases from 75-125ky (Scott and Stephens, 2014) to 0.7-1.5 My (Large and Marshall, 2015). The former value requires huge unsubstantiated hiatuses and the latter lies within chronostratigraphic estimates for the duration of the interval, confirming the assertion of Scott and Stephens that most time must lie within the coal.

By increasing the amount of time, the role of landscape in sustaining a body of water-saturated peat above depositional base level becomes essential. This also sustains the necessary sediment bypass, which when combined with long duration peat formation must have a profound influence on the supply and distribution of sediment within a basin. Basinwards, the period of peat accumulation should be marked by enhanced sediment supply and associated progradation for example thick shoreface sands. The association between thick shoreface sand deposition and extensive coastal plain coal formation is seen within many coal-bearing paralic systems. In Spitsbergen, the Todalen coals occur adjacent to the coeval Endalen Member shoreface sediments (Marshall, 2013). Thick shoreface sediments/barrier island deposition and extensive coastal peat formation can also be observed within the Cretaceous South-Western Interior Basin, USA (Pederson and Dehler, 2005), the Cretaceous Blackhawk Formation, USA (Jerrett et al., 2011a) and associated with extensive lignite deposits from the Gippsland Basin, Australia (Birch and Division, 2003).

The non-linearity of peat accumulation rates contrasts markedly with the linear peat production rate assumed in the sequence stratigraphic models. Given a finite accommodation rate, this non-linearity requires that peat accumulation will terminate if given sufficient time. However more importantly the range of subsidence rates over which significant long term peat accumulation leading to crustal storage may occur is reduced to <0.5 mm/yr. This in turn demonstrates that for many coal seams there is sufficient time for low rates of basin subsidence alone to contribute to maintaining the hydrologic landscapes necessary to sustain peat deposition in appropriate tectonic settings. Such low subsidence rates are encountered in many tectonic

settings (Allen & Allen 2013) and can be associated with significant accumulations of coal. For example: On extensional continental margins subsidence rates range from 0.2 mm/yr at the initiation of rifting, decreasing to less than 0.05 mm/yr during the flexural subsidence phase, over time scales of 10^6 – 10^7 years (Allen and Allen, 2013). Coal bearing examples of such continental margin basins include the Gippsland Basin (Birch and Division, 2003). Foreland basins subside at rates of 0.2–0.5 mm/yr, and cratonic basins at 0.01–0.04 mm/yr. Coal bearing examples of these include the Cretaceous coal bearing western interior foreland basins of the USA (Pederson and Dehler, 2005) and the cratonic coal bearing basins of Australia the coals of which are also notably oxidised which could be a consequence of particularly low rates of subsidence (Hunt and Smyth, 1987).

A key problem of extending peat accumulation models over long periods of up to 10^6 years is that rates of decay and coalification process operating over these periods are unknown. So while there is a good knowledge of decay rates over periods <10ka and diagenetic coalification rates over periods greater than 10My there is nothing between. The only certainty is the non-linearity in the mass balance between inputs and outputs. Extension of a peat accumulation model over much longer periods is however, a good test of whether or not a given model is reasonable. If it cannot enable any reasonable thicknesses of peat to be accommodated over periods $\gg 10$ ka at reasonable accommodation rates then it is unlikely to be a reasonable model. A simple example of the application of this test is the constant decay model of Clymo et al 1998, which if extended to long periods using best fit Holocene values places an absolute and nonsensical limit of <30 cm on the total thickness of coal that could ever be.

A consequence of the long-term view is that in the absence of accommodation, there must be a limit to the quantity of peat a given landscape can hold. Therefore, an important question is how quickly, and over what spatial and temporal scales, erosion will limit the size of the peatland carbon reservoir in different landscapes. The answers to this could have profound consequences for current and future management of the peatland. For example, it is likely that areas of upland peat will reach their natural erosional limits faster than lowland raised bogs. Evidence of this is the eroded state of upland blanket bog some of which appears to predate human influences (Tallis, 1985). Although the effects of overgrazing may confound this judgement (Tallis, 1985), it can be argued that areas closer to their natural limits will be more vulnerable to the effects of overgrazing and other environmental pressures. The inevitability of an erosional limit in landscapes where accommodation space is not being created also requires that we account for erosional losses when attributing a carbon flux to an area of peatland. It may also imply that in landscapes without the capacity to accommodate long-term accumulation of peat that the long-term management strategy should be to minimise rates of carbon loss and sustain vegetation cover rather than promote metastable growth above the natural limits imposed by erosion and mass movement. In terms of global carbon budgets, we can also question whether it is possible to have a significantly larger peatland carbon reservoir. So for example would it be possible to double the size of the peatland carbon reservoir within the current area of peatland and if not what would be the limiting processes?

The hydrological landscape requirements for sustained accumulation of peat mean that coal seam thickness, as deduced by Broadhurst and France, (1986) and Broadhurst and Simpson (1983), will be a sensitive indicator of syn-depositional deformation and landscape evolution within a basin. Using the approach above in combination with isopach maps it is possible to use

the variations in coal thickness to map the evolution of active structures and landscape within a basin. A specific example of this is the detailed work of Marshall (2013) who demonstrated the role of persistent structures in controlling landscape and the distribution and thickness of early Palaeocene coal within the Central Tertiary Basin, Svalbard. There are also numerous examples (e.g. Ferm and Staub, 1985; Read, 1989; Guion, 1987; Haszeldine, 1989; Greb et al 2001) linking coal thickness to syndepositional topography, differential compaction, and tectonics. However, they have not been interpreted in the context of a hydrological landscape over an appropriate timeframe. To fully demonstrate the application of this approach would require a complete re-analysis of data from a specific basin, which is beyond the scope of this paper.

5 Conclusions

To account for the crustal carbon contained within peat, coal and lignite requires considerably longer periods than previously considered. In turn, this requires consideration of the parameters of the hydrologic landscape that can sustain the conditions suitable for peat formation. This approach brings coal stratigraphic models in line with modern peat observations and can be applied to all peatland and coal settings on various scales. Furthermore this approach sets limits to the interpretation of the parameters of the hydrological landscape that enable long term accumulations of peat leading to coal. For example it casts severe doubts on the notion that thick peat accumulations form in areas of rapid subsidence. It also requires far lower and less variable accommodation rates to take peat into long term crustal storage and demonstrates that for many coal seams there is sufficient time for crustal deformation alone to sustain the necessary hydrologic landscapes.

The common stratigraphic assumption that coals are largely ephemeral features within a siliciclastic framework is a barrier to understanding environmental change, basin processes and the long-term storage of peatland carbon. The sustained periods of peat formation needed to account for the carbon in coal require recognition of a prolonged and fundamental shift in the syndepositional environment and processes of sediment transport and deposition both of which have basinwide consequences. The hydrologic landscape approach facilitates re-interpretation of these systems. It also raises questions as to the long-term validity of Holocene peat growth models and the ultimate limits to peat accumulation.

Acknowledgments

The authors have no conflict of interest. This work was jointly funded by the Faculty of Engineering University of Nottingham, a Natural Environment Research Council Studentship to Chris Marshall and the Store Norske Spitsbergen Kulkompani AS. The paper does not contain new data. The University Centre in Svalbard (UNIS) is thanked for providing office space, expertise and library access during research visits to Svalbard. M Waddington and R S Clymo provided are thanked for providing comments on the manuscript.

References

- Allen, P.A. & Allen, J.R. (2013). Basin Analysis: Principles and Application to Petroleum Play Assessment. 3rd Edition, Wiley-Blackwell p.632. ISBN: 978-0-470-67377-5.
- Alpern, B., & de Sousa, M. J. L. (2002). Documented international enquiry on solid sedimentary fossil fuels; coal: definitions, classifications, reserves-resources, and energy potential.

- International Journal of Coal Geology*, 50(1-4), 3-41. [https://doi.org/10.1016/S0166-5162\(02\)00112-X](https://doi.org/10.1016/S0166-5162(02)00112-X)
- Alexandrov, G. A., Brovkin, V. A., Kleinen, T., & Yu, Z. (2020). The capacity of northern peatlands for long-term carbon sequestration. *Biogeosciences*, 17(1), 47-54. <https://www.biogeosciences.net/17/47/2020/>
- Anderson, J. A. R. (1983). The tropical peat swamps of western Malesia. *Mires : swamp, bog, fen and moor : regional studies*. <https://ci.nii.ac.jp/naid/10019557618/en/>
- Baird, A. J., Morris, P. J., & Belyea, L. R. (2012). The DigiBog peatland development model 1: rationale, conceptual model, and hydrological basis. *Ecohydrology*, 5(3), 242-255. <https://doi.org/10.1002/eco.230>
- Birch, W. D., (ed) (2003). *Geology of Victoria*: Geological Society of Australia (Victoria Division). ISBN: 9781876125332
- Bohacs, K., & Suter, J. (1997). Sequence stratigraphic distribution of coaly rocks: Fundamental controls and paralic examples. *Aapg Bulletin-American Association of Petroleum Geologists*, 81(10), 1612-1639. <https://doi.org/10.1306/3B05C3FC-172A-11D7-8645000102C1865D>
- Borgmark, A. (2005). Holocene climate variability and periodicities in south-central Sweden, as interpreted from peat humification analysis. *Holocene*, 15(3), 387-395. <https://doi.org/10.1191/0959683605hl816rp>
- Briggs, J., Large, D. J., Snape, C., Drage, T., Whittles, D., Cooper, M., et al. (2007). Influence of climate and hydrology on carbon in an early Miocene peatland. *Earth and Planetary Science Letters*, 253(3-4), 445-454. <https://doi.org/10.1016/j.epsl.2006.11.010>
- Broadhurst, F. M., & France, A. A. (1986). Time Represented by Coal Seams in the Coal Measures of England. *International Journal of Coal Geology*, 6(1), 43-54. [https://doi.org/10.1016/0166-5162\(86\)90024-8](https://doi.org/10.1016/0166-5162(86)90024-8)
- Broadhurst, F. M., & Magraw, D. (1961). On a fossil tree found in an opencast coal site near Wigan, Lancashire. *Geological Journal*, 2(2), 155-158. <https://doi.org/10.1002/gj.3350020203>
- Broadhurst, F. M., & Simpson, I. M. (1983). Syntectonic Sedimentation, Rigs, and Fault Reactivation in the Coal Measures of Britain. *Journal of Geology*, 91(3), 330-337. <https://doi.org/10.1086/628775>
- Catuneanu, O. (2006). *Principles of Sequence Stratigraphy*. Elsevier Science. ISBN: 9780080473987
- Charman, D. (2002). *Peatlands and environmental change*. John Wiley and Sons, Chichester. ISBN: 978-0-470-84410-6
- Clymo, R. S. (1984). The Limits to Peat Bog Growth. *Philosophical Transactions of the Royal Society of London Series B-Biological Sciences*, 303(1117), 605-654. <https://doi.org/10.1098/rstb.1984.0002>
- Clymo, R. S., Turunen, J., & Tolonen, K. (1998). Carbon accumulation in peatland. *Oikos*, 81(2), 368-388. <https://doi.org/10.2307/3547057>
- Cowley, K., Looman, A., Maher, D. T., & Fryirs, K. (2018). Geomorphic controls on fluvial carbon exports and emissions from upland swamps in eastern Australia. *Science Of The Total Environment*, 618, 765-776. <https://doi.org/10.1016/j.scitotenv.2017.08.133>
- Dai, S., Bechtel, A., Eble, C. F., Flores, R. M., French, D., Graham, I. T., et al. (2020). Recognition of peat depositional environments in coal: A review. *International Journal of Coal Geology*, 219, 103383. <https://doi.org/10.1016/j.coal.2019.103383>

- Davies, R., Diessel, C., Howell, J., Flint, S., & Boyd, R. (2005). Vertical and lateral variation in the petrography of the Upper Cretaceous Sunnyside coal of eastern Utah, USA - implications for the recognition of high-resolution accommodation changes in paralic coal seams. *International Journal of Coal Geology*, 61(1-2), 13-33. <https://doi.org/10.1016/j.coal.2004.06.003>
- Diessel, C., Boyd, R., Wadsworth, J., Leckie, D., & Chalmers, G. (2000). On balanced and unbalanced accommodation/peat accumulation ratios in the Cretaceous coals from Gates Formation, Western Canada, and their sequence-stratigraphic significance. *International Journal of Coal Geology*, 43(1-4), 143-186. [https://doi.org/10.1016/S0166-5162\(99\)00058-0](https://doi.org/10.1016/S0166-5162(99)00058-0)
- Diessel, C.F.K. (1992). Coal bearing depositional systems. Springer, Berlin, 721 pp. <https://doi.org/10.1007/978-3-642-75668-9>
- Einsele, G. (2000). Sedimentary Basins: evolution, facies, and sediment budget, 2nd Edition. Springer-Verlag, Berlin. <https://doi.org/10.1007/978-3-662-04029-4>
- Evans, M., Warburton, J., & Yang, J. (2006). Eroding blanket peat catchments: Global and local implications of upland organic sediment budgets. *Geomorphology*, 79(1-2), 45-57. <https://doi.org/10.1016/j.geomorph.2005.09.015>
- Falcon-Lang, H. J. (2006). A history of research at the Joggins Fossil Cliffs of Nova Scotia, Canada, the world's finest Pennsylvanian section. *Proceedings of the Geologists Association*, 117, 377-392. [https://doi.org/10.1016/S0016-7878\(06\)80044-1](https://doi.org/10.1016/S0016-7878(06)80044-1)
- Fialkiewicz-Koziel, B., Smieja-Krol, B., Frontasyeva, M., Slowinski, M., Marcisz, K., Lapshina, E., et al. (2016). Anthropogenic- and natural sources of dust in peatland during the Anthropocene. *Sci Rep*, 6(1), 38731. <https://doi.org/10.1038/srep38731>
- Ferm, J. C., & Staub, J. R. (1985). Depositional Controls of Mineable Coal Bodies. *Sedimentology of Coal and Coal-Bearing Sequences*, 273-289. <https://doi.org/10.1002/9781444303797.ch15>
- Foulds, S. A., & Warburton, J. (2007). Wind erosion of blanket peat during a short period of surface desiccation (North Pennines, Northern England). *Earth Surface Processes and Landforms*, 32(3), 481-488. <https://doi.org/10.1002/esp.1422>
- Frolking, S., Roulet, N. T., Tuittila, E., Bubier, J. L., Quillet, A., Talbot, J., & Richard, P. J. H. (2010). A new model of Holocene peatland net primary production, decomposition, water balance, and peat accumulation. *Earth System Dynamics*, 1(1), 1-21. <https://doi.org/10.5194/esd-1-1-2010>
- Glaser, P. H., Siegel, D. I., Reeve, A. S., Janssens, J. A., & Janecky, D. R. (2004). Tectonic drivers for vegetation patterning and landscape evolution in the Albany River region of the Hudson Bay Lowlands. *Journal of Ecology*, 92(6), 1054-U1052. <https://doi.org/10.1111/j.0022-0477.2004.00930.x>
- Glaser, P. H., Siegel, D. I., Romanowicz, E. A., & Shen, Y. P. (1997). Regional linkages between raised bogs and the climate, groundwater, and landscape of north-western Minnesota. *Journal of Ecology*, 85(1), 3-16. <https://doi.org/10.2307/2960623>
- Gradzinski, R., Baryla, J., Doktor, M., Gmur, D., Gradzinski, M., Kedzior, A., et al. (2003). Vegetation-controlled modern anastomosing system of the upper Narew River (NE Poland) and its sediments. *Sedimentary Geology*, 157(3-4), 253-276. [https://doi.org/10.1016/S0037-0738\(02\)00236-1](https://doi.org/10.1016/S0037-0738(02)00236-1)
- Greb, S. F., Eble, C. F., Williams, D. A., & Nelson, W. J. (2001). Dips, ramps, and rolls - evidence for paleotopographic and syn-depositional fault control on the Western

- Kentucky No. 4 coal bed, Tradewater Formation (Bolsovian), Illinois Basin.
International Journal of Coal Geology, 45(4), 227-246. [https://doi.org/10.1016/S0166-5162\(00\)00023-9](https://doi.org/10.1016/S0166-5162(00)00023-9)
- Guion, P. D. (1987). The influence of a palaeochannel on seam thickness in the coal measures of Derbyshire, England. *International Journal of Coal Geology*, 7(3), 269-299. [https://doi.org/10.1016/0166-5162\(87\)90040-1](https://doi.org/10.1016/0166-5162(87)90040-1)
- Haszeldine, R. S. (1989). Coal reviewed: depositional controls, modern analogues and ancient climates. *Geological Society, London, Special Publications*, 41(1), 289. <https://doi.org/10.1144/GSL.SP.1989.041.01.20>
- Hokanson, K. J., Lukenbach, M. C., Devito, K. J., Kettridge, N., Petrone, R. M., and Waddington, J. M. (2016) Groundwater connectivity controls peat burn severity in the boreal plains. *Ecohydrol.*, 9: 574– 584. <https://doi.org/10.1002/eco.1657>.
- Holdgate, G. R., Kershaw, A. P., & Sluiter, I. R. K. (1995). Sequence stratigraphic analysis and the origins of Tertiary brown coal lithotypes, Latrobe Valley, Gippsland Basin, Australia. *International Journal of Coal Geology*, 28(2-4), 249-275. [https://doi.org/10.1016/0166-5162\(95\)00020-8](https://doi.org/10.1016/0166-5162(95)00020-8)
- Holz, M., Kalkreuth, W., & Banerjee, I. (2002). Sequence stratigraphy of paralic coal-bearing strata: an overview. *International Journal of Coal Geology*, 48(3-4), 147-179. [https://doi.org/10.1016/S0166-5162\(01\)00056-8](https://doi.org/10.1016/S0166-5162(01)00056-8)
- Hunt, J.W. & Smyth, M. (1989). Origin of inertinite-rich coals of Australian cratonic basins. *International Journal of Coal Geology*, 11 (1), 23-46, [https://doi.org/10.1016/0166-5162\(89\)90111-0](https://doi.org/10.1016/0166-5162(89)90111-0)
- Jackson, J.A. (1997) Glossary of geology: American Geological Institute, Fourth edition, 769 p.
- Jerrett, R. M., Davies, R. C., Hodgson, D. M., Flint, S. S., & Chiverrell, R. C. (2011a). The significance of hiatal surfaces in coal seams. *Journal of the Geological Society*, 168(3), 629-632. <https://doi.org/10.1144/0016-76492010-178>
- Jerrett, R. M., Flint, S. S., Davies, R. C., & Hodgson, D. M. (2011b). Sequence stratigraphic interpretation of a Pennsylvanian (Upper Carboniferous) coal from the central Appalachian Basin, USA. *Sedimentology*, 58(5), 1180-1207. <https://doi.org/10.1111/j.1365-3091.2010.01200.x>
- Kirby, E., & Whipple, K. X. (2012). Expression of active tectonics in erosional landscapes. *Journal of Structural Geology*, 44, 54-75. <https://doi.org/10.1016/j.jsg.2012.07.009>
- Large, D. J. (2007). A 1.16 Ma record of carbon accumulation in western European peatland during the Oligocene from the Ballymoney lignite, Northern Ireland. *Journal of the Geological Society*, 164, 1233-1240. <https://doi.org/10.1144/0016-76492006-148>
- Large, D. J., Jones, T. F., Briggs, J., Macquaker, J. H. S., & Spiro, B. F. (2004). Orbital tuning and correlation of 1.7 my of continuous carbon storage in an early Miocene peatland. *Geology*, 32(10), 873-876. <https://doi.org/10.1130/G20824.1>
- Large, D. J., & Marshall, C. (2015). Use of carbon accumulation rates to estimate the duration of coal seams and the influence of atmospheric dust deposition on coal composition. *Geological Society, London, Special Publications*, 404(1), 303-315. <https://doi.org/10.1144/sp404.15>
- Large, D. J., Spiro, B., Ferrat, M., Shopland, M., Kylander, M., Gallagher, K., et al. (2009). The influence of climate, hydrology and permafrost on Holocene peat accumulation at 3500 m on the eastern Qinghai-Tibetan Plateau. *Quaternary Science Reviews*, 28(27-28), 3303-3314. <https://doi.org/10.1016/j.quascirev.2009.09.006>

- Li, P. F., Irvine, B., Holden, J., & Mu, X. M. (2017). Spatial variability of fluvial blanket peat erosion rates for the 21st Century modelled using PESERA-PEAT. *Catena*, 150, 302-316. <https://doi.org/10.1016/j.catena.2016.11.025>
- Loisel, J., Yu, Z. C., Beilman, D. W., Camill, P., Alm, J., Amesbury, M. J., et al. (2014). A database and synthesis of northern peatland soil properties and Holocene carbon and nitrogen accumulation. *Holocene*, 24(9), 1028-1042. <https://doi.org/10.1177/0959683614538073>
- Marshall, C., Large, D. J., & Heavens, N. G. (2016). Coal-derived rates of atmospheric dust deposition during the Permian. *Gondwana Research*, 31, 20-29. <https://doi.org/10.1016/j.gr.2015.10.002>
- Marshall, C. (2013). *Palaeogeographic development and economic potential of the coal-bearing palaeocene Todalen Member, Spitsbergen*. (PhD), University of Nottingham. Retrieved from <http://eprints.nottingham.ac.uk/id/eprint/13794>
- Michaelsen, P., Henderson, R. A., Crosdale, P. J., & Mikkelsen, S. O. (2000). Facies architecture and depositional dynamics of the Upper Permian Rangal Coal Measures, Bowen Basin, Australia. *Journal of Sedimentary Research*, 70(4), 879-895. <https://doi.org/10.1306/2dc4093f-0e47-11d7-8643000102c1865d>
- Morris, P. J., Baird, A. J., & Belyea, L. R. (2012). The DigiBog peatland development model 2: ecohydrological simulations in 2D. *Ecohydrology*, 5(3), 256-268. <https://doi.org/10.1002/eco.229>
- Morris, P. J., Baird, A. J., & Belyea, L. R. (2013). The role of hydrological transience in peatland pattern formation. *Earth Surface Dynamics*, 1(1), 29-43. <https://doi.org/10.5194/esurf-1-29-2013>
- Page, S. E., Rieley, J. O., Shotyk, W., & Weiss, D. (1999). Interdependence of peat and vegetation in a tropical peat swamp forest. *Philosophical transactions of the Royal Society of London. Series B, Biological sciences*, 354(1391), 1885-1897. <https://doi.org/10.1098/rstb.1999.0529>
- Page, S. E., Wust, R. A. J., Weiss, D., Rieley, J. O., Shotyk, W., & Limin, S. H. (2004). A record of Late Pleistocene and Holocene carbon accumulation and climate change from an equatorial peat bog (Kalimantan, Indonesia): implications for past, present and future carbon dynamics. *Journal Of Quaternary Science*, 19(7), 625-635. <https://doi.org/10.1002/jqs.884>
- Payne, R. J., Ring-Hrubesh, F., Rush, G., Sloan, T. J., Evans, C. D., & Mauquoy, D. (2019). Peatland initiation and carbon accumulation in the Falkland Islands. *Quaternary Science Reviews*, 212, 213-218. <https://doi.org/10.1016/j.quascirev.2019.03.022>
- Pearsall, W. (1956). Two Blanket-Bogs in Sutherland. *Journal of Ecology*, 44(2), 493-516. doi:10.2307/2256833
- Pederson, J.L. & Dehler, C.M. (2005). Interior Western United States. Geological Society of America Field Guide. Geological Society of America. <https://doi.org/10.1130/0-8137-0006-X>
- Read, W. A. (1989). The influence of basin subsidence and depositional environment on regional patterns of coal thickness within the Namurian fluvio-deltaic sedimentary fill of the Kincardine Basin, Scotland. *Geological Society, London, Special Publications*, 41(1), 333. <https://doi.org/10.1144/GSL.SP.1989.041.01.23>
- Railsback L.B. (1995) Controls on Long-Term Global Rates of Coal Deposition, and the Link between Eustasy and Global Geochemistry. In: Haq B.U. (eds) Sequence

- Stratigraphy and Depositional Response to Eustatic, Tectonic and Climatic Forcing. Coastal Systems and Continental Margins, vol 1. Springer, Dordrecht
<https://doi.org/10.1007/978-94-015-8583-5>
- Rippon, J. H. (1998). The identification of syn-depositionally-active structures in the coal-bearing Upper Carboniferous of Great Britain. *Proceedings of the Yorkshire Geological Society*, 52(1), 73-93. <https://doi.org/10.1144/pygs.52.1.73>
- Schopf, J. M. (1956). A definition of coal. *Economic Geology*, 51(6), 521-527.
<https://doi.org/10.2113/gsecongeo.51.6.521>
- Schopf, J. M. (1966). Definitions of Peat and Coal and of Graphite That Terminates the Coal Series (Graphocite). *The Journal of Geology*, 74(5), 584-592.
- Scott, A. C., & Stephens, R. S. (2015). British Pennsylvanian (Carboniferous) coal-bearing sequences: where is the time? *Geological Society, London, Special Publications*, 404(1), 283-302. <https://doi.org/10.1144/sp404.14>
- Staub, J. R. (2002). Marine flooding events and coal bed sequence architecture in southern West Virginia. *International Journal of Coal Geology*, 49(2-3), 123-145.
[https://doi.org/10.1016/S0166-5162\(01\)00074-X](https://doi.org/10.1016/S0166-5162(01)00074-X)
- Tallis, J.H. (1985). Mass Movement and Erosion of a Southern Pennine Blanket Peat. *Journal of Ecology*, 73(1), 283-315. <https://doi.org/10.2307/2259784>
- Titheridge, D. G. (1993). The Influence of Half-Graben Syn-Depositional Tilting on Thickness Variation and Seam Splitting in the Brunner Coal Measures, New-Zealand. *Sedimentary Geology*, 87(3-4), 195-213. [https://doi.org/10.1016/0037-0738\(93\)90004-O](https://doi.org/10.1016/0037-0738(93)90004-O)
- Wadsworth, J., Diessel, C., & Boyd, R. (2010). The Sequence Stratigraphic Significance of Paralic Coal and Its Use as an Indicator of Accommodation Space in Terrestrial Sediments. *Application of Modern Stratigraphic Techniques: Theory and Case Histories*, 94, 201-219. <https://doi.org/10.2110/sepmssp.094.201>
- Watters, J. R., & Stanley, E. H. (2007). Stream channels in peatlands: The role of biological processes in controlling channel form. *Geomorphology*, 89(1-2), 97-110.
<https://doi.org/10.1016/j.geomorph.2006.07.015>
- Winston, R. B. (1994). Models of the Geomorphology, Hydrology, and Development of Domed Peat Bodies. *Geological Society of America Bulletin*, 106(12), 1594-1604.
[https://doi.org/10.1130/0016-7606\(1994\)106<1594:Motgha>2.3.Co;2](https://doi.org/10.1130/0016-7606(1994)106<1594:Motgha>2.3.Co;2)
- Winter, T. C. (2000). The vulnerability of wetlands to climate change: A hydrologic landscape perspective. *Journal of the American Water Resources Association*, 36(2), 305-311.
<https://doi.org/10.1111/j.1752-1688.2000.tb04269.x>
- Winter, T. C. (2001). The concept of hydrologic landscapes. *Journal of the American Water Resources Association*, 37(2), 335-349. <https://doi.org/10.1111/j.1752-1688.2001.tb00973.x>
- Yule, C. M., & Gomez, L. N. (2009). Leaf litter decomposition in a tropical peat swamp forest in Peninsular Malaysia. *Wetlands Ecology and Management*, 17, 231-241.
<https://doi.org/10.1007/s11273-008-9103-9>