

Turbulent mixing in the Indonesian throughflow exit passages, the Lesser Sunda waters

Turbulent mixing in the Indonesian throughflow exit passages, the Lesser Sunda waters
Adi Purwandana and Mochamad Furqon Azis Ismail
Research Center for Oceanography - Indonesian Institute of Sciences (RCO-LIPI)
Jl. Pasir Putih I Ancol Timur, Jakarta 14430, Indonesia
Corresponding author : adip001@lipi.go.id

Background
As the only low latitude passage connecting Pacific and Indian Oceans, Indonesian archipelago is an important role in controlling global ocean circulation, i.e. thermohaline circulation, through the input of warm, low salinity anomaly, that is eventually transported poleward out of the Indian Ocean within the Agulhas current system. Previous studies indicated that the existence of small eddies and narrow passages in the Indonesian seas will potentially mix the Pacific water mass with local Indonesian freshwater, before entering Indian Ocean.

Methods
We optimize the using of classical Thorpe method to infer TKE dissipation rate values from CTD profiles. This method was proved in a good agreement with microstructure measurement. The TKE dissipation rate is expressed as:
$$\epsilon = \frac{g}{N^2} \frac{d^2 \rho}{dz^2}$$

where ϵ is Thorpe length scale, defined as an rms

Results and Discussion
Water masses
The signature of Pacific water masses is highly evident in the Lesser Sunda waters. Our 63 grid averaging maps of salinity also indicate some fresher layers are observed in some straits in the Lesser Sunda waters. These are in the Lombok Strait (11° 27' S, 115° 0' E), the Sunda Strait (1° 12' S, 105° 42' E), and the straits of four straits (1° 22' S, 105° 42' E).
Consequently, there was increasing salinity of 0.1-0.2 in intermediate layer. It is likely due to intrusion of North Indian intermediate water (NINW).

TKE Dissipation Rate (ϵ)
The Lesser Sunda waters is characterized by strong dissipation rate. Very strong dissipation rate of greater than 10 $\text{m}^2 \text{s}^{-3}$ is observed in the Lombok Strait, Sunda Strait north of Sunda Strait, Sunda Strait and Sunda Strait.

Concluding Remarks
• A careful handling of CTD dataset by means of improved classical Thorpe method is a promising approach to map TKE dissipation rates and the location of direct measurements in the Indonesian seas.
• The Lesser Sunda waters with their narrow passages and shallow straits was proved to be an effective mixing hotspot of the IOT before entering the Indian Ocean.
• The dissipation rate inferred from high-resolution 10 m/20 m based by 60 m/s is consistent with our observational estimates in the hot spots of strong

References
1. Ahmadpoora, A., Whitard, R., Marlet, G., Wilkins, J., Spence, J., Kuhlmann, R., Jara, I., Supangat, A., 2006. Characteristics and variability of the Indonesian throughflow water at the surface straits. Deep Sea Res. Part I Oceanogr. Res. 53, 1942-1954. doi:10.1016/j.dsr.2009.08.004
2. Roussel-Labrousse, P., Goussin, Y., Fournier, S., Desmet, D., Mouton, C., Ahmadpoora, A., Jara, I., 2014. Contrasted turbulence intensities in the Indonesian Throughflow: a challenge for parameterizing energy dissipation rate. Deep Sea Res. Part I Oceanogr. Res. 100, 1-15. doi:10.1016/j.dsr.2014.04.015


Author Information | **Abstract** | **Contact Author** | **Get iPoster**

Adi Purwandana and Mochamad Furqon Azis Ismail

Research Center for Oceanography - Indonesian Institute of Sciences (RCO-LIPI)

Jl. Pasir Putih I Ancol Timur, Jakarta 14430, Indonesia

Corresponding author : adip001@lipi.go.id



PRESENTED AT:



JpGU - AGU Joint Meeting 2020

For a Borderless World of Geoscience
Japan Geoscience Union, American Geophysical Union

BACKGROUND

As the only low latitude passage connecting Pacific and Indian Oceans, Indonesian seas via its ITF (Indonesian throughflow) has an important role in controlling global ocean circulations, i.e thermohaline circulation, through the input of warm, low salinity anomaly, that is eventually transported poleward out of the Indian Ocean within the Agulhas current system.

Previous studies indicated that the existence of small straits and narrow passages in the Indonesian seas will potentially mix the Pacific water mass with local Indonesian fresher water, before entering Indian Ocean.

Here, we present the last mixing hotspot for the ITF before entering the Indian Ocean, i.e the Lesser Sunda waters which is the exit passages of the ITF.

Lack of direct turbulence measurements in the Indonesian seas has led us to apply an indirect Thorpe scale method to infer dissipation rates and diapycnal diffusivities using CTD profiles obtained from annual RCO-LIPI cruises.

METHODS

We optimize the using of classical Thorpe method to infer TKE dissipation rate values from CTD profiles. The method was proved in a good agreement with microstructure measurement. The TKE dissipation rate is expressed as,

$$\varepsilon_{Th-GM} = \begin{cases} 0.64L_T^2 N^3, & \text{when there is overturn} \\ \max\left(1 \times 10^{-10}, \varepsilon_0 \left(\frac{N^2}{N_0^2}\right)\right), & \text{when no overturn observed} \end{cases}$$

L_T is Thorpe length scale, defined as an rms value of vertical displacement for each turbulent patch to gain gravitationally stable density profile, N is buoyancy frequency, 1×10^{-10} is the lowest dissipation rate, defined based on the typical minimum dissipation rate observed by microstructure measurements, $\varepsilon_0 = 7 \times 10^{-10} \text{ m}^2 \text{ s}^{-3}$ and $N_0 = 3 \text{ cph}$ are the canonical Garret and Munk dissipation rate and respected buoyancy frequency reference, respectively. The final dissipation rate termed as ε_{Th-GM} will give a continuous profile of the dissipation rate as opposed to the standard Thorpe method.

Once the dissipation rate is estimated, turbulent diffusivity can be determined as,

$$K_{\rho Th-GM} = \Gamma \frac{\varepsilon_{Th-GM}}{N^2}$$


To inspect the spatial variability of hydrography and turbulent kinetic energy dissipation rates which is related to the potency of mixing events, we built maps by gridding horizontally the data over 0.5° and by vertical averaging over several depth intervals of 50-300 m, 300-500 m, 500-800 m, 800-1000 m, which represent the upper thermocline, lower thermocline, intermediate layer, and deeper layer.

RESULTS AND DISCUSSION

Water masses


The signature of Pacific water masses is highly eroded in the Lesser Sunda waters. Our 0.5° grid averaging maps of salinity also indicate some fresher layers are observed in some straits in the Lesser Sunda waters, those are in the Lombok Strait (115° - 117° E; 8° - 9° S), the Sape Strait (119° - 120° E; 8° - 9° S), and the straits of Alor islands (123° - 126° E; 8° - 9° S) at $\sigma_\theta = 22$ - 25 ; illustrating the incorporation of fresher surface water in the thermocline by the strong mixing in these spots.

Conversely, there was increasing salinity at $\sigma_\theta = 26$ - 27 or intermediate layer. It is likely due to intrusion of North Indian Intermediate water (NIIW).




TKE Dissipation Rate (ϵ)

The Lesser Sunda waters is characterized by strong dissipation rate. Very strong dissipation rate of greater than $10^{-7} \text{ m}^2 \text{ s}^{-1}$ is observed in the Lombok Strait, Sape Strait (north of Sumba Strait), Ombai Strait and Timor passage.




These strong TKE dissipation rates will potentially reduce the Pacific thermocline layer (North Pacific Subtropical Water, NPSW) signature.




Vertical Eddy Diffusivity (K_p)

We found an increasing value of K_p with depth due to the decrease of stratification at depth. Large vertical diffusivities greater than $10^{-3} \text{ m}^2 \text{s}^{-1}$ are observed in Ombai Strait.



The mean values are above typical open ocean value of $1 \times 10^{-5} \text{ m}^2 \text{s}^{-1}$. The median of the distribution shifts toward higher values with increasing depth as a result of the decreasing stratification. The highest values were observed in the deeper layer [800-2000] m with a mean and a median of $1.6 \times 10^{-3} \text{ m}^2 \text{s}^{-1}$ and $2.7 \times 10^{-4} \text{ m}^2 \text{s}^{-1}$, respectively.



Comparison with Model

We compared our observation based TKE dissipation rate with Nagai and Hibiya (2015) 3D Hydrostatic MITGC model. In both model and observations, the same trends are observed where the dissipation rates are relatively high in this region.

The ratio between the gridded observed dissipation rates and the gridded model-based estimates remains below an order of ten.

CONCLUDING REMARKS

- A careful handling of CTD dataset by means of improved classical Thorpe method is a promising approach to map TKE dissipation rates amid the lacking of direct measurements in the Indonesian seas.
- The Lesser Sunda waters with their narrow passages and shallow straits was proved to be an effective mixing hotspot of the ITF before entering the Indian Ocean.
- The dissipation rate inferred from high-resolution 3D MITGCM forced by M2 tides is consistent with our observational estimates in the hot spots of strong turbulence, suggesting the baroclinic tidal energy dissipation is the main driving mechanisms working in this area.
- Direct measurements using microstructure profiler is needed in the future to validate this indirect Thorpe method.

REFERENCES

1. Atmadipoera, A., Molcard, R., Madec, G., Wijffels, S., Sprintall, J., Koch-Larrouy, A., Jaya, I., Supangat, A., 2009. Characteristics and variability of the Indonesian throughflow water at the outflow straits. *Deep. Res. Part I Oceanogr. Res. Pap.* 56, 1942–1954. doi:10.1016/j.dsr.2009.06.004
2. Bouruet-Aubertot, P., Cuypers, Y., Ferron, B., Dausse, D., Ménage, O., Atmadipoera, A., Jaya, I., Olivier, M., Atmadipoera, A., Jaya, I., 2018. Contrasted turbulence intensities in the Indonesian Throughflow: a challenge for parameterizing energy dissipation rate. *Ocean Dyn.* 68, 1–75. doi:10.1007/s10236-018-1159-3
3. Garrett, C., Munk, W., 1975. Space-time scales of internal waves: A progress report. *J. Geophys. Res.* 80, 291–297. doi:10.1029/JC080i003p0
4. Nagai, T., Hibiya, T., 2015. Internal tides and associated vertical mixing in the Indonesian Archipelago. *J. Geophys. Res. C Ocean.* 3373–3390. doi:10.1002/2014JC010592
5. Purwandana, A., Cuypers, Y., Bouruet-Aubertot, P., Nagai, T., Hibiya, T., Atmadipoera, A.S., 2020. Spatial structure of turbulent mixing inferred from historical CTD datasets in the Indonesian seas. *Prog. Oceanogr.* 102312. doi:https://doi.org/10.1016/j.pocean.2020.102312