

Introduction

- Thunderstorms are characterised by lightning, thunder, rain showers, prevailing gusts, and hail, resulting in severe weather and the loss of lives, property, and farmland. Thunderstorms are the most destructive occurrence across eastern India during the pre-monsoon season (March-May) and are often pushed from the north-west to the south-east (Yamane and Hayashi, 2006; Tyagi, 2011; Das et al., 2014). Thunderstorm forecasting is required to avoid such problems.
- Several research have been conducted over the last century to better understand the various thunderstorm characteristics (Mukhopadhyay et al., 2005; Ghosh et al. 2008). Many studies have been published in which thunderstorm stability properties and associated thermodynamic indices have been studied (Schultz 1989; Kunz 2007; Tyagi et al. 2011; Sahu et al., 2020a; Sahu et al., 2020b). Many academics have investigated the operational effectiveness of various stability indices for thunderstorm prediction (Jacovides and Yonetani 1990; Lee and Passner 1993).
- The atmospheric static stability is measured via stability indices (Peppier 1988). The indices were computed using changes in dew-point temperature, air temperature at various levels, moisture, and dynamic driving forces in terms of wind shear. To predict the occurrence of thunderstorms, weather scientists employ thermodynamic characteristics and skill score analysis.
- The current research work employs different thermodynamic indices for spatial and temporal variations: the Boyden index (BOYD), Convective available potential energy (CAPE), Convective inhibition (CIN), Cross Totals index (CTI), Humidity index (HI), K index (KI), Lifted index (LI), Severe weather threat index (SWEAT), Total totals index (TTI), and Vertical totals index (VTI) over the eastern India.
- For temporal variations, we proposed suitable threshold values of these indices, and to identified some indices which performs best in predicting thunderstorm activity across the eastern India during the Pre-monsoon season for the study period.
- For Spatial variation we prepared composite maps to see the variations of these thermodynamic indices region wise for both thunderstorm and non-thunderstorm days and also predicted their trend. Here we shown the CAPE and CIN fluctuations and their trends to understand influence of climate change in changing the thunderstorm frequency and intensity.

Data and Methodology

- In this work we utilized radiosonde observations which is obtained from the University of Wyoming for temporal variation and ERA5 reanalysis datasets produced from European Centre for Medium-Range Weather Forecast (ECMWF) for spatial variation of these thermodynamic indices. Both 00 UTC and 12 UTC data was analyzed over the eastern Indian regions.
- The thunderstorm information used in this study was obtained from the India Meteorological Department (IMD), Pune.
- Thermodynamic indices was computed which is based on a formulation described in the form of a contingency table by Kunz (2007) and Tyagi et al. (2011).
- The Mann-Kendall test which is a non-parametric statistical test is used with 95% confidence level to comprehend the trends of these thermodynamic indices.

Results

Table 1: Threshold values of Indices according to maximum skill scores of Kolkata

Index	Threshold Values for 00 UTC					Threshold Values for 12 UTC				
	1987-1991	1992-1996	1997-2001	2002-2006	RSS	1987-1991	1992-1996	1997-2001	2002-2006	RSS
LI	≤ -1	≤ -3	≤ -3	≤ -3	0.216	≤ -3	≤ -4	≤ -5	≤ -4	0.250
SHOW	≤ 3	≤ 2	≤ 1	≤ 2	0.293	≤ 4	≤ 2	≤ 3	≤ 4	0.218
CAPE	≥ 762	≥ 558	≥ 791	≥ 1012	0.221	≥ 665	≥ 1248	≥ 1597	≥ 1474	0.249
CIN	≥ -267	≥ -195	≥ -272	≥ -272	0.056	≥ -227	≥ -247	≥ -179	≥ -179	0.028
VTI	≥ 22	≥ 22	≥ 24	≥ 29	0.013	≥ 24	≥ 23	≥ 21	≥ 21	-0.019
BOYD	≥ 93	≥ 96	≥ 97	≥ 95	0.009	≥ 94	≥ 96	≥ 97	≥ 96	-0.046
KI	≥ 24	≥ 29	≥ 27	≥ 20	0.231	≥ 15	≥ 26	≥ 25	≥ 23	0.095
TTI	≥ 44	≥ 43	≥ 44	≥ 44	0.220	≥ 41	≥ 46	≥ 44	≥ 39	0.144
CTI	≥ 15	≥ 15	≥ 17	≥ 18	0.270	≥ 10	≥ 19	≥ 17	≥ 12	0.179
SWEAT	≥ 234	≥ 219	≥ 169	≥ 163	0.257	≥ 87	≥ 215	≥ 198	≥ 167	0.163
BRN	≥ 21	≥ 22	≥ 20	≥ 21	0.184	≥ 18	≥ 28	≥ 26	≥ 24	0.189
HI	≤ 30	≤ 40	≤ 40	≤ 27	0.202	≤ 53	≤ 38	≤ 44	≤ 47	0.136
DPT	≥ 4	≥ 10	≥ 8	≥ 12	0.242	≥ 5	≥ 9	≥ 7	≥ 8	0.176
RH	≥ 47	≥ 55	≥ 47	≥ 36	0.170	≥ 35	≥ 32	≥ 35	≥ 30	0.037

Table 2: Threshold values of Indices according to maximum skill scores of Bhubaneswar


Index	Threshold Values for 00 UTC					Threshold Values for 12 UTC				
	1987-1991	1992-1996	1997-2001	2002-2006	RSS	1987-1991	1992-1996	1997-2001	2002-2006	RSS
LI	≤ -3	≤ -1	≤ -2	≤ -3	0.223	≤ -1	≤ -3	≤ -4	≤ -5	0.257
SHOW	≤ 3	≤ 10	≤ 3	≤ 0	0.256	≤ 6	≤ 3	≤ 1	≤ 2	0.196
CAPE	≥ 184	≥ 244	≥ 558	≥ 628	0.175	≥ 400	≥ 1594	≥ 975	≥ 1790	0.249
CIN	≥ -303	≥ -347	≥ -242	≥ -188	0.095	≥ -174	≥ -216	≥ -199	≥ -188	0.149
VTI	≥ 23	≥ 32	≥ 28	≥ 30	0.054	≥ 19	≥ 20	≥ 26	≥ 22	-0.029
BOYD	≥ 94	≥ 94	≥ 97	≥ 94	-0.032	≥ 96	≥ 95	≥ 97	≥ 94	-0.063
KI	≥ 23	≥ 16	≥ 18	≥ 28	0.189	≥ 21	≥ 26	≥ 30	≥ 28	0.103
TTI	≥ 42	≥ 33	≥ 42	≥ 49	0.207	≥ 38	≥ 41	≥ 49	≥ 41	0.143
CTI	≥ 16	≥ 13	≥ 11	≥ 15	0.246	≥ 11	≥ 18	≥ 17	≥ 19	0.185
SWEAT	≥ 139	≥ 228	≥ 131	≥ 200	0.181	≥ 100	≥ 108	≥ 168	≥ 119	0.140
BRN	≥ 3	≥ 34	≥ 28	≥ 15	0.124	≥ 11	≥ 11	≥ 28	≥ 50	0.167
HI	≤ 30	≤ 35	≤ 42	≤ 37	0.224	≤ 50	≤ 47	≤ 46	≤ 38	0.144
DPT	≥ 9	≥ 10	≥ 7	≥ 9	0.174	≥ 7	≥ 5	≥ 6	≥ 5	0.142
RH	≥ 41	≥ 44	≥ 40	≥ 43	0.125	≥ 47	≥ 41	≥ 36	≥ 40	0.050

Table 3: Threshold values of Indices according to maximum skill scores of Ranchi


Index	Threshold Values for 00 UTC				Threshold Values for 12 UTC			
	1996-2000	2001-2005	2006-2010	RSS	1996-2000	2001-2005	2006-2010	RSS
LI	≤ -3	≤ -2	≤ -2	0.403	≤ -2	≤ -2	≤ -3	0.374
SHOW	≤ 0	≤ 0	≤ -1	0.350	≤ 1	≤ 0	≤ 0	0.320
CAPE	≥ 494	≥ 682	≥ 692	0.407	≥ 486	≥ 604	≥ 730	0.423
CIN	≥ -290	≥ -258	≥ -427	-0.151	≥ -156	≥ -228	≥ -120	-0.060
VTI	≥ 29	≥ 25	≥ 21	0.068	≥ 27	≥ 26	≥ 24	-0.096
BOYD	≥ 96	≥ 96	≥ 95	0.013	≥ 97	≥ 96	≥ 95	-0.098
KI	≥ 27	≥ 29	≥ 26	0.248	≥ 29	≥ 32	≥ 33	0.205
TTI	≥ 49	≥ 48	≥ 43	0.317	≥ 46	≥ 48	≥ 47	0.241
CTI	≥ 29	≥ 25	≥ 21	0.348	≥ 27	≥ 26	≥ 24	0.328
SWEAT	≥ 141	≥ 194	≥ 157	0.319	≥ 153	≥ 186	≥ 160	0.274
BRN	≥ 8	≥ 5	≥ 10	0.348	≥ 12	≥ 8	≥ 18	0.306
DPT	≥ 5	≥ 8	≥ 11	0.158	≥ 2	≥ 7	≥ 10	0.261
RH	≥ 47	≥ 40	≥ 41	0.132	≥ 43	≥ 34	≥ 58	0.149

Summary


- The results for Bhubaneswar reveal that the daytime atmosphere during the pre-monsoon season had enhanced latent instability for thunderstorm days throughout the research. The findings suggest an increase in the extreme severity of severe weather occurrences over the site.
- For Kolkata, The threshold values for both 00 and 12 UTC are fluctuating. CAPE (increasing) and CIN (decreasing) threshold values are contrasting to each other.
- For Ranchi, the environment is favorable to thunderstorm occurrence all of the time, the threshold values reveal that the 12 UTC observations show larger possibilities of thunderstorm occurrence than the 00 UTC observations.
- The findings of temporal variations reveal that the threshold values of thermodynamic indices change in every 5-year interval at all three sites, but not in the same proportion.




CAPE variation over Eastern Indian states (Odisha, West Bengal and Jharkhand) at 00 UTC (top panel) and 12 UTC (bottom panel). The left-hand panel depicts TD variants, whereas the right-hand panel depicts NTD variations.



CIN variation over Eastern Indian states (Odisha, West Bengal and Jharkhand) at 00 UTC (top panel) and 12 UTC (bottom panel). The left-hand panel depicts TD variants, whereas the right-hand panel depicts NTD variations.



Study area, The eastern Indian states (Odisha, West Bengal, and Jharkhand) are color-coded on the India map inset.



CAPE and CIN trend across Odisha at 00 UTC (top panel) and 12 UTC (bottom panel). The left-hand panel depicts TD variants, whereas the right-hand panel depicts NTD variations.

References

- Das, S., Mohanty, U.C., Tyagi, A., Sikla, D.R., Joseph, P.V., Rathore, L.S., Habib, A., Baidya, S.K., Sonam, K. and Sarkar, A., 2014. The SAARC STORM: a coordinated field experiment on severe thunderstorm observations and regional modeling over the South Asian Region. *Bulletin of the American Meteorological Society*, 95(4), pp.603-617.
- Ghosh, A., Lohar, D. and Das, J., 2008. Initiation of Nor'wester in relation to mid-upper and low-level water vapor patterns on METEOSAT-5 images. *Atmospheric Research*, 87(2), pp.116-135.
- Jacovides, C.P. and Yonetani, T., 1990. An evaluation of stability indices for thunderstorm prediction in Greater Cyprus. *Weather and forecasting*, 5(4), pp.559-569.
- Kunz, M., 2007. The skill of convective parameters and indices to predict isolated and severe thunderstorms. *Natural Hazards and Earth System Sciences*, 7(2), pp.327-342.
- Lee, R.R. and Passner, J.E., 1993. The development and verification of TIPS: An expert system to forecast thunderstorm occurrence. *Weather and forecasting*, 8(2), pp.271-280.
- Mukhopadhyay, P., Singh, H.A.K. and Singh, S.S., 2005. Two severe nor'westers in April 2003 over Kolkata, India, using Doppler radar observations and satellite imagery. *Weather*, 60(12), pp.343-353.
- Peppier, R.A., 1988. A Review of Static Stability Indices and Related Thermodynamic Parameters. Illinois State Water Survey Division, SWS Miscellaneous Publication 104.
- Sahu, R.K., Dadich, J., Tyagi, B. and Vissa, N.K., 2020b. Trends of thermodynamic indices thresholds over two tropical stations of north-east India during pre-monsoon thunderstorms. *Journal of Atmospheric and Solar-Terrestrial Physics*, 214, p.105472.
- Sahu, R.K., Dadich, J., Tyagi, B., Vissa, N.K. and Singh, J., 2020a. Evaluating the impact of climate change in threshold values of thermodynamic indices during pre-monsoon thunderstorm season over Eastern India. *Natural Hazards*, 102(3), pp.1541-1569.
- Schultz, P., 1989. Relationships of several stability indices to convective weather events in northeast Colorado. *Weather and Forecasting*, 4(1), pp.73-80.
- Tyagi, B., Krishna, V.N. and Sanyal, A.N.V., 2011. Study of thermodynamic indices in forecasting pre-monsoon thunderstorms over Kolkata during STORM pilot phase 2006-2008. *Natural hazards*, 56(3), pp.491-498.
- Yamane, Y., Hayashi, T., Dewan, A.M., Akter, F. (2010) Severe local convective storms in Bangladesh: part I. *Climatol Atmos Res* 95(4):400-406.