

# A Harmonised Instrumental Earthquake Catalogue for Iceland and the Northern Mid-Atlantic Ridge

Kristján Jónasson<sup>1</sup>, Bjarni Bessason<sup>2</sup>, Ásdís Helgadóttir<sup>1</sup>, Páll Einarsson<sup>3</sup>, Gunnar B. Guðmundsson<sup>4</sup>, Bryndís Brandsdóttir<sup>3</sup>, Kristín S. Vogfjörd<sup>4</sup>, and Kristín Jónsdóttir<sup>4</sup>

<sup>1</sup>Faculty of Industrial Engineering, Mechanical Engineering and Computer Science, University of Iceland, Hjarðarhagi 2, Reykjavík, Iceland

<sup>2</sup>Faculty of Civil and Environmental Engineering, University of Iceland, Hjarðarhagi 2, Reykjavík, Iceland

<sup>3</sup>Science Institute, University of Iceland, Sturlugata 7, Reykjavík, Iceland

<sup>4</sup>Icelandic Meteorological Office, Bústaðavegur 7–9, Reykjavík, Iceland

**Correspondence:** K. Jónasson (jonasson@hi.is)

**Abstract.** A comprehensive catalogue of historical earthquakes, with accurate epicentres and harmonised magnitudes is a crucial resource for seismic hazard mapping. Here we update and combine catalogues from several sources to compile a catalogue of earthquakes in and near Iceland, in the years 1900–2019. In particular the epicentres are based on local information, whereas the magnitudes are based on teleseismic observations, primarily from international on-line catalogues. The most reliable epicentre information comes from the catalogue of the Icelandic Meteorological Office, but this is complemented with information from several technical reports, scientific publications, and newspaper articles. The catalogue contains 1281 moment magnitude ( $M_W$ )  $\geq 4$  events and the estimated completeness magnitude is  $M_W$  5.5 in the first years, going down to  $M_W$  4.5 for recent years. The largest magnitude is  $M_W$  7.0. Such merging of local data and teleseismic catalogues has not been done before for Icelandic earthquakes, and the result is an earthquake map with much more accurate locations than earlier maps. The catalogue also lists additional 5640 earthquakes on the Mid-Atlantic Ridge, north of  $43^\circ$ , with both epicentres and magnitudes determined teleseismically. When moment magnitudes are not available, proxy  $M_W$  values are computed using  $\chi^2$ -regression, normally on the surface-wave magnitude, but exceptionally on the body-wave magnitude. Magnitudes  $M_W \geq 4.5$  have associated uncertainty estimates. The actual combined seismic moment released in the Icelandic earthquakes is found to be consistent with the moment estimated using a simple plate motion model, indicating that the seismic activity of the catalogue period might be typical for any 120 year timespan. The catalogue is named ICEL-NMAR and it is available online on <http://data.mendeley.com>.

## 1 Introduction


Seismic hazard in Iceland is the highest in Northern Europe and is comparable to that in Southern Europe. The seismicity is caused by tectonic movements of the plate boundary of the North American plate and the Eurasian plate crossing the island, as well as volcanic activity (Einarsson, 1991, 2008). Based on historical records, faulting mechanisms, and tectonic context, it can be argued that earthquakes larger than about  $M_W$  7.2 are not to be expected (Halldórsson, 1992a). This is further supported by the limited thickness of the seismogenic part of the Icelandic crust, about 8–12 km (e. g. Stefánsson et al. 1993).

Since the settlement of Iceland in the 8th or 9th century A.D. destructive earthquakes have repeatedly been reported in local chronicles with descriptions of structural damage and fatalities (Sólnes et al., 2013). However, because of the low population density, the losses and number of deaths and injuries has been low and gained little global attention. The main characteristic of the seismicity are shallow ( $< 10$  km) strike-slip earthquakes as well as earthquakes related to volcanic activity. The first instrumentally recorded earthquakes in Iceland occurred in 1896 when six destructive earthquakes struck in South Iceland in a two week period (Ambraseys and Sigbjörnsson, 2000; Sigbjörnsson and Rupakhety, 2014). These events were recorded at several stations in Europe: England, France, Poland and Italy, equipped with rather primitive seismographs (Sólnes et al., 2013, p. 579–583). Damped seismographs, which could measure absolute ground motion, were introduced around the year 1900, allowing (later) magnitude computation. In 1909 a Mainka seismograph was installed in Reykjavik. It was operated until 1914, and again from 1925 when continuous operation was secured (IMO, 1924–2006).

The main motivation behind this study is to construct a catalogue with harmonised magnitudes (which are comparable in both time and space) and reassessed locations for Iceland to use in seismic hazard analysis. A selection criterion for inclusion is that the earthquake was instrumentally recorded by seismic centres outside Iceland and assigned a surface-wave, body-wave, or moment magnitude ( $M_S$ ,  $m_b$  or  $M_W$ ), and that it is listed either in the International Seismological Centre (ISC) Bulletin Event Catalog (ISC, 2020), or in the catalogue of Ambraseys and Sigbjörnsson (2000), which lists and reappraises internationally recorded earthquakes in the region  $62^\circ$ – $68^\circ$ N and  $12^\circ$ – $26^\circ$ W (Fig. 1), in the period 1896–1995. This catalogue will be referred to as the AMB-SIG catalogue. The new catalogue contains reappraised magnitudes and locations for earthquakes in the AMB-SIG region (referred to as ICEL) and the period 1900–2019, a total of 1281 earthquakes. Icelandic earthquakes are almost always less than 12 km deep, but the exact depth information is often not resolvable and therefore the catalogue does not include hypocentral depth.

The magnitudes are all copied or computed from ISC, AMB-SIG, or the Global Centroid Moment Tensor (GCMT) Catalog (GCMT, 2020).  $M_W$  values are provided for all earthquakes. They are of three types: (a) taken directly from the GCMT catalogue if available there (the golden standard), (b) averaged or copied from values in the ISC catalogue, or (c) proxy values computed with regression using  $M_S$  or  $m_b$ . For the regression, region-specific magnitude relationships were developed using data from a larger region, referred to as NMAR. This region follows the Northern Mid-Atlantic Ridge (Fig. 2), and includes all of the region AOI (Atlantic Ocean and Iceland) of Grünthal and Wahlström (2012). A byproduct of our study is therefore a catalogue of 6921 events in the whole NMAR region (including the 1281 ICEL events). Locations of events outside ICEL are copied directly from the ISC catalogue, and magnitudes are obtained in the same way as inside it. The magnitude range of the new catalogue is  $M_W$  4–7.08, as events  $M_W < 4$  were omitted.

For the whole catalogue period local information is crucial for improving earthquake locations. Before 1955, and also for several subsequent events, written sources often provide valuable location information. Since 1955, when three seismometers were installed in Iceland covering the primary seismic zones, locally computed epicenters may be assumed to be more accurate than teleseismic epicenters in international catalogues, which are off by tens of km. One of the innovations in the new catalogue is therefore to use such local data. The primary local sources on epicenters are a catalogue compiled at the Icelandic Meteorological Office (IMO), seismological bulletins, newsletters and reports published by the IMO and the University of




**Figure 1.** The ICCEL region, 62°–68°N and 12°–26°W. The figure shows place names in Iceland mentioned in the article. Towns and villages with 2020 population of at least 800 are also indicated as well as the Tjörnes Fracture Zone (TFZ), the South Iceland Seismic Zone (SISZ), and the Reykjanes Peninsula (RP)

Iceland Science Institute (UISI), journal articles with results of studies on Icelandic earthquakes, and contemporary accounts of earthquakes from newspapers. The origin times are generally taken from the IMO catalogue when available, otherwise from the international catalogues.

60 An early published list of instrumentally recorded earthquakes in Iceland and the surrounding oceans appeared in Gutenberg and Richter's book (1949), p. 196, 207, which lists 60 large earthquakes in the period 1910–1945 in the NMAR region, of these 8 are in the ICCEL region. Six years later Tryggvason (1955) compiled a list of earthquakes  $M \geq 5\frac{1}{4}$  in 1927–1945, 121 in NMAR, of these 22 are in ICCEL. Another global source for earthquakes in the first part of the 20th century is the International Seismological Summary (ISS), the predecessor of the ISC.

65 Since shortly after the IMO was established, it has been responsible for monitoring earthquakes in Iceland. From the beginning, accounts of earthquakes have been published in the IMO monthly newsletter *Veðráttan* (the Weather) (IMO, 1924–2006), in addition the Seismological Bulletin (IMO, 1926–1973) was compiled and distributed to seismological centres abroad, and



**Figure 2.** The NMAR region,  $44^{\circ}$ – $75^{\circ}$ N,  $0^{\circ}$ – $40^{\circ}$ W, and  $67^{\circ}$ – $73^{\circ}$ N,  $0^{\circ}$ – $17^{\circ}$ E. The small part in the eastern hemisphere is added to make the region include all of the AOI region of Grünthal and Wahlström (2012). The ICEL region is also marked in. Four main seismic zones are marked on the map, i. e. Charlie-Gibbs Seismic Zone (CGSZ), South-Iceland Seismic Zone (SISZ), Tjörnes Fracture Zone (TFZ), and Jan Mayen Fracture Zone (JMFZ). The displayed locations and magnitudes are those of the new catalogue.

since 1975 computerised earthquake catalogues have been kept, and made available to scientists working elsewhere. After 1965 earthquake research took off at the University of Iceland, and has flourished ever since with a number of case studies, as well as historical summaries.

The new century has seen a surge in the publication of local and global earthquake catalogues, and Iceland is not an exception. The aforementioned catalogue of Ambraseys and Sigbjörnsson (2000), covers the same ICEL region as the current study and lists 415 earthquakes with  $M_S$  and/or  $m_b$  magnitudes. The epicentres for a portion of these were reassessed, but for the

remaining ones, inaccurate teleseismically determined locations were given. Unfortunately this catalogue was only published  
75 in a very limited distribution, and it is not available online.

Grünthal and Wahlström (Grünthal and Wahlström, 2003) compiled a historical catalogue of earthquakes in Central and Northern Europe until 1993, with magnitudes and locations in Iceland taken from a data file obtained from the IMO. These data were compiled at the IMO independently of the IMO catalogue discussed in Sect. 2.2.1, and are still available on the IMO website ([hraun.vedur.is/ja/ymislegt/storskjalf.html](http://hraun.vedur.is/ja/ymislegt/storskjalf.html)). The locations are reasonably accurate, but the resulting  $M_W$  magnitudes  
80 are exaggerated compared to our results, by up to a whole magnitude unit for some of the early events (the average difference is 0.41 before 1970, 0.37 between 1970 and 1980, and 0.27 after 1980; 3rd quartiles 0.59, 0.47 and 0.36 respectively). The work on this catalogue continued with a number of subsequent projects (Grünthal et al., 2009; Grünthal and Wahlström, 2012; Grünthal et al., 2013), under several acronyms, CENEC (Central, Northern and northwestern European Catalogue), EMEC (European Mediterranean earthquake catalogue), SHARE (seismic hazard harmonization), and SHEEC (SHARE European  
85 earthquake catalogue).

For the Iceland region, all these projects adopted the original 2003 catalogue, adding data (locations and local magnitudes) after 1990 from IMO's catalogue. Among the products of these studies was the "SHARE" hazard map for Europe, where the hazard was greatly overestimated in some places in Iceland, among them in the Reykjavík capital area, where the estimated PGA for a 10% exceedance probability in 50 years is given as 0.4–0.5 g (Woessner et al., 2015). Several recent local studies  
90 estimate 10% 50 year PGA as 0.1–0.2 g in the Reykjavik area (Sólnes et al., 2004; SCI, 2010; Sólnes et al., 2013; D'Amico et al., 2016). The reason for the presumed overestimation is likely a combination of errors in the underlying catalogs and differences in modelling.

In 2010 the ISC initiated work on a global catalogue of large earthquakes since 1904, the ISC-GEM (Global Earthquake Model) Global Instrumental Earthquake Catalogue. The first version was released in 2013 and the work is ongoing, with  
95 version 7 being released in 2020 (Storchak et al., 2013; Di Giacomo et al., 2015). The catalogue contains 40 earthquakes in the ICEL region. It is not used as a source for the new catalogue but instead for quality check and comparison.

Panzer et al. (2016) compiled a catalogue of earthquakes in South-Iceland 1991–2013. It reports locations and magnitudes from IMO's database, cleaned and corrected, as well as proxy  $M_W$ -values based on regression of GCMT-magnitudes on the IMO data, like the CENEC/EMEC catalogues. It has more than 150 000 events with magnitudes down to  $M = 0$ . Unfortunately  
100 the IMO magnitudes are very inaccurate, at least when  $M_W \geq 4$  (Fig. 4), and thus this catalogue has not been used directly in the current work.

The next section discusses the primary sources used to compile the new catalogue. This is followed by a two sections describing how epicenters and magnitudes in the catalogue are determined. The final section contains details of the catalogue, including how to retrieve it, as well as a discussion of completeness magnitude, comparison with ISC-GEM, and comparison  
105 with the total moment of a simple plate motion model.

## 2 Sources and data

This section discusses the primary sources used to compile the new ICEL-NMAR catalogue. These sources consist of four teleseismic international catalogues, used primarily to obtain and/or compute magnitudes, and several types of local Icelandic sources used as a basis for event locations. The local sources include the catalogue of the IMO, scientific publications, seismo-  
110 logical bulletins, newsletters and technical reports, as well as newspaper articles. The section concludes with a few remarks on how individual events in different sources have been matched up.

### 2.1 Teleseismic catalogues

#### 2.1.1 The ISC Bulletin Event Catalog

The ISC database (2020) contains data on earthquake location and magnitude contributed by several seismological agencies  
115 from around the world. For each earthquake a single origin time (UTC) and location but multiple magnitude values are provided. The magnitudes are of several different types, but in the present work only  $M_S$ ,  $m_b$  and  $M_W$  are considered. Magnitudes coded as  $m_S$  and  $M_s$  are treated as  $M_S$ , and similarly for varying capitalization of  $m_b$ . In addition in the period 1955–1970 there are a few magnitude values marked as  $M$  and these are also treated as  $M_S$  cf. (Sykes, 1965). When both  $M$  and  $M_S$  values are available for an earthquake the difference is small. Each magnitude is either marked ISC, to signify that the value  
120 is computed by ISC themselves, or else it is marked with the abbreviation of a submitting agency. The ISC-marked values are referred to as reviewed, and according to Storchak et al. (2017), "seismic events are reprocessed resulting in more robust and reliable mb and MS magnitudes". Di Giacomo and Storchak (2016) say that ISC puts considerable effort into relocating earthquakes and recomputing their magnitudes. They also recommend that preference be given to three agencies, CTBTO (Comprehensive nuclear-Test-Ban Treaty Organization, also known as International Data Centre, IDC, Vienna), MOS (Geophys-  
125 ical Survey of Russian Academy of Sciences, Moscow), and USGS (United States Geological Survey). Among other important agencies is the Swiss Seismological Service, providing the ZUR-RMT (Zurich Moment Tensors) Braunmiller et al. (2002).

#### 2.1.2 The GCMT Earthquake Catalog

The GCMT catalogue (2020) contains data on seismic moment tensors with associated  $M_W$  magnitudes of large earthquakes ( $M_W \geq 5$ ) around the world, starting in 1976 (Dziewonski et al., 1981; Ekström et al., 2012). This is considered to be the most  
130 authoritative catalogue providing  $M_W$  (Di Giacomo and Storchak, 2016). There are 663 events in the NMAR region in this catalogue, and all but 7 of them are also in the ISC catalogue. The GCMT catalog gives  $M_W$  with two decimal places, while ISC gives only one, but apart from that most of the values match between the catalogs.

#### 2.1.3 The catalogue of Ambraseys and Sigbjörnsson

Ambraseys and Sigbjörnsson (2000) published an earthquake catalogue for Iceland or more specifically for the region shown  
135 in Fig. 1. The catalogue covers exactly one century, i. e. from 1896 to 1995, and lists 422 earthquakes. The catalogue is based

on teleseismic data from seismological bulletins, and information from books, journals, newspapers and reports. The authors recalculated surface-wave magnitudes ( $M_S$ ) and locations when possible. Ambraseys and Sigbjörnsson (2000) mention that the greatest outstanding problem was the epicentral accuracy, particularly for pre-1960 macroseismic and instrumental events. They specially remark that epicentres before 1918 reported by the British Association for Advancement of Science (BAAS, 1913–1917) are crude, as well as epicentres of events before 1950 reported by the ISC, although to a lesser degree (Ambraseys and Sigbjörnsson, 2000). This catalogue contains valuable information for the time period from 1900 to 1960 when fewer records are available from other catalogues.

#### 2.1.4 The USGS Earthquake Catalog

A simple online search in the USGS catalogue (2020) provides one magnitude value per earthquake ( $M_W$ ,  $M_S$  or  $m_b$ ), although several magnitude types are often computed. The remaining values are in the ISC database, labelled USGS. Corresponding magnitudes from the two sources are in almost all cases identical. However the locations in the USGS catalogue are different from those in the ISC catalogue, the difference frequently amounting to a few tens of kilometers.

### 2.2 Local sources and catalogues

#### 2.2.1 The catalogue of the Icelandic Meteorological Office

The Icelandic Meteorological Office (IMO) in Reykjavík has been responsible for monitoring earthquakes in Iceland since shortly after its foundation in 1920 when the Mainka seismograph mentioned in the introduction was reinstalled there in 1925. A second Mainka instrument was installed in 1927, also in Reykjavík. Data processing was conducted at the IMO and the results were published in Seismological Bulletins (IMO, 1926–1973) which were sent to several seismological agencies around the world. These results were mainly phase readings and reports of felt earthquakes along with a few locations.

After 1980 the IMO reanalyzed these data and combined them with other local and global sources, e. g. the University of Iceland (UI) reports discussed in the next subsection, and Kárník (1968), to mention a few. The resulting event locations and magnitudes form the basis of IMO's catalogue for the period 1926–1952.

In 1951–1952 three Sprengnether short-period seismographs, measuring all three components of motion, were installed in Reykjavík and the old seismographs were moved to Akureyri in North Iceland and to Vík in South Iceland (Fig. 1), and in the following two decades several more instruments were installed.

As detailed in the next subsection, the University of Iceland Science Institute (UISI) initiated several research projects involving seismic measurements after 1970. Many of these were in cooperation with the IMO, and at the same time IMO's network continued to expand. As before the resulting data were published in the Seismological Bulletins. The IMO catalogue 1952–1974 is based on these and a digital-only bulletin for 1974.

From 1975 to 1986 no bulletins were published, and to fill up this gap, phase readings from the UISI and the IMO stations were merged and reanalyzed to compute locations and magnitudes. This work was carried out at the IMO after 1990, and earthquakes of magnitude  $M_L > 3$  were entered into the IMO database. The database for this period is somewhat preliminary

and incomplete, as manual review is lacking. The period 1987–1990 is also in the IMO database, with results based on *Mánaðaryfirlit jarðskjálfta* (Monthly reports of earthquakes) (IMO, 1987–1990), published by the IMO in cooperation with the UI SI.

In 1991 a digital seismic system, the South Iceland Lowland (SIL) system was implemented by the IMO (Stefánsson et al., 1993; Bödvarsson et al., 1996). As the name implies, it began in South Iceland, but was gradually expanded to cover all geologically active areas in the country. In 2020 around 80 stations are in operation in the SIL-network. Even if the system did not cover the whole island to begin with, all events of magnitude  $M_L > 4$  occurring within a few tens of km offshore should be present for the whole period. Locations and local magnitudes are automatically computed by the system, all automatically located events are manually reviewed, and the location recomputed. The IMO catalogue from 1991 is based on the SIL system analysis.

### 2.2.2 Data from the University of Iceland Science Institute

Research on historical seismicity at the University of Iceland relies heavily on reports by Tryggvason (1978a,b, 1979) and Ottósson (1980). Tryggvason’s reports are based on the early seismographic observations at IMO and overseas for the years 1930–1960, augmented by felt reports and newspaper reports. Ottósson’s report on earthquakes during 1900–1930 is based on felt reports and newspapers, supported by rare teleseismic observations.

Technical advances and increasing interest in crustal activity following the Surtsey eruptions in 1963–1967 led to a proliferation of seismic observations in Iceland in the late 1960ies (Einarsson, 2018). Cooperation started between the UI SI and the Lamont-Doherty Earth Observatory (LDEO) at Columbia University in NY. A team from LDEO came to Iceland with several portable seismographs to study the background seismicity of the mid-Atlantic plate boundary (Ward, 1971). A network of six stations was operated on the Reykjanes Peninsula segment of the boundary during 1971–1976 (Björnsson et al., 2020), augmented by a dense network in the summers of 1971 and 1972 (Klein et al., 1973, 1977). The work continued by building an island-wide network of short-period, vertical component seismographs, designed and built at UI SI. The installation began in South Iceland in 1973 and the network was gradually expanded in the following years, to the Tjörnes Fracture Zone (TFZ) in North Iceland in 1974, and to other parts in 1975–1979. A telemetered network was installed in Central Iceland in 1985. These networks provided valuable data on major events such as the Krafla volcano-tectonic episode of 1975–1984 (Einarsson and Brandsdóttir, 1980; Brandsdóttir and Einarsson, 1979; Buck et al., 2006; Wright et al., 2012), the Hekla eruptions of 1980 and 1991 (Grönvold et al., 1983; Soosalu and Einarsson, 2002) and the Gjálp eruption in Central Iceland in 1996 (Einarsson et al., 1997), as well as the location of the major seismically active structures of Iceland (Einarsson, 1991). After 1991, the analog seismic stations were gradually replaced by the SIL-system discussed in the previous subsection. The last analog stations were dismantled in Central Iceland in 2010. Some of the data gathered by the seismic network discussed above, including epicentres, are documented in the *Skjálftabréf* (Earthquake letter) (UI SI, 1975–1988).

### 2.2.3 Newspapers

200 Newspapers are an important source on earthquakes in Iceland during the first part of the 20th century. The web page <http://timarit.is> provides search access to all newspapers published in Iceland during 1830–2016. News about earthquakes often provide direct or indirect information on their epicentres. In the current work we have used this data source extensively to check the correctness of the sources listed in the previous sections, and when deemed appropriate, to correct earthquake locations for the new catalogue.

### 205 2.3 Combining catalogues

All the catalogues that need to be combined for the current study have their own version of both origin time and location of each earthquake. Jones et al. (2000) and several later publications propose that two records that differ by less than 16 s and 100 km refer to the same earthquake. We have discovered that this is too strict, and use windows of 16 s and 320 km. Increasing the window to 25 s and 1000 km gave identical event pairings. Furthermore, the AMB-SIG catalogue only provides times to the  
210 nearest whole minute, so for that a 90 s time window is used. For each earthquake, the ISC-time, all available locations (ISC, AMB-SIG, IMO, other local sources), and all available magnitude values of different types ( $M_W$ ,  $M_S$ ,  $m_b$ ) and from different catalogues/contributors are entered into a data file. This file is then used for further processing as described below. It contains some smaller earthquakes that are absent from the final catalogue, as explained at the beginning of Section 4 below. The counts of events according to period, region, location source, and magnitude source, in Sections 3.1, 3.2 and 4.1, are however all made  
215 using the catalogue, instead of this data file, as we deem that information to be more relevant for the reader.

## 3 Earthquake locations in the ICEL region

When accurate instrumentally determined location of an earthquake is missing, which applies to a large part of the study period, several methods may be used to determine the epicentre. Sometimes the historical accounts, discussed in Sect. 2.2 provide quite accurate locations, especially in inhabited areas. For the past decades a major effort has been devoted to the mapping of surface  
220 expressions of earthquake faults in Iceland, and these often indicate the location of historical earthquakes (Einarsson, 2015). Furthermore, the main faults tend to produce microearthquakes detected with the SIL network. By relative locations, detailed maps of the subsurface faults can be produced (Slunga et al., 1995). Combining all these methods and adding expert judgement will normally give a much more accurate locations than those provided by the international catalogues, and the same holds for many of the locations in the IMO catalogues, even before 1990. One could say that we have reinterpreted the data with  
225 seismological and tectonic understanding that has been accumulating in recent years and decades.

The remainder of this section describes details of how this methodology has been applied for several subperiods of the study period.

### 3.1 The period until 1990

In the period 1900–1925 there are 22 earthquakes in the ICEL region listed in our final catalogue. All of these are in the AMB-SIG catalogue, and 4 are also in the ISC catalogue, originally coming from Gutenberg and Richter (1949). The authors have viewed all these earthquakes on a map, checked newspapers articles for contemporary accounts of them (using the web service timarit.is mentioned in Sect. 2.2.3), as well as scientific publications, in particular the report of Ottósson (1980). The result of this scrutiny is to use the AMB-SIG location for 14 earthquakes, the aforementioned report for one event, and adjusted the location of 6 events using the methodology described at the beginning of this section. In the new catalogue these location sources have been specified as “Amb-Sig”, “Report” and “New” respectively. Finally, for the 22 January 1910 earthquake we use the location provided by (Stefánsson et al., 2008), 20 km offshore North-Iceland. This source is marked as [1] in the catalogue, with details in an accompanying reference list.

In the period 1926–1954 there are 87 earthquakes in the catalogue, and their location has been scrutinised in the same way. Sometimes we can take into account that an origin time is within a known earthquake series. For this period additional data sources are the IMO catalogue (Sect. 2.2.1), as well as the reports of Tryggvason (1978a,b, 1979) which often provide direct epicentres. This results in using 37 AMB-SIG locations, 9 IMO locations (marked “IMetO” in the new catalogue), 33 locations from the reports, 4 computed as average of the most believable reported locations (marked “Average”), and 4 relocated (marked “New”).

In the period 1955–1990 there are 346 earthquakes in the catalogue. Having multiple local seismometers opens the possibility of computing locations from local measurements. Such locations have found their way into several of our sources, but the quality is variable. There are several journal articles stemming from this period providing locations for 39 earthquakes and our choice is to trust these. The relevant articles are listed in the reference list in the readme-file accompanying the catalogue, and specified as [2], [3], etc. in the catalogue itself. Some of the articles are also cited in Sect. 2.2.2 above. Available locations for the remaining 307 earthquakes were viewed on a map, up to 4 locations per earthquake: From AMB-SIG, IMO, ISC, and one of the earthquake reports, newsletters or bulletins. It transpired that none of these sources could be used as an overall first choice, but instead we had to select the most believable one in each case, or sometimes take an average or relocate. The result was to use AMB-SIG for 64 cases, the IMO catalogue for 94, ISC for 17, 3 from reports, 73 locations from the *Skjálftabréf* (Earthquake letter) (UISI, 1975–1988) (marked “Letter”), 15 averaged, and 40 relocated.

### 3.2 Earthquakes after 1990

For the period 1991–2019 the catalogue contains 826 earthquakes in the ICEL region. With the introduction of the SIL system described in Sect. 2.2.1, the quality of the local epicentre information vastly improved after 1990. We have viewed maps of these locations together with ISC and USGS locations, along with a background layer showing microearthquake activity. From this comparison it was evident that the errors in the teleseismic locations are in many cases tens of kilometers (c. f. Sect. 3.3). The SIL-locations are however accurate to a few km inside the station network, and they are judged to be more accurate than the teleseismic locations in the region  $63^{\circ}\text{N}$ – $67^{\circ}\text{N}$  and  $13^{\circ}\text{W}$ – $25^{\circ}\text{W}$ . Outside this region ISC locations are used for 118 events,

AMB-SIG for 2 and USGS for 1. Inside the region SIL locations are used for 703 events, ISC for 17 (these events are missing in the SIL-catalog), and finally there are 2 events located using published results as detailed in Section 5.2.

### 3.3 Uncertainty of earthquake locations

To get some indication of the uncertainty in event locations in the international catalogues we have looked at the variability between different catalogs, which can be considered as a proxy of the precision of the locations. For 293 events in both the AMB-SIG and the ISC catalogues (period 1910–1996), the median location difference is 10 km, the 90th percentile is 31 km and the maximum is 311 km. The difference does not seem to decrease markedly with time or with earthquake magnitude. A similar comparison between the ISC and the USGS catalogues in the whole NMAR-region (4186 events; 1973–2019) gave a median of 9 km, 90th percentile 25 km, and maximum 284. Comparison of ISC and SIL in the region defined in Section 3.2 (860 events; 1991–2019) gave median 4.4 km, 90th percentile 23 km, and maximum 150 km.

## 4 Earthquake sizes

Contrary to earthquake locations, where local information is crucial, estimating the size of larger earthquakes with teleseismic data is often easier and more reliable than using regional and local data. The dominant periods at teleseismic distances are longer and the structure is smoother due to attenuation of the higher frequencies (Wang et al., 2009; Karimiparidari et al., 2013; Yadav et al., 2009).

Modern earthquake catalogues generally provide moment magnitudes for all earthquakes larger than about  $M_W$  4. For earthquakes whose source mechanism and magnitude have not been modeled by moment tensor inversion of seismic data, regression on surface- or body-wave magnitudes is customarily used to obtain proxy  $M_W$  values, and this procedure is followed here. As mentioned in the introduction, earthquakes from the whole NMAR region are used to construct the  $M_S$ - $M_W$  and  $m_b$ - $M_W$  regression relationships, thus improving the accuracy of these relationships and at the same time getting a larger catalogue of 6921 earthquakes. The data file discussed in Sect. 2.3 above contains some earthquakes that are too small to be included in the catalogue, but are used in the regression in order to improve the relationship for small magnitudes.

For each earthquake there are usually several  $m_b$ -values, contributed by different agencies, and the same applies to  $M_S$ , and sometimes also  $M_W$ . These values must be appropriately averaged or selected before they can be used in the regression. This subtask is dealt with in the next subsection, followed by a subsection on uncertainty in the magnitude estimates in the context of previous studies. Subsection 4.3 discusses the proxy regression, and finally there are two short subsections on the uncertainty in the proxy and local magnitudes.

### 4.1 Best estimates of $M_W$ , $M_S$ and $m_b$

Segja að allar tölur hér að neðan séu án skjálftanna sem er sleppt enda meira relevant.

#### 290 4.1.1 Estimates of $M_W$

In the NMAR region 868 earthquakes in our final catalogue have modeled moment magnitudes, of these 148 are in the ICEL region. The GCMT catalogue is the golden standard for moment magnitudes, and available GCMT  $M_W$  values are used verbatim, 665 in total in the larger NMAR region. The magnitudes range from  $M_W$  4.51 to 7.08, stemming from the period 1976–2019. Additional 204 earthquakes have modeled  $M_W$ -values from other sources, all downloaded via ISC, 201 are  
 295 the ZUR-RMT from the Swiss Seismological Service, all stemming from the period 2000–2005 and 2 are from the USGS catalogue. In addition 61 earthquakes have both a GCMT value and a ZUR-RMT value, with the ZUR-RMT values on average 0.08 magnitudes higher (standard deviation 0.09). The common values are in the range 4.8–6.6 and a graph of  $M_{GCMT}$  against  $M_{ZUR-RMT}$  shows that the relationship is approximately linear with slope 1, which justifies using  $-0.08$  as an agency correction for ZUR-RMT. More precisely, we set  $M_{est} = M_{ZUR-RMT} - 0.08$ , and the estimated values are in the range 3.62–5.22. Gasperini  
 300 (2012) found a very similar result.

Similarly GCMT and USGS have 109 common events, with a correction of 0.00 and standard deviation of 0.08, and we set  $M_{est} = M_{USGS}$  for the 3 events. Other agencies which provide 35 additional  $M_W$  values in the ISC catalogue have been compared with the GCMT catalogue in the same way, but in all cases the standard deviation is too high to include them.

#### 4.1.2 Estimates of $M_S$

305 The final catalogue contains 5050  $M_S$  values for earthquakes in the NMAR region, of these 1080 in the ICEL region. This time the golden standard consists of reviewed values in the ISC catalogue. The situation is somewhat complicated by the fact that three important sources for magnitudes in the first half of the catalogue period have very little overlap with these reviewed values, so that corresponding agency corrections cannot be determined. In fact all sources have small overlap with ISC before 1965. The period has therefore been divided in two, 1900–1964, and 1965–2019.

310 Of the 317  $M_S$  values before 1965, 43 are ISC-reviewed. The remaining 274  $M_S$  values come from a total of 24 other sources, the most important being Ambraseys and Sigbjörnsson (2000), Sykes (1965) (PAL in the ISC catalogue), and the California Institute of Technology in Pasadena (PAS). For each of these earthquakes a direct average of available magnitudes is used.

Of the 4733  $M_S$ -values since 1965, 2828 are ISC-reviewed, again used unchanged. The remaining 1905 events have  $M_S$   
 315 values from a total of 33 sources. After pooling agencies with fewer than 20 events all sources have sufficient overlap with ISC to estimate an agency correction,  $\Delta_i$ , computed as the average of all available differences,  $\delta_i = M_{ISC} - M_i$ , where  $M_i$  is the magnitude estimated by agency  $i$ . When only one source is available,  $M_{est}$  is set to  $M_i + \Delta_i$ , but otherwise a weighted average is computed using

$$M_{est} = \sum_i w_i (M_i + \Delta_i), \quad (1)$$

320 where the  $w_i$  are normalised weights, and the sum is taken over all available  $M_i$ . If the  $\Delta_i$  are independent it is optimal to weigh with their inverse variance, and, even if not optimal, it is more robust to use the same weights when the  $\Delta_i$  are correlated

(Schmelling, 1995). To be precise,  $w_i = (1/\sigma_i^2) / \sum_i (1/\sigma_i^2)$ , where  $\sigma_i$  is the standard deviation of the available  $\delta_i$ . The lowest corrections (0.02–0.04) and the lowest standard deviations (0.10–0.16) are those for AMB-SIG, CTBTO, MOS and USGS. Of the 1905 events without reviewed ISC magnitudes, 93% are contributed by a single agency (the majority from CTBTO), and for (only) 7% of them Eq. (1) is used.

### 4.1.3 Estimates of $m_b$

The catalogue contains 6581 NMAR events with an  $m_b$  value, of these 1128 ICEL events. Again it is beneficial to split the period at year 1965. ISC-reviewed values are once more used when available, for 38 earthquakes out of 63 before 1965 and for 5262 out of 6545 since 1965. Of the 25 remaining earthquakes in the first period Ambraseys and Sigbjörnsson (2000) provide  $m_b$  for 17 events, USGS for 4 events, and for 4 events an average is taken. Of the 1283 remaining earthquakes in the second period there are 44 contributors of  $m_b$  values, the largest being CTBTO and USGS. Final  $m_b$  values are computed as for  $M_S$ : 82% have a single contributor and 18% use Eq. (1). Agency corrections and standard deviations are somewhat higher than for  $M_S$ , typically 0.1–0.2 and 0.15–0.25, respectively.

## 4.2 Uncertainty of magnitude estimates

### 4.2.1 A short survey of uncertainty estimates

Helffrich (1997) discusses the uncertainty of moment magnitudes in the GCMT and USGS catalogues, and his conclusion corresponds to a standard deviation in  $M_W$  of 0.05, 0.04, and 0.10, for deep, intermediate, and shallow events, respectively. Kagan (2003) studies the accuracy of earthquake catalogues extensively. Among his conclusions are the standard deviation of  $M_W$  for both the GCMT and USGS catalogues on the order of 0.05–0.09 for deep to shallow earthquakes, 0.07–0.11 for  $M_W$  6 to 8, and decreasing from 0.11 to 0.06 in the period 1980–2002. Werner (2008) models the magnitude accuracy of 25000 events during 1980–2006 with a Laplace-distribution. The confidence interval presented in the article corresponds to the confidence interval of a normal distribution with  $\sigma = 0.08$ . Finally, Gasperini et al. (2012) conclude with an even lower value,  $\sigma(M_W) = 0.07$ . Many of the estimates cited above are obtained by dividing the standard deviation of magnitude difference between the USGS and the GCMT catalogues by  $\sqrt{2}$ , on the assumption that the errors in them are independent and have the same variance. In reality the errors are probably correlated, so that the cited values may be underestimates of the actual uncertainties.

With a little handwaving Kagan (2003) estimates the uncertainty of  $M_S$  in the ISC catalogue to be about 0.2, and that of  $m_b$  to be about 0.25. In line with these numbers, Kagan also concludes that when  $M_S$  and/or  $m_b$  is turned into proxy  $M_W$ , the uncertainty is about 3–4 times higher than when  $M_W$  is found with moment tensor modeling. This reckoning is supported by both Werner (2008) and Gasperini et al. (2013).

## 4.2.2 Uncertainty of the best estimates

For earthquakes occurring before 1965, there is not enough data to compute the uncertainty objectively, so that a subjective estimate must be used: For this period the uncertainty in  $M_S$  has been set to 0.25, and that in  $m_b$  to 0.35.

After 1964, Eq. (1) is used. Let  $M$  denote the actual magnitude of an earthquake, and  $M_g$  its “golden standard” estimated magnitude (which may be unavailable),  $M_{\text{GCMT}}$  for moment magnitude and  $M_{\text{ISC}}$  for the other two magnitudes. Also, let  $d = M_g - M$ . The uncertainty in  $M_g$ , or standard deviation of  $d$ , is set to

$$\sigma_d = \begin{cases} 0.09 & \text{for moment magnitude} \\ 0.18 & \text{for surface-wave magnitude} \\ 0.23 & \text{for body-wave magnitude} \end{cases} \quad (2)$$

and these numbers are used directly when  $M_g$  is available and  $M_{\text{est}} = M_g$ . Keeping in mind that almost all the earthquakes in the NMAR region are shallow, these uncertainties are perhaps somewhat lower than those quoted in Sect. 4.2.1. However, the accuracy of the global catalogues has probably improved since the quoted studies were carried out, and, furthermore, these studies do not explicitly specify GCMT or reviewed ISC magnitudes.

When  $M_g$  is not available, and  $M_{\text{est}}$  is computed via Eq. (1) the error in the magnitude estimate may be partitioned into several terms:

$$\begin{aligned} M_{\text{est}} - M &= (M_{\text{est}} - M_g) + (M_g - M) \\ &= \sum w_i (M_i + \Delta_i - M_g) + d \\ &= \sum w_i (\Delta_i - \delta_i) + d \end{aligned}$$

using that the  $w_i$  sum to 1. Treating  $d$  and the  $\delta_i$  as random variables, and the  $\Delta_i$  as constants this gives,

$$\text{Var}(M_{\text{est}} - M) = \sigma_d^2 + \sum_i w_i^2 \text{Var} \delta_i + 2 \sum_{i < j} w_i w_j \text{Cov}(\delta_i, \delta_j) - 2 \sum_i w_i \text{Cov}(d, \delta_i)$$

The first term is given by Eq. (2), and  $\text{Var} \delta_i$  and  $\text{Cov}(\delta_i, \delta_j)$  can be approximated by  $\sigma_i^2$  and  $\sigma_{ij}$ , the data covariance of the available pairs  $(\delta_i, \delta_j)$ . Finally, for the last term, we have

$$w_i \text{Cov}(d, \delta_i) = r_i \sigma_d \sigma_i \quad (3)$$

where  $r_i$  is the correlation between  $d$  and  $\delta_i$ . A reasonable constraint is that this correlation is positive: If  $M_g$  overestimates  $M$ , why should  $M_i$  overestimate  $M$  even more? Another constraint is that the estimated variance in  $M_{\text{est}}$  is not smaller than when the golden standard  $M_g$  can be used. The second constraint corresponds to  $r_i = \sigma_i / (2\sigma_d)$ . Selecting the middle road with  $r_i = \sigma_i / (4\sigma_d)$  seems reasonable: it gives  $r_i$  in the range 0.11–0.64; on average 0.28. This choice corresponds to approximating the last term with  $\sum_i w_i^2 \sigma_i^2$ , and the uncertainty estimate:

$$\text{SD}(M_{\text{est}} - M) = \sqrt{\sigma_d^2 - \frac{1}{2} \sum_i w_i^2 \sigma_i^2 + 2 \sum_{i < j} w_i w_j \sigma_{ij}} \quad (4)$$

The root-mean-square (RMS) average uncertainty for all cases where Eq. (1) is used to estimate  $M_W$  is 0.113, for  $M_S$  it is 0.205, and for  $m_b$  0.302.

### 380 4.3 Proxy values for $M_W$

In the New Manual of Seismological Observatory Practice, Bormann et al. (2013) recommend the use of general orthogonal regression to convert between magnitude types when uncertainties in the types differ significantly, as when estimating  $M_W$  from  $M_S$  or  $m_b$ . They also recommend using a nonlinear relationship. An implementation of such a procedure is given by Gasperini et al. (2013) which is based on Stromeyer et al. (2004), and also by Di Giacomo et al. (2015), and we have chosen  
 385 to follow this procedure. A proxy  $M_W$  value is computed from  $M_S$  using

$$M_W^{\text{proxy}} = \exp(a + bM_S) + c, \quad (5)$$

where  $M_S$  is the best estimate of Sect. 4.1,  $a$ ,  $b$  and  $c$  are parameters determined by  $\chi^2$ -regression using Matlab's optimization toolbox and the formulae in Appendix B of Gasperini et al. (2013) (note that the two terms in curly braces in Eq. (B2) in the Appendix should be squared).

390 Bormann et al. (2013) and Di Giacomo et al. (2015) also recommend weighing data points in magnitude ranges with low data frequency higher (histogram equalization). We use a moderately weighted regression of this type: an earthquake with moment and surface-wave magnitudes  $M_W$  and  $M_S$  gets a weight of  $M_W + M_S - 2$ . The effect is that the largest earthquakes weigh about twice as much as the smallest ones.


There is freedom in the regression to fix one of the uncertainties,  $\sigma(M_S)$  or  $\sigma(M_W)$ , and it is also possible to fix their ratio.  
 395 If the ratio is taken as 2.0, as in Gasperini's article, the NMAR data gives  $\sigma(M_S) = 0.176$  and  $\sigma(M_W) = 0.0881$ .

Exactly the same method could be used to compute  $M_W$  from best estimates of  $m_b$ . However the NMAR dataset contains much fewer large earthquakes than the one used by Gasperini et al., so when this is attempted, the relationship turns out to be very slightly concave rather than convex (logarithmic rather than exponential). The nonlinearity is so slight that it can be ignored with a linear model. For earthquakes larger than about  $m_b = 5.75$  an  $M_S$  value is almost always available, and, as  
 400 explained below, preferred. Thus a model valid for  $m_b < 5.75$  is constructed and used:

$$M_W^{\text{proxy}} = a + bm_b, \quad (6)$$

Earthquakes in the Bárðarbunga caldera (Fig. 1) exhibit a different relationship between  $M_W$  and  $m_b$  than the rest of the data set: for the same  $M_W$ , their  $m_b$  is  $\sim 0.15$  higher. Therefore a separate model is used for these earthquakes. The relationship between  $M_W$  and  $M_S$  is also slightly different in the caldera than elsewhere, and for consistency separate models are also used  
 405 in this case. The ratio used by Gasperini et al.,  $\sigma(m_b)/\sigma(M_W) = 2.5$ , gives  $\sigma(m_b) = 0.225$  and  $\sigma(M_W) = 0.0900$ .

As one might expect the deviation in the  $M_S$  model is considerably lower than in the  $m_b$  model (Fig. 3). Thus  $M_S$  is used to compute a proxy  $M_W$  when it is available, for 4217 events in the NMAR region, of which 933 are in the ICEL region. In the absence of an  $M_S$  value the  $m_b$  relation must be used, for 2954 events in NMAR, of which 379 are in ICEL.  $M_S$  is available for almost all large earthquakes, the ones that are important for hazard assessment. Only three  $m_b > 5$ -values are used to compute  
 410 proxy  $M_W$  in the ICEL region and therefore the regression only uses data with  $m_b < 5.5$  (Fig. 3).



**Figure 3.** Magnitude pairs for earthquakes in the Northern Mid-Atlantic Ridge (NMAR) region 1976–2019, exponential relations for  $M_S$  and linear relations for  $m_b$ , all fitted with  $\chi^2$ -regression. There are 733  $M_W$ - $M_S$  pairs outside Bárðarbunga and 95 in it, and 744  $M_W$ - $m_b$  pairs outside and 97 in Bárðarbunga. Note that a few earthquakes with  $m_b < 3.5$ , and thus not included in the final catalogue, are used for the regression. A slight random jitter has been applied to the pairs to avoid superimposing different data points.

**Table 1.** Parameters of exponential and linear models for  $M_W$ , obtained with  $\sigma(M_W) = 0.09$ , c. f. Eqs. (5) and (6), RMSD is the root-mean-square deviation between the model and the y-coordinates of the data, and the last column gives the estimated  $\sigma(m_b)$  and  $\sigma(M_S)$ , respectively.

Model	$a$	$b$	$c$	RMSD	Uncertainty
non-caldera $M_W \sim M_S$	0.850	0.143	0.613	0.142	0.174
non-caldera $M_W \sim m_b$	0.070	1.041		0.257	0.225
caldera $M_W \sim M_S$	-1.401	0.383	3.657	0.072	0.009
caldera $M_W \sim m_b$	-0.585	1.139		0.153	0.111

To use a somewhat round number, and to have a single  $M_W$  uncertainty, the current work uses  $\sigma(M_W) = 0.09$  for all the models,  $m_b$  and  $M_S$ , in and outside Bárðarbunga (Fig. 3, Table 1). These uncertainty values are in good agreement with the results quoted in Sect. 4.2.1, perhaps somewhat lower, which might reflect that our data is more recent and that there is continuous improvement in the quality of the global catalogues.

415 To study possible change in the  $M_S$ - $M_W$  relationship or in the accuracy of the moment tensor  $M_W$  values, a separate modeling was tested for a few sub-periods. A slight, somewhat erratic, improvement in the accuracy was observed, but no significant change in the relationship. Thus it was decided to use a single model for the whole period.

#### 4.4 Uncertainty of the proxy magnitudes

Following Gasperini et al. (2013), the variance of  $M_W^{\text{proxy}}$  for an earthquake obtained with  $M_S$  regression may be estimated  
420 with:

$$\begin{aligned}\sigma_{\text{proxy}}^2 &= (f'(M_S)\sigma_{\text{MS}})^2 + \sigma(M_W)^2 \\ &= b^2 \exp(a + bM_S)^2 \sigma_{\text{MS}}^2 + \sigma(M_W)^2\end{aligned}\quad (7)$$

where  $\sigma_{\text{MS}}^2$  is the variance estimate for the earthquake, obtained as described in Sect. 4.2.2,  $\sigma(M_W) = 0.09$  as in Sect. 4.3,  $f$   
is the model function given in Eq. (5), and  $a$  and  $b$  are the regression parameters (Table 1). The values of  $\sigma_{\text{proxy}}$  computed with  
Eq. (7) are in the range 0.11–0.24, and their RMS-average is 0.15, indicating that only few earthquakes have uncertainty in the  
425 high end of the range. A similar procedure is used in the  $m_b$  regression case and the uncertainties given by the analog of Eq.  
(7) are in the range 0.26–0.53 (RMS-avg. 0.27). For the caldera models, the uncertainty ranges are 0.10–0.18 (RMS-average  
0.11) for  $M_S$  and 0.28–0.39 (RMS-avg. 0.28) for  $m_b$ . Note that because  $f'(M_S) < 1$  then  $\sigma_{\text{proxy}}$  will be smaller than  $\sigma_{\text{MS}}$ .

#### 4.5 Uncertainty in recent local magnitudes


The SIL system described in Sect. 2.2.1 provides two types of local magnitudes, denoted with  $M_L$  and  $M_{LW}$  (Rögnvaldsson  
430 and Slunga, 1993). To assess the uncertainty in these values  $\chi^2$ -regression has been applied, with modeled (non-proxy)  $M_W$ -  
magnitudes on the  $y$ -axis and  $M_L$  and  $M_{LW}$  on the  $x$ -axis with  $\sigma(M_W) = 0.09$ , as in Sect. 4.3 (with caldera earthquakes  
excluded). The resulting estimates are  $\sigma(M_L) = 0.47$  and  $\sigma(M_{LW}) = 0.57$ , far higher than the corresponding values 0.18 for  
 $M_S$  and 0.23 for  $m_b$ . Restricting the comparison to earthquakes onshore Iceland (24 events) gave an improved  $\sigma(M_L) = 0.22$   
but a worse  $\sigma(M_{LW}) = 0.75$ . In all cases there is a considerable negative bias of 0.6–1.4 magnitudes, more offshore (outside  
435 the SIL network) than onshore. One explanation for the large spread and bias of the local magnitudes is that the SIL system's  
analysis is optimised towards robust magnitude estimation of smaller earthquakes than those of this comparison. Figure 4  
shows the spread of the data, evidently in line with these estimates. It has no meaning to show the regression curves because  
of the high uncertainties.

### 5 Results and discussion

440 The primary results of this study is the ICEL-NMAR catalogue, described briefly in the next subsection. Section 5.2 discusses  
the completeness of the catalogue as a function of magnitude and time. Next is a section which compares the new catalogue with  
the ISC-GEM catalogue discussed in the introduction, and finally there is a section with a general discussion. The catalogue  
earthquakes within the region  $63^\circ$ – $67^\circ$ N and  $13^\circ$ – $25^\circ$ W are plotted in Fig. 5.

#### 5.1 The ICEL-NMAR Earthquake Catalogue

445 The new catalogue is available in the Mendeley Data Repository, as the *ICEL-NMAR Earthquake Catalogue, version 2*; see Data  
Availability section below. There are three files, `icel-nmar-v2.txt` with the actual earthquake data, `supporting-info-v2.txt`




**Figure 4.** Moment tensor modeled magnitudes ( $M_W$ ) and two types of local magnitudes computed by the SIL system (see Sect. 2.2.1). Earthquakes in the calderas Bárðarbunga and Katla have been excluded, but apart from that all events with both SIL- and  $M_W$ -magnitudes are included, 24 onshore and 146 offshore. The ZUR-RMT  $M_W$  values were computed by the Swiss Seismological Service 2000–2005.

with meta information, and `origin-time-v2.txt`. For each earthquake `icel-nmar-v2.txt` provides region (ICEL or NMAR), origin time, location,  $M_W$ , the  $M_W$  uncertainty estimated with Eq. (4) or (7) as appropriate, and information on how the  $M_W$  value is computed or what its source is. When available, similar information for  $M_S$  and  $m_b$  are given, and finally information on the origin time and location sources. For completeness, all available origin times are provided in `origin-time-v2.txt`. All events smaller than  $M_W$  4 were excluded and the uncertainty was not computed for  $M_W < 4.5$  because the regression accuracy is reduced at the lower magnitudes. Hypocentral depth is not provided in the catalogue. The brittle part of the Icelandic crust in most areas is less than 12 km thick, and earthquakes of any significance will rupture the whole thickness (Hjaltadóttir, 2010; Pedersen et al., 2003; Stefánsson et al., 1993).

## 455 5.2 Magnitude of completeness

To investigate the magnitude of completeness of the new harmonised catalogue for the whole NMAR region, two methods were used. Firstly, histograms with 10–30 year bins of the earthquake count with magnitudes exceeding different thresholds were created (Fig. 6), and secondly Gutenberg-Richter models were constructed for a few selected periods and minimum magnitudes. The histograms show that the catalogue appears to be complete for  $M_W \geq 6$  for the whole period, for  $M_W \geq 5.5$  since 1915, for  $M_W \geq 5$  since 1970, and for  $M_W \geq 4.5$  since 2000. Gutenberg-Richter modeling with simple declustering (Gardner and Knopoff, 1974) indicate a magnitude of completeness of 5.5 for the whole period, and 4.5 for the period after 1970 (data not shown). For the ICEL region similar histograms indicated a completeness magnitude of 5.5 for the whole period, 5 from 1915, and 4.5 from 1965.

It is interesting to compare the number of large events during the 20th century with lists of historical earthquakes in earlier centuries. Table 2 shows earthquakes with estimated magnitude  $\geq 6$  in Iceland or within 20 km offshore during 1700–1899, in




**Figure 5.** Earthquakes in or near Iceland during 1900–2019 listed in the new catalogue. For the first part of the period, location coordinates are often given in round numbers (tenths of degrees or even half or whole degrees). The map shows slightly jittered locations ( $\leq 3$  km; except when  $M_W > 5.75$ ) to avoid superimposing different events. The magnitude range for the smallest earthquakes is  $M_W$  4–4.25. For the other ranges the central value is specified, so that e. g.  $M_W \sim 4.5$  implies the range 4.25–4.75. The largest event is  $M_W$  7.00 in the TFZ at  $18^\circ$ W. The indicated tectonic features are explained in Figure 2.

total 17 events. In the new catalogue there are 8 earthquakes with  $M_W \geq 6$  in the 20th century in the same region, and 4 more in the first two decades of the 21st century.

In the final catalogue there are a few periods with disproportionately many earthquakes connected to tectonic activity (SISZ 2000 and 2008) and volcanic activity (Krafla region 1975–1976, Hengill 1994–1999, Bárðarbunga 2014–2015).

470 In the wake of large earthquakes it is possible that other events are triggered by their propagating waves. These secondary events can be missing from the international catalogues because their signal is lost in the coda of the primary event at teleseismic distances. An example of this are two events on the Reykjanes Peninsula triggered by the  $M_W$  6.52 South-Iceland event on 2000-06-17 15:40:41, occurring 26 and 30 seconds later, and 65 and 80 km farther west, respectively. The size of the first one was estimated to be  $M_L$  5.5 (Antonioli et al., 2006), and that of the second one  $M_W$  5.79 (Pagli et al., 2003). Our estimated  
 475  $M_W$  for the first event is 5.5, and both  $M_W$  values have been added to the new catalogue with uncertainties of 0.4 and 0.2, respectively. These are the only events not coming from one of the four teleseismic catalogues of Sect. 2.1.



**Figure 6.** Count of earthquakes in the NMAR region exceeding different  $M_W$  thresholds according to period.

### 5.3 Comparison with the ISC-GEM catalogue

Version 7.0 of the ISC-GEM catalogue was released in 2020. In the NMAR region it contains much fewer events than our new catalogue (168, with  $M_W$  in the range 5.42–7.00), and no local information is used to relocate them. Non-proxy  $M_W$  magnitudes in ISC-GEM and the current catalogue are identical, but in general the proxy values differ, both because ISC-GEM uses a different regression model and because the underlying  $M_S$  and  $m_b$  data may differ. The difference in the more important  $M_S$  regression curves is slight. Comparing Fig. 3 and the corresponding figure in (Di Giacomo et al., 2015) for  $M_S = 5$  the ISC-GEM curve is 0.06 higher, for  $M_S = 6$  it is 0.02 lower and for  $M_S = 7$  it is 0.05 lower.

There are 119 earthquakes with proxy  $M_W$  common to the catalogues, of these 30 in the ICEL region. Their ISC-GEM magnitudes are on average 0.06 lower than the ones presented here. The largest absolute difference is 0.47 and for 85 events the difference is less than 0.2. For the ICEL region the mean difference is 0.02, the largest absolute one is 0.26, and there are 24 events which differ by less than 0.2 magnitudes.

A few events which differ most were investigated, and it transpired that the explanation was usually a combined effect of the regression curve difference and the underlying data difference.

### 5.4 Cumulative seismic moment and the earthquake cycle

The question arises how representative the seismic activity of the catalogue period is for any period of 120 years. The answer depends on the length of the typical earthquake cycle. If the cycle is significantly longer than 120 years our sample may underestimate the seismicity greatly, e. g., if the period does not contain a characteristic maximum magnitude earthquake. Studies of South Iceland earthquakes indicate that we may be near this critical duration of the cycle. The study of Einarsson et al. (1981) gave an average time between major earthquake sequences of about 80 years, ranging between 45 and 112 years.

**Table 2.** Historical large earthquakes in Iceland in the 18th and 19th centuries. The magnitude estimates are based on the resulting damage (Halldórsson, 1992b; Stefánsson et al., 2008; Sólnes et al., 2013). The epicentral locations are approximate but overall the longitude is more accurate than the latitude since in most cases N-S surface faults have been mapped and linked to the largest events. Note that these earthquakes are not included in the new catalogue.

Date	Lat.	Lon.	$M_S$	$M_W^{\text{proxy}}$
1706, April	63.9	21.2	6.0	6.1
1732, Sept.	64.0	20.0	6.7	6.7
1734, March	63.9	20.8	6.8	6.8
1755, Sept.	66.1	17.6	7.0	7.0
1766, Sept.	63.9	21.2	6.0	6.1
1784, August	63.9	20.5	7.1	7.1
1784, August	63.9	21.0	6.7	6.7
1829, Feb.	63.9	20.0	6.0	6.1
1838, June	66.3	18.8	6.5	6.5
1872, April	66.1	17.4	6.5	6.5
1872, April	66.2	17.9	6.5	6.5
1885, Jan.	66.3	16.9	6.3	6.4
1896, August	64.0	20.1	6.9	6.9
1896, August	64.0	20.3	6.7	6.7
1896, Sept.	63.9	21.0	6.0	6.1
1896, Sept.	64.0	20.6	6.5	6.5
1896, Sept.	63.9	21.2	6.0	6.1

Stefánsson and Halldórsson (1988) concluded that the South Iceland Seismic Zone (SISZ) had a total release of accumulated strain in about 140 years. Decriem et al. (2010) estimated the accumulated strain by plate movements since the 1896–1912 earthquakes and compared to the released seismic moment during the earthquakes of 2000 and 2008. They found that only about half of the strain had been released by these events.

500 For comparison with our catalogue we estimate the potential seismic moment release in the two fracture zones, the SISZ and the TFZ, by a simplified geometric model of two transform faults parallel to the relative plate motion. The simplification is justified by the arguments of Sigmundsson et al. (1995), who showed that the seismic moment of many closely spaced, short transverse faults (bookshelf faults) is equivalent to that released by a single transform fault. We also assume that almost all the seismic moment is released by the transform zones and not by the divergent segments of the plate boundary or the magmatically  
505 induced seismicity. The length of the transform zones is taken as 180 km and 150 km for the South and North Iceland zones, respectively, i. e. the offset of the ridge axes. The width of the fault is taken to be the thickness of the seismogenic part of the crust, about 10 km, below which the slip is assumed to be aseismic. The spreading rate is 19 mm/yr, and the shear modulus

$20 \cdot 10^9$  Pa (McGarr and Barbour, 2018). The moment rate will then be:

$$20 \cdot 10^9 \times 19 \cdot 10^{-3} \times 330 \cdot 10^3 \times 10 \cdot 10^3 = 1.25 \cdot 10^{18} \text{ Nm/yr.} \quad (8)$$

510 This result can be compared with the total seismic moment released in Iceland during the catalogue period, which may be estimated using the catalogue data and the completeness information of Sect. 5.2. Such computation for all earthquakes  $\geq M_W$  4 in the area shown in Fig. 6, excluding the Reykjanes Ridge and Bárðarbunga, gives a total of  $1.61 \cdot 10^{20}$  Nm. Adding a simple correction for smaller events assuming the Gutenberg-Richter law with  $b = 1$  raises the estimate to  $1.64 \cdot 10^{20}$  Nm, corresponding to an annual rate of  $1.37 \cdot 10^{18}$  Nm/yr. This agrees quite (even surprisingly) well with the result of Eq. (8).

## 515 5.5 General discussion

We have constructed a new catalogue of earthquakes in Iceland and, as a byproduct, for the Northern Mid-Atlantic Ridge. A general criterion for entry into the catalogue is that an earthquake has been instrumentally recorded by agencies outside Iceland. Locations of events in the ICEL region (Fig. 1) have been reassessed and proxy  $M_W$  values for earthquakes without modeled moment magnitudes have been computed. The resulting moment magnitudes range from 4 to 7.08. For the ICEL region the  
520 catalogue is reasonably complete for  $M_W \geq 5.5$  for the whole period. There are 36 earthquakes of this size onshore or less than 20 km offshore, i. e. 2.8 per decade, and of these 10 have  $M_W \geq 6$ , i. e. 0.8 per decade.

To our knowledge, the map in Fig 5 is the first earthquake map of Iceland which is not substantially confounded by misplaced events. The locations of the two large TFZ-events marked with a star in Fig. 5 (the easternmost 1910 and the westernmost 1963) are still uncertain and controversial. Neither of them appears to have occurred on the best known structures, the Húsavík-Flatey  
525 fault or the Grímsey Oblique Rift. Stefánsson et al. (2008) suggest that the 1963 event originated on a NNE-striking fault offshore Skagafjörður, based on the distribution of recent earthquakes and the focal mechanism solutions of Stefánsson (1966) and Sykes (1967). They furthermore suggest that the 1910 event originated on the eastern margin of the Grímsey Shoal. We adopt these locations in our catalogue. Distribution of epicentres and recent bathymetric data support these suggestions (Einarsson et al., 2019).

530 The largest events occur in the two seismic zones, where the plate boundaries are parallel to the plate movements (Fig. 1 and 5). The distance from these events to the Reykjavik capital area, where 63% of the population live, is some tens of kilometers, and the same holds for Akureyri in North Iceland, with 5% of the population. However there are several towns and villages within the zones. An important future task is to carry out a detailed analysis of the seismic hazard both in these urban areas and elsewhere in Iceland. The new catalogue should prove to be an essential resource for such seismic hazard mapping.

535 *Data availability.* The international earthquake catalogues from USGS, GCMT and ISC are freely available online. In addition we used the catalogue of Ambraseys and Sigbjörnsson (2000), as well as scattered data on individual earthquakes from various printed sources, as detailed in Sect. 2. We also used the Icelandic Meteorological Office catalogue for the period 1926–2019. Work is currently underway to put at least part of this catalogue online. The new catalogue is available in the Mendeley data repository as the ICEL-NMAR Earthquake Catalogue (<http://dx.doi.org/10.17632/7zh6xg22cv.2>). Version 3, including the most recent events, is forthcoming.

540 *Author contributions.* K Jónasson and B Bessason prepared the manuscript with contributions from all authors, K Jónasson and Á Helgadóttir wrote software for data processing, P Einarsson, GB Gudmundsson, K Jónasson, B Bessason, B Brandsdóttir and Kristín Vogfjörd reappraised event locations.

*Acknowledgements.* We thank all the people and institutions who have set up seismometers, gathered data from these, and used them to compute earthquake locations and magnitudes. Without their contribution this work would not have been possible. The public datasets that we have used are available from three open web sources: the ISC Bulletin Event Catalog (2020); the GCMT Earthquake Catalog (2020); and the USGS Earthquake Catalog (2020). Finally, we use the earthquake catalogue of the Icelandic Meteorological Office for the period 1926–2019.

Contributions of Kristín Vogfjörd were supported by the RISE project, grant number 821115 under the EC Horizon2020 programme.

## References

- 550 Ambraseys, N. N. and Sigbjörnsson, S.: Re-appraisal of the seismicity in Iceland, Earthquake Engineering Research Centre, University of Iceland, Selfoss, Iceland, 2000.
- Antonioli, A., Belardinelli, M., Bizzarri, A., and Vogfjord, K.: Evidence of instantaneous dynamic triggering during the seismic sequence of year 2000 in south Iceland, *Journal of Geophysical Research: Solid Earth*, 111, 2006.
- BAAS: Seismological Committee, quarterly issues, British Association for the Advancement of Science, 1913–1917.
- 555 Björnsson, S., Einarsson, P., Tulinius, H., and Hjartardóttir, A. R.: Seismicity of the Reykjanes Peninsula 1971–1976, *Journal of Volcanology and Geothermal Research*, 391, 2020.
- Bormann, P., Wendt, S., and DiGiacomo, D.: Chapter 3: Seismic Sources and Source Parameters, in: *New Manual of Seismological Observatory Practice 2 (NMSOP-2)*, edited by Bormann, P., pp. 1–259, Deutsches GeoForschungszentrum GFZ, Potsdam, 2013.
- Brandsdóttir, B. and Einarsson, P.: Seismic activity associated with the September 1977 deflation of the Krafla central volcano in northeastern  
560 Iceland, *Journal of Volcanology and Geothermal Research*, 6, 197–212, 1979.
- Braunmiller, J., Kradolfer, U., Baer, M., and Giardini, D.: Regional moment tensor determination in the European–Mediterranean area—initial results, *Tectonophysics*, 356, 5–22, 2002.
- Buck, W. R., Einarsson, P., and Brandsdóttir, B.: Tectonic stress and magma chamber size as controls on dike propagation: Constraints from the 1975–1984 Krafla rifting episode, *Journal of Geophysical Research: Solid Earth*, 111, 2006.
- 565 Bödvarsson, R., Rögnvaldsson, S. T., Jakobsdóttir, S., Slunga, R., and Stefánsson, R.: The SIL data acquisitional and monitoring system, *Seismological Research Letters*, 67, 35–47, 1996.
- Decriem, J., Árnadóttir, T., Hooper, A., Geirsson, H., Sigmundsson, F., Keiding, M., Ófeigsson, B. G., Hreinsdóttir, S., Einarsson, P., LaFemina, P., , and Bennett, R. A.: The 2008 May 29 earthquake doublet in SW Iceland, *Geophysical Journal International*, 181, <https://doi.org/10.1111/j.1365-246X.2010.04565.x>, 2010.
- 570 Di Giacomo, D. and Storchak, D. A.: A scheme to set preferred magnitudes in the ISC Bulletin, *Journal of Seismology*, 20, 555–567, 2016.
- Di Giacomo, D., Bondár, I., Storchak, D. A., Engdahl, E. R., Bormann, P., and Harris, J.: ISC-GEM: Global instrumental earthquake catalogue (1900–2009), III. Re-computed MS and mb, proxy M W, final magnitude composition and completeness assessment, *Physics of the Earth and Planetary Interiors*, 239, 33–47, 2015.
- Dziewonski, A. M., Chou, T.-A., and Woodhouse, J. H.: Determination of earthquake source parameters from waveform data for studies of  
575 global and regional seismicity, *Journal of Geophysical Research: Solid Earth*, 86, 2825–2852, 1981.
- D’Amico, V., Albarello, D., Sigbjörnsson, R., and Rupakhety, R.: Seismic hazard assessment for Iceland in terms of macroseismic intensity using a site approach, *Bulletin of Earthquake Engineering*, 14, 1797–1811, 2016.
- Einarsson, P.: Earthquakes and present-day tectonism in Iceland, *Tectonophysics*, 189, 261–79, 1991.
- Einarsson, P.: Plate boundaries, rifts and transforms in Iceland, *Jökull*, 58, 35–58, 2008.
- 580 Einarsson, P.: Mechanisms of earthquakes in Iceland, in: *Encyclopedia of Earthquake Engineering*, edited by Beer, M., Patelli, E., Kougioumtzoglou, I. A., and Au, S. K., Springer, 2015.
- Einarsson, P.: Short-term seismic precursors to Icelandic eruptions 1973–2014, *Frontiers in Earth Science*, 6, 45, 2018.
- Einarsson, P. and Brandsdóttir, B.: Seismological evidence for lateral magma intrusion during the July 1978 deflation of the Krafla volcano in NE-Iceland, *Journal of Geophysics*, 47, 160–165, 1980.

- 585 Einarsson, P., Björnsson, S., Foulger, G., Stefánsson, R., and Skaftadóttir, T.: Seismicity pattern in the South Iceland seismic zone, in: Earthquake Prediction: An International Review, edited by Simpson, D. and Richards, P., vol. 4, pp. 141–151, Wiley Online Library, 1981.
- Einarsson, P., Brandsdóttir, B., Gudmundsson, M. T., Björnsson, H., Grönvold, K., and Sigmundsson, F.: Center of the Iceland hotspot experiences volcanic unrest, *Eos, Transactions American Geophysical Union*, 78, 369–375, 1997.
- Einarsson, P., Brandsdóttir, B., and Ásta Rut Hjartardóttir: Seismicity, faults, and bathymetry of the Tjörnes Fracture Zone, in: Proceeding of  
590 the Northquake 2019 Workshop, Husavik Academic Centre, Húsavík, Iceland, 2019.
- Ekström, G., Nettles, M., and Dziewoński, A. M.: The global CMT project 2004–2010: Centroid-moment tensors for 13,017 earthquakes, *Physics of the Earth and Planetary Interiors*, 200, 1–9, 2012.
- Gardner, J. K. and Knopoff, L.: Is the sequence of earthquakes in Southern California, with aftershocks removed, Poissonian?, *Bulletin of the Seismological Society of America*, 64, 1363–1367, 1974.
- 595 Gasperini, P., Lolli, B., Vannucci, G., and Boschi, E.: A comparison of moment magnitude estimates for the European-Mediterranean and Italian regions, *Geophysical Journal International*, 190, 1733–1745, 2012.
- Gasperini, P., Lolli, B., and Vannucci, G.: Body-Wave Magnitude  $m_b$  is a Good Proxy of Moment Magnitude  $M_W$  for Small Earthquakes ( $m_b < 4.5–5.0$ ), *Seismological Research Letters*, 84, 932–937, 2013.
- GCMT: Global Centroid Moment Tensor Earthquake Catalog, downloaded on 25 Nov. 2020, <https://www.globalcmt.org>, 2020.
- 600 Grönvold, K., Larsen, G., Einarsson, P., Thorarinsson, S., and Sæmundsson, K.: The Hekla eruption 1980–1981, *Bulletin Volcanologique*, 46, 349–363, 1983.
- Grünthal, G. and Wahlström, R.: An  $M_w$  based earthquake catalogue for central, northern and northwestern Europe using a hierarchy of magnitude conversions, *Journal of Seismology*, 7, 507–531, 2003.
- Grünthal, G. and Wahlström, R.: The European-Mediterranean earthquake catalogue (EMEC) for the last millennium, *Journal of Seismology*,  
605 16, 535–570, 2012.
- Grünthal, G., Wahlström, R., and Stromeyer, D.: The unified catalogue of earthquakes in central, northern, and northwestern Europe (CENEC)—updated and expanded to the last millennium, *Journal of Seismology*, 13, 517–541, 2009.
- Grünthal, G., Wahlström, R., and Stromeyer, D.: The SHARE European earthquake catalogue (SHEEC) for the time period 1900–2006 and its comparison to the European-Mediterranean earthquake catalogue (EMEC), *Journal of Seismology*, 17, 1339–1344, 2013.
- 610 Gutenberg, B. and Richter, C. F.: *Seismicity of the Earth*, Princeton University Press, Princeton, New Jersey, 1949.
- Halldórsson, P.: Seismic hazard assessment based on historical data and seismic measurements, in: Proceedings of the International Conference on Preparedness and Mitigation for Natural Disasters '92, pp. 53–63, Association of Chartered Engineers in Iceland, Reykjavík, 1992a.
- Halldórsson, P.: Um jarðskjálftasvæði Suðurlands (About the South Iceland seismic zone), *Árbók VFÍ (Icelandic Journal of Engineering)*, 4,  
615 226–239, 1992b.
- Helffrich, G. R.: How good are routinely determined focal mechanisms? Empirical statistics based on a comparison of Harvard, USGS and ERI moment tensors, *Geophysical Journal International*, 131, 741–750, 1997.
- Hjaltadóttir, S.: Use of relatively located microearthquakes to map fault patterns and estimate the thickness of the brittle crust in Southwest Iceland, Tech. Rep. VI-2010-003, Icelandic Meteorological Office, Reykjavík, 2010.
- 620 IMO: Veðráttan (the Weather), Icelandic Meteorological Office, Reykjavík, <https://timarit.is>, 1924–2006.
- IMO: Seismological Bulletin, Icelandic Meteorological Office, Reykjavík, 1926–1973.
- IMO: Mánaðaryfirlit jarðskjálfta (Monthly reports of earthquakes), Icelandic Meteorological Office, Reykjavík, 1987–1990.

- ISC: International Seismological Centre Bulletin Event Catalog, downloaded on 25 Nov. 2020, <http://www.isc.ac.uk/iscbulletin/search/>, 2020.
- 625 Jones, A., Michæl, A., Simpson, B., Jacob, S., and Oppenheimer, D.: Rapid distribution of earthquake information for everybody, *Seismological Research Letters*, 71, 355–358, 2000.
- Kagan, Y. Y.: Accuracy of modern global earthquake catalogs, *Physics of the Earth and Planetary Interiors*, 135, 173–209, 2003.
- Karimiparidari, S., Zare, M., Memarian, H., and Kijko, A.: Iranian earthquake, a uniform catalog with moment magnitudes, *Journal of Seismology*, 17, 897–911, 2013.
- 630 Klein, F. W., Einarsson, P., and Wyss, M.: Microearthquakes on the mid-Atlantic plate boundary on the Reykjanes Peninsula in Iceland, *Journal of Geophysical Research*, 78, 5084–5099, 1973.
- Klein, F. W., Einarsson, P., and Wyss, M.: The Reykjanes Peninsula, Iceland, earthquake swarm of September 1972 and its tectonic significance, *Journal of Geophysical Research*, 82, 865–888, 1977.
- Kárník, V.: Seismicity of the European area, Part 1, Academia, Praha, 1968.
- McGarr, A. and Barbour, A. J.: Injection-induced moment release can also be aseismic, *Geophysical Research Letters*, 45, 5344–5351, 2018.
- 635 Ottósson, K. G.: Jarðskjálftar á Íslandi 1900–1929 (Earthquakes in Iceland 1900–1929), Tech. Rep. RH-80-05, University of Iceland Science Institute, Reykjavík, 1980.
- Pagli, C., Pedersen, R., Sigmundsson, F., and Feigl, K. L.: Triggered fault slip on June 17, 2000 on the Reykjanes Peninsula, SW-Iceland captured by radar interferometry, *Geophysical Research Letters*, 30, 2003.
- 640 Panzera, F., Zechar, J. D., Vogfjörd, K. S., and Eberhard, D.: A Revised earthquake catalogue for South Iceland, *Pure and Applied Geophysics*, 173, 97–116, 2016.
- Pedersen, R., Jónsson, S., Árnadóttir, T., Sigmundsson, F., and Feigl, K. L.: Fault slip distribution of two June 2000  $M_w$ 6.5 earthquakes in South Iceland estimated from joint inversion of InSAR and GPS measurements, *Earth and Planetary Science Letters*, 213, 487–502, 2003.
- Rögnvaldsson, S. T. and Slunga, R.: Routine fault plane solutions for local networks: a test with synthetic data, *Bulletin of the Seismological Society of America*, 83, 1232–1247, 1993.
- 645 Schmelling, M.: Averaging correlated data, *Physica Scripta*, 51, 676, 1995.
- SCI: Icelandic National Annexes to Eurocodes, Standards Council of Iceland, Reykjavík, 2010.
- Sigbjörnsson, R. and Rupakhety, R.: A saga of the 1896 South Iceland earthquake sequence: magnitudes, macroseismic effects and damage, *Bulletin of earthquake engineering*, 12, 171–184, 2014.
- Sigmundsson, F., Einarsson, P., Bilham, R., and Sturkell, E.: Rift-transform kinematics in south Iceland: Deformation from Global Positioning System measurements, 1986 to 1992, *Journal of Geophysical Research: Solid Earth*, 100, 6235–6248, 1995.
- 650 Slunga, R., Rögnvaldsson, S. T., and Bödvarsson, R.: Absolute and relative locations of similar events with application to microearthquakes in southern Iceland, *Geophysical Journal International*, 123, 409–419, 1995.
- Soosalu, H. and Einarsson, P.: Earthquake activity related to the 1991 eruption of the Hekla volcano, Iceland, *Bulletin of Volcanology*, 63, 536–544, 2002.
- 655 Stefánsson, R.: Methods of focal mechanism studies with application to two Atlantic earthquakes, *Tectonophysics*, 3, 209–243, 1966.
- Stefánsson, R. and Halldórsson, P.: Strain release and strain build-up in the South Iceland seismic zone, *Tectonophysics*, 152, 267–276, 1988.
- Stefánsson, R., Bödvarsson, R., Slunga, R., Einarsson, P., Jakobsdóttir, S., Bungum, H., Gregersen, S., Havskov, J., Hjelme, J., and Korhonen, H.: Earthquake prediction research in the South Iceland Seismic Zone and the SIL project, *Bulletin of the Seismological Society of America*, 83, 696–716, 1993.

- 660 Stefánsson, R., Gudmundsson, G. B., and Halldórsson, P.: Tjörnes fracture zone. New and old seismic evidences for the link between the North Iceland rift zone and the Mid-Atlantic ridge, *Tectonophysics*, 447, 117–126, 2008.
- Storchak, D. A., Di Giacomo, D., Bondár, I., Engdahl, E. R., Harris, J., Lee, W. H. K., Villaseñor, A., and Bormann, P.: Public release of the ISC-GEM global instrumental earthquake catalogue (1900–2009), *Seismological Research Letters*, 84, 810–815, 2013.
- Storchak, D. A., Harris, J., Brown, L., Lieser, K., Shumba, B., Verney, R., Di Giacomo, D., and Korger, E. I. M.: Rebuild of the Bulletin of  
665 the International Seismological Centre (ISC), part 1: 1964–1979, *Geoscience Letters*, 4, 32, 2017.
- Stromeyer, D., Grünthal, G., and Wahlström, R.: Chi-square regression for seismic strength parameter relations, and their uncertainties, with applications to an  $M_W$  based earthquake catalogue for central, northern and northwestern Europe, *Journal of Seismology*, 8, 143–153, 2004.
- Sykes, L. R.: The seismicity of the Arctic, *Bulletin of the Seismological Society of America*, 55, 501–518, 1965.
- 670 Sykes, L. R.: Mechanism of earthquakes and nature of faulting on the mid-oceanic ridges, *Journal of Geophysical Research*, 72, 2131–2153, 1967.
- Sólnes, J., Sigbjörnsson, R., and Elíasson, J.: Probabilistic seismic hazard mapping of Iceland, in: *Proceedings of the 13th World conference on earthquake engineering*, Vancouver, BC, Canada, 2004.
- Sólnes, J., Sigmundsson, F., and Bessason, B., eds.: *Natural Hazard in Iceland, Volcanic Eruptions and Earthquakes*, University of Iceland  
675 Press and Natural Catastrophe Insurance of Iceland, Reykjavík, 2013.
- Tryggvason, E.: Jarðskjálftar á Íslandi og nyrzta hluta Atlantshafsins (Seismicity of Iceland and the surrounding oceans), *Náttúrufræðingurinn*, 25, 194–197, 1955.
- Tryggvason, E.: Jarðskjálftar á Íslandi 1930–1939 (Earthquakes in Iceland 1930–1939), Tech. Rep. RH-78-21, University of Iceland Science Institute, Reykjavík, 1978a.
- 680 Tryggvason, E.: Jarðskjálftar á Íslandi 1940–1949 (Earthquakes in Iceland 1940–1949), Tech. Rep. RH-78-22, University of Iceland Science Institute, Reykjavík, 1978b.
- Tryggvason, E.: Jarðskjálftar á Íslandi 1950–1959 (Earthquakes in Iceland 1950–1959), Tech. Rep. RH-79-06, University of Iceland Science Institute, Reykjavík, 1979.
- UISI: Skjálftabréf (Earthquake Letter), University of Iceland Science Institute, Reykjavík, <https://jardvis.hi.is/skjalftabref>, 1975–1988.
- 685 USGS: United States Geographical Survey Earthquake Catalog, downloaded on 25 Nov. 2020, <https://earthquake.usgs.gov/earthquakes/search/>, 2020.
- Wang, Q., Jackson, D. D., and Kagan, Y. Y.: California earthquakes, 1800–2007: A unified catalog with moment magnitudes, uncertainties, and focal mechanisms, *Seismological Research Letters*, 80, 446–457, 2009.
- Ward, P. L.: New interpretation of the geology of Iceland, *Geological Society of America Bulletin*, 82, 2991–3012, 1971.
- 690 Werner, M. J.: Magnitude uncertainties impact seismic rate estimates, forecasts, and predictability experiments, *Journal of Geophysical Research - Solid Earth*, 113, 1–20, 2008.
- Woessner, J., Laurentiu, D., Giardini, D., Crowley, H., Cotton, F., Grünthal, G., Valensise, G., Arvidsson, R., Basili, R., Demircioglu, M. B., et al.: The 2013 European seismic hazard model: key components and results, *Bulletin of Earthquake Engineering*, 13, 3553–3596, 2015.
- Wright, T. J., Sigmundsson, F., Pagli, C., Belachew, M., Hamling, I. J., Brandsdóttir, B., Keir, D., Pedersen, R., Ayele, A., Ebinger, C., et al.:  
695 Geophysical constraints on the dynamics of spreading centres from rifting episodes on land, *Nature Geoscience*, 5, 242–250, 2012.
- Yadav, R., Bormann, P., Rastogi, B. K., Das, M. C., and Chopra, S.: A homogeneous and complete earthquake catalog for northeast India and the adjoining region, *Seismological Research Letters*, 80, 609–627, 2009.